**Loss of mitochondrial pyruvate transport initiates cardiac glycogen accumulation and heart failure**

Rachel C. Weiss, B.S.,[1] Kelly D. Pyles, B.S.,[1] Kevin Cho, Ph.D.,[2]
Michelle Brennan, Ph.D.,[1] Jonathan S. Fisher, Ph.D.,[3] Gary J. Patti, Ph.D.,[2]
Kyle S. McCommis, Ph.D.[1]*

[1] Edward A. Doisy Department of Biochemistry & Molecular Biology, Saint Louis University School of Medicine
[2] Departments of Chemistry, Medicine, and Center for Mass Spectrometry and Metabolic Tracing, Washington University in St. Louis
[3] Department of Biology, Saint Louis University

*For correspondence:
Kyle S. McCommis, PhD
1100 South Grand Blvd.
St. Louis, MO 63104
kyle.mccommis@heath.slu.edu
Phone: 314-977-9233

K.S.M. maintained full access to all study data and takes responsibility for its integrity and that of the data analyses.

Short title: Cardiac glycogen accumulation from MPC-deficiency

**Abstract**

Heart failure involves metabolic alterations including increased glycolysis despite unchanged or decreased glucose oxidation. The mitochondrial pyruvate carrier (MPC) regulates pyruvate entry into the mitochondrial matrix, and cardiac deletion of the MPC in mice causes heart failure. How MPC deletion results in heart failure is unknown. We performed targeted metabolomics and isotope tracing in wildtype (fl/fl) and cardiac-specific Mpc2-/- (CS-Mpc2-/-) hearts after in vivo injection of U-$^{13}$C-glucose. Failing CS-Mpc2-/- hearts contained normal levels of ATP and phosphocreatine, yet these hearts displayed increased enrichment from U-$^{13}$C-glucose and increased glycolytic metabolite pool sizes. $^{13}$C enrichment and pool size was also increased for the glycogen intermediate UDP-glucose, as well as increased enrichment of the glycogen pool. Glycogen levels were increased ~6-fold in the failing CS-Mpc2-/- hearts, and glycogen granules were easily detected by electron microscopy. In young, non-failing CS-Mpc2-/- hearts, increased glycolytic $^{13}$C enrichment occurred, but glycogen levels remained low and unchanged compared to fl/fl hearts. Inhibiting glycogen synthase with MZ-101 reduced cardiac glycogen levels and improved heart failure. Feeding a ketogenic diet to CS-Mpc2-/- mice reversed the heart failure and normalized the cardiac glycogen and glycolytic metabolite accumulation. Cardiac glycogen levels were also elevated in mice infused with angiotensin-II, and both the cardiac hypertrophy and glycogen levels were improved by ketogenic diet. Thus, loss of MPC in the heart causes glycogen accumulation and heart failure, while inhibition of glycogen synthesis or a ketogenic diet can reverse both the glycogen accumulation and heart failure.

**Keywords:** heart failure, mitochondria, pyruvate, ketogenic diet

**Nonstandard Abbreviations and Acronyms**
2-DG, 2-deoxyglucose; $\alpha$KG, alpha-ketoglutarate; ADP, adenosine diphosphate; AMP,
adenosine monophosphate; ATP, adenosine triphosphate; AngII, angiotensin-II; Cr,
creatine; DHAP, dihydroxyacetone phosphate; FAO, fatty acid oxidation; FBP, fructose
bisphosphate; G6P, glucose-6-phosphate; GSD, glycogen storage disease; KD,
ketogenic diet; KEGG, Kyoto encyclopedia of genes and genomes; LF, low-fat; MPC,
mitochondrial pyruvate carrier; NAD+ and NADH, oxidized and reduced nicotinamide
adenine dinucleotide; NADP+ and NADPH, oxidized and reduced nicotinamide adenine
dinucleotide phosphate; PCr, phosphocreatine; PEP, phosphoenolpyruvate; P/M,
pyruvate/malate; R5P, ribose-5-phosphate; RT-qPCR, reverse transcription quantitative
PCR, Sedo7P, sedoheptulose 7-phosphate; UDP, uridine diphosphate; UHPLC, ultra-
high performance liquid chromatography

## Introduction

Currently, 6.7 million adults in the United States suffer from heart failure, which has risen from 6.0 million in just two years[1]. Those diagnosed with heart failure have a 5-year mortality rate of over 50%, which has remained unchanged in recent years despite existing treatment options[1]. Heart failure can arise from a multitude of causes, including ischemia/infarction, cardiac hypertrophy, chronic hypertension, or valvular diseases[1,2]. A better understanding of the development and pathophysiology of heart failure is needed to generate improved treatment options.

Since the heart lacks the ability to store ample amounts of metabolites or high energy phosphates, a continuous supply of nutrients is required to fuel its constant contractile work. Mitochondrial oxidative metabolism plays a large role in the production of high energy phosphates in the adult heart. The preferred fuel source of the heart are fatty acids, with fatty acid oxidation (FAO) accounting for ~70% of the adenosine triphosphate (ATP) produced[3,4]. Yet the healthy heart is metabolically flexible and can adapt to use alternative fuel sources to meet its energy demands, including the use of glucose, lactate, ketone bodies, and amino acids[3,4]. Glycolysis accounts for ~5% of ATP produced in the healthy heart[3,4]. However, during hypertrophy and heart failure metabolic changes occur, including significantly reduced FAO, increased glycolysis, but relatively decreased glucose oxidation[3-6]. This mismatch between increased glycolysis, yet unchanged or decreased glucose oxidation has been suggested to be important for stimulating cardiac hypertrophy and failure[7-9].

The mitochondrial pyruvate carrier (MPC) is responsible for transporting pyruvate, the end product of glycolysis or oxidized lactate, into the mitochondrial

matrix[10,11]. The mammalian MPC is a heterodimeric complex composed of MPC1 and

MPC2 proteins, both of which are necessary for MPC complex formation and pyruvate

transport[10,11]. Once pyruvate has been transported into the mitochondria, it can be

converted into acetyl-CoA and used in the citric acid cycle[10,11]. Recent studies by us and

others revealed that cardiac MPC gene and protein expression is significantly reduced

in humans with heart failure[12-15] and that deletion of MPC1[4,12,14] or MPC2[12,13] in mice

leads to spontaneous heart failure with age. However, it is still not understood how

reduction or deletion of MPC expression contributes to heart failure.

In the present study, we suggest that this heart failure is associated with

accumulation of glycogen in the MPC-deficient heart. Even with dilated cardiomyopathy,

cardiac-specific MPC2-/- (CS-Mpc2-/-) hearts contain normal levels of ATP and

phosphocreatine compared to wildtype littermates, suggesting that lack of pyruvate

oxidation does not simply result in "energetic stress". Glucose uptake is increased, and

glycolytic metabolites accumulate in both young, non-failing and failing CS-Mpc2-/-

hearts. However, glycogen accumulation is only observed in failing CS-Mpc2-/- hearts

and the heart failure and glycogen accumulation can be improved by inhibiting glycogen

synthase or providing a high-fat, low-carbohydrate ketogenic diet (KD). Lastly, we also

observe increased glycogen in hearts from angiotensin-II (AngII) infused mice, a

hypertrophic stimulus that reduces *Mpc* expression[12], and both the cardiac glycogen

and hypertrophy in this model are also normalized by KD feeding.

126
127  **Results**

128
**Heart failure in CS-Mpc2-/- mice is not due to energetic stress**

We and others have previously described spontaneous heart failure development

in MPC-deficient murine hearts[4,12-14], but what causes these hearts to fail is not

understood. In attempt to uncover insights we performed bulk RNA sequencing on CS-

Mpc2-/- hearts with dilated cardiomyopathy and age-matched fl/fl littermates (Figs. 1a-

c). This analysis uncovered 956 significantly differentially expressed genes in the failing

hearts compared to control, with ribosome and protein synthesis pathways upregulated

and multiple pathways related to fatty acid metabolism, pyruvate metabolism, and

amino acid catabolism downregulated (Fig. 1c). The downregulation of fatty acid

metabolism is a well-known metabolic alteration in failing hearts[2], including in these CS-

Mpc2-/- hearts[13]. Despite transcriptional downregulation of fatty acid, pyruvate, and

amino acid metabolic pathways, surprisingly CS-Mpc2-/- hearts maintained normal

levels of the high energy phosphates ATP and phosphocreatine as well as normal

phosphocreatine/creatine and phosphocreatine/ATP ratios (Figs. 1d-j). Although AMP

levels were slightly elevated and ADP levels trended to be elevated, the lack of a

decrease in ATP or phosphocreatine suggests normal energetic status of these failing

CS-Mpc2-/- hearts. Additionally, there were no significant differences in nicotinamide

nucleotides NAD+, NADH, NADP+, or NADP in failing CS-Mpc2-/- hearts (Figs. 1k-n),

suggesting that redox control of metabolic pathways is also normal. Altogether, these

results suggest that despite multiple decreased oxidative pathways, the heart failure in

CS-Mpc2-/- mice is not caused by overt bioenergetic stress.

**Failing CS-Mpc2-/- hearts display increased glycolysis**

As it has been well described that heart failure is associated with increased

glucose utilization, often with unchanged or decreased glucose oxidation[16], we

hypothesized that MPC deletion would mimic these metabolic changes. To evaluate

glucose metabolism, we injected mice intraperitoneally (i.p.) with U-$^{13}$C-glucose, excised

the hearts 30 min post injection, and assessed metabolite pool sizes and $^{13}$C

isotopologues by UHPLC-MS/MS. We first focused on fully $^{13}$C labeled glycolytic

metabolites (Fig. 2a) and their pool sizes (Supplementary Fig. 1). While glucose

enrichment and pool size were unaltered, glucose-6-phosphate (G6P) enrichment

showed a trend to be increased and all other downstream glycolytic metabolites

displayed significantly greater $^{13}$C enrichment and pool size in CS-Mpc2-/- hearts (Figs.

2b-g and Supplementary Figs. 1a-h). Additionally, CS-Mpc2-/- hearts displayed

increased enrichment and pool sizes of both lactate and alanine (Figs. 2h-i and

Supplementary Figs. 1i-j), indicative of decreased pyruvate carbon entry into the TCA

cycle. Altogether, these results suggest increased glycolytic metabolism in failing CS-

Mpc2-/- hearts.

Surprisingly, CS-Mpc2-/- hearts do not display significantly decreased enrichment

of TCA cycle metabolites (Supplementary Figs. 2a-l). Likewise, pool sizes of TCA cycle

metabolites were also unaltered in CS-Mpc2-/- hearts apart from α-ketoglutarate being

significantly decreased (Supplementary Figs. 2a-l). This normal TCA cycle enrichment

from U-$^{13}$C-glucose contrasts studies of MPC deletion or inhibition in other cell types,

such as hepatocytes[17], when provided U-$^{13}$C-glucose or U-$^{13}$C-pyruvate. However, while

one previous study identified decreased TCA cycle enrichment with MPC1 deletion in

isolated cardiomyocytes[14], another previous study observed normal or even significantly

increased enrichment of the TCA cycle in MPC1-/- hearts perfused with $^{13}$C-glucose[4]. It

has recently been suggested that mitochondrial lactate import and metabolism could

explain pyruvate carbon entry into MPC-/- cardiac mitochondria[18]. While the puzzling

$^{13}$C enrichment results do not support it, it is expected and likely that MPC deletion

reduces cardiac pyruvate oxidation, as both isolated mitochondria and permeabilized

cardiac muscle fibers display significantly reduced oxygen consumption when provided

with pyruvate/malate as substrates (Supplementary Fig. 3)[13]. Additionally, while

mitochondria from fl/fl hearts are sensitive to the MPC inhibitor UK-5099 which

decreases pyruvate respiration, there is little-to-no effect of UK-5099 in CS-Mpc2-/-

mitochondria or muscle fibers (Supplementary Fig. 3). Potential explanations for these

surprising TCA cycle $^{13}$C enrichment results are discussed in more detail below.

**CS-Mpc2-/- hearts do not increase glucose enrichment of the pentose phosphate**

**pathway, but do increase flux into glycerol-3-phosphate**

Several recent studies have suggested the importance of glucose flux through

the pentose phosphate pathway to fuel hypertrophied or failing hearts with amino acids,

RNA, and nucleotides[7,9]. Interestingly, in failing CS-Mpc2-/- hearts we observed no

significant increases in $^{13}$C enrichment of 6-phosphogluconate, ribose-5-phosphate,

sedoheptulose-7-phosphate, nor serine or glycine which can be end products of the

pentose phosphate pathway (Supplementary Figs. 4a-f). As described above, we also

identified no changes in NADP+ or NADPH (Figs. 1m-n), indicative of unaltered flux

through the pentose phosphate pathway. Several pentose phosphate pathway

metabolites displayed increased pool sizes in CS-Mpc2-/- hearts (Supplementary Figs.

4b-f), however the unaltered enrichment from U-$^{13}$C glucose suggests that CS-Mpc2-/-

hearts do not significantly increase glucose incorporation into the pentose phosphate

pathway. Lastly, we observed increased $^{13}$C enrichment and pool size of glycerol-3-

phosphate (Supplementary Figs. 1a,k-l), suggesting the failing CS-Mpc2-/- hearts

increasingly shuttle glucose carbons into glycerophospholipid synthesis.

**Failing CS-Mpc2-/- hearts accumulate glycogen**

205            Another potential fate of glucose is storage as glycogen, and indeed we[13] and

others[4] previously observed increased glycogen content in MPC-deficient hearts,

without expanding upon this finding. Here we confirmed again increased glycogen

concentrations in failing CS-Mpc2-/- hearts measured biochemically (Fig. 2j) as well as

by detection of abundant glycogen granules on transmission electron microscopy

images of failing CS-Mpc2-/- hearts (Fig. 2k). In hearts from U-$^{13}$C-glucose injected

mice, we observed significantly elevated $^{13}$C enrichment and pool size of UDP-glucose

(Figs. 2l-m), indicating increased flow of carbon into this glycogen intermediate. When

glucose was liberated from the cardiac glycogen, fl/fl hearts displayed essentially zero

$^{13}$C enrichment, while all but one of the CS-Mpc2-/- hearts displayed detectable

enrichment (Fig. 2n). Conversely, the percentage of completely unlabeled glucose

liberated from glycogen was significantly decreased in CS-Mpc2-/- hearts (Fig. 2o).

Taken together, these results suggest that loss of MPC in the heart results in increased

glycogen synthesis and accumulation of glycogen.

**Glucose uptake is increased, and glycogen synthase should be inhibited in CS-**

**Mpc2-/- hearts**

While the $^{13}$C-glucose enrichment studies suggest increased glycolysis in CS-Mpc2-/- hearts, the bolus of glucose injected makes it difficult to determine if cardiac glucose uptake is altered, and we observed similar labeling of glucose in fl/fl and CS-Mpc2-/- hearts in those studies (Fig. 2b). To more appropriately assess glucose uptake, left ventricle myocardial muscle fibers were incubated with $^{3}$H-2-deoxyglucose in the absence or presence of insulin. Basal glucose uptake in CS-Mpc2-/- myocardial fibers was nearly double that of fl/fl myocardial fibers and remained significantly higher with mild insulin-stimulation (Fig. 3a). Since heart failure is often associated with cardiac insulin-resistance[19], we assessed insulin signaling by blotting for phosphorylated and total AKT after i.p. insulin injection, which was not significantly altered in CS-Mpc2-/- mice (Fig. 3b). Lastly, we also investigated the regulatory enzymes for glycogen synthesis and degradation. Interestingly, in the failing CS-Mpc2-/- hearts (Figs. 3c-e), we observed significantly elevated phosphorylation of glycogen synthase at serine 640 (Figs. 3c,f), which is a strong inhibitory phosphorylation site[20]. Additionally, expression of glycogenin-1, which acts as a primer for glycogen synthase by initializing the glucosyl chain[21], was decreased in failing CS-Mpc2-/- hearts (Figs. 3c,g). We did not observe any difference in glycogen phosphorylase expression, therefore cannot suggest whether glycogenolysis, or the breakdown of glycogen, is altered in CS-Mpc2-/- hearts. Altogether, these results suggest that cardiac MPC loss increases glucose uptake and glycogen synthesis, despite the heart's attempt to inhibit glycogen synthesis by phosphorylation of glycogen synthase and decreased glycogenin-1 expression.

**Non-failing CS-Mpc2-/- hearts from young mice display increased glycolysis, but**

**no glycogen accumulation**

As increased glucose uptake and glycolysis are known to occur in failing hearts,

we investigated whether these metabolic changes were due to the presence of heart

failure in CS-Mpc2-/- hearts, or a direct consequence of MPC deletion. To assess this,

we repeated the U-[13]C-glucose enrichment experiment in 6-week-old fl/fl and CS-Mpc2-

/- mice before the onset of cardiac remodeling and dysfunction[13]. The [13]C enrichment of

all glycolytic metabolites was increased in these young, non-failing CS-Mpc2-/- hearts

(Figs. 4a-i). The pool size of glucose in these hearts was increased, but the pool size of

all glycolytic metabolites was unaltered in the non-failing CS-Mpc2-/- hearts

(Supplementary Figs. 5a-j). However, neither UDP-glucose enrichment nor pool size

were significantly altered in the non-failing CS-Mpc2-/- hearts (Figs. 4j-k), suggesting no

increase in glycogen synthesis. Indeed, glycogen concentrations were normal in these

young, non-failing CS-Mpc2-/- hearts (Fig. 4l) and unlike in the failing hearts, we did not

readily detect glycogen granules on electron micrographs (Fig. 4m). The increased

glycolysis in these non-failing CS-Mpc2-/- hearts did not appear to increase pentose

phosphate pathway flux as ribose-5-phosphate, and the enrichment of serine and

glycine and their pool sizes remained normal (Supplementary Figs. 5k-o). However,

enrichment into glycerol-3-phosphate was increased, while pool size remained

unchanged (Supplementary Figs. 5p,q), suggesting increased flow of glucose carbons

into glycerophospholipid synthesis. As within the failing CS-Mpc2-/- hearts, these

nonfailing CS-Mpc2-/- hearts also displayed normal enrichment and pool size of TCA

cycle intermediates after $^{13}$C-glucose injection (Supplementary Fig. 6). Cumulatively,

these data suggest that cardiac MPC deletion, on its own, increases glycolytic

metabolism, but only with the development of heart failure does glycogen accumulate.

**Inhibition of glycogen synthase improves heart failure in CS-Mpc2-/- mice**

To evaluate if glycogen accumulation had a causal relationship with heart failure

development, we next attempted to decrease cardiac glycogen by treating mice with a

recently developed glycogen synthase 1 (GYS1) inhibitor, MZ-101[22]. This potent small

molecule inhibitor was recently shown to decreased glycogen accumulation in both

skeletal muscle[22,23] and the heart[23] in mouse models of Pompe disease, a glycogen

storage disease also associated with cardiomyopathy. MZ-101 treatment had no

significant impact on blood glucose or lactate measurements (Figs. 5a,b). While vehicle-

treated CS-Mpc2-/- mice displayed cardiac glycogen accumulation, glycogen levels

were completely normalized in CS-Mpc2-/- mice treated with MZ-101 (Fig. 5c).

Somewhat surprisingly, cardiac glycogen levels were unaffected in fl/fl mice treated with

MZ-101, which was also observed previously in normal WT mice[23], likely due to the low

levels of glycogen in normal hearts. In slow twitch soleus skeletal muscle, glycogen

content in both fl/fl and CS-Mpc2-/- mice was significantly decreased by MZ-101 (Fig.

5d), whereas there was no significant impact on liver glycogen content (Fig. 5e), since

MZ-101 inhibits only GYS1 and not hepatic GYS2[22]. Dilated cardiomyopathy was

significantly reduced as indicated by reduced heart weight and histology (Figs. 5f,g).

Gene expression for natriuretic peptides, markers of cardiac failure, were elevated in

vehicle-treated CS-Mpc2-/- hearts, and reduced by MZ-101 (Figs. 5h,i). Lastly,

phosphorylation of glycogen synthase serine-640 was increased in vehicle-treated CS-

Mpc2-/- failing hearts and was significantly decreased by MZ-101 (Fig. 5j). All together

these data indicate that inhibition of glycogen synthase reduces glycogen accumulation

leading to the improvement of heart failure in CS-Mpc2-/- mice.

**Ketogenic diet normalizes the CS-Mpc2-/- heart failure and glycogen**

**accumulation**

Previous reports by us and others showed that feeding a ketogenic diet could

either prevent or reverse the heart failure in MPC-deficient hearts[4,13]. Here we again

normalized the CS-Mpc2-/- hearts within 3 weeks of ketogenic diet feeding (Fig. 6a).

This was associated with complete normalization of the glycogen content of the hearts

measured both biochemically and by electron microscopy (Figs. 6b,c). Phosphorylation

of glycogen synthase serine-640, which was dramatically increased in the failing CS-

Mpc2-/- hearts from low fat (LF)-fed mice, was also normalized by KD-feeding (Fig. 6d).

To evaluate glucose metabolism, we once again injected mice with U-$^{13}$C-glucose and

measured cardiac metabolite enrichment and pool sizes by mass spectrometry.

Interestingly, glucose enrichment was significantly increased in both fl/fl and CS-Mpc2-/-

mice fed a KD compared to mice fed LF diet (Fig. 6e), suggesting that these hearts that

have been chronically subjected to low blood glucose and insulin from KD feeding[13]

have the ability to enhance glucose uptake when provided with a bolus. This led to

increased glycolytic enrichment down to elevated enrichment of pyruvate, lactate, and

alanine in hearts from KD-fed mice (Supplementary Figs. 7a-c). However, $^{13}$C

enrichment of TCA cycle intermediates were very low in hearts of KD-fed mice,

suggesting that ketogenic diet strongly downregulates glucose carbon incorporation into

the TCA cycle and oxidation (Supplementary Figs 7d-l). The failing CS-Mpc2-/- hearts

from LF-fed mice once again displayed increased enrichment of the glucose within

glycogen, however the increased U-$^{13}$C-glucose uptake with KD-feeding also generally

increased $^{13}$C-enrichment of glycogen (Fig. 6f). However, the metabolite pool sizes of

glucose-6-phosphate, pyruvate, and the glycogen precursor UDP-glucose, paint the

clear picture of accumulation in the failing CS-Mpc2-/- heart, and normalization with KD-

feeding (Figs. 6g-i). Lastly, in confirmation of our biochemical assay and electron

micrographs, glucose within the glycogen pool measured by mass spectrometry was

significantly elevated in CS-Mpc2-/- hearts from LF-fed mice and normalized by KD-

feeding (Fig. 6j). All together, these data suggest that KD rescues the heart failure in

CS-Mpc2-/- mice likely by decreasing the need for pyruvate entry into the TCA cycle and

normalizing glycolytic metabolism and glycogen storage.

**AngII-induced cardiac hypertrophy increases glycogen and is improved by**

**ketogenic diet**

To evaluate if increased cardiac glycogen was characteristic of non-genetic

cardiomyopathy, we subjected mice to AngII infusion via subcutaneous osmotic pump in

combination with LF or KD-feeding. As expected, KD-feeding significantly lowered blood

glucose concentrations and increased plasma β-hydroxybutyrate levels in both saline

and AngII infused mice (Figs. 7a-b). AngII infused mice fed LF displayed significant

hypertrophy as shown by increase in HW/TL, which was significantly reduced by KD-

feeding (Fig. 7c). Cardiac glycogen more than doubled in AngII infused mice fed LF diet,

and again cardiac glycogen levels normalized by KD (Fig. 7d). AngII was previously

shown to reduce cardiac MPC1 and MPC2 expression[12], and we confirmed that both

*Mpc1* and *Mpc2* expression were significantly decreased by AngII in LF-fed mice (Figs.

7e-f). Interestingly, ketogenic diet significantly reduced cardiac *Mpc* expression

compared to LF (Figs. 7e-f), potentially contributing to the significant decrease of $^{13}$C-

glucose enrichment of the TCA cycle we observed with KD (Supplementary Fig. 7). Like

in the failing CS-Mpc2-/- hearts, there was a significant increase in phosphorylation of

glycogen synthase in the hearts of AngII infused mice fed LF diet (Figs. 7g-h),

suggesting a cellular attempt to inhibit glycogen synthesis. This phosphorylation of

glycogen synthase was also slightly increased in hearts from KD-fed mice (Figs. 7g-h),

likely due to the low dietary carbohydrate and decreased glycemia (Fig. 7a). All

together, these data suggest that AngII-induced cardiac hypertrophy decreases *Mpc*

expression and increases glycogen stores; however, this can be normalized through

KD-feeding which also improves the cardiac hypertrophy.

## Discussion

This work describes that decreases in MPC expression, or genetic deletion of the

MPC, increases cardiac glucose utilization, glycogen accumulation, and heart failure.

Altered cardiac metabolism is a hallmark of heart failure, generally consisting of

decreased mitochondrial oxidative capacity, particularly for fatty acids which the normal

heart prefers[2]. It is generally believed that this decrease in mitochondrial oxidation

creates an "energy starved" heart that then causes an increase in myocardial glucose

uptake and glycolysis to attempt to fuel the heart[16]. Our current study suggests this may

not represent the full story beyond these metabolic changes. In hearts deficient for the

mitochondrial pyruvate carrier, which limits the ability of pyruvate from glucose or lactate

to enter the mitochondria for oxidation, profound dilated cardiomyopathy develops

despite normal levels of high energy phosphates ATP and phosphocreatine. ATP was

also previously found to be normal in MPC1-/- hearts that had begun to fail[12]. Although

this indicates a lack of "energetic stress" or the "energy starved failing heart", glucose

uptake and glycolysis is increased in these CS-Mpc2-/- hearts. Previous studies

suggest that increased glucose utilization is important for fueling the hypertrophied or

failing heart, predominantly by increasing nucleotide and amino acid anabolism through

the pentose phosphate pathway[7,9]. Here, we suggest that accumulation of glycogen is

contributing to the heart failure in CS-Mpc2-/- hearts and in AngII-induced hypertrophy.

The heart has very little capacity to store nutrients or high energy phosphates[24],

making constant tissue perfusion critical. This includes very little storage of glucose as

glycogen, with the mouse heart typically containing <5 μmol glucosyl units per gram,

compared to 5-30 μmol/g in skeletal muscles and 50-200 μmol/g in the liver in the fed

condition[25]. Of course, skeletal muscle glycogen is important during exhaustive

exercise, and liver glycogen stores are critical for maintaining glycemia during fasting,

and these physiologic stressors can deplete the skeletal muscle and liver glycogen

pools. Interestingly, heart glycogen levels remain normal or are even elevated in the

fasted state[13,26-28], with this elevation likely due to the hearts switching to even more

predominant fat oxidation in the fasted state inhibiting glycolysis more than glucose

uptake[29]. High cardiac workloads increase glycogenolysis and may decrease cardiac

glycogen concentrations[30,31], however these perfused heart studies showed that

perfusion with ample substrates (glucose or lactate) abrogated the depletion of

glycogen. Thus, it is not well understood if cardiac glycogen levels are decreased or

maintained under higher workloads *in vivo*. Exhaustive exercise has been shown to

decrease cardiac glycogen content by 25%, which was rapidly repleted[32]. Overall, these

previous studies suggest that while cardiac glycogen levels are low, the consistent flux

of glucose in and out of the glycogen pool and relative maintenance of glycogen

concentrations indicates an important role.

Another interesting point from these previous studies is that glucose released

from cardiac glycogen is preferentially oxidized instead of converted to lactate[30,33]. This

would have significant effects in MPC-deficient hearts that cannot efficiently oxidize this

glycogen-derived pyruvate. This inability to oxidize glucose from glycogen may partially

explain the accumulation of glycogen in these MPC-deficient hearts, in addition to the

increased glucose uptake and incorporation of $^{13}$C-glucose into glycogen we observed.

That said, a limitation of our current study was the inability to assess rates of

glycogenolysis.

A puzzling finding from this study, and a previous study[4], is that enrichment of the

TCA cycle from $^{13}$C-glucose is not decreased in MPC-/- hearts (Supplementary Figs. 2

and 6), even though we observe significantly reduced respiration with pyruvate as

substrate in isolated mitochondria and permeabilized cardiac muscle fibers from CS-

Mpc2-/- hearts (Supplementary Fig. 3). There are several potential explanations for this

maintained TCA cycle enrichment from $^{13}$C-glucose. In this whole-heart method, $^{13}$C

enrichment of the TCA cycle would be maintained in the non-cardiomyocyte cells in the

heart which would not have *Mpc2* deleted by Mlc2vCre. This explanation is supported in

that decreased TCA cycle enrichment was observed in isolated cardiomyocytes from

MPC1-/- hearts[14]. Another possibility is that there could be a metabolic bypass of the

MPC, such as alanine transamination and alanine mitochondrial transport as we have

previously identified in hepatocytes[17], or mitochondrial import of lactate as recently

suggested[18]. This sort of bypass may be occurring to a degree, as we do observe much

greater reduction of pyruvate respiration in isolated mitochondria (devoid of cytosolic

enzymes) than we do when using permeabilized cardiac tissue (Supplementary Fig. 3).

Lastly, while we mostly think of these organic acids as TCA cycle metabolites, it is clear

many of these can be released from mitochondria and found in the cytosol and other

organelles[34]. Therefore, non-mitochondrial pools of these metabolites may somehow be

labeled by $^{13}$C-glucose in the CS-Mpc2-/- hearts. Overall, with the dramatic reduction in

mitochondrial pyruvate respiration and the pronounced pathologic phenotype of the CS-

Mpc2-/- hearts, we infer that pyruvate oxidation is reduced despite normal $^{13}$C

enrichment of TCA cycle metabolites in these studies.

Perhaps due to its low abundance, cardiac glycogen has not been well studied in

the context of heart failure. The one exception is the association between glycogen

accumulation and hypertrophic cardiomyopathy that occurs in multiple forms of

glycogen storage diseases (GSD). For example, GSD type-II, or Pompe disease, is

caused by mutation in the enzyme acid $\alpha$-glucosdiase, which catalyzes the lysosomal

degradation of glycogen[35]. This results in cardiac glycogen accumulation, up to 200-fold

in Pompe disease mouse models, and development of hypertrophic cardiomyopathy[36].

Currently, Pompe disease treatment involves enzyme replacement therapy, requiring

regular invasive and time-consuming infusions. Recent studies evaluated MZ-101, a

glycogen synthase 1 inhibitor, in mouse models of Pompe disease which significantly

reduced glycogen load in skeletal and cardiac muscle[22,23]. Other than reduction in

glycogen, cardiac endpoints were not evaluated in either study. These two studies

suggest the potential efficacy of "substrate reduction therapies" for Pompe disease[22,23].

Interestingly, both these mouse models of Pompe disease[22] and the CS-Mpc2-/- hearts

in which glycogen accumulates display increased phosphorylation of glycogen synthase

which should be inhibitory, yet glycogen synthase activity remains or is enhanced in

these pathologies. Thankfully, MZ-101 inhibits both the unphosphorylated and

phosphorylated forms of GYS1[22]. In summary, glycogen storage diseases clearly

support the connection between increased cardiac glycogen and heart failure, and this

current work suggests that cardiac glycogen reduction with MZ-101 successfully

improves heart failure caused by defective mitochondrial pyruvate metabolism.

This current work also describes the normalization of cardiac glycogen and

rescuing of heart failure by a KD in both the CS-Mpc2-/- model, and in AngII-induced

hypertrophy. We hypothesize that KD is improving these hearts by decreasing glycemia

and cardiac glucose metabolism, ultimately reducing the need for mitochondrial

pyruvate metabolism and the MPC as shown by very low $^{13}$C-glucose enrichment of the

TCA cycle in hearts from KD-fed mice (Supplementary Fig. 7). Significantly reduced

glucose oxidation, even with insulin stimulation, was recently observed in hearts from

mice fed a KD[37]. One interesting point to this current study however is that hearts from

KD-fed mice appeared to have increased uptake of the bolus $^{13}$C-glucose (Fig. 6e). This

is somewhat contrary to the KD-induced insulin resistance recently described[37], and

argues that the hearts of KD-fed mice can readily utilize glucose when challenged with a

bolus. Ultimately, this chronic decrease in glycemia and cardiac glucose metabolism

from ketogenic diet feeding allows the glycogen levels to return to normal. Thus, our

findings not only provide new insights into how reduced mitochondrial pyruvate

metabolism contributes to heart failure but also provides important implications for the

development of strategies to reduce cardiac glucose utilization and glycogen levels.

One could speculate this decrease in glycogen may also be a mechanism for the

profound cardioprotective effects of sodium-glucose cotransporter-2 (SGLT2) inhibitors,

as recently postulated that SGLT2 inhibitors improve heart failure by reducing

intermediates of glucose and lipid metabolism[38]. However, cardiac glycogen content

with SGLT2 inhibitor treatment requires further investigation.

**Methods**

462
**Animals**

All animal procedures were performed in accordance with National Institutes of

Health guidelines and approved by the Institutional Animal Care and Use Committee at

Saint Louis University. Generation of mice with *Mpc2* gene flanked by loxP sites, as well

as cardiac-specific deletion of *Mpc2* by crossing to Mlc2v-Cre knock-in mice (Jackson

Laboratory Strain #029465) have been described previously[13,17]. All mice used in these

experiments were from a C57BL6/J background. All experiments were performed with a

mixture of male and female littermate mice. Mice were housed in a climate-controlled

specific-pathogen free facility maintained at 22-24°C and 40-60% humidity in ventilated

cages with a 12-hour light/dark cycle. Ad libitum access to drinking water was provided

by automatic cage waterers. Food was also provided ad libitum. All mice were group-

housed, up to five mice per cage. Based on our previous study[13], the experiments in

young, non-failing CS-Mpc2-/- mice were performed at 6 weeks of age and experiments

in CS-Mpc2-/- mice with heart failure were performed at 14-30 weeks of age. For

ketogenic diet (KD) experiments, fl/fl and CS-Mpc2-/- mice were initially fed chow and

aged to 16 weeks to develop heart failure in CS-Mpc2-/- mice, then switched to either

low-fat (LF) control diet or KD for a duration of 3 weeks. The LF diet (F1515, BioServ)

composition is 68.9%kcal carbohydrate, 19.0%kcal protein, and 12.1%kcal fat.

Ketogenic diet (F3666, BioServ) composition is 1.8%kcal carbohydrate, 4.7%kcal

protein, and 93.5%kcal fat. Diets were randomly assigned for whole cages based on

genotypes contained within to result in roughly even numbers for each group. Unless

otherwise stated, at study culmination mice were fasted for 2-3 hours, and euthanized

by $CO_2$ asphyxiation. Blood was collected via cannulation of the inferior vena cava and

placed into EDTA-treated tubes on ice. Tissues were then excised, weighed, and snap

frozen in liquid nitrogen. Plasma was collected by centrifuging blood tubes at 8,000$g$ for

8 minutes at 4°C and freezing the plasma supernatant in liquid nitrogen. Plasma and

tissues were stored at -80°C until analyzed.

MZ-101 was purchased from Enamine US, Inc, and was dissolved in 45% H2O,

30% polyethylene glycol 200, 20% polypropylene glycol, and 5% N-methyl-2-pyrrolidone

(all from Sigma Aldrich) as performed previously [22]. Gavage dosing (100 mg/kg MZ-101

or similar volume of vehicle) was performed twice daily in 14-week-old CS-Mpc2-/- or

fl/fl littermate mice. Fresh gavage solutions were prepared weekly. Mice were treated for

2 weeks, then euthanized by $CO_2$ and plasma/tissues harvested as described above.

AngII infusion was performed in 9-week-old wildtype C57BL6/J male mice

purchased from the Jackson Laboratory (Strain #000664). Subcutaneous osmotic

pumps (#1004, Alzet) with either AngII (1.44mg/kg/day, Sigma Aldrich) or saline vehicle

were implanted for an infusion duration of three weeks similarly to as described

previously[39]. Mice were anesthetized by inhaled 2-3% isoflurane in 100% oxygen via a

nose cone, and the upper back area shaved with electric clippers and scrubbed with

7.5% betadine antiseptic. A ~1cm incision was made in the prepped area of the mouse's

back, and forceps inserted to create a subcutaneous pocket. After inserting the pump,

the wound was sealed with 2-3 surgical staples. Mice were treated with 1 mg/kg

subcutaneous buprenorphine-SR and several drops of lidocaine were applied to wound

edges for pain management. Mice were allowed to arouse while on a heat pad and

placed back into their home cage. Mice were observed daily, and surgical staples were

removed within 7-10 days. Mice were switched to LF or KD 3 days prior to pump

implant. Diet was assigned randomly for each cage, and then vehicle versus AngII

pumps were assigned for mice for each diet to result in roughly even numbers per

group. Mice were euthanized by $CO_2$ and plasma/tissues harvested as described

above.

**RNA sequencing and analysis**

Hearts used for RNA sequencing were excised from 6 fl/fl and 6 CS-Mpc2-/- 30-

week-old chow-fed mice. RNA was isolated from ~25 mg of frozen left ventricular tissue

using RNA-STAT (Tel-Test, Inc) and purified using the PureLink RNA Mini Kit with

DNase treatment (Invitrogen). RNA was then pooled between 2 different fl/fl and CS-

Mpc2-/- hearts, resulting in n=3 samples per group. RNA concentration and quality were

determined by 2100 Bioanalyzer RNA 6000 Nano Kit (Agilent) and depleted of rRNA

using RiboMinus Eukaryote Kit v2 (Ambion). DNA libraries were constructed using Ion

Total RNA-seq Kit v2 (Thermo Fisher Scientific) and sequenced on an Ion Torrent

Proton DNA sequencer (Thermo Fisher Scientific). Raw data were aligned to the mm10

genome. Gene counts for each sample were normalized by read depth, low expression

genes filtered from the data, and $t$-test performed for statistical comparison. Lists of

upregulated and downregulated differentially expressed genes (p < 0.05) between fl/fl

and CS-Mpc2-/- hearts were subjected to pathway-level overrepresentation analysis

using IMPaLA[40], which included Kyoto encyclopedia of genes and genomes (KEGG)

pathway analysis, among others. Heat map of differentially expressed genes was

generated with Shiny heatmap[41]. RNA sequencing files were uploaded to NCBI GEO

Series Number GSE268694.

**Gene expression analysis**

Relative gene expression was determined by reverse transcription quantitative

PCR (RT-qPCR). Total RNA was extracted from snap frozen tissues using RNA-STAT

(Tel-Test, Inc). [RNA] was assessed by Nanodrop 2000c (Thermo Fisher Scientific).

Then 1 $\mu$g of sample was reverse transcribed into complementary DNA by Superscript

VILO (Thermo Fisher Scientific) using a 2720 Thermal cycler (Applied Biosystems).

Relative quantification of target gene expression was measured in duplicate using

Power SYBR Green (Thermo Fisher Scientific) with a QuantStudio 3 PCR System

(Thermo Fisher Scientific). Target gene threshold cycle (Ct) values were normalized to

reference gene (*Rplp0*) Ct values by the $2^{-\Delta\Delta Ct}$ method. Oligonucleotide primer

sequences used in 5' to 3' were: *Rplp0* FWD: GCAGACAACGTGGGCTCCAAGCGAA,

*Rplp0* REV: GGTCCTCCTTGGTGAACACGAAGCCC, *Mpc1* FWD:

GACTTTCGCCCTCTGTTGCTA, *Mpc1* REV: GAGGTTGTACCTTGTAGGCAAAT, *Mpc2*

FWD: CCGCCGCGATGGCAGCTG, *Mpc2* REV: GCTAGTCCAGCACACACCAATCC,

*Nppa* FWD: AGTGCGGTGTCCAACACAGA, *Nppa* REV:

GACCTCATCTTCTACCGGCATCT, *Nppb* FWD: GCTGCTTTGGGCACAAGATAG, and

*Nppb* REV: GCAGCCAGGAGGTCTTCCTA.

**Western blotting and protein expression analysis**

Protein extracts were prepared by homogenizing ~30 mg of frozen ventricular

tissue in lysis buffer (15mM NaCl, 25 nM Tri Base, 1 mM EDTA, 0.2% NP-40, 10%

glycerol) supplemented with 1X cOmplete protease inhibitor cocktail (Millipore Sigma)

and a phosphatase inhibitor cocktail (1 mM $Na_3VO_4$, 1 mM NaF, and 1 mM PMSF) by

using 2-3 zirconia beads (Biospec products) and homogenization for 1 min with a Mini

Beadbeater (Biospec products). Protein concentrations were measured using MicroBCA

Assay kit (Thermo Fisher Scientific) and absorbance measured with a Synergy HTX

multi-mode plate reader (Biotek). Then 50 μg of protein lysate was electrophoresed on

criterion TGX 4-15% precast gel (Bio-Rad Laboratories) and transferred onto PVDF

membrane (Millipore). Membranes were blocked with 5% bovine serum albumin (BSA)

(#BAH65 Equitech-Bio, Inc) in Tris-buffered saline with Tween-20 (TBST) for at least 1

hour.

Primary antibodies were used at 1:1000 dilution in 5% BSA-TBST, with gentle

rocking at 4°C overnight. AKT (9272), phosphorylated AKT (serine 473) (4060), MPC1

(14462), MPC2 (46141), glycogen synthase (3886), and phosphorylated glycogen

synthase (serine 640) (47043) were from Cell Signaling. $\alpha$-tubulin (T6199) was from

Sigma. PYGM (ab231963) was from abcam. Glycogenin-1 (sc-271109) was from Santa

Cruz Biotechnology. After primary antibody incubation, membranes were washed 3X for

10 minutes in TBST and probed with IRDye secondary antibodies at 1:10,000 (Li-Cor

Biosciences, 926-32213 or 926-68072) in 5% BSA-TBST for 1 hour, then washed 3X 10

min. Membranes were imaged on an Odyssey CLx imaging system and analyzed with

Image Studio Lite software (Li-Cor Biosciences).

**575    Targeted metabolomics and $^{13}$C enrichment from glucose**

Mice were fasted for 2 hours then injected intraperitoneally (i.p.) with 1g/kg

uniformly labelled U-$^{13}$C-glucose (CLM-1396, Cambridge Isotope Labs) in saline. 30 min

later the mice were anesthetized with 1-2% isoflurane (Covetrus) in 100% oxygen via an

induction chamber then transferred to a nose cone. A thoracotomy was rapidly

performed and the beating heart snap-frozen using a metal clamp chilled in liquid

nitrogen. Hearts were stored at -80°C until processed. The heart tissue was pulverized

with a mortar and pestle chilled in a liquid nitrogen bath. To every 1 mg of pulverized

tissue, 40 µl of 2:2:1 acetonitrile:methanol:water was added. The samples were then

briefly vortexed, placed in a liquid nitrogen bath for 1 minute, and then sonicated for 10

minutes at 25°C. This was repeated 3X. Samples were stored at -20°C overnight. The

following day, the samples were centrifuged at 14,000*g* for 10 min at 4°C, and

supernatants added to glass liquid chromatography vials with autosampler inserts and

stored at -80°C until analyzed.

Ultra-high-performance liquid chromatography (UHPLC)-MS/MS was performed

with a Vanquish Flex UHPLC system interfaced with an Orbitrap ID-X Tribid Mass

spectrometer (Thermo Fisher Scientific) similar to as previously described[42]. Hydrophilic

interaction liquid chromatography (HILIC) separation was performed using a HILICON

iHILIC-(P) Classic column (Tvistevagen, Umea, Sweden) with the following

specifications: 100 × 2.1 mm, 5 µm. Mobile-phase solvents were composed of

A = 20 mM ammonium bicarbonate, 0.1% ammonium hydroxide and 2.5 µM medronic

acid in water:acetonitrile (95:5) and B = acetonitrile:water (95:5). The column

compartment was 45°C for all experiments. The following linear gradient was applied at

a flow rate of 250 µL min $^{-1}$: 0–1 min: 90% B, 1–12 min: 90-35% B, 12–12.5 min: 35-

25% B, 12.5–14.5 min: 25% B. The column was re-equilibrated with 20 column volumes

of 90% B. The injection volume was 4 µL for all experiments.

Data were collected with the following settings: spray voltage, 3.5kC (positive) /

−2.8 kV (negative); sheath gas, 50; auxiliary gas, 10; sweep gas, 1; ion transfer tube

temperature, 300°C; vaporizer temperature, 200°C; mass range, 67–1,000 Da,

resolution, 120,000 (MS1), 30,000 (MS/MS); maximum injection time, 100 ms; isolation

window, 1.6 Da. LC/MS data were processed and analyzed with the open-source

Skyline software[43]. Natural abundance correction of $^{13}$C for tracer experiments was

performed with AccuCor[44]. All studies contained 3 quality control samples and a blank

sample, and only metabolites with sufficient abundance and fractional enrichment over

the limit of detection were included in analysis. These methods were used to analyze

the pool size and $^{13}$C enrichment of glucose and glycolytic metabolites, organic acids

and amino acids, and the pool sizes of nucleotide phosphates and phosphocreatine and

creatine. Data are presented as the % enrichment of the specified $^{13}$C isotopologue, or

as the normalized peak area for each metabolite's pool size.

**Transmission electron microscopy and gross histology**

The papillary muscle from hearts (under 1 mm$^3$) was excised and fixed in 2.5%

glutaraldehyde in 0.1M sodium cacodylate, pH 7.4 overnight. Samples were then post-

fixed with 1% osmium tetroxide for 1 h and then serial dehydration with alcohol (50%,

75%, 90%, and 100%) was performed and then embedded in Epon 12. Ultrathin

sections were then post-stained with uranyl acetate and lead citrate. Samples were

imaged on a JEOL 1400 Plus transmission electron microscope equipped with an AMT

camera.

Hearts from vehicle and MZ-101 treated mice were cut into a ~1 mm short axis

slice with a razor blade and fixed in 10% neutral buffered formalin. Slices were then

processed, embedded, sectioned, and stained by H&E by the Saint Louis University

Advanced Spatial Biology and Research Histology Core.

**Cardiac glycogen assay**

Cardiac glycogen concentrations were measured similarly to as preformed

previously[13]. Briefly, 15-30 mg of heart tissue were boiled in 30% KOH at 100°C for 30

min. Then, 2% $Na_2SO_4$ and 100% EtOH were added, and the tubes were vortexed.

Samples were boiled again to help aid in the precipitation of glycogen and then tubes

were centrifuged at 16,000$g$ for 5 min and supernatant aspirated. The pellet was

dissolved in 200 μL $H_2O$ and 800 μL of 100% ethanol was added. The samples were

then boiled for 5 minutes, and recentrifuged at 16,000$g$ for 5 min. This wash step was

repeated 3X. The final pellets were suspended in 200 μL of 0.3 mg/ml amyloglucosidase

(Roche) in 0.2 M sodium acetate (pH 4.8). Serial dilutions of 10 mg/ml oyster glycogen

(Sigma) were prepared as standards. Samples and standards were then incubated in a

40°C water bath for 3 hours. Samples and standards were then diluted 1:1 with $H_2O$,

and the samples were added to a 96-well plate. Glucose assay buffer (0.3 M

triethanolamine, 4 mM $MgCl_2$, 2 mM ATP, 0.9 mM $NADP^+$, and 5 μg/mL glucose-6-

phosphate dehydrogenase (Roche)) was added to each well, and the absorbance was

measured at 340 nm with a Synergy HTX multi-mode plate reader (Biotek). Then 1 μg

hexokinase (Roche) was added to each well and the plate was incubated at room

temperature in the dark for 30 minutes, and absorbance measured again at 340 nm.

The glycogen concentration was calculated from the difference in absorbance readings

plotted in relation to the oyster glycogen standards.

Glycogen [13]C enrichment and pool size measurement was performed similarly as

described previously[22]. Briefly, glycogen from cardiac tissue of mice injected with U-[13]C-

glucose as described above was precipitated and digested by 0.3 mg/mL

amyloglucosidase in 0.2 M sodium acetate as described for the biochemical glycogen

assay. Following the 3 hour digestion with amyloglucosidase, the samples were added

to a 2:2:1 acetonitrile:methanol:water mixture maintaining a 1:9 ratio of sample:

extraction solvent. The samples were vortexed for 1 min then stored at -20°C for 1 hour.

Then the samples were centrifuged at 14,000$g$ for 10 min at 4°C. The supernatant was

then transferred to glass LC vials for UHPLC-MS/MS analysis as described above.

**Myocardial glucose uptake assay**

Glucose uptake was measured similarly to methods used for skeletal muscle and

papillary muscle as described previously[45-47]. Mice were anesthetized with 50 mg/kg

sodium pentobarbital intraperitoneal (i.p.) injection, the heart removed, and two ~2-4 mg

left ventricular muscle strips collected from each heart. Individual cardiac strips were

placed into a well of a 24-well plate containing Krebs-Henseleit Buffer recovery solution

with non-glucose fuel sources (4.5 mM lactate, 0.5 mM pyruvate, and 200 µM BSA-

palmitate), 34.8 mM mannitol, and 0.1% radioimmunoassay-grade BSA at 35°C.

Incubation buffers were pre-gassed with 95% $O_2$:5% $CO_2$, and each well of the 24-well

plate had its own gas supply. After 60 min in recovery buffer, samples were moved to

another well containing the same solution $\pm$ 30 µU/mL (submaximal) insulin for 30 min.

Samples were then moved to another well containing glucose transport medium $\pm$ 30

µU/mL insulin. Glucose transport assay medium was the same as recovery medium but

also contained 1 mM 2-deoxyglucose (2-DG), 2 µCi/mL $^3$H-2-DG, and 0.3 mCi/mL $^{14}$C-

mannitol at 30°C. After 10 min in the glucose transport assay medium, samples were

clamp frozen. Samples were homogenized in ice-cold buffer containing protease and

phosphatase inhibitors (50 mM HEPES, pH 7.4, 2 mM $Na_3VO_4$, 150 mM NaF,10 µg/mL

leupeptin, 10 µg/mL aprotinin, 0.5 µg/mL pepstatin and 1 mM phenylmethylsulfonyl

fluoride) and centrifuged at 4°C for 10 min at 13,500$g$. Supernatant protein was

assessed by the Bradford method, and aliquots were mixed with scintillation fluid

(Ultima Gold XR, Perkin Elmer, Boston, MA, USA) before counting of disintegrations per

minute (DPM, TriCarb 3110TR, Perkin Elmer). The DPM of $^{14}$C mannitol was used to

measure extracellular volume and correct for DPM of $^{3}$H-2-DG in the extracellular fluid.

**Insulin signaling assessment**

Insulin signaling was evaluated as described previously[48]. Briefly, mice were

fasted for 3 hours, then injected i.p. with 5mU/g insulin or saline vehicle. 10 minutes

after injection, mice were euthanized by $CO_2$ and the hearts collected and snap frozen

in liquid nitrogen and stored at -80°C. Western blotting was then performed for

phosphorylated and total AKT as described above.

**Plasma ketone body assay**

Plasma ketone concentrations were measured using a β-hydroxybutyrate

colorimetric assay kit (Cayman Chemical). Samples were incubated in the dark for 25

min and absorbance measured at 445 nm with a Synergy HTX multi-mode plate reader

(Biotek) and plotted against a standard curve for quantification.

**Mitochondrial isolation and respirometry**

Mitochondria were isolated from freshly excised hearts by ten passes of a glass-

on-glass Dounce homogenizer on ice in 4 mL of buffer containing 250 mM sucrose, 10

mM Tris, and 1 mM EDTA, pH 7.4. Homogenates were centrifuged at 1,000$g$ x 5 min at

4°C, and the supernatant transferred to a new tube and centrifuged at 10,000$g$ x 10 min

at 4°C. This mitochondria-enriched pellet was washed with fresh homogenization buffer

and recentrifuged at 10,000$g$ x 10 min at 4°C twice. The mitochondrial pellet was then

solubilized in ~200 $\mu$L of Mir05 respiration buffer consisting of 0.5 mM EGTA, 3 mM

$MgCl_2$, 60 mM lactobionic acid, 20 mM taurine, 10 mM $KH_2PO_4$, 20 mM HEPES, 110

mM sucrose, and 1g/L fatty-acid free BSA, pH 7.1. Mitochondrial protein concentration

was measured by MicroBCA (Thermo Fisher Scientific).

50 $\mu$g of mitochondria were added to each chamber of an Oxygraph O2k

FluoRespirometer (Oroboros Instruments), and respiration was stimulated with 5 mM

pyruvate, 2 mM malate, and then coupled with 2.5 mM ADP+$Mg^{2+}$. Lastly, 5 $\mu$M UK-

5099 was added to the chambers to inhibit the MPC and pyruvate respiration.

Respiration was also performed on saponin-permeabilized left-ventricular muscle

fibers using the Oroboros O2k similar to as previously performed[13]. Respiration was

again stimulated with 5 mM pyruvate, 2 mM malate, and then coupled with 2.5 mM

ADP+$Mg^{2+}$, and inhibited with 5 $\mu$M UK-5099. Respiration data was recorded and

analyzed with DatLab version 7.4.0.1 software, and is presented as oxygen

consumption rate normalized to mitochondrial mass or tissue mass added to the

chamber.

**Statistical analysis**

All data are presented as dot plots with mean±S.D. All data were graphed and

statistically analyzed with GraphPad Prism version 10.4.2. An unpaired, two-tailed

Student's *t*-test with Welch correction was used for comparison of two groups.

Experiments with more than two groups were analyzed using a two-way analysis of

variance (ANOVA) with Tukey post-hoc multiple-comparisons test. A $P$ value of less than

0.05 was considered statistically significant.

**Data availability**

Data supporting the findings of this study are available from the corresponding author

upon reasonable request. The RNA sequencing data generated in this study have been

deposited in NCBI GEO Series Number GSE268694.

**Acknowledgements**

R.C.W. is supported by a Doisy Scholar Fellowship from the Doisy Fund of the Edward

A. Doisy Department of Biochemistry and Molecular Biology at Saint Louis University

School of Medicine. This work was supported by NIH R00-HL136658 to K.S.M and R35

ES028365 to G.J.P.

**Disclosures**

G.J.P. has a research collaboration agreement with Thermo Fisher Scientific and is a

scientific advisor for Cambridge Isotope Laboratories. K.S.M. receives research support

from Cirius Therapeutics and serves as a consultant for BioGenerator Ventures. All

other authors declare no conflicts of interest.

**Author Information**
744
745 Authors and Affiliations
746
747 **Edward A. Doisy Department of Biochemistry and Molecular Biology, Saint Louis**
748 **University School of Medicine, St. Louis, MO, USA**

749 Rachel Weiss, Kelly Pyles, Michelle Brennan, & Kyle McCommis
750
751 **Department of Biology, Saint Louis University, St. Louis, MO, USA**
752 Jonathan Fisher
753
**Departments of Chemistry, Medicine, and the Center for Mass Spectrometry and**
**Metabolic Tracing, Washington University in St. Louis, St. Louis, MO, USA**
Kevin Cho & Gary Patti

Contributions
R.C.W, J.S.F, G.J.P., and K.S.M. conceived and designed the experiments. R.C.W.,
K.D.P., K.C., M.B., and J.S.F. performed the experiments and analyzed the data. R.C.W
and K.S.M. wrote the manuscript. All authors read, edited, and approved the
manuscript.
763
764
765

**References**

1    Tsao, C. W. *et al.* Heart Disease and Stroke Statistics-2023 Update: A Report From the
American Heart Association. *Circulation* **147**, e93-e621,
doi:10.1161/CIR.0000000000001123 (2023).
2    Lopaschuk, G. D., Karwi, Q. G., Tian, R., Wende, A. R. & Abel, E. D. Cardiac Energy
Metabolism in Heart Failure. *Circ Res* **128**, 1487-1513,
doi:10.1161/CIRCRESAHA.121.318241 (2021).
3    De Jong, K. A. & Lopaschuk, G. D. Complex Energy Metabolic Changes in Heart Failure
With Preserved Ejection Fraction and Heart Failure With Reduced Ejection Fraction. *Can*
*J Cardiol* **33**, 860-871, doi:10.1016/j.cjca.2017.03.009 (2017).
4    Zhang, Y. *et al.* Mitochondrial pyruvate carriers are required for myocardial stress
adaptation. *Nat Metab* **2**, 1248-1264, doi:10.1038/s42255-020-00288-1 (2020).
5    Brahma, M. K. *et al.* Increased Glucose Availability Attenuates Myocardial Ketone Body
Utilization. *J Am Heart Assoc* **9**, e013039, doi:10.1161/JAHA.119.013039 (2020).
6    Zhabyeyev, P. *et al.* Pressure-overload-induced heart failure induces a selective reduction
in glucose oxidation at physiological afterload. *Cardiovasc Res* **97**, 676-685,
doi:10.1093/cvr/cvs424 (2013).
7    Karlstaedt, A., Khanna, R., Thangam, M. & Taegtmeyer, H. Glucose 6-Phosphate
Accumulates via Phosphoglucose Isomerase Inhibition in Heart Muscle. *Circ Res* **126**, 60-
74, doi:10.1161/CIRCRESAHA.119.315180 (2020).
8    Davogustto, G. E. *et al.* Metabolic remodeling precedes mTORC1-mediated cardiac
hypertrophy. *J Mol Cell Cardiol* **158**, 115-127, doi:10.1016/j.yjmcc.2021.05.016 (2021).
9    Ritterhoff, J. *et al.* Metabolic Remodeling Promotes Cardiac Hypertrophy by Directing
Glucose to Aspartate Biosynthesis. *Circ Res* **126**, 182-196,
doi:10.1161/CIRCRESAHA.119.315483 (2020).

10   Bricker, D. K. *et al.* A mitochondrial pyruvate carrier required for pyruvate uptake in
792           yeast, Drosophila, and humans. *Science* **337**, 96-100, doi:10.1126/science.1218099
793           (2012).

794   11   Herzig, S. *et al.* Identification and functional expression of the mitochondrial pyruvate
795           carrier. *Science* **337**, 93-96, doi:10.1126/science.1218530 (2012).

796   12   Fernandez-Caggiano, M. *et al.* Mitochondrial pyruvate carrier abundance mediates
797           pathological cardiac hypertrophy. *Nat Metab* **2**, 1223-1231, doi:10.1038/s42255-020-
798           00276-5 (2020).

799   13   McCommis, K. S. *et al.* Nutritional modulation of heart failure in mitochondrial pyruvate
800           carrier-deficient mice. *Nat Metab* **2**, 1232-1247, doi:10.1038/s42255-020-00296-1
801           (2020).

802   14   Cluntun, A. A. *et al.* The pyruvate-lactate axis modulates cardiac hypertrophy and heart
803           failure. *Cell Metab* **33**, 629-648 e610, doi:10.1016/j.cmet.2020.12.003 (2021).

804   15   Lopez-Vazquez, P. *et al.* Reduced mitochondrial pyruvate carrier expression in hearts
805           with heart failure and reduced ejection fraction patients: ischemic vs. non-ischemic
806           origin. *Front Cardiovasc Med* **11**, 1349417, doi:10.3389/fcvm.2024.1349417 (2024).

807   16   Tran, D. H. & Wang, Z. V. Glucose Metabolism in Cardiac Hypertrophy and Heart Failure. *J*
808           *Am Heart Assoc* **8**, e012673, doi:10.1161/JAHA.119.012673 (2019).

809   17   McCommis, K. S. *et al.* Loss of Mitochondrial Pyruvate Carrier 2 in the Liver Leads to
810           Defects in Gluconeogenesis and Compensation via Pyruvate-Alanine Cycling. *Cell Metab*
811           **22**, 682-694, doi:10.1016/j.cmet.2015.07.028 (2015).

812   18   Cluntun, A. A. *et al.* Direct mitochondrial import of lactate supports resilient
813           carbohydrate oxidation. *bioRxiv*, doi:10.1101/2024.10.07.617073 (2024).

814   19   Riehle, C. & Abel, E. D. Insulin Signaling and Heart Failure. *Circ Res* **118**, 1151-1169,
815           doi:10.1161/CIRCRESAHA.116.306206 (2016).

816   20   Wang, Y. & Roach, P. J. Inactivation of rabbit muscle glycogen synthase by glycogen
817           synthase kinase-3. Dominant role of the phosphorylation of Ser-640 (site-3a). *J Biol*
818           *Chem* **268**, 23876-23880 (1993).

819   21   Pitcher, J., Smythe, C. & Cohen, P. Glycogenin is the priming glucosyltransferase required
820           for the initiation of glycogen biogenesis in rabbit skeletal muscle. *Eur J Biochem* **176**,
821           391-395, doi:10.1111/j.1432-1033.1988.tb14294.x (1988).

822   22   Ullman, J. C. *et al.* Small-molecule inhibition of glycogen synthase 1 for the treatment of
823           Pompe disease and other glycogen storage disorders. *Sci Transl Med* **16**, eadf1691,
824           doi:10.1126/scitranslmed.adf1691 (2024).

825   23   Gaspar, R. C. *et al.* Small molecule inhibition of glycogen synthase I reduces muscle
826           glycogen content and improves biomarkers in a mouse model of Pompe disease. *Am J*
827           *Physiol Endocrinol Metab* **327**, E524-E532, doi:10.1152/ajpendo.00175.2024 (2024).

828   24   Doenst, T., Nguyen, T. D. & Abel, E. D. Cardiac metabolism in heart failure: implications
829           beyond ATP production. *Circ Res* **113**, 709-724, doi:10.1161/CIRCRESAHA.113.300376
830           (2013).

831   25   Barre, L. *et al.* Genetic model for the chronic activation of skeletal muscle AMP-activated
832           protein kinase leads to glycogen accumulation. *Am J Physiol Endocrinol Metab* **292**,
833           E802-811, doi:10.1152/ajpendo.00369.2006 (2007).

26   Evans, G. The glycogen content of the rat heart. *J Physiol* **82**, 468-480,
doi:10.1113/jphysiol.1934.sp003198 (1934).
27   Reichelt, M. E., Mellor, K. M., Curl, C. L., Stapleton, D. & Delbridge, L. M. Myocardial
glycophagy - a specific glycogen handling response to metabolic stress is accentuated in
the female heart. *J Mol Cell Cardiol* **65**, 67-75, doi:10.1016/j.yjmcc.2013.09.014 (2013).
28   Schneider, C. A., Nguyen, V. T. & Taegtmeyer, H. Feeding and fasting determine
postischemic glucose utilization in isolated working rat hearts. *Am J Physiol* **260**, H542-
548, doi:10.1152/ajpheart.1991.260.2.H542 (1991).
29   Depre, C., Vanoverschelde, J. L. & Taegtmeyer, H. Glucose for the heart. *Circulation* **99**,
578-588, doi:10.1161/01.cir.99.4.578 (1999).
30   Goodwin, G. W., Taylor, C. S. & Taegtmeyer, H. Regulation of energy metabolism of the
heart during acute increase in heart work. *J Biol Chem* **273**, 29530-29539,
doi:10.1074/jbc.273.45.29530 (1998).
31   McConville, P., Lakatta, E. G. & Spencer, R. G. Greater glycogen utilization during 1- than
2-adrenergic receptor stimulation in the isolated perfused rat heart. *Am J Physiol*
*Endocrinol Metab* **293**, E1828-1835, doi:10.1152/ajpendo.00288.2007 (2007).
32   Conlee, R. K. & Tipton, C. M. Cardiac glycogen repletion after exercise: influence of
synthase and glucose 6-phosphate. *J Appl Physiol Respir Environ Exerc Physiol* **42**, 240-
244, doi:10.1152/jappl.1977.42.2.240 (1977).
33   Goodwin, G. W., Ahmad, F. & Taegtmeyer, H. Preferential oxidation of glycogen in
isolated working rat heart. *J Clin Invest* **97**, 1409-1416, doi:10.1172/JCI118561 (1996).
34   Martinez-Reyes, I. & Chandel, N. S. Mitochondrial TCA cycle metabolites control
physiology and disease. *Nat Commun* **11**, 102, doi:10.1038/s41467-019-13668-3 (2020).
35   Puzzo, F. *et al.* Rescue of Pompe disease in mice by AAV-mediated liver delivery of
secretable acid alpha-glucosidase. *Sci Transl Med* **9**, doi:10.1126/scitranslmed.aam6375
(2017).
36   Huang, J. Y. *et al.* CRISPR-Cas9 generated Pompe knock-in murine model exhibits early-
onset hypertrophic cardiomyopathy and skeletal muscle weakness. *Sci Rep* **10**, 10321,
doi:10.1038/s41598-020-65259-8 (2020).
37   Ho, K. L. *et al.* The ketogenic diet does not improve cardiac function and blunts glucose
oxidation in ischemic heart failure. *Cardiovasc Res*, doi:10.1093/cvr/cvae092 (2024).
38   Packer, M. SGLT2 inhibitors: role in protective reprogramming of cardiac nutrient
transport and metabolism. *Nat Rev Cardiol* **20**, 443-462, doi:10.1038/s41569-022-00824-
4 (2023).
39   Yang, T. *et al.* Cell-Specific Actions of the Prostaglandin E-Prostanoid Receptor 4
Attenuating Hypertension: A Dominant Role for Kidney Epithelial Cells Compared With
Macrophages. *J Am Heart Assoc* **11**, e026581, doi:10.1161/JAHA.122.026581 (2022).
40   Kamburov, A., Cavill, R., Ebbels, T. M., Herwig, R. & Keun, H. C. Integrated pathway-level
analysis of transcriptomics and metabolomics data with IMPaLA. *Bioinformatics* **27**,
2917-2918, doi:10.1093/bioinformatics/btr499 (2011).
41   Khomtchouk, B. B., Hennessy, J. R. & Wahlestedt, C. shinyheatmap: Ultra fast low
memory heatmap web interface for big data genomics. *PLoS One* **12**, e0176334,
doi:10.1371/journal.pone.0176334 (2017).

42 Ferguson, D. *et al.* Mitochondrial pyruvate carrier inhibition initiates metabolic crosstalk
to stimulate branched chain amino acid catabolism. *Mol Metab* **70**, 101694,
doi:10.1016/j.molmet.2023.101694 (2023).
43 Adams, K. J. *et al.* Skyline for Small Molecules: A Unifying Software Package for
Quantitative Metabolomics. *J Proteome Res* **19**, 1447-1458,
doi:10.1021/acs.jproteome.9b00640 (2020).
44 Su, X., Lu, W. & Rabinowitz, J. D. Metabolite Spectral Accuracy on Orbitraps. *Anal Chem*
**89**, 5940-5948, doi:10.1021/acs.analchem.7b00396 (2017).
45 Fisher, J. S. *et al.* Glucose transport rate and glycogen synthase activity both limit skeletal
muscle glycogen accumulation. *Am J Physiol Endocrinol Metab* **282**, E1214-1221,
doi:10.1152/ajpendo.00254.2001 (2002).
46 Ching, J. K. *et al.* Impaired insulin-stimulated glucose transport in ATM-deficient mouse
skeletal muscle. *Appl Physiol Nutr Metab* **38**, 589-596, doi:10.1139/apnm-2012-0175
(2013).
47 Binsch, C. *et al.* Deletion of Tbc1d4/As160 abrogates cardiac glucose uptake and
increases myocardial damage after ischemia/reperfusion. *Cardiovasc Diabetol* **22**, 17,
doi:10.1186/s12933-023-01746-2 (2023).
48 Kamm, D. R. *et al.* Novel insulin sensitizer MSDC-0602K improves insulinemia and fatty
liver disease in mice, alone and in combination with liraglutide. *J Biol Chem* **296**, 100807,
doi:10.1016/j.jbc.2021.100807 (2021).
897
898
899

**Figure Legends**

**Fig. 1: Heart Failure in CS-Mpc2-/- mice is not due to energetic stress. a** Heart weight/tibia length (HW/TL) in CS-Mpc2-/- and fl/fl littermates, n=6. **b** Representative gross images of excised fl/fl and CS-Mpc2-/- hearts. Scale bar = 5 mm. **c** Heatmap of differentially expressed genes identified by RNA sequencing of hearts from fl/fl and CS-Mpc2-/- littermates, n=3. Most prominently dysregulated KEGG pathways upregulated and downregulated are listed. **d-n** mass spectrometry normalized peak area pool sizes of adenosine triphosphate (ATP) **(d)**, adenosine diphosphate (ADP) **(e)**, adenosine monophosphate (AMP) **(f)**, phosphocreatine **(g)**, creatine **(h)**, phosphocreatine/creatine (PCr/Cr) ratio **(i)**, PCr/ATP ratio **(j)** oxidized nicotinamide adenine dinucleotide (NAD$^+$) **(k)**, reduced nicotinamide adenine dinucleotide (NADH) **(l)**, oxidized nicotinamide adenine dinucleotide phosphate (NADP$^+$) **(m)**, and reduced nicotinamide adenine dinucleotide phosphate (NADPH) **(n)** in hearts of fl/fl and CS-Mpc2-/- littermates that had been injected i.p. with U-$^{13}$C-glucose, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

**Fig. 2: Failing CS-Mpc2-/- hearts display increased glycolysis and accumulate glycogen. a** Schematic of U-$^{13}$C-glucose labeling of glycolytic metabolites and glycogen, created with BioRender. **b-i** $^{13}$C % enrichment of glucose **(b)**, glucose-6-phosphate (G6P) **(c)**, fructose bisphosphate (FBP) **(d)**, dihydroxyacetone phosphate (DHAP) **(e)**, phosphoenolpyruvate (PEP) **(f)**, pyruvate **(g)**, lactate **(h),** and alanine **(i)** in hearts from fl/fl and CS-Mpc2-/- mice after injection with U-$^{13}$C-glucose, n=11-12. **j** Cardiac glycogen levels measured biochemically in fl/fl and CS-Mpc2-/- littermates, n=16. **k** Representative transmission electron microscopy images at 5,000X magnification of the cardiac papillary muscle of fl/fl and CS-Mpc2-/-, white arrowhead denotes glycogen granules. Scale bars = 1 μm. **l-m** $^{13}$C % enrichment (**l**) and pool size (**m**) of uridine diphosphate (UDP)-glucose**,** n=11-12. **n-o** $^{13}$C % enrichment of the glucose within glycogen for fully labeled M6 **(n),** and fully unlabeled M0 **(o)**, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction. See also Supplementary Figs. 1-4.

**Fig. 3: Glucose uptake is increased, and glycogen synthesis should be inhibited in CS-Mpc2-/- hearts. a** Glucose uptake assessed by 2-deoxyglucose in the present or absence of insulin in cardiac muscle fibers from fl/fl and CS-Mpc2-/- littermates, n=9-12. **b** Western blot images of phosphorylated (serine-473) and total protein kinase B (AKT) and the ratio of quantified densitometry for pAKT/AKT from hearts of fl/fl and CS-Mpc2-/- mice injected with either saline vehicle or 5mU/g insulin. **c-g** Western blot images of phosphorylated glycogen synthase (pGS) serine-640, total glycogen synthase (GS), muscle glycogen phosphorylase (PYGM), glycogenin-1, mitochondrial pyruvate carrier (MPC) 1, MPC2, and $\alpha$-tubulin (**c**), and the normalized densitometry quantification for MPC1/tubulin **(d)**, MPC2/tubulin **(e)**, pGS/total GS **(f)**, and glycogenin-1/tubulin **(g)** in hearts of fl/fl and CS-Mpc2-/- littermates, n=4. Data are presented as mean±SD. Data in **a** were evaluated by two-way analysis of variance (ANOVA) with Tukey post-hoc

944 multiple-comparisons test. Data in **b** were evaluated by unpaired, two-tailed Student's t-
945 test with Welch correction due to only n=2 in saline groups. Data in **d-g** were evaluated
946 by unpaired, two-tailed Student's t-test with Welch correction.
947
**Fig. 4: Non-failing CS-Mpc2-/- hearts from young mice display increased**
**glycolysis, but no glycogen accumulation. a-k** $^{13}$C % enrichment of glucose **(a)**,
gluocse-6-phosphate (G6P) **(b)**, fructose bisphosphate (FBP) **(c)**, dihydroxyacetone
phosphate (DHAP) **(d)**, 2/3-phosphoglycerate (PG) **(e)**, phosphoenolpyruvate (PEP) **(f)**,
pyruvate **(g)**, lactate **(h)**, alanine **(i)**, and uridine diphosphate (UDP)-glucose **(j)**, and the
UDP-glucose normalized pool size (**k**) in hearts of young non-failing CS-Mpc2-/- and fl/fl
littermates injected i.p. with U-$^{13}$C-glucose,  n=8-10. **l** Heart glycogen content measured
biochemically in young non-failing CS-Mpc2-/- and fl/fl littermates, n=6-7. **m**
Representative transmission electron microscopy images at 15,000X magnification of
the cardiac papillary muscle in young fl/fl and CS-Mpc2-/- littermates. Scale bars = 1
μm. Data are presented as mean±SD, and were analyzed by unpaired, two-tailed
Student's t-test with Welch correction. See also Supplementary Figs. 5-6.
960
**Fig. 5: Inhibition of glycogen synthase with MZ-101 reduces glycogen and**
**improves heart failure in CS-Mpc2-/- mice. a-b** Blood glucose **(a)** and blood lactate
**(b)** measured after a 2 hour fast from fl/fl and CS-Mpc2-/- mice treated with either
vehicle or 100 mg/kg MZ-101 twice daily for 2 weeks, n=7-12. **c-e** Heart **(c)**, soleus **(d)**,
and liver **(e)** glycogen content measured biochemically in fl/fl and CS-Mpc2-/- mice
treated with either Veh or MZ-101, n=7-12. **f** Heart weight to tibia length ratio (HW/TL) of
fl/fl and CS-Mpc2-/- mice treated with either Veh or MZ-101, n=7-12. **g** Representative
H&E stained images of mid-ventricular short axis sections of fl/fl and CS-MPC2-/- mice
treated with either Veh or MZ-101. Scale bars = 1 mm. **h-i** Cardiac gene expression for
Nppa and Nppb markers of heart failure from fl/fl and CS-Mpc2-/- mice treated with
either vehicle or 100 mg/kg MZ-101 twice daily for 2 weeks, n=7-12. **j** Representative
western blot images for phosphorylated and total glycogen synthase 1 and α-Tubulin,
and normalized quantified densitometry for phosphorylated/total glycogen synthase, n=4
per group. Data are presented as mean±SD, and were analyzed by unpaired, two-tailed
Student's t-test with Welch correction.
976
**Fig. 6: Ketogenic diet normalizes the CS-Mpc2-/- heart failure and glycogen**
**accumulation. a** Heart weight to tibia length ratio (HW/TL) of fl/fl and CS-Mpc2-/- mice
fed either low fat (LF) or ketogenic diet (KD), n=4-7. **b** Heart glycogen content measured
biochemically in fl/fl and CS-Mpc2-/- mice fed either a LF diet or KD, n=4-7. **c**
Representative transmission electron microscopy images at 5,000X magnification of the
cardiac papillary muscle of fl/fl and CS-Mpc2-/- mice fed LF or KD. Arrowheads denote
glycogen granules. Scale bars = 1 μm. **d** Western blot images of phosphorylated
glycogen synthase (serine 640) and total glycogen synthase from hearts of fl/fl or CS-
Mpc2-/- mice fed either LF or KD, n=2. **e-j** $^{13}$C % enrichment of glucose **(e)** and the
glucose within glycogen **(f)**, and the normalized pool size of glucose-6-phosphate (G6P)
**(g)**, pyruvate **(h)**, uridine diphosphate (UDP)-glucose **(i)**, and glycogen **(j)** in fl/fl and CS-
Mpc2-/- mice that were fed LF or KD and injected i.p. with U-$^{13}$C-glucose, n=6-7. Data

are presented as mean±SD, and were analyzed by two-way analysis of variance (ANOVA) with Tukey post-hoc multiple-comparisons test. See also Supplementary Fig. 7.

**Fig. 7: AngII-induced cardiac hypertrophy increases glycogen and is improved by ketogenic diet. a** and **b** Blood glucose (**a**) and plasma β-hydroxybutyrate (β-HB) concentrations (**b**) of saline vehicle or AngII-infused mice fed low fat (LF) diet or ketogenic diet (KD), n=7-10. **c** Heart weight to tibia length ratio (HW/TL) of saline vehicle and AngII infused mice fed LF or KD, n=7-10. **d** Cardiac glycogen measured biochemically from mice infused with saline or AngII fed either LF or KD, n=7-10. **e** and **f** Normalized gene expression for *Mpc1* (**e**) and *Mpc2* (**f**) in hearts from mice infused with saline or AngII fed LF or KD, n=7-10. **g** and **h** representative western blot images of phosphorylated glycogen synthase (pGS1), total glycogen synthase (GS1), glycogenin-1, and $\alpha$-tubulin (**g**), and the normalized densitometry quantified for pGS/GS (**h**) from hearts of mice infused with saline or AngII fed LF or KD, n=4. Data are presented as mean±SD, and were analyzed by two-way analysis of variance (ANOVA) with Tukey post-hoc multiple-comparisons test.

**Supplementary Fig. 1: Glycolytic pool sizes in failing MPC-/- hearts.** Related to Fig. 2. **a** Schematic pathway of glycolysis and accessory glucose pathways. **b-l** Normalized pool size of glucose **(b)**, glucose-6-phosphate (G6P) **(c)**, fructose bisphosphate (FBP) **(d)**, dihydroxyacetone phosphate (DHAP) **(e)**, 2/3-phosphoglycerate (PG) **(f)**, phosphoenolpyruvate (PEP) **(g)**, pyruvate **(h)**, lactate **(i)**, alanine **(j)**, and the $^{13}$C % enrichment of glycerol 3-phosphate (Glycerol 3P) **(k)** and glycerol 3P pool **(l)** of failing CS-Mpc2-/- hearts and fl/fl littermates injected with U-$^{13}$C-glucose, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

**Supplementary Fig. 2: TCA cycle enrichment and pool sizes in failing CS-Mpc2-/- hearts.** Related to Fig. 2. **a-l** the $^{13}$C % enrichment and pool sizes of citrate **(a-b)**, aconitate **(c-d)**, $\alpha$-ketoglutarate (KG) **(e-f)**, succinate **(g-h)**, fumarate **(i-j)**, and malate **(k-l)** in fl/fl and failing CS-Mpc2-/- hearts of mice injected i.p. with U-$^{13}$C-glucose, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

**Supplementary Fig. 3: MPC deletion reduces pyruvate oxidation.** Related to Fig. 2. **a-b** Oxygen consumption rates (OCR) measured from isolated cardiac mitochondria (**a**) and permeabilized cardiac muscle fibers (**b**) from hearts of fl/fl and CS-Mpc2-/- littermates stimulated with pyruvate/malate (P/M), P/M plus adenosine diphosphate (ADP), and P/M, ADP, and the MPC inhibitor UK-5099 (5 $\mu$M), n=4-5. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

**Supplementary Fig. 4: No major changes in the pentose phosphate pathway in failing MPC-/- hearts.** Related to Fig. 2. **a** Schematic of $^{13}$C enrichment of the pentose

phosphate pathway from U-$^{13}$C-glucose. Metabolites in grey were not measured. **b-f** $^{13}$C
% enrichment and normalized pool size of 6-phosphogluconate (PG) **(b)**, ribose-5-
phosphate (R5P) **(c)**, sedoheptulose 7-phosphate (Sedo7P) **(d)**, serine **(e)**, and glycine
**(f)**, in hearts from fl/fl and failing CS-Mpc2-/- mice injected with U-$^{13}$C-glucose, n=11-12.
Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed
Student's t-test with Welch correction.
1040
**Supplementary Fig. 5: Glycolytic pool sizes in young, non-failing MPC-/- hearts.**
Related to Fig. 4. **a** Schematic of glycolysis and accessory glucose pathways. **b-q**
Normalized pool sizes of glucose **(b)**, glucose-6-phosphate (G6P) (**c**), fructose
bisphosphate (FBP) (**d**), dihydroxyacetone phosphate (DHAP) **(e)**, 2/3
phosphoglycerate (PG) **(f)**, phosphoenolpyruvate (PEP) **(g)**, pyruvate **(h)**, lactate **(i)**,
alanine (j), ribose-5-phosphate (R5P) **(k)**, and the $^{13}$C enrichment and pool size of
serine **(l-m)**, glycine **(n-o)**, and glycerol 3-phosphate (glycerol 3P) **(p-q)** in young non-
failing CS-Mpc2-/- and fl/fl littermates injected with U-$^{13}$C-glucose, n=8-10. Data are
presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test
with Welch correction.
1051
**Supplementary Fig. 6: TCA cycle enrichment and pool sizes in young, non-failing**
**CS-Mpc2-/- hearts.** Related to Fig. 4. **a-l** $^{13}$C % enrichment and normalized pool sizes
of citrate (**a-b**), aconitate (**c-d**), $\alpha$-ketoglutarate ($\alpha$KG) (**e-f**), succinate (**g-h**), fumarate (**i-**
**j**), and malate (**k-l**), in hearts of young, nonfailing CS-Mpc2-/- and fl/fl littermates
injected with U-$^{13}$C-glucose, n=8-10. Data are presented as mean±SD. Data were
evaluated by unpaired, two-tailed Student's t-test with Welch correction.
1058
**Supplementary Fig. 7: Ketogenic diet decreases TCA cycle enrichment from**
**glucose.** Related to Fig. 6. **a-l** $^{13}$C % enrichment of pyruvate (**a**), lactate (**b**), alanine (**c**),
citrate (**d**), aconitate (**e**), $\alpha$-ketoglutarate ($\alpha$KG) (**f**), glutamate (**g**), glutamine (**h**),
succinate (**i**), fumarate (**j**), malate (**k**), and aspartate (**l**) in hearts from fl/fl and CS-Mpc2-
/- fed either low-fat (LF) or ketogenic diet (KD) and injected with U-$^{13}$C-glucose, n=6-7.
Data are presented as mean±SD. Data were evaluated by two-way analysis of variance
(ANOVA) with Tukey post-hoc multiple-comparisons test.

Figure 1.

**Fig. 1: Heart Failure in CS-Mpc2-/- mice is not due to energetic stress. a** Heart weight/tibia length (HW/TL) in CS-Mpc2-/- and fl/fl littermates, n=6. **b** Representative gross images of excised fl/fl and CS-Mpc2-/- hearts. Scale bar = 5 mm. **c** Heatmap of differentially expressed genes identified by RNA sequencing of hearts from fl/fl and CS-Mpc2-/- littermates, n=3. Most prominently dysregulated KEGG pathways upregulated and downregulated are listed. **d-n** mass spectrometry normalized peak area pool sizes of adenosine triphosphate (ATP) **(d)**, adenosine diphosphate (ADP) **(e)**, adenosine monophosphate (AMP) **(f)**, phosphocreatine **(g)**, creatine **(h)**, phosphocreatine/creatine (PCr/Cr) ratio **(i)**, PCr/ATP ratio **(j)** oxidized nicotinamide adenine dinucleotide (NAD$^+$) **(k)**, reduced nicotinamide adenine dinucleotide (NADH) **(l)**, oxidized nicotinamide adenine dinucleotide phosphate (NADP$^+$) **(m)**, and reduced nicotinamide adenine dinucleotide phosphate (NADPH) **(n)** in hearts of fl/fl and CS-Mpc2-/- littermates that had been injected i.p. with U-$^{13}$C-glucose, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

# Figure 2.

**Fig. 2: Failing CS-Mpc2-/- hearts display increased glycolysis and accumulate glycogen. a** Schematic of U-$^{13}$C-glucose labeling of glycolytic metabolites and glycogen, created with BioRender. **b-i** $^{13}$C % enrichment of glucose **(b)**, glucose-6-phosphate (G6P) **(c)**, fructose bisphosphate (FBP) **(d)**, dihydroxyacetone phosphate (DHAP) **(e)**, phosphoenolpyruvate (PEP) **(f)**, pyruvate **(g)**, lactate **(h),** and alanine **(i)** in hearts from fl/fl and CS-Mpc2-/- mice after injection with U-$^{13}$C-glucose, n=11-12. **j** Cardiac glycogen levels measured biochemically in fl/fl and CS-Mpc2-/- littermates, n=16. **k** Representative transmission electron microscopy images at 5,000X magnification of the cardiac papillary muscle of fl/fl and CS-Mpc2-/-, white arrowhead denotes glycogen granules. Scale bars = 1 μm. **l-m** $^{13}$C % enrichment (**l**) and pool size (**m**) of uridine diphosphate (UDP)-glucose**,** n=11-12. **n-o** $^{13}$C % enrichment of the glucose within glycogen for fully labeled M6 **(n),** and fully unlabeled M0 **(o)**, n=11-12. Data are presented as mean±SD. Data were evaluated by unpaired, two-tailed Student's t-test with Welch correction. See also Supplementary Figs. 1-4.

# Figure 3.

**Fig. 3: Glucose uptake is increased, and glycogen synthesis should be inhibited in CS-Mpc2-/- hearts. a** Glucose uptake assessed by 2-deoxyglucose in the present or absence of insulin in cardiac muscle fibers from fl/fl and CS-Mpc2-/- littermates, n=9-12. **b** Western blot images of phosphorylated (serine-473) and total protein kinase B (AKT) and the ratio of quantified densitometry for pAKT/AKT from hearts of fl/fl and CS-Mpc2-/- mice injected with either saline vehicle or 5mU/g insulin. **c-g** Western blot images of phosphorylated glycogen synthase (pGS) serine-640, total glycogen synthase (GS), muscle glycogen phosphorylase (PYGM), glycogenin-1, mitochondrial pyruvate carrier (MPC) 1, MPC2, and $\alpha$-tubulin (**c**), and the normalized densitometry quantification for MPC1/tubulin **(d)**, MPC2/tubulin **(e)**, pGS/total GS **(f)**, and glycogenin-1/tubulin **(g)** in hearts of fl/fl and CS-Mpc2-/- littermates, n=4. Data are presented as mean±SD. Data in **a** were evaluated by two-way analysis of variance (ANOVA) with Tukey post-hoc multiple-comparisons test. Data in **b** were evaluated by unpaired, two-tailed Student's t-test with Welch correction due to only n=2 in saline groups. Data in **d-g** were evaluated by unpaired, two-tailed Student's t-test with Welch correction.

Figure 4.

**Fig. 4: Non-failing CS-Mpc2-/- hearts from young mice display increased glycolysis, but no glycogen accumulation. a-k** $^{13}$C % enrichment of glucose **(a)**, gluocse-6-phosphate (G6P) **(b)**, fructose bisphosphate (FBP) **(c)**, dihydroxyacetone phosphate (DHAP) **(d)**, 2/3-phosphoglycerate (PG) **(e)**, phosphoenolpyruvate (PEP) **(f)**, pyruvate **(g)**, lactate **(h)**, alanine **(i)**, and uridine diphosphate (UDP)-glucose **(j)**, and the UDP-glucose normalized pool size **(k)** in hearts of young non-failing CS-Mpc2-/- and fl/fl littermates injected i.p. with U-$^{13}$C-glucose, n=8-10. **l** Heart glycogen content measured biochemically in young non-failing CS-Mpc2-/- and fl/fl littermates, n=6-7. **m** Representative transmission electron microscopy images at 15,000X magnification of the cardiac papillary muscle in young fl/fl and CS-Mpc2-/- littermates. Scale bars = 1 μm. Data are presented as mean±SD, and were analyzed by unpaired, two-tailed Student's t-test with Welch correction. See also Supplementary Figs. 5-6.

# Figure 5.

**a** Blood Glucose (mg/dL) — Veh, MZ-101 groups (fl/fl, Mpc2-/-)
p = 0.9729, p = 0.7462, p = 0.4022, p = 0.8199

**b** Blood Lactate (mM) — Veh, MZ-101
p = 0.0079, p = 0.1516, p = 0.9060

**c** Heart Glycogen (mmol/g) — Veh, MZ-101
p < 0.0001, p = 0.0002, p = 0.9987

**d** Soleus Glycogen (mmol/g) — Veh, MZ-101
p > 0.9999, p = 0.0230, p = 0.0202, p > 0.9999

**e** Liver Glycogen (mmol/g) — Veh, MZ-101
p = 0.3963, p = 0.5029, p = 0.9962, p = 0.9972

**f** HW/TL (mg/mm) — Veh, MZ-101
p < 0.0001, p = 0.0223, p = 0.0055

**g** Vehicle: fl/fl, CS-Mpc2-/- ; MZ-101: fl/fl, CS-Mpc2-/-

**h** *Nppa* — Normalized Expression — Veh, MZ-101
p < 0.0001, p = 0.0227, p = 0.5258

**i** *Nppb* — Normalized Expression — Veh, MZ-101
p = 0.0021, p = 0.0237, p = 0.6852

**j** Vehicle: fl/fl, MPC2-/- ; MZ-101: fl/fl, MPC2-/-
pGS (S640), GS, $\alpha$-Tubulin

Norm. pGS/Total GS — Veh, MZ-101
p = 0.0013, p = 0.0024, p = 0.1724

**Fig. 5: Inhibition of glycogen synthase with MZ-101 reduces glycogen and improves heart failure in CS-Mpc2-/- mice. a-b** Blood glucose **(a)** and blood lactate **(b)** measured after a 2 hour fast from fl/fl and CS-Mpc2-/- mice treated with either vehicle or 100 mg/kg MZ-101 twice daily for 2 weeks, n=7-12. **c-e** Heart **(c)**, soleus **(d)**, and liver **(e)** glycogen content measured biochemically in fl/fl and CS-Mpc2-/- mice treated with either Veh or MZ-101, n=7-12. **f** Heart weight to tibia length ratio (HW/TL) of fl/fl and CS-Mpc2-/- mice treated with either Veh or MZ-101, n=7-12. **g** Representative H&E stained images of mid-ventricular short axis sections of fl/fl and CS-MPC2-/- mice treated with either Veh or MZ-101. Scale bars = 1 mm. **h-i** Cardiac gene expression for *Nppa* and *Nppb* markers of heart failure from fl/fl and CS-Mpc2-/- mice treated with either vehicle or 100 mg/kg MZ-101 twice daily for 2 weeks, n=7-12. **j** Representative western blot images for phosphorylated and total glycogen synthase 1 and $\alpha$-Tubulin, and normalized quantified densitometry for phosphorylated/total glycogen synthase, n=4 per group. Data are presented as mean±SD, and were analyzed by unpaired, two-tailed Student's t-test with Welch correction.

# Figure 6.

**a** HW/TL (mg/mm) — LF: p < 0.0001; KD: p = 0.9880; between diets: p < 0.0001

**b** Heart Glycogen (mmol/g) — LF: p < 0.0001; KD: p = 0.9993; between diets: p < 0.0001

**c** Low-fat diet: fl/fl, CS-MPC2$^{-/-}$ | Ketogenic diet: fl/fl, CS-MPC2$^{-/-}$

**d** Low-fat diet (fl/fl, MPC2$^{-/-}$) | Ketogenic diet (fl/fl, MPC2$^{-/-}$)
p-Glycogen Synthase (S640)
Glycogen Synthase

**e** Glucose M6 — % Enrichment; LF: p = 0.9822; KD: p = 0.9974; p = 0.0007; p = 0.0024

**f** Glycogen M6 — % Enrichment; LF: p = 0.0421; KD: p = 0.2641; p = 0.2081; p = 0.7510

**g** G6P pool — Normalized Peak Area; LF: p = 0.0385; KD: p > 0.9999; p = 0.2576

**h** Pyruvate pool — Normalized Peak Area; LF: p = 0.0039; KD: p = 0.9998; p = 0.0497

**i** UDP-glucose pool — Normalized Peak Area; LF: p = 0.0006; KD: p = 0.9863; p = 0.0030

**j** Glycogen pool — Normalized Peak Area; LF: p < 0.0001; KD: p = 0.9919; p < 0.0001

fl/fl ○   CS-Mpc2-/- ○

**Fig. 6: Ketogenic diet normalizes the CS-Mpc2-/- heart failure and glycogen accumulation. a** Heart weight to tibia length ratio (HW/TL) of fl/fl and CS-Mpc2-/- mice fed either low fat (LF) or ketogenic diet (KD), n=4-7. **b** Heart glycogen content measured biochemically in fl/fl and CS-Mpc2-/- mice fed either a LF diet or KD, n=4-7. **c** Representative transmission electron microscopy images at 5,000X magnification of the cardiac papillary muscle of fl/fl and CS-Mpc2-/- mice fed LF or KD. Arrowheads denote glycogen granules. Scale bars = 1 $\mu$m. **d** Western blot images of phosphorylated glycogen synthase (serine 640) and total glycogen synthase from hearts of fl/fl or CS-Mpc2-/- mice fed either LF or KD, n=2. **e-j** $^{13}$C % enrichment of glucose **(e)** and the glucose within glycogen **(f)**, and the normalized pool size of glucose-6-phosphate (G6P) **(g)**, pyruvate **(h)**, uridine diphosphate (UDP)-glucose **(i)**, and glycogen **(j)** in fl/fl and CS-Mpc2-/- mice that were fed LF or KD and injected i.p. with U-$^{13}$C-glucose, n=6-7. Data are presented as mean±SD, and were analyzed by two-way analysis of variance (ANOVA) with Tukey post-hoc multiple-comparisons test. See also Supplementary Fig. 7.

Figure 7.

**Fig. 7: AngII-induced cardiac hypertrophy increases glycogen and is improved by ketogenic diet. a** and **b** Blood glucose (**a**) and plasma β-hydroxybutyrate (β-HB) concentrations (**b**) of saline vehicle or AngII-infused mice fed low fat (LF) diet or ketogenic diet (KD), n=7-10. **c** Heart weight to tibia length ratio (HW/TL) of saline vehicle and AngII infused mice fed LF or KD, n=7-10. **d** Cardiac glycogen measured biochemically from mice infused with saline or AngII fed either LF or KD, n=7-10. **e** and **f** Normalized gene expression for *Mpc1* (**e**) and *Mpc2* (**f**) in hearts from mice infused with saline or AngII fed LF or KD, n=7-10. **g** and **h** representative western blot images of phosphorylated glycogen synthase (pGS1), total glycogen synthase (GS1), glycogenin-1, and $\alpha$-tubulin (**g**), and the normalized densitometry quantified for pGS/GS (**h**) from hearts of mice infused with saline or AngII fed LF or KD, n=4. Data are presented as mean±SD, and were analyzed by two-way analysis of variance (ANOVA) with Tukey post-hoc multiple-comparisons test.