A VLP vaccine platform comprising the core protein of hepatitis B virus with

N-terminal antigen capture.

Kaniz Fatema*, Joseph S. Snowden*, Alexander Watson, Lee Sherry, Neil A.

Ranson, Nicola J. Stonehouse# and David J. Rowlands#

Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology,

Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, United Kingdom

* These authors contributed equally to the study.

Address correspondence to David J. Rowlands, D.J.Rowlands@leeds.ac.uk and Nicola J.

Stonehouse, N.J.Stonehouse@leeds.ac.uk

Keywords: vaccine, platform, virus-like particle, ClearColi, HBcAg, Junín virus

Word count

Abstract: 179; Body text: 3,450 (including references)

Abstract

Nanoparticle presentation systems offer the potential to develop new vaccines rapidly in response to emerging diseases, a public health need that has become increasingly evident in the wake of the COVID-19 pandemic. Previously, we reported a nanoparticle scaffold system termed VelcroVax. This was constructed by insertion of a high affinity SUMO binding protein (Affimer), able to recognise a SUMO peptide tag, into the major immunodominant region of VLPs assembled from a tandem (fused dimer) form of hepatitis B virus (HBV) core protein (HBc). Here we describe a modified form of VelcroVax, comprising monomeric HBc with the Affimer inserted at the N-terminus (termed N-VelcroVax). In contrast to the tandem form of VelcroVax, N-VelcroVax VLPs were expressed well in *E. coli.* The VLPs effectively bound SUMO-tagged Junín virus glycoprotein, gp1 as assessed by structural and serological analyses. Cryo-EM characterisation of N-VelcroVax complexed with a SUMO-Junín gp1 showed continuous density attributable to the fused Affimer, in addition to evidence of target antigen capture. Collectively, these data suggest that N-VelcroVax has potential as a versatile next generation vaccine scaffold.

Introduction

Virus vaccines developed by conventional means, e.g. attenuation or chemical inactivation, have been highly effective in reducing disease burdens in both humans and domesticated animals. Furthermore, advances in molecular biology and immunology in recent decades have expanded the options available for development of novel vaccines against intractable diseases and against newly emerging pathogens. Newer approaches to vaccine development include recombinant expression of protein subunits, recombinant live viral vectors, DNA, mRNA and nanoparticle vaccine platforms [1-3]. The original nanoparticle vaccines were virus-like particles (VLPs) produced by recombinant expression of self-assembling viral structural proteins. The first VLP vaccine to be licenced for human use was the hepatitis B surface antigen vaccine (HBsAg), which has been used to immunise many millions of people since its introduction in the 1980s [4, 5]. More recently, VLP vaccines have been successfully developed to immunise against human papillomavirus. These VLP vaccines provide immunity against the viruses from which the VLPs originated [6]. However, the principle of using VLPs as presentation systems for antigens derived from heterologous pathogens has been exemplified by the licencing of Mosquirix, which comprises HBsAg modified to include antigenic components of the malaria parasite [7]. In addition to viral structural proteins, other self-assembling proteins have been shown to produce nanoparticles suitable for chemical modification to provide candidate vaccine platforms [8, 9].

VLPs lack viral genetic material and therefore have no replicative ability and are intrinsically safe. Additionally, due to their structural resemblance to native viruses, VLPs inherently possess "self-adjuvanting" immunogenic properties to elicit a strong immune response. Moreover, VLPs are able to elicit both humoral and cellular immune responses initiating the production of specific antibodies and activating cytotoxic T-cells [10-13]. In addition to acting as antigens themselves, VLPs can be modified to display foreign epitopes [14-16]. A key

advantage of using VLP-based vaccine platforms is the ability to present antigens in a dense,

highly repetitive manner, which leads to cross-linking of B cell receptors (BCRs) and B cell

activation. Many studies have shown that VLP-based vaccines containing multivalent antigens

elicit robust B cell activation and signalling and induce more effective immune responses than

monovalent antigens [11, 13, 17, 18].

The capsid (or core) protein of hepatitis B virus (HBc) readily forms icosahedral VLPs when

expressed in a wide range of prokaryotic and eukaryotic systems, which are inherently highly

immunogenic. Consequently, the HBc VLP system has long been investigated as a potential

vaccine platform [14, 15, 19-21]. HBc protein monomers comprise either 183 or 185 amino

acids and dimerise to produce structural subunits which assemble to form icosahedral VLPs with

$T = 3$ (90 dimers) or $T = 4$ (120 dimers) morphologies [22]. The dimeric assembly results in

the formation of 4 helix bundles (spikes) which project from the particle surface and a sequence

at the tips of the spikes comprises the major immunodominant region (MIR). The MIR has

been used as a preferred site for insertion of foreign epitopes because of its immunodominance

[16, 23, 24] but this also poses challenges. There is a potential for steric clashes when two

closely located copies of the inserted sequence are present at the tips of the spikes [23, 25].

Furthermore, insertion of large or hydrophobic sequences can interfere with assembly and

therefore abrogate VLP formation [26, 27]. We previously addressed this problem by fusing

two copies of HBc protein to produce tandem HBc VLPs, which allows the insertion of a

foreign sequence into one MIR while leaving the other MIR in the tandem partner unmodified,

resulting in the presence of only one foreign epitope per spike [23]. This approach allowed us

to develop VelcroVax, incorporating an antigen capture sequence, an Affimer (a non-antibody

high affinity artificial binding protein), inserted at a single MIR per spike [16]. Although these

constructs form VLPs when expressed in eukaryotic systems, such as yeast or plants, they do

not assemble efficiently when expressed in *E.coli*. Consequently, here we have developed an

alternative antigen capture platform compatible with expression in *E.coli*. The MIR is the most

exposed part of the VLP and consequently is the immunodominant region [19, 23]. Therefore,

in order to reduce the inherent anti-hepatitis B virus immunogenicity of the VLPs, we modified

the natural HBc protein sequence by inserting 5 additional amino acids within the MIR. This

construct (which we refer to as wild type (wt) HBc 190) was modified to include an Affimer

fused to the N-terminus and termed N-VelcroVax. The resulting VLPs were able to bind a viral

glycoprotein (Junín gp1) bearing the appropriate affinity tag as demonstrated by ELISA,

gradient centrifugation and cryo-EM.

Results

Expression of N-VelcroVax in *E. coli*

We previously generated a vaccine platform system termed VelcroVax in which an anti-SUMO Affimer was inserted into the MIR of one of the HBc monomers within a fused HBc dimer. The Affimer was presented on the surface of assembled VLPs and was able to bind SUMO-tagged target antigens [16]. Here, we have modified a monomeric HBc protein containing a 5 amino acid insertion into the MIR, termed wt HBc 190 (Fig.1A), by introducing the anti-SUMO Affimer sequence (Fig. 1C) at the N-terminus to produce N-VelcroVax (Fig 1B). This protein was expressed in *E.coli*, using the ClearColi BL21 (DE3) strain to minimise contamination with bacteria-derived pyrogens. Both wt HBc 190 and N-VelcroVax were efficiently expressed, as demonstrated by Western Blot analysis using anti-HBc (10E11) antibodies (Fig. 1D), with proteins of 22 kDa and 35 kDa detected, as expected. A minor band apparent in the N-VelcroVax sample is likely to be a degradation product.

N-VelcroVax assembles into VLPs.

After confirmation of expression of N-VelcroVax in ClearColi BL21 (DE3) *E. coli*, cell extracts were purified by differential ultra-centrifugation including separation of assembled VLPs on sucrose gradients. Gradient fractions were then analysed by Coomassie blue staining and Western blot which confirmed the presence of N-VelcroVax VLPs (Fig. 2A) [16]. The purity of the sucrose gradient separated VLPs can be assessed from the lack of visible contaminating proteins in the Coomassie stained gel. Examination of the peak fraction (fraction 5) by negative stain transmission electron microscopy (TEM) showed VLPs of the expected morphology (Fig. 2B).

Complexes of SUMO-Junín gp1 and N-VelcroVax VLP co-sediment during sucrose gradient centrifugation.

We examined the SUMO-binding properties of N-VelcroVax VLPs using a SUMO-tagged gp1

protein of Junín virus, which we had previously shown to bind to the Affimer when presented

at the MIR of tandem VelcroVax [16]. For the initial investigation of binding, N-VelcroVax

VLPs were mixed with SUMO-Junín gp1 at a 1:1 molar ratio overnight at 4°C and separated

by 15 - 45% sucrose gradient ultracentrifugation. The resulting fractions were analysed by

Western blot using anti-HBc (10E11) and anti-Junín antibodies to detect the positions of N-

VelcroVax (Fig. 3A) and SUMO-Junín gp1 (Fig. 3B) respectively in the gradients. Both HBc

N-VelcroVax (35 kDa) and SUMO-Junín gp1 (54 kDa) were present in the same gradient

fractions, indicative of a non-covalent interaction (Fig. 3). As shown in Fig. 3A, the majority

of the SUMO-Junín gp1 co-sedimented with the VLPs, although some unbound material

(29.8 +/- 3.8 %) was detected in the top gradient fractions.

Evaluation of SUMO-Junín gp1 binding to N-VelcroVax by ELISA.

The SUMO-Junín gp1 interaction with HBc N-VelcroVax was assessed by ELISA. SUMO-

Junín gp1 and N-VelcroVax were mixed before loading onto ELISA plates. Molar ratios ranged

from 1:1 to 5:1 (HBc monomer:gp1). After washing, the wells were interrogated with anti-HBc

or anti-SUMO-Junín gp1 antisera. The results indicated that N-VelcroVax VLPs clearly bound

SUMO-Junín gp1 with maximum binding occurring when the components were mixed at a 1:1

ratio (Fig. 4). No binding of SUMO-Junín gp1 to wt HBc 190 VLPs was detected, as expected

(Fig. 4).

Structural characterisation of N-VelcroVax and N-VelcroVax:SUMO-Junín gp1 particles.

N-VelcroVax samples were vitrified for cryo-EM data collection (Fig. S1, Table S1). This

confirmed two different sizes of icosahedral particles. Reconstructions of each particle stack

with icosahedral symmetry imposed revealed that these corresponded to $T = 3$ and $T = 4$

configurations, as expected.

Interestingly, both reconstructions (resolved to ~3.5 Å and ~3.3 Å, respectively) showed

continuous densities located between the four-helix bundles on the surface of the VLPs (Fig.

5). Each of these densities was of appropriate dimension to accommodate an Affimer and was

located close to the residue at the N-terminus of a wt HBc 190 monomer, as expected.

Unstructured internal density was also observed, possibly representing non-specific binding of

nucleic acid derived from the *E.coli* expression system as reported previously [28] (Fig. 5).

Next, we attempted to visualise N-VelcroVax VLPs in complex with a SUMO-tagged antigen,

specifically, the gp1 glycoprotein of Junín virus (SUMO-gp1). As initial screening by negative

stain TEM of N-VelcroVax:SUMO-gp1 complexes revealed significant aggregation, an

alternative on-grid binding approach was pursued (i.e. separate samples of N-VelcroVax and

SUMO-gp1 were applied directly to the grid in sequence). This showed no evidence of

aggregation of N-VelcroVax particles and visual inspection suggested that the boundaries of

the VLPs had become less well defined, indicative of decoration with SUMO-gp1 molecules

(Fig. 6A).

This on-grid binding approach was subsequently used to prepare samples for cryo-EM analysis

of N-VelcroVax:SUMO-gp1 (Fig. S2, Table S1). Processing of cryo-EM data yielded density

maps for $T = 3$ and $T = 4$ VLPs resolved to 3.0 Å and 3.5 Å, respectively. Initial inspection of

density maps revealed no additional density attributable to bound SUMO-gp1, although there

appeared to be a subtle change in the orientations of the Affimers, with these being closer

together in the N-VelcroVax:SUMO-gp1 complex map compared to the unliganded N-

VelcroVax map (Fig. 6B).

Hypothesising that the SUMO-gp1 was perhaps bound at a low occupancy, we attempted to

resolve density for the glycoprotein by performing focussed classification, as described

previously [29-32]. A mask was applied to a single Affimer including the volume expected to

contain SUMO-gp1 for both $T = 3$ and $T = 4$ VLPs (Fig. S3). Interestingly, several focussed

classes revealed low resolution density continuous with Affimer density, suggesting that

SUMO-gp1 had been captured by N-VelcroVax Affimers but was only present at a low

occupancy (Fig. 6C), in contrast to the ELISA and sucrose density gradient data, which

suggested that the occupancy by SUMO-gp1 was higher. This is most likely due to the marked

differences in the methods used for complex formation for the solution-based and cryo-EM

studies. Interestingly, density corresponding to the SUMO-gp1-binding Affimer in asymmetric

reconstructions of focussed classes had reoriented upwards, suggesting that movement of the

Affimer was necessary to accommodate the size of SUMO-gp1 (Fig. 6D).

Discussion

The development of vaccine platforms suitable for multimeric presentation of antigenic components of important pathogens, so maximising their immune recognition, is an attractive approach toward novel vaccine development. To this end, we produced recombinant HBc VLPs (N-VelcroVax) with an anti-SUMO Affimer genetically fused at the N-terminus of HBc190 protein. Although HBc VLPs have been expressed in a number of expression systems, e.g., insect cells, mammalian cells, plants, yeast [6, 11, 13, 24, 33], using low cost expression systems would make the VLPs accessible more widely, especially for lower-to-middle income countries (LMICs). Hence, we opted for a cost-effective *E. coli* expression system which offers simple, fast and inexpensive production, ease of in-process control and high productivity [6, 13, 33-36]. However, contamination with high levels of endotoxin derived from the *E. coli* cell wall is a major drawback of this expression system [35, 37-39]. To address this problem, we used ClearColi BL21 (DE3) *E. coli* cells for expression of N-VelcroVax. A previous study showed that using ClearColi BL21 (DE3) *E. coli* for expression of apolipoprotein A and heat shock protein 70 significantly reduced endotoxin level as measured by LAL assay (10 μg/mL) and did not trigger endotoxic responses in HEK-Blue hTLR4 cells [37]. The endotoxin content of N-VelcroVax expressed in ClearColi BL21 (DE3) *E. coli* was 10.2 EU/mL as assessed by LAL assay, which is half the maximum recommended level of endotoxin acceptable for subunit vaccines [35, 37, 40].

Examination by negative stain TEM showed that the morphology of the N-VelcroVax VLPs was as expected [41, 42]. Cryo-EM analysis of the VLPs revealed particles in both $T = 3$ and $T = 4$ configurations, as expected. In our earlier analyses, Affimer density was not visible, likely as a result of the location of the Affimers on flexible linkers at the tips of four-helix bundles at the MIR [16]. However, for N-VelcroVax, in which Affimers are attached at the N-terminus and spatially constrained between the four-helix bundles, Affimer density was

resolved. This suggests that the Affimers in the N-VelcroVax system are more rigidly

constrained in their position compared to when presented at the MIR region.

N-VelcroVax was also visualised in complex with SUMO-gp1. Although SUMO-gp1 density

was not resolved in symmetric reconstructions, focussed classification revealed low resolution

density continuous with Affimer density. This finding is suggestive of low occupancy, with the

majority of unoccupied sites leading to the 'averaging out' of SUMO-gp1 density in symmetric

reconstructions. Low occupancy is likely a result of the on-grid binding method used during

grid preparation to prevent particle aggregation. Indeed, both sucrose density gradient and

ELISA assays demonstrated a much more efficient interaction in solution. Interestingly, an

asymmetric reconstruction derived from particles belonging to a single focussed class revealed

that the corresponding Affimer had undergone a major reorientation in order to accommodate

the target antigen, indicative of some steric hindrance.

Conclusion

Together with our earlier report of Affimer presentation at the MIR, these data demonstrate the

flexibility of the HBc VLP system as a vaccine scaffold and open the possibility of a dual

presentation system using both the N-terminus and the MIR for antigen display.

Materials and methods

Expression constructs

An anti-SUMO-Affimer sequence was genetically fused at the N-terminus of wt HBc 190 to produce N-VelcroVax (Fig. 2). This was subcloned into the pET29b expression vector using *Nde*1 and *Xho*1 and transformed into *E. coli* DH5α chemically competent cells for plasmid propagation.

For the target protein, Junín gp1, a SUMO-His6 tag [16] was genetically fused to the N-terminus of the Junín virus glycoprotein 1 to form SUMO-Junín GP1 (kindly supplied by Thomas A. Bowden, University of Oxford, UK). This protein was expressed in mammalian cells.

Expression and purification of N-VelcroVax

ClearColi BL21 (DE3) *E. coli* was transformed with the N-VelcroVax plasmid and used for protein expression. N-VelcroVax plasmid was mixed with ClearColi BL21 (DE3) *E. coli* cells and incubated on ice for 30 min before being transferred to a 42°C heat block for 50 seconds. Heat-shock transformed cells were then plated on LB-Agar plates (30 μg/mL kanamycin) and incubated overnight at 37°C. A single colony was picked and cultured in 10 mL starter LB culture (30 μg/mL kanamycin) overnight at 37°C and shaking at 200 rpm. The preculture was diluted 1:1000 into 500 mL LB (30 μg/mL kanamycin) and cultured at 37°C, 200 rpm until OD_{600} ~0.6 – 0.8. Protein expression was induced with isopropyl β-D-1-thiogalactopyranoside (IPTG) (0.1 μM) and incubated at 18°C, 200 rpm for a further 7 hours. The culture was centrifuged at $4000 \times g$ for 10 minutes and the pellet was lysed with chemical lysis buffer. N-VelcroVax VLPs were purified by sedimentation through a 30% sucrose cushion followed by 15-45% sucrose gradient ultracentrifugation. Fractions (1 mL) were collected manually (top down) and fractions containing the VLPs were analysed by SDS-PAGE 12% gel where the

protein bands were visualised by Coomassie blue R250 staining. The presence of HBcAg-reactive proteins was assessed by Western blot with MAb 10E11 using standard protocols. The protein content of fractions was assessed directly by BCA assay (Pierce, ThermoFisher Scientific).The concentration of purified N-VelcroVax was measured using bicinchoninic acid (BCA) assay (Pierce) which revealed a yield of protein ~ 6.5 mg/L and endotoxin content of 10.2 EU/mL (Pierce LAL Chromogenic Endotoxin Quantitation kit; Thermo Scientific). Purified N-VelcroVax VLPs were stored at 4°C.

Enzyme-linked immunosorbent assay (ELISA)

A non-competitive indirect ELISA was developed to assess the binding of N-VelcroVax vaccine scaffold to capture SUMO-tagged proteins using 96-well ELISA plates (Greiner Technologies, Bio-One, UK). The plate was coated with N-VelcroVax at 2.5 µg/mL (125 ng/well) and incubated overnight at 4 °C. The plate was blocked with blocking buffer (PBS + 2% skimmed milk) for 1 h at room temperature (RT) and then incubated with SUMO-tagged Junín gp1 for 1 hour at RT. Specific primary antibodies (mouse monoclonal anti-HBc 1 (10E11, Abcam ab8639), anti-Junín gp1 (NR 2567)) were added and incubated for 1 hour at RT. The bound primary antibodies were detected with HRP-conjugated goat anti-mouse IgG and incubated for 1 hour at RT. Subsequently, chromogenic substrate TMB (ThermoFisher, Waltham, MA, USA) was added and incubated for 20 minutes at RT. The plates were washed with wash buffer (PBS + 1% Tween 20) four times between each step. The reaction was stopped by 2 M H_2SO_4 (50 µL/well) and plates then analysed using a Biotech PowerWave XS2 plate reader.

Negative-stain electron microscopy

Samples were prepared for negative-stain transmission EM by application to carbon-coated 300-mesh copper grids (Agar Scientific, UK) that had been glow discharged in air at 10 mA

for 30 seconds immediately before use. Where N-VelcroVax or SUMO-Junín gp1 was imaged

alone, 3 µL of sample was applied directly to the grid surface for 30 seconds before washing

and staining. To probe the interaction of N-VelcroVax with SUMO-Junín gp1, the ligand was

applied directly to the grid surface for 30 seconds after first applying N-VelcroVax and blotting

away excess fluid. Following sample application, excess fluid was wicked away before grids

were washed twice with 10 µL dH$_2$O. To stain the sample, 10 µL 1-2% uranyl acetate solution

was applied to the grid and immediately removed by blotting, before an additional 10 µL 1-2%

uranyl acetate solution was applied for 30 seconds. Grids were blotted to leave a thin film of

stain and left to air dry.

Imaging was performed using an FEI Tecnai F20 transmission EM (operating at 200 kV with

a field emission gun), equipped with an FEI CETA camera (Astbury Biostructure Laboratory,

University of Leeds). Data were collected at various defocus values (−2.0 µm to −5.0 µm) at a

nominal magnification of 25,000× giving an object sampling of 0.418 nm/pixel.

Cryo-electron microscopy

N-VelcroVax (in the presence or absence of target antigen) was vitrified using a LEICA EM GP

plunge freezing device (Leica Microsystems, Wetzlar, Germany) in preparation for imaging by

cryo-EM. Lacey carbon 400-mesh copper grids coated with a <3-nm continuous carbon film

(Agar Scientific, UK) were glow discharged in air at 10 mA for 30 seconds prior to the

application of 3 µL N-VelcroVax (or N-VelcroVax:SUMO-gp1). For N-VelcroVax:SUMO-gp1,

an on-grid interaction approach was used, such that following the application of VLP, excess

fluid was manually blotted away and 3 µL SUMO-gp1 was applied. The sample was incubated

on the grid surface for 30 seconds at 80% relative humidity (8°C) before excess fluid was blotted

for a duration between 1 – 4 seconds and the grid vitrified in liquid ethane (cooled to −179°C by

liquid nitrogen). Grids were transferred to liquid nitrogen for storage prior to clipping and

imaging with an FEI Titan Krios TEM (Astbury Biostructure Laboratory, University of Leeds).

Data collection was performed in integrating mode at a magnification of 75,000× (calibrated

object sampling of 1.065 Å/pixel) with the microscope operating at 300 kV (a full set of data

collection parameters is provided in Table S1).

Image Processing

Image processing was performed in Relion-3.0 and Relion-3.1 [43-45]. Following motion

correction and CTF estimation of micrographs [45], a subset of manually picked particles was

subjected to 2D classification, and class averages were used as templates for automated particle

picking. Initial particle stacks generated by automated picking were down-sampled (2× for N-

VelcroVax, 4× or 5× for N-VelcroVax:SUMO-gp1) before several rounds of 2D classification

to progressively remove junk particles and separate $T = 3$ and $T = 4$ VLPs. To reduce

computational load, the first 2D classification was performed with CTFs ignored until the first

peak, and maximum signal limited to 200. $T = 3$ and $T = 4$ particles were separately re-extracted

without down-sampling for independent 3D refinement (with I1 symmetry imposed) of each VLP

configuration, based on initial models generated *de novo* in Relion. For each reconstruction, map

quality was improved with iterations of CTF refinement and Bayesian polishing. Final

refinements were performed with a solvent-excluding mask and flattened Fourier shell

correlation (FSC) calculations. This was followed by mask-based sharpening, determination of

nominal resolution using the gold-standard FSC criterion (FSC = 0.143), and calculation of local

resolution for each map.

Focussed classification was performed using 2× down-sampled data to limit the computational

load, following a previously described protocol [29]. Briefly, a cylindrical mask was generated

in SPIDER [46] and placed manually over a single Affimer and the space distal to it, expected to

contain density for SUMO-gp1, using UCSF Chimera [47]. Relion was used to apply a soft edge

to the mask, and to generate symmetry-expanded stacks of particles based on the orientational

information generated during separate symmetrised 3D refinements for $T = 3$ and $T = 4$ VLPs.

Symmetry expanded particle stacks were subjected to 3D classification without alignments, with

the focussed mask applied. Asymmetric reconstructions were performed using Relion following

focussed classification.

Acknowledgments

We thank other members of the Stonehouse/Rowlands group at the University of Leeds for their insightful contributions. We also thank Professor Tom Bowden and Dr Guido Paesen, University of Oxford who provided SUMO-Junín gp1.

Funding

This work was supported by a Wellcome PhD studentship to J.S.S. (102174/B/13/Z). Electron microscopy was performed in the Astbury Biostructure Laboratory, which was funded by the University of Leeds and Wellcome (108466/Z/15/Z).

Conflict of Interest

The authors declare that there is no conflict of interest.

Authors and Contributions

K.F., J.S.S., N.J.S., and D.J.R. conceived and designed the experiments. K.F. and L.S. generated the N-VelcroVax sequence. K.F. introduced this into ClearColi BL21 (DE3) cells, generated material for the characterisation of N-VelcroVax and performed serological characterisation. J.S.S. performed negative stain EM and cryo-EM. J.S.S. and A.W. processed and analysed cryo-EM data. N.A.R., D.J.R. and N.J.S. provided supervision. K.F., J.S.S., L.S., D.J.R., & N.J.S. wrote and edited the manuscript. Funding was secured for this research by N.A.R and N.J.S.

References

1. Amanna, I. and M. Slifka, *Vaccination Strategies against Highly Variable Pathogens.* Current Topics in Microbiology and Immunology. Springer International Publishing, 2018: p. 1-30.
2. Delany, I., R. Rappuoli, and E. De Gregorio, *Vaccines for the 21st century.* EMBO molecular medicine, 2014. **6**(6): p. 708-720.
3. Nandy, A. and S.C. Basak, *Bioinformatics in design of antiviral vaccines.* Encyclopedia of Biomedical Engineering, 2019: p. 280.
4. Krugman, S. and M. Davidson, *Hepatitis B vaccine: prospects for duration of immunity.* The Yale journal of biology and medicine, 1987. **60**(4): p. 333.
5. Brenzel, L., et al., *Vaccine-preventable diseases.* Disease control priorities in developing countries, 2006. **2**: p. 389-412.
6. Gu, Y., et al., *Characterization of an Escherichia coli-derived human papillomavirus type 16 and 18 bivalent vaccine.* Vaccine, 2017. **35**(35): p. 4637-4645.
7. Laurens, M.B., *RTS, S/AS01 vaccine (Mosquirix™): an overview.* Human vaccines & immunotherapeutics, 2020. **16**(3): p. 480-489.
8. Nguyen, B. and N.H. Tolia, *Protein-based antigen presentation platforms for nanoparticle vaccines.* npj Vaccines, 2021. **6**(1): p. 70.
9. Tariq, H., et al., *Virus-like particles: Revolutionary platforms for developing vaccines against emerging infectious diseases.* Frontiers in microbiology, 2022. **12**: p. 790121.
10. Crisci, E., J. Bárcena, and M. Montoya, *Virus-like particles: the new frontier of vaccines for animal viral infections.* Veterinary immunology and immunopathology, 2012. **148**(3-4): p. 211-225.
11. Nooraei, S., et al., *Virus-like particles: preparation, immunogenicity and their roles as nanovaccines and drug nanocarriers.* Journal of nanobiotechnology, 2021. **19**(1): p. 1-27.
12. Ulrich, R., et al., *Core particles of hepatitis B virus as carrier for foreign epitopes.* Advances in virus research, 1998. **50**: p. 141-182.
13. Huang, X., et al., *Escherichia coli-derived virus-like particles in vaccine development.* npj Vaccines, 2017. **2**(1): p. 1-9.
14. Gregson, A.L., et al., *Phase I trial of an alhydrogel adjuvanted hepatitis B core virus-like particle containing epitopes of Plasmodium falciparum circumsporozoite protein.* PloS one, 2008. **3**(2): p. e1556.
15. Kutscher, S., et al., *Design of therapeutic vaccines: hepatitis B as an example.* Microbial biotechnology, 2012. **5**(2): p. 270-282.
16. Kingston, N.J., et al., *VelcroVax: a "Bolt-On" Vaccine Platform for Glycoprotein Display.* Msphere, 2023. **8**(1): p. e00568-22.
17. Kushnir, N., S.J. Streatfield, and V. Yusibov, *Virus-like particles as a highly efficient vaccine platform: diversity of targets and production systems and advances in clinical development.* Vaccine, 2012. **31**(1): p. 58-83.
18. Pumpens, P. and E. Grens, *HBV core particles as a carrier for B cell/T cell epitopes.* Intervirology, 2001. **44**(2-3): p. 98-114.
19. Whitacre, D.C., B.O. Lee, and D.R. Milich, *Use of hepadnavirus core proteins as vaccine platforms.* Expert review of vaccines, 2009. **8**(11): p. 1565-1573.
20. Pang, E.L., et al., *Epitope presentation of dengue viral envelope glycoprotein domain III on hepatitis B core protein virus-like particles produced in Nicotiana benthamiana.* Frontiers in Plant Science, 2019. **10**: p. 455.
21. Ravin, N., et al., *Plant-produced recombinant influenza vaccine based on virus-like HBc particles carrying an extracellular domain of M2 protein.* Biochemistry (Moscow), 2012. **77**(1): p. 33-40.
22. Petrovskis, I., et al., *Production of the HBc Protein from Different HBV Genotypes in E. coli. Use of Reassociated HBc VLPs for Packaging of ss-and dsRNA.* Microorganisms, 2021. **9**(2): p. 283.

23. Peyret, H., et al., *Tandem fusion of hepatitis B core antigen allows assembly of virus-like*
particles in bacteria and plants with enhanced capacity to accommodate foreign proteins. PloS
one, 2015. **10**(4): p. e0120751.
24. Stephen, S.L., et al., *Recombinant expression of tandem-HBc virus-like particles (VLPs)*, in *Virus-*
Derived Nanoparticles for Advanced Technologies. 2018, Springer. p. 97-123.
25. Wang, W., et al., *Major immunodominant region of hepatitis B virus core antigen as a delivery*
vector to improve the immunogenicity of the fusion antigen ROP2-SAG1 multiepitope from
Toxoplasma gondii in Mice. Viral immunology, 2017. **30**(7): p. 508-515.
26. Shepherd, D.A., et al., *Using ion mobility spectrometry–mass spectrometry to decipher the*
conformational and assembly characteristics of the hepatitis B capsid protein. Biophysical
journal, 2013. **105**(5): p. 1258-1267.
27. Karpenko, L., et al., *Insertion of foreign epitopes in HBcAg: how to make the chimeric particle*
assemble. Amino acids, 2000. **18**: p. 329-337.
28. Zlotnick, A., et al., *Localization of the C terminus of the assembly domain of hepatitis B virus*
capsid protein: implications for morphogenesis and organization of encapsidated RNA.
Proceedings of the National Academy of Sciences, 1997. **94**(18): p. 9556-9561.
29. Scheres, S.H., *Processing of structurally heterogeneous cryo-EM data in RELION.* Methods in
enzymology, 2016. **579**: p. 125-157.
30. Conley, M.J., et al., *Calicivirus VP2 forms a portal-like assembly following receptor*
engagement. Nature, 2019. **565**(7739): p. 377-381.
31. Snowden, J.S., et al., *Dynamics in the murine norovirus capsid revealed by high-resolution cryo-*
EM. PLoS Biology, 2020. **18**(3): p. e3000649.
32. Snowden, J.S., et al., *Structural insight into Pichia pastoris fatty acid synthase.* Scientific
Reports, 2021. **11**(1): p. 9773.
33. Naskalska, A. and K. Pyrć, *Virus like particles as immunogens and universal nanocarriers.* Polish
journal of microbiology, 2015. **64**(1).
34. Zhu, J., *Mammalian cell protein expression for biopharmaceutical production.* Biotechnology
advances, 2012. **30**(5): p. 1158-1170.
35. Schwarz, H., et al., *Residual endotoxin contaminations in recombinant proteins are sufficient*
to activate human CD1c+ dendritic cells. PLoS One, 2014. **9**(12): p. e113840.
36. Kniskem, P.J., et al., *Unusually high-level expression of a foreign gene (hepatitis B virus core*
antigen) in Saccharomyces cerevisiae. Gene, 1986. **46**(1): p. 135-141.
37. Mamat, U., et al., *Detoxifying Escherichia coli for endotoxin-free production of recombinant*
proteins. Microbial cell factories, 2015. **14**(1): p. 1-15.
38. Demain, A.L. and P. Vaishnav, *Production of recombinant proteins by microbes and higher*
organisms. Biotechnology advances, 2009. **27**(3): p. 297-306.
39. Khow, O. and S. Suntrarachun, *Strategies for production of active eukaryotic proteins in*
bacterial expression system. Asian Pacific journal of tropical biomedicine, 2012. **2**(2): p. 159-
162.
40. Brito, L.A. and M. Singh, *Commentary: acceptable levels of endotoxin in vaccine formulations*
during preclinical research. Journal of pharmaceutical sciences, 2011. **100**(1): p. 34-37.
41. Lua, L.H., et al., *Bioengineering virus-like particles as vaccines.* Biotechnology and
bioengineering, 2014. **111**(3): p. 425-440.
42. Heider, S. and C. Metzner, *Quantitative real-time single particle analysis of virions.* Virology,
2014. **462**: p. 199-206.
43. Scheres, S.H., *RELION: implementation of a Bayesian approach to cryo-EM structure*
determination. Journal of structural biology, 2012. **180**(3): p. 519-530.
44. Zivanov, J., et al., *New tools for automated high-resolution cryo-EM structure determination*
in RELION-3. Elife 7. iCP (E). 2018, PA28αβ-iCP-PA28αβ.
45. Zheng, S.Q., et al., *MotionCor2: anisotropic correction of beam-induced motion for improved*
cryo-electron microscopy. Nature methods, 2017. **14**(4): p. 331-332.

46.   Frank, J., et al., *SPIDER and WEB: processing and visualization of images in 3D electron*
microscopy and related fields. Journal of structural biology, 1996. **116**(1): p. 190-199.
47.   Pettersen, E.F., et al., *UCSF Chimera—a visualization system for exploratory research and*
analysis. Journal of computational chemistry, 2004. **25**(13): p. 1605-1612.
48.   Wynne, S., R. Crowther, and A. Leslie, *The crystal structure of the human hepatitis B virus*
capsid. Molecular cell, 1999. **3**(6): p. 771-780.
49.   Tiede, C., et al., *Affimer proteins are versatile and renewable affinity reagents.* Elife, 2017. **6**:
p. e24903.

Figures

Figure 1: Characterisation of N-VelcroVax. (A) X-ray crystal structure of HBc monomer indicating the N-terminus, C-terminus and major immunodominant region (MIR) (PDB: 1QGT; ref. [48]). **(B)** wt HBc 190 construct (contains no anti-SUMO-Affimer) and N-VelcroVax, indicating the N-terminal insertion of the anti-SUMO-Affimer. **(C)** X-ray crystal structure of an Affimer selected against human SUMO protein (PDB: 5ELJ). Loop 1 and Loop 2 are the variable regions [49]. **(D)** Small-scale expression of HBc VLPs: Western Blot of wt HBc 190 and N-VelcroVax expressed in ClearColi BL21 (DE3) *E. coli* cells detected with mouse monoclonal anti-HBc (10E11). The figure is a representative example of three separate experiments.

Figure 2. Purification and characterisation of N-VelcroVax particles. **(A)** Coomassie blue staining (upper panel) and Western blot (lower panel) of gradient purified N-VelcroVax particles, expressed in ClearColi BL21 (DE3) *E. coli* cells detected with mouse monoclonal anti-HBc (10E11). **(B)** Negative stain TEM analysis of N-VelcroVax. The VLPs were stained using 2% uranyl acetate. The scale bar shows 200 nm. The figure is a representative example of three separate experiments.

Figure 3. Evaluating the binding of N-VelcroVax to SUMO-Junín gp1. N-VelcroVax and
SUMO-Junín gp1 were mixed and incubated overnight before separation on a sucrose density
gradient. Gradient fractions were analysed by Western blot using either mouse monoclonal
anti-HBc (10E11, A) or anti-Junín gp1 (NR 2567, B). The figure is a representative example
of three separate experiments.

Figure 4. Evaluation of the interaction between N-VelcroVax and SUMO-Junín gp1 by ELISA. N-VelcroVax and SUMO-Junín gp1 were mixed in molecular ratios ranging from 1:1 to 1:5 and binding determined by ELISA. Anti-HBc 10E11 (1:1000) and anti-Junín gp1 NR 2567 (1:32,000) were used to detect HBc VLPs and SUMO-Junín gp1 respectively. Anti-mouse antibody conjugated with HRP was added as secondary antibody. TMB chromogenic substrate was used to detect HRP. The optical density at 450 nm (OD 450 nm) is represented in arbitrary units ($n = 3$).

Figure 5. Characterisation of N-VelcroVax VLPs by cryo-EM. Density maps for $T = 4$ (left) and $T = 3$ (right) N-VelcroVax reconstructions, filtered according to local resolution. Each density map is shown as a complete isosurface representation and central cross-section at both low (approx. 2σ, Affimer densities visible) and high (approx. 4σ, Affimer densities not visible) contour level. Cross-sectional views show density, potentially nucleic acid bound non-specifically to the interior surface of the VLPs. Maps are coloured by radial distance (Å).

Figure 6. Structural characterisation of N-VelcroVax in complex with SUMO-gp1. (A) Representative micrographs from negative stain TEM analysis of N-VelcroVax alone (left), SUMO-gp1 alone (centre), or N-VelcroVax:SUMO-gp1 following on-grid interaction (right). Scale bars show 200 nm. **(B)** Isosurface representations of cryo-EM density maps for the $T = 4$ configuration of unliganded N-VelcroVax (left) and N-VelcroVax:SUMO-gp1 (right), shown at ~2 σ and coloured radially. Affimers show a subtle change in relative positioning, as indicated by the white arrow. **(C)** Asymmetric reconstruction of particles from a single focussed class following focussed classification of N-VelcroVax:SUMO-gp1 $T = 4$ particles, showing additional low-resolution density corresponding to bound SUMO-gp1. All focussed classes are shown in Fig. S3. **(D)** Density from an asymmetric reconstruction of $T = 4$ N-VelcroVax:SUMO-gp1 following focussed classification (purple) is suggestive of a reorientation of the Affimer compared to its position in the unliganded N-VelcroVax density map (grey). Fitted atomic models (orange – Affimers, green – remainder of N-VelcroVax) are shown for illustrative purposes.

Figure S1. Processing of N-VelcroVax cryo-EM dataset. (A) Representative micrograph from N-VelcroVax data collection. Scale bar shows 100 nm. (B) 2D classes containing the most particles following automated particle picking. Data were two-fold down-sampled prior to classification. (C-E) Density map for T = 3 VLP of N-VelcroVax filtered according to local resolution, including (C) central section (D) whole VLP and (E) enlarged views of the asymmetric unit. (F) FSC plot for T = 3 N-VelcroVax reconstruction. (G-I) Density map for T = 4 VLP of N-VelcroVax. (J) FSC plot for T = 4 N-VelcroVax reconstruction. All reconstructions shown at ~4 σ and coloured according to radial distance or local resolution, as indicated.

Figure S2. Processing of N-VelcroVax:SUMO-gp1 cryo-EM dataset. (A) Representative
micrograph from N-VelcroVax:SUMO-gp1 data collection. Scale bar shows 100 nm. **(B)** 2D
classes containing the most particles following automated particle picking. Data was five-fold
down-sampled prior to classification. **(C-E)** Density map for $T = 3$ VLP filtered according to
local resolution, including **(C)** central section **(D)** whole VLP and **(E)** enlarged views of the
asymmetric unit. **(F)** FSC plot for $T = 3$ N-VelcroVax:SUMO-gp1 reconstruction. **(G-I)**
Density map for $T = 4$ VLP. **(J)** FSC plot for $T = 4$ N-VelcroVax:SUMO-gp1 reconstruction.
All reconstructions shown at ~3 σ and coloured according to radial distance or local resolution,
as indicated.

Figure S3. Focussed classification of N-VelcroVax:SUMO-gp1 cryo-EM dataset. (A, B) Density observed in all 10 focussed classes from focussed classification of **(A)** $T = 3$ N-VelcroVax:SUMO-gp1 and **(B)** $T = 4$ N-VelcroVax:SUMO-gp1. The number above each class indicates the proportion of sub-particles that were assigned. The position of the mask (grey) is shown for reference. Classes are shown oriented from the viewpoint indicated by the white arrows. **(C-E)** Asymmetric reconstructions using particles contained in the focussed classes indicated by dashed boxes in (A) and (B), filtered by local resolution.

    Table S1: N-VelcroVax and N-VelcroVax:SUMO-Junín gp1 cryo-EM data collection and

528   **processing parameters.**

Sample	N-VelcroVax		N-VelcroVax:SUMO-gp1	
Microscope	FEI Titan Krios		FEI Titan Krios	
Detector mode	Linear		Linear	
Camera	Falcon III		Falcon IV	
Voltage (kV)	300		300	
Pixel size (Å)	1.065		1.065	
Nominal magnification	75,000×		75,000×	
Exposure time (s)	1.0		1.3	
Total dose (e$^-$/Å^2)	43		54.7	
Number of fractions	30		40	
Defocus range (μm)	−0.5 to −2.9		−0.5 to −2.9	
Number of micrographs	12,797		23,966	
Acquisition software	Thermo Scientific EPU		Thermo Scientific EPU	
	T = 3	**T = 4**	**T = 3**	**T = 4**
EMDB ID	EMD-50832	EMD-50833	EMD-50834	EMD-50835
PDB ID	PDB-9FWE	PDB-9FWF	N/A	N/A
Number of particles contributing to map	40,254	56,416	298,802	57,537
Map resolution (FSC = 0.143) (Å)	3.5	3.3	3.0	3.5
Map resolution range around atom positions (Å)	3.2 − 4.2	3.1 − 5.4	2.8 − 3.6	3.3 − 6.0
Map sharpening B factor (Å^2)	-187	-179	-203	-223