Title

Impact of Maximal Overexpression of a Non-toxic Protein on Yeast Cell Physiology

Authors

Yuri Fujita[1], Shotaro Namba[1], Yoshiaki Kamada[2,3], Hisao Moriya[4]

Affiliations

[1]Graduate School of Environmental, Life, Natural Science and Technology, Okayama University

[2]National Institute for Basic Biology, Okazaki, Japan

[3]Graduate University for Advanced Studies (SOKENDAI), Hayama, Kanagawa, Japan

[4]Faculty of Environmental, Life, Natural Science and Technology, Okayama University

Competing interests

The authors declare that there are no competing interests.

Author contribution

YF and HM conceived and designed the experiments, YF and YK performed experiments, YF, SN, and HM analyzed the data, and YF and HM wrote the paper.

Summary

While it is recognized that excess expression of non-essential proteins burdens cell growth, the physiological state of cells under such stress is largely unknown. This is because it is challenging to distinguish between adverse effects arising from the properties of the expressed excess protein (cytotoxicity) and those caused solely by protein overexpression. In this study, we attempted to identify the model protein with the lowest cytotoxicity in yeast cells by introducing a new neutrality index. We found that a non-fluorescent fluorescent protein (mox-YG) and an inactive glycolytic enzyme (Gpm1-CCmut) showed the lowest cytotoxicity. These proteins can be expressed at levels exceeding 40% of total protein while maintaining yeast growth. The transcriptome of cells expressing mox-YG to the limit indicated that the cells were in a nitrogen source requirement state. Proteome analysis revealed increased mitochondrial protein and decreased ribosome abundance, similar to the inactivated state of the TORC1 pathway. The decrease in ribosome abundance was presumably due to defective nucleolus formation, partially rescued by a mutation in the nuclear exosome. These findings suggest that massive overexpression of excess protein, termed protein burden, causes nitrogen source starvation, a metabolic shift toward more energy-efficient respiration, and a ribosomal biosynthesis defect due to an imbalance between ribosomal protein and rRNA synthesis in the nucleolus.

Introduction

Expression levels of proteins are optimized to maximize the fitness of organisms, and the evolutionary principle that determines those levels is considered to be demand and constraint (Bruggeman et al., 2020) (Figure 1A). Demand is the evolutionary pressure that works to increase the expression of a protein to meet the requirements of its function, and constraint is the evolutionary pressure that works to decrease the expression of a protein to avoid the negative effects produced by excess protein. Constraints appear as disturbances in cellular function when the protein is overexpressed (Figure 1A, the red arrow). The mechanisms of constraining expression levels, or defects due to overexpression, can be roughly classified into four major categories (Moriya, 2015) (Figure 1B). "Resource overload" (also called "burden") (Kastberg et al., 2022) is a disorder in which intracellular resources used for protein processing, such as synthesis, localization, degradation, and modification of proteins, are depleted or monopolized by the processing of that protein, resulting in a disruption of processing of other proteins (Dong et al., 1995; Eguchi et al., 2018; Kintaka et al., 2020, 2016; Stoebel et al., 2008). This phenomenon occurs in proteins with high expression levels. "Stoichiometry imbalance" is a disorder triggered by an imbalance of amounts between constituents within complexes, leading to untimely activation or inactivation (pathway modulation), degradation (resource overload), aggregation (promiscuous interaction), etc. (Kaizu et al., 2010; Makanae et al., 2013; Papp et al., 2003). This occurs due to the overexpression of proteins that form complexes. "Pathway modulation" is a disorder in which pathways are untimely activated or inactivated. This occurs especially with overexpression of proteins involved in pathway regulation, such as metabolic enzymes, kinases and phosphatases, and transcription factors (Prelich, 2012; Youn et al., 2017). "Promiscuous interaction" is a disorder in which weak interactions are enhanced by mass action to form complexes or aggregates that are not normally formed. This damages cellular function by inactivating essential proteins or overloading degradation resources, and is caused by overexpression of intrinsically disordered proteins or liquid-liquid phase-separating proteins (Bolognesi et al., 2016; Ma et al., 2010; Vavouri et al., 2009).

Figure 1. Schematic diagrams illustrating the constraints on protein expression levels.

A) Protein expression levels are determined by demand and constraints (created by the authors, inspired

by the work by Keren *et al*. (Keren et al., 2016). The existence of this evolutionary principle is revealed

by the relationship between fitness (growth rate) and expression level when the expression level is

altered, and in general, the native expression level provides the highest level of fitness for the organism

(Bruggeman et al., 2020). The existence of constraints is revealed by a decrease in fitness upon

overexpression (red arrow). Constraints determine the expression limit of the protein, i.e., the

expression level at which growth inhibition occurs (dotted lines), but there can be multiple constraints

for a single protein. The constraint that exists at the highest expression limit is protein burden.

B) Four major mechanisms that constrain protein expression levels. The figure outlines the fate of

proteins and shows what adverse effects occur upon overexpression (red arrows). The resource overload

that occurs within the synthesis process (transcription and translation) is specifically referred to as

protein burden. This diagram is a more concise redrawing of the author's earlier one (Moriya, 2015).

That asterisk means differentially localized proteins.

C) A barrel model for resource overload (modified from the author's earlier one) (Kintaka et al., 2020).

The size of the barrels represents the capacity of each process, and the arrows represent the fate of the

proteins processed there. The expression limit of a protein is defined by the lowest-capacity process

that processes the protein. Among these processes, synthesis, which processes all proteins, is considered

to have the largest capacity. Thus, proteins processed only by synthesis would have the highest

expression limits, and their overexpression would overload synthesis (i.e., cause protein burden).

Previous studies by the authors, conducted in this context, estimated that protein burden occurs at more

than 15% of total protein (Eguchi et al., 2018).

D) Difference in burden due to gene structure (codon optimization). The amount of transcription

required to achieve the same protein expression limit depends on the degree of codon optimization. This

may change whether transcription or translation results in a higher burden.

E) An ideal framework for protein burden studies. See text for details.

As the expression level of a protein increases, it eventually encounters one of the barriers created by

either these specified constraints or others that are unknown, all of which are related to the function and

physicochemical properties of the protein, and damage cellular function. The expression level at which

this occurs is referred to as the "expression limit" in this paper (see Figure 1A). If the constraints could

be removed from the protein, the expression limit would increase. For example, if components of a

complex are simultaneously expressed to resolve a stoichiometry imbalance, the expression limits of

those proteins will be increased (Kaizu et al., 2010; Makanae et al., 2013). In this manner, when (one)

constraint is removed from a protein, the expression limit would increase until the next constraint barrier

is hit (Figure 1A, dotted lines). The amount of expression limit created by a constraint barrier depends

on the type of constraint. Stoichiometry imbalances, pathway modulations, and promiscuous

interactions would create different expression limits for each protein. This is because they are

determined by the activity, regulatory mechanism, and physicochemical properties of each protein. On

the other hand, resource overload is presumed to be determined to some extent by the capacities of the

processes in the cell (Kintaka et al., 2020) (Figure 1C). Essentially, the synthesis, in which all proteins

are processed, should have the largest capacity, while processes such as transport, folding, and

degradation, in which only a fraction of proteins are processed, should have smaller capacities. In fact,

adding transport or degradation signals to a cytosolic protein lowers the expression limit (Kintaka et al.,

2016). In that light, the final constraint barrier that emerges when all possible constraints are removed

is the overloading of the synthesis process.

Overloading of the synthesis process is specifically referred to as "protein burden" or "protein cost"

(Eguchi et al., 2018; Kafri et al., 2016). It has long been the subject of study, mainly as an effect on

growth produced by excess (gratuitous) proteins that are not necessary for cellular function (Bruggeman

et al., 2020; Scott et al., 2010). This idea was first proposed in studies of the cost of unnecessary

induction of the *lac* operon in *Escherichia coli* (Koch, 1983). In recent years, much attention has been

paid to the actual mechanisms by which massive expression of excess protein inhibits growth and the

response of cells (Farkas et al., 2018; Kafri et al., 2016; Metzl-Raz et al., 2020). During protein burden,

it is presumed that depletion of the materials, machinery, and factors necessary for transcription and

translation occurs, along with competition from these monopolies. In particular, amino acids and

aminoacyl-tRNAs are major candidates for depletion (Kastberg et al., 2022). In addition, ribosomes,

which are crucial for protein synthesis, themselves require significant resources for their synthesis and have therefore been considered major candidates for depletion during protein burden, as they compete for these limited resources (Metzl-Raz et al., 2017). Mechanistic analysis of protein burden in budding yeast, a model eukaryotic cell, has been intensively conducted by Barkai and colleagues, who reported that transcription is rate-limiting in phosphate starvation and translation in nitrogen source starvation (Kafri et al., 2016), and that basic transcription factors are limiting resources (Metzl-Raz et al., 2020). In addition, a cellular response in the form of increased cell size and enhanced protein synthesis capacity has been observed (Kafri et al., 2016). Thus, the previously expected decrease in ribosomes has not been detected.

As mentioned above, to investigate protein burden, we need to overexpress proteins with minimal constraints. This is because overexpressing proteins with additional constraints cannot overload the synthetic process, as the additional constraints reduce the expression limits (Figure 1A). Also, the cellular damage and cellular response generated by the constraint mask the protein burden response. However, it remains unclear which proteins are not subject to constraints other than protein burden. Proteins that would induce protein burden have been determined by assuming that they would be minimally constrained based on their properties, and this assumption has been confirmed experimentally. Model proteins selected in this manner in yeast include fluorescent proteins (FPs) and certain glycolytic enzymes (Eguchi et al., 2018; Farkas et al., 2018; Kafri et al., 2016; Kintaka et al., 2020, 2016). FPs do not have a specific function or interaction partner in yeast cells and are not actively sorted into cellular compartments other than the cytosol. On the other hand, glycolytic enzymes are naturally expressed at very high levels and are thus considered to be less constrained. Notably, these proteins exhibit an expression limit of approximately 15% of total protein in yeast cells (Eguchi et al., 2018). However, it is not clear whether these proteins are free from constraints other than protein burden. This is because there may be unknown constraints, and there is no evidence that the measured expression limit is the "true limit" at which protein burden occurs.

In addition to the protein (amino acid sequence) itself, there is another aspect to be considered when

selecting a model for protein burden. That is the nucleotide sequence of the gene used for expression,

i.e., transcriptional, and translational regulatory sequences, codon usage, and so on (Figure 1D). For

example, if a protein is to be expressed to the limit by a less codon-optimized and thus less

translationally efficient mRNA, a large amount of mRNA would be required, which would induce a

greater burden on transcription. Conversely, a similar attempt by an mRNA with high translation

efficiency would induce a greater burden on translation. These are the constraints inherent in protein

synthesis itself. Although there is some evidence that protein synthesis in nature is economically

designed to optimally load transcription and translation (Frumkin et al., 2017; Hausser et al., 2019;

Mahima and Sharma, 2023), it still needs to be determined what the genetic design should be to

minimize the burden.

Considering the above circumstance, a realistic approach to protein burden requires more than just

drawing straightforward conclusions from the overexpression of a single model protein (or gene).

Rather, it requires experimentally identifying proteins with the highest possible expression limits,

investigating the negative impacts of their overexpression, and, if any constraints are revealed,

examining whether their removal can increase expression limits. Through successive iterations, this

process gradually deepens our understanding of protein burden (Figure 1E). In each iteration, the least

constrained protein (and its gene) will be generated and utilized as a benchmark for further investigation

into proteins with high (or low) expression limits and the adverse effects of their overexpression. This

will (unintentionally) elucidate a new mechanism of constraint. Indeed, the enhanced green fluorescent

protein (EGFP), which we have used as a model for protein burden (Kintaka et al., 2020, 2016; Makanae

et al., 2013), was found to form a chaperone-entrapped aggregate upon overexpression and trigger the

heat shock response due to its low folding ability (Namba et al., 2022). Its overexpression also causes

abnormal yeast cell morphology through proteasomal stress as it contains cysteines. The moxGFP

(mox), which improved folding properties and lacks cysteine residues (Costantini et al., 2015), showed

a higher expression limit and no longer induced abnormal morphologies. However, even when

overexpressing mox, the anticipated depletion and competition for factors, or cellular responses to them

have not been observed (Namba et al., 2022).

In this study, we found that a mutation causing loss of fluorescence in mox, and other fluorescent

proteins increased the expression limit up to 3-fold, to more than 40% of total protein. Introducing a

neutrality index (NI) to assess protein constraint, we found that non-fluorescent mox (mox-YG) and

enzymatically inactive Gpm1 (Gpm1-CCmut) are currently the least constrained. Upon overexpression

of mox-YG, depletion of amino acid or nitrogen sources, decreased ribosomal expression, and a

metabolic shift from glycolysis to oxygen respiration were observed. Thus, physiological states, as

anticipated with such protein burdens, are only generated by massive overexpression of proteins with

significantly lower constraints. Some of the responses found appear to be due to inactivation of the

TORC1 pathway, suggesting that eukaryotic cells use this regulatory pathway to buffer protein burden.

In the burdened cells, the nucleolus was dysplastic, and mutations in the nuclear exosome partially

restored this. Thus, the abnormal formation of liquid-liquid phase-separated organelles due to dilution

may be one of the defects caused by the strong protein burden.

Results

Fluorescent property is a constraint on the expression of fluorescent proteins.

To evaluate the expression limits of FPs, we overexpressed FPs to levels that induced growth inhibition. Gene constructs, in which the *TDH3* promoter controls the target protein genes, were incorporated into plasmids designed for the genetic-tug-of-war (gTOW) method (Figures 2A and Figure 2—figure supplement 1A) (Moriya et al., 2006). Subsequently, yeast cells were transformed and cultured in a medium depleted of leucine. The maximum growth rate (MGR) of cells was calculated from serial measurements of optical density at 660 nm (OD_{660}). Protein levels were determined through the analysis of proteins from cells in the logarithmic growth phase (Figure 2—figure supplement 1B). For each protein, we also analyzed a mutant in which the Tyr residue forming the fluorophore was replaced with Gly (YG mutation), resulting in a loss of fluorescence. This is because we unexpectedly found that this mutation substantially increased the expression limits of FPs (see below).

Overexpression of the FPs and mutants all caused growth inhibition (Figure 2B, C). EGFP and mox were expressed at about 15% of the total protein as previously reported (Kintaka et al., 2016), but mCherry (mChe) and YG mutants were expressed at higher levels. Mox-YG expressed 44% of the total protein (Figures 2D, E). Our previous work had evaluated only expression limits (Eguchi et al., 2018), but we realized that we could not properly assess the constraints on each protein if the growth rates at the expression limit were different. For example, the mChe's expression limit (20%) is higher than that of sfGFP (17%), but the relative MGRs are 20% and 80%, respectively, which is much lower for mChe. The relationship between %MGR and %protein level when the control is set at 100% is shown in the scatter plot in Figure 2F. The lower left indicates more constraint as lower expression levels lead to growth inhibition, and the upper right indicates less constraint as higher expression levels do not cause growth inhibition. Here, we describe "unconstrained" as "neutral" and calculate the neutrality index (NI) as the product of %MGR and %protein level.

Figure 2. Evaluation of expression constraint (or neutrality) of fluorescent proteins.

A) Experimental setup of the analysis. Target proteins were expressed under the control of the *TDH3*

promoter (*TDH3$_{pro}$*) using the multicopy plasmid pTOW40836. The cells were pre-cultured under –Ura

conditions, and then cultured in –LeuUra conditions.

B) Growth curves of the vector control and cells overexpressing FPs and their mutants in synthetic

medium (–LeuUra). The solid or dotted lines show the average calculated from three biological

replicates. Growth curves with the standard deviations (SD) of replicates are shown in Figure 2—figure

supplement 2.

C) MGR of cells overexpressing FPs and their mutants (percent over the vector control).

D) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing FPs. All proteins

were separated after staining with a fluorescent dye.

E) Protein level of the target protein (percent over the total protein). The amount was calculated from

the intensity of the target bands separated by SDS-PAGE of **D**. See Figure 2—figure supplement 1B

for the method of FP and total protein quantification. Note that EGFP-YG and mCherry may not be

accurately measured due to dimerization and cleavage of the protein, respectively.

F) Relationship between MGR and the protein level. MGR data are the same as in **C**, and the protein

level data are the same as in **E**. The neutrality indexes (NIs), calculated from the products of MGR and

protein levels, are indicated by the lines. If the investigated proteins have the same NI, they are located

on the same dotted line segment.

G) The neutrality index for the FPs and mutants. The neutrality index is the product of %MGR (/Vector)

and %protein level (/total protein).

The bars and error bars in **C**, **E**, and **G** show the means and SDs calculated from three biological

replicates. The raw data is shown with dot plots.

Note that the regime of NI in Figure 2F shows a theoretical neutrality distribution but does not show

the relationship between growth rate and the expression level when the expression level of a protein

with a particular neutrality is increased or decreased. For example, 20% of the growth rate would be

maintained when the expression of mox-YG (NI = 2,400) is increased to 100%, but this is not realistic.

We also note that in this study, we calculated NI using the percentage of the target protein among total

cellular proteins separated by SDS-PAGE. Therefore, to calculate NI, the expression limit of the target

protein must be sufficiently high to be detectable by this method. On the other hand, if detection

sensitivity is improved, for example by using Western blotting, it becomes possible to estimate the NI

even for proteins under strong constraints—i.e., those that exhibit cytotoxicity. In fact, when we

recalculated the NI for EGFPs fused with various localization signals from our previous study (Kintaka

et al. 2016), all values were found to be below 100 (Figure 2−figure supplement 3).

The NI of each protein indicates that introducing a non-fluorescent mutation in FPs uniformly increases

neutrality (Figure 2G), suggesting that "having fluorescence" is a constraint on the expression of FPs.

The most neutral proteins were sfGFP-YG and mox-YG, with NIs of about 2,400. Notably, despite

significant differences in the degree of growth inhibition and expression limits between the two proteins,

their NIs are nearly identical (Figures 2F, G). They differ only in two amino acid residues (Cys - Ser)

(Figure 2—figure supplement 4), which seem to generate differences in expression efficiency. However,

this does not appear to affect their NIs.

We also investigated the impact of expression efficiency on NI with another FP, mChe. The mChe used

in this study is codon optimized (the codon adaptation index (CAI) = 0.584) (Keppler-Ross et al., 2008),

while the mChe used by Kafri et al. for protein cost analysis (here referred to as mChe-Kafri) is not

codon optimized (CAI = 0.236) (Kafri et al., 2016). When overexpressed, mChe-Kafri (and mChe-

Kafri-YG) had lower expression and growth inhibition than mChe (and mChe-YG) (Figures 2—figure

supplement 5A-D). This may be due to the lower translation rate of mChe-Kafri mRNA. However, their

NIs were not significantly different, especially for the YG mutation (Figures 2—figure supplement 5E,

F). Thus, the constraints indicated by NI are apparently not affected by translation efficiency. In the

study using this mCherry-Kafri, it was also reported that the more mChe-Kafri was expressed, the more

the cells enlarged (Kafri et al., 2016). However, no cell enlargement was observed with mox-YG, which

can be expressed more than mCherry. Interestingly, the mChe-YG cells did not show enlargement

(Figures 2—figure supplement 6A, B), indicating that cell enlargement is a phenomenon that occurs

specifically when fluorescent mChe is overexpressed.

It should be noted that the growth inhibition caused by the overexpression of mox-YG in this study is a

burden on translation rather than on transcription. This is because overexpression of a frameshift mutant

(moxFS), in which four nucleotides were inserted directly after the ATG of the mox-YG gene, abolished

the notable growth delay (Figures 2—figure supplement 6C, D).

Constraint on fluorescent protein expression arises from fluorophore properties.

Next, we further analyzed the possibility that fluorescence constrains the expression of FPs. When yeast

was grown in the dark, the degree of growth inhibition of mox was the same (Figure 2—figure

supplement 7A) and the expression limit of mox was rather increased (Figure 2—figure supplement

7B). Phototoxicity does indeed exist when expressing mox with fluorescence. This is because, when

exposed to strong light, cells overexpressing mox show a reduced growth rate (Figure 2—figure

supplement 7C-E) and abnormalities in their internal structures (Figure 2—figure supplement 7F).

However, the expression limits of mox and mox-YG were almost unchanged by the light (Figure 2—

figure supplement 7G, H). Thus, the constraint was not due to phototoxicity itself.

The number of Tyr residues was also irrelevant because NI was like that of mox-YG when a single Tyr

was added to the C-terminus of mox-YG (Figure 2—figure supplement 8A-E). Reactive oxygen species

(H_2O_2) are generated during fluorophore formation. When H_2O_2 was added to the medium, stronger

growth inhibition occurred only in mox (Figure 2—figure supplement 8F-H). This suggests that the

reactive oxygen species generated during fluorophore formation are responsible for the constraint.

However, this possibility is not supported by the omics analysis described later (see Discussion). Mox-

YG, like mox, was present throughout the cytoplasm and did not cause any characteristic localization

or aggregation (Figure 2—figure supplement 9).

During this study, we examined a moxGFP (mox-T203I) in which Thr203 was accidentally replaced by

Ile (Figure 2—figure supplement 10). The fluorescence of this mutant is reduced to 78% of mox

(Figures 2—figure supplement 10A-C). The growth rate of the cells expressing this mox-T203I was

like that of mox and mox-YG (Figure 2—figure supplement 10E). Still, the expression limit was

intermediate between mox and mox-YG (Figure 2—figure supplement 10F), resulting in the NI

intermediate between mox and mox-YG (Figure 2—figure supplement 10G). This result further

supports that fluorescence is related to constraints. However, as mentioned above, fluorescence

emission itself should not be toxic (Figure 2—figure supplement 7). We next focused on the formation

process of the fluorophore: the tyrosine of the GFP fluorophore undergoes phenol and phenolate state

changes, with Thr203 stabilizing the phenol form and substitution to Ile reducing the phenolate form

(Kummer et al., 2000). Furthermore, in enhanced GFPs such as mox, substitution of Ser65 before Tyr66

for Thr anchors the phenolate form. To investigate the possibility that this is a factor determining the

limit of mox, we produced mox with Thr65 replaced with Ser and examined its expression limit. The

mox-T65S mutant was expressed 12% more than mox (Figures 2—figure supplement 10H, I),

suggesting that the phenol-type fluorophore can be a limiting factor for the expression level of the

fluorescent protein, while the concrete mechanism is still unclear.

Overexpression of mox-YG causes an amino acid starvation response.

To understand the cell physiology under protein burden, we next performed transcriptome (RNA-seq)

analysis of the cells overexpressing mox-YG, the least constrained protein. Although the expression

limit of mox-YG is about 3-fold higher than that of mox (Figure 2E), the growth rates at the expression

limits of both are not largely different (Figure 2C). Thus, by comparing transcriptomes upon their

overexpression, we may be able to distinguish and extract the general transcriptional response due to

reduced growth rates from the (protein burden) response associated with the massive expression of a

less constrained protein.

The expression changes under mox and mox-YG overexpression were calculated as fold change (FC)

over the vector control (Figures 3—figure supplement 1A, B). The overall trend of expression changes

was similar ($r = 0.64$, Figure 3A). This similar variation may include responses associated with the expression system and reduced growth rate. On the other hand, there were also different transcriptional responses between the two. Gene categories that showed significant expression changes in either mox or mox-YG overexpression are shown in Figure 3B. In addition to the enhanced ribosome synthesis and decreased glycolysis observed in previous mox overexpression (Namba et al., 2022), enhanced oxidative phosphorylation was detected. Interestingly, "Ribosome biogenesis", which was elevated in mox overexpression, showed a rather decreasing trend in mox-YG overexpression (Figure 3B). Indeed, *IFH1*, which is involved in ribosomal gene transcription (Martin et al., 2004), was upregulated in mox, but downregulated in mox-YG (Figure 3A). Different trends were also observed for "amino acid metabolism" (Figure 3B). The amino acid transporter genes *AGP1* and *GAP1* were upregulated only in mox-YG overexpression (Figure 3A, C). Expressions of specific amino acid and nitrogen source transporter genes were strongly induced in mox-YG (Figure 3C). The expression of glucose transporter genes was also very different (Figure 3C). These results suggest that mox and mox-YG induce similar but distinct transcriptional responses, particularly that mox-YG overexpression may result in depletion of amino acid and nitrogen sources and decreased ribosome synthesis.

The transcriptional responses in mox-YG overexpression obtained by RNA-seq were confirmed by transcriptional reporter analysis (Figure 3D and Figure 3—figure supplement 2). In this reporter analysis, we also tested overexpression of a frameshift mutant (moxFS) that imposes a burden on transcription but not on translation (Figure 2—figure supplement 6C, D). As a result, most of the responses observed under mox-YG overexpression were not observed under moxFS overexpression, suggesting that these transcriptional responses are likely caused by a burden on translation. The only exception was *TDH3pro*, whose activity decreased even under moxFS overexpression. Since *TDH3pro* is used for driving overexpression in this system, this result suggests that transcriptional competition may occur as a result of overexpression. Notably, in our gTOW experiments, leucine depletion is employed to induce overexpression; however, we consider it unlikely that leucine depletion *per se* is responsible for the induction of *GAP1*, because *GAP1* expression remained unchanged in the vector control even under gradual leucine depletion (Figure 3—figure supplement 3).

Figure 3. Transcriptional response of mox and mox-YG overexpression.

A) Comparison of transcriptional responses of mox and mox-YG overexpression. Genes that showed common or different transcriptional responses in mox and mox-YG overexpression are shown in

indicated colors. Genes that showed characteristic responses are also shown. FDR: false discovery rate.

r: Pearson correlation coefficient.

B) Gene groups that showed significant transcriptional changes. Among the KEGG orthology level 3

categories, gene groups in the category that were significantly up-regulated or down-regulated (FDR <

0.05) in mox or mox-YG overexpression are shown in the violin plots. Genes with significant expression

changes (FDR < 0.05) within the same category are shown by swarm plots. Asterisks indicate significant

differences (FDR < 0.05) between mox and mox-YG comparisons in gene groups belonging to the same

category. Comparisons for all categories are shown in Figure 3—figure supplement 1C.

C) Expression changes in representative amino acid, ammonium, and glucose transporters.

D) Verification of promoter activity by the reporter assay. Constructs used for promoter analysis with

transcription reporters and quantitative results of RFP fluorescence values (arbitrary unit) for each

promoter. "moxFS" means the frameshift mutant of mox-YG is overexpressed. Bars and error bars

indicate the mean and SD of maximum fluorescence values for RFP calculated from four biological

replicates. Raw data are shown as dot plots. The p-values were calculated by performing Welch's t-test

and applying Bonferroni correction. "n.s." means $p > 0.05$. Detailed constructs and time series data for

promoters other than those shown here are in Figure 3—figure supplement 2.

Overexpression of mox-YG causes a proteomic response that partially overlaps with TORC1

inactivation.

Protein burden is an overload on synthetic resources, and thus, when this overload occurs strongly, it is

expected that there would be a reduction in the amount of proteins other than those being overexpressed

due to competition for synthetic resources. Indeed, in cells overexpressing mox-YG, the amounts of

proteins other than mox-YG were reduced (Figure 4A). Interestingly, the total amount of protein per

cell remained almost constant, with the amount of other proteins decreasing as mox-YG was

overexpressed. This trend was generally true for the overexpression of other FP (Figure 4—figure

supplement 1). This suggests that there is a mechanism (or constraint) that maintains the overall cellular

protein levels.

The maintenance of total cellular protein levels under overexpression conditions may be due to

constraints such as intracellular space limitations or limitations in the availability of nutrients in the

medium. To investigate whether nutrient limitation contributes to this phenomenon, we attempted

overexpression of mox and mox-YG in YPD medium, which is considered to be more nutrient-rich than

the SC medium. Since leucine depletion could not be used to drive overexpression in this context, we

employed an alternative strategy using the aureobasidin A resistance marker (*AUR1d*) to increase

plasmid copy number (Figure 4—figure supplement 2A). In this experiment, mox-YG was expressed

at approximately 20% of the total cellular protein, with a corresponding increase in total protein content

per cell (Figure 4—figure supplement 2C, D). This result suggests that in nutrient-rich YPD medium,

cells may possess the capacity to buffer excess protein expression by increasing the total proteome.

However, under these expression conditions, no growth inhibition was observed (Figure 4—figure

supplement 2B), and thus the induction of mox-YG was not sufficiently strong. We therefore consider

it necessary to investigate the regime under conditions of stronger overexpression in future studies.

To examine changes in protein expression, proteomic analysis of mox-YG overexpressing cells (in the

SC medium) was performed. The distribution of intensities obtained from the analysis is shown in the

Proteomaps in Figure 4B. In mox-YG overexpressing cells, mox-YG accounted for 40% of the total

protein expression. Figure 4A suggests that in mox-YG overexpressing cells, proteins other than mox-

YG are overall reduced to about 60% compared to the vector. Therefore, we considered it necessary to

normalize the data overall to accurately represent changes in the per-cell proteome. However, in this

study, we used data without such correction to focus on how the composition of the proteome, other

than mox-YG, changed (see Materials and Methods and Figures 4—figure supplement 3A-C for more

details).

Figure 4C illustrates the categories that showed significant changes upon mox-YG overexpression (all

categories are shown in Figure 4—figure supplement 4). Elevated mitochondrial proteins, such as

oxidative phosphorylation and TCA cycle, and decreased ribosomal protein and ribosomal synthesis

were observed. Interestingly, the proteomic changes observed in mox-YG overexpressing cells showed

little correlation with the corresponding transcriptomic changes (Figure 4—figure supplement 3D). The only notable exception was oxidative phosphorylation, which was upregulated at both the transcript and protein levels (Figure 3B and 4C). In contrast, for example, ribosomal proteins were upregulated at the transcript level but markedly downregulated at the protein level (Figure 3B and 4C). This discrepancy may indicate a cellular state in which transcriptional responses are uncoupled from protein expression as a result of protein burden.

As mentioned above, RNA-seq results suggested nitrogen source starvation in mox-YG overexpression. Therefore, we investigated whether this response is influenced by the regulation through the TORC1 pathway, a primary mechanism responsive to nitrogen sources. By comparing expression changes due to rapamycin (a TORC1 inhibitor) treatment (Gowans et al., 2018) and mox-YG overexpression, we found that about 40% of the proteins with altered expression were congruent with the group of proteins regulated by TORC1 (Figures 4—figure supplement 5A, B). Thus, the proteome expression changes due to mox-YG overexpression may include responses due to inactivation of the TORC1 pathway as well as other responses. Figure 4C shows which categories had significant expression changes when separated into rapamycin-responsive (Rap) and rapamycin-non-responsive (Non-Rap) gene populations. Carbohydrate metabolism, elevated TCA cycle, and decreased ribosome synthesis were suggested to be responses due to TORC1 inactivation, while endocytosis, cytoskeleton, protein transport and lipid synthesis, and glycolysis were suggested to be due to responses other than TORC1 inactivation (Figure 4—figure supplement 6). Oxidative phosphorylation included both Rap responses (12/32) and non-responses (20/32) (Figure 4—figure supplement 6C). Indeed, mox-YG overexpression reduced the sensitivity to rapamycin compared to the vector control (Figure 4—figure supplement 7A-E), supporting the notion that TORC1 activity was diminished. However, we were unable to obtain direct evidence demonstrating TORC1 inactivation. We examined the phosphorylation status of Atg13 (Kamada et al. 2000), a known TORC1 target, by Western blotting. In mox-YG overexpressing cells, however, the total expression level of Atg13 was markedly reduced (Figure 4—figure supplement 7F). Therefore, it remains inconclusive how the decreased expression of TORC1 targets affects downstream signaling under these conditions.

Above transcriptome and proteome analyses suggested that mox-YG overexpression induces

mitochondrial development and a metabolic shift from fermentation to oxidative respiration. To verify

mitochondrial development, we performed fluorescence microscopy. Indeed, mitochondria labeled with

Rip1-mScarlet-I, a component of the oxidative phosphorylation pathway, were significantly more

developed in mox-YG overexpressing cells (Figure 4D, 4—figure supplement 8). Furthermore, oxygen

consumption was increased in these cells (Figure 4E, 4—figure supplement 9), supporting the notion

that they had undergone a metabolic shift toward oxidative respiration.

Figure 4. Overexpression of mox-YG causes a proteome response like that of TORC1 inactivation.

A) Ratio of the total protein level over the vector control. The colored area indicates the amount of overexpressed protein. The amount was calculated from the intensity of the target bands separated by SDS-PAGE. Bars and error bars indicate the mean and SD of protein levels calculated from three biological replicates. The p-values were calculated by performing Welch's t-test and applying Bonferroni correction. "n.s." means $p > 0.05$.

B) Visualization of the proteome of the vector control and mox-YG overexpressing cells using Proteomaps (Liebermeister et al., 2014). Each polygon represents the mass fraction of the protein within the proteome. Similar colors indicate similar or closely related pathways and proteins.

C) Violin plots showing groups of genes in the KEGG orthology level 3 category whose expression was increased or decreased by mox-YG overexpression. Gene groups with significantly altered expression (FDR < 0.05) in mox-YG overexpression are shown in red. Genes with significant expression changes (FDR < 0.05) within the same category are shown by swarm plots. "+" and "–" indicate the categories with significantly increased or decreased expression, respectively, when separated into Rapamycin-responsive (Rap) and Rapamycin-non-responsive (Non-Rap) gene populations. Comparisons for all categories are shown in Figures 4—figure supplement 6A, B. Published data (Gowans et al., 2018) was used for comparison with the rapamycin response genes.

D) Fluorescence microscopy image showing the localization of Rip1-mScarlet-I. The vector control and mox-YG overexpressing cells were observed in the log phase. Stationary phase cells, bright-field images, quantitative results, and statistical analyses are shown in Figure 4—figure supplement 8.

E) Oxygen consumption rate in vector control and mox-YG overexpressing cells (arbitrary units, AU). Antimycin A was added as a control for respiratory inhibition. Measurements were conducted with three biological replicates. Bars and error bars indicate the mean and SD from three biological replicates. Raw data are shown as dot plots. The p-values were calculated by performing Welch's t-test and applying Bonferroni correction. "n.s." indicates $p > 0.05$. The measurement data are shown in Figure 4—figure supplement 9.

Overexpression of mox-YG causes a dysplastic nucleolus, which is alleviated by mutations in the nuclear exosome.

Ribosomes are a crucial factor for protein burden because they are themselves protein synthesizers

while being synthesized in the highest amounts in the cell. Therefore, we further focused on the behavior

of ribosomes in mox-YG cells and expression changes were investigated in detail for each ribosomal

protein (Figure 5—figure supplement 1). While the majority of the 84 ribosomal proteins detected (45

of them significantly) decreased, six (Rps12, Rps31, Ppp0, Rpp2a, Rpl5, and Rpl12) showed significant

increases. Interestingly, all decreased ribosomal proteins are assembled into ribosomes in the nucleolus,

while all increased ribosomal proteins are assembled into ribosomes in the cytoplasm (Lafontaine,

2015). They constitute the "beak" of the 40S subunit and the "P-stalk" and the "central protuberance

(pc)" of the 60S subunit (Figure 5A). Electron micrographs showed an abnormal nucleolar formation

of mox-YG overexpressing cells (Figures 5B and Figure 5—figure supplement 3), in addition to reduced

ribosome density in the cytoplasm (Figure 5—figure supplement 2A). This abnormal nucleolar

formation was also confirmed by fluorescence microscopy images of Nsr1 (Nucleolin), the major

protein of the nucleolus (Figure 5—figure supplement 2B-D).

Based on these results, we hypothesized that overexpression of mox-YG creates an imbalance between

the amount of ribosomal protein synthesized in the cytoplasm and the amount of rRNA synthesized in

the nucleolus (Figure 5C, D). The imbalance may lead to the creation of misassembled ribosomes,

which are eventually degraded. We speculated that the primary enzyme degrading the misassembled

ribosome should be the nuclear exosome. This is because our previous genome-wide mutation analysis

identified exosome mutants (*dis3-1, mtr3-ts, mtr4-1, rpn42-ph, rpn45-ph, and ski6-ph*) as primary

mitigators of protein burden (Kintaka et al., 2020). We confirmed that the *mtr4-1* mutant does not show

growth retardation upon mox-YG overexpression (Figure 5E). In fact, in *mtr4-1* cells, there was no

difference in growth rate between the vector control and mox-YG overexpression (Figure 5—figure

supplement 5B), but rather a shorter lag time for the mox-YG overexpression than the vector control

(Figure 5E). We next quantified protein upon mox-YG overexpression in *mtr4-1* cells (Figure 5F).

Compared to the wild-type (Vector), the total protein content in *mtr4-1* (Vector) cells was reduced to

78%, which was likely related to the decreased growth. In the *mtr4-1* mox-YG overexpressing cells, the

amount of other proteins decreased (to 54%), overwhelmed by the 40% expression of mox-YG.

However, the overall protein content increased compared to the *mtr4-1* vector control, with a

composition like the wild-type mox-YG overexpressing cells. Therefore, in *mtr4-1* cells, even though

mox-YG overexpression reduced the amount of other proteins (to 54%) compared to the vector control

(78%), the growth rate was not reduced (Figure 5—figure supplement 5B). This suggests that mox-YG

overexpression restores the effect of growth inhibition caused by *mtr4-1*. Next, we observed the

composition of the nucleolus in *mtr4-1* (Figures 5G and Figure 5—figure supplement 5D, E). The

failure of nucleolus formation observed upon mox-YG overexpression was partially, but significantly,

rescued in the *mtr4-1* strain. This supports our hypothesis that misassembled ribosomes are degraded

by the exosome (Figure 5D), and with low exosome activity, nucleolus formation (and efficient

ribosome assembly) would be possible even with overexpression of mox-YG (Figure 5—figure

supplement 4D).

While experimenting with mox-YG overexpressing cells, we found that these cells stop growth earlier

(lower final density) at higher temperatures (38°C) than the vector controls (Figure 5H and Figure 5—

figure supplement 6A). Cells at this time did not show cell death such as cell bursting (Figure 5—figure

supplement 6B). Recently, it was reported that intracellular osmotic pressure increases at high

temperatures, resulting in less formation of liquid-liquid phase-separated structures such as nucleoli,

which are recovered when cells are placed in hyperosmotic conditions (Watson et al., 2023). We

hypothesized that the growth inhibition of mox-YG overexpressing cells at high temperatures may be

due to the enhanced failure of nucleolus formation, and thus cultured mox-YG overexpressing cells

under hyperosmotic pressure. At 30°C, hyperosmolarity caused the same decrease in growth for the

vector control and mox-YG overexpression (Figures 5I and Figure 5—figure supplement 6C). On the

other hand, at 38°C, hyperosmolarity caused a decrease in growth only in the vector control and no

further decrease in mox-YG overexpression cells (Figures 5J and Figure 5—figure supplement 6D).

The same trend was observed regarding the size of the nucleolus (Figures 5—figure supplement 6E, F).

Under the conditions with added sorbitol and under mox-YG overexpression, the size of the nucleoli

was smaller than the control at 30°C, whereas, at 38°C, it was rather larger than the control. This

suggests that mox-YG overexpression and hyperosmolarity may mitigate each other's growth-inhibitory

effects, and one explanation may be that hyperosmolarity suppresses the abnormal organization of the

nucleolus caused by mox-YG overexpression.

Figure 5. mox-YG overexpression causes abnormal nucleolus formation.

A. 3D structure of the ribosome. PDB model 6GQV (Pellegrino et al., 2018) is used for the base model and colored using ChimeraX software (Meng et al., 2023). Ribosomal proteins with increased expression upon mox-YG overexpression are shown with the named structures containing them. See also Figure 5—figure supplement 1 for the quantitative data.

B) Electron microscope images of the vector control and mox-YG overexpressing cells. The arrow in the image points to the nucleolus structure. N: nucleus. Images of other observed cells are shown in Figure 5—figure supplement 3.

C, D) Model diagrams showing the hypothetical situation of the wild type (**C**) and mox-YG overexpressing cells (**D**). In WT cells, enough ribosomal proteins (RPs) are produced, resulting in RP-assembled rRNAs and the formation of nucleolus with normal morphology. On the other hand, in mox-YG overexpressing cells, the amount of RP is reduced due to translation competition, increasing misassembled rRNA. As a result, degradation of rRNA by exosomes may be accelerated, resulting in abnormal nucleolus morphology.

E) Growth curves of WT and *mtr4-1* mutant cells with the vector control or upon overexpression of mox-YG at 30˚C. The solid or dotted lines show the average calculated from three biological replicates. Growth curves with SDs of replicates are shown in Figure 5—figure supplement 5A.

F) Ratio of total protein levels of WT and *mtr4-1* strains with the vector or upon mox-YG overexpression, calculated the total protein level of WT cells with the vector as 100%. Gray bars indicate expression levels of proteins other than mox-YG; green shaded bars indicate mox-YG expression levels. Error bars were SDs calculated from three biological replicates. The *p*-values were calculated by performing Welch's *t*-test. "n.s." means $p > 0.05$.

G) Fluorescence microscopy image of nucleolus-localized protein Nsr1-mScarlet-I of the WT and *mtr4-1* mutant cells with the vector or mox-YG overexpression. Bright field and merged images, and quantification of the nucleolus size are shown in Figures 5—figure supplement 5D and 5E.

H) Growth curves of the cells with the vector or under mox-YG overexpression cultivated at 30°C or 38°C. The solid or dotted lines show the average calculated from three biological replicates. Growth curves with SDs of replicates are shown in Figure 5—figure supplement 6A.

I, J) Final OD of the cell culture with or without 1M sorbitol at 30°C (**I**) or 38°C (**J**). The bars and error bars show the means and SDs calculated from three biological replicates. The red dotted line indicates the final OD estimated from the product of (mox-YG without sorbitol) / (Vector without sorbitol) and (Vector with sorbitol) / (Vector without sorbitol). Growth curves with SDs of replicates are shown in Figures 5—figure supplement 6C, D.

The Gpm1 mutant shows a similar NI to mox-YG, yet yields distinct phenotypes upon overexpression.

Thus far, our investigation has focused on heterologous fluorescent proteins. Finally, we turned to endogenous yeast proteins to identify those with the lowest cytotoxicity and to examine the phenotypic consequences of their overexpression. We previously investigated the expression limits of a group of glycolytic enzymes in *S. cerevisiae*, with Gpm1 and its catalytic center mutant (CCmut) having the highest expression limits (Eguchi et al., 2018). Here, we evaluated the neutrality of the wild types and their CC-mutants of Tdh3 and Gpm1 (Figure 6A and Figure 6—figure supplement 1). The growth rates at the expression limits were higher in the CCmut for both Tdh3 and Gpm1, and in Gpm1, the CCmut grew two-fold faster (Figures 6—figure supplement 1A, C). The protein levels of wild types and mutants of both proteins were not significantly different (Figures 6—figure supplement 1D, E). The calculated NIs were highest for Gpm1-CCmut, which was equivalent to that of mox-YG (Figures 6A). We previously concluded that the activities of glycolytic enzymes generally do not affect their expression limits (Eguchi 2018), but their neutralities suggest that their activities can become constraints in some way. This may be due to a disturbance in metabolism or a property of the active enzyme, such as an effect of reactive amino acids (His or Cys, Figure 6—figure supplement 2).

Although Gpm1 and Gpm1-CCmut exhibited NIs comparable to that of mox-YG, suggesting that they are similarly unconstrained, notable differences in cell morphology were observed (Figure 6B, 6—figure supplement 3). Cells overexpressing Gpm1 appeared enlarged and spherical, while those overexpressing Gpm1-CCmut did not show such hypertrophy but were still rounder than vector controls or mox-YG overexpressing cells. The transcriptional responses observed in mox-YG overexpressing

cells, including the induction of *GAP1*, a marker of the amino acid starvation response, were not

detected in Gpm1-CCmut overexpression (Figure 6C and Figure 6—figure supplement 4). In Gpm1-

CCmut overexpressing cells, similar to mox-YG overexpression, the levels of proteins other than

Gpm1-CCmut were reduced, despite a slight, statistically non-significant increase in total protein

amount (Figure 6—figure supplement 1F). While mitochondrial development was observed in Gpm1-

CCmut cells using Rip1–mScarlet-I labeling (Figure 6—figure supplement 5), no increase in oxygen

consumption was detected (Figure 6D and Figure 6—figure supplement 6). The nucleolus, visualized

with Nsr1–mScarlet-I, appeared similarly shrunken in Gpm1-CCmut overexpressing cells as in mox-

YG overexpressing cells (Figure 6E and Figure 6—figure supplement 7). These results suggest that the

phenotypes observed in mox-YG overexpressing cells include both mox-YG–specific effects and

responses commonly associated with protein burden. Alternatively, as indicated by the abnormal cell

morphology, Gpm1-CCmut may possess unknown constraints or cytotoxic properties not captured by

the neutrality index, which could obscure the detection of phenotypes typically associated with protein

burden.

Figure 6. The Gpm1 mutant shows a similar NI to mox-YG, yet yields distinct phenotypes upon

overexpression.

A) The neutrality index for Gpm1, Tdh3, and their CCmuts. Data on growth rates and protein expression

levels used for the neutrality index calculation are presented in Figure 6—figure supplement 1.

B) Top images: representative microscopic images of cells overexpressing Gpm1 or Gpm1–CCmut.

Bottom graph: Distribution of cell axis ratios upon overexpression of Gpm1 and Gpm1–CCmut,

compared with the vector control. The p–values were calculated by the Mann–Whitney U–test.

Statistical analysis of cell size and comparison with mox-YG overexpressing cells are shown in Figure

6—figure supplement 3.

C) Promoter activity reporter assay. Constructs used for promoter analysis with transcription reporters

and quantitative results of RFP fluorescence values (arbitrary unit) for each promoter. Time series data

are in Figure 6—figure supplement 4.

D) Oxygen consumption rate in vector control, mox-YG, and Gpm1-CCmut overexpressing cells

(arbitrary units, AU). Antimycin A was added as a control for respiratory inhibition. The measurement

data are shown in Figure 6—figure supplement 6.

E) Fluorescence microscopy images of nucleolus with Nsr1–mScarlet–I. Log phase cells ($OD_{660} = 1.0$)

with the control vector or under Gpm1 or Gpm1–CCmut overexpression were observed. Statistical

analysis of nucleolar size is shown in Figure 6—figure supplement 7.

The bars and error bars in **A**, **C** and **D** show the means and SDs calculated from three biological

replicates. The raw data is shown with dot plots. The p-values were calculated by performing Welch's

t-test and applying Bonferroni correction. "n.s." means $p > 0.05$.

Discussion

In this study, we found that the fluorescent properties of FPs constrain their expression limits. This fluorescence-induced cytotoxicity appears as different phenotypes depending on the FPs; an excess of fluorescent mCherry causes cell enlargement, but such a phenotype was not seen with green fluorescent proteins. On the other hand, the cause of the cytotoxicity is not clear currently. Fluorescence itself does not seem to constrain the expression limit, because the mox expression limit remained the same when cultured in the dark or under strong light (Figure 2—figure supplement 7). We also believe that H_2O_2 generated during fluorophore maturation (Craggs, 2009) is unlikely. In the presence of H_2O_2, mox overexpressing cells grew slightly slower than mox-YG overexpressing cells (Figure 2—figure supplement 8H). However, no transcriptional response to oxidative stress, including the catalase gene (*CTT1*), was seen with mox overexpression (FDR < 0.187, over the vector control, Data S1). Because *CTT1* expression was elevated in mox-YG overexpression (FDR < 0.006, over the vector control, Data S1), presumably due to enhanced respiration, mox-YG overexpression might rather increase oxidative stress tolerance. Conversely, "inactive" proteins might accumulate more without interfering with intracellular functions because both inactive fluorescent proteins and inactive glycolytic enzymes showed higher neutrality (Figures 2 and 6). However, our current data did not clarify the mechanism, and investigating more proteins and their variants would be required.

The currently least cytotoxic proteins, non-fluorescent mox-YG and inactive Gpm1 express more than 40% of the total protein, whereas yeast cells can maintain growth under these conditions (Figures 2 and 6). This is higher than the prediction in *E. coli*, which stops growing when 37% of the excess protein is expressed (Bruggeman et al., 2020). This may be due to the very low cytotoxicity of the proteins expressed in this study, or it may be because eukaryotic cells have greater capacity than prokaryotic cells. In the mox-YG overexpressing cells, we observed a previously expected but unobserved cellular physiology of the protein burden: depletion of protein synthesis resources (i.e., nitrogen source; Figure 3) and a decrease in other proteins (glycolytic enzymes and ribosomes; Figure 4), possibly due to competition for synthesis. In addition, metabolic conversion to mitochondria and failure of nucleolus

formation were observed (Figures 4 and 5). As a matter of fact, the nitrogen source depletion response

(Figure 3) itself is not self-evident. Because the total protein content in the mox-YG overexpressing

cells is the same as that in the vector control (Figure 4A), the intracellular nitrogen source used for

protein synthesis should not be depleted. Then, why does the nitrogen source depletion response occur?

Excess mox-YG may promote wasteful consumption of amino acids and nitrogen, or there may be

pathways that sense the amount of "useful proteins" and their ability to be synthesized, or there may be

depletion of specific amino acids in mox-YG excess. Another possibility is that there may be some

mechanism that maintains the amount of total protein in the cell even when the nitrogen source is

depleted. Depletion of nitrogen sources probably results in the inactivation of the TORC1 pathway

(Figure 4C), metabolic conversion from fermentation to respiration, and ribosome depression to an

economic mode. In *E. coli*, the ppGpp pathway plays this role (Shachrai et al., 2010). While the above

is thought to be a physiological response to protein burden, a defect in ribosome synthesis in the

nucleolus was also suggested (Figure 5B). This could be due to a conflict between nuclear rRNA

synthesis and cytoplasmic protein synthesis or to the negative effects of the inability to create a liquid-

liquid phase-separated structure due to dilution effects (see below). The partial restoration of nucleolus

formation by a mutation of the exosome (Figure 5G) supports this possibility.

When one type of protein is expressed as much as 40% of the total, as in the present study, the

cytoplasmic crowding, viscosity, pH, etc. may change due to the nature of the protein and the effect of

the reduced protein by competition. These may alter the movement of substances. It is known that when

ribosomes are reduced, the fluidity of large proteins changes (Delarue et al., 2018). Water activity-

osmotic pressure may change (Watson et al., 2023), and pH may also be affected. These may affect the

formation of liquid-liquid phase-separated structures (Delarue et al., 2018; Watson et al., 2023). The

failure of nucleolus formation observed in mox-YG overexpression (Figure 5B) may be due to such a

condition change. In fact, the growth of mox-YG overexpressing cells is significantly impaired at high

temperatures (Figure 5H), where liquid-liquid phase separation is less likely to occur (Watson et al.,

2023). In contrast, the effect appears to be reduced at high osmolarity (Figure 5J), where it is more

likely to occur (Watson et al., 2023). A mutation of the nuclear exosome restored nucleolus formation

(Figure 5G), which might be due to the reformation of the liquid-liquid phase-separated structure (i.e.,

nucleolus) by increasing the concentration of rRNA. These have not been considered phenomena that

occur with protein overexpression. The engineering of physicochemical properties of the cytoplasm

thus may be possible by mass expression of proteins with specific properties.

In this study, in addition to mox-YG, inactive Gpm1 was also expressed in large amounts as well (Figure

6). Both are globular proteins with relatively small molecular weight, but it is not yet clear what

characteristics are necessary for them not to be cytotoxic. Further investigation of the neutrality of

various proteins will clarify these features. In addition, while the overexpression of Gpm1-CCmut also

caused nucleolar shrinkage similar to that observed in mox-YG overexpressing cells, it did not induce

other phenotypes such as nitrogen starvation or enhanced respiration (Figure 6). Therefore, it remains

unclear which of the phenotypes observed in mox-YG overexpressing cells are general consequences

of "protein burden" and which are specific to mox-YG. As noted in the Introduction (Figure 1E), it is

essential to continue investigating multiple proteins that appear to be "unconstrained" in order to

address this issue. In this study, expression limits were investigated, but no experiments were performed

to gradually increase expression levels. For example, it would be meaningful to investigate at what

expression level of mox-YG a shift to the economy mode or failure of nucleolus formation occurs. Also,

due to the limitations of the experimental system, it may not be possible to express mox-YG to its true

limit. The expression of one specific protein cannot be 100% of the total protein, and there is always a

limit. As the number of ribosomes continues to decrease due to competition for synthesis, the excess

proteins cannot be synthesized either. Thus, there must be a final equilibrium point. A more powerful

and controllable expression system would be needed to investigate this. Although the translation system

appears to be the primary target of burden in mox-YG overexpression (Figure 2—figure supplement

6C, D), the bottleneck in the expression system may also differ depending on gene design (Figure 1D)

(Hausser et al., 2019).

Materials and Methods

Strains, growth conditions, and yeast transformation

BY4741 (*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) (Brachmann et al., 1998) was used as the host strain for the experiments. Yeast culture and transformation were performed as previously described (Amberg et al., 2005). A synthetic complete (SC) medium without uracil (Ura) or leucine (Leu), as indicated, was used for yeast culture. Other strains used in this study are listed in the Key Resource Table.

Plasmid construction

The plasmids were constructed by homologous recombination activity of yeast cells (Oldenburg 1997), and their sequences were verified by DNA sequencing. Plasmids used in this study are listed in the Key Resource Table.

Genetic tug-of-war

To overexpress a target protein to a level that causes growth inhibition (in this study this is referred to as "expression limit"), we used gTOW (Figure 2—figure supplement 1A) (Moriya et al., 2012, 2006). The gene of the target protein is incorporated into the gTOW plasmid (here, pTOW40836) and introduced into yeast cells lacking *ura3* and *leu2* genes. The selection marker in this case is uracil (–Ura conditions). Since this plasmid has the replication origin of 2 μm plasmid (*2 μ ORI*), this plasmid becomes multicopy in the cell (about 30 in the case of the vector). When yeast cells carrying the plasmid are transferred to a medium without leucine and uracil (–Leu/Ura conditions), the copy number of the plasmid in the cells increases (to about 120 in the case of the vector). This is because *leu2-89* on the plasmid is a *LEU2* allele with a large deletion in its promoter, so cells with higher plasmid copy numbers will grow faster in –Leu/Ura conditions. In other words, *leu2-89* acts as a bias to raise the plasmid copy number in –Leu/Ura conditions. If overexpression of the target protein inhibits growth, the plasmid copy number will be lower than the copy number that gives rise to the critical expression of that protein.

In other words, the target gene acts as a bias to lower the plasmid copy number. The resulting tug-of-

war between the selection biases created by the two genes leads to a target protein being expressed at

close to the critical expression level in –Leu/Ura conditions (the green-colored area in Figure 2—figure

supplement 1A). By using a version of the aureobasidin A resistance gene (*AUR1*) with a deletion in its

promoter (*AUR1d*) as a bias to increase plasmid copy number, experiments based on the same concept

as gTOW can also be conducted in rich medium (YPD + 500 ng/mL aureobasidin A). In this case, the

plasmid pTOW-AUR1d is used (Figure 4—figure supplement 2A).

Measuring growth rate

Cellular growth was measured by monitoring OD_{660}, every 10 min using a compact rocking incubator

(TVS062CA, ADVANTEC). The max growth rate (MGR) was calculated as described previously

(Moriya *et al.* 2006). Average values, SD, and *p*-values of Welch's *t*-test were calculated from at least

three biological replicates. The growth data of Figures 2—figure supplement 10A was measured by

monitoring OD_{595}, respectively, every 10 min using a microplate reader (Infinite F200, TECAN).

Protein analysis

Yeast cells overexpressing a target protein were cultivated in SC–Leu/Ura medium. The total protein

was extracted from log-phase cells ($OD_{660} = 1.0$) with a NuPAGE LDS sample buffer (Thermo Fisher

Scientific) after 0.2 mol/l NaOH treatment (Kushnirov 2000). For each analysis, the total protein

extracted from 1 optical density (OD) unit of cells with OD_{660} was used. For total protein visualization,

the extracted total protein was labeled with Ezlabel FluoroNeo (ATTO), as described in the

manufacturer's protocol, and separated by 4–12% sodium-dodecyl sulfate acrylamide gel

electrophoresis (SDS-PAGE). Proteins were detected and measured using the LAS-4000 image

analyzer (GE Healthcare) in SYBR–green fluorescence detection mode and Image Quant TL software

(GE Healthcare). Protein quantification was performed as shown in Figure 2—figure supplement 1B.

Average values, SD, and *p*-values of Welch's *t*-test were calculated from three biological replicates.

Western blotting of Atg13

Cells overexpressing FPs were collected at OD_{660} = 1 or 2 and fixed with 100 µL of ice–cold alkaline

solution (0.2 N NaOH and 0.5% β–mercaptoethanol). After 5 min of incubation on ice, 1 mL of ice–

cold acetone was added to the sample and incubated at −20˚C to precipitate proteins. The protein

samples were precipitated with a microfuge (15,000 rpm, 5 min), air–dried, suspended in 100 µL of

SDS–PAGE sample buffer, and incubated at 65˚C for 15 min. The samples were thoroughly dissolved

by sonication and subjected to SDS–PAGE (40 mA, 1h). After ponceau S staining, proteins are

transferred to the PVDF membrane (Millipore #IPVH304F0) and blocked in 5% skim milk/TBST.

Atg13 was detected using a rabbit anti–Atg13 antibody (1:3000) and peroxidase-conjugated goat anti-

rabbit secondary antibodies (Jackson ImmunoResearch, #111–035–003, 1:10,000). Chemiluminescent

substrates (Millipore, #WBLUF0100) and Light–Capture II (ATTO) were used for signal detection.

Measuring GFP fluorescence

GFP fluorescence (Ex485nm/Em535nm) was measured every 10 min using a microplate reader (Infinite

F200, TECAN). Average values, SD, and p-values of Welch's t-test were calculated from eight

biological replicates.

RNA-seq analysis

BY4741 cells overexpressing a target protein were cultured in SC–Leu/Ura medium and harvested at

the logarithmic growth phase. RNA extraction was performed according to (Köhrer and Domdey, 1991).

The purified RNA was quality checked by BioAnalyzer (Agilent) or MultiNA (Shimazu), and the

concentration was measured by Qubit (Thermo Fisher Scientific). Purified RNA was stored at −80°C

until subsequent experiments. cDNA libraries were prepared using the TruSeq Stranded Total RNA kit

(Illumina), using half the protocol of the TruSeq RNA library prep kit. Three biological replicates were

analyzed for all strains. The sequences were checked for read quality by FastP (Chen et al., 2018) and

then aligned using Hisat2 (Kim et al., 2019). The aligned data were formatted into bam files by Samtools

(Li et al., 2009) and quantified by StringTie (Pertea et al., 2015). Finally, expression level variation

analysis was performed by EdgeR (Robinson et al., 2010). The raw data were deposited into DDBJ

(accession number: PRJDB18064). The processed data (Data_S1) included the average expression

levels (log10CPM) of 6685 genes in the vector control, the changes in expression (log2FC) upon

overexpression of mox or mox-YG compared to the vector control, and the significance values of

expression change (mox_FDR, mox-YG_FDR). Figures 3—figure supplement 1A, B is the

visualization of this data using volcano plots. This data is used in subsequent data analyses below.

Reporter assay

The strains used for the reporter assay were created by introducing the promoter region of each gene

and the sequence of mScarlet-I, a type of RFP, into the *FCY1* locus of the genome as shown in Figure

3—figure supplement 2A. Correct integration was checked by histidine prototrophy and 5-FC resistance

of the strain. The cell density (OD_{595}) and RFP fluorescence (Ex535nm/Em590nm) of the created strains

were measured every 10 min using a microplate reader (Infinite F200PRO, TECAN). Average and SD

of max RFP fluorescence, and *p*-values of Welch's *t*-test were calculated from eight biological replicates.

Proteome analysis

The vector control and mox-YG overexpressing cells (three biological replicates) were cultured in 5

mL of SC–LeuUra medium with shaking and cells were collected at $OD_{660} = 1$. Cells were washed twice

with PBS, frozen on dry ice, and transported to Kazusa DNA Research Institute. The samples were

prepared as described in the previous study (Hughes et al., 2019; Kawashima et al., 2022). The cells

were dissolved in 100 mM Tris-HCl (pH 8.0) containing 4% SDS and 20 mM NaCl using BIORUPTOR

BR-II (SONIC BIO Co., Kanagawa, Japan). The 20 μg of extracted proteins were quantified using

Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific, WA, USA) at 500 ng/μL. The protein

extracts were reduced with 20 mM tris (2-carboxyethyl) phosphine for 10 min at 80°C followed by

alkylation with 30 mM iodoacetamide for 30 min at room temperature in the dark. Protein purification

and digestion were performed using the sample preparation (SP3) method (Kawashima et al., 2022).

The tryptic digestion was performed using 500 ng/μL Trypsin/Lys-C Mix (Promega, Madison, WI,

USA) for overnight at 37°C. The digests were purified using GL-Tip SDB (GL Sciences, Tokyo, Japan)

according to the manufacturer's protocol. The peptides were dissolved again in 2% ACN containing

0.1% TFA and quantified using BCA assay at 150 ng/µL. The digested peptides were loaded directly

using a 75 µm × 12 cm nanoLC nano-capillary column (Nikkyo Technos Co., Ltd., Tokyo, Japan) at

40°C and then separated with a 30-minute gradient (mobile phase A = 0.1% FA in water, B = 0.1% FA

in 80% ACN) consisting of 0 min 8% B, 30 min 70% B at a flow rate of 200 nL/min using an UltiMate

3000 RSLCnano LC system (Thermo Fisher Scientific). The eluted peptides were detected using a

quadrupole Orbitrap Exploris 480 hybrid mass spectrometer (Thermo Fisher Scientific) with normal

window DIA. The MS1 scan range was set as a full scan with m/z 490-745 at a mass resolution of

15,000 to set an Auto Gain Control (AGC) target for MS1 as 3×10^6 and a maximum injection time of

23 msec. The MS2 were collected at more than m/z 200 at 30,000 resolutions to set an AGC target of 3

$\times 10^6$, maximum injection time of "auto", and fixed normalized collision energy of 28%. The isolation

width for MS2 was set to 4 m/z, and for the 500-740 m/z window pattern, an optimized window

arrangement was used in Scaffold DIA (Proteome Software, Inc., Portland, OR, USA). The raw data

were searched against an *in silico* predicted spectral library using DIA-NN (Demichev et al., 2020).

(version:1.8.1, https://github.com/vdemichev/DiaNN) First, an *in-silico* predicted spectral library was

generated from the human protein sequence database (UniProt id UP000005640, reviewed, canonical,

20381 entries) using DIA-NN. The DIA-NN search parameters were as follows: protease, trypsin;

missed cleavages, 1; peptide length range, 7-45; precursor charge range, 2-4; precursor mass range,

495-745; fragment ion m/z range, 200-1800; mass accuracy, 10 ppm; static modification, cysteine

carbamidomethylation; enabled "Heuristic protein interferences," "Use isotopologues," "MBR," and

"No shared spectra." Additional commands were set as follows: "mass acc cal 10," "peak translation,"

and "matrix spec q". The protein identification threshold was set at < 1% for both peptide and protein

false discovery rates (FDRs).

The signal intensity data for 4241 proteins (Data_S2) was used for further data analysis below.

Data analysis

The analysis and visualization of RNA-seq data and proteome data were conducted using custom

Python code, and coding and execution were carried out in the Collaboratory. In the RNAseq data,

6,685 protein-coding genes were detected (Data_S1). Among these, 4,291 genes assigned to various

categories of KEGG Orthology level 3 (Kanehisa et al., 2023) were placed into 77 categories. Analysis

was performed on 47 categories that contained more than 10 gene elements. We investigated whether

the expression of genes in each category, as a group, was significantly higher or lower upon

overexpression of mox and mox-YG compared to the vector control. Specifically, mox/Vector (log2FC)

and mox-YG/Vector (log2FC) were calculated along with their FDR values, and each gene was assigned

to a KEGG orthology level 3 category. Next, whether the gene groups within each category showed

significantly higher or lower expression changes compared to gene groups outside of their category was

tested using the Mann-Whitney U-test. Significant differences were indicated in the graphs with a red

border and red stripes. The graphs showed the overall distribution of genes within each category using

violin plots, and only genes showing significant differences in mox/Vector (log2FC) and mox-

YG/Vector (log2FC) were highlighted in swarm plots. An asterisk indicated significant differences on

the graphs when the expression changes (log2FC) between mox/Vector and mox-YG/Vector were

significant by the Mann-Whitney U-test. The significance threshold was set at FDR < 0.05 for all. Only

categories where either mox or mox-YG overexpression showed significant expression changes were

extracted and presented as Figure 3B, and all 47 categories were shown in Figure 3—figure supplement

1C.

The proteome analysis was conducted using the signal intensity data (Data_S2). This data includes

proteins that were detected only in some samples under certain conditions or replicates, especially those

with low signals or large expression variations. Additionally, the total protein signal sum excluding

mox-YG was almost the same between the vector control and mox-YG overexpression (Figure 4—

figure supplement 3A). This result contradicts the SDS−PAGE findings, which indicate that the total

protein mass per cell, including mox-YG, is the same between the vector control and mox-YG

overexpression (Figure 4A). Therefore, it is considered necessary to adjust the total protein signal sum,

including mox-YG, to match that of the vector control to understand the proteome change per cell. In

this study, however, we used the most conservative approach by removing proteins with zero values

and not applying any overall corrections (Figure 4—figure supplement 3B), which allowed the analysis

of 3,588 proteins. Although this may not correctly analyze "per cell" proteome change, it enables us to

capture the change in proteome composition (excluding mox-YG), thus providing information about

cellular responses. If the total correction is applied, only 138 proteins were increased (Figure 4—figure

supplement 3C), most of which are mitochondrial proteins.

For the detected 3,588 proteins above, we compared the increase or decrease of protein groups within

each category of KEGG Orthology Level 3 with those outside the category. Analysis was performed on

45 categories, each containing more than 10 proteins, to investigate whether the expression of protein

groups in each category was significantly higher or lower under mox-YG overexpression compared to

the vector control. Specifically, the average log2FC and its FDR values were calculated using data from

three replicates for mox-YG/Vector, and each protein was assigned to a KEGG orthology level 3

category. Next, whether the protein groups within each category showed significantly higher or lower

expression changes compared to protein groups outside of their category was tested using the Mann-

Whitney *U*-test. Significant differences were indicated in the graphs with a red border and red stripes.

The graphs showed the overall distribution of proteins within each category using violin plots, and only

proteins showing significant differences in mox-YG/Vector (log2FC) were highlighted in swarm plots.

The significance threshold was set at FDR < 0.05 for all. Only categories where mox-YG

overexpression showed significant expression changes were extracted and presented in Figure 4C and

all 47 categories were shown in Figure 4—figure supplement 6. Published data (Gowans et al., 2018)

was used to separate protein groups into rapamycin-responsive and non-responsive, and similar analysis

and visualization were conducted (Figures 4—figure supplement 6A, B).

Fluorescence microscopic observation

Cells overexpressing target proteins were cultured in SC–LeuUra medium and observed under an

inverted microscope (DMI 6000 B, Leica) at log phase ($OD_{660} = 1.0$). Images were acquired with Leica

Application Suite and processed with Leica Application Suite X software (Leica Microsystems). Cells

in brightfield images were separated from the images using yeast_segmentation-master_v3 (Lu et al.,

2019). Cell morphological traits were assessed using parameters obtained when analysed using

CellProfiler (Ver. 4.2.6) (Carpenter et al., 2006). "AreaShape_Area" was used to analyse FP-

overexpressing cell size (arbitrary unit; AU) in Figure 2—figure supplement 6B, Nsr1-GFP in Figure

5—figure supplement 2C and Nsr1-mScarlet-I in Figure 5—figure supplement 5E, 6F, and Figure 6—

figure supplement 7C. For analysis of the fluorescence intensity of Rip1-mScarlet-I in Figure 4—figure

supplement 8B and Figure 6—figure supplement 5A. "Intensity_ IntegratedIntensity" was used.

Intensity_MeanIntensity" was used to analyze the fluorescence intensity of Nsr1-GFP in Figure 5—

figure supplement 2D and Figure 6—figure supplement 7C. Ten images per experiment were used for

analysis. The axis ratio in Figure 6—figure supplement 3 was calculated as the ratio of

"AreaShape_MajorAxisLength" to "AreaShape_MinorAxisLength" in the data

Measurement of oxygen consumption

Cells overexpressing the target protein were collected at $OD_{660} = 1$. The cells (0.05 OD unit) were

treated with oxygen probe in SC–LeuUra and 50μM antimycin A in SC–LeuUra. Oxygen consumption

was measured according to the instruction manual of the Extracellular OCR Plate Assay Kit (Dojindo,

#E297). The fluorescence intensity (Ex500nm/Em650 nm) was recorded at 10 min intervals using a

microplate reader (Infinite F200, TECAN). The oxygen consumption rate (AU) was calculated as the

slope of a linear approximation of the kinetic data.

Electron microscopic observation

Cells grown to log phase ($OD_{600} = 1.0$) at 30°C in SC–LeuUra medium were collected and transported

to Tokai Electron Microscope Co. Transported samples were sandwiched between copper plates, flash-

frozen with liquefied propane, and then dehydrated with anhydrous ethanol. cells were fixed with a 5:5

mixture of propylene oxide and resin (Quetol-651, Nissin EM Co.). Fixed cells were cut with an

ultramicrotome (Ultracut UCT, Leica) to prepare 80 nm thick sections, stained with 2% uranyl acetate

and lead stain solution (Sigma-Aldrich Co.), and observed with a transmission electron microscope

(JEM-1400 Plus, JEOL Ltd.).

Data availability

All raw data, analysis code, and omics analysis pipelines used for generating the figures have been

deposited in GitHub (https://github.com/hisaomlab/Fujita_eLife_2025).

Acknowledgments

931 We thank the members of the Moriya laboratory (Okayama University) for their helpful discussions.

Funding

934 This work was partly supported by the Nagase Science and Technology Foundation, Institute for

935 Fermentation, Osaka (IFO) General Research Funding, JSPS KAKENHI Grant Numbers 22K19294,

936 and 24K02313, JST SPRING Japan Grant Number JPMJSP2126.

References

Amberg DC, Burke DJ, Burke D, Strathern JN. 2005. Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual. CSHL Press.

Bolognesi B, Lorenzo Gotor N, Dhar R, Cirillo D, Baldrighi M, Tartaglia GG, Lehner B. 2016. A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity upon Increased Protein Expression. *Cell Rep* **16**:222–231.

Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD. 1998. Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**:115–132.

Bruggeman FJ, Planqué R, Molenaar D, Teusink B. 2020. Searching for principles of microbial physiology. *FEMS Microbiol Rev* **44**:821–844.

Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang JH, Lindquist RA, Moffat J, Golland P, Sabatini DM. 2006. CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol* **7**:R100.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**:i884–i890.

Costantini LM, Baloban M, Markwardt ML, Rizzo M, Guo F, Verkhusha VV, Snapp EL. 2015. A palette of fluorescent proteins optimized for diverse cellular environments. *Nat Commun* **6**:7670.

Craggs TD. 2009. Green fluorescent protein: structure, folding and chromophore maturation. *Chem Soc Rev* **38**:2865–2875.

Delarue M, Brittingham GP, Pfeffer S, Surovtsev IV, Pinglay S, Kennedy KJ, Schaffer M, Gutierrez JI, Sang D, Poterewicz G, Chung JK, Plitzko JM, Groves JT, Jacobs-Wagner C, Engel BD, Holt LJ. 2018. mTORC1 Controls Phase Separation and the Biophysical Properties of the Cytoplasm by Tuning Crowding. *Cell* **174**:338–349.e20.

Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M. 2020. DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. *Nat Methods* **17**:41–44.

Dong H, Nilsson L, Kurland CG. 1995. Gratuitous overexpression of genes in Escherichia coli leads to

growth inhibition and ribosome destruction. *J Bacteriol* **177**:1497–1504.

Eguchi Y, Makanae K, Hasunuma T, Ishibashi Y, Kito K, Moriya H. 2018. Estimating the protein

burden limit of yeast cells by measuring the expression limits of glycolytic proteins. *Elife* **7**.

doi:10.7554/eLife.34595

Farkas Z, Kalapis D, Bódi Z, Szamecz B, Daraba A, Almási K, Kovács K, Boross G, Pál F, Horváth P,

Balassa T, Molnár C, Pettkó-Szandtner A, Klement É, Rutkai E, Szvetnik A, Papp B, Pál C. 2018.

Hsp70-associated chaperones have a critical role in buffering protein production costs. *Elife* **7**.

doi:10.7554/eLife.29845

Frumkin I, Schirman D, Rotman A, Li F, Zahavi L, Mordret E, Asraf O, Wu S, Levy SF, Pilpel Y. 2017.

Gene Architectures that Minimize Cost of Gene Expression. *Mol Cell* **65**:142–153.

Gowans GJ, Schep AN, Wong KM, King DA, Greenleaf WJ, Morrison AJ. 2018. INO80 Chromatin

Remodeling Coordinates Metabolic Homeostasis with Cell Division. *Cell Rep* **22**:611–623.

Hausser J, Mayo A, Keren L, Alon U. 2019. Central dogma rates and the trade-off between precision

and economy in gene expression. *Nat Commun* **10**:68.

Hughes CS, Moggridge S, Müller T, Sorensen PH, Morin GB, Krijgsveld J. 2019. Single-pot, solid-

phase-enhanced sample preparation for proteomics experiments. *Nat Protoc* **14**:68–85.

Kafri M, Metzl-Raz E, Jona G, Barkai N. 2016. The Cost of Protein Production. *Cell Rep* **14**:22–31.

Kaizu K, Moriya H, Kitano H. 2010. Fragilities caused by dosage imbalance in regulation of the

budding yeast cell cycle. *PLoS Genet* **6**:e1000919.

Kamada Y, Funakoshi T, Shintani T, Nagano K, Ohsumi M, Ohsumi Y. 2000. Tor-mediated induction

of autophagy via an Apg1 protein kinase complex. J Cell Biol 150:1507–1513.

Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. 2023. KEGG for taxonomy-

based analysis of pathways and genomes. *Nucleic Acids Res* **51**:D587–D592.

Kastberg LLB, Ard R, Jensen MK, Workman CT. 2022. Burden Imposed by Heterologous Protein

Production in Two Major Industrial Yeast Cell Factories: Identifying Sources and Mitigation

Strategies. *Front Fungal Biol* **3**:827704.

Kawashima Y, Nagai H, Konno R, Ishikawa M, Nakajima D, Sato H, Nakamura R, Furuyashiki T,

Ohara O. 2022. Single-Shot 10K Proteome Approach: Over 10,000 Protein Identifications by Data-Independent Acquisition-Based Single-Shot Proteomics with Ion Mobility Spectrometry. *J Proteome Res* **21**:1418–1427.

Keppler-Ross S, Noffz C, Dean N. 2008. A new purple fluorescent color marker for genetic studies in Saccharomyces cerevisiae and Candida albicans. *Genetics* **179**:705–710.

Keren L, Hausser J, Lotan-Pompan M, Vainberg Slutskin I, Alisar H, Kaminski S, Weinberger A, Alon U, Milo R, Segal E. 2016. Massively Parallel Interrogation of the Effects of Gene Expression Levels on Fitness. *Cell* **166**:1282–1294.e18.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**:907–915.

Kintaka R, Makanae K, Moriya H. 2016. Cellular growth defects triggered by an overload of protein localization processes. *Scientific Reports*. doi:10.1038/srep31774

Kintaka R, Makanae K, Namba S, Kato H, Kito K, Ohnuki S, Ohya Y, Andrews BJ, Boone C, Moriya H. 2020. Genetic profiling of protein burden and nuclear export overload. *Elife* **9**. doi:10.7554/eLife.54080

Koch AL. 1983. The protein burden of lac operon products. *J Mol Evol* **19**:455–462.

Köhrer K, Domdey H. 1991. [27] Preparation of high molecular weight RNAMethods in Enzymology. Academic Press. pp. 398–405.

Kummer AD, Wiehler J, Rehaber H, Kompa C, Steipe B, Michel-Beyerle ME. 2000. Effects of Threonine 203 Replacements on Excited-State Dynamics and Fluorescence Properties of the Green Fluorescent Protein (GFP). *J Phys Chem B* **104**:4791–4798.

Lafontaine DLJ. 2015. Noncoding RNAs in eukaryotic ribosome biogenesis and function. *Nat Struct Mol Biol* **22**:11–19.

Liebermeister W, Noor E, Flamholz A, Davidi D, Bernhardt J, Milo R. 2014. Visual account of protein investment in cellular functions. *Proc Natl Acad Sci U S A* **111**:8488–8493.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:2078–2079.

Lu AX, Zarin T, Hsu IS, Moses AM. 2019. YeastSpotter: accurate and parameter-free web segmentation

for microscopy images of yeast cells. *Bioinformatics* **35**:4525–4527.

Mahima, Sharma AK. 2023. Optimization of ribosome utilization in Saccharomyces cerevisiae. *PNAS*

Nexus **2**:gad074.

Makanae K, Kintaka R, Makino T, Kitano H, Moriya H. 2013. Identification of dosage-sensitive genes

in Saccharomyces cerevisiae using the genetic tug-of-war method. *Genome Res* **23**:300–311.

Ma L, Pang CNI, Li SS, Wilkins MR. 2010. Proteins Deleterious on Overexpression Are Associated

with High Intrinsic Disorder, Specific Interaction Domains, and Low Abundance. *J Proteome Res*

9:1218–1225.

Martin DE, Soulard A, Hall MN. 2004. TOR regulates ribosomal protein gene expression via PKA and

the Forkhead transcription factor FHL1. *Cell* **119**:969–979.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. UCSF

ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**:e4792.

Metzl-Raz E, Kafri M, Yaakov G, Barkai N. 2020. Gene Transcription as a Limiting Factor in Protein

Production and Cell Growth. *G3* **10**:3229–3242.

Metzl-Raz E, Kafri M, Yaakov G, Soifer I, Gurvich Y, Barkai N. 2017. Principles of cellular resource

allocation revealed by condition-dependent proteome profiling. *Elife* **6**. doi:10.7554/eLife.28034

Moriya H. 2015. Quantitative nature of overexpression experiments. *Mol Biol Cell* **26**:3932–3939.

Moriya H, Makanae K, Watanabe K, Chino A, Shimizu-Yoshida Y. 2012. Robustness analysis of

cellular systems using the genetic tug-of-war method. *Mol Biosyst* **8**:2513–2522.

Moriya H, Shimizu-Yoshida Y, Kitano H. 2006. In vivo robustness analysis of cell division cycle genes

in Saccharomyces cerevisiae. *PLoS Genet* **2**:e111.

Namba S, Kato H, Shigenobu S, Makino T, Moriya H. 2022. Massive expression of cysteine-containing

proteins causes abnormal elongation of yeast cells by perturbing the proteasome. *G3* **12**.

doi:10.1093/g3journal/jkac106

Oldenburg K. 1997. Recombination-mediated PCR-directed plasmid construction in vivo in yeast.

Nucleic Acids Research. doi:10.1093/nar/25.2.451

Papp B, Pál C, Hurst LD. 2003. Dosage sensitivity and the evolution of gene families in yeast. *Nature*

424:194–197.

Pellegrino S, Demeshkina N, Mancera-Martinez E, Melnikov S, Simonetti A, Myasnikov A, Yusupov

M, Yusupova G, Hashem Y. 2018. Structural Insights into the Role of Diphthamide on Elongation

Factor 2 in mRNA Reading-Frame Maintenance. *J Mol Biol* **430**:2677–2687.

Peralta D, Bronowska AK, Morgan B, Dóka É, Van Laer K, Nagy P, Gräter F, Dick TP. 2015. A proton

relay enhances H2O2 sensitivity of GAPDH to facilitate metabolic adaptation. *Nat Chem Biol*

11:156–163.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables

improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**:290–295.

Prelich G. 2012. Gene overexpression: uses, mechanisms, and interpretation. *Genetics* **190**:841–854.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics* **26**:139–140.

Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010. Interdependence of cell growth and

gene expression: origins and consequences. *Science* **330**:1099–1102.

Shachrai I, Zaslaver A, Alon U, Dekel E. 2010. Cost of unneeded proteins in E. coli is reduced after

several generations in exponential growth. *Mol Cell* **38**:758–767.

Stoebel DM, Dean AM, Dykhuizen DE. 2008. The cost of expression of Escherichia coli lac operon

proteins is in the process, not in the products. *Genetics* **178**:1653–1660.

Vavouri T, Semple JI, Garcia-Verdugo R, Lehner B. 2009. Intrinsic protein disorder and interaction

promiscuity are widely associated with dosage sensitivity. *Cell* **138**:198–208.

Watson JL, Seinkmane E, Styles CT, Mihut A, Krüger LK, McNally KE, Planelles-Herrero VJ, Dudek

M, McCall PM, Barbiero S, Vanden Oever M, Peak-Chew SY, Porebski BT, Zeng A, Rzechorzek

NM, Wong DCS, Beale AD, Stangherlin A, Riggi M, Iwasa J, Morf J, Miliotis C, Guna A, Inglis

AJ, Brugués J, Voorhees RM, Chambers JE, Meng Q-J, O'Neill JS, Edgar RS, Derivery E. 2023.

Macromolecular condensation buffers intracellular water potential. *Nature* **623**:842–852.

White MF, Fothergill-Gilmore LA. 1992. Development of a mutagenesis, expression and purification

system for yeast phosphoglycerate mutase. Investigation of the role of active-site His181. *Eur*

J Biochem **207**:709–714.

Youn J-Y, Friesen H, Nguyen Ba AN, Liang W, Messier V, Cox MJ, Moses AM, Andrews B. 2017.

Functional Analysis of Kinases and Transcription Factors in Saccharomyces cerevisiae Using an

Integrated Overexpression Library. *G3* **7**:911–921.

Figure supplements

1080

Figure 2—figure supplement 1. Methods used for overexpression and protein expression

quantification.

A) The genetic tug-of-war (gTOW) method. See Materials and Methods for details.

B) Protein quantification method. All proteins present in the cells were labeled with a fluorescent dye

(Ezlabel FluoroNeo) and separated by SDS-PAGE. The protein level of the target protein (Protein level)

was calculated by subtracting the band at the same position of the vector control (Orange) from the band

of the target protein on the gel (Purple), which was estimated from the molecular weight, and dividing

the result by the total protein (Green). Other protein (Other protein level) was calculated by subtracting

Protein level (%) from the value obtained by normalizing the total protein (Green) of the overexpression

strain by the total protein (Red) of the vector control.

1091

1092   **Figure 2—figure supplement 2. Comparison of growth of fluorescent protein overexpressing**

1093   **strains**

Growth curves of the cells with the vector (pTOW40836) and the cells overexpressing fluorescent proteins and their non-fluorescent YG mutants in synthetic medium (–LeuUra). The solid or dotted lines show the average calculated from three biological replicates. Shaded zones indicate their standard deviations. **A**: EGFP and EGFP-YG, **B**: sfGFP and sfGFP-YG, **C**: moxGFP (mox) and mox-YG, **D**: mCherry (mChe) and mChe-YG.

Figure 2—figure supplement 3. Neutrality indexes of constrained proteins

A) MGRs of cells overexpressing indicated proteins.

B) Protein levels of the indicated target proteins (percent over the total proteins).

C) The neutrality index for the indicated proteins.

MTS3–EGFP: EGFP targeted to mitochondria; cultured in SC–LeuUra; expressed by the *PYK1* promoter. NES–EGFP: EGFP with nuclear export signal; cultured in SC–Ura; expressed by the *HXT7* promoter. ss–moxGFP: moxGFP targeted to the endoplasmic reticulum; cultured in SC–Ura; expressed by the *PYK1* promoter. EGFP_K: the EGFP result of Kintaka *et al*. (Kintaka et al., 2016); cultured in SC–LeuUra; expressed by the *PYK1* promoter. EGFP_F: the EGFP result of this paper.

In **A** and **B**, the data was obtained from Kintaka *et al*. (Kintaka et al., 2016), except mox-YG and EGFP_F (same as Figure 2 in this paper). The bars and error bars show the means and SDs calculated from three biological replicates.

moxGFP

```
  1 M  S  K  G  E  E  L  F  T  G  V  V  P  I  L  V  E  L  D  G  20
  1 ATGTCTAAAGGTGAAGAATTATTCACTGGTGTTGTCCCAATTTTGGTTGAATTAGATGGT 60

 21 D  V  N  G  H  K  F  S  V  R  G  E  G  E  G  D  A  T  N  G  40
 61 GATGTTAATGGTCACAAATTTTCTGTCcgtGGTGAAGGTGAAGGTGATGCTACTaAtGGT 120

 41 K  L  T  L  K  F  I  S  T  T  G  K  L  P  V  P  W  P  T  L  60
121 AAATTGACCTTAAAATTTATTTcTACTACTGGTAAATTGCCAGTTCCATGGCCAACCTTA 180

 61 V  T  T  L  T  Y  G  V  Q  S  F  S  R  Y  P  D  H  M  K  R  80
181 GTCACTACTTTAACTTATGGTGTTCAATcTTTTTCTAGATACCCAGATCATATGAAACgt 240

 81 H  D  F  F  K  S  A  M  P  E  G  Y  V  Q  E  R  T  I  S  F 100
241 CATGACTTTTTCAAGTCTGCCATGCCAGAAGGTTATGTTCAAGAAAGAACTATTTcTTTC 300

101 K  D  D  G  T  Y  K  T  R  A  E  V  K  F  E  G  D  T  L  V 120
301 AAAGATGACGGTACtTACAAGACCAGAGCTGAAGTCAAGTTTGAAGGTGATACCTTAGTT 360

121 N  R  I  E  L  K  G  I  D  F  K  E  D  G  N  I  L  G  H  K 140
361 AATAGAATCGAATTAAAAGGTATTGATTTTAAAGAAGATGGTAACATTTTAGGTCACAAA 420

141 L  E  Y  N  F  N  S  H  N  V  Y  I  T  A  D  K  Q  K  N  G 160
421 TTGGAATACAACTtcAACTCTCACAATGTTTACATCActGCTGACAAACAAAAGAATGGT 480

161 I  K  A  N  F  K  I  R  H  N  V  E  D  G  S  V  Q  L  A  D 180
481 ATCAAAGcTAACTTCAAAATTAGACACAACgtTGAAGATGGTTCTGTTCAATTAGCTGAC 540

181 H  Y  Q  Q  N  T  P  I  G  D  G  P  V  L  L  P  D  N  H  Y 200
541 CATTATCAACAAAATACTCCAATTGGTGATGGTCCAGTCTTGTTACCAGACAACCATTAC 600

201 L  S  T  Q  S  R  L  S  K  D  P  N  E  K  R  D  H  M  V  L 220
601 TTATCCACTCAATCTcgtTTATCCAAAGATCCAAACGAAAAGAGAGACCACATGGTCTTG 660

221 L  E  F  V  T  A  A  G  I  T  H  G  M  D  E  L  Y  K  *  239
661 TTAGAATTTGTTACTGCTGCTGGTATTACCCATGGTATGGATGAATTGTACAAATAA 717
```

sfGFP

```
  1 M  S  K  G  E  E  L  F  T  G  V  V  P  I  L  V  E  L  D  G  20
  1 ATGTCTAAAGGTGAAGAATTATTCACTGGTGTTGTCCCAATTTTGGTTGAATTAGATGGT 60

 21 D  V  N  G  H  K  F  S  V  R  G  E  G  E  G  D  A  T  N  G  40
 61 GATGTTAATGGTCACAAATTTTCTGTCcgtGGTGAAGGTGAAGGTGATGCTACTaAtGGT 120

 41 K  L  T  L  K  F  I  C  T  T  G  K  L  P  V  P  W  P  T  L  60
121 AAATTGACCTTAAAATTTATTTgTACTACTGGTAAATTGCCAGTTCCATGGCCAACCTTA 180

 61 V  T  T  L  T  Y  G  V  Q  C  F  S  R  Y  P  D  H  M  K  R  80
181 GTCACTACTTTAACTTATGGTGTTCAATgTTTTTCTAGATACCCAGATCATATGAAACgt 240

 81 H  D  F  F  K  S  A  M  P  E  G  Y  V  Q  E  R  T  I  S  F 100
241 CATGACTTTTTCAAGTCTGCCATGCCAGAAGGTTATGTTCAAGAAAGAACTATTTcTTTC 300

101 K  D  D  G  T  Y  K  T  R  A  E  V  K  F  E  G  D  T  L  V 120
301 AAAGATGACGGTACtTACAAGACCAGAGCTGAAGTCAAGTTTGAAGGTGATACCTTAGTT 360

121 N  R  I  E  L  K  G  I  D  F  K  E  D  G  N  I  L  G  H  K 140
361 AATAGAATCGAATTAAAAGGTATTGATTTTAAAGAAGATGGTAACATTTTAGGTCACAAA 420

141 L  E  Y  N  F  N  S  H  N  V  Y  I  T  A  D  K  Q  K  N  G 160
421 TTGGAATACAACTtcAACTCTCACAATGTTTACATCActGCTGACAAACAAAAGAATGGT 480

161 I  K  A  N  F  K  I  R  H  N  V  E  D  G  S  V  Q  L  A  D 180
481 ATCAAAGcTAACTTCAAAATTAGACACAACgtTGAAGATGGTTCTGTTCAATTAGCTGAC 540

181 H  Y  Q  Q  N  T  P  I  G  D  G  P  V  L  L  P  D  N  H  Y 200
541 CATTATCAACAAAATACTCCAATTGGTGATGGTCCAGTCTTGTTACCAGACAACCATTAC 600

201 L  S  T  Q  S  R  L  S  K  D  P  N  E  K  R  D  H  M  V  L 220
601 TTATCCACTCAATCTcgtTTATCCAAAGATCCAAACGAAAAGAGAGACCACATGGTCTTG 660

221 L  E  F  V  T  A  A  G  I  T  H  G  M  D  E  L  Y  K  *  239
661 TTAGAATTTGTTACTGCTGCTGGTATTACCCATGGTATGGATGAATTGTACAAATAA 717
```

1112

1113   **Figure 2—figure supplement 4. Base and amino acid sequences of moxGFP and sfGFP.**

1114   **Amino acids relevant to the text are shown in red.**

1115

 Figure 2—figure supplement 5. Comparative evaluation of mCherry (mCherry-Kafri) used in

previous studies and this study.

A) Growth curves of cells overexpressing WT and YG mutants of mCherry used in Kafri *et al.* (Kafri et al. 2016) in synthetic medium (–LeuUra). The solid or dotted lines show the average calculated from three biological replicates.

B) Ratio of MGR of cells overexpressing FPs and their mutants over the vector control.

C) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing FPs. All proteins were separated after staining with a fluorescent dye.

D) Protein level of the target protein (percent over the total protein). The amount was calculated from the intensity of the target bands separated by SDS-PAGE of **D**. Note that mChe may not be accurately measured due to molecular cleavage of the protein.

E) Relationship between MGR and the protein level. MGR data are the same as in **B**, and the protein level data are the same as in **D**. The neutrality indexes (NIs), calculated from the products of MGR and %protein levels, are indicated by the lines. If the investigated proteins have the same NI, they are located on the same dotted line segment.

F) The neutrality index for the FPs and mutants. The neutrality index is the product of %MGR (/Vector) and %protein level (/total protein).

The bars and error bars in **B**, **D**, and **F** show the means and SDs calculated from three biological replicates. The raw data is shown with dot plots.

1135

Figure 2—figure supplement 6. Overexpression of mox-YG does not cause cell enlargement or

burden on transcription.

A) Representative microscopic images of cells overexpressing indicated target proteins or moxFS (a

frameshift mutant of mox).

B) Violin plots showing the cell size quantified from microscopic images. The number of cells used for

the quantification is shown.

C) Growth curves of the vector control and cells overexpressing moxFS and mox-YG in synthetic

medium (–LeuUra). The solid or dotted lines show the average calculated from three biological

replicates.

D) MGR of cells overexpressing moxFS and mox-YG (percent over the vector control). The bars and

error bars show the means and SDs of the MGR calculated from three biological replicates. The raw

data is shown with dot plots. The *p*-value of Welch's *t*-test is shown. n.s: not significant.

1148

1149   **Figure 2—figure supplement 7. Phototoxicity is not the constraint of mox expression.**

A) Growth curves of the vector control and cells overexpressing mox and mox-YG grown under room light or in the dark (Dark) in synthetic medium (–LeuUra). The solid or dotted lines show the average calculated from three biological replicates.

B) Protein level of mox and mox-YG (percent over the total protein) under room light or in the dark (Dark) in synthetic medium (–LeuUra).

C) Experimental setup for the cultivation of cells under strong light. Strong light was irradiated using a halogen lighting source (Megalight 100, HOYA-SCHOTT). Among the three cultures of cells overexpressing mox and mox-YG, culture #3 was closest to the light, followed by #2 and #1 in decreasing order. Cells were cultured in SC–LeuUra medium. Note that the light does not affect the OD_{660} measurement itself in this setup.

D) Growth curves of the cells overexpressing mox and mox-YG grown under the strong light. Growth curves of cells at the positions (#1, #2, and #3) in **C** are shown.

E) Color of the culture of cells overexpressing mox cultivated at the positions in **C**. The photo was taken after 40 hrs cultivation.

F) Fluorescence microscopy images of the cells cultivated under strong light. The cells were cultivated at the position in **C** for 40 hrs.

G) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing mox and mox-YG cultured under the light. Cells were cultured at the position in **C** to $OD_{660} = 1$ and analyzed. All proteins were separated after staining with a fluorescent dye.

H) Protein level of mox and mox-YG level in **G** (percent over the total protein).

1170

Figure 2—figure supplement 8. Effect of Tyr and oxidative stress for the expression limits of mox

and mox-YG.

A) Growth curves of cells with the vector (Vector) and overexpressing mox, mox-YG, and mox-YG

with one Tyr added to the C-terminal (mox-YG+Y) (n = 1 for each).

B) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing FPs. All proteins

were separated after staining with a fluorescent dye.

C-E) Quantification of MGR (**C**), the protein level (**D**), and the neutrality index (**E**) of cells

overexpressing mox (FP), mox-YG (Y66G), and mox-YG+Y (Y66G+Y). The MGR was calculated

from the growth curve in **A**. The protein level was calculated from the band in **B**. The neutrality index

was calculated as the product of %MGR in **C** and the %protein level in **D**. The results for n = 1 are

shown.

F, G) Growth curves of the vector control (Vector) and cells overexpressing mox and mox-YG in

synthetic medium (**F**) and medium with 2 mM H_2O_2 (**G**). Solid or dotted lines indicate mean values

calculated from three biological replicates.

H) Passing point when the time for Vector, mox, and mox-YG overexpressing cell cultures to reach

$OD_{660} = 1$ (the intersection of the dotted line in **F** and **G** with the respective growth curves).

1187

1188 **Figure 2—figure supplement 9. Overexpressed mox-YG does not show any specific localization.**

1189 Fluorescence microscopy images of the cells with the vector (Vector) and overexpressing mox and

1190 mox-YG in a strain expressing a GFP-binding protein (GBP) fused to mScarlet-I under the control of

1191 *ACT1* promoter.

1192

Figure 2—figure supplement 10. Mutational analysis of mox.

A, B) Curves representing time series data of cell growth (gray, OD_{595}) and GFP fluorescence (green)

overexpressing mox and mox-T203I mutant in the synthetic medium (–LeuUra). Solid lines are means

calculated from eight biological replicates.

C) Bar graph comparing the maximum value of GFP fluorescence obtained from the GFP fluorescence

curves shown in **A** and **B**. GFP fluorescence values for the vector control and mox-YG overexpressing

cells were also shown as controls. Average values, SD, and *p*-values of Welch's *t*-test were calculated

from eight biological replicates.

D) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing modified GFP.

All proteins were separated after staining with a fluorescent dye.

E) MGR of cells overexpressing modified moxGFP and their mutants (percent over the vector control).

F) Protein levels of the mox variants (percent over the total protein). The amount was calculated from

the intensity of the target bands separated by SDS-PAGE of **D**.

G) The neutrality index for the modified GFP. The neutrality index was calculated as the product of

the %MGR in **E** and the %protein level in **F**.

H) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing mox-T65S mutant

and mox of WT. All proteins were separated after staining with a fluorescent dye.

I) Protein levels of mox-T65S mutant and mox (percent over the total protein). The amount was

calculated from the intensity of the target bands separated by SDS-PAGE of **H**.

The bars and error bars in **E**, **F**, **G**, and **I** show the means and SDs calculated from three biological

replicates, except the T203I mutant (n = 2). The raw data is shown with dot plots.

1214

Figure 3—figure supplement 1. Comparison of the transcriptional response of mox and mox-YG

overexpressing cells.

A, B) Transcriptome changes in mox (**A**) and mox-YG (**B**) overexpressing cells. Significantly altered

ORFs (FDR < 0.05) compared to the vector control are shown in red. The number of ORFs that were

significantly increased and decreased are shown.

C) Violin plots showing transcriptome results for each category of KEGG orthology level 3. Categories

that were significantly up or down-regulated (FDR < 0.05) are indicated by red violin plots. Genes that

showed significant expression changes (FDR < 0.05) within the same category are indicated by swarm

plots. Asterisks indicate significant differences (FDR < 0.05) between mox and mox-YG comparisons

for groups of genes in the same category.

1225

Figure 3—figure supplement 2. Verification of promoter activity by the reporter assay.

A) The constructs used for promoter analysis by the transcriptional reporter included a sequence of RFP

(mScarlet-I) behind each promoter, a terminator for *TDH3*, and a sequence incorporating *spHIS5MX* as

a selection marker, which was integrated into the genomic *FCY1* locus. "moxFS" means the frameshift

mutant of mox-YG is overexpressed.

B) Curves representing time series data of cell growth (gray, OD_{595}) and mScarlet-I fluorescence (red,

arbitrary unit) overexpressing mox-YG and moxFS in SC–LeuUra medium for each promoter. Solid

lines are averages calculated from four biological replicates.

C) Quantitative results of mScarlet-I fluorescence values for each promoter. Bars and error bars indicate

the mean and SD of maximum fluorescence values (arbitrary unit) for mScarlet-I calculated from four

biological replicates. Raw data are shown as dot plots. The *p*-values were calculated by performing

Welch's *t*-test and applying the Bonferroni correction. "n.s." means $p > 0.05$.

Figure 3—figure supplement 3. Depletion of leucine in the gTOW experiment does not induce the

***GAP1* promoter.**

A) Reporter assay of the *GAP1* promoter in media with different concentrations of leucine. Curves

representing time series data of cell growth (gray, OD595) and mScarlet–I fluorescence (red, arbitrary

unit) overexpressing mox–YG. Solid lines are averages calculated from four biological replicates.

B) Quantification of the maximum growth rate (MGR) and maximum RFP fluorescence derived from

the results in A. Bar graphs indicate the mean maximum RFP fluorescence, and error bars represent the

SD. Line graphs indicate the MGR, with error bars representing SD. Means and SDs were calculated

from four biological replicates.

1248   **C**) Scatter plots of maximum fluorescence values and MGRs created using the values in **B**. Note that

1249   *GAP1* induction was observed only in mox-YG overexpression, even under conditions where the MGR

1250   was reduced to around three due to leucine depletion (the dotted line).

1251

Figure 4—figure supplement 1. Change in total protein levels upon overexpression of target

proteins.

Ratio of total protein content of cells overexpressing FPs over the vector control. Gray bars and error

bars indicate expression levels and SDs for proteins other than FPs; non-gray bars and error bars indicate

expression levels and SDs for each overexpressed protein. The mean and SD are calculated from three

biological replicates. The *p*-values were calculated by performing Welch's *t*-test and applying

Bonferroni correction. "n.s." means $p > 0.05$.

1259

Figure 4—figure supplement 2. Evaluation of expression constraint of fluorescent proteins under

nutrient–rich medium.

A) Experimental setup of the analysis. Target proteins were expressed under the control of the *TDH3*

promoter (*TDH3pro*) using the multicopy plasmid pTOW-AUR1d. This plasmid uses the aureobasidin

resistance gene *AUR1d*, which has a short promoter region, replacing the *leu2–89* used in the plasmid

shown in Figure 2A.

B) Growth curves of the vector control and cells overexpressing FPs in YPD included 500 ng/mL

aureobasidin A. The solid or dotted lines show the average calculated from three biological replicates.

Shaded zones indicate their standard deviations.

C) Gel images of SDS–PAGE–separated proteins extracted from cells overexpressing FPs. All proteins

were separated after staining with a fluorescent dye.

D) Ratio of total protein content of cells overexpressing FPs over the vector control in YPD added

aureobasidin A and SC–LeuUra. Gray bars and error bars indicate expression levels and SDs for

proteins other than; non–gray bars and error bars indicate expression levels and SDs for each

overexpressed protein. The mean and SD are calculated from three biological replicates. The *p*-values

were calculated by performing Welch's *t*-test.

1276

Figure 4—figure supplement 3. Comparison of normalized or not-normalized proteome results.

A) Bar graph showing the sum of signals for each gene detected by proteome analysis. Black or light

gray bars indicate the sum of signals other than mox-YG, and green-shaded bars indicate mox-YG

signals. The error bar indicates the SD calculated from three biological replicates.

B) Volcano plot of proteome results excluding proteins not detected in either condition.

C) Volcano plot of **B**'s results normalized so that the sum of mox-YG signals equals the sum of the

vector control signals. Significantly changed proteins (FDR < 0.05) are shown in red. The number of

genes whose expression decreased or increased during mox-YG overexpression compared to the vector

control is indicated in the upper left or upper right of the graph.

D) Comparison of proteome and transcriptome changes in cells overexpressing mox-YG.

The transcriptome data used are the same as those in Figure 3—figure supplement 1B, and the proteome

data are the same as those in Figure 4—figure supplement 3B. The Pearson correlation coefficient (*r*)

is shown.

1290

Figure 4—figure supplement 4. Violin plots showing groups of proteins whose expression was changed by mox-YG overexpression.

Protein groups with significantly altered expression (FDR < 0.05) upon mox-YG overexpression over the vector control are shown in red violin plots. Proteins with significant expression changes (FDR < 0.05) within the same category are shown by swarm plots.

1296

Figure 4—figure supplement 5. Cells overexpressing mox-YG exhibit proteomic changes that

overlap with those observed in rapamycin-treated cells.

A) Volcano plots showing genes whose expression levels were altered by rapamycin treatment in the

proteome results of mox-YG overexpressing cells.

B) Venn diagram representing the overlap of the proteome during mox-YG overexpression and

rapamycin treatment. The number of genes in each area is shown.

Published data (Gowans et al., 2018) was used for comparison with the rapamycin response genes.

1304

Figure 4—figure supplement 6. Comparison of groups of proteins whose expression levels are

changed by mox-YG overexpression and rapamycin treatment.

1305

1306

A, B) Violin plots showing protein expression changes upon mox-YG overexpression for each KEGG

ontology level 3 category. Plots with red borders represent categories that changed significantly (FDR

< 0.05). Proteins with significantly (FDR < 0.05) altered expression within a category are shown in the

swarm plot. The left side of the violin plot shows the analysis with all the data. The right side of the

plot shows the analysis with only rapamycin-responsive protein (**A**) and rapamycin-non-responsive

protein (**B**).

C) Bar graph showing the results for proteins in the "Oxidative phosphorylation" category. Proteins

whose expression is also increased during rapamycin treatment (Rap-up) are marked.

1315

Figure 4—figure supplement 7. Assessment of TORC1 pathway activity in mox-YG overexpressing cells

A, **B**) Growth curves of the vector control (**A**) or mox–YG overexpressing (**B**) cells under different rapamycin concentrations. The solid lines show the average calculated from four biological replicates.

C, **D**) Max growth rate of the vector control (**C**) or mox–YG overexpressing (**D**) cells under different

rapamycin concentrations. The bars and error bars show the means and SDs of the MGR calculated

from four biological replicates. The p–values were calculated by performing Welch's t–test.

E) Line graph showing the change in MGR with increasing rapamycin concentration, using 0 ng/mL of

rapamycin concentration as reference. Rapamycin concentrations with significant differences ($p < 0.05$)

between the vector control and mox–YG are marked with * above the mox–YG marker. The p–values

were calculated by performing Welch's t–test. In the vector control, the addition of 100 ng/mL

rapamycin caused significant growth inhibition, whereas in mox-YG overexpressing cells, the

inhibitory effect was limited. This highlights a decreased sensitivity to rapamycin in mox-YG

overexpressing cells.

F) Detection of Atg13 phosphorylation in the vector control and mox-YG overexpressing cells cultured

in SC–LeuUra medium to the indicated optical density (OD_{660}). Total protein was visualized by ponceau

S staining (left), and Atg13 was detected by Western blotting using an anti-Atg13 antibody (right).

Atg13 phosphorylated by TORC1 appears as a broad band (indicated by a line).

1334

Figure 4—figure supplement 8. Observation of mitochondria in mox-YG overexpressing cells

A) Fluorescence microscopy image showing the localization of Rip1-mScarlet-I. The vector control and

mox-YG overexpressing cells were observed in the log phase and stationary phase. The upper panel is

a bright-field (BF) image, and the lower panel is a merged image of the BF and fluorescence images.

B) Violin plots showing the quantified fluorescence values of Rip1-mScarlet-I from fluorescence

microscopy images taken in the log phase and stationary phase. The number (n) of observed cells is

shown. The p-values were calculated by Welch's t-test.

1342

1343 **Figure 4—figure supplement 9. Time-course analysis of oxygen consumption in mox-YG**

1344 **overexpressing cells.**

1345 **A**, **B**) Line graphs showing changes in the fluorescence of an oxygen-sensitive probe, measured in

1346 synthetic medium (**A**) and in medium containing 50 μM antimycin A (**B**). Solid lines represent the mean

1347 values from three biological replicates; Error bars indicate standard deviations.

1348

Figure 5—figure supplement 1. Expression changes of ribosomal proteins upon mox-YG

overexpression.

Bar graphs show the changes in the expression of ribosomal proteins in the proteome analysis of **Figure**

4. Significantly changed proteins are boxed in red. Structure names containing increased proteins are

shown. Note that when the amino acid sequences of paralogs are identical, they cannot be distinguished

by proteomic analysis, and the protein abundance of both members of the paralog pair is represented

under the name of only one.

1356

Figure 5—figure supplement 2. Microscopic analysis of the cells under mox-YG overexpression.

A) Electron microscopic images of the cells with the vector or under mox-YG overexpression. The

white rectangular area in each left image is magnified to show the cytoplasmic density (shown in each

right image).

B) Fluorescence microscopy image showing the size of nucleolus with Nsr1-EGFP. Log phase cells

(OD_{660} = 1.0) with the vector or under mox-YG overexpression were observed.

C, D) Quantification of the size of Nsr1-GFP fluorescent region (**C**) and fluorescence value of Nsr1-

GFP (**D**) in the cells with the vector or mox-YG overexpression. The number (n) of observed cells is

shown. The *p*-values were calculated by Welch's *t*-test.

1366

1367

A Vector

B mox-YG

1368

1369 **Figure 5—figure supplement 3. Electron microscopic images of the cells with the vector (A) or**

1370 **under mox-YG overexpression (B).**

1371

Figure 5—figure supplement 4. Model diagrams showing the hypothetical situation of the cells.

A) In the wild type cells, enough ribosomal proteins (RPs) are produced, resulting in RP-assembled

rRNAs and the formation of nucleolus with normal morphology.

B) In mox-YG overexpressing cells, the amount of RP is reduced due to translation competition,

increasing misassembled rRNA. As a result, degradation of rRNA by exosomes may be accelerated,

resulting in abnormal nucleolus morphology and reduction of ribosomes.

C) In exosome-deficient cells like the *mtr4-1* cells, errors in rRNA processing occur due to the reduction

of degradation by exosomes, resulting in the reduction of ribosomes and growth defects.

D) In mox-YG overexpressing, exosome-deficient cells, degradation of misassembled rRNA is reduced,

and nucleolus formation and ribosome biosynthesis are partially recovered.

1382

1383   **Figure 5—figure supplement 5. Positive genetic interaction between exosome mutation (*mtr4-1*)**

1384   **and mox-YG overexpression**

A) Growth curves of the wild type (WT) and *mtr4-1* mutant cells with the vector or under overexpression of mox-YG in SC–LeuUra medium at 30˚C. The solid or dotted lines show the average calculated from three biological replicates. Shaded zones indicate their SDs.

B) MGR of the wild type (WT) and *mtr4-1* mutant cells with the vector or under overexpression of mox-YG (percent over the vector control). The bars and error bars show the means and SDs calculated from three biological replicates. The raw data is shown with dot plots. The *p*-value of Welch's *t*-test is shown. n.s.: $p > 0.05$.

C) Gel images of SDS-PAGE-separated proteins extracted from the wild type (WT) and *mtr4-1* mutant cells with the vector or under overexpression of mox-YG. All proteins were separated after staining with a fluorescent dye.

D) Observation of nucleolus. The construct to observe nucleolus (Nsr1-mScarlet-I fusion gene) was integrated into the genomic *FCY1* locus. Fluorescence microscopy image of nucleolus-localized protein Nsr1-mScarlet-I of the WT and *mtr4-1* mutant cells with the vector or mox-YG overexpression.

E) Quantification of the size of Nsr1-mScarlet-I fluorescent region in WT and *mtr4-1* cells with the vector or mox-YG overexpression. The number (n) of observed cells is shown. The *p*-values were calculated by Welch's *t*-test.

1401

Figure 5—figure supplement 6. Comparison of growth of glycolytic enzymes overexpressing

strains.

A, **C**, **D**) Growth curves of cells with the vector or under mox-YG overexpression in SC–LeuUra

medium at the indicated temperature. In the indicated culture, sorbitol (1 M) is added. The solid or

dotted lines show the average calculated from three biological replicates. Shaded zones indicate their

standard deviations.

B) Bright-field microscopic images of cells with the vector or under mox-YG overexpression cultured

at the indicated temperature.

E) Fluorescence microscopy image showing the size of the Nsr1-mScarlet-I nucleolus, observed in the

vector control or mox-YG over-expressing cells (OD_{660} = 1.0) when cultured at 38°C or in sorbitol-

enriched medium.

F) Quantification results of the Nsr1-mScarlet-I regions observed in the eight conditions indicated in **E**.

Number of cells observed (n) is shown. The *p*-value was calculated by the Mann-Whitney *U*-test.

1415

Figure 6—figure supplement 1. Evaluation of the neutrality of Tdh3 and Gpm1.

A, **B**) Growth curves of cells with the vector (Vector) and overexpressing Gpm1 and Tdp3 and their

catalytic center mutants (CCmut) in synthetic medium (–LeuUra). The solid or dotted lines show the

average calculated from three biological replicates. Shaded zones indicate their SD.

C) MGR of cells overexpressing indicated proteins (percent over the vector control).

D) Gel images of SDS-PAGE-separated proteins extracted from cells overexpressing indicated proteins.

All proteins were separated after staining with a fluorescent dye.

E) Protein levels of the target protein (percent over the total protein). The amount was calculated from

the intensity of the target bands separated by SDS-PAGE of **D**.

F) Ratio of total protein content of cells overexpressing glycolytic enzymes over the vector control.

Gray bars and error bars indicate expression levels and SDs for proteins other than glycolytic enzymes;

non-gray bars and error bars indicate expression levels and SDs for each overexpressed protein.

The bars and error bars in **C, E**, and **F** show the means and SDs calculated from three biological

replicates. The raw data is shown with dot plots.

Gpm1

```
   1  M  P  K  L  V  L  V  R  H  G  Q  S  E  W  N  E  K  N  L  F  20
 959 ATGCCAAAGTTAGTTTTAGTTAGACACGGTCAATCCGAATGGAACGAAAAGAACTTATTC 1018

  21  T  G  W  V  D  V  K  L  S  A  K  G  Q  Q  E  A  A  R  A  G  40
1019 ACCGGTTGGGTTGATGTTAAATTGTCTGCCAAGGGTCAACAAGAAGCCGCTAGAGCCGGT 1078

  41  E  L  L  K  E  K  K  V  Y  P  D  V  L  Y  T  S  K  L  S  R  60
1079 GAATTGTTGAAGGAAAAGAAGGTCTACCCAGACGTCTTGTACACTTCCAAGTTGTCCAGA 1138

  61  A  I  Q  T  A  N  I  A  L  E  K  A  D  R  L  W  I  P  V  N  80
1139 GCTATCCAAACTGCTAACATTGCTTTGGAAAAGGCTGACAGATTATGGATTCCAGTCAAC 1198

  81  R  S  W  R  L  N  E  R  H  Y  G  D  L  Q  G  K  D  K  A  E  100
1199 AGATCCTGGAGATTGAACGAAAGACATTACGGTGACTTACAAGGTAAGGACAAGGCTGAA 1258

 101  T  L  K  K  F  G  E  E  K  F  N  T  Y  R  R  S  F  D  V  P  120
1259 ACTTTGAAGAAGTTCGGTGAAGAAAAATTCAACACCTACAGAAGATCCTTCGATGTTCCA 1318

 121  P  P  P  I  D  A  S  S  P  F  S  Q  K  G  D  E  R  Y  K  Y  140
1319 CCTCCCCCAATCGACGCTTCTTCTCCATTCTCTCAAAAGGGTGATGAAAGATACAAGTAC 1378

 141  V  D  P  N  V  L  P  E  T  E  S  L  A  L  V  I  D  R  L  L  160
1379 GTTGACCCAAATGTCTTGCCAGAAACTGAATCTTTGGCTTTGGTCATTGACAGATTGTTG 1438

 161  P  Y  W  Q  D  V  I  A  K  D  L  L  S  G  K  T  V  M  I  A  180
1439 CCATACTGGCAAGATGTCATTGCCAAGGACTTGTTGAGTGGTAAGACCGTCATGATCGCC 1498

 181  A  A  G  N  S  L  R  G  L  V  K  H  L  E  G  I  S  D  A  D  200
1499 GCTgcCGGTAACTCCTTGAGAGGTTTGGTTAAGCACTTGGAAGGTATCTCTGATGCTGAC 1558

 201  I  A  K  L  N  I  P  T  G  I  P  L  V  F  E  L  D  E  N  L  220
1559 ATTGCTAAGTTGAACATCCCAACTGGTATTCCATTGGTCTTCGAATTGGACGAAAACTTG 1618

 221  K  P  S  K  P  S  Y  Y  L  D  P  E  A  A  A  A  G  A  A  A  240
1619 AAGCCATCTAAGCCATCTTACTACTTGGACCCAGAAGCTGCCGCTGCTGGTGCCGCTGCT 1678

 241  V  A  N  Q  G  K  K  *  248
1679 GTTGCCAACCAAGGTAAGAAATAA 1702
```

Tdh3

```
   1  M  V  R  V  A  I  N  G  F  G  R  I  G  R  L  V  M  R  I  A  20
 959 ATGGTTAGAGTTGCTATTAACGGTTTCGGTAGAATCGGTAGATTGGTCATGAGAATTGCT 1018

  21  L  S  R  P  N  V  E  V  V  A  L  N  D  P  F  I  T  N  D  Y  40
1019 TTGTCTAGACCAAACGTCGAAGTTGTTGCTTTGAACGACCCATTCATCACCAACGACTAC 1078

  41  A  A  Y  M  F  K  Y  D  S  T  H  G  R  Y  A  G  E  V  S  H  60
1079 GCTGCTTACATGTTCAAGTACGACTCCACTCACGGTAGATACGCTGGTGAAGTTTCCCAC 1138

  61  D  D  K  H  I  I  V  D  G  K  K  I  A  T  Y  Q  E  R  D  P  80
1139 GATGACAAGCACATCATTGTCGATGGTAAGAAGATTGCTACTTACCAAGAAAGAGACCCA 1198

  81  A  N  L  P  W  G  S  S  N  V  D  I  A  I  D  S  T  G  V  F  100
1199 GCTAACTTGCCATGGGGTTCTTCCAACGTTGACATCGCCATTGACTCCACTGGTGTTTTC 1258

 101  K  E  L  D  T  A  Q  K  H  I  D  A  G  A  K  K  V  V  I  T  120
1259 AAGGAATTAGACACTGCTCAAAAGCACATTGACGCTGGTGCCAAGAAGGTTGTTATCACT 1318

 121  A  P  S  S  T  A  P  M  F  V  M  G  V  N  E  E  K  Y  T  S  140
1319 GCTCCATCTTCCACCGCCCCAATGTTCGTCATGGGTGTTAACGAAGAAAAATACACTTCT 1378

 141  D  L  K  I  V  S  N  A  S  S  T  T  N  C  L  A  P  L  A  K  160
1379 GACTTGAAGATTGTTTCCAACGCTTCTTcTAccACCAACTGTTTGGCTCCATTGGCCAAG 1438

 161  V  I  N  D  A  F  G  I  E  E  G  L  M  T  T  V  H  S  L  T  180
1439 GTTATCAACGATGCTTTCGGTATTGAAGAAGGTTTGATGACCACTGTCCACTCTTTGACT 1498

 181  A  T  Q  K  T  V  D  G  P  S  H  K  D  W  R  G  G  R  T  A  200
1499 GCTACTCAAAAGACTGTTGACGGTCCATCCCACAAGGACTGGAGAGGTGGTAGAACCGCT 1558

 201  S  G  N  I  I  P  S  S  T  G  A  A  K  A  V  G  K  V  L  P  220
1559 TCCGGTAACATCATCCCATCCTCCACCGGTGCTGCTAAGGCTGTCGGTAAGGTCTTGCCA 1618

 221  E  L  Q  G  K  L  T  G  M  A  F  R  V  P  T  V  D  V  S  V  240
1619 GAATTGCAAGGTAAGTTGACCGGTATGGCTTTCAGAGTCCCAACCGTCGATGTCTCCGTT 1678

 241  V  D  L  T  V  K  L  N  K  E  T  T  Y  D  E  I  K  K  V  V  260
1679 GTTGACTTGACTGTCAAGTTGAACAAGGAAACCACCTACGATGAAATCAAGAAGGTTGTT 1738

 261  K  A  A  A  E  G  K  L  K  G  V  L  G  Y  T  E  D  A  V  V  280
1739 AAGGCTGCCGCTGAAGGTAAGTTGAAGGGTGTTTTGGGTTACACCGAAGACGCTGTTGTC 1798

 281  S  S  D  F  L  G  D  S  H  S  S  I  F  D  A  S  A  G  I  Q  300
1799 TCCTCTGACTTCTTGGGTGACTCTCACTCTTCCATCTTCGATGCTTCCGCTGGTATCCAA 1858

 301  L  S  P  K  F  V  K  L  V  S  W  Y  D  N  E  Y  G  Y  S  T  320
1859 TTGTCTCCAAAGTTCGTCAAGTTGGTCTCCTGGTACGACAACGAATACGGTTACTCTACC 1918

 321  R  V  V  D  L  V  E  H  V  A  K  A  *  333
1919 AGAGTTGTCGACTTGGTTGAACACGTTGCCAAGGCTTAA 1957
```

1430

1432 The amino acid substitutions introduced to eliminate enzyme activity are shown in red. The H182A

1433 mutation in Gpm1 was referred to the study by White *et al.* (White and Fothergill-Gilmore, 1992), and

1434 the C150S mutation in Tdh3 was referred to the study by Peralta et al. (Peralta et al., 2015).

1435

Figure 6—figure supplement 3. Overexpression of Gpm1 and its CC-mutant induces abnormal

cell morphology.

A) Representative microscopic images of cells overexpressing Gpm1 or Gpm1–CCmut.

B) Violin plots showing cell size quantified from microscopic images. The number of cells used for

the quantification is shown. The *p*–values were calculated by the Mann–Whitney *U*–test.

C) Distribution of cell axis ratios upon overexpression of mox–YG, Gpm1, and Gpm1–CCmut,

compared with the vector control. The *p*–values were calculated by the Mann–Whitney *U*–test. "n.s."

means $p > 0.05$.

1444

Figure 6—figure supplement 4. Transcriptional response under mox–YG and Gpm1 CC-mutant

overexpression.

A) Curves representing time series data of cell growth (gray, OD_{595}) and mScarlet–I fluorescence (red,

arbitrary unit) overexpressing mox–YG and Gpm1–CCmut in SC–LeuUra + 50 mg/mL leucine medium

for each promoter. Solid lines are averages calculated from four biological replicates.

B) Quantitative results of mScarlet–I fluorescence values for each promoter. Bars and error bars indicate

the mean and SD of maximum fluorescence values (arbitrary unit) for mScarlet–I calculated from four

biological replicates. Raw data are shown as dot plots. The *p*–values were calculated by performing

Welch's *t*–test and applying the Bonferroni correction. "n.s." means *p* > 0.05.

Figure 6—figure supplement 5. Microscopic observation of mitochondria in cells overexpressing

Gpm1 or Gpm1-CCmut.

A) Fluorescence microscopy image showing the size of nucleolus with Rip1–mScarlet–I. Log phase

cells (OD_{660} = 1.0) with the vector or under Gpm1 or Gpm1–CCmut overexpression were observed.

B) Quantification of the fluorescence value of Rip1–mScarlet–I in the cells with the vector or Gpm1 or

Gpm1–CCmut overexpression. The number of cells used for the quantification is shown. The *p*–values

were calculated by Mann–Whitney *U* test.

1462

Figure 6—figure supplement 6. Time-course analysis of oxygen consumption in Gpm1-CCmut

overexpressing cells.

A, **B**) Line graphs showing changes in the fluorescence of an oxygen-sensitive probe, measured in

synthetic medium (**A**) and in medium containing 50 μM antimycin A (**B**). Solid lines represent the mean

values from three biological replicates; error bars indicate standard deviations.

1468

Figure 6—figure supplement 7. Microscopic observation of nucleoli in cells overexpressing Gpm1

or Gpm1-CCmut.

A) Fluorescence microscopy images showing the size of nucleolus with Nsr1–mScarlet–I. Log phase

cells (OD_{660} = 1.0) with the vector or under Gpm1 or Gpm1–CCmut overexpression were observed.

B, **C**) Quantification of the size of Nsr1–mScarlet–I fluorescent region (**B**) and fluorescence value of

Nsr1–mScarlet–I (**C**) in the cells with the vector or Gpm1 or Gpm1–CCmut overexpression. The

number of cells used for the quantification is shown. The *p*–values were calculated by Mann–Whitney

U test.