Extreme diversity of phage amplification rates and phage-antibiotic interactions revealed by PHORCE

Yuval Mulla[1,2,*,‡], Janina Müller[1,*], Denny Trimcev[1], and Tobias Bollenbach[1,3,‡]

1. Institute for Biological Physics, University of Cologne, 50931 Cologne, Germany
2. Molecular Microbiology Section, A-LIFE, AIMMS, Vrije Universiteit, 1081 BT Amsterdam, The Netherlands
3. Center for Data and Simulation Science, University of Cologne, 50931 Cologne, Germany

* These authors contributed equally.

‡ Correspondence: y.mulla@vu.nl, t.bollenbach@uni-koeln.de

Abstract

Growth rate plays a fundamental role in microbiology and serves as an important proxy for fitness in evolution. While high-throughput measurements of bacterial growth rates are easily performed in any microbiology lab, similar methods are lacking for bacteriophages. This gap hinders systematic comparisons of important phage phenotypes, such as their amplification rate in bacterial populations and their bactericidal effect, across different phages and environmental conditions. Here, we show that the amplification rate of lytic phages can be quantified by analyzing bacterial population growth and collapse dynamics under phage predation using a parsimonious mathematical model – an approach termed PHORCE (Phage-Host Observation for Rate estimation from Collapse Events). We found that the resulting phage amplification rate captures the bactericidal effect independent of initial phage and bacterial population sizes and across different growth conditions. Using high-throughput PHORCE, we found that the amplification rates of *Escherichia coli* phages vary widely by more than three orders of magnitude. Furthermore, our approach suggests that phage–antibiotic interactions are predominantly determined by the antibiotic, and not by the phage. In particular, the ribosome-inhibiting antibiotic doxycycline generally showed antagonism with phage amplification, whereas the DNA-damaging antibiotic nitrofurantoin was synergistic. This framework provides a means to quantitatively characterize phage phenotypes and may facilitate future high-throughput phage screens for antibacterial applications.

Introduction

Advances in metagenomic sequencing have revealed a staggering genomic diversity of phages[1]. A major challenge now is to understand how these diverse phages affect microbial communities, particularly through their ability to kill their bacterial hosts. While there are standardized measures of antibiotic activity, such as the minimal inhibitory concentration (MIC)[2], a similar metric for phage antibacterial activity is lacking. This is a pressing problem for potential applications of phages in therapeutics, the success of which depends on the bactericidal effect of phages[3–5]. This problem extends to the assessment of phage–antibiotic

combinations. Indeed, a recent retrospective study showed that phage-antibiotic synergy
(PAS) appears to be critical to the success of phage therapies[6]. The identification of
synergistic combinations has particular potential, but due to the lack of quantitative measures
of phage activity, no consensus quantitative definition of phage–antibiotic synergy has been
established. This situation limits a rigorous comparison between treatment options. Clearly,
new approaches to quantify phage activity, ideally in high throughput, are urgently needed.

While bacterial population size can be monitored dynamically during growth via optical
density, phages must be isolated from their hosts at each time point before quantification[7–9]
to determine their kinetic properties[10]. Genomically encoded fluorescent markers could solve
this problem, but remain challenging to engineer non-invasively in phages and are therefore
rarely used. A promising approach to circumvent this problem is to use growth curves in
phage predation assays to infer phage amplification parameters (latent period, adsorption
rate, and burst size) through mechanistic mathematical models[9,11,12]. However, these
approaches rely on time-resolved measurements of phage concentrations[9,11,12], which
severely limits throughput. In contrast, a number of empirical quantifications of phage activity
depend only on bacterial growth curves[13–17]. While phage predation experiments are
relatively straightforward, commonly used ad hoc measures (e.g. the area under the growth
curve[17]) are sensitively dependent on the initial phage and bacterial concentrations.
Consequently, cumbersome and error-prone normalization steps are required to enable
meaningful comparisons between experiments[18]. Such ad hoc measures are particularly
difficult to interpret when comparing phage activity at different bacterial growth rates[18]. As a
result, rigorous quantification of phage activity at inhibitory, and therefore clinically relevant,
antibiotic concentrations remains a challenge[19].

Here, we present PHORCE, a new approach to determine the bacterial concentration-
dependent phage amplification rate directly from bacterial growth curves. This approach
allows us to study the phenotypic diversity of lytic phages and to detect phage-antibiotic
interactions. We show that the phage amplification rate is a phage-intrinsic kinetic property
in that it is insensitive to the initial phage and bacterial concentrations. Importantly, the
approach does not require time-resolved phage concentration measurements, opening the
door to high-throughput applications. We illustrate the power of PHORCE by measuring
amplification rates for the entire BASEL (BActeriophage SElection for your Laboratory)
collection of *Escherichia coli* phages[20], revealing that they differ by more than three orders
of magnitude and correlate with phage taxonomy. Finally, we show how PHORCE allows for
a rigorous definition of phage–antibiotic interactions based on the antibiotic dependence of
the phage amplification rate.

Results

PHORCE captures phage-host dynamics

We aimed to quantitatively understand the bactericidal activity of phages in a growing
bacterial population. To use an assay that could be readily scaled up, we infected an
exponentially growing culture of *E. coli* bacteria (K-12 BW25113) with a lytic phage (Bas04,
a highly active lytic phage from a publicly available phage library[20]). The bacteria carry a
plasmid encoding luciferase genes for accurate growth rate determination (Methods). We
measured bioluminescence over time as a proxy for bacterial concentration (Fig. 1a). We

tracked population size using bioluminescence rather than optical density because both are
proportional to biomass at steady state, but bioluminescence provides a superior signal-to-
noise ratio[21]. Indeed, both read-outs report similar dynamics (Pearson's ρ=0.99, p=10[-65],
Suppl. Fig. 1). We observed that the initial growth of bacteria with and without phages is
similar, in particular exhibiting the same growth rate (Fig. 1a). This is in contrast to most
antibiotics, which rapidly decrease the bacterial growth rate at sufficient concentrations.
However, after eight hours, the bacterial population suddenly collapsed in the presence of
phage (orange), while the no-phage control (blue) maintained its exponential growth. This
sudden collapse, which has been observed in many previous studies using lytic phages[13–15],
is due to phage predation and therefore depends on the bactericidal effect of a phage: At the
time of collapse, the phage population is large enough to lyse bacteria at a higher rate than
they can grow.

Successful phage predation consists of two critical steps: First, a phage adsorbs to the
bacterial host. After a latency period required for phage replication within the host, the second
step occurs: bursting of the infected bacterium and release of progeny phage particles. To
model this process, we first investigated which of these two steps (adsorption or latency) is
rate-limiting for the infection cycle under our conditions. Importantly, the adsorption rate
depends on the bacterial concentration, whereas the latent period typically does not. We
reasoned that this distinction could be exploited to identify the rate-limiting step in our assay:
If only latency is limiting, then the collapse time should be independent of the absolute
concentrations of bacteria and phages, and only depend on their ratio (Supplementary
Information). To test this directly, we used the growth curve assay to measure collapse times
for a wide range of initial bacterial densities (Fig. 1b) while maintaining a fixed bacterium-to-
phage ratio (multiplicity of infection, MOI). Instead of a constant collapse time, we observed
a precise logarithmic decrease in collapse time with initial bacterial concentration over three
orders of magnitude (Fig. 1b, c). These results indicate that, under our conditions, adsorption
is rate-limiting for the phage infection cycle (Supplementary Information).

Trying to quantitatively explain our observations, we analyzed a minimal mathematical model
of phage predation[12] (Fig. 2a). This model consists of coupled differential equations
describing the bacterial density $b(t)$ growing at a rate r_{bac}, the phage density $p(t)$ adsorping
to bacteria at a rate k_{adsorb}. With each phage binding cycle, n_{burst} phages are created and
one bacterium disappears (Supplementary Information):

$$\frac{\mathrm{d}b}{\mathrm{d}t} = r_{\mathrm{bac}}b(t) - k_{\mathrm{adsorb}}p(t)b(t)$$

114   (Eq. 1a)

$$\frac{\mathrm{d}p}{\mathrm{d}t} = n_{\mathrm{burst}}k_{\mathrm{adsorb}}p(t)b(t)$$

115   (Eq. 1b)

116 In this regime of adsorption-limited phage predation, we derive an analytical result for the
117 collapse time t_{col} (Supplementary Information):

$$t_{\mathrm{col}} = \frac{1}{r_{\mathrm{bac}}} \log\left(1 + \frac{r_{\mathrm{bac}}}{r_{\mathrm{pha}}b_0} \log\left(\frac{p_\infty}{p_0}\right)\right)$$

118
119   (2)

Here, $r_{\text{pha}} = n_{\text{burst}} k_{\text{adsorp}}$ is the product of the adsorption constant k_{adsorb} and the burst size
n_{burst} (Supplementary Information), which plays the role of an amplification rate for phages
in the adsorption-limited regime (see below). p_0 and p_∞ denote the initial and final phage
concentrations, respectively, b_0 is the initial bacterial concentration, and r_{bac} is the bacterial
growth rate. Note that mathematically, the product of r_{pha} and the bacterial concentration
gives a specific population growth rate, analogous to the well-established bacterial growth
rate. In essence, r_{pha} quantifies how fast the phage population can grow in a bacterial
population and conveniently characterizes the double-exponential amplification of phages in
an exponentially growing bacterial population, regardless of the bacterial concentration.
Therefore, we refer to r_{pha} simply as the *phage amplification rate*. Unlike parameters
previously used to characterize phage amplification, the phage amplification rate thus
defined is independent of bacterial and phage concentrations[13–15,18].

Our result for the collapse time (Eq. 1) provides an efficient way to test whether the
mathematical model is consistent with the dynamics observed in phage predation
experiments. To this end, we next performed experiments in which we systematically varied
initial bacterial densities over two orders of magnitude (Fig. 2b,c, colors) and initial phage
densities over three orders of magnitude (Fig. 2b,c, columns) and quantified the collapse
times for the two-dimensional array of all 80 pairwise combinations. Next, we estimated the
fold increase in phage population size p_∞/p_0 from subsequent measurements of bacterial
growth curves (Ref. [9], Methods, Suppl. Fig. 2). This rich, quantitative dataset enables a
powerful test of the model's prediction of collapse time, since the phage amplification rate
remains as the sole free parameter in Eq. 1.

We found a quantitative agreement between the model and the experimental data over the
entire range of initial conditions for a single value of the phage amplification rate (Fig. 2d,
ρ=0.9). These results support that the mathematical model faithfully captures the essence of
the phage predation dynamics over several orders of magnitude of initial phage and bacterial
densities – up to the point of collapse (Suppl. Fig. 3). Post-collapse dynamics are largely
dominated by resistance evolution (Suppl. Fig. 4) and beyond the scope of this model. While
we have only tested lytic phage activity, simulations suggest that the model works similarly
well for lysogenic phages, as lysogeny predominantly affects the post-collapse dynamics
(Suppl. Fig. 4 and Ref. [9]). Furthermore, we found that the model works not only for *E. coli* lab
strains (Fig. 2d), but also for a clinical isolate of *P. aeruginosa* infected with a therapeutically
used phage (Suppl. Fig. 5). While the latent period may also contribute to the dynamics,
assuming adsorption as the sole rate-limiting step already quantitatively explains our results.
This approach also illustrates that the phage amplification rate can be readily measured
without the need for considerably more laborious time-resolved measurements of phage
concentration[9,12,22]. We have termed this approach to quantify phage amplification rates
PHORCE (Phage-Host Observation for Rate estimation from Collapse Events).

Phage amplification rates show great diversity

To test the broader applicability of PHORCE, we applied it to the entire BASEL collection of
69 different lytic *E. coli* phages[20] (Fig. 3a). These phages are ideally suited for this purpose
since they all infect the *E. coli* strain used in this study, are thought to represent natural *E. coli*
phage diversity, and are well characterized phenotypically and genetically[20]. Bacterial growth
curves for similar initial phage and bacterial densities showed that the collapse time is

strongly dependent on the phage used, as expected (Fig. 3b, Suppl. Fig. 6). We confirmed
the main observations made above with the Bas04 phage (Fig. 1-2): The collapse time
generally decreases with the initial bacterial concentration at a fixed multiplicity of infection
(Suppl. Fig. 7a), and the minimal model of phage predation in the adsorption-limited regime
faithfully captures the dynamics for the entire BASEL collection using the phage amplification
rate as the sole fit parameter (Fig. 3c). These results demonstrate that PHORCE is applicable
to diverse lytic phages and can be readily used as a method for quantifying phage
amplification rates in high-throughput assays.

Applying PHORCE to all phages in the BASEL collection revealed an enormous diversity of
phage amplification rates. Rates varied by more than three orders of magnitude among the
phages (Fig. 3d), and the true diversity is even greater because some phage amplification
rates were below the detection limit of our technique. To put this diversity into perspective,
the amplification rates of the host of these phages (i.e., different *E. coli* strains) vary by less
than an order of magnitude in common growth media, and even the growth-rate ratio between
the fastest-growing known bacterium, *Vibrio natriegens (*10 min doubling time[23]), and the
notoriously slow-growing *Mycobacterium tuberculosis (*24h doubling time[24]) is more than ten
times smaller than the observed range of phage amplification rates. We expect the diversity
of phage amplification rates to increase even further when other bacterial hosts are
considered.

We found that the phage genus partially determines the phage amplification rate. In
particular, *Vequintavirus* members amplify significantly slower than other phages (p=3×10[-4],
Fig. 3e). However, we did not find any relationship between amplification rate and other
phage characteristics, such as genome size or primary/terminal receptor (Suppl. Table 1).
Of particular interest is phage morphology, which has traditionally been used to classify
phages. In contrast to the phage genus, phage morphology had no predictive value for phage
amplification rate (Suppl. Fig. 7b). This lack of association between morphology and
phenotype further supports the plausibility of the modern phage classification based on
sequence rather than morphology.

How does the phage amplification rate compare to other measures of phage efficacy? A
common measure of phage efficacy is the total number of phages present in the culture at
the end of a phage predation experiment, e.g. determined by a plaque assay[25,26]. While this
phage yield is clearly distinct from the phage amplification rate, it is possible that in practice
these two quantities are highly correlated and measure essentially the same underlying
property. We have already shown that the phage amplification rate faithfully captures the
collapse time of bacterial populations across a wide range of initial conditions (Fig. 2d, Fig.
3c), but phage yield did not correlate at all with this key quantity (Suppl. Fig. 7c). These
results suggest that, in addition to conceptual advantages such as independence of initial
phage and bacterial densities, phage amplification rate can be a useful measure to quantify
phage efficacy for potential therapeutic applications aimed at eradicating bacterial
populations.

Phage-antibiotic interactions are largely determined by antibiotics

Since PHORCE can disentangle phage amplification rates from bacterial growth rates, we
reasoned that it could be suitable for characterizing interactions between phages and

antibiotics. Antibiotics can interfere with the ability of phages to infect bacteria; a case in point
are aminoglycosides which have recently been shown to be potent inhibitors of phage
infection[26]. Our approach should readily detect such phage-antibiotic interactions, which are
analogous to drug interactions[21]. A key difference is that drug interactions between antibiotics
are typically defined in terms of the drugs' quantitative effects on the bacterial growth rate[21].
This approach is not directly applicable to phage-antibiotic interactions, because phages do
not simply lead to exponential growth with a lower rate, but rather to more complicated
population dynamics (Fig. 1). In a plausible definition of phage–antibiotic interactions,
antagonism leads to a decrease in phage amplification rate in the presence of the antibiotic
and synergy to an increase; in a neutral interaction, the antibiotic does not affect the phage
amplification rate. The fact that the phage amplification rate is independent of initial
conditions and easy to measure is valuable for the rigorous definition and quantification of
phage-antibiotic interactions.

To test this idea, we measured phage amplification rates in the presence of antibiotics at
different concentrations. We first focused on the phage Bas04 and a fine-resolution
concentration gradient of the ribosome-inhibiting antibiotic doxycycline. We set up several
parallel precultures at different antibiotic concentrations to ensure steady state growth,
before adding phages. The phage-induced collapse time showed a clear monotonic increase
with doxycycline concentration, while the bacterial population size (measured by
luminescence intensity, Methods) at which the collapse occurred remained nearly constant
upon drug addition (Fig. 4a, blue). The phage amplification rate derived from these data
(Suppl. Eq. 6) decreased with increasing antibiotic concentration (Fig. 4b, blue), revealing a
clear antagonistic interaction between Bas04 and doxycycline. This antagonism is not a
general consequence of lowering the bacterial growth rate because the interaction of this
phage with the DNA-damaging drug nitrofurantoin is strikingly different: Here, the collapse
time remained nearly constant and the bioluminescence intensity at the time of the collapse
increased rapidly with drug concentration. The phage amplification rate increased
monotonically with the drug concentration (Fig. 4a,c, orange), revealing a clear case of
synergy between Bas04 and nitrofurantoin. These results show that antibiotics can drastically
affect phage amplification in a synergistic or antagonistic manner, which can be detected
using PHORCE.

To determine whether the observed interactions between antibiotics and phages are specific
to each phage or reflect more general effects of the drugs on phage infection, we analyzed
phage–antibiotic interactions for a diverse selection of phages. We measured the
amplification rate of nine different phages from the BASEL collection in fine-resolution
concentration gradients of nitrofurantoin and doxycycline. Similar to Bas04 above, these
experiments showed clear effects of the antibiotics on the dynamics of bacterial population
collapse for most phages (Suppl. Fig. 8). To systematically detect phage-antibiotic
interactions, we analyzed the correlation between phage amplification rate and drug
concentration for each phage-antibiotic combination (Fig. 4b). Since we do not know the
functional form (e.g., linearity) of the antibiotic-dependent phage amplification rate, we used
Kendall's τ as the interaction score, with $\tau > 0$ indicating synergy, and $\tau < 0$ antagonism,
and $p < 0.05$ as the significance threshold. The interaction score strictly identifies only the
direction (synergy or antagonism), but we found that it correlates with the strength of the
interaction (Kendall's $\tau=0.5$, $p=0.002$, Suppl. Fig. 9a).

For doxycycline, this analysis revealed three cases of neutrality (p<0.05) and no case of
synergy; the majority of interactions (6 out of 9) were antagonistic (Fig. 4d). In contrast, the
majority of phages (6 out of 9) showed synergy with nitrofurantoin while not a single case of
antagonism was detected for this drug (Fig. 4d). Thus, phages that showed any interaction
at all generally synergized with nitrofurantoin but were antagonized by doxycycline.
Additionally, the interaction scores for both drugs were not correlated across different phages
(Kendall's τ=0.1, p=0.6; Fig. 4e). Taken together, these results indicate that phage-antibiotic
interactions are primarily determined by the antibiotic and less by the phage.

Previous studies of phage-antibiotic interactions have mostly focused on synergy and often
relied on endpoint measurements of phage yield[25,26]. We tested the extent to which these
methods correlate with our approach, which is based on the dynamics of bacterial population
growth and collapse. While there is a clear difference in the bacterial growth curves between
nitrofurantoin and doxycycline for Bas04 (Fig. 4a), the phage yield decreases similarly with
increasing concentration for both antibiotics (Fig. 4c). In fact, we found no correlation
between the antibiotic-dependence of phage yield and the antibiotic-dependence of phage
amplification rate across all phages and drug concentrations for both antibiotics (Kendall's τ
=0.05, p=0.9, Suppl. Fig. 9b). Together with the lack of correlation between phage
amplification rate and phage yield across the entire BASEL collection (Suppl. Fig. 7c), these
results show that the phage amplification rate is a distinct property of phages that is largely
independent of the commonly used phage yield. While endpoint measurements of phage
yield are certainly relevant in other contexts, the phage amplification rate quantitatively
captures the bactericidal effect of phages across diverse conditions (Fig. 2d, Fig. 3c) and is
thus a key quantity relevant to phage-antibiotic synergy.

Discussion

We have shown that phage amplification under our conditions is limited by phage–bacteria
adsorption (Fig. 1), enabling characterization of phage bactericidal effect by a single
parameter, the phage amplification rate. This rate is readily measured with PHORCE and is
insensitive to the initial phage and bacterial concentrations (Fig. 2). Applying PHORCE to the
BASEL collection of diverse *E. coli* phages revealed that the phage amplification rate varies
by more than three orders of magnitude (Fig. 3). Using this parameter to quantitatively
characterize phage-antibiotic interactions, we observed that doxycycline antagonized many
phages, while nitrofurantoin often synergized with them (Fig. 4).

Similar to the specific growth rate of bacteria, which is one of the cornerstones of
microbiology, phage amplification rate provides a key phage characteristic that can be
directly compared across different phages, hosts, and growth conditions. This is difficult to
achieve with empirical quantifications[13–15,18]. While plaque size measurements have been
used to detect bactericidal effect[27], they are hard to standardize across experiments and
laboratories due to their sensitivity to initial conditions[28]. In addition, PHORCE does not
require time-resolved measurements of phage concentrations, resulting in greatly increased
throughput compared to previous mechanistic models[9,11,12]. Previous studies on phage-
antibiotic synergy have often relied on the phage yield[25,26], which does not correlate with
bactericidal effect under our conditions (Suppl. Fig. 7c, Suppl. Fig. 9b). Instead, we propose
to use the phage amplification rate as the primary measure of a phage's bactericidal effect,

similar to how the Minimal Inhibitory Concentration (MIC) is used to assess the bactericidal
effect of antibiotics.

The phage amplification rate depends on the culture conditions. For example, we see a
strong dependence on antibiotic concentration (Fig. 4), and we also expect strong
differences between, e.g., rich and minimal media or at different temperatures. This is similar
to the situation with antibiotic susceptibility testing, where the minimal inhibitory
concentration is strongly affected by growth conditions[2]. Using preclinical infection models,
it will be interesting to see which culture conditions best predict *in vivo* activity as a first step
towards mimicking patient conditions. In addition, it will be interesting to perform cross-
laboratory validation of reproducibility under similar culture conditions.

Notably, we found no correlation between the phage amplification rate and the phage yield
(Suppl. Fig. 7c). While one might intuitively expect that a higher phage amplification rate
would lead to a higher phage yield, it should be noted that a faster collapse of the bacterial
population caused by a higher phage amplification rate also implies a lower bacterial
population at the time of collapse, and thus a lower phage yield. For future work, it will be
interesting to model and experimentally verify the dependence of the phage yield on
microscopic phage kinetics. With such a model in hand, we can also begin to understand any
antibiotic-dependent phage yield.

PHORCE relies on the assumption of adsorption-limited phage dynamics, which we verified
experimentally (Fig. 1c). This assumption holds for all *E. coli* phages we tested (Suppl. Fig.
7) and also for a *P. aeruginosa* phage (Suppl. Fig. 5). However, this does not necessarily
apply to all growth conditions or all bacteria. For example, for slow-growing bacteria such as
pathogenic mycobacteria, the binding rate does not necessarily decrease, but the slower
bacterial metabolism is likely to increase the latent period. Therefore, it will be interesting in
future work to test the validity of our model for slow-growing bacteria.

PHORCE also relies on a detectable collapse of growth curves in liquid medium. However,
some phages show plaques on bacterial lawns but do not noticeably affect liquid growth
curves. In the *E. coli* phage collection we tested, this was the case for 11 out of 69 phages.
We speculate that this occurs because these phages have such a low amplification rate that
the critical phage density to cause a collapse is not reached before the bacterial culture
reaches stationary phase. For such phages, it will be interesting to develop a similar model
for extracting phage kinetic parameters from experimental data, using time-resolved plaque
growth data instead of growth curves in liquid media.

In a mathematical model of phage therapy, the key parameter predicting bacterial clearance
is the ratio of the phage clearance rate through the body to the product of the phage burst
size and the adsorption rate[29]. Our work shows how this product of burst size and adsorption
rate, which we defined as the phage amplification rate, can be readily measured in simple
laboratory experiments. The rate at which phages are cleared from the body cannot be
measured *in vitro*; future work should focus on measuring these rates *in vivo* to select phages
with the highest potential for achieving bacterial clearance.

While our model accurately captures the dynamics up to the point of collapse, it does not
capture the rate of the collapse nor does it capture the regrowth of resistant mutants (Suppl.

Fig. 3). Recently, it has been shown that the collapse rate is determined by the latent period[9],
suggesting that this parameter can also be extracted from bacterial growth curves. Finally,
resistance evolution is a major problem in phage therapy[30]. Therefore, an important future
direction is to quantitatively extract the rate of resistance evolution from the post-collapse
growth curves, so that the optimal phages can be quantitatively evaluated for both their lytic
activity and how rapidly they induce resistance.

The enormous diversity in phage amplification rates we observed (Fig. 3) has a number of
implications. Phages play a vital role as bacterial predators in various ecosystems such as
the ocean, gut, and soil, for which recent studies have extensively characterized phage-host
networks of infectivity[31–33]. Our findings highlight large variation in bactericidal effect among
phages. This implies that phage-host interactions occupy a continuum of activity, rather than
the simple binary yes–no classification often used[31–33]. This continuum of phage activity also
has potential implications for phage therapy. While some personalized phage therapies
consider quantitative measures of bactericidal effect[34], off-the-shelf cocktails typically rely on
binary measures of phage-host interactions[35–37], in part due to a lack of easy-to-use
techniques for comparing phage activity across hosts. This is a missed opportunity, as
highlighted by a strong correlation between phage amplification rate and therapeutic efficacy
recently observed in a *Drosophila melanogaster* infection model[18]. Measuring phage
amplification rates has traditionally required laborious time-resolved measurements of phage
concentration[5]. Our high-throughput approach eliminates the need for such time-consuming
measurements, using standard equipment that can be seamlessly integrated into product
development pipelines.

Phage cocktail design can be further aided by detecting phage-phage interactions. The
model can provide a quantitative null hypothesis of what a bacterial growth curve should
look like in the presence of two different phages, based on the individual phage-bacteria
growth curves, assuming no phage-phage interactions. Any deviations from the model
predictions would indicate interactions. In future work, we plan to extend the model for
multi-phage predation and perform experiments to detect phage-phage interactions, which
can then be analyzed using e.g. RNA-seq.

Using PHORCE, we observed a notable pattern: for the phage-antibiotic pairs we tested,
phage–antibiotic interactions are more strongly determined by the antibiotic rather than by
the phage itself (Fig. 4d-e). This suggests a mechanism whereby antibiotics have a broad
effect on the ability of phages to replicate inside bacterial cells. For example, doxycycline
inhibits ribosomes[21], which likely halts phage production because phages cannot hijack them
as usual[38]. This could explain the widespread antagonism observed with doxycycline (Fig. 4).
Nitrofurantoin's mechanism of action is more intricate and involves the reduction of this
prodrug to reactive intermediates that damage various cellular components; ultimately,
resulting in decreased transcription[6]. We speculate that this reduced transcription and
subsequent reduced load on the protein synthesis machinery may facilitate phage replication
by hijacking this machinery. Investigating the underlying mechanisms of phage–antibiotic
interactions is an interesting avenue for future work. As a starting point, it would be valuable
to explore whether phage–antibiotic interactions are generally determined by the drug's
mode of action, by extending the approach presented here to measure a comprehensive
phage-antibiotic interaction network for a wider range of phages and a diverse set of
antibiotics.

Our approach has applications beyond phage–antibiotic interactions and phage cocktail
design. Phages are typically classified as generalists or specialists based on their ability to
infect a wide or narrow range of hosts, respectively[40]. A fundamental question is whether
generalism is associated with a fitness cost. Measurements of phage amplification rate are
well suited to directly address such fitness tradeoffs. Since amplification rate is a key aspect
of phage fitness, we see another application of PHORCE in evolutionary biology. Phages,
with their large numbers, short generation times, and compact genomes are a popular model
system in laboratory evolution[41]. While their evolution is nowadays relatively easy to
characterize genotypically, there is a lack of quantitative methods to assess their phenotypic
evolution. PHORCE can fill this gap. Finally, the unprecedented ease with which phage
amplification rates can be measured (Suppl. Info "Requirements to determine the phage
amplification rate") opens up the possibility of systematically investigating quantitative
relationships between these amplification rates and various aspects of phage immunity[42] and
bacterial physiology[43].

Materials and Methods

Bacterial strains and bacteriophages

All collapse curve experiments were performed with *E. coli* K-12 BW25113 transformed with a kanamycin-resistance-bearing plasmid (pCS-λ) carrying luciferase genes used to determine the growth rate[21]. Previously, we have shown that this plasmid is stably maintained in the bacteria for many generations, even in the absence of the selective antibiotic[21]. The two-dimensional (2D) gradient experiments (Figs. 1-2, Suppl. Fig. 1-2) employed the Bas04 bacteriophage, sourced from the BASEL collection[20]. For the antibiotic-phage interactions (Fig. 4, Suppl. Fig. 8-9), a subset of nine phages (Bas01; Bas04; Bas10; Bas14; Bas23; Bas30; Bas37; Bas42; Bas63) was chosen to represent a wide diversity of phage amplification rates and phage families. Additionally, *E. coli* K-12 MG1655 ΔRM, an isolation strain from the BASEL collection, was used for conducting plaque assays to determine phage titers and for the generation of phage stock solutions. Purified phage stocks were stored at 4°C, whereas bacterial stocks were stored at -70°C with the addition of glycerol (17% in final solution). Both were stored in CryoPure tubes (Sarstedt, 72.377.992).

Growth media

For all collapse curve and reinoculation experiments, Lysogeny Broth (LB) Lennox medium (Sigma Aldrich, L3022) enriched with 5mM $MgSO_4$ (Sigma Aldrich, M7506) was utilized. The plaque assays employed the double agar overlay assay (DAOA) technique[7], which required an underlay of 1.5% LB agar (Sigma Aldrich, L2897) and an overlay of 0.5% LB agar, both supplemented with 5mM $MgSO_4$. In the DAOA process, bacterial overnight culture was added to the overlay to an optical density (OD) of 0.4 within the soft agar.

The antibiotics used were doxycycline (Sigma Aldrich, D9891) and nitrofurantoin (Sigma Aldrich, N7878). The doxycycline stock solution was prepared by dissolving it in DMSO (Sigma Aldrich, D8418), while nitrofurantoin was dissolved in DMF (Sigma Aldrich, D4551). Both solutions were then filter-sterilized and stored at −20°C in the dark. Aliquots were thawed at room temperature before use.

Two-dimensional gradient of initial phage and bacterial concentration

We set up phage predation assays with a two-dimensional gradient of initial bacterial and phage concentrations (Fig. 2). For the bacterial preculture, a frozen overnight culture stock was diluted 1:500 in fresh medium and incubated for 2-3 hours in a shaking incubator (Innova 44, Eppendorf New Brunswick, 30°C, 225 rpm). We use frozen stocks of overnight cultures for simplicity and because experience in our lab shows that this generally gives nearly identical results and usually increases reproducibility. Note that we also ensure that the bacterial cultures are in a steady state of exponential growth by preculturing for several hours. A gradient of phage titers was prepared in the same medium using dilutions by a factor of 2. The bacterial concentration gradient was established orthogonally, also using dilutions by a factor of 2 of the preculture, and added on top of the phage gradient in a white microtiter plate (LumiNunc F96-MicroWell, Thermo Fisher Scientific, 732-2697). The resulting final culture volume was 140 µl per well, and phage dilutions ranged from 5×10^{-3} to 10^{-7} from a phage stock concentration of 2.6×10^8 pfu/ml in ten steps and bacterial dilutions ranged from

10^{-2} to 8×10^{-5} in eight steps from a preculture of 1.4×10^8 bacteria/ml (Fig. 2). Plates were
sealed with transparent foil (TopSeal-A Plus, PerkinElmer, 6050185) and placed in a plate
reader (Synergy H1, BioTek Inc., VT, USA) at 30°C and 807 rpm, with bioluminescence
measurements taken every 5 minutes. Dual acquisition of both optical density and
bioluminescence (Fig. 1, Suppl. Fig. 1) was achieved by performing experiments in black
plates with a transparent bottom (CELLSTAR Ref. No. 655087); in this experiment, data
points were acquired every 20 minutes. In addition, the cell density of the preculture was
determined using a Coulter counter as described below.

Antibiotic concentration gradients

We set up phage predation assays in concentration gradients of two different antibiotics,
doxycycline and nitrofurantoin (Fig. 4, Suppl. Fig. 8-9). For each antibiotic, an eight-step
concentration gradient was constructed using a 1.5-fold serial dilution, with maximum
concentrations between 3 and 4 µg/ml. Precultures were inoculated in these antibiotic
concentration gradients by diluting frozen stocks between 1:1500 and 1:3000 and incubated
for 4 hours at 30°C (225 rpm for doxycycline and 807 rpm for nitrofurantoin). After this
incubation, the precultures were re-diluted (5-fold to 10-fold) in prewarmed medium
containing the respective antibiotic concentration and transferred to a white microtiter plate.
The cell density of the precultures was determined using a Coulter counter (see below). For
each phage-antibiotic interaction measurement, one of the ten tested phages was added to
each antibiotic concentration at a 10^{-5} dilution. One column of the antibiotic concentration
gradient plate served as the phage-free control. The plate was sealed with transparent foil
and incubated in a plate reader at 30°C and 807 rpm, monitoring bioluminescence every 5
minutes for 17 to 20 hours. Note that the determination of collapse time is insensitive to any
potential effect of the antibiotic on the bioluminescence signal per cell.

Phage purification from bacterial culture

For phage purification, 130 µl of a phage-bacteria mixture was combined with 130 µl of
medium and approximately 15 µl of chloroform (Sigma Aldrich, 372978) in each well of a
polypropylene 96-well deep well plate as chloroform dissolves polystyrene plates. The plate
was then subjected to vigorous shaking at approximately 1200 rpm using a Titramax 1000
platform shaker (Heidolph, 544-12200-00) for a duration of 5 to 10 minutes. Following this,
an additional 1040 µl of medium was introduced to each well, and the plate was shaken
again, this time at a lower speed of 600 rpm, for 2 minutes to ensure thorough mixing. In the
next step, the plate was centrifuged at 2580 rpm for 30 to 40 minutes, aiming to sediment
the cell lysate at the bottom of the wells. After centrifugation, a 150 µl aliquot of the
supernatant from each well was carefully transferred to a new 96-well microtiter plate. Due
to the addition of medium during the purification process, the resulting product corresponds
to a 1:10 dilution of the pure phage lysate.

Measurement of final phage concentrations

We measure final phage concentrations (Fig. 4c, Suppl. Fig. 7c) by purifying the product of
the phage predation assays (see section "Phage purification"), reinoculating it with bacteria
and comparing the collapse times with a dilution series of the respective phage stock. For
initiating the bacterial preculture, 2 ml of medium was inoculated with 4 µl of a frozen
overnight culture stock in a standard glass test tube. This mixture was then incubated in a

shaking incubator at 30°C and 225 rpm for 2-3 hours. Following incubation, the preculture
was diluted in a range from 1:60 to 1:300 with fresh medium and aliquoted to a microtiter
plate, dispensing 138.6 µl into each well. Each well then received 1.4 µl of either the phage
purification product (at a 1:10 dilution) for reinoculation or an equivalent volume of fresh
medium for control wells. To construct calibration curves, one column per phage strain was
used, starting with a phage stock dilution of 1:100 in the top well and proceeding with a 1:3
dilution series down the column. The plate was subsequently sealed and bioluminescence
was measured as described above. Because phage yield is measured after an overnight
incubation, resistant bacteria sometimes emerge. However, this does not have a major effect
on the phage yield (Suppl. Fig. 10).

Quantification of cell densities

For Coulter counter measurements, bacterial culture volumes ranging from 10 to 100 µl were
diluted in 12 ml of Isotone II diluent in a Coulter counter cuvette (Beckman Coulter, 580015).
The diluted samples were then vortexed at a medium speed. After vortexing, the samples
were transferred to a Multisizer 3 Coulter counter (Beckman Coulter), equipped with a 10 µm
aperture. The samples were allowed to stabilize for 1–2 minutes before initiating the
measurement. In each measurement, a 500 µl sample is analyzed by the device.

Spot assay

Plates for the spot assay were prepared using the Double Agar Overlay Assay (DAOA)
technique (see section "Growth media"). First, the underlay medium was microwaved until
liquefied and then cooled to 60°C, where it was held overnight. Subsequently, 38 ml of this
medium was poured into an Nunc OmniTray single-well plate (Thermo Fisher Scientific,
242811) and left to dry under a laminar flow hood for 15 minutes. For the overlay, bacterial
overnight culture was added to the overlay medium at 60°C, to achieve an OD of 0.4. Five
ml of this mixture was then layered over the solidified underlay in the omnitray plate. Finally,
phage samples were diluted in a 1:10 dilution series and 5 µl droplets from each dilution (10^{-5}
to 10^{-9}) were carefully applied to the overlay using a MINI 96 electronic 96-channel pipette
(Integra, 4803). Plaque-forming unit (PFU) counting was conducted manually for all dilutions
where plaques did not overlap.

Statistics

511 Statistical tests (Pearson, Kendall rank, and Wilcoxon signed rank) were implemented using
512 Python Scipy.stats (v1.11.1, resp. pearsonr, kendalltau and willcoxon). For multiple testing
513 correction, we used Python statsmodels (v0.14.0). Bootstrapping[44] was performed by
514 randomly resampling the time series 1,000 times using a custom Python script before
515 extracting the collapse times.

Analysis of bacterial growth curves

517 The bacterial growth rate was extracted from bioluminescence curves of bacterial cultures
518 without phages. Curves were fitted in the range of t>0.5 h and bioluminescence < 10^6 AU. A
519 first order polynomial was fitted (Levenberg-Marquardt method implemented with Python's
520 optimize.curve_fit, SciPy 1.11.1) to the bioluminescence values in log space to ensure that
521 the low concentration part of the curve also contributed to the fit. The slope of the curve was

taken as the bacterial growth rate, whereas the intersection was used to estimate the initial bacterial concentration (see below). The collapse time of phage predation curves was extracted by using Python's find_peaks (SciPy 1.11.1), using a width of four data points (20 minutes) for smoothing. Collapse points with bioluminescence values above 3×10^5 AU were excluded as they were outside of the steady state exponential growth regime. When multiple initial conditions were present of the same phage in the same growth conditions (Fig. 2d, Fig. 3c-d, Suppl. Fig. 7), Eq. 2 was fitted using optimize.curve_fit, SciPy 1.11.1 to extract the phage amplification rate from multiple growth curves. Otherwise, the phage amplification rate was calculated from a single growth curve using:

$$r_{\mathrm{pha}} = r_{\mathrm{bac}}\frac{\log(p_\infty/p_0)}{b_0\exp(t_{\mathrm{col}}r_{\mathrm{bac}}) - b_0}$$

(3)

r_{bac} was extracted from a bacterial growth curve in the absence of the phage, as described above. For phage amplification rates in the presence of an antibiotic (e.g. Fig. 4), we measured r_{bac} in the presence of the antibiotic at the relevant concentration, without any phage. b_0 was measured using a Coulter counter. We determined the ratio p_∞/p_0 in a high throughput manner inspired by Ref. [9]: we purified the phage from the experiment of interest (e.g. in the presence of antibiotic), diluted it by a factor of d (typically 60-300) and added bacteria from a fresh stock to measure the collapse time. For each phage, we create a calibration curve by measuring the collapse time at various phage stock dilutions (p_{stock}/p_0) against the same fresh bacterial stock. We fit the calibration curve with the following equation:

$$t_{\mathrm{col,stock}} = a\,\log(p_{\mathrm{stock}}/p_0) + b$$

(4)

We then extract the relative phage concentration using:

$$\frac{p_\infty}{p_{\mathrm{stock}}} = e^{(t_{\mathrm{col}}-b)/a}d$$

(5)

Lastly, by multiplying $p_\infty/p_{\mathrm{stock}}$ with the dilution factor of the first growth experiment (p_{stock}/p_0), we obtain p_∞/p_0.

The standard errors of the phage amplification rate (Fig. 1c, 4b) and the phage concentration (Fig. 4c) were determined by bootstrapping the growth curves 1000× and determining the collapse time to estimate their standard deviations. This error was then propagated to either the phage amplification rate or the relative phage stock concentration (Eq. 3 or 5, respectively).

Notably, the phage amplification rate only logarithmically depends on $p_\infty/p_{\mathrm{stock}}$. Since the experimental determination of p_∞/p_0 is relatively laborious to perform for many phages simultaneously, even with the above method, we tested whether the phage amplification rate could be approximated by assuming $p_\infty/p_0 \approx p_{\mathrm{stock}}/p_0$. We tested this for nine different phages (Suppl. Fig. 9c) and found that this assumption gives very similar results to explicitly measuring p_∞/p_0 (Pearson's ρ=0.95, p=10^{-22}). Therefore, when measuring the amplification rate of the entire BASEL collection (Fig. 3), we omitted the measurement of p_∞/p_0 and instead assumed $p_\infty/p_0 = p_{\mathrm{stock}}/p_0$.

For the measurements of antibiotic-dependent phage amplification rates (Fig. 4, Suppl. Fig. 9) we had many different initial conditions. While plate reader experiments can be performed in high throughput on 96-well plates, measuring the initial bacterial concentration (b_0) required single sample Coulter counter experiments. Therefore, we measured to what extent the initial bioluminescence intensity of growth curves (see above) could replace the Coulter counter experiments. For a limited set of replicates (1×8 concentrations of nitrofurantoin and 2×8 concentrations of doxycycline) and nine different phages, we measured the phage amplification rate by measuring b_0 with the Coulter counter and p_∞/p_0 with subsequent bacterial growth curves. We compared the amplification rate determined from both of these measured parameters (abscissa, Suppl. Fig. 9c) with the amplification rate where b_0 is semiquantitatively approximated by the extrapolated bioluminescence intensity at $t = 0$ and approximated $p_\infty/p_0 \approx p_\text{stock}/p_0$ (ordinate, Suppl. Fig. 9c). We found a clear correlation between both quantities (Pearson r=0.67, p=10[-44]) and therefore performed more replicate experiments (five per condition) using the much less laborious approximation of phage amplification rate (x-axis) for full characterization (Fig. 4b,d). It should be noted that while the Coulter counter provides a quantitative readout of bacterial concentration, bioluminescence provides only a semi-quantitative readout of bacterial concentration. However, this is sufficient for measuring antibiotic-dependent changes in amplification rate.

References

1. Dion, M. B., Oechslin, F. & Moineau, S. Phage diversity, genomics and phylogeny. *Nat. Rev. Microbiol.* **18**, 125–138 (2020).

2. Belanger, C. R. & Hancock, R. E. W. Testing physiologically relevant conditions in minimal inhibitory concentration assays. *Nat. Protoc.* **16**, 3761–3774 (2021).

3. Abedon, S. T. Automating Predictive Phage Therapy Pharmacology. *Antibiotics* **12**, 1423 (2023).

4. Abedon, S. T. How Simple Maths Can Inform Our Basic Understanding of Phage Therapy. *Clin. Infect. Dis.* **77**, S401–S406 (2023).

5. Lindberg, H. M., McKean, K. A. & Wang, I.-N. Phage fitness may help predict phage therapy efficacy. *Bacteriophage* **4**, e964081 (2014).

6. Pirnay, J.-P. *et al.* Personalized bacteriophage therapy outcomes for 100 consecutive cases: a multicentre, multinational, retrospective observational study. *Nat. Microbiol.* 1–20 (2024).

7. Kropinski, A. M., Mazzocco, A., Waddell, T. E., Lingohr, E. & Johnson, R. P. Enumeration of bacteriophages by double agar overlay plaque assay. *Methods Mol.*

Biol. Clifton NJ **501**, 69–76 (2009).

8. Dharmaraj, T. *et al.* Rapid assessment of changes in phage bioactivity using dynamic

light scattering. *PNAS Nexus* **2**, pgad406 (2023).

9. Geng, Y., Nguyen, T. V. P., Homaee, E. & Golding, I. Using bacterial population

dynamics to count phages and their lysogens. *Nat. Commun.* **15**, 7814 (2024).

10.   Kropinski, A. M. Practical Advice on the One-Step Growth Curve. *Methods Mol.*

Biol. Clifton NJ **1681**, 41–47 (2018).

11.   Nabergoj, D., Modic, P. & Podgornik, A. Effect of bacterial growth rate on

bacteriophage population growth rate. *MicrobiologyOpen* **7**, e00558 (2018).

12.   Styles, K. M., Brown, A. T. & Sagona, A. P. A Review of Using Mathematical

Modeling to Improve Our Understanding of Bacteriophage, Bacteria, and Eukaryotic

Interactions. *Front. Microbiol.* **12**, 724767 (2021).

13.   Konopacki, M., Grygorcewicz, B., Dołęgowska, B., Kordas, M. & Rakoczy, R.

PhageScore: A simple method for comparative evaluation of bacteriophages lytic

activity. *Biochem. Eng. J.* **161**, 107652 (2020).

14.   Storms, Z. J., Teel, M. R., Mercurio, K. & Sauvageau, D. The Virulence Index: A

Metric for Quantitative Analysis of Phage Virulence. *PHAGE* **1**, 27–36 (2020).

15.   Rajnovic, D., Muñoz-Berbel, X. & Mas, J. Fast phage detection and quantification:

An optical density-based approach. *PLOS ONE* **14**, e0216292 (2019).

16.   Keith, M. *et al.* Predictive phage therapy for Escherichia coli urinary tract infections:

Cocktail selection for therapy based on machine learning models. *Proc. Natl. Acad. Sci.*

121, e2313574121 (2024).

17.   Gencay, Y. E. *et al.* Engineered phage with antibacterial CRISPR–Cas selectively

reduce E. coli burden in mice. *Nat. Biotechnol.* **42**, 265–274 (2024).

18.   Blazanin, M., Vasen, E., Jolis, C. V., An, W. & Turner, P. E. Theoretical validation of

growth curves for quantifying phage-bacteria interactions. *BioRxiv* 2023.06.29.546975

(2023).

19.   Bulssico, J., Papukashvill, I., Espinosa, L., Gandon, S. & Ansaldi, M. Phage-

antibiotic synergy: Cell filamentation is a key driver of successful phage predation. *PLOS*

Pathog. **19**, e1011602 (2023).

20.   Maffei, E. *et al.* Systematic exploration of Escherichia coli phage–host interactions

with the BASEL phage collection. *PLOS Biol.* **19**, e3001424 (2021).

21.   Kavčič, B., Tkačik, G. & Bollenbach, T. Mechanisms of drug interactions between

translation-inhibiting antibiotics. *Nat. Commun.* (2020).

22.   Nabergoj, D., Modic, P. & Podgornik, A. Effect of bacterial growth rate on

bacteriophage population growth rate. *MicrobiologyOpen* **7**, e00558 (2018).

23.   Weinstock, M. T., Hesek, E. D., Wilson, C. M. & Gibson, D. G. Vibrio natriegens as

a fast-growing host for molecular biology. *Nat. Methods* **13**, 849–851 (2016).

24.   Groll, A. von, Martin, A., Portaels, F., Silva, P. E. A. da & Palomino, J. C. Growth

kinetics of Mycobacterium tuberculosis measured by quantitative resazurin reduction

assay: a tool for fitness studies. *Braz. J. Microbiol.* **41**, 300 (2010).

25.   Kebriaei, R. *et al.* Optimization of Phage-Antibiotic Combinations against

Staphylococcus aureus Biofilms. *Microbiol. Spectr.* **11**, e04918-22 (2023).

26.   Kever, L. *et al.* Aminoglycoside Antibiotics Inhibit Phage Infection by Blocking an

Early Step of the Infection Cycle. *mBio* **13**, e00783-22 (2022).

27.   Stachurska, X., Roszak, M., Jabłońska, J., Mizielińska, M. & Nawrotek, P. Double-

Layer Agar (DLA) Modifications for the First Step of the Phage-Antibiotic Synergy (PAS)

Identification. *Antibiotics* **10**, 1306 (2021).

28.   Mitarai, N., Brown, S. & Sneppen, K. Population Dynamics of Phage and Bacteria in

Spatially Structured Habitats Using Phage λ and Escherichia coli. *J. Bacteriol.* **198**,

1783–1793 (2016).

29.   Leung, C. Y. (Joey) & Weitz, J. S. Modeling the synergistic elimination of bacteria

by phage and the innate immune system. *J. Theor. Biol.* **429**, 241–252 (2017).

30.   Steen, B. A. van der, Gralka, M. & Mulla, Y. Rapid resistance evolution against

phage cocktails. *BioRxiv* (2024).

31.   Piel, D. *et al.* Phage–host coevolution in natural populations. *Nat. Microbiol.* **7**,

1075–1086 (2022).

32.   Göller, P. C. *et al.* Multi-species host range of staphylococcal phages isolated from

wastewater. *Nat. Commun.* **12**, 6965 (2021).

33.   Kauffman, K. M. *et al.* Resolving the structure of phage–bacteria interactions in the

context of natural diversity. *Nat. Commun.* **13**, 372 (2022).

34.   Gelman, D. *et al.* Clinical Phage Microbiology: a suggested framework and

recommendations for the in-vitro matching steps of phage therapy. *Lancet Microbe* **2**,

e555–e563 (2021).

35.   Lood, C., Haas, P.-J., van Noort, V. & Lavigne, R. Shopping for phages? Unpacking

design rules for therapeutic phage cocktails. *Curr. Opin. Virol.* **52**, 236–243 (2022).

36.   Guerrero-Bustamante, C. A., Dedrick, R. M., Garlena, R. A., Russell, D. A. &

Hatfull, G. F. Toward a Phage Cocktail for Tuberculosis: Susceptibility and

Tuberculocidal Action of Mycobacteriophages against Diverse Mycobacterium

tuberculosis Strains. *mBio* **12**, e00973-21 (2021).

37.   Gaborieau, B. *et al.* Prediction of strain level phage–host interactions across the

Escherichia genus using only genomic information. *Nat. Microbiol.* **9**, 2847–2861 (2024).

38.   Gerovac, M. *et al.* Phage proteins target and co-opt host ribosomes immediately

upon infection. *Nat. Microbiol.* **9**, 787–800 (2024).

39.   Dos Santos, C., Dos Santos, L. S. & Franco, O. L. Fosfomycin and nitrofurantoin:

classic antibiotics and perspectives. *J. Antibiot. (Tokyo)* **74**, 547–558 (2021).

40.   Chevallereau, A., Pons, B. J., van Houte, S. & Westra, E. R. Interactions between

bacterial and phage communities in natural environments. *Nat. Rev. Microbiol.* **20**, 49–

62 (2022).

41.   Borin, J. *et al.* Rapid bacteria-phage coevolution drives the emergence of multiscale

networks. *Science* **382**, 674–678 (2023).

42.   Millman, A. *et al.* An expanded arsenal of immune systems that protect bacteria

from phages. *Cell Host Microbe* **30**, 1556-1569.e5 (2022).

43.   Scott, M. & Hwa, T. Shaping bacterial gene expression by physiological and

proteome allocation constraints. *Nat. Rev. Microbiol.* **21**, 327–342 (2023).

44.   Beran, R. An Introduction to the Bootstrap. in *The Science of Bradley Efron:*

Selected Papers (eds. Morris, C. N. & Tibshirani, R.) 288–294 (Springer, New York, NY,

2008). doi:10.1007/978-0-387-75692-9_15.

Acknowledgments

We thank Alexander Harms for sharing the BASEL collection, Jean-Paul Pirnay for sharing
the Pseudomonas phage and Antiinfective Intelligence for sharing the *Pseudomonas* clinical
isolate; Gerrit Ansmann, Theresa Fink, Michael Lässig, and Rob Lavigne for feedback on the
manuscript; and the entire Bollenbach group for fruitful discussions. This work was supported
in part by German Research Foundation (DFG) standalone grant BO 3502/2-1 and
Collaborative Research Centre (SFB) 1310. Y.M. was supported by a Marie Curie Fellowship
from the European Union and a Humboldt research fellowship. Fig. 2a was created with
Biorender.com.

Author contributions

691   Conceptualisation: Y.M. and T.B.; Investigation: Y.M., J.M., D.T.; Mathematical model: D.T.,
692   Y.M.; Writing original draft: Y.M.; writing - review and editing: Y.M. J.M., D.T., T.B.;
693   Supervision: T.B.; Funding acquisition: T.B.

Fig. 1 Phage-induced bacterial population collapse is limited by phage-bacteria adsorption. a) Exemplary growth curve of *E. coli* in the absence and presence of a phage (Bas04), measured by bioluminescence (Methods). **b)** Growth curves measured at different initial bacterial concentrations b_0 (3×10^3 - 3×10^6 ml^{-1} from red to black; colors as in **c** at a fixed multiplicity of infection (MOI, i.e. phage-to-bacteria ratio) of 0.4. **c)** Collapse times from **b**; the observed logarithmic decrease (Pearson's ρ=0.995, p=10^{-10}) suggests that the phage-amplification kinetics is adsorption-limited (Supplementary Information). Error bars show standard errors from bootstrapping the growth curves. We obtained similar results with bioluminescence as with optical density (Suppl. Fig. 1).

Fig. 2 A mathematical model of phage infection allows the extraction of key phage amplification parameters (PHORCE). a–b) *E. coli* growth curves measured by bioluminescence (Methods) in the presence of the phage Bas04 at all combinations of eight different initial bacterial concentrations ($1.1\times10^4 - 1.4\times10^6\,\text{ml}^{-1}$, orange to blue) and ten different initial phage concentrations ($2.5\times10^3 - 1.3\times10^6\,\text{ml}^{-1}$, light to dark). Due to the large number of growth curves, only every other phage concentration is shown. Circles indicate the collapse points. **c)** Schematic of minimal model of phage amplification (Supplementary Information). We approximate the process by a single parameter: The phage amplification rate $r_{\text{pha}} = n_{\text{burst}} k_{\text{adsorb}}$. **d)** Comparison of observed collapse times and collapse times from the model. Using Eq. 2, we fit all collapse times measured in panel **a** simultaneously with a single value of r_{pha} as the only free parameter; the model is consistent with the experimental data (r^2=0.93).

Fig. 3 High-throughput PHORCE reveals an extremely wide range of phage amplification rates for a collection of diverse lytic phages. a) Schematic of 69 phages in the BASEL collection showing the phage morphology and taxonomic classification (colors). **b)** Example growth curves in the presence of different phages (colors) at the same initial phage ($1\times10^5\,\mathrm{ml}^{-1}$) and bacterial concentration ($4\times10^5\,\mathrm{ml}^{-1}$) with different collapse times. **c)** The fitted and measured collapse times for all BASEL phages, similar to Fig. 2d, show good agreement (Pearson's $\rho=0.95$, $p=10^{-129}$), regardless of the number of fitted data points (color). **d)** Distribution of the phage amplification rates for the entire BASEL collection (Methods). **e)** Vequintavirus members have a significantly lower amplification rate ($p=8\times10^{-5}$, Mann–Whitney U test with Bonferroni correction for multiple testing) than other phages. The detection limit for the phage amplification rate was $3\times10^{-10}\,\mathrm{ml\,h^{-1}}$, as slower growing phages did not cause a collapse within the exponential growth regime of the bacteria. This was the case for 7 of 12 *Vequintavirus* members and 4 of 57 others; these phages were excluded from the analysis.

Fig. 4 Phage-antibiotic interactions are determined by the antibiotic, not the phage. a)
Growth curves of bacteria and phages (Bas04) in the presence of doxycycline (left) and
nitrofurantoin (right). **b)** Phage amplification rate of Bas04 at different concentrations of
nitrofurantoin (orange) and doxycycline (blue). While the phage amplification rate increases
with nitrofurantoin (synergy), it decreases with doxycycline (antagonism). Error bars show
propagated standard errors from bootstrapping the growth curves (typically smaller than the
marker size). **c)** As **b**, but for phage yield (Methods). In contrast to the phage amplification
rate, the yield is not markedly different between the two drugs. Error bars show propagated
standard errors from bootstrapping the growth curves. **d)** Interaction score of nine different
phages with both nitrofurantoin (orange) and doxycycline (blue); interaction score is defined
as Kendall's τ of the antibiotic-dependent phage amplification rate (as in **c**). The dashed line
is at p=0.05; positive correlations indicate synergy, negative values indicate antagonism.
Nitrofurantoin shows significantly more synergistic interactions than doxycycline (p=0.008,
Wilcoxon signed rank test). **e)** Scatterplot comparing interaction scores of different phages
with nitrofurantoin to those with doxycycline. No correlation was found between the two drugs
(p=0.9, Kendalls τ), showing that the antibiotic, but not the phage, is predictive of the

749   interaction. Dashed line indicates identity. Phages in **d,e**: Bas01 (■); Bas04 (●); Bas10 (**+**);

750   Bas14 (▼); Bas23 (✘); Bas30 (★); Bas37 (♠); Bas42 (♦); Bas63 (▲).

 # Supplementary Information

 ## Model of phage amplification kinetics

 Adsorption-limited regime

Phage amplification kinetics are modeled as a two-step process: Phages must first adsorb
to bacteria at a rate k_{adsorb}, after which they cause lysis of bacteria after time τ_{latency} (latency
period). With each cycle, n_{burst} phages are created and one bacterium disappears. Note that
the burst size is a composite parameter of the product of the number of phages released
multiplied by the frequency of replicative progeny (e.g. for phages with an efficiency of plating
below 1). We first consider the adsorption-limited regime ($k_{\mathrm{adsorb}}b \ll 1/\tau_{\mathrm{latency}}k_{\mathrm{burst}}$). The
condition for the collapse time of the bacterial population is that the bacterial death rate due
to the phage is equal to the growth rate of the bacteria, which implies that the collapse time
is implicitly given by:

$$p(t_{\mathrm{col}}) = \frac{r_{\mathrm{bac}}}{k_{\mathrm{adsorb}}}.$$

(Suppl. Eq. 1)

Next, we calculate the collapse time of the bacterial population by neglecting the phage
contribution in (Main text Eq. 1a). In this approximation: $b(t) = b_0 e^{r_{\mathrm{bac}}t}$.

For a known initial bacterial concentration b_0, we can obtain the phage concentration as a
function of time by integrating (Main text Eq. 1b). In this regime, the phage population grows
super-exponentially from its initial value p_0:

$$p(t) = p_0 e^{\frac{n_{\mathrm{burst}}}{r_{\mathrm{bac}}} k_{\mathrm{adsorb}} b_0 (\exp(r_{\mathrm{bac}}t)-1)}.$$

769 (Suppl. Eq. 2)

770 We then insert in the collapse condition (Suppl. Eq. 1) and solve for t_{col}, to obtain

$$t_{\mathrm{col}} = \frac{1}{r_{\mathrm{bac}}} \log\left(1 + \frac{r_{\mathrm{bac}}}{k_{\mathrm{adsorb}}n_{\mathrm{burst}}b_0}\left(\log\left(\frac{r_{\mathrm{bac}}}{p_0\,k_{\mathrm{adsorb}}}\right)\right)\right).$$

771 (Suppl. Eq. 3)

772 We define the phage amplification rate as $r_{\mathrm{pha}} = n_{\mathrm{burst}}k_{\mathrm{adsorb}}$ and eliminate the dependence
773 on k_{adsorb} by approximating the phage concentration at the onset of collapse as
$p(t_{\mathrm{col}}) = \frac{r_{\mathrm{bac}}}{k_{\mathrm{adsorb}}} \approx p_\infty$
774 , with the final phage concentration p_∞ (phage yield). We have
775 validated this approximation experimentally (Suppl. Fig. 10). Using this approximation, we
776 obtain the collapse time in (Eq. 2):

$$t_{\mathrm{col}} = \frac{1}{r_{\mathrm{bac}}} \log\left(1 + \frac{r_{\mathrm{bac}}}{r_{\mathrm{pha}}b_0} \log\left(\frac{p_\infty}{p_0}\right)\right).$$

777 (Suppl. Eq. 4)

778 Finally, we use the measured collapse time to infer the phage amplification rate by solving
779 Suppl. Eq. 4 for r_{pha}:

$$r_{\text{pha}} = \frac{r_{\text{bac}}}{b_0 \left(e^{r_{\text{bac}} t_{\text{col}}} - 1 \right)} \log \left(\frac{p_\infty}{p_0} \right).$$

780   (Suppl. Eq. 5)

781   We find good agreement between this model and the experimental data (Fig. 2d, Fig. 3c).

Latency-limited regime

784   So far, we have considered the adsorption-limited regime where $k_{\text{adsorb}} b(t) \ll 1/\tau_{\text{latency}}$. We
785   now consider the opposite regime, where latency bursting is limiting:

$$\frac{\mathrm{d}b}{\mathrm{d}t} = r_{\text{bac}} b(t) - p(t)/\tau_{\text{latency}}$$

786   (Suppl. Eq. 6)

$$\frac{\mathrm{d}p}{\mathrm{d}t} = p(t) n_{\text{burst}}/\tau_{\text{latency}}$$

787   (Suppl. Eq. 7)

788   In this regime, the dynamics can be solved analytically, yielding

$$b(t) = b_0 e^{r_{\text{bac}} t} - \frac{p_0 \left(e^{n_{\text{burst}} t/\tau_{\text{latency}}} - e^{r_{\text{bac}} t} \right)}{n_{\text{burst}} - r_{\text{bac}}/\tau_{\text{latency}}}$$

789   (Suppl. Eq. 8)

$$p(t) = p_0 e^{n_{\text{burst}} t/\tau_{\text{latency}}}$$

790   (Suppl. Eq. 9)

792   Next, we calculate the collapse time ($\mathrm{d}b/\mathrm{d}t|_{t=t_{col}} = 0$). We find that the population collapses
793   at a fixed multiplicity of infection ($\mathrm{MOI} = p/b$):

$$p_{\text{col}}/b_{\text{col}} = \mathrm{MOI}_{\text{col}} = r_{\text{bac}} \tau_{\text{latency}}$$

794   (Suppl. Eq. 10)

796   Imposing the collapse condition we obtain for the collapse time:

$$t_{\text{col}} = \frac{\log \left(r_{\text{bac}} \tau_{\text{latency}} \left(\frac{1}{n_{\text{burst}}} + \frac{1}{\mathrm{MOI}_0} - r_{\text{bac}} \tau_{\text{latency}} \right) \right)}{\frac{n_{\text{burst}}}{\tau_{\text{latency}}} - r_{\text{bac}}}$$

(Suppl. Eq. 11)

Therefore, we find that the collapse time in the latency-limited regime depends only on the
initial MOI, and not on the absolute phage and bacterial concentrations separately. This
result is inconsistent with the experimental results in Fig. 1c and Suppl. Fig. 7a, as both show
a monotonic decrease in collapse time as a function of the initial bacterial concentration at a
fixed initial MOI. In principle, the experimental system could be in the intermediate regime
where $k_{\text{adsorb}} b(t)$ is on the order of $1/\tau_{\text{latency}}$ for at least for some duration of the experiment.
However, our experimental results indicate that this situation can only occur relatively late (if
at all), in the last few infection cycles before the collapse. The fact that the adsorption-limited
case (Suppl. Eq. 4) already shows good agreement with the experimental data (Fig. 2d, Fig.
3c) supports that it captures the essence of the dynamics up to the collapse and is thus
sufficient as a minimal model. Therefore, we do not consider more complex models with
inherently more free parameters.

Comparison of full growth curves between model and experiments

We further validated the model dynamics by comparing it with the experimentally measured
bacterial growth curves in Fig. 2. This requires a joint estimate of r_{pha} and k_{adsorb}. We obtain
an estimate of k_{adsorb} by taking advantage of the high degree of variation in the initial
conditions, noting that a high initial phage concentration must imply a fitness cost for the
bacterial population. Indeed, the ratio of the bacterial concentration at the time of collapse to
the initial bacterial concentration depends on the initial phage concentration (Suppl. Fig. 10).
We then computed r_{pha} using main text Eq. 2 for each initial condition, and took the average
over the initial conditions. Using k_{adsorb} and r_{pha} we integrated main text Eqs. 1a,b. We
integrated the dynamics using the 10$^{\text{th}}$ data point as the initial condition to avoid initial noise
in the experimental data.

Model extension including resistance and lysogeny

To investigate the effect of resistance and lysogeny on bacterial growth curves, we extended
the mathematical model (Suppl. Fig. 4). First, we described the number of bacteria sensitive
to phage infection:

$$\frac{\mathrm{d}b_{\text{sens}}}{\mathrm{d}t} = b_{\text{sens}}(t)(r_{\text{bac}} - r_{\text{res}}) - p(t)k_{\text{adsorp}}b_{\text{sens}}(t)$$

825   (Suppl. Eq. 12)

827   where r_{res} is the rate at which sensitive bacteria evolve phage resistance. For simplicity, we
828   assume that resistant bacteria grow at the same rate as sensitive bacteria and are completely
829   unaffected by the phage:

$$\frac{\mathrm{d}b_{\text{res}}}{\mathrm{d}t} = r_{\text{bac}}b_{\text{res}}(t) + r_{\text{res}}b_{\text{sens}}(t)$$

830   (Suppl. Eq. 13)

831   Finally, we included lysogeny by assuming that a fraction of the bacteria, ϕ_{lys}, is lysogenized:

$$\frac{\mathrm{d}p}{\mathrm{d}t} = p(t)n_{\text{burst}}k_{\text{adsorp}}b_{\text{sens}}(t)(1 - \phi_{\text{lys}})$$

833   The lysogenized bacteria are insensitive to further phage predation, and for simplicity we
834   assume that they grow at the same rate as sensitive bacteria:

$$\frac{\mathrm{d}b_{\text{lys}}}{\mathrm{d}t} = r_{\text{bac}}b_{\text{lys}}(t) + p(t)k_{\text{adsorp}}\phi_{\text{lys}}b_{\text{sens}}(t)$$

Note that we retrieve the limiting cases of no resistance evolution and no lysogeny with
$r_{\text{res}} = 0$ and $\phi_{\text{lys}} = 0$, respectively.

Requirements to determine the phage amplification rate

To ease the process of implementing PHORCE, we here summarize the requirements to
determine the phage amplification rate. The phage amplification rate can be measured from
a single growth curve, following Suppl. Eq. 5. One needs to measure the following quantities:

Collapse time (t_{col}): one can extract this from phage predation assay. It is important that the bacteria follow steady state growth throughout the entire experiment. Therefore, one needs to preculture the bacterial population in the relevant medium (e.g. incl. antibiotics) for at least 5 cell divisions before adding phage. The time point at which the phage is added is taken as time zero.

Bacterial growth rate r_{bac}: The bacterial growth rate can be extracted from the same curve as the collapse time. However, the curve is relatively short for phages with an early collapse time. Therefore, for each condition we measure the growth curves with phages to extract the collapse time and without phage to extract the bacterial growth rate.

Initial bacterial concentration b_0: There are two ways to estimate the initial bacterial concentration: either by quantitatively measuring the concentration or by semi-quantitatively estimating it from the growth curve. Both are detailed below.

Option 1: To quantitatively measure the bacterial concentration, a sample needs to be taken from the bacterial preculture just before adding phages. From this sample, the bacterial concentration can be measured using colony counting, a coulter counter, flow cytometry, or similar.

Option 2: The initial bacterial concentration can be semi-quantitatively estimated from the bacterial growth curves by extrapolating the bacterial growth curve (either optical density or bioluminescence) to time zero. This value can then be used as a semi-quantitative proxy for initial bacterial density. Using this approach, different phages can be compared semi-quantitatively with each other within the same growth conditions. Between growth conditions, the relationship between optical density (or especially bioluminescence) and bacterial density becomes unclear and therefore requires verification per condition (e.g. Suppl. Fig. 9c).

Ratio between the final and initial phage concentration p_{inf}/p_0: Lastly, there are three options to estimate the ratio between the final and initial phage concentrations: with plaque assay, via growth curves or by assuming the stock concentration of phages is the same as the phage concentration at the end.

Option 1: One can quantitatively measure the phage concentration of both the initial and final phage concentration using the plaque assay.

Option 2: One can use bacterial growth curves to estimate phage concentrations (Ref. [9] and Suppl. Fig. 2). There is no need to quantify the stock concentration using the plaque assay as only the ratio, and not the absolute numbers, of the final and initial phage concentration (p_{inf}/p_0) is required.

Option 3: We noticed that the final phage concentration is typically close to the phage stock concentration. Furthermore, the phage amplification rate only weakly depends on the ratio between the final and initial phage concentration. Therefore, one can approximate the ratio $p_{\text{inf}}/p_0 \approx p_{\text{stock}}/p_0$, which is the dilution factor of the stock solution when inoculating the bacterial sample. Note that this approximation likely works best when the phage stock solution was made under similar conditions (bacterial host, culturing conditions etc.) as the conditions in which the phage amplification rate is assessed.

Suppl. Fig. 1 Comparison of collapse times obtained from bioluminescence and
optical density measurements. a) Example of an *E. coli* growth curve in the absence (blue)
and presence (orange) of a phage (Bas04), as measured by optical density (Methods).
Initially there was condensation on the lid of the 96-well plate, resulting in an artifactually
higher optical density for the first half hour. **b)** The collapse time of phage Bas04 measured
with both bioluminescence and optical density for the same samples. All combinations of
seven MOIs (0.0004 – 0.3, 3× steps) and 11 bacterial concentrations (3×10^3 – 3×10^6, 2×
steps) were measured. Both methods show close agreement (Pearson's $\rho=0.99$, $p=10^{-65}$).

891
Suppl. Fig. 2 Calculation of the phage concentration from growth curves. a) Bacterial
growth curves were measured in the presence of several different dilutions (1/800, 1/1600
and 1/3200, respectively orange, pink and blue) of the Bas04 phage stock used in Figs. 1
and 2 at a fixed initial bacterial concentration (4×10^5 ml^{-1}) to quantify the phage
concentration-dependent collapse time. **b)** We fit these collapse times (t_{col}) with a logarithmic
decay (black line, see Ref. [9] for details) to obtain a calibration curve that we use to determine
unknown phage concentrations in samples containing the phage Bas04.

Suppl. Fig. 3 Comparison of full growth curves between model and experiments. Comparison between measured bacterial growth curves (black) and model trajectories (red, dotted) as a function of the different initial bacterial concentrations (1.1×10^4 – 1.1×10^6 ml^{-1}, left to right) and ten different initial phage concentrations (2.5×10^3 – 1.3×10^6 ml^{-1}, top to bottom). The procedure for calculating the model trajectories is explained in section *Comparison of full growth curves between model and experiments*. Overall, the model trajectories agree well with the measured bacterial growth curves up to collapse and capture the collapse time. The quality of the fit depends on the initial conditions: the higher the initial bacterial concentration, the further the system is from the adsorption-limited regime, and the worse the agreement with the model.

910
Suppl. Fig. 4 Resistance and lysogeny affect only post-collapse dynamics. Results of
a mathematical model including phage resistance and lysogeny (Suppl. Eqs. 12-15). When
resistance (orange dashed line) and lysogeny (green dashed line) are included, the dynamics
up to the collapse time and including the initial collapse are not noticeably affected. The curve
for resistance and lysogeny is not shown because it is indistinguishable from the curve for
lysogeny alone. For the simulations, we used an initial bacterial density of 10^5 cfu/ml, an initial
phage density of 10^5 pfu/ml, a bacterial growth rate of 1 h^{-1}, a resistance frequency of 10^{-8},
an adsorption rate of 10^{-9} cfu/ml/h, a burst size of 100, and a lysogeny frequency of 10^{-6},
which are all realistic parameters[9]. The y-axis shows the total bacterial population (sum of
the sensitive, resistant, and lysogenized populations).

Suppl. Fig. 5 PHORCE also captures the collapse time for a *Pseudomonas aeruginosa*
strain and its phage. We measured growth curves of a *Pseudomonas aeruginosa* clinical
isolate in the presence of a therapeutically used phage (Methods) to test whether PHORCE
also works for bacteria other than *E. coli* laboratory strains. **a)** Bacterial growth curves from
a two-dimensional gradient of initial bacterial densities (serially diluted by a factor of 3, from
left to right: 460 - 3.3×10^5 cfu/ml) and initial phage densities (serially diluted by a factor of 3,
from black to red: 30 - 1.7×10^6 pfu/ml). **b)** Collapse time versus initial bacterial density for
the experiment in **a**. Similar to our results for *E. coli* (Fig. 1c), the collapse time decreases
logarithmically with the bacterial density for a fixed initial bacterial-to-phage ratio (CFU/PFU
= 0.2). **c)** The PHORCE model (Eq. 1) quantitatively captures the dependence of the collapse
time on the initial bacterial and phage densities, using the phage amplification rate as the
only free parameter. The fitted phage amplification rate was 1.4×10^{-6} ml h^{-1}, which is
comparable to the strongest BASEL phage. Such a high rate is plausible because this phage
was selected for its strong activity.

Suppl. Fig. 6 Raw bacterial growth curves in the presence of the BASEL phages. Each graph shows a time-resolved bacterial growth curve, measured using bioluminescence (Methods), in the presence of a different phage. Green to black shows different initial bacterial concentrations (resp. 480, 1.44×10^4 and 4.32×10^5 ml^{-1}) at a fixed initial MOI and red to black shows different phage concentrations at a fixed initial bacterial concentration (resp. 8.1×10^5, 2.7×10^4 and 900× diluted from the phage stock solutions). The first column shows the BASEL phages 1–12, the second column shows 13–24, and so on (top to bottom).

Suppl. Fig. 7 Phage amplification characteristics of the BASEL collection. Analysis of the growth curves in Suppl. Fig. 6. **a)** The collapse time was measured at three bacterial densities, keeping the ratio of phages to bacteria (multiplicity of infection, MOI) fixed, for the entire BASEL collection. $t_{\mathrm{col},1-3}$ indicate time of collapse at resp. the lowest, middle and highest bacterial density. Each data point represents a different phage. All data points below the two solid lines show a monotonic increase in collapse time as a function of the bacterial concentration. The dashed line indicates where the collapse time increases with the logarithm of the bacterial concentration. The vast majority of data points lie below the two solid lines and cluster around the dashed line, suggesting that the amplification kinetics of all BASEL phages are predominantly adsorption-limited. **b)** Phage amplification rates for different phage morphotypes, 3/6 podoviruses, 0/34 siphoviruses and 8/29 myoviruses had a phage amplification rate below the detection limit (3×10^{-10} ml h^{-1}). None of the morphotypes differ significantly from the others (p=0.08, Kruskal-Wallis) and any differences become even more insignificant if the vequintavirus members (Fig. 3e) are excluded from the myoviruses (p=0.87, Kruskal-Wallis). **c)** Comparison between the overnight yield (y-axis) and the phage amplification rate (x-axis) for each phage in the BASEL collection shows no significant correlation (τ=-0.04, p=0.6, Kendall rank correlation) **d)** Comparison between the phage amplification rate (x-axis) and its approximation by assuming $p_\infty/p_0 = p_{\mathrm{stock}}/p_0$ (y-axis). Each data point shows the phage amplification rate determined from a single amplification curve (nine phages, three replicates per condition). Both results are in close agreement (Pearson's ρ=0.95, p=10^{-22}).

Suppl. Fig. 8 Bacterial growth curves in the presence of antibiotics and phages.
Bioluminescence as a function of time. Each row shows a different phage from the BASEL
collection (phages from top to bottom: Bas01, Bas04, Bas10, Bas14. Bas17, Bas23, Bas30,
Bas37, Bas42, Bas55, Bas63), each column shows a different antibiotic (left: nitrofurantoin,
orange; right, doxycycline, blue) at different concentrations (0.02 - 0.2 × MIC from dark to
colored, where black is no antibiotic).

Suppl. Fig. 9 Phage-antibiotic interactions. a) The degree of correlation between the phage amplification rate and the antibiotic concentration was assessed using either a Kendall correlation test (x-axis) or a Pearson correlation test (y-axis). Both scores are significantly correlated with each other (Kendall's τ =0.5, p=0.002). **b)** No correlation was found between the interaction score based on the phage yield and the interaction score based on the amplification rate (p=0.6, Kendall rank correlation). **c)** The phage amplification rate was calculated either by explicitly measuring the initial bacterial concentration and the final phage concentration (x-axis) or by approximating the initial bacterial concentration with the initial bioluminescence value and assuming $p_\infty/p_0 = p_{stock}/p_0$ (y-axis) (Methods). The explicit and approximate approaches yield similar results (Pearson's ρ=0.67, p=10^{-44}). On both axes, the antibiotic-dependent phage amplification rate was normalized to the phage amplification rate in the absence of the drug. For both doxycycline and nitrofurantoin, nine phages and eight drug concentrations were measured, respectively.

Suppl. Fig. 10 Dependence of phage yield on time of measurement. Results of an experiment with a two-dimensional gradient of initial phage (Bas14) and bacterial concentration, similar to Fig. 2. **a)** Bacterial growth curves in the two-dimensional gradient. Different columns show different initial bacterial densities (left to right: 1:100 - 1:12 800 dilutions from a 1:100 diluted overnight culture pre-cultured for 2 h using 7 steps of 5-fold serial dilutions); different colors show different initial phage concentrations (black to red: 0.001 - 6.25×10⁻⁵ relative to the stock concentration using 4 steps of 5-fold serial dilutions). **b)** Phage yield versus collapse time. We took phage samples at t = 8 h (blue) and at t = 21 h (orange) and measured the relative phage concentration by reinoculation (Suppl. Fig. 2). At these time points, the bioluminescence drops slightly as the cultures are removed from the plate reader and therefore cool down slightly before being returned to the temperature-controlled plate reader. Since the samples have different initial conditions, these yields (blue data points) correspond to different time points relative to the collapse time: the x-axis shows the time of the first phage concentration measurement (t = 8 h) minus the collapse time. **c)** Ratio of phage amplification rate calculated from the phage yields measured after 8 h and after 21 h.

1004

1005   **Suppl. Table 1 BASEL collection metadata.** The table lists all details regarding the
1006   taxonomic classification, isolation/source, host receptors, and genomic features of all *E. coli*
1007   phages used in this study. The data were taken from Ref. [20] and complemented with the
1008   phage amplification rates and phage stock concentrations characterized in this work.