A Constitutive Heterochromatic Region Shapes Genome Organization and Impacts Gene Expression in *Neurospora crassa*

Andrew T. Reckard[1], Abhishek Pandeya[2]*, Jacob M. Voris[1]*, Carlos G. Gonzalez Cruz[1],

Oluwatosin Oluwadare[2], and Andrew D. Klocko[1‡]

[1]Department of Chemistry & Biochemistry, University of Colorado Colorado Springs, Colorado Springs, CO 80918, USA

[2]Department of Computer Science, University of Colorado Colorado Springs, Colorado Springs, CO 80918, USA

*Equal Contribution

[‡]Corresponding Author: Andrew D. Klocko, email: aklocko@uccs.edu

Author Contributions: A.T.R. and A.D.K. designed research; A.T.R., A.P., J.V., C.G.C, O.O. and A.D.K. performed research; A.T.R., A.P., J.V., C.G.C., O.O. and A.D.K. analyzed data; and A.T.R., A.P., J.V., C.G.C., O.O. and A.D.K. wrote the paper.

Abstract

Genome organization is essential for proper function, including gene expression. In metazoan

genome organization, chromatin loops and Topologically Associated Domains (TADs) facilitate

local gene clustering, while chromosomes form distinct nuclear territories characterized by

compartmentalization of silent heterochromatin at the nuclear periphery and active euchromatin in

the nucleus center. A similar hierarchical organization occurs in the fungus *Neurospora crassa*

where its seven chromosomes form a Rabl conformation, where heterochromatic centromeres and

telomeres independently cluster at the nuclear membrane, while interspersed heterochromatic loci

in Neurospora aggregate across megabases of linear genomic distance for forming TAD-like

structures. However, the role of individual heterochromatic loci in normal genome organization

and function is unknown. Here, we examined the genome organization of a Neurospora strain

harboring a ~47.4 kilobase facultative (temporarily silent) heterochromatic region deletion, as well

as the genome organization of a strain deleted of a 110.6 kilobase permanently silent constitutive

heterochromatic region. While the facultative heterochromatin deletion had little effect on local

chromatin structure, the constitutive heterochromatin deletion altered local TAD-like structures,

gene expression, and the predicted 3D genome structure by qualitatively repositioning genes into

the nucleus center. Our work elucidates the role of individual heterochromatic regions for genome

organization and function.

Keywords: genome organization, chromosome conformation, heterochromatin, *Neurospora*

crassa, gene expression

Introduction

The correct organization of a genome, including proper chromosome folding, is necessary for the function of DNA-templated processes, including gene expression regulation (1–5). Indeed, improper DNA folding altering transcriptional patterns occurs in human cancer cells (6–8). Eukaryotic genomes are hierarchically folded in nuclei, where DNA molecules are wrapped about histone octamers to form chromatin fibers that are extruded into globules ("loops") by cohesin and compacted into Topologically Associated Domains (TADs) (9–17). TADs of similar transcriptional activity are then compartmentalized into the nucleus (10, 11, 17). Specifically, the more open, gene-rich, and actively transcribed euchromatin primarily localizes to the nucleus center, while the silent, often gene-poor, and compacted heterochromatin is confined to the nuclear periphery (18–20). Heterochromatin segregation may drive chromatin compartmentalization, although transcriptional activity might force silent genomic loci to the nuclear periphery (19). In metazoans, the formation of individual chromosome territories typified by reduced inter-chromosomal contacts leads to the clustering of intra-chromosomal heterochromatic regions (21, 22). In contrast, the independent aggregation of the heterochromatic centromeres and telomeres into the Rabl conformation promotes extensive intra- and inter-chromosomal heterochromatic contacts across fungal chromosomes (9, 23–25). All told, heterochromatin bundling is a conserved feature of eukaryote genome organization.

Fungi are excellent models for studying genome organization. Resembling metazoans, fungal genomes are partitioned into euchromatin and heterochromatin, and the heterochromatin of many filamentous fungi, including *Neurospora crassa*, is subdivided into permanently silent constitutive heterochromatin and temporarily silent facultative heterochromatin (26–28). The adenine and thymine (AT)-rich, gene-poor, and repetitive constitutive heterochromatic regions, comprising

~16% of the Neurospora genome, are marked by the tri-methylation of lysine 9 on histone H3

(H3K9me3) that is catalyzed by the histone methyltransferase DIM-5 (KMT1) (27, 29–31).

Specific H3K9me3-marked regions found within the seven Neurospora chromosomes ("Linkage

Groups" [LGs]) include the regional centromeres, the 14 telomeres, and ~200 interspersed

heterochromatic regions (27, 31). Neurospora facultative heterochromatin is delineated by the

subtelomeric di- or tri-methylation of lysine 27 on histone H3 (H3K27me2/3), which is catalyzed

by the SET-7 (KMT6) histone methyltransferase of the Polycomb Repressive Complex 2 (PRC2)

(32–34). In the Neurospora three-dimensional (3D) genome organization, as assessed by in nucleus

chromosome conformation capture coupled with high-throughput sequencing (*in situ* Hi-C), robust

intra- and inter-chromosomal heterochromatic interactions are readily observed across megabases

of linear genomic distance with few heterochromatin-euchromatin contacts, highlighting the

compartmentalization of fungal chromatin (9, 23, 24). The clustering of heterochromatic regions

forms the anchors of TAD-like structures; similar structures are observed in other ascomycete

fungi, suggesting the use of heterochromatin to organize fungal genomic DNA may be conserved

(9, 23–25, 35). Further, subtelomeric facultative heterochromatin enriched with H3K27me2/3

mediates the interaction between chromosome ends and the nuclear periphery (36). Together, both

heterochromatin types are necessary to organize the Neurospora genome into the Rabl

chromosome conformation.

To understand the importance of individual facultative and constitutive heterochromatic

regions on 3D genome organization in Neurospora, we performed chromatin-specific *in situ* Hi-C

on a strain harboring a deletion of a ~47.4 kilobase (kb) H3K27me2/3 domain on the LG VI left

arm (33), as well as a strain in which we deleted a ~110 kb H3K9me3 domain on the LG II right

arm. While the H3K27me2/3 deletion minimally impacts genome organization, the H3K9me3

deletion ("ΔLGIIK9het25") reorganizes the heterochromatin bundle and alters regional TADs

across LG II without affecting deposition of H3K9me3 on nearby AT-rich loci. The 3D structure

prediction of individual ΔLGIIK9het25 strain chromosomes is qualitatively different than those of

its wild type (WT) counterpart, and numerous genes became upregulated in the ΔLGIIK9het25

strain, possibly due to altered gene positioning within the 3D genome. We conclude that a

constitutive heterochromatic region organizes the *Neurospora crassa* genome for proper fungal

genome function.

Results

Loss of a H3K27me2/3-enriched region minimally affects the local chromatin landscape.

Since interactions between regions of similar heterochromatin type comprise many of the

strongest intra-chromosomal contacts in the Neurospora genome (23, 24), we examined how

deletions of either type, facultative or constitutive heterochromatin, could impact fungal

heterochromatin clustering. Starting with facultative heterochromatin, we examined a previously

constructed strain, N4933, originally used to assess position-dependent signals for H3K27me2/3

deposition in Neurospora genome (33). N4933 harbors a 47.4 kb deletion of LG VIL, from

basepairs 175,121 to 222,557, which removes fourteen H3K27me2/3-enriched genes ("ΔK27").

To assess how ΔK27 impacts genome organization, we performed two *in situ* Hi-C replicates with

*Dpn*II to capture gene-rich chromatin interactions within DNA of higher guanine and cytosine

basepair percentages, including those in facultative heterochromatic regions, in the nucleus. Our

replicates show highly similar Hi-C contact matrices across LG VI and are highly correlated

(Figure S1), allowing replicate merging into a single Hi-C dataset with 9.2 million (M) valid reads.

For comparison, we processed an identical number of valid Hi-C reads from our WT *in situ Dpn*II

Hi-C dataset (23). Our initial observation is that, despite having nearly identical valid reads, the

ΔK27 dataset has a slight bias in capturing more local contacts, most likely due to Hi-C experiment

variability, which concomitantly reduces inter-chromosomal contacts. Indeed, decreased inter-

centromeric contacts are observed in raw datasets due to fewer centromeric *Dpn*II sites (Figures 1,

S2-S3). Inter-centromeric interactions are present upon Knight-Ruiz (KR) (37) correction (Figure

S3), indicating capture of biologically relevant structures. Upon directly comparing WT and ΔK27

contact matrices, the decrease in moderate and long-range contacts is evident (Figure S3).

Therefore, we focused our analyses on the local LG VIL changes that manifest with the

H3K27me2/3 region deletion.

On WT LG VI, we observe consistent off-diagonal contacts that inversely decrease with

genomic distance, in addition to strong inter-telomeric contacts and the insulation of centromeric

DNA from local euchromatin, in both raw and KR corrected heatmaps (Figure 1A). A higher

resolution (5kb) heatmap of the 500 kb surrounding the region deleted in N4933 shows a gradual

decrease of contacts from the diagonal (Figure 1A). In contrast, the ΔK27 deletion Hi-C, mapped

to the Neurospora reference genome version 14 ("nc14") (23), appears as a white cross emanating

from the deletion locus indicating a gap in mappable Hi-C reads (Figure 1B), since the DNA in

this region was not present for nuclear ligation capture. This gap occurs in both the raw and KR

corrected contact probability heatmaps, although the base of the diagonal on the KR corrected

heatmap has some signal (Figures 1B-C), perhaps due to correctional bias. Importantly, stronger

off diagonal signal at the boundary intersections of the facultative heterochromatin deletion appear

(Figures 1B-C, black arrowheads), showing how new interactions occur between the

H3K27me2/3-enriched chromatin flanking the ΔK27 deletion. Interestingly, another N4933 locus

devoid of Hi-C signal is observed over the *sly1-1* transposase gene NCU09969 (Figure 1C, gray

arrowhead), suggesting the ΔK27 strain had a transposition event. Regarding telomere clustering,

the ΔK27 strain is still capable of forming intra- and inter-chromosomal subtelomeric interactions

(Figures 1D-E), with strong contacts between chromosome ends originating from the terminal

~100kb chromatin enriched with H3K27me2/3. We conclude a deletion of a H3K27me2/3-

enriched region minimally impacts the organization of the remaining LGVIL facultative

heterochromatin.

Deletion of a ~110 kb constitutive heterochromatic region alters silent region clustering.

In Neurospora nuclei, gene-poor AT-rich constitutive heterochromatin is compartmentalized

from gene-rich euchromatin or facultative heterochromatin (23, 38). To ask if altered constitutive

heterochromatin composition could have consequences on genome organization and function, we

deleted a 110,609 bp H3K9me3-enriched region on LG IIR, aptly called "LGIIK9het25" as this

region is the 25th constitutive heterochromatic region from the LG II left telomere. We chose

LGIIK9het25 for deletion for a high probability of observing changes to genome organization by

deleting a large AT-rich region without compromising chromosome function by removing the

centromere. Using split-marker homologous recombination in a D*mus-52* strain (39), we replaced

the 110.6 kb LGIIK9het25 with a hygromycin resistance gene (hygromycin phosphotransferase;

hph) controlled by a strong constitutive *trpC* promoter (a P$_{trpC}$*::hph* construct; Figure S4A). We

note a similar-sized DNA replacement could have more accurately represented the LG II

chromosome length, but we wanted to avoid cryptic promoters, repetitive sequences for RIP (27,

40, 41), or AT-rich DNA sequences forming novel constitutive heterochromatic regions within the

inserted DNA. We successfully obtained hygromycin-resistant primary transformants containing

the LGIIK9het25 deletion (DLGIIK9het25); in sum, we removed 109,208 bp from the Neurospora

genome, given that $P_{trpC}::hph$ inserts 1,401 bp. A hygromycin-resistant progeny from a sexual

cross, xKL9AR-17, was characterized by PCR analysis as harboring both the DLGIIK9het25 allele

(Figure S4B) and a wild type $mus-52^+$ allele. The final strain, NKL2 (genotype: mat A;

D$LGIIK9het25::hph$), had no noticeable growth defect from a WT strain (Figure S4C-D).

We performed chromatin-specific Hi-C (23) using the restriction enzyme DpnII to monitor the

contacts of gene-rich euchromatin, while MseI Hi-C captures interactions between AT-rich, gene-

poor constitutive heterochromatic regions. Our DpnII and MseI DLGIIK9het25 Hi-C replicates

show highly reproducible and correlated interactions (Figure S5), allowing us to merge replicates

into enzyme-specific datasets, with the NKL2 DpnII dataset containing 20.6M valid reads and the

NKL2 MseI dataset comprising 12.2M valid reads. All Hi-C datasets display the inter-

chromosomal centromeric interactions independent of telomere clusters, indicating our Hi-C

captured biologically relevant structures (Figures S6-S7). As before, we selected nearly identical

valid read numbers from previously published WT DpnII and MseI datasets for comparisons (23).

Compared to a WT DpnII Hi-C dataset (Figure 2A), the deleted LGIIK9het25 region appears

as a large white "cross" devoid of Hi-C interactions on LGII in the NKL2 DpnII contact probability

matrix (Figure 2B). Looking specifically at a 500 kb region surrounding LGIIK9het25, the WT

dataset has extensive interactions within and around LGIIK9het25, while the NKL2 has contact

probability loss, as no interactions are observed in the raw interaction heatmap and sparse contacts

at the diagonal in the KR corrected dataset (Figure 2A-B), which may be due to a few reads

mapping to repetitive DNA sequences at this locus. The Hi-C contacts surrounding DLGIIK9het25

appear slightly enhanced, as the flanking euchromatin moderately gains contacts in the NKL2

DpnII dataset, relative to a WT dataset (Figure 2A-B). Similar contact probability changes occur

in WT and NKL2 MseI datasets, where in the WT dataset the LGIIK9het25 region extensively

contacts internal heterochromatin yet is isolated from surrounding euchromatin, but the NKL2

dataset completely loses heterochromatic contacts from the LGIIK9het25 region yet gains flanking

euchromatin interactions (Figure 2C-D). Direct comparison of the WT and NKL2 *Dpn*II and *Mse*I

datasets surrounding LGIIK9het25 shows the contact probability gain between flanking

euchromatic regions when a separating heterochromatic region is lost (Figure 2E-F, arrowheads).

This surrounding euchromatin may gain local, on-diagonal interactions, as shown in the *Mse*I

comparison (Figure 2F), possibly due to either reduced heterochromatic interactions or a local

contact capture bias in the NKL2 dataset. Between LG I and LG II, the LGIIK9het25 deletion

causes a loss of inter-heterochromatic region interactions, including from the largest non-

centromeric heterochromatic region on LG I (Figure 2G, white arrowhead), while the LG II

centromere loses inter-centromeric contacts but modestly gains interactions with interspersed

heterochromatic regions (Figure 2G, black arrowhead). Similar gains in inter-heterochromatic

region interactions are evident across the genome in WT vs. NKL2 *Mse*I Hi-C dataset comparisons

(Figure S8), possibly from the absence of LGIIK9het25 contacts influencing euchromatin

interactions. As evident in WT vs. NKL2 *Dpn*II comparisons, the inter-chromosomal contact

strength between centromere-proximal euchromatin and more-distant euchromatin on

chromosome arms is increased, yet chromosome-internal euchromatic contacts are reduced (Figure

S9), possibly due to enhanced centromere-interspersed heterochromatin bundling. Similarly, both

DLGIIK9het25 Hi-C datasets have depleted inter-centromeric contacts; KR correction reduces

*Dpn*II contact biases (Figures S8-S9). All told, the LGIIK9het25 deletion moderately impacts the

organization of the entire Neurospora genome.

The LGIIK9het25 deletion minimally alters H3K9me3 deposition but alters TAD-like structures.

Upon deleting LGIIK9het25, the possibility exists that genome function, specifically

H3K9me3 deposition on nearby AT-rich loci, may be impacted. To assess if constitutive

heterochromatic region loss limits histone post-translational modification enrichment, we

performed H3K9me3 Chromatin Immunoprecipitation-sequencing (ChIP-seq) on the NKL2

strain. Our two H3K9me3 ChIP-seq replicates were highly reproducible (Figure S10) and were

merged into a single Reads Per Kilobase per Million (RPKM) normalized file for direct

comparison to previously published WT H3K9me3 ChIP-seq datasets, although using spike-in

DNA controls may be best practice for assessing strain differences (42, 43). We mapped the ChIP-

seq reads to either the nc14 WT genome or an "NKL2 reference genome" which replaced the AT-

rich DNA of LGIIK9het25 with the $P_{trpC}::hph$ sequence; enhanced images show low levels of

background signal across the $P_{trpC}::hph$ sequence in an NKL2 H3K9me3 ChIP-seq dataset that are

absent in WT H3K9me3 data (Figure S4E).

H3K9me3 ChIP-seq enrichment exhibits similar patterns across LG II, independent of the

reference genome used for mapping reads. For ChIP-seq data mapped to the WT nc14 genome,

identical H3K9me3-enriched regions are observed, except peaks to the right of LGIIK9het25 are

shifted by ~110kb from the deletion site (Figure 3A), given that genomic elements positions within

bedgraph/bigwig enrichment count files are shifted based on input bam file read mapping position.

Qualitatively, the NKL2 H3K9me3 peaks are higher (Figure 3A), possibly reflecting increased

H3K9me3 purification from the remaining constitutive heterochromatic regions or enhanced

sequencing depth limiting background signal. When H3K9me3 ChIP-seq data from both strains

was mapped to the NKL2 reference genome, the loss of LGIIK9het25 signal from the WT dataset

is apparent, while the remaining constitutive heterochromatic regions have identical LG II genomic

positions (Figure 3B).

To further assess local chromatin changes with the LGIIK9het25 deletion, we examined

Topologically Associated Domain (TAD)-like structure positions on LG IIR. The TAD-like

structures in WT and NKL2 *Dpn*II datasets, mapped to the nc14 (WT) genome, show striking

differences across euchromatin, with larger TAD-like structures predicted with the ΔLGIIK9het25

allele, as opposed to the smaller TAD-like structures in WT datasets (Figure 4A), although TAD

structure prediction can vary (44). To more clearly understand the strain-specific chromatin folding

changes without the ΔLGIIK9het25 signal gap, we predicted TAD-like structures WT and NKL2

*Dpn*II and *Mse*I datasets mapped to the NKL2 reference genome. Here, clear changes to TAD-like

structures are observed at the deletion site, with the euchromatin in the *Dpn*II dataset having larger

TAD-like structures in the NKL2 strain, while the NKL2 ΔLGIIK9het25 *Mse*I strain has smaller

TAD-like structures on the diagonal (Figure 4B, black arrowheads), with this difference likely

stemming from the chromatin monitored – and the density of restriction sites – in each Hi-C

experiment. The flanking TAD-like structures are nearly identical in both datasets, except for two

larger TADs-like structures in the NKL2 *Mse*I data (Figure 4B, open arrowheads). Interestingly, at

the $P_{trpC}::hph$ insertion, two novel yet small TAD-like structures are observed in both NKL2

datasets, combined with additional stronger Hi-C interactions in *Mse*I data (Figure 4B, purple line),

suggesting the constitutive P_{trpC} promoter could influence local contacts – and possibly regulate

expression – of nearby genes. Overall, the deletion of the constitutive heterochromatic region

LGIIK9het25 alters the local folding of the euchromatin surrounding the deletion site.

The LGIIK9het25 deletion alters the three-dimensional folding of Neurospora chromosomes.

It remained possible that a large constitutive heterochromatic region deletion would alter

chromosome folding. To assess the 3D folding of LG II, we computationally predicted the LG II

3D structure using WT and NKL2 ΔLGIIK9het25 Hi-C datasets. We chose to merge all *Dpn*II and

*Mse*I data together from each strain to maximize the possible Hi-C contacts across their genomes;

we did not combine *Dpn*II and *Mse*I data at ratios equivalent to the Neurospora chromatin

distribution, as previously performed (23). With the maximal contact probability merged matrices,

we iteratively generated LG II 3D structures using 3DMax (45) customized for biologically

relevant structures in fungal genomes.

In the WT strain, LG II has an elongated structure with condensed chromosome arms, a more

isolated centromere, telomeres localized to a single "side", and extensive intervening chromatin

loops (Figure 5A). Similar 3D structure prediction changes are observed on the other six WT

chromosomes (Figure S11). The LGIIK9het25 region forms a loop and associates with nearby

chromatin loops (Figure 5A), consistent with the presence of nine, moderately sized

heterochromatic regions in the terminal 800 kb of the LG II right arm (Figure 3A). In contrast, the

chromosome arms within the LG II 3D structure when LGIIK9het25 is deleted are more compact

and form a wider structure, despite similar centromere isolation, telomere locations, and looping

of intervening chromatin (Figure 5B). In fact, the terminal right arm of NKL2 LG II does not form

any loops in this 3D structure prediction, suggesting that the requirement for telomere clustering

impacts the chromosome folding prediction in the NKL2 strain (Figure 5B). Similar folding

differences occur on each LG in NKL2, with some chromosomes - specifically LG I and LG IV -

having drastically altered 3D structures relative to WT LG folding (Figures S11, S12). Together,

the deletion of a ~110kb constitutive heterochromatic region modifies the predicted 3D structures

of Neurospora LGs.

The LGIIK9het25 deletion increases gene expression due to altered 3D genome folding.

To understand if constitutive heterochromatin could influence genome function through gene

expression, we assessed changes in polyadenylation messenger RNA sequencing (polyA mRNA-

seq) in WT and NKL2 ΔLGIIK9het25 strains. As a control for complete loss of constitutive

heterochromatin function, we examined gene expression changes in a Δ*dim-5* strain, which is

devoid of H3K9me3 genome-wide (29). Volcano plots that display the significant gene expression

changes ($log_2 > 3.0$ or < -3.0; adjusted p-value < 0.001) in the NKL2 strain relative to the WT

strain show 84 genes are significantly upregulated and no genes significantly downregulated,

(Figure 6A, left; Supplementary File S1), consistent with the removal of a repressive

heterochromatic region; additional genes could be considered differentially regulated with less

stringent cutoff values. In contrast, the Δ*dim-5* strain has 175 differentially regulated genes

compared to WT, with genes having both increased and decreased transcription (Figure 6A, right;

Supplementary File S2), including the strong reduction of the gene NCU04402 encoding DIM-5

(Figure 6B), highlighting our polyA mRNA-seq dataset accuracy. Many Δ*dim-5* differentially

expressed genes associate with altered peaks of H3K27me2/3 (Figure S13), as this facultative

heterochromatin mark relocates to constitutive heterochromatic regions in a Δ*dim-5* strain (46, 47).

Unique genes are found in the ΔLGIIK9het25 and Δ*dim-5* datasets, with only 25 overlapping

differentially regulated genes. Notably, ΔLGIIK9het25 causes less drastic expression changes

relative to the Δ*dim-5* polyA mRNA-seq (Figure 6A-B). Further, there are no differentially

expressed genes in the ΔLGIIK9het25 strain wtih putative functions in chromatin or transcriptional

regulation (Supplementary File S3), although numerous hypothetical genes without defined

functions are present, preventing a complete assessment of possible genes involved in regulating

transcription. Interestingly, gene expression changes in the ΔLGIIK9het25 strain occur on each

chromosome (Figures 6B, S14), with many genes near H3K9me3-marked heterochromatic

regions. Specifically, the gene NCU08696, directly adjacent to LGIIK9het25, is downregulated in

the Δ*dim-5* strain but upregulated in ΔLGIIK9het25 (Figure 6B, zoomed region), the latter possibly

due to its proximity to the strong P$_{trpC}$ promoter. While longer chromosomes would be expected to

have more differentially expressed genes, the short ~4.3 Mb LG VII has more genes with changed

expression (n = 24 differentially expressed genes [DEGs]) than either the ~10 Mb LG I (n = 20

DEGs) or the 6.0 Mb LG IV (n =11 DEGs) (Figures 6B and S14, Supplementary File S2). We

conclude that a constitutive heterochromatic region controls gene expression across the fungal

genome.

To assess if altered 3D chromosome folding could explain the transcriptional changes in a

ΔLGIIK9het25 strain, we colored bins harboring differentially expressed genes on models of

NKL2 LGs. Qualitatively, the positions of the bins containing the differentially expressed genes

are modified on the 3D chromosome structures in the ΔLGIIK9het25 strain relative to the WT

strain (Figures 6C-D, S15, S16). Specifically, on LG II, the bins with differentially expressed genes

appear more "enclosed" within the WT structure, while these same bins are repositioned to another

"face" of the NKL2 LG II structure (Figure 6C-D). Similar qualitative bin movements altering

gene positions are readily observed on other LGs (Figure S16).

Finally, we examined the positions of ΔLGIIK9het25 differentially expressed genes relative to

the 3D folding predicted for the WT and NKL2 genomes. Our 3D genome models maintain the

biologically relevant centromere cluster independent of the bundling of telomeres (Figure S17).

The WT genome model forms an elongated structure with the LGIIK9het25 region close to the

structure bottom (Figure 7A), consistent with the "stretched" conformation of individual

chromosomes (Figures 5, S15-S16) and a previous report (48), despite different programs

predicting each 3D structure. The NKL2 genome deleted of LGIIK9het25 is wider and less

elongated (Figure 7B), similar to the individual NKL2 chromosomes. Intriguingly, the 20 kb bins

containing the ΔLGIIK9het25 differentially expressed genes are qualitatively oriented towards the

WT genome periphery, while the same bins are more centrally localized in the NKL2 genome,

consistent with the increased gene expression that occurs with the LGIIK9het25 deletion. We

conclude that the loss of a single heterochromatic region alters the 3D conformation of the genome

and disrupts normal genome function in the Neurospora nucleus.

Discussion

Here, we examined if facultative and constitutive heterochromatic regions organize the fungal

genome by performing Hi-C on strains harboring silent region deletions. Our results suggest that

gene-rich facultative heterochromatic loci are plastic in nature, as a facultative heterochromatin

deletion simply fuses nearby H3K27me2/3-enriched regions within the linear chromosome, which

minimally impacts facultative heterochromatin clustering. Specifically, the subtelomere bundling

at the terminal ~150 kb of each LG is maintained in the facultative heterochromatin deletion strain.

Subtelomere clustering may be compromised by larger deletions or only deletions targeting the

terminal ~150 kb of a chromosome. Consistent with these hypotheses, genome-wide loss of

H3K27me2/3 in a Neurospora Δ*set-7* (Δ*kmt-6*) strain causes subtelomere cluster dissociation from

the nuclear periphery (36), suggesting at least some subtelomeric H3K27me2/3 is needed for

nuclear membrane association. While larger or telomere-proximal deletions of facultative

heterochromatin could assess subtelomere clustering requirements, Neurospora may endure

alterations to gene-rich genomic loci, assuming no essential genes are deleted, given the absence

of wide-ranging topological or physiological consequences. Since most genes enriched with

H3K27me2/3 are novel genes with a hypothetical function (32), Neurospora may tolerate

subtelomeric structural variants if facultative heterochromatin is deposited. In contrast, some

Fusarium species require H3K27me2/3 for normal growth (49), meaning facultative

heterochromatic structural variants may be selected against in Fusarium.

We also assessed the role of a constitutive heterochromatic region for organizing the

Neurospora genome. The most recent Neurospora reference genome (nc14) (23) has ~200

interspersed H3K9me3-enriched regions distributed across its seven chromosomes, and apart from

the regional centromeres and telomeres of each LG required for chromosome function, the

remaining interspersed heterochromatic regions have often been dismissed as "junk DNA".

However, when we delete the largest, non-centromeric heterochromatic region on LG II, genome

organization and function are impacted. Specifically, the remaining heterochromatic regions have

altered clustering when LGIIK9het25 is deleted, as increases in intra- and inter-chromosomal

heterochromatic region contacts are observed, including between interspersed heterochromatic

regions and centromeres. These data suggest the Neurospora heterochromatin bundle is dynamic,

with no "defined" interactions mediated by specific DNA binding motifs in each heterochromatic

region. Thus, heterochromatic region aggregation mediated by Liquid Liquid Phase Separation or

a chromatin binding protein, such as Heterochromatin Protein-1 (HP1) (50–52) may allow shifting

contacts between constitutive heterochromatic regions. Since HP1 loss does not radically disrupt

the Neurospora heterochromatin bundle (24), perhaps a combination of redundant factors drive

silent chromatin bundling in fungi. Similarly, HP1-independent pericentromere clustering is

observed in Drosophila (53). Future experiments may elucidate additional mechanisms necessary

for heterochromatin clustering in eukaryotes.

The LGIIK9het25 deletion also showed changes in the local chromatin structure and the

folding of the 3D genome. In a manner synonymous to the deletion of the H3K27me2/3 region,

the loss of LGIIK9het25 caused the flanking chromatin to simply "fuse", as shown by interaction

gains between neighboring euchromatin bins. However, removing the intervening permanently

silent region causes the predicted TADs within this newly fused chromatin to change – forming a

larger TAD-like structure comprised of unique chromatin loops, suggesting the loss of

LGIIK9het25 alters the regional chromatin looping. Since interspersed heterochromatic regions in

Neurospora and other filamentous fungi act as "anchors" at the bases of TAD-like structures (9,

23–25, 35), in a mechanism tantamount to CTCF forming metazoan loops (10), removing a

heterochromatic "boundary element" would allow distant euchromatic regions to interact within a

single TAD-like structure. Complicating matters is how fungal TAD-like structures have extensive

inter-TAD interactions (23–25), meaning that the non-uniform TAD-like structures in Neurospora

may minimize the functional consequences of altering loop structures. Further, it would be

expected that TAD-like structures would differentially form on fungal chromosomes with different

patterns of interspersed heterochromatic regions. While fungi and metazoans employ different

chromatin folding mechanisms, it is possible that a deletion of a metazoan constitutive

heterochromatic region would also disrupt TADs by changing the highly isolated TAD-internal

interactions, in a similar manner to our observations in fungi. Here, loss of a H3K9me3-enriched

region from a metazoan genome could alter TAD compartmentalization or misregulate gene

expression if several smaller TADs are merged into a larger TAD. Unfortunately, the consequences

to metazoan genome function would be more dire, given how TAD boundary dysfunction

compromises proper gene expression (6–8). In fact, our chromosome and genome structure

predictions suggest local TAD changes could manifest into a completely unique folding pattern for

the overall genome. Here, the ΔLGIIK9het25 deletion caused each chromosome, and the entire

genome, to form a conformation qualitatively distinct from a wild type genome structure. While

these models are merely predictions and should be interpreted lightly, it is impactful that the

repositioning of a single locus has the potential to alter the 3D folding of the whole genome. Thus,

despite the species-specific differences in TADs or loops, it is hypothetically possible that syntenic

changes of linear chromosomal DNA could alter a species' genome organization and function.

Importantly, we show a novel function for a constitutive heterochromatic region, as the

ΔLGIIK9het25 strain is important for downregulating gene expression in a WT straiin, consistent

with the repressive nature of H3K9me3. The gene expression changes we observe when a single

constitutive heterochromatic region is deleted are not at the same extent as if all H3K9me3 is

removed in a Δ*dim-5* strain, with numerous genes both strongly up- and down-regulated (36),

suggesting the LGIIK9het25 deletion may minorly affect genome function. Overall, this single

constitutive heterochromatic region may subtly repress gene expression in the Neurospora

genome, which was hypothesized when strong gene – constitutive heterochromatin contacts were

observed in a high-resolution Neurospora Hi-C dataset (23). However, it is difficult to imagine that

the single LGIIK9het25 region deletion directly causes these gene expression changes genome

wide. More likely, the ΔLGIIK9het25 allele alters the positioning of the heterochromatin bundle –

and the nearby euchromatin – at the nuclear periphery, which repositions some gene promoters for

increased transcription machinery accessibility. This hypothesis is supported by modeling the gene

expression changes on the ΔLGIIK9het25 chromosomes and across the genome and reasonably

explains gene activation in the NKL2 strain. Of course, altered promoters can also drive gene

expression changes, as observed by the insertion of the strong P_{trpC} promoter on LG II caused a

local increase in Hi-C contacts and the production of mRNA from gene NCU08696 proximal to

the LGIIK9het25 region. In fact, NCU08696, and others close to constitutive heterochromatic

regions, are downregulated in a Δ*dim-5* strain devoid of H3K9me3. Further, the proper expression

of the Neurospora *met-8* gene requires the immediately adjacent interspersed constitutive

heterochromatic region (54). Together, these data support a hypothetical role for constitutive

heterochromatin in regulating gene expression. In fact, the local deposition of H3K9me3 at a gene

promoter can repress gene expression in metazoans, as the control of mouse embryonic stem cell

identity requires the co-deposition of H3K9me3 and H3K36me3 for transcriptional repression

(55). Thus, in addition to the canonical repression of transposable elements and repetitive

sequences in eukaryotic genomes (56–59), H3K9me3 directly regulates gene transcription. We

propose heterochromatin-mediated gene regulation also occurs through 3D organization, where

genes interact with H3K9me3-enriched regions. Specifically, genes would associate with

constitutive heterochromatic regions to localize to the more repressive periphery for reducing

transcription, and should these genes move to the more active nuclear center, transcription would

increase. Currently, it is unknown how the constitutive heterochromatin – gene interaction occurs,

but aggregation of HP1 bound to H3K9me3 at promoters with the constitutive heterochromatin

bundle is an attractive model. Fungi may be more apt to employ this transcriptional regulation

mechanism, as opposed to their metazoan counterparts, given how fungal TAD-like structures have

more promiscuous contacts and how the Rabl chromosome conformation has extensive inter-

chromosomal contacts (9, 23–25, 35). Recently, the histone methyltransferase SETDB1, which

catalyzes H3K9me3 across metazoan genomes, was shown to maintain gene expression in addition

to driving chromatin compartmentalization (60), suggesting that constitutive heterochromatic

regions have additional, underappreciated roles in both genome organization and function across

diverse species. Further research in multiple species should help elucidate novel functional roles

for this "junk DNA" that is pervasive yet evolutionarily retained in genomes from fungi to

metazoans.

Materials and Methods

Strains and growth conditions

   Neurospora crassa strains WT N150 (derived from the strain 74-OR23-IVA [Fungal Genetics Stock Center #2489]), N2930 (*mat A his-3; Δmus52::bar*[+]), N3944 (mat x; *Δdim-5::bar*[+]), and N4933 (*mat a*; Δ47.4 kb::*hph*[+]) were gifts from Eric U. Selker (University of Oregon). All strains were grown with 1x Vogels minimal medium + 1.5% sucrose and supplements at 32°C (61). All strains are available upon request.

   The ΔLGIIK9het25 (NKL2) strain replacing 110,609 basepairs (bp) of AT-rich DNA from H3K9me3-enriched region #25 on LG II (between bp 3,602,033 and 3,712,857 of Neurospora reference genome version 14 [nc14]) (23) with 1,401 bp of the P_{trpC}::*hph* hygromycin resistance cassette (deleting a total of 109,208 bp of DNA from the Neurospora genome) was created by split marker gene replacement. Briefly, the ~1,000 bp upstream of LGIIK9het25 was PCR amplified with oligonucleotides ("oligos") oKL14 and oKL15, while the ~1,000 bp downstream of LGIIK9het25 was amplified with oligos oKL18 and oKL19, by Phusion DNA polymerase (cat# F530-L; ThermoFisher Scientific) using the manufacturer's conditions; oligos oKL15 and oKL18 had ten 5' complementary nucleotides specific to the P_{trpC} promoter and *hph* gene 3' end, respectively. Oligo sequences are provided in Table S1. DNA fragments were gel purified using the GenCatch Gel Extraction Kit (cat# 2260250; Epoch Life Sciences), and split marker fragments fusing the LGIIK9het25 upstream to the P_{trpC} promoter ("UP"; oKL14 and 2955) or fusing the *hph* gene to the LGIIK9het25 downstream ("DOWN"; oligos 2954 and oKL19) were PCR amplified with the upstream and downstream fragments, and an *hph* gene containing plasmid (p3xFLAG::hph::LoxP) (62) using LA Taq (cat# RR002M; Takara), and gel purified. The UP and

DOWN split marker fragments were transformed into N2930 by electroporation (BioRad

MicroPulser Electroporator, cat# 1652100EDU). Hygromycin resistant colonies were selected on

1x Vogels+FGS media with 100 µg/mL hygromycin (cat# ant-hg-5; Invivogen). Genomic DNA

was isolated from resultant colonies, using standard protocols (63), and proper hygromycin

cassette integration at LGIIK9het25 borders was performed by PCR using oKL49 and 2955 for

the left border, 2954 and oKL50 for the right border, and oKL49 and oKL50 for the entire

integrated $P_{trpC}::hph$ gene. Positive integrant strain tKL1A-2 was crossed to N150 (xKL9AR) and

genomic DNA was isolated from hygromycin resistant progeny; similar PCR reactions were

performed to confirm the presence of the ΔLGIIK9het25 allele in individual cross progeny. PCR

with oligos oKL55 and oKL56 determined Δ*mus-52::bar* allele absence; strain mating type was

established by crosses with strains of known mating types. The final homokaryotic strain NKL2

from cross progeny xKL9AR-17 contains only the ΔLGIIK9het25 deletion (NKL2 genotype: *mat*

A; ΔLGIIK9het25::hph).

Hi-C library construction and bioinformatic analyses

In situ Hi-C was performed as previously described (23, 24). Hi-C libraries were Illumina

sequenced at the Genomics and Cell Characterization Core Facility (GC3F; University of Oregon),

either on a Hi-Seq 2500 as PE100 reads or a NovaSeq 6000 as PE59 reads. For comparison to

mutant Hi-C datasets, WT *Dpn*II and *Mse*I *in situ* Hi-C datasets (NCBI Gene Expresssion Omnibus

(GEO) accession number GSE173593) (23) were generated by extracting the number of total reads

from WT datasets using the sed command to provide approximately equal numbers of valid reads

as mutant datasets. Reads were mapped with bowtie2 (64) to the *Neurospora crassa* version 14

(nc14) reference genome (23), the output sam files were used to build the initial Hi-C contact

matrix at a high [1 kb] resolution with hicExplorer software package (65), and the output contact

matrix was used to analyze replicate dataset similarity (hicCorrelate), generate lower resolution

contact matrices (hicMergeMatrixBins), compare WT and mutant datasets (hicCompareMatrices),

KR correct matrices (hicCorrectMatrix), assess TADs (hicFindTADs [using --

correctForMultipleTesting false discovery rate, --thresholdComparisons 0.3, and --delta 0.15

values to make TAD bed files] and hicPlotTADs), and generate images (hicPlotMatrix). Individual

chromosome 3D structure prediction was performed with 3DMax (https://github.com/BDM-

Lab/3DMax)(45) while entire Neurospora 3D genome structure prediction was performed by

LorDG (66), both using custom scripts that are available on the Oluwadare lab GitHub website

(https://github.com/OluwadareLab/Fungi_3DGenome), tailored to fungal chromosome biology.

Input contact matrices for structure prediction were made by merging the individual *Dpn*II and

*Mse*I matrices (hicSumMatrices) and converting the .h5 output format to NxN text files with a

custom script (23) (https://github.com/Klocko-Lab). Output PDB files were displayed on

ChimeraX (67), individual bins with differentially expressed genes or chromosomal features were

manually colored, and high-resolution images were used for figure construction. PDB files are

available on Zenodo (10.5281/zenodo.11245552).

Chromatin Immunoprecipitation-sequencing (ChIP-seq) library construction and bioinformatic

analyses

ChIP-seq was performed as previously described (38), using anti-H3K9me3 (Active Motif;

cat# 39161, lot# 30220003) and anti-H3K27me2/3 (Active Motif; cat# 39535, lot# 23120014)

antibodies. All ChIP-seq libraries were Illumina sequenced at the GC3F on a NovaSeq 6000 as

SR118 reads. Fastq dataset files were mapped to nc14 with bowtie2 (64), converted to bam files,

sorted, and indexed with samtools (68), made into bigwig or bedgraph files with Deeptools (69),

and displayed on Integrative Genomics Viewer (IGV) (70), with exported png images from IGV

used for figure creation. Previously published NCBI GEO or the NCBI Sequence Read Archive

(SRA) datasets for WT H3K9me3 (GEO accession numbers GSE68897 and GSE98911), WT

H3K27me2/3 (GEO accession numbers GSE68897 and GSE100770), and Δ*dim-5* H3K27me2/3

(GEO accession number GSE68897 and SRA accession number SRP058573)(31, 46, 47, 71), were

mapped to the nc14 genome (23) and used for bigwig/bedgraph file creation for image generation.

Total RNA isolation, polyadenylated RNA-seq library construction, and bioinformatic analyses

Total RNA was isolated from overnight liquid cultures of Neurospora strains grown in 1x

Vogels, 1.5% sucrose plus supplements (32, 36). Cultures were harvested by vacuum filtration,

placed in 2.0mL screw-top tubes, and 300µL of acid washed glass beads suspended in dH_2O (150-

212µm, Sigma Aldrich cat# G1145-10G), 350µL Acid Phenol:chloroform (5:1; ThermoFisher cat#

AM9720), and 350µL NETS (300mM NaCl, 1mM EDTA, 10mM Tris pH 7.5, 0.2% SDS) were

added. Samples were vortexed for 10 minutes at maximum speed and centrifuged at 4°C for 5

minutes at 13k rpm. The aqueous layer was transferred to two tubes (~250µL each), 650µL of cold

100% EtOH was added, and RNA was precipitated on ice for 10 minutes and pelleted by

microcentrifugation at 4°C for 10 minutes at 13k rpm. The pellet was washed with cold 70% EtOH

(made with DECP-treated dH_2O), the supernatant was removed, and pellets were allowed to air

dry for 15 minutes and resuspended in 50µL DEPC-treated dH_2O. Insoluble material was removed

by centrifugation, and the initial RNA concentration was measured by Nanodrop. Total RNA

(10µg) was resuspended in 50µL of DEPC-treated dH_2O. Total RNA cleanup was performed by

the Zymo RNA clean and concentrator kit (Zymo Research cat# R1013), per the manufacturer's

protocol; the 15µL elution was divided into 4µL for Qubit concentration (RNA HS assay kit;

ThermoFisher Scientific, cat# Q32852) and TapeStation (Agilent) quality control analysis, while

the remaining 11µL was stored at -80oC. Only total RNA samples with high ribosomal RNA to

messenger RNA (mRNA) ratios were used for library construction. Polyadenylated mRNA (polyA

mRNA) was selected with the NEBNext Poly(A) mRNA Magnetic Isolation Module (New

England Biolabs [NEB], cat# E7490L) and polyA mRNA sequencing libraries were generated with

the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (NEB, cat# E7760L and

E7765L) per the manufacturer's protocols; quadruplicate polyA mRNA-seq replicate libraries

were constructed for each strain. All polyA mRNA-seq libraries were Illumina sequenced at the

GC3F on a NovaSeq 6000 as SR118 reads. Raw fastq files were mapped to the nc14 genome using

HiSat2 (72), and gene count files were generated with htseq-count (73). Differential Expression

analysis between strains was performed with DESeq2 (74) in R/Rstudio (75, 76), selecting

significantly changed genes as those with expression $\log_2 > 3.0$ or $\log_2 < -3.0$ and an adjusted p-

value < 0.001; bed files for display on IGV were generated from the DESeq2 output gene lists in

Microsoft Excel, and volcano plots of $\log_2$ fold change vs. adjusted p-value were generated in R/R

studio.

Data Availability

547 All genomic datasets have been deposited to the NCBI GEO under the SuperSeries accession

548 number GSE269235. Individual datasets have the subseries numbers GSE269228 (ChIP-seq),

549 GSE269229 (Hi-C), and GSE269230 (polyA mRNA-seq).

Acknowledgements

The authors thank Sara Rodriguez, Yulia Shtanko, and Ashley Scadden for Hi-C library

generation help; Sara Hanson (Colorado College) for TapeStation use and RNA-seq protocol

advice; the University of Oregon GC3F for Illumina library sequencing service, and Klocko lab

members and UCCS Department of Chemistry & Biochemistry colleagues for helpful comments

and discussions. Funding was provided by a UCCS Undergraduate Research Academy award to

A.T.R., start-up funds from the UCCS College of Engineering and Applied Science to O.O., a

Maximizing Investigators Research Award grant (1R35GM150402-01) to O.O., start-up funds

from the UCCS College of Letters, Arts, and Sciences to A.D.K., and a NIH Academic Research

Enhancement Award grant (1R15GM140396-01) to A.D.K.

Declaration of Interest Statement

563   The authors report there are no competing interests to declare.

References

1. Cremer T, Cremer C. 2001. Chromosome territories, nuclear architecture and gene regulation in
mammalian cells. Nat Rev Genet 2:292–301.

2. Misteli T. 2007. Beyond the Sequence: Cellular Organization of Genome Function. Cell
128:787–800.

3. Dekker J. 2008. Gene Regulation in the Third Dimension. Science 319:1793–1794.

4. Sexton T, Cavalli G. 2015. The Role of Chromosome Domains in Shaping the Functional
Genome. Cell 160:1049–1059.

5. Bonev B, Cavalli G. 2016. Organization and function of the 3D genome. Nat Rev Genet 17:661–
678.

6. Flavahan WA, Drier Y, Liau BB, Gillespie SM, Venteicher AS, Stemmer-Rachamimov AO,
Suvà ML, Bernstein BE. 2016. Insulator dysfunction and oncogene activation in IDH mutant
gliomas. Nature 529:110–114.

578 7. Dixon JR, Xu J, Dileep V, Zhan Y, Song F, Le VT, Yardımcı GG, Chakraborty A, Bann DV,
579 Wang Y, Clark R, Zhang L, Yang H, Liu T, Iyyanki S, An L, Pool C, Sasaki T, Rivera-Mulia JC,
580 Ozadam H, Lajoie BR, Kaul R, Buckley M, Lee K, Diegel M, Pezic D, Ernst C, Hadjur S, Odom
581 DT, Stamatoyannopoulos JA, Broach JR, Hardison RC, Ay F, Noble WS, Dekker J, Gilbert DM,
582 Yue F. 2018. Integrative detection and analysis of structural variation in cancer genomes. Nat
583 Genet 50:1388–1398.

584 8. Northcott PA, Lee C, Zichner T, Stütz AM, Erkek S, Kawauchi D, Shih DJH, Hovestadt V,
585 Zapatka M, Sturm D, Jones DTW, Kool M, Remke M, Cavalli FMG, Zuyderduyn S, Bader GD,
586 VandenBerg S, Esparza LA, Ryzhova M, Wang W, Wittmann A, Stark S, Sieber L, Seker-Cin H,
587 Linke L, Kratochwil F, Jäger N, Buchhalter I, Imbusch CD, Zipprich G, Raeder B, Schmidt S,
588 Diessl N, Wolf S, Wiemann S, Brors B, Lawerenz C, Eils J, Warnatz H-J, Risch T, Yaspo M-L,
589 Weber UD, Bartholomae CC, Kalle C von, Turányi E, Hauser P, Sanden E, Darabi A, Siesjö P,
590 Sterba J, Zitterbart K, Sumerauer D, Sluis P van, Versteeg R, Volckmann R, Koster J, Schuhmann
591 MU, Ebinger M, Grimes HL, Robinson GW, Gajjar A, Mynarek M, Hoff K von, Rutkowski S,
592 Pietsch T, Scheurlen W, Felsberg J, Reifenberger G, Kulozik AE, Deimling A von, Witt O, Eils
593 R, Gilbertson RJ, Korshunov A, Taylor MD, Lichter P, Korbel JO, Wechsler-Reya RJ, Pfister SM.
594 2014. Enhancer hijacking activates GFI1 family oncogenes in medulloblastoma. Nature 511:428–
595 34.

596 9. Torres DE, Reckard AT, Klocko AD, Seidl MF. 2023. Nuclear genome organization in fungi:
597 From gene folding to Rabl chromosomes. FEMS Microbiol Rev
598 https://doi.org/10.1093/femsre/fuad021.

599 10. Rao SSP, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL,
600 Machol I, Omer AD, Lander ES, Aiden EL. 2014. A 3D Map of the Human Genome at Kilobase
601 Resolution Reveals Principles of Chromatin Looping. Cell 159:1665–1680.

602 11. Lieberman-Aiden E, Berkum NL van, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I,
603 Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke
604 A, Stamatoyannopoulos J, Mirny LA, Lander ES, Dekker J. 2009. Comprehensive Mapping of
605 Long-Range Interactions Reveals Folding Principles of the Human Genome. Science 326:289–
606 293.

607 12. Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. 1997. Crystal structure of the
608 nucleosome core particle at 2.8 Å resolution. Nature 389:251–260.

609 13. Luger K, Richmond TJ. 1998. DNA binding within the nucleosome core. Curr Opin Struc Biol
610 8:33–40.

611 14. Rao SSP, Huang S-C, Hilaire BGS, Engreitz JM, Perez EM, Kieffer-Kwon K-R, Sanborn AL,
612 Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J, Turner D, Ye Z, Omer
613 AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander ES, Aiden EL. 2017. Cohesin Loss
614 Eliminates All Loop Domains. Cell 171:305-320.e24.

15. Mizuguchi T, Fudenberg G, Mehta S, Belton J-M, Taneja N, Folco HD, FitzGerald P, Dekker
J, Mirny L, Barrowman J, Grewal SIS. 2014. Cohesin-dependent globules and heterochromatin
shape 3D genome architecture in *S. pombe*. Nature 516:432–435.

16. Campos EI, Reinberg D. 2009. Histones: Annotating Chromatin. Annu Rev Genet 43:559–
599.

17. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B. 2012. Topological
Domains in Mammalian Genomes Identified by Analysis of Chromatin Interactions. Nature
485:376–380.

18. Kind J, Pagie L, de Vries SS, Nahidiazar L, Dey SS, Bienko M, Zhan Y, Lajoie B, de Graaf
CA, Amendola M, Fudenberg G, Imakaev M, Mirny LA, Jalink K, Dekker J, van Oudenaarden A,
van Steensel B. 2015. Genome-wide Maps of Nuclear Lamina Interactions in Single Human Cells.
Cell 163:134–147.

19. Falk M, Feodorova Y, Naumova N, Imakaev M, Lajoie BR, Leonhardt H, Joffe B, Dekker J,
Fudenberg G, Solovei I, Mirny LA. 2019. Heterochromatin drives compartmentalization of
inverted and conventional nuclei. Nature 570:395–399.

20. Schoenfelder S, Sexton T, Chakalova L, Cope NF, Horton A, Andrews S, Kurukuti S, Mitchell
JA, Umlauf D, Dimitrova DS, Eskiw CH, Luo Y, Wei C-L, Ruan Y, Bieker JJ, Fraser P. 2010.
Preferential associations between co-regulated genes reveal a transcriptional interactome in
erythroid cells. Nat Genet 42:53–61.

21. Hoencamp C, Dudchenko O, Elbatsh AMO, Brahmachari S, Raaijmakers JA, Schaik T van,
Cacciatore ÁS, Contessoto VG, Heesbeen RGHP van, Broek B van den, Mhaskar AN, Teunissen
H, Hilaire BGS, Weisz D, Omer AD, Pham M, Colaric Z, Yang Z, Rao SSP, Mitra N, Lui C, Yao
W, Khan R, Moroz LL, Kohn A, Leger JSt, Mena A, Holcroft K, Gambetta MC, Lim F, Farley E,
Stein N, Haddad A, Chauss D, Mutlu AS, Wang MC, Young ND, Hildebrandt E, Cheng HH,
Knight CJ, Burnham TLU, Hovel KA, Beel AJ, Mattei P-J, Kornberg RD, Warren WC, Cary G,
Gómez-Skarmeta JL, Hinman V, Lindblad-Toh K, Palma FD, Maeshima K, Multani AS, Pathak
S, Nel-Themaat L, Behringer RR, Kaur P, Medema RH, Steensel B van, Wit E de, Onuchic JN,
Pierro MD, Aiden EL, Rowland BD. 2021. 3D genomics across the tree of life reveals condensin
II as a determinant of architecture type. Science 372:984–989.

22. Imakaev M, Fudenberg G, McCord RP, Naumova N, Goloborodko A, Lajoie BR, Dekker J,
Mirny LA. 2012. Iterative correction of Hi-C data reveals hallmarks of chromosome organization.
Nat Methods 9:999–1003.

23. Rodriguez S, Ward A, Reckard AT, Shtanko Y, Hull-Crew C, Klocko AD. 2022. The genome
organization of *Neurospora crassa* at high resolution uncovers principles of fungal chromosome
topology. G3 Genes Genomes Genetics https://doi.org/10.1093/g3journal/jkac053.

24. Galazka JM, Klocko AD, Uesaka M, Honda S, Selker EU, Freitag M. 2016. Neurospora
chromosomes are organized by blocks of importin alpha-dependent heterochromatin that are
largely independent of H3K9me3. Genome Res 26:1069–1080.

25. Torres DE, Kramer HM, Tracanna V, Fiorin GL, Cook DE, Seidl MF, Thomma BPHJ. 2023.
Implications of the three-dimensional chromatin organization for genome evolution in a fungal
plant pathogen. Nat Commun 15:1701.

26. Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, FitzHugh W, Ma L-J,
Smirnov S, Purcell S, Rehman B, Elkins T, Engels R, Wang S, Nielsen CB, Butler J, Endrizzi M,
Qui D, Ianakiev P, Bell-Pedersen D, Nelson MA, Werner-Washburne M, Selitrennikoff CP,
Kinsey JA, Braun EL, Zelter A, Schulte U, Kothe GO, Jedd G, Mewes W, Staben C, Marcotte E,
Greenberg D, Roy A, Foley K, Naylor J, Stange-Thomann N, Barrett R, Gnerre S, Kamal M,
Kamvysselis M, Mauceli E, Bielke C, Rudd S, Frishman D, Krystofova S, Rasmussen C,
Metzenberg RL, Perkins DD, Kroken S, Cogoni C, Macino G, Catcheside D, Li W, Pratt RJ,
Osmani SA, DeSouza CPC, Glass L, Orbach MJ, Berglund JA, Voelker R, Yarden O, Plamann M,
Seiler S, Dunlap J, Radford A, Aramayo R, Natvig DO, Alex LA, Mannhaupt G, Ebbole DJ,
Freitag M, Paulsen I, Sachs MS, Lander ES, Nusbaum C, Birren B. 2003. The genome sequence
of the filamentous fungus *Neurospora crassa*. Nature 422:859–868.

27. Lewis ZA, Honda S, Khlafallah TK, Jeffress JK, Freitag M, Mohn F, Schübeler D, Selker EU.
2009. Relics of repeat-induced point mutation direct heterochromatin formation in *Neurospora*
crassa. Genome Res 19:427–437.

28. Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger MR, Altemose
N, Uralsky L, Gershman A, Aganezov S, Hoyt SJ, Diekhans M, Logsdon GA, Alonge M,
Antonarakis SE, Borchers M, Bouffard GG, Brooks SY, Caldas GV, Chen N-C, Cheng H, Chin
C-S, Chow W, Lima LG de, Dishuck PC, Durbin R, Dvorkina T, Fiddes IT, Formenti G, Fulton
RS, Fungtammasan A, Garrison E, Grady PGS, Graves-Lindsay TA, Hall IM, Hansen NF, Hartley
GA, Haukness M, Howe K, Hunkapiller MW, Jain C, Jain M, Jarvis ED, Kerpedjiev P, Kirsche
M, Kolmogorov M, Korlach J, Kremitzki M, Li H, Maduro VV, Marschall T, McCartney AM,
McDaniel J, Miller DE, Mullikin JC, Myers EW, Olson ND, Paten B, Peluso P, Pevzner PA,
Porubsky D, Potapova T, Rogaev EI, Rosenfeld JA, Salzberg SL, Schneider VA, Sedlazeck FJ,
Shafin K, Shew CJ, Shumate A, Sims Y, Smit AFA, Soto DC, Sović I, Storer JM, Streets A,
Sullivan BA, Thibaud-Nissen F, Torrance J, Wagner J, Walenz BP, Wenger A, Wood JMD, Xiao
C, Yan SM, Young AC, Zarate S, Surti U, McCoy RC, Dennis MY, Alexandrov IA, Gerton JL,
O'Neill RJ, Timp W, Zook JM, Schatz MC, Eichler EE, Miga KH, Phillippy AM. 2022. The
complete sequence of a human genome. Science 376:44–53.

29. Tamaru H, Zhang X, McMillen D, Singh PB, Nakayama J, Grewal SI, Allis CD, Cheng X,
Selker EU. 2003. Trimethylated lysine 9 of histone H3 is a mark for DNA methylation in
Neurospora crassa. Nat Genet 34:75–79.

30. Lewis ZA, Adhvaryu KK, Honda S, Shiver AL, Knip M, Sack R, Selker EU. 2010. DNA
Methylation and Normal Chromosome Behavior in Neurospora Depend on Five Components of a
Histone Methyltransferase Complex, DCDC. Plos Genet 6:e1001196.

31. Klocko AD, Uesaka M, Ormsby T, Rountree MR, Wiles ET, Adhvaryu KK, Honda S, Selker
EU. 2019. Nucleosome Positioning by an Evolutionarily Conserved Chromatin Remodeler
Prevents Aberrant DNA Methylation in Neurospora. Genetics 211:563–578.

32. Jamieson K, Rountree MR, Lewis ZA, Stajich JE, Selker EU. 2013. Regional control of histone
H3 lysine 27 methylation in Neurospora. Proc National Acad Sci 110:6027–6032.

33. Jamieson K, McNaught KJ, Ormsby T, Leggett NA, Honda S, Selker EU. 2018. Telomere
repeats induce domains of H3K27 methylation in Neurospora. Elife 7:e31216.

34. Wiles ET, Selker EU. 2017. H3K27 methylation: a promiscuous repressive chromatin mark.
Curr Opin Genet Dev 43:31–37.

35. Winter DJ, Ganley ARD, Young CA, Liachko I, Schardl CL, Dupont P-Y, Berry D, Ram A,
Scott B, Cox MP. 2018. Repeat elements organise 3D genome structure and mediate transcription
in the filamentous fungus *Epichloë festucae*. Plos Genet 14:e1007467.

36. Klocko AD, Ormsby T, Galazka JM, Leggett NA, Uesaka M, Honda S, Freitag M, Selker EU.
2016. Normal chromosome conformation depends on subtelomeric facultative heterochromatin in
Neurospora crassa. Proc National Acad Sci 113:15048–15053.

37. Knight PA, Ruiz D. 2013. A fast algorithm for matrix balancing. Ima J Numer Anal 33:1029–
1047.

38. Scadden AW, Graybill AS, Hull-Crew C, Lundberg TJ, Lande NM, Klocko AD. 2023. Histone
deacetylation and cytosine methylation compartmentalize heterochromatic regions in the genome
organization of *Neurospora crassa*. Proc Natl Acad Sci 120:e2311249120.

39. Ninomiya Y, Suzuki K, Ishii C, Inoue H. 2004. Highly efficient gene replacements in
Neurospora strains deficient for nonhomologous end-joining. P Natl Acad Sci Usa 101:12248–
12253.

40. Selker E. 1990. Premeiotic Instability Of Repeated Sequences In *Neurospora crassa*. Annu
Rev Genet 24:579–613.

41. Gladyshev E, Kleckner N. 2014. Direct recognition of homology between double helices of
DNA in *Neurospora crassa*. Nat Commun 5:3509–3509.

42. Bonhoure N, Bounova G, Bernasconi D, Praz V, Lammers F, Canella D, Willis IM, Herr W,
Hernandez N, Delorenzi M, Consortium TC, Hernandez N, Delorenzi M, Deplancke B, Desvergne
B, Guex N, Herr W, Naef F, Rougemont J, Schibler U, Andersin T, Cousin P, Gilardi F, Gos P,
Lammers F, Raghav S, Villeneuve D, Fabbretti R, Vlegel V, Xenarios I, Migliavacca E, Praz V,
David F, Jarosz Y, Kuznetsov D, Liechti R, Martin O, Delafontaine J, Cajan J, Gustafson K, Krier
I, Leleu M, Molina N, Naldi A, Rib L, Symul L, Bounova G. 2014. Quantifying ChIP-seq data: a
spiking method providing an internal reference for sample-to-sample normalization. Genome Res
24:1157–68.

43. Orlando DA, Chen MW, Brown VE, Solanki S, Choi YJ, Olson ER, Fritz CC, Bradner JE,
Guenther MG. 2014. Quantitative ChIP-Seq Normalization Reveals Global Modulation of the
Epigenome. Cell Reports 9:1163–1170.

44. Zufferey M, Tavernari D, Oricchio E, Ciriello G. 2018. Comparison of computational methods
for the identification of topologically associating domains. Genome Biol 19:217.

45. Oluwadare O, Zhang Y, Cheng J. 2018. A maximum likelihood algorithm for reconstructing
3D structures of human chromosomes from chromosomal contact data. BMC Genom 19:161.

46. Jamieson K, Wiles ET, McNaught KJ, Sidoli S, Leggett N, Shao Y, Garcia BA, Selker EU.
2016. Loss of HP1 causes depletion of H3K27me3 from facultative heterochromatin and gain of
H3K27me2 at constitutive heterochromatin. Genome Res 26:97–107.

47. Basenko EY, Sasaki T, Ji L, Prybol CJ, Burckhardt RM, Schmitz RJ, Lewis ZA. 2015.
Genome-wide redistribution of H3K27me3 is linked to genotoxic stress and defective growth. Proc
National Acad Sci 112:E6339–E6348.

48. Poinsignon T, Gallopin M, Grognet P, Malagnac F, Lelandais G, Poulain P. 2023. 3D models
of fungal chromosomes to enhance visual integration of omics data. NAR Genom Bioinform
5:lqad104.

49. Connolly LR, Smith KM, Freitag M. 2013. The Fusarium graminearum Histone H3 K27
Methyltransferase KMT6 Regulates Development and Expression of Secondary Metabolite Gene
Clusters. Plos Genet 9:e1003916.

50. Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA,
Redding S, Narlikar GJ. 2017. Liquid droplet formation by HP1α suggests a role for phase
separation in heterochromatin. Nature 547:236–240.

51. Freitag M, Hickey PC, Khlafallah TK, Read ND, Selker EU. 2004. HP1 Is Essential for DNA
Methylation in Neurospora. Mol Cell 13:427–434.

52. Zenk F, Zhan Y, Kos P, Löser E, Atinbayeva N, Schächtle M, Tiana G, Giorgetti L, Iovino N.
2021. HP1 drives *de novo* 3D genome reorganization in early Drosophila embryos. Nature
593:289–293.

53. Stutzman AV, Hill CA, Armstrong RL, Gohil R, Duronio RJ, Dowen JM, McKay DJ. 2024.
Heterochromatic 3D genome organization is directed by HP1a- and H3K9-dependent and
independent mechanisms. Mol Cell https://doi.org/10.1016/j.molcel.2024.05.002.

54. Yang S, Li W, Qi S, Gai K, Chen Y, Suo J, Cao Y, He Y, Wang Y, He Q. 2014. The highly
expressed methionine synthase gene of *Neurospora crassa* is positively regulated by its proximal
heterochromatic region. Nucleic Acids Res 42:6183–95.

55. Barral A, Pozo G, Ducrot L, Papadopoulos GL, Sauzet S, Oldfield AJ, Cavalli G, Déjardin J.
2022. SETDB1/NSD-dependent H3K9me3/H3K36me3 dual heterochromatin maintains gene
expression profiles by bookmarking poised enhancers. Mol Cell 82:816-832.e12.

56. Padeken J, Zeller P, Towbin B, Katic I, Kalck V, Methot SP, Gasser SM. 2019. Synergistic
lethality between BRCA1 and H3K9me2 loss reflects satellite derepression. Genes Dev 33:436–
451.

57. Padeken J, Methot SP, Gasser SM. 2022. Establishment of H3K9-methylated heterochromatin
and its functions in tissue differentiation and maintenance. Nat Rev Mol Cell Biol 23:623–640.

58. Herberg S, Simeone A, Oikawa M, Jullien J, Bradshaw CR, Teperek M, Gurdon J, Miyamoto
K. 2015. Histone H3 lysine 9 trimethylation is required for suppressing the expression of an
embryonically activated retrotransposon in *Xenopus laevis*. Sci Rep 5:14236.

59. Zhao S, Lu J, Pan B, Fan H, Byrum SD, Xu C, Kim A, Guo Y, Kanchi KL, Gong W, Sun T,
Storey AJ, Burkholder NT, Mackintosh SG, Kuhlers PC, Edmondson RD, Strahl BD, Diao Y,
Tackett AJ, Raab JR, Cai L, Song J, Wang GG. 2023. TNRC18 engages H3K9me3 to mediate
silencing of endogenous retrotransposons. Nature 623:633–642.

60. Keenan CR, Coughlan HD, Iannarella N, Fierro AT del, Keniry A, Johanson TM, Chan WF,
Garnham AL, Whitehead LW, Blewitt ME, Smyth GK, Allan RS. 2024. Suv39h-catalyzed
H3K9me3 is critical for euchromatic genome organization and the maintenance of gene
transcription. Genome Res 34:556–571.

61. Davis RH. 2000. Neurospora: contributions of a model organism. Oxford University Press.

62. Honda S, Selker EU. 2009. Tools for Fungal Proteomics: Multifunctional Neurospora Vectors
for Gene Replacement, Protein Expression and Protein Purification. Genetics 182:11–23.

63. Miao VPW, Freitag M, Selker EU. 2000. Short TpA-rich segments of the ζ-η region induce
DNA methylation in *Neurospora crassa*. J Mol Biol 300:249–273.

64. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods
9:357–359.

65. Ramírez F, Bhardwaj V, Arrigoni L, Lam KC, Grüning BA, Villaveces J, Habermann B,
Akhtar A, Manke T. 2018. High-resolution TADs reveal DNA sequences underlying genome
organization in flies. Nat Commun 9:189.

66. Trieu T, Cheng J. 2017. 3D genome structure modeling by Lorentzian objective function.
Nucleic Acids Res 45:1049–1058.

67. Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE.
2021. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein
Sci 30:70–82.

68. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R, Subgroup 1000 Genome Project Data Processing. 2009. The Sequence Alignment/Map format
and SAMtools. Bioinformatics 25:2078–2079.

69. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F,
Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res 44:W160-5.

70. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.
2011. Integrative genomics viewer. Nat Biotechnol 29:24–6.

71. Klocko AD, Summers CA, Glover ML, Parrish R, Storck WK, McNaught KJ, Moss ND,
Gotting K, Stewart A, Morrison AM, Payne L, Hatakeyama S, Selker EU. 2020. Selection and
Characterization of Mutants Defective in DNA Methylation in *Neurospora crassa*. Genetics
216:genetics.303471.2020.

72. Zhang Y, Park C, Bennett C, Thornton M, Kim D. 2020. Rapid and accurate alignment of
nucleotide conversion sequencing reads with HISAT-3N. Genome Res 31:1290–1295.

73. Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinform Oxf Engl 31:166–9.

74. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15:550.

75. Team Rs. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.
http://www.rstudio.com/.

76. Team RC. 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Figure Legends

Figure 1. A strain harboring a 47.4 kilobase deletion of a H3K27me2/3-enriched domain has

minimal genome organization change. (A-B) Contact probability heatmaps of *Dpn*II

(euchromatin-specific) *in situ* Hi-C datasets at 20 kb resolution across Linkage Group VI (LG VI)

of the (A) wild type (WT) or (B) N4933 strains, the latter deleted of a Δ47.4kb H3K27me2/3-

enriched region ("ΔK27"). Enhanced heatmaps show the terminal 500 kb of LG VI, harboring the

deleted H3K27me2/3 region, at 5 kb resolution. Each Hi-C image displayed has the raw contact

probability heatmap shown above the diagonal and the Knight Ruiz (KR) corrected contact

probability heatmap, for reducing inherent contact probability biases (37), displayed below the

diagonal. Contact probability scale bars shown below the images. Images of wild type H3K9me3

(green) and H3K27me2/3 (purple) ChIP-seq tracks, displayed on IGV (70), above and to the right.

The purple line in (A) shows the WT location of the H3K27me2/3-enriched region that was deleted

in the ΔK27 strain, while the black arrowhead in (B) shows the deletion site in the N4933 strain.

(C) KR corrected contact probability heatmaps of the *Dpn*II *in situ* Hi-C datasets of WT and ΔK27

strains primarily showing on-diagonal contacts at 5 kb bin resolution. Wild type H3K9me3 ChIP-

seq (green), H3K27me2/3 ChIP-seq (purple), and gene (gray) tracks shown below. Contact

probability scalebar to the left. The black arrowhead shows the gain in contacts between fused

euchromatic DNA flanking the ΔK27 deletion. The gray arrowhead shows a loss of contact

probability at a putative transposable element (gene NCU09969; *sly-1*). (D-E) Contact probability

heatmaps of telomere interactions within WT and ΔK27 strains, as (D) shows intra-chromosomal

(e.g., the contacts between the left and right telomeres of LG VI), while (E) shows inter-

chromosomal (e.g., the contacts between the left telomere of LG VI and the right telomere of LG

V) telomere contacts. Telomere schematic, with green circle showing the position of the telomere

repeats, H3K27me2/3 ChIP-seq (purple), and gene (gray) tracks shown above and to the right. The

red "X" shows the H3K27me2/3 domain that is deleted, while the black arrowhead shows the

depletion of contacts at that deletion site.

Figure 2. Deletion of a ~110 kb H3K9me3-enriched domain alters constitutive

heterochromatic region clustering. (A-D) Contact probability heatmaps of (A-B) *Dpn*II

(euchromatin-specific) and (C-D) *Mse*I (heterochromatin specific) *in situ* Hi-C datasets at 20 kb

resolution across LG II, , as in Figure 1A-B, of the (A, C) wild type (WT) or (B, D) NKL2 strains,

the latter deleted of the 25th H3K9me3-enriched region from the LG II left telomere

("ΔLGIIK9het25"). The purple line on panel A shows the deleted LGIIK9het25 region. Enhanced

heatmaps at 5 kb resolution show a 500 kb region of LG II surrounding the ΔLGIIK9het25 allele,

as in Figure 1A-B. (E-F) Images showing the $\log_2$ change in Hi-C contact probability between WT

and ΔLGIIK9het25 (E) *Dpn*II (euchromatin-specific) or (F) *Mse*I (heterochromatin-specific)

datasets at 5 kb resolution across a 500 kb region of LG II surrounding the ΔLGIIK9het25 allele.

Scalebar of contact probability changes is shown below. (G) Images showing the change in inter-

chromosomal *Mse*I (heterochromatin-specific) Hi-C contact probability between the entire LG I

chromosome and the (top) LG II centromere or (bottom) LGIIK9het25 interspersed

heterochromatic region.

Figure 3. The LGIIK9het25 deletion does not compromise the deposition of H3K9me3 at

other gene-poor constitutive heterochromatic regions in the NKL2 strain. (A-B) Images of

WT (dark green) or NKL2 ΔLGIIK9het25 (light green) H3K9me3 ChIP-seq and gene (gray)

tracks, displayed on IGV, of the entire LG II chromosome or enhanced regions. H3K9me3 ChIP-

seq reads were mapped to the (A) WT nc14 reference genome or (B) the NKL2 ΔLGIIK9het25

reference genome. Numbers in the enhanced panel in Figure 3A count the heterochromatic regions

from the LG II left telomere, while the black arrowheads show the position at which the DNA

sequence has been deleted in the NKL2 ΔLGIIK9het25 strain, and the purple line shows the ~110

kb of DNA that is missing from the NKL2 genome (causing all peaks to be shifted by ~110 kb

starting at the black arrowhead). The red arrow in panel B shows the position of the inserted

P_{trpC}::*hph* gene in the NKL2 reference genome.

Figure 4. The LGIIK9het25 deletion alters the predicted TAD-like structures in the surrounding chromatin. (A-B) Contact probability heatmaps of *Dpn*II (KR corrected) or *Mse*I (raw) *in situ* Hi-C datasets of WT and ΔLGIIK9het25 strains primarily showing on-diagonal contacts at 5 kb bin resolution, mapped to either the (A) nc14 WT reference genome or the (B) NKL2 ΔLGIIK9het25 reference genome. Wild type H3K9me3 ChIP-seq (green) and gene (gray) tracks shown below. The contact probability scalebar is shown to the left of each heatmap. Black triangular lines show predicted TAD-like structures, as calculated by the program hicFindTADs, in the defined regions. In panel B, the black arrowheads indicate predicted TAD-like structures surrounding the LGIIK9het25 constitutive heterochromatic region deletion that change, while the open arrowheads show altered TAD-like structures in more distant, flanking euchromatin. The red arrow shows the position of the $P_{trpC}::hph$ gene in the NKL2 reference genome, while the purple line shows increased chromatin contacts surrounding the P_{trpC} promoter.

Figure 5. The predicted 3D folding of a single chromosome is changed with the LGIIK9het25 deletion. (A-B) Images depicting the predicted 3D modeling of LG II from (A) WT or (B) NKL2 ΔLGIIK9het25 strains at 20 kb resolution. The green highlighted region shows the 20 kb bins covering the LGIIK9het25 region in the WT structure. The enhanced image shows the folding of the chromatin at the right arm of LG II surrounding the LGIIK9het25 region in a WT strain. The arrow shows the LGIIK9het25 deletion location.

Figure 6. The LGIIK9het25 region deletion causes increased gene expression genome wide. (A) Volcano plots showing the $\log_2$ fold change from WT expression level of all Neurospora genes

verses adjusted p-values from (left) the NKL2 ΔLGIIK9het25 strain or (right) the Δ*dim-5* strain.

Points colored red are genes that have significant expression changes, with $\log_2$ fold changes > 3.0

or < -3.0 and adjusted p-values < 0.001. The number of significantly changed genes in each strain

indicated in red at the top of the plot. (B) Images of IGV tracks showing WT H3K9me3 ChIP-seq

(green), genes (gray), or the positions of differentially expressed genes with $\log_2$ fold changes >

3.0 ("up" genes) or < -3.0 ("down" genes) and adjusted p-values < 0.001. The purple "X" symbols

show the LGIIK9het25 constitutive heterochromatic region deleted in the NKL2 strain, while the

black arrows show the positions of genes NCU08696 on LG II or *dim-5* (NCU04402) on LG IV.

(C-D) Images depicting the predicted 3D modeling of LG II from (C) WT or (D) NKL2

ΔLGIIK9het25 strains at 20 kb resolution. The green highlighted region shows the 20 kb bins

encompassing the LGIIK9het25 region in the WT structure, while red highlights show the 20 kb

bins containing differentially expressed genes in the NKL2 strain, relative to a WT strain; the same

bins are marked on both structures. The arrow in panel D shows the LGIIK9het25 deletion

location.

Figure 7. The predicted 3D structure of the entire genome, as well as the positions of

differentially expressed genes, are altered when the LGIIK9het25 region is deleted. (A-B)

Images depicting the predicted 3D modeling of the entire genome of (A) WT or (B) NKL2

ΔLGIIK9het25 strains. The model on the left shows the side view of the genome structure, which

is rotated up 90° to obtain the right model, which shows the bottom of the structure from the

centromere cluster. The green highlighted region shows the 20 kb bins covering the LGIIK9het25

region in the WT structure, while the red regions highlight the 20 kb bins containing differentially

expressed genes in the NKL2 ΔLGIIK9het25 strain.

A Linkage Group VI
WT Hi-C *Dpn*II digest ~9M valid reads, raw

H3K9me3
H3K27me2/3

WT Hi-C *Dpn*II ~9M valid reads, KR correct

raw

bins

KR

Distance (kb)

5 kb Resolution Bin Reads (log$_{10}$)

Distance (Mb)

20 kb Resolution Bin Reads (log$_{10}$)

B Linkage Group VI
ΔK27 N4933 Hi-C *Dpn*II, raw

H3K9me3
H3K27me2/3

ΔK27 N4933 Hi-C *Dpn*II KR correct

raw

bins

KR

Distance (kb)

5 kb Resolution Bin Reads (log$_{10}$)

Distance (Mb)

20 kb Resolution Bin Reads (log$_{10}$)

Figure 1

C Linkage Group VI

0 kb   100   200   300   400   500 kb

WT N150
*Dpn*II Hi-C

5 kb Resolution Bin Reads (log$_{10}$)

ΔK27 N4933
*Dpn*II Hi-C

5 kb Resolution Bin Reads (log$_{10}$)

WT H3K27me2/3

WT H3K9me3

genes

D LGVI L tel   LGVI L tel

WT H3K27me2/3
genes

LGVI R tel

WT N150   ΔK27 N4933

E LGVI L tel   LGVI L tel

LGV R tel

WT N150   ΔK27 N4933

A Linkage Group II - 20 kb Resolution
WT *Dpn*II Hi-C ~21M valid reads, raw

B Linkage Group II - 20 kb Resolution
NKL2 ΔLGIIK9het25 *Dpn*II Hi-C, raw

Figure 2

C Linkage Group II - 20 kb Resolution
WT *Mse*I Hi-C ~12M valid reads, raw

D Linkage Group II - 20 kb Resolution
NKL2 ΔLGIIK9het25 *Mse*I Hi-C, raw

E WT v NKL2 ΔLGIIK9het25 *Dpn*II Hi-C

F WT v NKL2 ΔLGIIK9het25 *Mse*I Hi-C

G WT v NKL2 ΔLGIIK9het25 *Mse*I Hi-C

Figure 3

Figure 4

A

RNA-seq comparison WT vs. NKL2

n = 84

-log₁₀(adjusted p-value)

log₂ Fold Change of all Neurospora genes

log₂ Fold Change > 3.0 or < -3.0; adjusted p-value < 0.001

RNA-seq comparison WT vs. Δdim-5

n = 175

log₂ Fold Change of all Neurospora genes

log₂ Fold Change > 3.0 or < -3.0; adjusted p-value < 0.001

B

3.5 Mb   4.1 Mb

WT H3K9me3   NCU08696   1000   0

NKL2 - 4x up

genes

4x up

Δdim-5   4x down

Linkage Group II

0 Mb   4 Mb

WT H3K9me3   1000   0

NKL2 - 4x up

genes

Δdim-5   4x up   4x down

Linkage Group IV

0 Mb   6 Mb

WT H3K9me3   1000   0

NKL2 - 4x up

genes

Δdim-5   4x up   4x down

dim-5 (NCU04402)

C

WT LG II

D

NKL2 LG II

20 kb bins with Differentially Expressed Genes

Figure 6

A

B

Figure 7

WT genome

NKL2 genome

90°

90°

20 kb bins with Differentially Expressed Genes