1

Title: Understanding organism-habitat relationships and critically evaluating reference areas is

key to marine protected area assessment

Authors: Conner Jainese[1], Peter M. Carlson[1], Katelin Seeto[1], Lyndsey McNeill[1], Kelly Sivertson[1],

Jennifer E. Caselle[1]

Affiliation: Marine Science Institute, University of California Santa Barbara[1]

Corresponding Author:

Conner Jainese, connerjainese@ucsb.edu

University of California, Santa Barbara

Marine Science Institute, Bld. 520

Santa Barbara, California

Abstract

      Marine Protected Areas (MPAs) have been implemented globally as a conservation tool to improve the health and function of marine ecosystems. Research has focused on assessing MPA effectiveness, however certain habitats and communities are often avoided because they are difficult or expensive to monitor. Mesophotic (30-100m) rocky reef fish communities are a valuable commercial and recreational resource that is highly targeted but often overlooked in monitoring due to depth restricted sampling. We used two MPAs in California's statewide protection network, along with parried reference sites, to test how protection status along with environmental conditions influenced the abundance and biomass of three highly targeted species with varying life histories and habitat preferences. Depth and habitat were strong predictors for all groups; ocean whitefish (*Caulolatilus princeps*), California sheephead (*Semicossyphus pulcher*), and targeted rockfish (*Sebastes spp*). The pattern of these effects however, differed between the species groups and the influence of protection was mixed. This work highlights how species with high habitat affinities benefit differently from protection, as a function of depth and habitat representation within the MPA/reference pair. To accurately evaluate MPAs, and the network as a whole, researchers must recognize organism-habitat relationships and incorporate them when assessing conservation efforts.

Introduction

Marine Protected Areas (MPAs) have been implemented globally as a means of protecting marine ecosystems from anthropogenic pressures (Gleason et al. 2013; Sullivan-Stack et al. 2022; Maestro et al. 2019). The primary goal of most MPAs is to conserve and maintain marine biodiversity, with secondary benefits, such as increased catch rates outside of protection zones and economic stability for local communities, also predicted to occur (Murray et al. 1999; Sala and Giakoumi 2018). There is strong evidence that an MPA network, opposed to a single MPA, is more effective at broadly protecting a diverse suite of species across various life stages and habitats (Gaines et al. 2010; Grorud-Colvert et al. 2014). Well-designed MPA networks are able to incorporate nursery areas, account for ontogenetic habitat shifts, recognize different species-habitat associations, and protect vulnerable spawning grounds (Grüss et al. 2014; Olson et al. 2019). Monitoring all of these habitats is difficult and it is rare that all habitats within a single MPA or across an MPA network are effectively monitored. Monitoring priority is often given to those habitats more easily or cheaply accessed (shallow water reefs, grass beds, shorelines, etc.), while more challenging habitats are often data-poor (Field et al. 2007; Noble-James et al. 2023).

Mesophotic (30-100m) reefs represent one habitat that has been historically difficult to monitor and a growing body of research acknowledges our relatively incomplete understanding of these communities globally (Kahng et al. 2017; James et al. 2017; Cerrano et al. 2019). Mesophotic reefs support a diverse suite of commercially valuable species and it is estimated that a significant portion of commercial groundfish harvest falls within this depth zone (Miller et al. 2017; Jacquemont et al. 2024). As a result, these fishes are left under-surveyed and relatively data poor, while simultaneously being

highly targeted. Along the west coast of California many of the mesophotic fisheries are

actively managed, with enforcement agencies frequently revising depth restrictions and

species-specific take limits. Active management requires monitoring data from these

depths and a combination of fisheries-dependent and independent data has been shown to

provide a robust measure of stock size (Dennis et al. 2015; Howard et al. 2023). Baited

remote underwater video (BRUV) systems have opened new opportunities for better

understanding mesophotic reefs in temperate and tropical ecosystems by providing a

fisheries-independent, non-destructive sampling technique that can operate at depths

beyond recreational or even most scientific SCUBA diving limits (Brown et al. 2022;

Andradi-Brown et al. 2016; Brokovich et al. 2008).

Mesophotic reefs represent an ecologically important habitat that contains distinct

invertebrate and fish communities (Williams et al. 2019; Honeyman et al. 2023; Bell et al.

2024). Below 30m depth, solar irradiance begins to attenuate, limiting photosynthesis and

algal growth that characterizes shallower reefs. At these depths, structure forming sessile

invertebrate, many themselves of conservation concern, such as gorgonians, anemones,

calcareous algae, and bryozoans dominate the benthos (Ponti et al. 2018). In the Northern

hemisphere, the mesophotic temperate reef fish community is dominated by Rockfishes

(genus *Sebastes*), a diverse assemblage of ecologically and economically important fishes

that includes >100 recognized species (Love et al. 2002; Love and Yoklavich 2006; Love

2011). The mesophotic depth zone offers a relatively stable set of environmental conditions

that can function as nursery habitat and a refuge space for species that were historically

found along wider depth gradients (Baillon et al. 2012; Kahng et al. 2017; Giraldo-Ospina

et al. 2020). These habitats also provide an important ontogenetic bridge between shallow

nursery habitats and deeper adult habitats (Love 2011; Swadling et al. 2022). Finally, these

deep habitats might be critical refuges from the effects of climate change, such as increases

in sea water temperature (MacDonald et al. 2016; Pereira et al. 2018). Together these

factors make mesophotic reefs important contributors to marine biodiversity and a

necessary component of an effective MPA network.

One of the expected benefits of MPAs is the buildup of targeted species within its

boundaries and consequently, an MPA's performance is often measured by its ability to

enhance the biomass and abundance of fisheries targets (Caselle et al. 2015; Lenihan et al.

2022). Recreational and commercial fisheries often have a number of targeted species that

utilize a wide breadth of habitats across the seascape (Region 2003; Scholz et al. 2006).

Effective MPA networks should protect habitats ranging from sand, sand-rock ecotone,

rocky reefs, and pelagic zones, while incorporating these habitats across an extensive depth

range (Young and Carr 2015). Despite the fact that MPAs are an ecosystem conservation

tool, it is unlikely any MPA can achieve a 'one-size fits all' design, instead, any single

MPA will likely be more effective at protecting some species compared to others. Factors

such as species movement, the availability of preferred habitat, oceanographic conditions,

predator and prey abundances, and recruitment dynamics will all influence an MPA's

ability to enhance particular targeted species (Moffitt et al. 2011).

With the purpose of better understanding the effects of MPAs on mesophotic rocky

reef fish communities, we implemented the use of BRUV camera systems at two

geographically distinct MPAs (and nearby reference areas) within California's Northern

Channel Islands (Figure 1): Carrington Point State Marine Reserve (SMR) at Santa Rosa

Island and Anacapa Island SMR/State Marine Conservation Area (SMCA). Straddled

across a biogeographic transition zone, where cool southern-flowing waters of the

California Current meet warmer northern-flowing waters from the Southern California

Counter Current (Horn and Allen 1978), these two MPAs are characteristically distinct,

occupying opposite ends of a strong environmental gradient that exists across the Northern

Channel Islands. The primary variation along this gradient is in sea surface temperature

(SST), accompanied by other related factors including productivity, frequency of

disturbance/wave exposure, algal species persistence, and level of human interaction

(Harms and Winant 1998; Hamilton et al. 2010; Caselle et al. 2015).

We utilized stereo BRUV surveys because they are well suited for capturing fish

community data across this under-sampled depth range and are useful for a variety of habitat

types (Cappo et al. 2004; Heyns-Veale et al. 2016; Whitmarsh et al. 2017). Additional

advantages of assessing fish communities using BRUVs include its non-destructive nature as a

sampling technique (essential for MPA monitoring), the ability to collect highly accurate fish

measurements for biomass calculations, the creation of a video record that can be re-analyzed to

extract novel information or correct for inter-observer biases, the ability to observe species

targeted by fishing and non-targeted species, and the opportunity to collect visual data on

species habitat associations (Cappo et al. 2006; Langlois et al. 2010; Bennett et al. 2016).

BRUV systems have become a widely implemented technique in many parts of the world as a

non-destructive, fishery independent data collection tool for monitoring marine protected areas

(Rees et al. 2004; Langlois et al. 2006; Kelaher et al. 2014) but are still in the early stages of use

in California.

Using BRUV survey data from Carrington Pt. SMR and Anacapa SMCA/SMR

and nearby reference areas we measured the importance of habitat type, depth, and

protection status on the abundance and biomass of targeted fish species. These two focal

MPAs are nested into larger networks; first, a network within the Channel Islands (Figure

1B), which itself is nested into a large California statewide network of protected areas.

Previous studies have shown individual variation in MPA performance across the network

for shallow reefs and kelp forests (Caselle et al. 2015; Ziegler et al. 2023). Here we use

two MPAs to demonstrate the potential for individual MPA variation in performance in

presumably more stable, deeper water habitats.

Methods

Sampling Design

Our BRUV design was modified from existing designs used extensively to monitor a

variety of marine habitats in Australia (Willis and Babcock 2000; Goetze et al. 2021; Harvey et

al. 2021). We deployed BRUVs using a stratified random approach, using geological habitat

maps and bathymetry to target rocky reef habitat within our sampling depths (30-100m).

Because the Carrington Pt. SMR has very little habitat >50m deep, survey depths at this site

were generally limited to this maximum depth, while surveys at Anacapa extended to 100m

depth. A map showing each MPA/reference zone polygon can be found in Figure S1.

Sampling effort included both MPA and reference sites (where fishing is allowed)

on any single day to control for environmental variability between sampling days.

Sampling took place from August to October for four consecutive years (2019-2022). In

2020, we also sampled in June at Anacapa Island as a response to the shutdown of

commercial passenger fishing vessels (CPFV) at that location due to COVID-19. We

tested for seasonal or COVID related affects and finding none, this data was included in

all analyses unless otherwise noted. A map of all sampling locations, by year is available

in Figure S2. A more detailed description of BRUV sampling methods and tools can be

found in supplemental materials Methods S1.

Video Processing

We analyzed video files using observation logging and 3D measurement software

SeaGIS EventMeasure (www.seagis.com.au). We identified all observed fishes to the

lowest taxonomic group possible. In order to quantify the relative abundance of fishes we

recorded MaxN, the maximum number of individuals of a species present in a single video

frame (Willis and Babcock 2000) using the first 30 minutes of each BRUV survey (Harasti

et al. 2015), for every species observed. Without the ability to discern individuals or

distinguish multiple sightings of the same fish, MaxN is a conservative estimate of relative

abundance that has become the standard metric for BRUV surveys (Langlois et al. 2020).

We also measured the total length for every fish observed in the MaxN video

frame. In some instances, we were not able to accurately measure all individuals in a

particular MaxN frame due to the camera angle or obstructions blocking one of the

cameras. In this case, we measured as many individuals as possible from the MaxN frame.

We estimated the biomass of individual fishes using an allometric length-weight

conversion: $W = aTL^b$, where parameters a and b are species specific constants, TL is total

length in cm, and W is weight in grams. We obtained length-weight fitting parameters

from the literature and FishBase (Froese and Pauly 2000). Because not all fish in the MaxN

frame can be measured (some individuals are not within view of both cameras) we used the

mean of all individual weights in the MaxN frame, multiplied by the MaxN for each

individual species to estimate biomass. We classified species as targeted by fishing or non-

targeted by fishing using information from the CA Department of Fish and Wildlife as well

as local knowledge of the authors.

In order to understand the influence of habitat on MPA effectiveness, and because

we cannot control the precise view of the forward facing BRUV cameras, we classified

habitat based on substrate characteristics within the field of view of the cameras. The four

habitat classifications were "Hard", "Mixed-Hard", "Mixed-Soft", and "Soft" (Categories

defined in Table S1).

To standardize the quality of videos used in analysis, we quantified the "usability"

of each video. All videos were assigned a "usability" score that considered the amount of

visible substrate and video length (Table S2). Any videos where the BRUV unit did not

remain upright for the entire 30 min (i.e., scores of 3 or 4) were removed from analyses.

Focal fish species

In addition to groupings of all targeted species and all non-targeted species, we also

analyzed patterns of biomass and MaxN for specific focal species. Ocean whitefish, the most

abundant species observed in our study, are a medium size (39-56 cm TL at sexual maturity)

tilefish (*Malacanthidae*) common in Southern California and Baja, Mexico. Typically residing at

the periphery of rocky reefs, ocean whitefish primarily feed on small invertebrates often in

nearby sandy flats (Love 2011). This species is a staple for the recreational fishery in the heart of

its geographic range and is likely to benefit from fishing protection.

California sheephead, which appeared on nearly half of our surveys, are the largest and

only targeted wrasse (*Labridae*) species that inhabits the study area. California sheephead are not

only highly targeted by recreational anglers, but also by a smaller live-fish commercial fishery

(Love 2011). As a protogynous hermaphrodite (female at birth with the potential to transition to

male later in life), fishing pressure has a more complex effect on the sex/size structure and the

reproductive potential of a local population (Hamilton et al. 2007).

With 21 unique species of rockfish observed in this study, we more closely

examined a sub-group of rockfish that we considered "targeted" and are more likely to

respond to protection from fishing. This grouping excludes small-bodied rockfish that are

infrequently caught and almost never retained by anglers such as halfbanded (*S.*

semicinctus) and calico rockfish (*S. dallii*). Three targeted species: vermilion (*S. miniatus*),

copper (*S. caurinus*), and blue rockfish (*S. mystinus*) made up over 70% of the targeted

rockfish observed on our surveys. Vermilion and copper rockfish are long-lived (50-60

years) high value species that have a significant recreational and commercial fishery,

increasing their potential to benefit from protection. Blue rockfish are a schooling, largely

planktivorous species, that is of moderate value to recreational and commercial fishing

(Love 2011).

Abundance and Biomass Modeling

Generalized linear models (GLM) were used to understand how differences in habitat

type, depth, and MPA status influenced fish abundance and biomass. To account for species-

specific habitat preference and other potential traits (e.g. movement patterns) we ran models on

targeted rockfish, ocean whitefish, and California sheephead separately. We also ran separate

models for each island because the depth ranges differed (Carrington Pt. SMR ranged from 30m

- 60m depth; Anacapa SMCA/SMR ranged from 30m – 90m depth). Predictor variables were

MPA status, depth, and habitat type, as well as the interaction between MPA status and habitat

type to test if the effect of MPA status is similar across different habitat types.

All statistical models and model diagnostics were run using R version 4.3.0. Data from

each species and metric (biomass and MaxN) combination was zero inflated, and the fitdistrplus

package and descdist() function were used to determine the best fit distribution. Each data

combination was fit to a negative binominal distribution and the models were run using the

glm.nb function in the lme4 package. Randomized quantile residuals were calculated and tested

for overdispersion using the DHARMa package and all dispersions statistics were <1. The

'visreg' package was used to visualize how each model parameter influenced a particular metric

and represents each model estimate when all other predictors are held constant. We standardized

the visualizations by setting MPA status to 'MPA', habitat type to 'Hard', and depth as the

median depth from a particular island. Each regression plot shows a model prediction, 95%

confidence interval, and the partial residuals of the model. We calculated the precent deviance

explained for each model to understand how well our predictors explained the variation in the

data.

Population Size Structure

To understand how protection from fishing alters the demographics of targeted fish

populations, we plotted the size density histograms of the three most common targeted rockfishes

(copper, vermillion, and blue rockfish), ocean whitefish, and California sheephead.

<u>Results</u>

We completed 280 and 341 BRUV surveys at Carrington Pt. and Anacapa Island

respectively for a total of 621 surveys. Of those 621 surveys, 493 were classified as 'usable'

(79% usable) and were included in analyses (Table S3) In general, two conditions resulted in

unusable video; these were very strong currents and/or highly rugose habitat causing the BRUV

system to not land upright, or tip over at some point during the video survey.

Habitat Description

We found differences in amount and type of rock at the two islands. When rock was

present at Anacapa, the habitat was characterized by a patchy reef system with most video

classified as 'mixed soft' or 'mixed hard' (Fig. 2, Table S4). In addition, the Anacapa MPA

contained more 'soft' habitat classifications than the other island/MPA groupings. Carrington

Pt. was characterized by more consistent hard bottom, with most videos classified as 'hard' or

'mixed hard' (Figure 2, Table S4).

Similar to the video observations, benthic habitat maps showed a disparity between

the amount of rocky substrate present in the Anacapa MPA and reference area, with 3-5

times more eligible grid cells in the reference area. Although the reference area was twice

as large as the MPA, there still appears to be proportionally more abundant hard bottom

habitat in the reference area. This is particularly true for the deepest zone (70-100m),

which had 5 times as many eligible grid cells (MPA = 33 vs Reference = 167, Table S5) in

the reference area. Carrington Pt. had a relatively similar number of eligible grid cells

within the 30m - 50m depth bin, with the reference area only being slightly larger (Table

S5). Overall, the benthic habitat maps used to identify eligible grid cells generally agreed

with the observed habitat data from BRUV surveys, with Carrington Pt containing more

consistent hard bottom and Anacapa containing more sand-rock ecotone habitat.

Species Observed

We observed a total of 65 fish species on the BRUV surveys across four years and

both locations. Total summed MaxN and frequency of occurrence for each species are

shown in Table S6. A single species, ocean whitefish, accounted for 32% of the summed

MaxN values. These fish had the highest abundance across our study and were present on

84% of all surveys. Two species, copper rockfish and California sheephead had both high

MaxN values and high frequency of occurrence at both islands. Copper rockfish occurred

on 48% of surveys while California sheephead occurred on 49% of surveys (Table S6).

Four schooling species: jack mackerel (*Trachurus symmetricus*), blacksmith (*Chromis*

punctipinnis), halfbanded rockfish, and blue rockfish were abundant when observed but

occurred less frequently (4-27% of all surveys).

The BRUVs were able to observe species rarely seen in other surveys in California

(e.g., SCUBA, ROV, submersible). These included species of concern including giant sea

bass (*Stereolepis gigas*) (50 individual observations), bocaccio (*Sebastes paucispinis*) (79

individuals) and bat rays (*Myliobatis californica*) (11 individuals) as well as various

species of the more cryptic flatfishes (including California halibut (*Paralichthys*

californicus) and Pacific sanddabs (*Citharichthys sordidus*)). Although many species were

documented by only a single individual, this type of presence-only data can be useful for

documenting climate-induced range shifts, invasions, and habitat associations.

Abundance and Biomass

Targeted Rockfish -

At both islands, targeted rockfishes showed similar patterns with habitat type and

depth being significant predictors of both biomass and abundance. Generally, both

biomass and abundance increased as the availability of hard substrate increased, with

'hard' vs. 'mixed soft' and 'soft' being highly significant (all eight comparisons across

island, metric, hard vs. mixed hard, and hard vs. soft had a p-value <0.001; Figure 3 &

Table S7). At Anacapa, depth was a significant predictor for targeted rockfish abundance

and biomass, with both metrics increasing with depth (MaxN p-value = <0.001; biomass

p-value = <0.001; Figure 3 & Table S7). The patterns at Carrington Pt. were less

consistent, with depth only being a significant predictor of biomass (p-value = 0.02,

Figure 3 & Table S7), most likely related to the limited depth range at this location

compared to Anacapa. Interestingly, the interaction between soft habitat and MPA status

was almost significant for targeted rockfish biomass at Anacapa. Here, soft habitat was the

only habitat type predicted to have greater targeted rockfish biomass inside the MPA

compared to the reference area (p-value = 0.06, estimate = -1.43; Table S7 & Figure S3A).

This interaction term indicates that the model estimate for soft bottom habitats in the MPA

was greater compared to reference areas, while in other habitats (e.g., 'Hard', 'Mixed

Hard', and 'Mixed-soft') the reference area estimate was larger. The only significant MPA

effect, beyond the influence of habitat and depth, was for targeted rockfish biomass inside

the Carrington Pt. MPA (p-value = 0.03; Figure 3 & Table S7).

Ocean Whitefish -

Depth was a strong predictor of ocean whitefish biomass and abundance with higher

abundance and biomass found in shallower waters (p-value = < 0.001 for all comparisons of

MaxN and biomass at both islands; Figure 4 & Table S8). The only significant habitat predictor

was for soft habitats at Carrington Pt., with both abundance and biomass significantly lower on

sandy, soft bottom habitats compared to hard substrate (p-value = 0.05; Figure 4 & Table

S8). We did not find a significant interaction term between MPA status and habitat type for

either ocean whitefish metric (Figure S4 & Table S8). Anacapa showed a significant MPA effect,

with the abundance and biomass of ocean whitefish greater inside the MPA (MaxN p-value =

0.02; biomass p-value = 0.01; Figure 4 & Table S8).

California Sheephead -

Similar to ocean whitefish, depth was found to be a strong predictor of California

sheephead biomass and abundance, with higher levels found in the shallower waters (Carrington

Pt. and Anacapa MaxN p-value <0.001; Carrington Pt. biomass p-value = <0.01; Anacapa

biomass p-value = <0.001; Figure 5 & Table S9). California sheephead were significantly less

abundant on two habitat types at Anacapa, 'mixed soft' and 'soft', compared to 'hard' substrate

('mixed soft' p-value = 0.03; 'soft' p-value = <0.01; Figure 5 & Table S9), and had significantly

lower biomass on 'soft' vs 'hard' habitat at Anacapa ('soft' p-value = 0.05; Figure 5 & Table

S9). At Carrington Pt., habitat alone was not found to be a significant predictor of sheephead

abundance or biomass. We did find a significant interaction term for sheephead biomass and

abundance between 'soft' habitat and MPA status at Carrington Pt., however these results had

very few observations of sheephead on soft habitat across protection zones, limiting inference

from this result. At Anacapa, we found a significant negative MPA effect for the abundance and

biomass of sheephead, with greater values in the reference area (MaxN p-value = 0.01, estimate

= 1.01; biomass p-value = 0.03, estimate = 1.09; Figure 5 & Table S9).

Deviance Explained

Across all species and both metrics, the model framework performed slightly better at

Anacapa compared to Carrington Pt., with the deviance explained ranging from 0.46 – 0.22 at

Anacapa and 0.22 – 0.17 at Carrington Pt. (Tables S7 – S9). This is likely related to the larger

available depth range at Anacapa and that depth was a significant predictor of abundance and

biomass for all three species in this study. The highest deviance explained was for California

sheephead at Anacapa (DE = 0.46 for MaxN; DE = 0.45 for Biomass; Table S9) and is most

likely related to the smaller range in MaxN and biomass values associated with the California

sheephead data compared to the larger range associated with targeted rockfish and ocean

whitefish.

Focal Species Size Structure

We plotted the density of individual fish total lengths in MPAs and reference areas

for the five most common targeted species in our study; copper rockfish (n = 506),

vermilion rockfish (n = 459), blue rockfish (n = 1760), ocean whitefish (n = 5767), and

California sheephead (n = 603) (Figure 6). Copper rockfish were generally larger inside the

MPAs at both Anacapa and Carrington Pt., with the MPA containing more individuals

>400mm TL (Figure 6A). Vermilion rockfish had a bimodal size structure with multiple

size classes present at Carrington Pt. and Anacapa (Figure 6B). Inside the Anacapa MPA,

the population size structure is dominated by smaller sized individuals (<20cm), with

proportionally fewer adult fishes. While the population size structure patterns for

vermillion rockfish at Carrington Pt. appear to be more related to protection from fishing,

with proportionally larger individuals inside the MPA. The smaller-bodied schooling blue

rockfish had similar population size structure across protection zones at Carrington Pt. and

Anacapa (Figure 6C). Ocean whitefish and California Sheephead showed similar patterns

at Anacapa and Carrington, with a greater proportion of larger individuals inside the MPA

at each location (Figure 6D & E).

Discussion

In this study we show that environmental drivers, such as depth and habitat availability,

need to be considered when assessing the ability of MPAs to enhance the biomass and

abundance of targeted species. MPA effectiveness, in both tropical and temperate marine

ecosystems, has been linked to benthic features such as the amount of consolidated hard bottom,

reef complexity, and habitat continuity (Dames et al. 2020). Individual species have been shown

to respond differently to such features, hence species-habitat associations must be considered

during MPA implementation, when defining MPA goals, and when assessing performance.

Studies have shown that a lack of preferred habitat inside the protection zone can limit the ability

of MPAs to enhance species and functional diversity (Navarro-Martínez et al. 2023). MPAs are

designed as ecosystem management tools, with broad goals of protecting biodiversity, yet

individual species can have widely varying habitat requirements. Many MPA design processes,

especially for MPA networks, try to encompass habitat replication and representation (Roberts et

al. 2003; Halpern et al. 2010; Saarman et al. 2013), but in practice, it is impossible to design

optimally for every species. Thus, accounting for habitat in post implementation monitoring and

choice of reference areas, can help to set realistic expectations for MPA performance.

The Carrington Pt. MPA and adjacent reference area had similar amounts of continuous

hard substrate, which indicated that the greater biomass of targeted rockfishes in the MPA was

likely related to protection more than habitat. Rockfish (family *Sebastes*) are generally

characterized as a demersal, rock associated and heavily fished family, and it appears that the

Carrington Pt. MPA is well designed for their protection. Inside the Anacapa MPA, there was a

notable lack of continuous hard substrate, particularly at depth, that appears to limit the

performance of the MPA at that location for this rock-associated fish group and makes

comparisons with the paired reference area challenging. However, the Anacapa MPA is likely

more effective at enhancing the ocean whitefish population, a more mobile species that is

associated with the sand-rock ecotone (Bellquist et al. 2008). Habitat maps and *in situ*

observations from Anacapa MPA and reference area showed that these locations were dominated

by a mix of soft, sandy habitat and isolated patch reefs. Ocean whitefish abundance and biomass

was also strongly associated with depth, but unlike targeted rockfish, ocean whitefish are more

abundant in shallower areas, with the majority of observations occurring shallower than 60m.

The examples above, comparing very different fish species groups, demonstrate how

scientific guidelines and a network approach to MPA design can help meet ecosystem goals and

protect populations of different species across a variety of habitats. The creation of networks,

with numerous reserves spread across various habitats, has become the standard by which to

structure marine no-take zones, both at the local and global scale. The California Marine Life

Protection Act (MLPA) established the network of MPAs throughout California with the goals of

protecting ecosystem biodiversity and protecting and rebuilding populations of key species. This

process relied on MPA design guidelines from a scientific advisory committee and chief among

those guidelines was 'habitat representation' throughout the network (Saarman et al. 2013). At

the network level, this was achieved by including a variety of major habitat types such as; rocky

intertidal zones, estuaries, and shallow and deep subtidal habitat. Here we show that a more

nuanced treatment of habitat within an MPA can provide critical information on individual MPA

performance for key species (Young and Carr 2015).

By incorporating habitat availability in and out of the MPAs and studying deeper zones

than previously explored, we found several results that differed from prior studies and/or our

expectations. Contrary to our predictions and prior studies of shallower reefs in the area (Caselle

et al. 2015; Honeyman et al. 2023), California sheephead abundance and biomass at these deeper

depths was significantly greater in the Anacapa reference area which is open to fishing. Active

tracking data has shown that California sheephead use a variety of habitats, but tend to favor

rocky-reefs, spending 50% - 70% of their time on hard, consolidated substrate (Topping et al.

2005). The deep-water habitat within the Anacapa MPA is lacking this hard substrate, which is

more abundant in the shallow kelp forests of the Anacapa MPA. This illustrates the importance

of surveying the full range of depths for species of interest when evaluating MPA performance

or advising fisheries management. A more comprehensive approach to MPA design and

management can be achieved by filling gaps in our knowledge of how MPAs may differentially

enhance fish communities at all depths and habitat types.

To measure the ecological effects of a marine protected areas, especially without data for

that site prior to MPA implementation, researchers often use a nearby fished area (reference site)

as a control by which hypotheses about the cessation of fishing can be tested. Selection of these

'control' sites must weigh numerous factors such as; habitat type, depth range, species

composition, and seasonal oceanographic characteristics, when deciding on location and scale.

Often the easiest way to broadly control for various environmental factors is by placing the

reference area in close proximity to the MPA. However, unlike strictly controlled lab

experiments, MPAs are expected to affect control areas by design. Adjacent placement of

reference areas, while often meeting goals of controlling for environmental conditions and

perhaps even habitat, creates a scientific design challenge, with the likelihood that reference sites

will be affected by the presence of a nearby MPA through the redistribution of fishing effort, and

potential "spillover" of fish coming from the MPA (Caselle et al. 2015; Di Lorenzo et al. 2016).

Anacapa Island presents an interesting case study in reference site selection. Previous

studies of very deep water habitats using Remotely Operated Vehicles excluded the Anacapa

MPA from analyses, citing a lack of appropriate reference area (Karpov et al. 2012). This

exclusion comes despite containing one of the oldest MPAs in the Northern Channel Islands

network, with the Anacapa eastern SMR established in 1978. Here we chose the reference

location as immediately offshore of, and deeper than, long-standing reference sites used for

shallower rocky reef and kelp forest studies (Caselle et al. 2015; White et al. 2021). While shown

to be appropriate for shallow rock reef, we found that in deeper waters, the reference area

contained more deep rock than in the MPA, potentially limiting our ability to detect an MPA

effect on rock-associated species (e.g., targeted rockfish, California sheephead). However, both

reference area and MPA contained similar amounts of deep sand-rock ecotone habitat favored by

ocean whitefish, the most numerous species of fish in our study. Hence tradeoffs exist where

reference site selection must incorporate species-habitat associations, assess MPA goals, work to

inform management decisions, and are beholden to the realities of field studies.

A consistent MPA effect was observed in the size structure of three targeted rockfish

species, ocean whitefish and California sheephead, with the greater proportion of larger

individuals inside the protection zone. This is not surprising as one of the earliest effects of a

cessation of fishing is to allow larger individuals to grow and persist (Mumby et al. 2021;

Bejarano et al. 2019; Taylor and McIlwain 2010). The notable exception was for vermillion

rockfish inside the Anacapa MPA, which contained fewer large individuals than the reference

area. Interestingly, the size structure for vermilion rockfish at both study sites reveal distinct

cohorts at two size classes. This likely indicates periodic strong recruitment/survivorship for this

species and provides an opportunity to track these cohorts over time. Repeated annual sampling

such as done here allows the tracking of these types of population demographics through time to

better understand year to year variability in recruitment dynamics for valuable fishery species.

The ability to collect size structure data, with tools such as stereo BRUVs, can provide important

additional insight into the effects of protection on these populations and is particularly important

for the management of long-lived species like rockfishes.

The ability of MPAs to conserve and rebuild fish communities has been shown to

improve with the protection of habitat across a wide range of depths (Curley et al. 2002; Goetze

et al. 2021). Many species of fish utilize different depth zones and habitats across life stages,

often with juvenile fish recruiting to relatively protected shallow nursery habitats and moving to

deeper offshore habitats as they mature (Gibson et al. 2002; Aburto-Oropeza et al. 2009; Li et al.

2022). This ontogenetic movement is common with a number of the rockfish species observed in

this study (Love et al. 1991) and should be considered in the context of MPA performance. The

availability of suitable habitat along the ontogenetic depth profile likely influences the

distribution of rockfish at varying life stages. Additionally, seasonal spawning migrations across

depth gradients are observed globally and emphasize the significance of providing protection

across this gradient (Luo et al. 2020; Lombardo et al. 2020; Thorburn et al. 2021). We observed

a distinct break in suitable rockfish habitat within the Anacapa MPA complex, with very little

deep habitat (>70m) present. Not only does this reduce the overall available habitat for adult

rockfish, it likely reduces the potential for this MPA to provide a valuable ontogenetic bridge for

species with high habitat affinity. An MPA that contains contiguous habitats within a species full

depth range is better equipped to protect that species at all stages of life, enhancing the benefits

experienced by protection from fishing.

The protection of connected depth gradients is also likely to provide resilience for fish

populations faced with environmental stressors including seasonal changes in water temperature,

short-term temperature anomalies (marine heat waves), and long-term changes resulting from

global climate change. In an ever-warming global ocean, it has become necessary for fish

communities to adapt to rising sea temperatures, often through range shifts in latitude or depth

(Perry et al. 2005; Dahms and Killen 2023; Chaikin et al. 2022). Access to deep refuge habitat

can improve the resilience of some fish communities and allow them to survive shallow water

stressors (MacDonald et al. 2016; Pereira et al. 2018). Inversely, global climate change has been

shown to reduce deep-water dissolved oxygen, creating upward pressure on fish communities

escaping oxygen minimum zones (OMZ) (Ross et al. 2020). This has been observed around

Southern California's Channel Islands, with some rockfish communities shifting to shallower

depths as a response to OMZ shoaling (Meyer-Gutbrod et al. 2021). Spatial protections such as

MPAs can provide corridors of protected habitat across depth zones to allow for this type of

adaptive migratory process. The combined effects of large-scale environmental stressors and

species-specific life histories on habitat use across depth gradients have the potential to alter an

MPAs effectiveness in the near and long-term.

We used the same model framework across two species and one species group to

understand the importance of MPA status, habitat type, and depth in determining patterns of

abundance and biomass. Across all models, the model performance indicated that these factors

were important in explaining the observed patterns, however other aspects of the system could be

incorporated to more accurately evaluate conservation efforts. Future studies will work to

integrate factors such as distance to reserve edge, presence of other ephemeral habitats (e.g.

ophiuroid or urchin barrens), and oceanographic parameters. BRUVs are particularly well suited

to adaptive monitoring programs, in that they are deployable in multiple habitats, relatively

inexpensive to operate and provide a video record that can be reassessed over time (Langlois et

al. 2010; Whitmarsh et al. 2017).

The California MPA network will continue to work towards protecting marine

biodiversity and populations of key fished species across the state. Our results demonstrate the

realized benefits of a well-designed and well enforced protection network, with a nuanced look

at how two geographically distinct MPAs might effectively protect different fisheries targets. We

also highlight the importance of acknowledging and describing the limitations of particular

MPAs when assessing an MPA network to align expectations with realistic outcomes. Continued

monitoring efforts will provide valuable ecological knowledge as MPAs age and management

efforts adapt to preserve fisheries in the face of climate uncertainty.

Acknowledgements:

This work would not be possible without funding support from the California Ocean Protection

Council and California Seagrant (Grant Numbers R/MPA-43 and R/MPA- 48). Special thanks to

C. Honeyman, A. Parsons-field, J. Kolda, B. Brock, B. Behar, P. Campbell, W. Horstmeyer, N.

Castaneda, S. Jaeger, K. McCaffrey, S. Urgoiti Crespo, G. Hansen, J. Harding, and M.

O'Connell for assistance with fieldwork. Thank you to Y. Chen, Y. Sha, T. Hozdic, G.

Simmons, J. Zounes, and J. Eisaguirre for video analysis. Thank you to S. Gaines, D. McCauley,

R. Starr, J. Lindholm, and J. Todd for advice and insightful discussions. Thanks to A. Giraldo

Ospina and C. Free for help with data analysis.

References

Aburto-Oropeza O, Dominguez-Guerrero I, Cota-Nieto J, Plomozo-Lugo T (2009) Recruitment and ontogenetic habitat shifts of the yellow snapper (Lutjanus argentiventris) in the Gulf of California. Mar Biol 156:2461–2472. https://doi.org/10.1007/s00227-009-1271-5

Andradi-Brown DA, Macaya-Solis C, Exton DA, et al (2016) Assessing Caribbean Shallow and Mesophotic Reef Fish Communities Using Baited-Remote Underwater Video (BRUV) and Diver-Operated Video (DOV) Survey Techniques. PLOS ONE 11:e0168235. https://doi.org/10.1371/journal.pone.0168235

Baillon S, Hamel J-F, Wareham VE, Mercier A (2012) Deep cold-water corals as nurseries for fish larvae. Frontiers in Ecology and the Environment 10:351–356. https://doi.org/10.1890/120022

Bejarano S, Pardede S, Campbell SJ, et al (2019) Herbivorous fish rise as a destructive fishing practice falls in an Indonesian marine national park. Ecological Applications 29:e01981. https://doi.org/10.1002/eap.1981

Bell JJ, Micaroni V, Harris B, et al (2024) Global status, impacts, and management of rocky temperate mesophotic ecosystems. Conservation Biology 38:e13945. https://doi.org/10.1111/cobi.13945

Bellquist LF, Lowe CG, Caselle JE (2008) Fine-scale movement patterns, site fidelity, and habitat selection of ocean whitefish (*Caulolatilus princeps*). Fisheries Research 91:325–335. https://doi.org/10.1016/j.fishres.2007.12.011

Bennett K, Wilson S k., Shedrawi G, et al (2016) Can diver operated stereo-video surveys for fish be used to collect meaningful data on benthic coral reef communities? Limnology and Oceanography: Methods 14:874–885. https://doi.org/10.1002/lom3.10141

Brokovich E, Einbinder S, Shashar N, et al (2008) Descending to the twilight-zone: changes in coral reef fish assemblages along a depth gradient down to 65 m. Marine Ecology Progress Series 371:253–262. https://doi.org/10.3354/meps07591

Brown K, Monk J, Williams J, et al (2022) Depth and benthic habitat influence shallow and mesophotic predatory fishes on a remote, high-latitude coral reef. PLOS ONE 17:e0265067. https://doi.org/10.1371/journal.pone.0265067

Cappo M, Harvey E, Shortis M (2006) Counting and measuring fish with baited video techniques-an overview. In: Australian Society for Fish Biology Workshop Proceedings. Australian Society for fish biology Tasmania, pp 101–114

Cappo M, Speare P, De'ath G (2004) Comparison of baited remote underwater video stations (BRUVS) and prawn (shrimp) trawls for assessments of fish biodiversity in inter-reefal areas of the Great Barrier Reef Marine Park. Journal of Experimental Marine Biology and Ecology 302:123–152. https://doi.org/10.1016/j.jembe.2003.10.006

Caselle JE, Rassweiler A, Hamilton SL, Warner RR (2015) Recovery trajectories of kelp forest animals are rapid yet spatially variable across a network of temperate marine protected areas. Sci Rep 5:14102. https://doi.org/10.1038/srep14102

Cerrano C, Bastari A, Calcinai B, et al (2019) Temperate mesophotic ecosystems: gaps and perspectives

of an emerging conservation challenge for the Mediterranean Sea. The European Zoological Journal 86:370–388. https://doi.org/10.1080/24750263.2019.1677790

Chaikin S, Dubiner S, Belmaker J (2022) Cold-water species deepen to escape warm water temperatures. Global Ecology and Biogeography 31:75–88. https://doi.org/10.1111/geb.13414

Curley BG, Kingsford MJ, Gillanders BM (2002) Spatial and habitat-related patterns of temperate reef fish assemblages: implications for the design of Marine Protected Areas. Mar Freshwater Res 53:1197–1210. https://doi.org/10.1071/mf01199

Dahms C, Killen SS (2023) Temperature change effects on marine fish range shifts: A meta-analysis of ecological and methodological predictors. Global Change Biology 29:4459–4479. https://doi.org/10.1111/gcb.16770

Dames V, Bernard A, Floros C, et al (2020) Zonation and reef size significantly influence fish population structure in an established marine protected area, iSimangaliso Wetland Park, South Africa. Ocean & Coastal Management 185:105040. https://doi.org/10.1016/j.ocecoaman.2019.105040

Dennis D, Plagányi É, Van Putten I, et al (2015) Cost benefit of fishery-independent surveys: Are they worth the money? Marine Policy 58:108–115. https://doi.org/10.1016/j.marpol.2015.04.016

Di Lorenzo M, Claudet J, Guidetti P (2016) Spillover from marine protected areas to adjacent fisheries has an ecological and a fishery component. Journal for Nature Conservation 32:62–66. https://doi.org/10.1016/j.jnc.2016.04.004

Field SA, O'connor PJ, Tyre AJ, Possingham HP (2007) Making monitoring meaningful. Austral Ecology 32:485–491. https://doi.org/10.1111/j.1442-9993.2007.01715.x

Froese R, Pauly D (2000) FishBase 2000: Concepts Designs and Data Sources. WorldFish

Gaines SD, White C, Carr MH, Palumbi SR (2010) Designing marine reserve networks for both conservation and fisheries management. Proceedings of the National Academy of Sciences 107:18286–18293. https://doi.org/10.1073/pnas.0906473107

Gibson RN, Robb L, Wennhage H, Burrows MT (2002) Ontogenetic changes in depth distribution of juvenile flatfishes in relation to predation risk and temperature on a shallow-water nursery ground. Marine Ecology Progress Series 229:233–244. https://doi.org/10.3354/meps229233

Giraldo-Ospina A, Kendrick GA, Hovey RK (2020) Depth moderates loss of marine foundation species after an extreme marine heatwave: could deep temperate reefs act as a refuge? Proceedings of the Royal Society B: Biological Sciences 287:20200709. https://doi.org/10.1098/rspb.2020.0709

Gleason M, Fox E, Ashcraft S, et al (2013) Designing a network of marine protected areas in California: Achievements, costs, lessons learned, and challenges ahead. Ocean & Coastal Management 74:90–101. https://doi.org/10.1016/j.ocecoaman.2012.08.013

Goetze JS, Wilson S, Radford B, et al (2021) Increased connectivity and depth improve the effectiveness of marine reserves. Global Change Biology 27:3432–3447. https://doi.org/10.1111/gcb.15635

Grorud-Colvert K, Claudet J, Tissot BN, et al (2014) Marine Protected Area Networks: Assessing Whether the Whole Is Greater than the Sum of Its Parts. PLOS ONE 9:e102298.

https://doi.org/10.1371/journal.pone.0102298

Grüss A, Robinson J, Heppell SS, et al (2014) Conservation and fisheries effects of spawning aggregation marine protected areas: What we know, where we should go, and what we need to get there. ICES Journal of Marine Science 71:1515–1534. https://doi.org/10.1093/icesjms/fsu038

Halpern BS, Lester SE, McLeod KL (2010) Placing marine protected areas onto the ecosystem-based management seascape. Proceedings of the National Academy of Sciences 107:18312–18317. https://doi.org/10.1073/pnas.0908503107

Hamilton SL, Caselle JE, Malone DP, Carr MH (2010) Incorporating biogeography into evaluations of the Channel Islands marine reserve network. PNAS 107:18272–18277. https://doi.org/10.1073/pnas.0908091107

Hamilton SL, Caselle JE, Standish JD, et al (2007) Size-Selective Harvesting Alters Life Histories of a Temperate Sex-Changing Fish. Ecological Applications 17:2268–2280. https://doi.org/10.1890/06-1930.1

Harasti D, Malcolm H, Gallen C, et al (2015) Appropriate set times to represent patterns of rocky reef fishes using baited video. Journal of Experimental Marine Biology and Ecology 463:173–180. https://doi.org/10.1016/j.jembe.2014.12.003

Harms S, Winant CD (1998) Characteristic patterns of the circulation in the Santa Barbara Channel. Journal of Geophysical Research: Oceans 103:3041–3065. https://doi.org/10.1029/97JC02393

Harvey ES, McLean DL, Goetze JS, et al (2021) The BRUVs workshop – An Australia-wide synthesis of baited remote underwater video data to answer broad-scale ecological questions about fish, sharks and rays. Marine Policy 127:104430. https://doi.org/10.1016/j.marpol.2021.104430

Heyns-Veale ER, Bernard ATF, Richoux NB, et al (2016) Depth and habitat determine assemblage structure of South Africa's warm-temperate reef fish. Mar Biol 163:158. https://doi.org/10.1007/s00227-016-2933-8

Honeyman C, Carlson P, Jainese C, et al (2023) Correspondence among multiple methods provides confidence when measuring marine protected area effects for species and assemblages. Journal of Applied Ecology 60:2699–2712. https://doi.org/10.1111/1365-2664.14515

Horn MH, Allen LG (1978) A Distributional Analysis of California Coastal Marine Fishes. Journal of Biogeography 5:23–42. https://doi.org/10.2307/3038105

Howard RA, Ciannelli L, Wakefield WW, Haltuch MA (2023) Comparing fishery-independent and fishery-dependent data for analysis of the distributions of Oregon shelf groundfishes. Fisheries Research 258:106553. https://doi.org/10.1016/j.fishres.2022.106553

Jacquemont J, Loiseau C, Tornabene L, Claudet J (2024) 3D ocean assessments reveal that fisheries reach deep but marine protection remains shallow. Nat Commun 15:4027. https://doi.org/10.1038/s41467-024-47975-1

James LC, Marzloff MP, Barrett N, et al (2017) Changes in deep reef benthic community composition across a latitudinal and environmental gradient in temperate Eastern Australia. Marine Ecology Progress Series 565:35–52. https://doi.org/10.3354/meps11989

Kahng S, Copus JM, Wagner D (2017) Mesophotic Coral Ecosystems. In: Rossi S, Bramanti L, Gori A, Orejas C (eds) Marine Animal Forests: The Ecology of Benthic Biodiversity Hotspots. Springer International Publishing, Cham, pp 185–206

Karpov KA, Bergen M, Geibel JJ (2012) Monitoring fish in California Channel Islands marine protected areas with a remotely operated vehicle: the first five years. Marine Ecology Progress Series 453:159–172. https://doi.org/10.3354/meps09629

Kelaher BP, Coleman MA, Broad A, et al (2014) Changes in Fish Assemblages following the Establishment of a Network of No-Take Marine Reserves and Partially-Protected Areas. PLOS ONE 9:e85825. https://doi.org/10.1371/journal.pone.0085825

Langlois T, Chabanet P, Pelletier D, Harvey E (2006) Baited underwater video for assessing reef fish populations in marine reserves. Fisheries Newsletter-South Pacific Commission 118:53

Langlois T, Goetze J, Bond T, et al (2020) A field and video annotation guide for baited remote underwater stereo-video surveys of demersal fish assemblages. Methods in Ecology and Evolution 11:1401–1409. https://doi.org/10.1111/2041-210X.13470

Langlois TJ, Harvey ES, Fitzpatrick B, et al (2010) Cost-efficient sampling of fish assemblages: comparison of baited video stations and diver video transects. Aquatic Biology 9:155–168. https://doi.org/10.3354/ab00235

Lenihan HS, Fitzgerald SP, Reed DC, et al (2022) Increasing spillover enhances southern California spiny lobster catch along marine reserve borders. Ecosphere 13:e4110. https://doi.org/10.1002/ecs2.4110

Li L, Hollowed AB, Cokelet ED, et al (2022) Three-dimensional ontogenetic shifts of groundfish in the Northeast Pacific. Fish and Fisheries 23:1221–1239. https://doi.org/10.1111/faf.12679

Lombardo SM, Adams AJ, Danylchuk AJ, et al (2020) Novel deep-water spawning patterns of bonefish (Albula vulpes), a shallow water fish. Mar Biol 167:187. https://doi.org/10.1007/s00227-020-03799-3

Love MS (2011) Certainly more than you want to know about the fishes of the Pacific Coast: a postmodern experience. Really Big Press

Love MS, Carr MH, Haldorson LJ (1991) The ecology of substrate-associated juveniles of the genusSebastes. Environ Biol Fish 30:225–243. https://doi.org/10.1007/BF02296891

Love MS, Yoklavich M, Thorsteinson LK (2002) The rockfishes of the northeast Pacific. Univ of California Press

Love MS, Yoklavich MM (2006) CHAPTER 10. Deep Rock Habitats. In: CHAPTER 10. Deep Rock Habitats. University of California Press, pp 253–266

Luo J, Ault JS, Ungar BT, et al (2020) Migrations and movements of Atlantic tarpon revealed by two decades of satellite tagging. Fish and Fisheries 21:290–318. https://doi.org/10.1111/faf.12430

MacDonald C, Bridge TCL, Jones GP (2016) Depth, bay position and habitat structure as determinants of coral reef fish distributions: Are deep reefs a potential refuge? Marine Ecology Progress Series

561:217–231. https://doi.org/10.3354/meps11953

Maestro M, Pérez-Cayeiro ML, Chica-Ruiz JA, Reyes H (2019) Marine protected areas in the 21st century: Current situation and trends. Ocean & Coastal Management 171:28–36. https://doi.org/10.1016/j.ocecoaman.2019.01.008

Meyer-Gutbrod E, Kui L, Miller R, et al (2021) Moving on up: Vertical distribution shifts in rocky reef fish species during climate-driven decline in dissolved oxygen from 1995 to 2009. Global Change Biology 27:6280–6293. https://doi.org/10.1111/gcb.15821

Miller RR, Field JC, Santora JA, et al (2017) Spatial valuation of California marine fisheries as an ecosystem service. Can J Fish Aquat Sci 74:1732–1748. https://doi.org/10.1139/cjfas-2016-0228

Moffitt EA, Wilson White J, Botsford LW (2011) The utility and limitations of size and spacing guidelines for designing marine protected area (MPA) networks. Biological Conservation 144:306–318. https://doi.org/10.1016/j.biocon.2010.09.008

Mumby PJ, Steneck RS, Roff G, Paul VJ (2021) Marine reserves, fisheries ban, and 20 years of positive change in a coral reef ecosystem. Conservation Biology 35:1473–1483. https://doi.org/10.1111/cobi.13738

Murray SN, Ambrose RF, Bohnsack JA, et al (1999) No-take Reserve Networks: Sustaining Fishery Populations and Marine Ecosystems. Fisheries 24:11–25. https://doi.org/10.1577/1548-8446(1999)024<0011:NRN>2.0.CO;2

Navarro-Martínez ZM, de la Guardia E, Szelistowski WA, Angulo-Valdés JA (2023) Habitats diversity and MPA regulations are insufficient in promoting healthy coral reef fish assemblages in Punta Francés National Park (Cuba). Regional Studies in Marine Science 60:102826. https://doi.org/10.1016/j.rsma.2023.102826

Noble-James T, Bullimore R, McBreen F, et al (2023) Monitoring benthic habitats in English Marine Protected Areas: Lessons learned, challenges and future directions. Marine Policy 157:105852. https://doi.org/10.1016/j.marpol.2023.105852

Olson AM, Hessing-Lewis M, Haggarty D, Juanes F (2019) Nearshore seascape connectivity enhances seagrass meadow nursery function. Ecological Applications 29:e01897. https://doi.org/10.1002/eap.1897

Pereira PHC, Macedo CH, Nunes J de ACC, et al (2018) Effects of depth on reef fish communities: Insights of a "deep refuge hypothesis" from Southwestern Atlantic reefs. PLOS ONE 13:e0203072. https://doi.org/10.1371/journal.pone.0203072

Perry AL, Low PJ, Ellis JR, Reynolds JD (2005) Climate Change and Distribution Shifts in Marine Fishes. Science 308:1912–1915. https://doi.org/10.1126/science.1111322

Ponti M, Turicchia E, Ferro F, et al (2018) The understorey of gorgonian forests in mesophotic temperate reefs. Aquatic Conservation: Marine and Freshwater Ecosystems 28:1153–1166. https://doi.org/10.1002/aqc.2928

Rees M, Heyward A, Cappo M, et al (2004) Ningaloo Marine Park-Initial survey of seabed biodiversity in intermediate and deeper waters (March 2004). Australian Government, Australian Institute of

Marine Science and Natural Heritage Trust

Region M (2003) Review of some California fisheries for 2002: market squid, sea urchin, Dungeness crab, prawn, coastal pelagic finfish, albacore, ocean salmon, nearshore live-fish, Pacific herring, and recreational

Roberts CM, Andelman S, Branch G, et al (2003) Ecological Criteria for Evaluating Candidate Sites for Marine Reserves. Ecological Applications 13:199–214. https://doi.org/10.1890/1051-0761(2003)013[0199:ECFECS]2.0.CO;2

Ross T, Du Preez C, Ianson D (2020) Rapid deep ocean deoxygenation and acidification threaten life on Northeast Pacific seamounts. Global Change Biology 26:6424–6444. https://doi.org/10.1111/gcb.15307

Saarman E, Gleason M, Ugoretz J, et al (2013) The role of science in supporting marine protected area network planning and design in California. Ocean & Coastal Management 74:45–56. https://doi.org/10.1016/j.ocecoaman.2012.08.021

Sala E, Giakoumi S (2018) No-take marine reserves are the most effective protected areas in the ocean. ICES Journal of Marine Science 75:1166–1168. https://doi.org/10.1093/icesjms/fsx059

Scholz A, Steinback C, Mertens M, et al (2006) Commercial fishing grounds and their relative importance off the Central Coast of California. Report submitted to the California Marine Life Protection Act Initiative (May 4)

Sullivan-Stack J, Aburto-Oropeza O, Brooks CM, et al (2022) A Scientific Synthesis of Marine Protected Areas in the United States: Status and Recommendations. Front Mar Sci 9:. https://doi.org/10.3389/fmars.2022.849927

Swadling DS, Knott NA, Taylor MD, et al (2022) Seascape connectivity of temperate fishes between estuarine nursery areas and open coastal reefs. Journal of Applied Ecology 59:1406–1416. https://doi.org/10.1111/1365-2664.14157

Taylor BM, McIlwain JL (2010) Beyond abundance and biomass: effects of marine protected areas on the demography of a highly exploited reef fish. Marine Ecology Progress Series 411:243–258. https://doi.org/10.3354/meps08672

Thorburn J, Wright PJ, Lavender E, et al (2021) Seasonal and Ontogenetic Variation in Depth Use by a Critically Endangered Benthic Elasmobranch and Its Implications for Spatial Management. Front Mar Sci 8:. https://doi.org/10.3389/fmars.2021.656368

Topping DT, Lowe CG, Caselle JE (2005) Home range and habitat utilization of adult California sheephead, Semicossyphus pulcher (Labridae), in a temperate no-take marine reserve. Marine Biology 147:301–311. https://doi.org/10.1007/s00227-005-1573-1

White JW, Yamane MT, Nickols KJ, Caselle JE (2021) Analysis of fish population size distributions confirms cessation of fishing in marine protected areas. Conservation Letters 14:e12775. https://doi.org/10.1111/conl.12775

Whitmarsh SK, Fairweather PG, Huveneers C (2017) What is Big BRUVver up to? Methods and uses of baited underwater video. Rev Fish Biol Fisheries 27:53–73. https://doi.org/10.1007/s11160-016-

9450-1

Williams J, Jordan A, Harasti D, et al (2019) Taking a deeper look: Quantifying the differences in fish assemblages between shallow and mesophotic temperate rocky reefs. PLOS ONE 14:e0206778. https://doi.org/10.1371/journal.pone.0206778

Willis TJ, Babcock RC (2000) A baited underwater video system for the determination of relative density of carnivorous reef fish. Mar Freshwater Res 51:755–763. https://doi.org/10.1071/mf00010

Young M, Carr M (2015) Assessment of Habitat Representation across a Network of Marine Protected Areas with Implications for the Spatial Design of Monitoring. PLOS ONE 10:e0116200. https://doi.org/10.1371/journal.pone.0116200

Ziegler SL, Johnson JM, Brooks RO, et al (2023) Marine protected areas, marine heatwaves, and the resilience of nearshore fish communities. Sci Rep 13:1405. https://doi.org/10.1038/s41598-023-28507-1

Figure 1. Maps showing the study region and the two focal MPAs in the Santa Barbara Channel, Northern Channel Islands (NCI). A) shows the location of the NCI along the West Coast of North America. B) shows the NCI with red polygons delineating the NCI MPAs within the state-wide California Marine Protected Area Network. Panel C) shows Carrington Pt. MPA and associated reference area while D) shows Anacapa MPA complex and associated reference area. Both panels C) and D) have dark grey shading representing mapped hard bottom and 10m bathymetry lines in light grey. Locations of BRUV surveys by year are shown in Figure S2.

Figure 2. Stack bar plot showing the proportion of observed habitat by depth bin from Anacapa and Carrington Point. Habitat was classified from BRUV surveys and was categorized based on visible habitat in which a survey was conducted.

Figure 3. Partial model predictions from a negative binomial generalized linear model used to test the effect of MPA Status, depth, and habitat on the MaxN and biomass of targeted rockfishes for Anacapa (A) and Carrington Point (B). The blue line and shaded region represent the model predictions and 95% CI for a particular predictor variable when all other variables are held constant. The points represent the partial residuals of the model. The habitat codes 'M H' and 'M S' represent mixed hard and mixed soft habitats respectively.

Figure 4. Partial model predictions from the negative binomial generalized linear model used to test the effect of MPA Status, depth, and habitat on the MaxN and biomass of ocean whitefish for Anacapa (A) and Carrington Point (B). The blue line and shaded region

represent the model predictions and 95% CI for a particular predictor variable when all other variables are held constant. The points represent the partial residuals of the model. The habitat codes 'M H' and 'M S' represent mixed hard and mixed soft habitats respectively.

Figure 5. Partial model predictions from the negative binomial generalized linear model used to test the effect of MPA Status, depth, and habitat on the MaxN and biomass of sheephead for Anacapa (A) and Carrington Point (B). The blue line and shaded region represent the model predictions and 95% CI for a particular predictor variable when all other variables are held constant. The points represent the partial residuals of the model. The habitat codes 'M H' and 'M S' represent mixed hard and mixed soft habitats respectively.

Figure 6. Species-specific density histograms for the five targeted fish species; Copper Rockfish (A), Vermillion rockfish (B), Blue rockfish (C), ocean whitefish (D), and California sheephead (E) from Anacapa and Carrington Pt. The red density histograms represent data from within the MPA, while blue density histograms represent the data form reference areas open to fishing.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6