1   ORIGINAL RESEARCH

2   Francesca R. Napoli et al

Microphthalmia and disrupted retinal development due to a

LacZ knock-in/knock-out allele at the *Vsx2* locus

Francesca R. Napoli[1], Xiaodong Li[1], Alan A. Hurtado[2], Edward M. Levine[1,3]*

[1] Department of Ophthalmology and Visual Sciences

Vanderbilt University Medical Center, Nashville TN 37232

[2] Program in Chemical and Physical Biology

Vanderbilt University, Nashville TN 37232

[3] Department of Cell and Developmental Biology

Vanderbilt University, Nashville TN 37232

*Corresponding Author: ed.levine@vumc.org

Abstract

Visual System Homeobox 2 (*Vsx2*) is a transcription factor expressed in the developing retina that regulates tissue identity, growth, and fate determination. Several mutations in the *Vsx2* gene exist in mice, including a spontaneous nonsense mutation and two targeted missense mutations originally identified in humans. Here, we expand the genetic repertoire to include a *LacZ* reporter allele (*Vsx2^LacZ^*) designed to express beta-Galactosidase (b-GAL) and simultaneously disrupt *Vsx2* function (knock-in/knock-out). The retinal expression pattern of b-GAL is concordant with VSX2, and the mutant allele is recessive. *Vsx2^LacZ^* homozygous mice have congenital bilateral microphthalmia accompanied by defects in retinal development including ectopic expression of non-retinal genes, reduced proliferation, delayed neurogenesis, aberrant tissue morphology, and an absence of bipolar interneurons - all hallmarks of *Vsx2* loss-of-function. Unexpectedly, the mutant VSX2 protein is stably expressed, and there are subtle differences in eye size and early retinal neurogenesis when compared to the null mutant, *ocular retardation J*. The perdurance of the mutant VSX2 protein combined with subtle deviations from the null phenotype leaves open the possibility that *Vsx2^LacZ^* allele is not a complete knock-out. The *Vsx2^LacZ^* allele exhibits loss-of-function characteristics and adds to the genetic toolkit for understanding *Vsx2* function.

Keywords: LacZ, reporter knock-in, Vsx2, Chx10, retina, proliferation, lineage fidelity, neurogenesis, immunohistology

Introduction

Visual System Homeobox (VSX) genes encode paired-like homeodomain transcription factors required for visual system development in both vertebrates and invertebrates. Two paralogous genes, *Vsx1* and *Vsx2* (formerly *Chx10*), are present in vertebrates and have distinct roles in eye growth and retinal formation during development. In humans, clinical features of *Vsx1* disruption include keratoconus and visual deficits whereas *Vsx2* disruption include non-syndromic congenital bilateral microphthalmia and blindness. *Vsx2* mutations also cause non-syndromic microphthalmia in other vertebrates. Understanding how *Vsx2* functions and is regulated has provided important insights into ocular development.

Vsx2 expression is activated in the nascent retinal domain during the morphological transition from the optic vesicle to optic cup, remains expressed in retinal progenitor cells (RPCs) through development, and becomes restricted to bipolar cells and Müller glia in the adult retina.[1,2] Consistent with onset of expression, reductions in eye size and defects in retinal development commence soon after optic cup formation in murine *Vsx2* mutants[1,3] and at an equivalent stage in *Vsx2* mutant human organoids.[4] In mice, retinal defects include ectopic expression of non-retinal genes and differentiation programs more akin to neighboring tissues (lineage infidelity), slowed proliferation, delayed onset of neurogenesis, disrupted laminar morphogenesis, and a lack of bipolar cells.[1,3,5–12]

Multiple *Vsx2* alleles encoding structural or regulatory variants exist in mice (informatics.jax.org; Quick Search term: Vsx2; Category: Alleles), but most phenotypic characterizations of *Vsx2* disruption in the eye have been done with three protein

coding variants. The most extensively studied is the *ocular retardation J* allele (*orJ;*

MGI:1856112), a spontaneous mutation that replaces tyrosine at position 176 with a

premature stop codon (Fig. 1A).[1,5] Although *Vsx2* mRNA remains expressed in the *orJ*

mutant retina, VSX2 protein is not detected, and the *orJ* allele is considered a functional

null.[1,3,10] The other two alleles are gene targeted missense mutations that are

homologous to human variants associated with congenital microphthalmia and

blindness: a glutamine substitution for arginine at amino acid position 200 (*R200Q*;

MGI:5449355), and a tryptophan substitution of arginine at position 227 (*R227W*;

MGI:5449356; (Fig. 1A)).[3] Like the *orJ* allele, both are recessive, and their phenotypes

are consistent with disrupted *Vsx2* function although they differ in severity. The *R200Q*

mutant closely phenocopies *orJ,* while the *R227W* mutant is more severe as revealed

by even smaller eyes, increased pigmentation of the retina, and a failure to initiate

neurogenesis.[3] The *R200Q* and *R227W* alleles also differ from *orJ* in that the mutant

proteins are expressed.[3] The *R200Q* mutation is in the homeodomain and abrogates

DNA binding to canonical sites.[3,13] The *R227W* mutation is in the CVC domain, a

conserved stretch of approximately 50 amino acids that is a defining feature of the VSX

gene family.[3] Although its mechanism of action remains elusive, the CVC domain also

appears to promote DNA binding and sumoylation.[14,15] Consistent with its role in DNA

binding, the *R227W* protein exhibits weakened binding to canonical VSX2 binding

sites.[3]

Here, we expand the genetic toolkit for *Vsx2* with a new targeted allele that

replaces exon 3 with an IRES-*LacZ* expression cassette (*Vsx2^LacZ^*). We show that b-

GAL protein expression from the *LacZ* cassette is a reliable reporter of endogenous

VSX2 expression, and that the ocular phenotype of the *Vsx2^{LacZ}* homozygous mutant

(*LacZ* mutant; Z/Z) exhibits the hallmarks of *Vsx2* disruption in a manner similar to the

orJ mutant. However, unlike *orJ*, the predicted truncated protein is expressed and there

are subtle differences in the *LacZ* mutant phenotype – leaving open the possibility that

the allele is not a complete knockout. This allele adds to the repertoire of *Vsx2* mutants

in which protein variants are expressed and provides an additional resource for probing

Vsx2 function *in vivo.*

Results

The VSX2-encoding portion of the *Vsx2^{LacZ}* mRNA is solely composed of exons 1

and 2, therefore lacking the homeodomain and CVC domain. This is followed by an

internal ribosome entry sequence (IRES) and *LacZ* cDNA, which encodes the

Escherichia coli b-GAL protein (Fig. 1A). The *Vsx2^{LacZ}* (*LacZ*) allele was generated by

Cre recombination of the *Vsx2^{tm1a(EUCOMM)Wtsi}* allele (MGI ID:4453691),[16] and genomic

DNA PCR confirmed germline transmission (Fig. 1B). Western blot analysis of E15.5

retinal lysates probed with an antibody specific to the N-terminal portion of the human

VSX2 (aa1-131; see Table 1 for antibodies and dilutions) revealed that like the *orJ*

allele, full length VSX2 is not expressed from the *LacZ* allele (Fig. 1C; top row).

Unexpectedly, the predicted, truncated VSX2 protein from the *LacZ* allele was detected,

indicating stable expression (Fig. 1C; second row). Detection of b-GAL confirmed that

the gene targeting strategy was successful and that the IRES was sufficient to produce

two distinct proteins from the same transcript (Figure 1C; fourth row). Immunohistology

revealed that the mutant protein lacked nuclear localization, consistent with the

retention of the nuclear export signal (NES) in exon 2 and the loss of the nuclear

localization signal (NLS) in exon 3 (Fig. 1D,E).[17] We also confirmed that b-GAL

accurately reports VSX2 expression, indicated by their extensive colocalization

throughout development and easily visualized in the inner nuclear layer (INL) of *LacZ*

heterozygous retinas (*LacZ* het; *+/Z*) at P28. (Fig. 1F-I; Supplemental Fig. 1A-F).

Finally, the stereotypical microphthalmic phenotype was evident and significant at E12.5

(Fig. 1J-L; see figure legends and Supplemental Table 1 for statistics; see Methods for

quantification methods and statistical analyses).

We next performed immunohistology to characterize the *LacZ* mutant phenotype

in the retina. Establishment of optic cup regionalization appeared intact at E10.5 as

evidenced by VSX2 expression in the retina, LHX2 and PAX6 in the RPE and retina,

OTX1/2 and MITF in the RPE (Fig. 2A-E,G-K). The RPC and proliferation markers

PCNA and CCND1 also appeared to be normal in the E10.5 *LacZ* mutant retina

(Supplemental Fig. 2A-D). However, MITF was ectopically expressed in the retina (Fig.

2E,K) and PAX2 expression extended into the dorsal retina in the *LacZ* mutant (Fig.

2F,L; arrows). At E11.5, VSX2, LHX2, and PAX6 expression remained unaltered (Fig.

2M-O,S-U), but the ectopic expression of MITF (Fig. 2E,K) and dorsal expression of

Pax2 (Fig. 2R,X; arrows) were still evident, and OTX1/2 expression in the central retina

was elevated in the *LacZ* mutant (Fig. 2P,V). These altered expression patterns are

consistent with compromised retinal identity. Further evidence of altered retinal identity

was observed at E12.5 with continued ectopic expression of OTX1/2 in the peripheral

retina, most notably on the ventral side (Fig. 4E,K; arrows). From E11.5 to E15.5, the

RPC population, as indicated by PCNA, is maintained (Fig. 3A-F; Supplemental Fig.

2E,F), but proliferation progressively decreased, most notably in the peripheral retina,

as assessed by reduced CCND1 expression (Fig. 3G-L; Supplemental Fig. 2G,H;

arrows) and EdU incorporation (Fig. 3M-R; Supplemental Fig. 2I,J; arrows). This early

embryonic phenotype emulates those of the *orJ*, *R200Q,* and *R227W* mutants.[1,3]

Neurogenesis and establishment of the inner and outer neuroblastic layers

initiate at E11.5 in the central retina and progress out to the retinal peripheral margin by

E15.5. This is readily observed in the *LacZ* het retina at E12.5, E13.5, and E15.5 using

the expression of neurogenesis and cell-type specific markers including TUBB3, a

marker of cytoskeletal maturation in early post mitotic neuronal precursors, ATOH7, a

marker expressed by neurogenic progenitors preceding cell cycle exit, BHLHB5 and

POU4F2, markers of amacrine and RGCs respectively, and OTX1/2, a marker of

neurogenic RPCs biased towards photoreceptor fates as well as photoreceptor

precursors (Fig. 4A-E;M-Q; Supplemental Fig. 2K,M,O,Q,S). In contrast, none of these

neurogenesis markers were detected in the central retina of the *LacZ* mutant at E12.5

(Fig. 4G-K). OTX1/2 was expressed in the peripheral ventral retina (Fig. 4E,K; arrows),

but as mentioned above, this expression is associated with compromised retinal

identity.[7] By E15.5 in the *LacZ* mutant, all markers were expressed in patterns

consistent with neurogenesis, although stained cells were restricted to the central retina

(Fig. 4S-W; arrows). Correlated with the delayed neurogenesis is reduced expression of

the neurogenic competence marker SOX2 at E12.5 (Fig. 4F,L), and restricted

expression in a small central region of the *LacZ* mutant retina at E13.5 (Supplemental

Fig.2 S,T; arrows) and E15.5 (Fig. 4R,X; arrows).[17] These observations are consistent

with a delay in the onset of retinal neurogenesis, to a similar extent as the *orJ* and

R200Q mutants.

We observed variation in the severity of the microphthalmic phenotype of *LacZ*

mutant embryos at E15.5, ranging from similar to more severe than in *orJ* embryos (Fig.

5A). Although the mean reduction in eye size of the *LacZ* and *orJ* mutants were similar

in magnitude when compared to their respective controls (Fig. 5B, Supplemental Table

1), variability was greater in the *LacZ* mutants as determined by the coefficients of

variation (CV) for each genotype (Fig. 5C; see Methods for formula). We also observed

variation in the extent of neurogenesis, as assessed by TUBB3 expression, between

LacZ mutants (Fig. 5D-G). While neurogenic length and density (defined in methods)

were similar in the *LacZ* and *orJ* mutants (Fig. 5H,I; Supplemental Table 1), the extent

of variation in these measurements were similar except for neurogenic length in the

LacZ mutants, which had a larger CV compared to the other genotypes (Fig. 5J,K).

Linear regression analysis revealed that neurogenic density and eye size were highly

correlated (Fig. 5L), but this was not the case for neurogenic length and eye size (Fig.

5M). These observations suggest a link between eye size and neuron production, but

not between eye size and progression of the neurogenic wave from the central to

peripheral retina.

Given the variation in neurogenesis at E15.5, we hypothesized that differences

between the *LacZ* and *orJ* mutant phenotypes began earlier. We therefore focused on

the onset of neurogenesis, which exhibits a near stereotypical delay of two days in the

orJ retina, initiating at approximately E13.5.[8] Expression of TUBB3 (Fig. 5N; box and

inset) and OTX1/2 (Supplemental Fig. 2R; arrow) was observed at E13.5 in the central

retina of the *LacZ* mutant, but cells were sparse compared to the *LacZ* het

(Supplemental Fig. 2Q). Similarly, cells expressing the neurogenic progenitor marker

ATOH7 were present but sparse in the *LacZ* mutant retina (Supplemental Fig. 2K,L;

arrow), whereas POU4F2, and BHLHB5 were not detected (Supplemental Fig. 2M-P).

We therefore examined TUBB3 staining in retinas from the *LacZ* and *orJ* mutants at

E13.25. Indeed, neurogenesis was not observed at E13.25 in the *LacZ* mutant retina

(n=7) but was in the *orJ* mutant retina (n=4; Fig. 5O-Q). The sum of these observations

suggests that the *LacZ* mutant exhibits subtle increases in phenotypic severity

compared to the *orJ* mutant (see discussion).

Elevated MITF expression in the retina is a central driver of the ocular

phenotypes in *orJ, R200Q,* and *R227W* mutant mice, and a semi-dominant negative

allele of *Mitf* (*Mitf^mi*) is sufficient to significantly reduce phenotypic severity.[3,8,9,18] We

predicted a similar outcome with the *LacZ* allele, and to test this, we crossed *LacZ*

mutant mice with our *Vsx2^orJ/orJ*; *MITF^mi/+* line to generate *Vsx2^orJ/Z*; *MITF^mi/+* embryos. At

both E12.5 and E15.5, *Vsx2^orJ/Z*; *MITF^mi/+* eyes were significantly larger than those of

Vsx2^orJ/Z; *MITF^+/+* (Fig. 6). This suggests that, like the other *Vsx2* mutants, *Mitf* has a

prominent role in causing microphthalmia in the *LacZ* mutant retina.

The effects of early *Vsx2* dysfunction are compounded in the postnatal retina as

seen by severe hypocellularity and tissue disorganization. Additionally, due to the

specific postnatal requirement of *Vsx2* for the specification and differentiation of the last

neuronal cell class, bipolar cells, they are not generated.[1,6,11,19] These features are also

evident in the *LacZ* mutant retina. At P0 (Fig. 7), the *LacZ* mutant retina is reduced to a

1-3 cell-thick, epithelium consisting of dispersed RPCs and/or newly forming Müller glia

(EdU, MCM6, SOX9) and differentiating retinal neurons (TUBB3) including RGC's

(POU4F2), amacrine cells (BHLHB5), and photoreceptors (RECOVERIN, OTX1/2). At

P10 (Fig. 8), the thin apical epithelium, composed largely of photoreceptors

(RECOVERIN, OTX1/2) remains intact while an additional population of disorganized

aberrantly proliferating RPCs or Müller glia (EdU, MCM6, Sox9), and neurons (TUBB3)

including RGCs (POU4F2), and amacrine cells (BHLHB5) arises basal to the organized

epithelium. Signs of degeneration emerge at P14 (Supplemental Fig. 3), as the cellular

organization at the apical surface becomes disrupted and hypocellular regions, possibly

plexiform outgrowths, arise in the disorganized basal cell layer. By P28 (Fig. 9), the

retina has severely degenerated - consisting of disorganized retinal progenitor cells, and

specified Müller glia, RGCs, amacrine cells, and photoreceptors. Consistent with the

other *Vsx2* mutants, bipolar cells are not observed in the *LacZ* mutant retina at P14

(Supplemental Fig. 3Q-T) or P28 (Fig. 10) as revealed by PKCα (PKC) and VSX1.

Interestingly, sparse PKC-positive cells were observed at P14, which could suggest

limited bipolar cell formation, but this antibody also detects amacrine cells and likely

accounts for the pattern observed at P28.[6]

Discussion

Based on the observations reported here, we propose that the *LacZ* allele

exhibits loss-of-function properties with strong phenotypic similarities to the *orJ* null

allele, which lacks protein expression, and the *R200Q* allele, which produces a DNA

binding-deficient protein.[3] All three mutants exhibit bilateral microphthalmia, reduced

retinal proliferation, delayed neurogenesis, an absence of bipolar cells, and are affected

by ectopic Mitf activity.[3,8,9,12,18] Because the mutant VSX2 protein is expressed, we

cannot conclude that the *LacZ* allele is a complete knockout, and consistent with this,

we observed subtle differences in the timing of neurogenesis onset in the *LacZ* mutant

retinas compared to *orJ* mutant, and greater variability amongst *LacZ* mutants at E15.5

when compared to that of *orJ* mutants in both eye size and parameters of neurogenesis.

There are two reasons that could account for these differences. One is that the

mutant protein is not inert (see below). The other is that the *LacZ* mutant phenotype is

sensitive to its genetic background, which has been documented for the *orJ* mutant

phenotype. The *orJ* mutation was discovered in the 129svimj background, but when

outcrossed, phenotypic variations occur.[5,11,20] For example, increased pigmentation

and more severe microphthalmia resulted when *orJ* mice were bred with mice with

mixed genetic backgrounds including C57Bl6 mice.[21] We also observed similar

increases in phenotypic severity in an outcross with Black Swiss mice (unpublished). In

contrast, less severe phenotypes were observed when *orJ* mice were crossed to

p27Kip1 (*Cdkn1b*) knockout mice on a C57BL/6 genetic background[6] or outcrossed with

wild type *Mus castaneous*.[11] Relevant here is that our characterization of the *LacZ*

allele was done in a mixed C57BL/6;129sv mixed genetic background which could

contribute to variations in phenotypic severity.

It is also possible that the mutant VSX2 protein has activity, but what this activity

might be is not clear. The mutant coding sequence extends from amino acid 1 through

151 (out of 361 or 380; https://www.ncbi.nlm.nih.gov/protein/NP_031727) and lacks the

NLS, homeodomain and CVC domain, making direct DNA binding unlikely. The NES is

retained, however, and could contribute to non-nuclear localization.[17] Polyubiquitination

and subsequent degradation of zebrafish Vsx1 protein is enhanced by the CVC domain

and a conserved, putative PEST sequence located close to the CVC domain, and their

absence could explain the apparent stability of the mutant Vsx2 protein.[22] Interactions of

the mutant protein with other proteins is possible, but domains or motifs have not been

identified in the retained N-terminal portion of VSX2 other than putative disordered

regions**.** It is also possible that the mutant protein has novel or neomorphic activity

because it is no longer localized to the nucleus, but how this might alter the phenotype

is hard to reconcile because the allele is not dominant. Interestingly, the *R227W* allele

exhibits properties of a recessive neomorph.[3] We attributed this to the ability of the

R227W mutant protein, retaining weak DNA binding, to activate a cryptic transcriptional

circuit that increased non-retinal gene expression, including *Mitf,* beyond that observed

in the *orJ* mutant retina.[3] Further work is needed to determine if the truncated VSX2

protein from the *LacZ* allele has activity.

Limitations of the study:

It is unclear whether the subtle differences in the phenotypes between the *LacZ*

and *orJ* mutants are due to the expression of the mutant protein in the *LacZ* mutant,

differences in genetic background, or both. One approach to address this issue is to

minimize differences in genetic background by generating a congenic 129sv/imj strain

carrying the *LacZ* allele. It's also unclear if the mutant protein has activity or whether a

similar truncated protein with pathogenicity could exist in humans. This limitation could

be addressed by generating CRISPR mutations in human iPSCs that are differentiated

into retinal organoids. Finally, although eye size is improved in the combined *LacZ/orJ*

mutant with the *mi* mutation, the retinal phenotype was not characterized as part of this

study. *Mi* mutant mice with the *LacZ* allele in a homozygous state would be needed to

compare the effects of reduced Mitf activity on the *LacZ* mutant phenotype compared to

the *orJ* mutant phenotype.

In summary, the *LacZ* allele is a new addition to the allelic series of *Vsx2*

mutations, and its functional properties make it useful for further probing *Vsx2* function

in the *in vivo* context.

Experimental Procedures

Animals

279   The *Vsx2^{LacZ}* line was generated from ES cells containing the *Vsx2$^{tm1a(EUCOMM)Wtsi}$*

280 allele. Germline founders were crossed with CMV-Cre mice (RRID:IMSR_JAX:006054)

281 to excise the *LoxP* flanked region containing the Neo cassette and *Vsx2* exon 3 to

282 create the *Vsx2^{LacZ}* strain. *Vsx2^{orJ}* mice (RRID:IMSR_JAX:000395) and *Mitf^{mi}* mice

283 (RRID:IMSR_JAX001573) are maintained in the laboratory.

Tissue collection

Embryos and pups were generated using single night matings, with the day that

the vaginal plug was observed considered E0.5. *Vsx2^{LacZ}* embryos were collected at

E10.5, E11.5, E12.5, E13.5, E15.5 and *Vsx2^{LacZ}* pups were collected at P0, P10, P14,

and P28. All embryos and pups were staged using whole-body attributes.[23–25]

To assess proliferation, EdU was delivered via intra-peritoneal injection at a dose

of 30 µg/g body weight at two hours prior to euthanasia of dam for embryonic dissection

and 45 minutes prior to euthanasia for postnatal pups. Dams and pups from P10 and

older were euthanized using CO_2 and cervical dislocation; P0 pups were anesthetized

using hypothermia and thereafter euthanized via decapitation. At E15.5, embryo eyelids

were removed for imaging to allow for accurate eye size quantification.

Tissue preparation

Embryo heads and postnatal pup eyes were fixed at room temperature in 4%

PFA-PBS solution for 45 minutes or 1 hour, respectively. After several PBS washes, a

sucrose cryopreservation series was performed before the samples were frozen in

TissueTek OCT (Sakura Finetech) and stored at -80°C. Tissue was sectioned in twelve

micrometer sections at -20°C with a Leica CM1950 cryostat and adhered to Superfrost

Plus slides (Fisher Scientific). The sections were then dried for 1-2 hours before storage

at -20°C and -80°C.

PCR

DNA was extracted from embryonic tissue samples by adding 75μL alkaline lysis

buffer to sample tube, heating to 95°C for 10 minutes, cooling for 10 minutes, followed

by adding 75μL of neutralization buffer. DNA extractions were stored at -20°C.

In a 10:1 ratio, 9μL of master mix was added to 1μL extracted DNA and run in a

ProFlex thermocycler (Life Technologies).

A 1.5% agarose gel was used to run the PCR product alongside a Quick-Load

1kb Plus DNA Ladder at 70 V for 95 minutes. A UVP Gel Doc-It[TS3] Imager was used to

image the gel.

The PCR and gel electrophoresis confirming germline transmission of the LacZ

allele is a representative image from three replicates.

Western Blot

Freshly dissected E15.5 retinal tissue was flash frozen in liquid nitrogen and

stored at -80°C. Lysates were prepared with RIPA buffer and EDTA-free protease

inhibitor (Millipore Sigma) at a volume of 100 µL per 50 mg of tissue. Lysates for

heterozygotes contained 2 retinas per sample (1 animal) and 4 retinas per sample (2

animals) for homozygotes. BCA assay (Quick Start Bradford, Bio rad) was used to

determine protein concentration using a wavelength of 595 nm. 20 ug of protein per

lane was loaded in a 20 µL volume. Gels were run for 40 min at 90 V and then 40 min at

110 V. Proteins were transferred using an iBlot2 (Fisher Scientific) system with a setting

of 15V in 7 mins. Blots were incubated for 1 hour at room temperature in 1X PBS, pH

7.4, 0.1% Tween 20 containing 5% bovine serum albumin (PBST/B), followed by

incubation with constant rocking overnight at 4°C with primary antibody diluted in

PBST/B (Table 1). Excess antibody was removed, and blots were washed 3 times for

10 minutes each with PBST/B, then incubated for 1 hour at room temperature with

secondary antibody diluted in PBST (Table 1). Excess antibody was removed and blots

were washed 6 times for 10 minutes each with PBST. Chemiluminescence HRP

substrate (Immobilon Western, Millipore) was used for signal detection and images

were captured on an Amersham Imager 600 (General Electric). The western blot

presented is a representative image from 3 replicates.

333 *Immunohistochemistry and EdU detection*

To remove OCT, slides were washed in PBS and incubated in blocking solution

(PBST (0.05% Tween 20) containing 2% normal donkey serum (PBST/DS)) for 1 hour

at room temperature. Slides were then incubated overnight at 4°C in primary antibody

diluted in PBST/DS (Table 1). Excess antibody was removed, slides were washed 3

times for 5 minutes each with PBST, followed by incubation for 1 hour at room

temperature in a dark humidified chamber with secondary antibodies diluted in

PBST/DS (Table 1). Excess antibody was removed, and slides were washed 3 times for

10 minutes each with PBST. EdU incorporation was detected using AlexaFluor 568 or

647 Click-iT Cell Reaction Kit (Invitrogen-Molecular Probes, Eugene, Oregon). DAPI

(1:40000 in PBS) was used to label nuclei. After final washes in dH_2O, slides were air

dried and coverslipped with Fluoromount and stored at 4°C until imaging. All

immunohistology expression patterns are representative images from a sample size of

more than three, collected from at least two different litters for each stage and genotype.

Quantifications

Eye size was quantified using the pentagon-ROI tool of Fiji on embryo pictures

taken on a Nikon C-FIDH SMZ1270. Eye size was defined by the area of the two-

dimensional circumference of the pigmented eye (sphere). Area was measured in

$pixels^2$ and converted to μm^2. Neurogenic length, as defined by the linear length of the

retinal basal surface that TUBB3-positive cells occupy, was quantified using the

freehand line-ROI tool of Fiji and the percentage calculated by dividing the neurogenic

length by the total length of the retinal basal surface and multiplying by 100. Neurogenic

density, as defined by the number of TUBB3-positive pixels divided by the total number

of pixels in the neurogenic region (the apicobasal region of the retina containing

TUBB3-positive cells), is used as a proxy for cells within the neurogenic region. This

was quantified using Fiji by creating an ROI outlining the TUBB3-positive region of the

retina and clearing the non-ROI regions using Edit>Clear Outside. The neurogenic

region was then made binary using Threshold tool, then the number of positive (white)

and negative (black) pixels were quantified using Analyze>Histogram. The neurogenic

density was then calculated using the ratio of positive pixels to total pixels in the

neurogenic region.

Statistical Analysis

Prism was used to run all statistical analyses and to create graphical

representations. Unpaired two-tailed T-tests were used for comparisons limited to 2

conditions (Fig. 1L; Fig. 5B,H,I; Fig. 6C,F). The coefficient of variation within a sample

group was calculated using the formula CV=(Standard Deviation / Sample Mean)*100

(Fig. 5C,J,K). One way ANOVAs accompanied by post-hoc analyses (Brown-Forsythe,

Bartlett's, Tukey's multiple comparison tests) were run using Prism on the E15.5 eye

size, neurogenic length, and neurogenic density data to thoroughly compare the

heterozygous and homozygous phenotypes of the *LacZ* and *orJ* alleles (Fig. 5B,H,I).

Linear regression was used to describe the relationship between eye size and

measures of neurogenesis (neurogenic density (Fig. 5L) and neurogenic length (Fig.

5M). Supplemental Table 1 lists the results of all statistical analyses.

Imaging/Microscopy

Whole embryos were imaged using Nikon C-FIDH SMZ1270. Immunofluorescent

staining at E10.5-E11.5 and postnatal stages were imaged using a Nikon Eclipse E600

while those at E12.5-E15.5 were imaged using a Zeiss LSM710 confocal microscope

with 20X lens to capture a high-resolution image of the relatively larger ROI. Image

histograms were adjusted using Fiji. Images were resized and trimmed using Adobe

Photoshop. Figures were made using Adobe Illustrator.

Acknowledgments

Funding was provided by the National Eye Institute (R01-EY013760; P30-EY008126)

and an unrestricted grant from Research to Prevent Blindness, Inc. AAH was supported

by the National Science Foundation Graduate Research Fellows Program (1937963).

The Zeiss LSM710 confocal microscope used for this study was acquired with funding

from NIH (S10-RR027396). We thank members of the Levine and Fuhrmann

laboratories for their insights and feedback.

Ethics Statement

All procedures and experiments with mice were approved by the Vanderbilt Institutional

Animal Care and Use Committee and conformed to the ARVO guidelines for the use of

animals in vision research. The research was conducted in concordance with the NIH

guidelines for Responsible Conduct in Research.

Disclosures

The authors report no conflicts of interest in this work.

References

1. Burmeister M, Novak J, Liang MY, et al. Ocular retardation mouse caused by Chx10 homeobox null allele: impaired retinal progenitor proliferation and bipolar cell differentiation. *Nat Genet*. 1996;12(4):376-384. doi:10.1038/ng0496-376

2. Liu ISC, Chen J de, Ploder L, et al. Developmental expression of a novel murine homeobox gene (Chx10): Evidence for roles in determination of the neuroretina and inner nuclear layer. *Neuron*. 1994;13(2):377-393. doi:10.1016/0896-6273(94)90354-9

3. Zou C, Levine EM. Vsx2 Controls Eye Organogenesis and Retinal Progenitor Identity Via Homeodomain and Non-Homeodomain Residues Required for High Affinity DNA Binding. Desplan C, ed. *Plos Genet*. 2012;8(9):e1002924. doi:10.1371/journal.pgen.1002924

4. Phillips MJ, Perez ET, Martin JM, et al. Modeling human retinal development with patient-specific iPS cells reveals multiple roles for VSX2. *Stem Cells*. Published online 2014:n/a-n/a. doi:10.1002/stem.1667

5. Theiler K, Varnum DS, Nadeau JH, Stevens LC, Cagianut B. A new allele of ocular retardation: early development and morphogenetic cell death. *Anat Embryol*. 1976;150(1):85-97. doi:10.1007/bf00346288

6. Green ES, Stubbs JL, Levine EM. Genetic rescue of cell number in a mouse model of microphthalmia: interactions between Chx10 and G1-phase cell cycle regulators. *Development*. 2003;130(3):539-552. doi:10.1242/dev.00275

7. Sigulinsky CL, German ML, Leung AM, Clark AM, Yun S, Levine EM. Genetic chimeras reveal the autonomy requirements for Vsx2 in embryonic retinal progenitor cells. *Neural Dev*. 2015;10(1):12. doi:10.1186/s13064-015-0039-5

8. Leung AM, Rao MB, Raju N, et al. A framework to identify functional interactors that contribute to disrupted early retinal development in Vsx2 ocular retardation J mice. *Dev Dyn*. Published online 2023. doi:10.1002/dvdy.629

9. Horsford DJ, Nguyen MTT, Sellar GC, Kothary R, Arnheiter H, McInnes RR. Chx10 repression of Mitf is required for the maintenance of mammalian neuroretinal identity. *Development*. 2005;132(1):177-187. doi:10.1242/dev.01571

10. Rowan S, Chen CMAM, Young TL, Fisher DE, Cepko CL. Transdifferentiation of the retina into pigmented cells in ocular retardation mice defines a new function of the homeodomain gene Chx10. *Development*. 2004;131(20). doi:10.1242/dev.01300

11. Bone Larson C, Basu S, Radel JD, et al. Partial rescue of the ocular retardation phenotype
by genetic modifiers. *J Neurobiol*. 2000;42(2):232-247. doi:10.1002/(sici)1097-
4695(20000205)42:2<232::aid-neu7>3.0.co;2-4

12. Bharti K, Liu W, Csermely T, Bertuzzi S, Arnheiter H. Alternative promoter use in eye
development: the complex role and regulation of the transcription factor MITF. *Development*.
2008;135(6):1169-1178. doi:10.1242/dev.014142

13. Percin EF, Ploder LA, Yu JJ, et al. Human microphthalmia associated with mutations in the
retinal homeobox gene CHX10. *Nat Genet*. 2000;25(4):397-401. doi:10.1038/78071

14. Kurtzman AL, Schechter N. Ubc9 interacts with a nuclear localization signal and mediates
nuclear localization of the paired-like homeobox protein Vsx-1 independent of SUMO-1
modification. *Proc Natl Acad Sci USA*. 2001;98(10):5602-5607. doi:10.1073/pnas.101129698

15. Dorval KM, Bobechko BP, Ahmad KF, Bremner R. Transcriptional activity of the paired-
like homeodomain proteins CHX10 and VSX1. *J Biol Chem*. 2005;280(11):10100-10108.
doi:10.1074/jbc.m412676200

16. Skarnes WC, Rosen B, West AP, et al. A conditional knockout resource for the genome-wide
study of mouse gene function. *Nature*. 2011;474(7351):337-342. doi:10.1038/nature10163

17. Knauer SK, Carra G, Stauber RH. Nuclear Export Is Evolutionarily Conserved in CVC
Paired-Like Homeobox Proteins and Influences Protein Stability, Transcriptional Activation, and
Extracellular Secretion. *Mol Cell Biol*. 2005;25(7):2573-2582. doi:10.1128/mcb.25.7.2573-
2582.2005

18. Konyukhov B, Sazhina M. Interaction of the genes of ocular retardation and microphthalmia
in mice. *Folia Biol (Praha)*. 1966;12(2):116-123.
http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=4958088&retmode=r
ef&cmd=prlinks

19. Goodson NB, Kaufman MA, Park KU, Brzezinski JA. Simultaneous deletion of Prdm1 and
Vsx2 enhancers in the retina alters photoreceptor and bipolar cell fate specification, yet differs
from deleting both genes. *Development*. 2020;147(13):dev190272. doi:10.1242/dev.190272

20. Wong G, Conger SB, Burmeister M. Mapping of genetic modifiers affecting the eye
phenotype of ocular retardation (Chx10or-J) mice. *Mamm Genome*. 2006;17(6):518-525.
doi:10.1007/s00335-005-0159-z

21. Rowan S, Cepko CL. Genetic analysis of the homeodomain transcription factor Chx10 in the
retina using a novel multifunctional BAC transgenic mouse reporter. *Dev Biol*. 2004;271(2):388-
402. doi:10.1016/j.ydbio.2004.03.039

22. Kurtzman AL, Gregori L, Haas AL, Schechter N. Ubiquitination and degradation of the
zebrafish paired-like homeobox protein VSX-1. *J Neurochem*. 2000;75(1):48-55.
doi:10.1046/j.1471-4159.2000.0750048.x

23. Wong MD, Eede MC van, Spring S, et al. 4D atlas of the mouse embryo for precise
morphological staging. *Development*. 2015;142(20):3583-3591. doi:10.1242/dev.125872

24. Theiler K. *The House Mouse: Atlas of Embryonic Development*. Springer, Berlin; 1989.
doi:10.1007/978-3-642-88418-4

25. Kaufman MH. *The Atlas of Mouse Development*. Elsevier Science; 1992.
https://books.google.com/books?id=Lo5pAAAAMAAJ

26. Clark AM, Yun S, Veien ES, et al. Negative regulation of Vsx1 by its paralog Chx10/Vsx2 is
conserved in the vertebrate retina. *Brain Res*. 2008;1192:99-113.
doi:10.1016/j.brainres.2007.06.007

Table 1: Primary Antibodies

NAME	SPECIES[1]	DILUTION[2]	SOURCE	CATALOG #	RRID
ATOH7	Rb	1:200 (IH)	Novus Biologicals	NBP1-88639	AB_11034390
b-GAL	Ms	1:200 (IH) 1:1000 (WB)	Developmental Studies Hybridoma Bank	40-1a-s	AB_528100
BHLHB5	Rb	1:200 (IH)	Santa Cruz Biotechnology	sc-6045	AB_2065343
CCND1	Rb	1:400 (IH)	Abcam	ab16663 (SP4)	AB_443423
LHX2	Rb	1:300 (IH)	GeneTex	GTX129241	AB_2783558
MCM6	Gt	1:300 (IH)	Santa Cruz Biotechnology	sc-9843	AB_2142543
MITF	Ms	1:800 (IH)	Exalpha Biologicals Inc.	X1405M	-
OTX1/2	Gt	1:1000 (IH)	GeneTex	AF1979	AB_2157172
PAX2	Rb	1:100 (IH)	Covance Laboratories	PRB-276P	AB_291611
PAX6	Rb	1:300 (IH)	BioLegend	901301	AB_2565003
PCNA	Ms	1:500 (IH)	Santa Cruz Biotechnology	sc-56	AB_628110
PKC	Rb	1:200 (IH)	Sigma-Aldrich	P-4334	AB_477345
POU4F2	Gt	1:200 (IH)	Santa Cruz Biotechnology	sc-6026	AB_673441
RECOVERIN	Rb	1:100 (IH)	Chemicon International	AB5585	AB_2253622
SOX2	Rb	1:400 (IH)	Abcam	ab97959	AB_2341193
SOX9	Rb	1:1000 (IH)	EMD Serono Inc	AB5535	AB_2239761
TUBB3	Rb	1:40K (IH)	BioLegend	802001	AB_2564645
VSX1	Rb	1:250 (IH)	Levine Lab[26]	-	-
VSX2 - NT	Sh	1:400 (IH) 1:1000 (WB)	Exalpha Biologicals Inc.	X1180P	AB_2314191
ATP5A1	Rb	1:2000 (WB)	Proteintech	14676-1-AP	

[1] Gt, goat; Ms, mouse; Rb, rabbit; Sh, sheep
[2] IH, immunohistology; WB, western blot

484   **Table 2: Secondary Antibodies**

NAME	DILUTION[1]	SOURCE	CATALOG #	RRID
Alexa Fluor™ 488 donkey anti-rabbit IgG (H+L)	1:1000 (IH)	Invitrogen	A21206	AB_2535792
Alexa Fluor™ 568 donkey anti-rabbit IgG (H+L)	1:1000 (IH)	Invitrogen	A10042	AB_2534017
Alexa Fluor™ 647 donkey anti-rabbit IgG (H+L)	1:1000 (IH)	Invitrogen	A31573	AB_2536183
Alexa Fluor® 488 Affinipure Donkey Anti-Mouse IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	715-545-150	AB_2340846

Alexa Fluor® 594 Affinipure Donkey Anti-Mouse IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	715-585-150	AB_2340854
Alexa Fluor® 647 Affinipure Donkey Anti-Mouse IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	715-605-150	AB_2340862
Alexa Fluor® 488 Affinipure Donkey Anti-Goat IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	705-545-147	AB_2336933
Alexa Fluor® 594 Affinipure Donkey Anti-Goat IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	705-585-147	AB_2340433
Alexa Fluor® 647 Affinipure Donkey Anti-Goat IgG (H+L)	1:1000 (IH)	Jackson ImmunoResearch Laboratories Inc.	705-605-147	AB_2340437
Peroxidase AffiniPure™ Donkey Anti-Goat IgG (H+L)	1:1000 (WB)	Jackson ImmunoResearch Laboratories Inc.	705-035-147	AB_2313587
Anti-Rabbit IgG (H+L), HRP Conjugate	1:1000 (WB)	Promega	W4011	AB_430833
Anti-Mouse IgG (H+L), HRP Conjugate	1:1000 (WB)	Promega	W4021	AB_430834
[1] IH, immunohistology; WB, western blot				

Abbreviations:

b-GAL, beta-Galactosidase; C, cornea; CM, ciliary margin; D, dorsal; E, embryonic;

EdU, 5-Ethynyl-2'-deoxyuridine; GCL, ganglion cell layer; het, heterozygous; INL, inner

nuclear layer; IRES, internal ribosome entry sequence; LacZ, *Escherichia coli LacZ*

gene; L, lens; *orJ*, *ocular retardation J*; ONL, outer nuclear layer; P, postnatal; R, retina;

RGC, retinal ganglion cell; RPC, retinal progenitor cell; RPE, retinal pigmented

epithelium; V, ventral.

Figure Legends

Figure 1. *Vsx2^{LacZ}* recombination confirmation and gross phenotype. (A)

Schematic of *Vsx2^{LacZ}* allele and resultant mRNA following Cre recombination. **(B)**

Genomic DNA PCR shows mendelian inheritance of *Vsx2* wildtype and *LacZ*

sequences confirming germline transmission. **(C)** Western blot analysis of E15.5 retinal

lysates. The N-terminal specific VSX2 antibody was used in the top 2 rows to detect full

length and truncated VSX2 proteins. B-GAL antibody was used in 4[th] row to detect b-

GAL expression from *LacZ allele*. VSX2 and b-GAL antibodies were used on separate

blot. The probing for ATP5A1 serves as a loading control for each blot. **(D-E)**

Immunohistology at E15.5 reveals continued yet differentially localized expression of

VSX2 in the *LacZ* mutant (retinas are contained within the dashed lines). **(F-I)**

Recombination reporter, b-GAL, is co-expressed with VSX2 in the *LacZ* heterozygote

INL at P28. **(J-L)** Microphthalmia phenotype of the *LacZ* mutant is observable and

significant at E12.5 (****$p<0.0001$, T-test, sample sizes: 9 (*Z/+*), 10 (*Z/Z*)).

Abbreviations: ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion cell

layer. Scale bars: (D-H) 100 μm; (J,K) 2000 μm.

Figure 2. Regionalization of the early embryonic *LacZ* mutant eye. (A-L) Markers of

eye field specification and regionalization at E10.5. All markers exhibit similar

expression patterns in the *LacZ* mutant compared to the *LacZ* het except for MITF,

which is ectopically expressed in the retina (E,K) and PAX2, which is expanded into the

dorsal retina (arrows; F,L). **(M-X)** Markers of eye field specification and regionalization

at E11.5. MITF continues to be ectopically expressed (Q,W) and PAX2 extends into the

dorsal retina of the *LacZ* mutants (arrows; R,X). All other markers were similar between

the *LacZ* het and mutant. Abbreviations: R, retina; RPE, retinal pigment epithelium; CM,

ciliary margin; L, lens; C, cornea. Scale bar: 100 μm.

Figure 3. Proliferation is reduced in the embryonic *LacZ* mutant retina. (A-F) The

general proliferation marker PCNA remains expressed throughout the *LacZ* mutant

retina, similar to the *LacZ* het at E11.5, E12.5, and E15.5. In contrast, the patterns of

CCND1 expression **(G-L)** and EdU incorporation **(M-R)** became progressively restricted

to the central retina in the *LacZ* mutants compared to *LacZ* hets, most notably at E12.5

and E15.5. Arrows in I-L and O-R point to the regions where staining is absent in the

LacZ mutant retina compared to the *LacZ* het. Scale bar: 100 µm.

Figure 4. Retinal neurogenesis is delayed and centrally restricted in the *LacZ*

mutant. Immunohistology assessing cytoskeletal maturation with TUBB3 **(A,G,M,S)**, the

presence of neurogenic progenitors with ATOH7 **(B,H,N,T)**, retinal ganglion cells with

POU4F2 **(C,I,O,U)**, amacrine cells with BHLHB5 **(D,J,P,V)** and photoreceptor

precursors with OTX1/2 (**E,K,Q,W)**, reveal a delay in neurogenesis in the *LacZ* mutant

retina, with markers lacking at E12.5 and centrally restricted at E15.5 (arrows in S-W).

OTX1/2 staining also reveals ectopic staining in the ventral peripheral retina at E12.5

(compare red arrows in E and K), which is associated with compromised retinal identity.

(F,L,R,X) Expression of neurogenic competence regulator SOX2 in RPCs is reduced at

E12.5 (F,L) and centrally restricted at E15.5 in the *LacZ* mutant retina (R,X; arrow

denotes region of persistent SOX2 expression). Scale bar: 100 µm.

Figure 5. Timing of neurogenesis initiation and E15.5 variation in *LacZ* mutant

deviate from the *orJ* mutant phenotype. (A) Microphthalmia was similar in the *LacZ*

and *orJ* mutants at E15.5, but severity was variable in the *LacZ* mutants (compare iii

and iv). **(B,C)**. Quantifications of eye size revealed similar degrees of severity (B;

****p<0.0001; ANOVA; sample sizes: 11 (*Z/+*), 7 (*Z/Z*), 11 (*orJ/+*), 6 (*orJ/orJ*)), but the

coefficient of variation (CV) was highest in the *LacZ* mutants compared to the other

genotypes (C). **(D-G)** TUBB3 expression, indicating neurogenesis, is centrally restricted

and seemingly covaries with microphthalmia severity in the *LacZ* mutant at E15.5. **(H,I)**

Neurogenic density (H); and length (I) are similarly decreased in the *LacZ* and *orJ*

mutants (****$p<0.0001$; ***$p<0.001$; $^{ns}p>0.05$; ANOVA; sample sizes: 5 (*Z/+* and *orj/+*

combined (Het)), 7 (*Z/Z*), 4 (*orJ/orJ*)). **(J,K)**. CVs for neurogenic density were similar

across genotypes (J), but the CV for neurogenic length was specifically increased in the

LacZ mutant (K). **(L)** Linear regression analysis of eye size and neurogenic density in

LacZ mutant retinas at E15.5 ($R^2=0.944$; $F=83.44$; $p=0.0003$; sample size: 7). **(M)**

Linear regression analysis of eye size and neurogenic length in *LacZ* mutant retinas at

E15.5 ($R^2=0.364$; $F=2.856$; $p=0.1518$; sample size: 7). **(N-Q)** Neurogenesis initiates at

E13.5 in *LacZ* mutant (compare N to P). In contrast, neurogenesis initiates in the *orJ*

retina by E13.25 (Q). Scale bars: (A) 500 µm; (D-G; N-Q) 100 µm.

Figure 6. Functional rescue of embryonic *LacZ* mutant microphthalmic phenotype

with *Mitf^{mi/+}* allele. The semi-dominant negative allele *MITF^{mi/+}* reduced the severity of

microphthalmia in *Vsx2^{orJ//Z}* mutants at E12.5 **(A-C)** and E15.5 **(D-F)**. ****$p<0.0001$; T-

test; sample sizes for E12.5: 10 (*orJ/Z;+/+*),20 (*orJ/Z;mi/+*); sample sizes for E15.5: 14

(*orJ/Z;+/+*), 22 (*orJ/Z;mi/+*)). Scale bar: 1000 µm.

Figure 7. **Immunohistology at P0.** The *LacZ* mutant retina at P0 is reduced to a thin

epithelium, containing interspersed **(A-D,G,H)** proliferating RPCs (EdU, MCM6, SOX9).

(E-P) Similarly, markers of differentiating cells (TUBB3, POU4F2, BHLHB5, OTX1/2,

RECOVERIN are expressed in a disorganized manner. In addition to marking RPCs,

SOX9 could indicate precocious Müller glia differentiation. Scale bar: 100 µm.

Figure 8. Immunohistology at P10. (A-L) A disorganized population of proliferating

cells (EdU, MCM6), new neurons (TUBB3, POU4F2, BHLHB5), and Müller glia (SOX9),

persist in the *LacZ* mutant retina. (**M-P)** In contrast, the photoreceptor markers OTX1/2

and RECOVERIN suggest the presence of an organized apical epithelium. Scale bar:

100 µm.

Figure 9. Immunohistology at P28. The *LacZ* mutant retina at P28 is severely

degenerated and disorganized. Lacking cellular organization, spherical **(A-D)** aberrantly

proliferating cells are interspersed with **(E-P)** progenitors (MCM6, SOX9), differentiating

neuronal cells (TUBB3), Müller glia (SOX9), retinal ganglion cells (POU4F2), amacrine

cells (BHLHB5), and photoreceptors (OTX1/2, RECOVERIN). Scale bar: 100 µm.

Figure 10. Bipolar cells are not detected in the *LacZ* mutant retina. (A-D) The rod-

bipolar marker PKC is not detected in the *LacZ* mutant retina at P10, but basally

displaced cells are observed at P28. This antibody also detects amacrine cells at this

age. **(E,F)** The cone-bipolar marker VSX1 is not detected in the *LacZ* mutant retina at

P10 or P28. Scale bar: 100 µm.

Supplemental Figure 1. Expression of VSX2 and b-GAL in postnatal *LacZ* het

retina. (A,B) Immunohistology reveals b-GAL to be an accurate reporter of VSX2

expression with both expressed throughout the *LacZ* het retina at P0. **(C-F)** At P10 and

P14, VSX2 and b-GAL are restricted to the INL, consistent with expression in bipolar

cells and Müller glia. All VSX2 and b-GAL stains are from the same animal for each

respective stage and genotype. Scale bar: 100 μm.

Supplemental Figure 2. Immunohistology at E10.5 and E13.5. (A-D) Expression of

the proliferation markers PCNA, CCND1, and pHH3 in the *LacZ* mutant retina are

normal at E10.5. **(E-J)** At E13.5, PCNA extends throughout the *LacZ* mutant retina

(E,F), whereas CCND1 and EdU are limited to the central retina (G-J). Arrows in G-J

point to areas where staining is absent in the *LacZ* mutant compared to the *LacZ* het.

(K,L) ATOH7 expression is detected in a few cells in the central retina of the *LacZ*

mutant (arrow). **(M-P)** Retinal ganglion cells (POU4F2) and amacrine cells (BHLHB5)

are not detected in the *LacZ* mutant retina at E13.5. **(Q,R)** OTX1/2 expression is

detected in a few cells in the central retina of the *LacZ* mutant (arrow). **(S,T)** SOX2

expression is reduced in the peripheral retina of the *LacZ* mutant. Arrows in S and T

point to areas where staining is absent in the *LacZ* mutant compared to the *LacZ* het.

Scale bars: 100 μm.

Supplemental Figure 3. *LacZ* mutant retina at P14. (A-F) As the P14 retina continues

to aberrantly proliferate (EdU, MCM6) and give rise to new neurons (TUBB3), signs of

degeneration emerge. **(C-P)** The previously organized apical epithelium composed

largely of progenitors (MCM6, SOX9), Müller glia (SOX9), and photoreceptors (OTX1/2,

RECOVERIN) begins to seemingly lose its integrity as amongst the disorganized basal

cell population, consisting of progenitors (MCM6, SOX9), Müller glia (SOX9), retinal

ganglion cells (POU4F2) and amacrine cells (BHLHB5), hypocellular regions, possibly

plexiform outgrowths, encroach. **(Q,R)** PKC expressing cells are present in the *LacZ*

mutant retina, but their identity is not known, with possibilities being amacrine cells or

remnant bipolar cells. **(S,T)** VSX1 expressing cells are not detected in the *LacZ* mutant

retina. Scale Bar: 100 µm.

Supplemental Figure 4. Uncropped images for western blots in Figure 1C. (A) Blot

1 probed with N-terminal VSX2 antibody, which detects the full length (VSX2 WT) and

truncated (VSX2 Mut) proteins. The last 4 lanes in panel A were used for the first two

rows in Fig. 1C. **(B)** Blot 1 stripped and re-probed with antibody to a validated reference

protein ATP5A1. The last 4 lanes in panel B were used for the third row in Fig. 1C. **(C)**

Blot 2 probed with b-GAL antibody. Spacings between lanes were cropped and the

bands at 100 kDa correspond to the fourth row in Fig. 1C. **(D)** Blot 2 was stripped and

reprobed with ATP5A1 antibody. Spacings between lanes were cropped and the bands

at 50 kDa correspond to the fifth row in Fig. 1C.

A

FRT

| En2 SA | IRES | LacZ | pA |

Vsx2^LacZ Gene

Homeodomain   CVC Domain

orJ   R200Q   R227W

EXON 1 | INTRON 1 | EXON 2 | INTRON 2 | EXON 3 | INTRON 3 | EXON 4 | INTRON 4 | EXON 5

Vsx2^LacZ mRNA

5' UTR | EXON 1 | EXON 2 | IRES | LacZ | 3'UTR & pA

B

+/+   +/Z   Z/Z

700 bp — *LacZ*

300 bp — *Vsx2* WT

C

+/+   +/Z   Z/Z   orJ

50 kDa — VSX2 WT
25 kDa — VSX2 Mut
50 kDa — ATP5A1
100 kDa — bGal
50 kDa — ATP5A1

D +/Z   **E** Z/Z

VSX2

F VSX2   **G** bGal   **H** VSX2/bGal   **I**

+/Z

ONL
INL
GCL

J +/Z   **K** Z/Z

E12.5

L

4×10⁵
3×10⁵
2×10⁵
1×10⁵

Eye Size (μm²)

****

+/Z   Z/Z
E12.5