MANUSCRIPT TITLE: Mosquito population dynamics is shaped by the interaction among

larval density, season, and land use

AUTHORS: Nicole Solano[1,2], Emily C. Herring[3], Carl W. Hintz[4], Philip M. Newberry[1,2],

Annakate M. Schatz[1,2], Joseph W. Walker[2†], Craig W. Osenberg[1], Courtney C. Murdock[1,2‡]

AFFILIATIONS:

1. Odum School of Ecology, University of Georgia, Athens, GA, USA

2. Center for Ecology of Infectious Diseases, University of Georgia, Athens, GA, USA

3. Department of Population Health, College of Veterinary Medicine, University of

Georgia, Athens, GA, USA

4. Department of Applied Ecology, North Carolina State University, Raleigh, NC, USA

Corresponding Author: Correspondence to Nicole Solano (nicole.n.solano@gmail.com)

KEY WORDS

Aedes albopictus; Density-Dependence*;* Fitness*;* Per Capita Growth Rate*;* Seasonality*;* Life

History Traits

† Current address: Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT, USA

‡ Current address: Department of Entomology, College of Agriculture and Life Sciences, Cornell University, Ithaca, NY, USA

<u>DATA AVAILABILITY STATEMENT</u>

The author is still cleaning the code that support the findings of this study for publication

purposes, but data and code will be openly available on Figshare if the paper is accepted.

ABSTRACT (English)

Understanding how variation in key abiotic and biotic factors interact at spatial scales

relevant for mosquito fitness and population dynamics is crucial for predicting current and future

mosquito distributions and abundances, and the transmission potential for human pathogens.

However, studies investigating the effects of environmental variation on mosquito traits have

investigated environmental factors in isolation or in laboratory experiments that examine

constant environmental conditions that often do not occur in the field. To address these

limitations, we conducted a semi-field experiment in Athens, Georgia using the invasive Asian

tiger mosquito (*Aedes albopictus*). We selected nine sites that spanned natural variation in

impervious surface and vegetation cover to explore effects of the microclimate (temperature and

humidity) on mosquitoes. On these sites, we manipulated conspecific larval density at each site.

We repeated the experiment in the summer and fall. We then evaluated the effects of land cover,

larval density, and time of season, as well as interactive effects, on the mean proportion of

females emerging, juvenile development time, size upon emergence, and predicted per capita

population growth (i.e., fitness). We found significant effects of larval density, land cover, and

season on all response variables. Of most note, we saw strong interactive effects of season and

intra-specific density on each response variable, including a non-intuitive decrease in

development time with increasing intra-specific competition in the fall. Our study demonstrates

that ignoring the interaction between variation in biotic and abiotic variables could reduce the

accuracy and precision of models used to predict mosquito population and pathogen transmission

dynamics, especially those inferring dynamics at finer-spatial scales across which transmission

and control occur.

ABSTRACT (Spanish)

Para poder predecir la distribución y abundancia de las poblaciones de mosquitos y la

transmisión potencial de patógenos a humanos, es crucial comprender cómo factores abióticos y

bióticos clave para el éxito reproductivo y la dinámica poblacional de los mosquitos interactúan a

escalas relevantes. Sin embargo, los estudios que han investigado los efectos de variables

ambientales en las características demográficas de los mosquitos han considerado su efecto de

forma aislada o en experimentos de laboratorio bajo condiciones ambientales constantes que, a

menudo, no reflejan lo que ocurre en el campo. Para abordar estas limitaciones, llevamos a cabo

un experimento de semi-campo en Athens, Georgia, utilizando el mosquito invasor tigre asiático

(*Aedes albopictus*). Seleccionamos nueve sitios que abarcaban variaciones naturales en la

superficie impermeable y cobertura vegetal para explorar los efectos del microclima

(temperatura y humedad) en los mosquitos. También manipulamos la densidad de larvas de tigre

asiático en dos experimentos que fueron realizados en el verano y otoño. Evaluamos los efectos

de la cobertura vegetal, la densidad de larvas, la temporada climática, y la interacción entre estas

variables en la proporción de hembras que emergieron, el tiempo de desarrollo de las larvas, el

tamaño al momento de la emergencia, y el crecimiento demográfico per cápita previsto (éxito

reproductivo). Encontramos efectos significativos de la densidad de larvas, la variación de la

cobertura vegetal y la estación del año en todas las variables de respuesta. Más notablemente,

observamos un fuerte efecto de la interacción entre la temporada climática y la densidad de

larvas en todas las variables de respuesta, incluyendo una disminución no intuitiva en el tiempo

de desarrollo con el aumento de la competencia intraespecífica en el otoño. Nuestro estudio

evidencia que ignorar la interacción entre variables abióticas y bióticas podría reducir la

exactitud y precisión de los modelos utilizados para predecir las dinámicas de las poblaciones de

mosquitos, y por tanto, de la transmisión de patógenos. Esto, especialmente en modelos que

infieren estas dinámicas a escalas espaciales más finas, en las cuales ocurre la transmisión y el

control.

<u>INTRODUCTION</u>

Arthropod-borne viruses (arboviruses) pose significant global health and economic

burdens (Espinal et al. 2019; Girard et al. 2020; Ogunlade et al. 2021). Some of the most notable

arboviruses (dengue, Chikungunya, yellow fever, Ross River, and Zika) are transmitted by *Aedes*

mosquitoes and often cause chronic morbidity and high mortality for people infected. *Aedes-*

borne arboviruses continue to expand, which is partially attributed to increasing urbanization,

climate change, and the geographical spread of *Aedes* mosquitoes (Bhatt et al. 2013; Wilder-

Smith et al. 2017). To better anticipate areas and times of seasons most at risk for vector-borne

pathogen transmission, target vector control efforts, and reduce the incidence of these pathogens,

it is critical to understand how global change influences mosquito population dynamics.

Multiple environmental factors (abiotic and biotic) affect mosquito demography, vital

rates (e.g., larval development, growth, and survival), and population dynamics. Key abiotic

factors include variation in temperature, relative humidity, and precipitation, while biotic factors

include variation in intra- and inter-species competition, availability of resources, and presence /

density of biological enemies (predators and parasites). In addition, these environmental factors

can exhibit considerable spatial heterogeneity (Cohen et al. 2016). When this spatial variation is

ignored, predictions of mosquito population dynamics averaged over space or time can lead to

considerable bias if the effects of these factors are nonlinear (Bozinovic et al. 2011; Martin and

Huey 2008; Ruel and Ayres 1999). As a result, mathematical models might overlook the

influence of hot spots or ecological traps on mosquito population dynamics.

Given the effects that global change (e.g., urbanization and climate change) could have

on the environment and thus mosquito-borne pathogen transmission and public health, numerous

studies have focused on the effects of abiotic factors, especially temperature, on mosquitoes

(Mordecai et al. 2019; Nik Abdull Halim et al. 2022; Reinhold et al. 2018; Shocket et al. 2020).

Temperature has non-linear (unimodal) effects on mosquito traits and their ability to transmit

pathogens (Mordecai et al. 2017). Shifts in temperature can alter mosquito distributions,

consequently changing the spatial and temporal patterns of environmental suitability for

mosquito-borne pathogen transmission (Tesla et al. 2018; Miazgowicz et al. 2020; Wimberly et

al. 2020). Climate change is expected to decrease suitability for transmission in some areas

while increasing it in others (Lafferty and Mordecai 2016; Ryan et al. 2021; 2023). Therefore,

the incorporation of abiotic factors (e.g., temperature and precipitation) into predictive models

has advanced our understanding of mosquito population dynamics (Alto and Bettinardi 2013;

Evans et al. 2019; Murdock et al. 2017), pathogen transmission, and overall disease risk

(Messina et al. 2016; Mordecai et al. 2013; Zhang et al. 2017).

The effect of biotic factors, however, is generally excluded from species distribution

models and mechanistic models of mosquito-borne disease transmission, despite the fact that at

fine scales (e.g., a backyard) biotic factors likely determine mosquito abundances (Beale et al.

2014; Ewing et al. 2019; Golding et al. 2015; Leach et al. 2016). In the field, mosquito larvae

cohabitate with conspecifics and/or heterospecifics at varying densities. The competition and

resource limitation that results from these interactions reduces larval nutrition and can have

lasting effects on mosquito survival, reproduction, and generation time by slowing development,

reducing adult size, and lowering teneral reserves (e.g., lipids and proteins) of adult mosquitoes

going through final stages of maturation) (Briegel 2003). There has been a large body of work

investigating the effects of conspecific larval densities on various metrics of mosquito fitness

(e.g., development rate, body size, survival, reproduction). Increases in larval densities typically

decrease mosquito fitness and its components (Breaux et al. 2014; Schneider et al. 2000;

Yoshioka et al. 2012); similarly, larvae reared in environments with high nutrient availability

tend to emerge as larger, more fecund adults than larvae reared in low nutrient environments.

Differences in mosquito body condition can also translate into differences in vector competence

(Alto and Bettinardi 2013; Alto et al. 2005; 2007) and vectorial capacity (Moller-Jacobs et al.

2014; Shapiro et al. 2016).

However, there has been little investigation of the potential interactive effects of abiotic

and biotic factors on mosquito population dynamics, which likely occur in the field. For

example, the relationship between mosquito body size and fecundity (number of eggs produced

in the first gonotrophic cycle) is modulated by temperature (Ezeakacha and Yee 2019; Reiskind

and Zarrabi 2012) and food quality (Costanzo et al. 2018). Interactions in the field could be

mediated by temperature-dependent growth of microbial populations, for example, that

mosquitoes feed on in the juvenile environment (Ratkowsky et al. 1982), which in turn could

exacerbate or dampen the effects of competition on mosquito population dynamics. Finally,

most studies that have considered the interaction between abiotic (i.e., temperature, relative

humidity) and biotic factors (i.e., intraspecific competition) on mosquito life history traits and

population dynamics have been conducted in the lab under constant environmental conditions

that may not be reflective of field environments. Instead, abiotic variables like temperature, for

example, vary diurnally, seasonally, and across space (e.g., due to variation in land use and land

cover) (Wimberly et al. 2020; Murdock et al. 2017; Evans et al. 2018; 2019).

To improve our understanding of how field-relevant variation in key abiotic and biotic

factors potentially interact at spatial scales relevant for mosquito fitness and population

dynamics, we conducted a semi-field experiment using the Asian tiger mosquito (*Aedes*

albopictus) system in Athens, Georgia during summer and fall 2017. *Ae. albopictus* is a highly

invasive mosquito in the United States and other areas of the world, as well as a competent

vector of arboviruses including dengue and chikungunya (Benedict et al. 2007; Bonizzoni et al.

2013; Gratz 2004; Paupy et al. 2009). We selected nine sites that spanned natural variation in

impervious surface and vegetation cover, which was expected to lead to variation in the

microclimate experienced by mosquitoes based on established relationships between land use,

land cover, temperature, and relative humidity (Murdock et al. 2017; Wimberly et al. 2020). On

these sites we also manipulated conspecific larval density. We then measured the effects of

variation in land cover, larval density, time of season, and potential interactions on larval

development and survival, adult size and emergence, and predicted per capita population growth

(i.e., fitness) of *Ae. albopictus*. Understanding the drivers of spatial and temporal heterogeneity

of *Ae. albopictus* abundance is crucial for predicting and managing its future invasive potential

and understanding population dynamics of *Ae. albopictus* and other species with similar ecology

(e.g., *Aedes aegypti)*, which play an important role in human arbovirus transmission.

METHODS

Experimental design

We conducted the study between June 2017 and October 2017 in Athens-Clarke County,

Georgia, USA. Following previous work (Evans et al. 2018; Murdock et al. 2017), we used an

impervious surface and vegetation cover map to select nine 30 x 30 m sites – three each with low

(0-5%), intermediate (6-40%), and high (41-100%) areal cover of impervious surfaces, which we

refer to as rural, suburban, and urban land use types respectively (Figure S1) (Evans et al. 2019;

Murdock et al. 2017). Within each of the nine sites, we placed three trays in full shade separated

by approximately 6-9 m. Each tray contained four wide-mouth mason jars containing 300mL of

leaf infusion (see Murdock et al. 2017). Each jar was assigned a density treatment (0, 30, 60, or

120 *Ae. albopictus* first instar larvae from a laboratory colony). We covered the top of the jars

with mesh netting to prevent emerging adults from escaping (and other adults from ovipositing)

and staked a wire cage with a vinyl roof over the trays to the ground to prevent changes in

habitat volume from rainfall, nutrients due to falling vegetation, and loss from animal

disturbance.

To understand how seasonal variation in abiotic factors may alter relationships between

conspecific larval density and traits governing mosquito population dynamics, we conducted this

experiment in the summer (June 11 - July 25) and fall (September 8 - October 18) 2017. Jars

were visited daily, at the same time of day, until no emergence was observed within the previous

24 hours. During each visit and for each jar we recorded the developmental stage of a typical

mosquito (i.e., larvae, pupae, adult), number of emerged adults, and any observations that might

have affected the results (e.g., fallen tree branch). If adults emerged, we collected them with a

mouth aspirator, placed them into a 50 mL falcon tube labeled with the respective jar ID, and

then placed the tubes on ice. We transported mosquitoes to the laboratory and stored them at -

20°C until we determined their sex (Darsie and Ward 2005) and wing length.

Microclimate variables

To measure variation in microclimate, we deployed one data logger (Monarch

Instruments, Amherst, NH, USA: Radio Frequency Identification (RFID) Temperature Track-It

Logger) to each tray in a separate jar filled with leaf infusion but no larvae. We also uniformly

distributed six data loggers throughout each site at approximately 0.5-1.0 m above the ground

and in full shade (Murdock et al. 2017). Loggers recorded air or water temperature and relative

humidity every ten minutes. Microclimate data were downloaded monthly and were summarized

by recording the daily minimum (T_{min}), daily maximum (T_{max}), and daily average (T_{mean}) for

each logger. We also calculated the diurnal range in temperature (T_{range}) and relative humidity

(R_{range}) as the difference between the observed daily minimum and maximum for each logger

(e.g., $T_{range} = T_{max} - T_{min}$; $R_{range} = R_{max} - R_{min}$).

Mosquito life history traits

In this study, we only report female data since females (not males) are responsible for egg

production, population growth rates, and pathogen transmission. Therefore, emergence,

development time, wing length, fecundity, and population growth rates all refer to female

individuals (unless otherwise noted). To calculate emergence, we divided the total number of

females that emerged by half the total number of 1st instar larvae placed into the habitat

(assuming half of the larvae were female). Juvenile development time was defined as the day an

adult female emerged (since the start of the experiment). One wing was removed from each

individual, mounted onto a glass slide using clear nail polish, and measured with a dissecting

microscope and micrometer. Wings were measured from the alular notch to the apex of the

wing, excluding fringe setae (Packer and Corbet 1989). Finally, we used the data collected in

this experiment to estimate the per capita growth rates of mosquito populations in response to

our experimental treatments by using Livdahl & Sugihara's (1984) equation 7:

206
$$r' = \frac{ln\left(\frac{1}{N_0} \sum_x A_x \; f(w_x)\right)}{D + \frac{\sum_x xA_x \; f(w_x)}{\sum_x A_x \; f(w_x)}} \tag{1}$$

where N_0 is the initial number of females in a cohort (which we assume to be half the initial

number of larvae in the jar (i.e., 15, 30, or 60)), A_x is the number of females that emerged on day

x, D is the time to reproduction following emergence (which we assume to be 14 days: Lounibos

et al. 2002). The fecundity $f(w_x)$ of females that emerged on day x, with an average wing length

of w_x, is estimated from the linear relationship between wing length and number of eggs laid

from Lounibos et al. (2002):

213
$$f(w_x) = -121.240 + (78.02 * w_x) \tag{2}$$

Data Analyses

To assess how daily microclimate conditions (mean, minimum and maximum

temperature: T_{mean}, T_{min}, T_{max}; and relative humidity: R_{mean}, R_{min}, R_{max}) varied with season and

land use, we used generalized linear mixed models (GLMMs) with lognormal distributions and

log link functions. All models included season, land-use type, and their interaction as fixed

effects with site, logger, and day as random effects. We also used GLMMs to determine how

emergence, larval development rate, wing length, and per capita growth rate varied across

seasons, land use types, and density treatments. Those models all had season, land use type,

density treatment, and their interactions as fixed effects, with site, tray, and jar as random effects.

All analyses were conducted in R version 4.3.3 (R Core Team 2024). Models were fit using

the *glmmTMB* package (Brooks et al. 2017), and corrected Akaike Information Criterion (AICc)

values were used to determine which models described the data best. One-way analysis of

variance (ANOVA) was conducted for all models using the *car* package (Fox and Weisberg

2019). If models showed statistically significant effects, we performed a post-hoc analysis using

Tukey's multiple pair-wise comparison of means [*emmeans* package (Lenth 2023)] to identify

differences between independent variables. We assessed the scaled residuals of models for

overdispersion and divergence from normality using the DHARMa package (Hartig 2022).

Figure 1. Temperature (ºC) (top panels) and relative humidity (%) (bottom panels) at three sites

in each of three land use types (rural = blue, suburban = gray, and urban = red) in two seasons

(summer 2017 [left panels] and fall 2017 [right panels]). Mean microclimates across the three

land use types are in solid lines, with daily maxima and minima indicated by the dashed lines.

RESULTS

Microclimate

Five of the 54 temperature loggers placed in larval environments failed (three in the

summer and two in the fall), and 36 of the 108 loggers placed in terrestrial environments failed

or were stolen (16 in the summer and 20 in the fall). However, microclimate variation across

season and land use types was well represented by the remaining loggers and failure rates were

roughly equivalent across sites. Diurnal temperature and relative humidity metrics all varied

significantly with season (Table S1). Summer temperatures (i.e., mean daily T_{min}, T_{mean}, and

T_{max}) were higher than fall temperatures (Figure 1, Tables 1 & S1). Relative humidity (mean

daily R_{min}, R_{mean}, and R_{max}) was also greater in the summer than fall (although R_{max} was only

significantly greater in summer urban sites) (Figure 1, Tables 1 & S1). Seasonal variation in

diurnal ranges for each abiotic factor also occurred. Mosquitoes in the summer experienced

lower fluctuations throughout the day in temperature (T_{range}) and relative humidity (R_{range}) than in

the fall (Table 1). Finally, we observed significant effects of land use on diurnal temperature and

relative humidity metrics (Table S1). Urban sites had significantly warmer daily average

minimum temperatures (T_{min}) than rural sites overall (Table 1) and were less humid than

suburban and rural sites (i.e., lower R_{min}, R_{mean}, and R_{max}: Table 1). As a result, urban sites

experienced a wider diurnal range in relative humidity (R_{range}) than rural sites (Table 1).

In addition to the main effects of season and land use on mosquito microclimate

variables, we observed a significant interaction between season and land use on average daily

T_{max} and T_{range} (Table S1). T_{max} was lower in suburban sites in the summer, but higher in the fall,

compared to rural and urban sites (Table 1). Urban sites also had marginally lower average

diurnal temperature ranges (T_{range}) than rural sites, but only in the fall season (Table 1). We did

not observe an interactive effect of season and land use on metrics of relative humidity (Table

S1).

Emergence

Of the 5670 first instar *Ae. albopictus* placed into the field during each season, a total of

3156 (56%) mosquitoes (males and females combined) successfully emerged during the summer

while only 1233 (38%) mosquitoes (males and females combined) emerged in the fall. We

observed significant main effects of season and conspecific density on the proportion of female

first instar larvae that emerged as adults (i.e., survived through the larval and pupal stages; Table

S2). Females began emerging earlier in the summer trial (day 9) than in the fall trial (day 13),

and there was a significantly higher proportion of first instar larvae that survived to emerge as

adult females from the low density treatment relative to medium and high density treatments

(Table 2).

We also observed significant interactions between season and density as well as land use

type and density on adult female emergence (Table S2). For example, there was no significant

difference in the survival of female larvae in the low-density treatments in the summer and fall;

however, the negative effects of increasing conspecific density on the larval survival was much

greater in the fall (Figure 2a, Table 3). Further, the proportion of females that emerged from

larval habitats in each conspecific density treatment varied across land use, with the proportion

of females that emerged at low conspecific densities being highest in suburban sites. Habitats

with intermediate conspecific densities on urban sites experienced the highest survival. Finally,

for larval habitats with high conspecific densities, rural sites had the highest juvenile survival

(Figure 3).

Figure 2. Mixed effects model prediction of the (**a**) the proportion of female *Ae. albopictus*

larvae that emerged (i.e., survival probability), (**b**) mean female *Ae. albopictus* development time

(days), (**c**) mean wing length (mm), and (**d**) per capita intrinsic growth rate, r' (days^{-1}) for each

density treatment (low = 30 larvae, medium = 60 larvae, high = 120 larvae) across two seasons

(summer = red, fall = orange). Each box represents the interquartile range (25th to 75th

percentile), with the median (black horizontal line), whiskers extend to the extremes or 1.5 times

the interquartile range beyond the interquartile (whichever is smaller), and the dots indicate any

values beyond the whiskers.

Figure 3. Mixed effects model prediction of the proportion of female *Ae. albopictus* larvae that

emerged from each density treatment (low = 30 larvae, medium = 60 larvae, high = 120 larvae)

across each land use type (rural = blue, suburban = gray, urban = red). Each box represents the

interquartile range (25th to 75th percentile), with the median (black horizontal line), whiskers

extend to the extremes or 1.5 times the interquartile range beyond the interquartile (whichever is

smaller), and the dots indicate any values beyond the whiskers.

Female Development Time

Both time of season and conspecific larval density significantly affected the average

amount of time required for adult female emergence (Table S2), with females developing more

quickly in the summer relative to the fall and from the low-density treatment (Tables 2 & S2,

Figure 2b). However, a significant interaction between season and larval density demonstrates

that the main effect of density is being driven by the relationship between mean female

development time (time to emergence) and larval density in the summer. Females reared in the

low-density treatment were the fastest to develop in the summer but were the slowest to develop

in the fall (Figure 2b, Table 3).

Wing Length

We observed significant main effects of season and density on female wing length (Table

S2). Females reared in the summer had longer wings compared to the fall, and females

developing in the high conspecific larval treatment had significantly shorter wings than

mosquitoes from the low-density treatment (Figure 2c). We did not observe a significant

interaction between these two factors on female wing size (Table S2).

Per Capita Population Growth Rate

Integrating these responses revealed strong effects of season, conspecific larval density,

and the interaction between these two factors on the per capita population growth rate, r' (Figure

2d, Table S2). In general, the per capita growth rate was greater in the summer relative to the

fall and at higher conspecific densities (Table 2). A significant interaction between season and

density was observed due to greater declines in per capita growth rate with increasing

conspecific density in the fall relative to the summer (Figure 2d, Table 3). Additionally, we

observed negative per capita growth rates at the highest conspecific density in the fall, while all

other combinations of factors led to positive projected per capita growth (Figure 2d).

DISCUSSION

To date, the majority of studies investigating the effects of various abiotic and biotic

factors on mosquito fitness, population dynamics, vector competence, and transmission potential

investigate the effect of varying a single environmental variable such as temperature (e.g., Liu-

Helmersson et al. 2014; Agyekum et al. 2022; Monteiro et al. 2007), conspecific density (e.g.,

Hawley 1985; Sauers et al. 2022; Reiskind and Lounibos 2009), and food resources (e.g., Levi et

al. 2014; Telang and Wells 2004). Further, these studies are often conducted in laboratory

environments across a range of constant conditions that explore a wider range of environmental

conditions than what might be experienced in the field. However, mosquito populations in the

field will be subject to the effects of many different environmental variables that fluctuate

diurnally and seasonally, and likely interact. Previous work in the *Ae. albopictus* system

suggests that microclimate variation in the field (rather than coarser environmental

manipulations in the lab) can translate into significant variation in mosquito life history traits,

population dynamics, and arbovirus transmission potential (Evans et al. 2018; 2019; Murdock et

al. 2017; Wimberly et al. 2020). Our experimental study builds on these results by showing that

variation in microclimate conditions (temperature and relative humidity) with season and land

use can interact with variation in key biotic variables (conspecific density) to affect metrics of

mosquito fitness (larval survival and development time, female size at emergence, and per capita

growth rates). The results from this study have broad implications for our understanding and

ability to predict the effects of current and future environmental variation on mosquito fitness,

population dynamics, and the environmental suitability for mosquito-borne disease transmission.

The effects of land use on microclimate, metrics of mosquito fitness, and population

dynamics

Similar to previous studies conducted in the same region (Athens, Georgia) (Wimberly et

al. 2020; Murdock et al. 2017; Evans et al. 2018), urban sites had higher minimum temperatures

than rural sites and were drier than non-urban land use types. This is consistent with the Urban

Heat Island (UHI) effect, where nighttime temperatures (analogous to T_{min}) are 1-2.5°C higher

(Hibbard et al. 2017) than surrounding areas due to the higher percentage of impervious surfaces

in urban areas that absorb, store, and re-emit heat (e.g., asphalt and concrete). However, in non-

urban areas, pervious materials that absorb yet do not re-emit heat (e.g., mulch, soil, vegetation),

have higher evapotranspiration rates, and provide ample shading resulting in lower temperatures

and higher relative humidity than urban areas (Oke 1982; Bowler et al. 2010; Huang et al. 1987;

Jabbar et al. 2023). Despite the variation in microclimate observed across land use, this variation

for the most part did not affect metrics of mosquito fitness or predicted per capita growth rates.

This is somewhat different from previous studies in this system (Murdock et al. 2017; Evans et

al. 2018) and others (Waldock et al. 2013; Li et al. 2014) that found microclimate variation

across land use to significantly affect the proportion of adults emerging, mosquito development

rate, and the ability of mosquitoes to transmit dengue-2 virus. However, this is not necessarily

surprising as the previous effects observed were subtle and likely due to the fact that Athens-

Clarke County is a small city in Georgia, with less variation in impervious surface cover,

vegetation cover, and ambient temperatures than seen in larger cities in the U.S. and worldwide

(Peng et al. 2011; Oke 1973; Zhou et al. 2017; Imhoff et al. 2010). We did observe effects of

land use on the relationship between conspecific density and the proportion of adult females

emerging, suggesting the effects of intra-specific competition could potentially be modified

across an urban gradient. However, while significant, these effects were subtle.

The effects of seasonality on microclimate, metrics of mosquito fitness, and population

dynamics

Seasonal variation in microclimate was much larger than what we observed across land

use, with temperature and relative humidity being higher during the summer field trial relative to

the fall. Our finding of a higher relative humidity in the summer compared to the fall is

consistent with a previous study in the same system (Evans et al. 2018), probably due to higher

levels of precipitation during the summer. As a result, we observed strong seasonal effects on

the proportion of females emerging from the larval environment, the juvenile development time,

their body size upon emergence, and predicted per capita growth rates from larval habitats. This

is not unexpected considering mosquitoes are ectothermic, and trait performance has been shown

to have non-linear relationships with temperature variation across a diversity of mosquito species

(reviewed by: Mordecai et al. 2019). For example, laboratory work conducted under a range of

constant temperatures predicts that *Ae. albopictus* larval survival and mosquito development

rates should be optimized at ~25°C and ~35°C, respectively (Mordecai et al. 2017). Thus, it is

unsurprising that we generally saw faster larval development times and increased adult female

emergence in the warmer summer field trial (with temperatures ranging from 25-25.7°C) than the

cooler fall (with temperatures ranging from 21.1-21.6°C) across our sites.

As a result of these seasonal effects on metrics of mosquito fitness, we also observed

higher per capita growth rates in the summer relative to the fall. These results support the

findings of previous work in this system, that found similar seasonal trends in adult female

emergence, juvenile development times on these sites, and predicted mosquito population

dynamics (Evans et al. 2018; 2019, but see Murdock et al. 2017 which saw reduced female

emergence in the summer relative to the fall overall). Interestingly, despite warmer summer

temperatures and the increased juvenile development times in the summer relative to the fall,

mosquitoes emerged at larger sizes. This is counter to what would be expected from previous

work in this system (Murdock et al. 2017, but see Evans et al. 2018 that found no effect of

season on wing size) and other studies (Ezeakacha and Yee 2019; Lounibos et al. 2002; Briegel

and Timmermann 2001), but could be due to seasonality modifying the effect of conspecific

density on the juvenile development time and larval survival (proportion of adult females

emerging).

Seasonality shapes the relationship between intra-specific competition and metrics of

mosquito fitness and population dynamics

The effects of increasing conspecific larval density on larval survival, larval development

time, size upon emergence, and predicted per capita growth rates have been documented

extensively to date. In general, our results support previous research that shows increasing

conspecific larval densities result in decreased juvenile survival, smaller adults upon emergence,

and decreased per capita growth rates (Walker et al. 1991; Sauers et al. 2022; Lord 1998).

However, our study highlights a critical and novel role of seasonality and land use in modifying

the effects of conspecific larval density on mosquito fitness and population dynamics.

In both the summer and fall trial, we observed a negative effect of increasing conspecific

larval density on larval survival (i.e., the proportion of adult females emerging). However, this

trend was much stronger during the fall trial. This could be due to a hurricane that occurred at

the beginning of the study (day three) when larvae were in their most vulnerable stage (i.e., first

instar) and/or due to cooler fall temperatures being below the thermal optimum for *Ae.*

albopictus larval survival (~25°C, Mordecai et al. 2017). For example, mean temperatures in the

fall were <15°C for three consecutive days. These temperatures, which can result in high *Ae.*

albopictus larval and pupal mortality (reviewed in Waldock et al. 2013), combined with the

additional stress of developing under high intra-specific competition, could work synergistically

to increase juvenile mortality.

Contrary to our hypothesis and the multitude of prior studies (e.g., Sauers et al. 2022;

Lord 1998; Ezeakacha and Yee 2019), larvae developed faster with higher conspecific densities

in the fall. Several factors could explain this result. First, the timing of larval mortality is known

to affect the amount of resources available to the surviving larvae (Neale and Juliano 2019;

McIntire and Juliano 2018; Evans et al. 2022). If mortality of early instar larvae was elevated by

cooler fall temperatures and high intra-specific densities, the surviving larvae would have

experienced reduced intra-specific competition and increased access to food resources, both of

which allow larvae to develop faster (Evans et al. 2022; Neale and Juliano 2019; Teng and

Apperson 2000). Stage-specific overcompensation and the Hydra effect have been detected in

Ae. albopictus when early larval mortality occurred (McIntire and Juliano 2018). Another

possible explanation may be survival bias. Since very few females survived in the fall from the

medium-density treatment (7.2%) and even fewer from the high-density treatment (1.2%), the

results are biased to those few individuals that did survive; these survivors might have been

444 individuals of higher fitness quality allowing them to persist and develop in more competitive

445 conditions. A third explanation could be that larvae facultatively alter their development rate

446 based on a stressor such as low temperatures and/or low quantity of nutritional resources due to

447 increased competition. While not demonstrated before in *Ae. albopictus*, previous work in the

448 *Ae. triseriatus* and *Ae. communis* systems have shown larvae can speed up development rates in

449 environments subject to evaporation (Chodorowski 1969; Juliano and Stoffregen 1994).

450  Based on the extensive literature documenting relationships between mosquito wing size

451 (i.e. size upon emergence) with variation in conspecific density (Alto et al. 2005; Lord 1998;

452 Evans et al. 2021) or temperature (Kingsolver and Huey 2008; Rueda et al. 1990), we observed

453 unexpected, seasonal effects on these relationships. While wing size upon emergence decreased

454 with increasing conspecific density as previous work has shown, this effect was stronger in the

455 warmer summer than the cooler fall. Temperature effects have also been well studied with

456 warmer temperatures resulting in smaller wings and body size–"hotter is smaller"–and cooler

457 temperatures resulting in larger wings and body size–"cooler is bigger" hypothesis (Ezeakacha

458 and Yee 2019; Kingsolver and Huey 2008; Rueda et al. 1990). Our results contradict the "cooler

459 is bigger" hypothesis since mosquitoes developing in the summer had larger wings than those

460 reared in the fall. Only a few studies have demonstrated smaller wing lengths at cooler

461 temperatures (Reiskind and Zarrabi 2012; Lounibos et al. 2002). The result could be attributed

462 to slower microbial growth in the fall (due to cooler temperatures) limiting the amount of food

463 available to larvae combined with anabolism being less efficient at low temperatures (Reiskind

464 and Janairo 2018). Thus, females in the summer may have access to more microbially rich

465 habitats regardless of larval density treatment as well as a greater capacity to incorporate

466 nutrients into tissue, allowing them to develop more quickly in the summer and emerge as larger

adults. Finally, like development time, mosquito size upon emergence converges at high

densities regardless of season, suggesting that high amounts of intra-specific competition could

eventually outweigh seasonal variation in abiotic factors such as temperature and relative

humidity on mosquito body size and condition.

In general, while per capita growth rates declined as larval conspecific densities increased

(see also: Lord 1998; Edgerly and Livdahl 1992; Reiskind and Lounibos 2009) we found the per

capita growth rates in the summer to be higher and less affected by intra-specific competition

than in the fall. This is likely because mosquitoes in the summer (relative to the fall) survived

better, developed more rapidly, and emerged as larger adults with higher predicted fecundity

despite increases in intra-specific competition. Most studies to date have considered the effects

of temperature or conspecific density on growth rates in isolation, without considering the

possible interaction. Those that do consider the effects of temperature on the relationship

between intra-specific competition and per capita growth rates have generally found intra-

specific competition to intensify as temperatures warm (Evans et al. 2021; Teng and Apperson

2000), which is not what we observed in this study.

CONCLUSION

This study has several important implications. It is generally thought that variation in

biotic factors (e.g., intra-specific and inter-specific competition, quality and quantity of habitat,

biological enemies) will vary more and thus exert their effects on organismal fitness and

population dynamics at fine-spatial scales relative to abiotic factors (e.g., temperature,

precipitation, relative humidity) that have more regional effects (Wiens 1989; McGill 2010;

Russell et al. 2006; Sandel and Smith 2009). This study demonstrates that while variation in

abiotic factors at fine-spatial scales (e.g., microclimate) had relatively little effect, we do see

important effects of season, which represents variation in abiotic factors but at a longer temporal

scale. The experimentally imposed variation in density (a biotic factor) had much greater effects

on mosquito traits and interacted with the abiotic factors. Thus, ignoring the interaction between

variation in biotic and abiotic variables could reduce the accuracy and precision of models used

to predict mosquito population and disease transmission dynamics, especially those inferring

dynamics at finer-spatial scales across which transmission and control occur. By not accounting

for these interactions, mismatches in model predictions could result in unexpected variation in

mosquito densities spatially and seasonally, as well as unexplained carry-over effects of how

these variables interact to affect mosquito fitness and vectorial capacity (Moller-Jacobs et al.

2014; Shapiro et al. 2016; Alto et al. 2007).

AUTHOR CONTRIBUTIONS

CCM designed the experiment. NS, ECH, CWH, PMN, AMS, and JWW conducted fieldwork

for the study. NS conducted the statistical analyses, with guidance from CCM and CWO. NS

drafted the manuscript with guidance from CCM and CWO. All authors were involved in

reviewing and editing the manuscript. All authors read and approved the final manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank members of the Murdock lab (UGA and Cornell), Osenberg lab (UGA), and Drake lab

(UGA) for their helpful discussions and field support. We particularly thank Sydney Habegger,

Luis Hernandez, and Abigail LeCroy for their assistance with fieldwork, mounting, and

measuring wings, Juliana Hoyos and Maria Luisa Müller Theissen for their feedback on the

Spanish abstract, and Ashutosh Pathak for his thoughtful discussion and guidance with the

statistical analyses. We also thank the property owners who provided us access to the study

sites.

FUNDING INFORMATION

This work was supported by the National Science Foundation Research Experiences for

Undergraduates (Grant no: 1659683), the National Science Foundation Research Traineeship:

Interdisciplinary Disease Ecology Across Scales (Grant no: 1545433), the National Institutes of

Health Office of Research Infrastructure Programs (Grant No: 2T35OD010433-11), and the

start-up resources provided by the University of Georgia.

<u>REFERENCES</u>

Agyekum, Thomas P., John Arko-Mensah, Paul K. Botwe, Jonathan N. Hogarh, Ibrahim Issah,

Duah Dwomoh, Maxwell K. Billah, Samuel K. Dadzie, Thomas G. Robins, and Julius N.

Fobil. 2022. "Effects of Elevated Temperatures on the Growth and Development of Adult

Anopheles Gambiae (s.l.) (Diptera: Culicidae) Mosquitoes." *Journal of Medical*

Entomology 59 (4): 1413–20. https://doi.org/10.1093/jme/tjac046.

Alto, Barry W., and David Bettinardi. 2013. "Temperature and Dengue Virus Infection in

Mosquitoes: Independent Effects on the Immature and Adult Stages." *The American*

Journal of Tropical Medicine and Hygiene 88 (3): 497. https://doi.org/10.4269/ajtmh.12-

0421.

Alto, Barry W., L. Philip Lounibos, Stephen Higgs, and Steven A. Juliano. 2005. "Larval

Competition Differentially Affects Arbovirus Infection in Aedes Mosquitoes." *Ecology*

86 (12): 3279–88. https://doi.org/10.1890/05-0209.

Alto, Barry W., L. Philip Lounibos, Christopher N. Mores, and Michael H. Reiskind. 2007.

"Larval Competition Alters Susceptibility of Adult Aedes Mosquitoes to Dengue

Infection." *Proceedings of the Royal Society B: Biological Sciences* 275 (1633): 463.

https://doi.org/10.1098/rspb.2007.1497.

Beale, Colin M., Mark J. Brewer, and Jack J. Lennon. 2014. "A New Statistical Framework for

the Quantification of Covariate Associations with Species Distributions." *Methods in*

Ecology and Evolution 5 (5): 421–32. https://doi.org/10.1111/2041-210X.12174.

Benedict, Mark Q., Rebecca S. Levine, William A. Hawley, and L. Philip Lounibos. 2007.

"Spread of The Tiger: Global Risk of Invasion by The Mosquito Aedes Albopictus."

Vector-Borne and Zoonotic Diseases 7 (1): 76–85.

https://doi.org/10.1089/vbz.2006.0562.

Bhatt, Samir, Peter W. Gething, Oliver J. Brady, Jane P. Messina, Andrew W. Farlow, Catherine

L. Moyes, John M. Drake, et al. 2013. "The Global Distribution and Burden of Dengue."

Nature 496 (7446): 504–7. https://doi.org/10.1038/nature12060.

Bonizzoni, Mariangela, Giuliano Gasperi, Xioaguang Chen, and Anthony A. James. 2013. "The

Invasive Mosquito Species Aedes Albopictus: Current Knowledge and Future

Perspectives." *Trends in Parasitology* 29 (9): 460–68.

https://doi.org/10.1016/j.pt.2013.07.003.

Bowler, Diana E., Lisette Buyung-Ali, Teri M. Knight, and Andrew S. Pullin. 2010. "Urban

Greening to Cool Towns and Cities: A Systematic Review of the Empirical Evidence."

Landscape and Urban Planning 97 (3): 147–55.

https://doi.org/10.1016/j.landurbplan.2010.05.006.

Bozinovic, Francisco, Daniel A. Bastías, Francisca Boher, Sabrina Clavijo-Baquet, Sergio A.

Estay, and Michael J. Angilletta. 2011. "The Mean and Variance of Environmental

Temperature Interact to Determine Physiological Tolerance and Fitness." *Physiological*

and Biochemical Zoology 84 (6): 543–52. https://doi.org/10.1086/662551.

Breaux, Jennifer A., Molly K. Schumacher, and Steven A. Juliano. 2014. "What Does Not Kill

Them Makes Them Stronger: Larval Environment and Infectious Dose Alter Mosquito

Potential to Transmit Filarial Worms." *Proceedings of the Royal Society B: Biological*

Sciences 281 (1786): 20140459. https://doi.org/10.1098/rspb.2014.0459.

Briegel, H., and S. E. Timmermann. 2001. "Aedes Albopictus (Diptera: Culicidae):

Physiological Aspects of Development and Reproduction." *Journal of Medical*

Entomology 38 (4): 566–71. https://doi.org/10.1603/0022-2585-38.4.566.

Briegel, Hans. 2003. "Physiological Bases of Mosquito Ecology." *Journal of Vector Ecology:*

Journal of the Society for Vector Ecology 28 (1): 1–11.

Brooks, Mollie E., Kasper Kristensen, Koen J. van Benthem, Arni Magnusson, Casper W. Berg,

Anders Nielseon, Hans J. Skaug, Martin Maechler, and Benjamin M. Bolker. 2017.

"glmmTMB Balances Speed and Flexibility Among Packages for Zero-Inflated

Generalized Linear Mixed Modeling." *The R Journal* 9 (2): 378–400.

https://doi.org/10.32614/RJ-2017-066.

Chodorowski, A. 1969. "The Desiccation of Ephemeral Pools and the Rate of Development in

Aedes Communis Larvae." *Polish Archive of Hydrobiology* 16:79–91.

Cohen, Jeremy M., David J. Civitello, Amber J. Brace, Erin M. Feichtinger, C. Nicole Ortega,

Jason C. Richardson, Erin L. Sauer, Xuan Liu, and Jason R. Rohr. 2016. "Spatial Scale

Modulates the Strength of Ecological Processes Driving Disease Distributions."

Proceedings of the National Academy of Sciences 113 (24): E3359–64.

https://doi.org/10.1073/pnas.1521657113.

Costanzo, Katie S., Katie M. Westby, and Kim A. Medley. 2018. "Genetic and Environmental

Influences on the Size-Fecundity Relationship in Aedes Albopictus (Diptera: Culicidae):

Impacts on Population Growth Estimates?" *PLOS ONE* 13 (8): e0201465.

https://doi.org/10.1371/journal.pone.0201465.

Darsie, Richard F., and Ronald A. Ward. 2005. *Identification and Geographical Distribution of*

the Mosquitos of North America, North of Mexico. University Press of Florida.

https://search.ebscohost.com/login.aspx?direct=true&AuthType=ip,shib&db=cat06564a

&AN=uga.9931074443902959&site=eds-live&custid=uga1.

Edgerly, Janice S., and Todd P. Livdahl. 1992. "Density-Dependent Interactions Within a

Complex Life Cycle: The Roles of Cohort Structure and Mode of Recruitment." *Journal*

of Animal Ecology 61 (1): 139–50. https://doi.org/10.2307/5517.

Espinal, Marcos A., Jon K. Andrus, Barbara Jauregui, Stephen Hull Waterman, David Michael

Morens, Jose Ignacio Santos, Olaf Horstick, Lorraine Ayana Francis, and Daniel Olson.

2019. "Emerging and Reemerging Aedes-Transmitted Arbovirus Infections in the Region

of the Americas: Implications for Health Policy." *American Journal of Public Health* 109

(3): 387. https://doi.org/10.2105/AJPH.2018.304849.

Evans, Katherine G., Zoey R. Neale, Brendan Holly, Cecilia C. Canizela, and Steven A. Juliano.

2022. "Survival-Larval Density Relationships in the Field and Their Implications for

Control of Container-Dwelling Aedes Mosquitoes." *Insects* 14 (1): 17.

https://doi.org/10.3390/insects14010017.

Evans, Michelle V., John M. Drake, Lindsey Jones, and Courtney C. Murdock. 2021. "Assessing

Temperature-Dependent Competition between Two Invasive Mosquito Species."

Ecological Applications 31 (5): e02334. https://doi.org/10.1002/eap.2334.

Evans, Michelle V., Carl W. Hintz, Lindsey Jones, Justine Shiau, Nicole Solano, John M. Drake,

and Courtney C. Murdock. 2019. "Microclimate and Larval Habitat Density Predict

Adult Aedes Albopictus Abundance in Urban Areas." *The American Journal of Tropical*

Medicine and Hygiene 101 (2): 362–70. https://doi.org/10.4269/ajtmh.19-0220.

Evans, Michelle V., Justine C. Shiau, Nicole Solano, Melinda A. Brindley, John M. Drake, and

Courtney C. Murdock. 2018. "Carry-over Effects of Urban Larval Environments on the

616   Transmission Potential of Dengue-2 Virus." *Parasites & Vectors* 11 (1): 426.

617   https://doi.org/10.1186/s13071-018-3013-3.

618 Ewing, David A., Bethan V. Purse, Christina A. Cobbold, Stefanie M. Schäfer, and Steven M.

619   White. 2019. "Uncovering Mechanisms behind Mosquito Seasonality by Integrating

620   Mathematical Models and Daily Empirical Population Data: Culex Pipiens in the UK."

621   *Parasites & Vectors* 12 (1): 1–19. https://doi.org/10.1186/s13071-019-3321-2.

622 Ezeakacha, Nnaemeka F., and Donald A. Yee. 2019. "The Role of Temperature in Affecting

623   Carry-over Effects and Larval Competition in the Globally Invasive Mosquito Aedes

624   Albopictus." *Parasites & Vectors* 12 (1): 123. https://doi.org/10.1186/s13071-019-3391-

625   1.

626 Fox, John, and Sanford Weisberg. 2019. *An R Companion to Applied Regression*. 3rd ed.

627   Thousand Oaks CA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

628 Girard, Marc, Christopher B. Nelson, Valentina Picot, and Duane J. Gubler. 2020. "Arboviruses:

629   A Global Public Health Threat." *Vaccine* 38 (24): 3989.

630   https://doi.org/10.1016/j.vaccine.2020.04.011.

631 Gratz, N. G. 2004. "Critical Review of the Vector Status of Aedes Albopictus." *Medical and*

632   *Veterinary Entomology* 18 (3): 215–27. https://doi.org/10.1111/j.0269-

633   283X.2004.00513.x.

634 Hartig, Florian. 2022. "DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

635   Regression Models." R. https://CRAN.R-project.org/package=DHARMa.

636 Hawley, W. A. 1985. "The Effect of Larval Density on Adult Longevity of a Mosquito, Aedes

637   Sierrensis: Epidemiological Consequences." *Journal of Animal Ecology* 54 (3): 955–64.

638   https://doi.org/10.2307/4389.

Hibbard, K.A, F.M. Hoffman, D. Huntzinger, T.O. West, D.J. Wuebbles, D.W. Fahey, K.A.

Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock. 2017. "Changes in Land Cover

and Terrestrial Biogeochemistry." Climate Science Special Report: Fourth National

Climate Assessment, Volume I. Washington, DC: U.S. Global Change Research

Program. https://doi.org/10.7930/J0416V6X.

Huang, Y. J., H. Akbari, H. Taha, and A. H. Rosenfeld. 1987. "The Potential of Vegetation in

Reducing Summer Cooling Loads in Residential Buildings." *Journal of Applied*

Meteorology and Climatology 26 (9): 1103–16. https://doi.org/10.1175/1520-

0450(1987)026<1103:TPOVIR>2.0.CO;2.

Imhoff, Marc L., Ping Zhang, Robert E. Wolfe, and Lahouari Bounoua. 2010. "Remote Sensing

of the Urban Heat Island Effect across Biomes in the Continental USA." *Remote Sensing*

of Environment 114 (3): 504–13. https://doi.org/10.1016/j.rse.2009.10.008.

Jabbar, Hajer Khaled, Mustafa N. Hamoodi, and Amjed N. Al-Hameedawi. 2023. "Urban Heat

Islands: A Review of Contributing Factors, Effects and Data." *IOP Conference Series:*

Earth and Environmental Science 1129 (1): 012038. https://doi.org/10.1088/1755-

1315/1129/1/012038.

Juliano, SA, and TL Stoffregen. 1994. "Effects of Habitat Drying on Size at and Time to

Metamorphosis in the Tree Hole Mosquito *Aedes Triseriatus*." *Oecologia* 97 (3): 369–76.

https://doi.org/10.1007/bf00317327.

Kingsolver, Joel G., and Raymond B. Huey. 2008. "Size, Temperature, and Fitness: Three

Rules." *Evolutionary Ecology Research* 10:251–68.

Lafferty, Kevin D., and Erin A. Mordecai. 2016. "The Rise and Fall of Infectious Disease in a

Warmer World." *F1000Research* 5 (August):F1000 Faculty Rev-2040.

https://doi.org/10.12688/f1000research.8766.1.

Leach, Katie, W. Ian Montgomery, and Neil Reid. 2016. "Modelling the Influence of Biotic

Factors on Species Distribution Patterns." *Ecological Modelling* 337 (October):96–106.

https://doi.org/10.1016/j.ecolmodel.2016.06.008.

Lenth, R. 2023. "Emmeans: Estimated Marginal Means, Aka Least-Squares Means." R package.

R.

Levi, Tal, Eitan Ben-Dov, Preeti Shahi, Dov Borovsky, and Arieh Zaritsky. 2014. "Growth and

Development of *Aedes Aegypti* Larvae at Limiting Food Concentrations." *Acta Tropica*

133 (May):42–44. https://doi.org/10.1016/j.actatropica.2014.02.001.

Li, Yiji, Fatmata Kamara, Guofa Zhou, Santhosh Puthiyakunnon, Chunyuan Li, Yanxia Liu,

Yanhe Zhou, Lijie Yao, Guiyun Yan, and Xiao-Guang Chen. 2014. "Urbanization

Increases Aedes Albopictus Larval Habitats and Accelerates Mosquito Development and

Survivorship." *PLOS Neglected Tropical Diseases* 8 (11): e3301.

https://doi.org/10.1371/journal.pntd.0003301.

Liu-Helmersson, Jing, Hans Stenlund, Annelies Wilder-Smith, and Joacim Rocklöv. 2014.

"Vectorial Capacity of Aedes Aegypti: Effects of Temperature and Implications for

Global Dengue Epidemic Potential." *PLOS ONE* 9 (3): e89783.

https://doi.org/10.1371/journal.pone.0089783.

Lord, Cynthia C. 1998. "Density Dependence in Larval Aedes Albopictus (Diptera: Culicidae)."

Journal of Medical Entomology 35 (5): 5.

Lounibos, L. P., S. Suárez, Z. Menéndez, N. Nishimura, R. L. Escher, S. M. O'Connell, and J. R.

Rey. 2002. "Does Temperature Affect the Outcome of Larval Competition between

Aedes Aegypti and Aedes Albopictus?" *Journal of Vector Ecology: Journal of the*

Society for Vector Ecology 27 (1): 86–95.

Martin, Tara Laine, and Raymond B. Huey. 2008. "Why 'Suboptimal' Is Optimal: Jensen's

Inequality and Ectotherm Thermal Preferences." *The American Naturalist* 171 (3): E102–

18. https://doi.org/10.1086/527502.

McGill, Brian J. 2010. "Matters of Scale." *Science* 328 (5978): 575–76.

https://doi.org/10.1126/science.1188528.

McIntire, Kristina M., and Steven A. Juliano. 2018. "How Can Mortality Increase Population

Size? A Test of Two Mechanistic Hypotheses." *Ecology* 99 (7): 1660–70.

https://doi.org/10.1002/ecy.2375.

Messina, Jane P., Moritz UG Kraemer, Oliver J. Brady, David M. Pigott, Freya M. Shearer,

Daniel J. Weiss, Nick Golding, et al. 2016. "Mapping Global Environmental Suitability

for Zika Virus." eLife. eLife Sciences Publications Limited. April 19, 2016.

https://doi.org/10.7554/eLife.15272.

Miazgowicz, K. L., M. S. Shocket, S. J. Ryan, O. C. Villena, R. J. Hall, J. Owen, T. Adanlawo,

et al. 2020. "Age Influences the Thermal Suitability of Plasmodium Falciparum

Transmission in the Asian Malaria Vector Anopheles Stephensi." *Proceedings of the*

Royal Society B, July. https://doi.org/10.1098/rspb.2020.1093.

Moller-Jacobs, Lillian L., Courtney C. Murdock, and Matthew B. Thomas. 2014. "Capacity of

Mosquitoes to Transmit Malaria Depends on Larval Environment." *Parasites & Vectors*

7 (1): 1–12. https://doi.org/10.1186/s13071-014-0593-4.

Monteiro, Laura C. C., José R. B. de Souza, and Cleide M. R. de Albuquerque. 2007. "Eclosion

Rate, Development and Survivorship of Aedes Albopictus (Skuse)(Diptera: Culicidae)

under Different Water Temperatures." *Neotropical Entomology* 36 (6): 966–71.

https://doi.org/10.1590/s1519-566x2007000600021.

Mordecai, Erin A., Jamie M. Caldwell, Marissa K. Grossman, Catherine A. Lippi, Leah R.

Johnson, Marco Neira, Jason R. Rohr, et al. 2019. "Thermal Biology of Mosquito-Borne

Disease." *Ecology Letters* 22 (10): 1690–1708. https://doi.org/10.1111/ele.13335.

Mordecai, Erin A., Jeremy M. Cohen, Michelle V. Evans, Prithvi Gudapati, Leah R. Johnson,

Catherine A. Lippi, Kerri Miazgowicz, et al. 2017. "Detecting the Impact of Temperature

on Transmission of Zika, Dengue, and Chikungunya Using Mechanistic Models." *PLOS*

Neglected Tropical Diseases 11 (4): e0005568.

https://doi.org/10.1371/journal.pntd.0005568.

Mordecai, Erin A., Krijn P. Paaijmans, Leah R. Johnson, Christian Balzer, Tal Ben-Horin, Emily

de Moor, Amy McNally, et al. 2013. "Optimal Temperature for Malaria Transmission Is

Dramatically Lower than Previously Predicted." Edited by Peter Thrall. *Ecology Letters*

16 (1): 22–30. https://doi.org/10.1111/ele.12015.

Murdock, Courtney C., Michelle V. Evans, Taylor D. McClanahan, Kerri L. Miazgowicz, and

Blanka Tesla. 2017. "Fine-Scale Variation in Microclimate across an Urban Landscape

Shapes Variation in Mosquito Population Dynamics and the Potential of Aedes

Albopictus to Transmit Arboviral Disease." Edited by Samuel V. Scarpino. *PLOS*

Neglected Tropical Diseases 11 (5): e0005640.

https://doi.org/10.1371/journal.pntd.0005640.

Neale, Joseph T., and Steven A. Juliano. 2019. "Finding the Sweet Spot: What Levels of Larval

Mortality Lead to Compensation or Overcompensation in Adult Production?" *Ecosphere*

10 (9): e02855. https://doi.org/10.1002/ecs2.2855.

Nik Abdull Halim, Nik Muhammad Hanif, Nazri Che Dom, Rahmat Dapari, Hasber Salim, and

Nopadol Precha. 2022. "A Systematic Review and Meta-Analysis of the Effects of

Temperature on the Development and Survival of the Aedes Mosquito." *Frontiers in*

Public Health 10. https://www.frontiersin.org/articles/10.3389/fpubh.2022.1074028.

Ogunlade, Samson T., Michael T. Meehan, Adeshina I. Adekunle, Diana P. Rojas, Oyelola A.

Adegboye, and Emma S. McBryde. 2021. "A Review: Aedes-Borne Arboviral Infections,

Controls and Wolbachia-Based Strategies." *Vaccines* 9 (1).

https://doi.org/10.3390/vaccines9010032.

Oke, T. R. 1973. "City Size and the Urban Heat Island." *Atmospheric Environment* 7 (8): 769–

79. https://doi.org/10.1016/0004-6981(73)90140-6.

———. 1982. "The Energetic Basis of the Urban Heat Island." *Quarterly Journal of the Royal*

Meteorological Society 108 (455): 1–24. https://doi.org/10.1002/qj.49710845502.

Packer, M. J., and Philip S. Corbet. 1989. "Size Variation and Reproductive Success of Female

Aedes Punctor (Diptera: Culicidae)." *Ecological Entomology* 14 (3): 297–309.

https://doi.org/10.1111/j.1365-2311.1989.tb00960.x.

Paupy, C., H. Delatte, L. Bagny, V. Corbel, and D. Fontenille. 2009. "Aedes Albopictus, an

Arbovirus Vector: From the Darkness to the Light." *Microbes and Infection*, Forum on

Chikungunya, 11 (14): 1177–85. https://doi.org/10.1016/j.micinf.2009.05.005.

Peng, Shushi, Shilong Piao, Philippe Ciais, Pierre Friedlingstein, Catherine Ottle, François-Marie

Bréon, Huijuan Nan, Liming Zhou, and Ranga B. Myneni. 2011. "Surface Urban Heat

Island Across 419 Global Big Cities." Research-article. ACS Publications. American

Chemical Society. World. December 21, 2011. https://doi.org/10.1021/es2030438.

R Core Team. 2024. "R: A Language and Environment for Statistical Computing." Vienna,

Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Ratkowsky, D A, J Olley, T A McMeekin, and A Ball. 1982. "Relationship between

Temperature and Growth Rate of Bacterial Cultures." *Journal of Bacteriology* 149 (1): 1–

5. https://doi.org/10.1128/jb.149.1.1-5.1982.

Reinhold, Joanna M., Claudio R. Lazzari, and Chloé Lahondère. 2018. "Effects of the

Environmental Temperature on Aedes Aegypti and Aedes Albopictus Mosquitoes: A

Review." *Insects* 9 (4): 158. https://doi.org/10.3390/insects9040158.

Reiskind, M. H., and L. P. Lounibos. 2009. "Effects of Intraspecific Larval Competition on

Adult Longevity in the Mosquitoes Aedes Aegypti and Aedes Albopictus." *Medical and*

Veterinary Entomology 23 (1): 62–68. https://doi.org/10.1111/j.1365-2915.2008.00782.x.

Reiskind, Michael H., and Ali A. Zarrabi. 2012. "Is Bigger Really Bigger? Differential

Responses to Temperature in Measures of Body Size of the Mosquito, Aedes

Albopictus." *Journal of Insect Physiology* 58 (7): 911–17.

https://doi.org/10.1016/j.jinsphys.2012.04.006.

Reiskind, Michael, and M. Shawn Janairo. 2018. "Tracking Aedes Aegypti (Diptera: Culicidae)

Larval Behavior Across Development: Effects of Temperature and Nutrients on

Individuals' Foraging Behavior." *Journal of Medical Entomology* 55 (5): 1086–92.

https://doi.org/10.1093/jme/tjy073.

Rueda, L. M., K. J. Patel, R. C. Axtell, and R. E. Stinner. 1990. "Temperature-Dependent

Development and Survival Rates of Culex Quinquefasciatus and Aedes Aegypti (Diptera:

Culicidae).” *Journal of Medical Entomology* 27 (5): 892–98.

https://doi.org/10.1093/jmedent/27.5.892.

Ruel, Jonathan J., and Matthew P. Ayres. 1999. “Jensen’s Inequality Predicts Effects of

Environmental Variation.” *Trends in Ecology & Evolution* 14 (9).

https://doi.org/10.1016/s0169-5347(99)01664-x.

Russell, Roly, Spencer A. Wood, Gary Allison, and Bruce A. Menge. 2006. “Scale,

Environment, and Trophic Status: The Context Dependency of Community Saturation in

Rocky Intertidal Communities.” *The American Naturalist* 167 (6): E158–70.

https://doi.org/10.1086/504603.

Ryan, Sadie J., Colin J. Carlson, Blanka Tesla, Matthew H. Bonds, Calistus N. Ngonghala, Erin

A. Mordecai, Leah R. Johnson, and Courtney C. Murdock. 2021. “Warming

Temperatures Could Expose More than 1.3 Billion New People to Zika Virus Risk by

2050.” *Global Change Biology* 27 (1): 84–93. https://doi.org/10.1111/gcb.15384.

Ryan, Sadie J., Catherine A. Lippi, Oswaldo C. Villena, Aspen Singh, Courtney C. Murdock,

and Leah R. Johnson. 2023. “Mapping Current and Future Thermal Limits to Suitability

for Malaria Transmission by the Invasive Mosquito Anopheles Stephensi.” *Malaria*

Journal 22 (1): 104. https://doi.org/10.1186/s12936-023-04531-4.

Sandel, Brody, and Adam B. Smith. 2009. “Scale as a Lurking Factor: Incorporating Scale-

Dependence in Experimental Ecology.” *Oikos* 118 (9): 1284–91.

https://doi.org/10.1111/j.1600-0706.2009.17421.x.

Sauers, Logan A., Kelsey E. Hawes, and Steven A. Juliano. 2022. “Non-Linear Relationships

between Density and Demographic Traits in Three Aedes Species.” *Scientific Reports* 12

(May):8075. https://doi.org/10.1038/s41598-022-11909-y.

Schneider, P., W. Takken, and P. J. Mccall. 2000. "Interspecific Competition between Sibling

Species Larvae of Anopheles Arabiensis and An. Gambiae." *Medical and Veterinary*

Entomology 14 (2): 165–70. https://doi.org/10.1046/j.1365-2915.2000.00204.x.

Shapiro, Lillian L. M., Courtney C. Murdock, Gregory R. Jacobs, Rachel J. Thomas, and

Matthew B. Thomas. 2016. "Larval Food Quantity Affects the Capacity of Adult

Mosquitoes to Transmit Human Malaria." *Proceedings of the Royal Society B: Biological*

Sciences, July. https://doi.org/10.1098/rspb.2016.0298.

Shocket, Marta S., Christopher B. Anderson, Jamie M. Caldwell, Marissa L. Childs, Lisa I.

Couper, Songhee Han, Mallory J. Harris, et al. 2020. "Environmental Drivers of Vector-

Borne Diseases," December. https://doi.org/10.1093/oso/9780198853244.003.0006.

Telang, Aparna, and Michael A Wells. 2004. "The Effect of Larval and Adult Nutrition on

Successful Autogenous Egg Production by a Mosquito." *Journal of Insect Physiology* 50

(7): 677–85. https://doi.org/10.1016/j.jinsphys.2004.05.001.

Teng, Hwa-Jen, and Charles S. Apperson. 2000. "Development and Survival of Immature Aedes

Albopictus and Aedes Triseriatus (Diptera: Culicidae) in the Laboratory: Effects of

Density, Food, and Competition on Response to Temperature." *Journal of Medical*

Entomology 37 (1): 40–52. https://doi.org/10.1603/0022-2585-37.1.40.

Tesla, Blanka, Leah R. Demakovsky, Erin A. Mordecai, Sadie J. Ryan, Matthew H. Bonds,

Calistus N. Ngonghala, Melinda A. Brindley, and Courtney C. Murdock. 2018.

"Temperature Drives Zika Virus Transmission: Evidence from Empirical and

Mathematical Models." *Proceedings of the Royal Society B*, August.

https://doi.org/10.1098/rspb.2018.0795.

Waldock, Joanna, Nastassya L. Chandra, Jos Lelieveld, Yiannis Proestos, Edwin Michael,

George Christophides, and Paul E. Parham. 2013. "The Role of Environmental Variables

on Aedes Albopictus Biology and Chikungunya Epidemiology." *Pathogens and Global*

Health 107 (5): 224–41. https://doi.org/10.1179/2047773213Y.0000000100.

Walker, Edward D., Daniel L. Lawson, Richard W. Merritt, William T. Morgan, and Michael J.

Klug. 1991. "Nutrient Dynamics, Bacterial Populations, and Mosquito Productivity in

Tree Hole Ecosystems and Microcosms." *Ecology* 72 (5): 1529–46.

https://doi.org/10.2307/1940953.

Wiens, J. A. 1989. "Spatial Scaling in Ecology." *Functional Ecology* 3 (4): 385–97.

https://doi.org/10.2307/2389612.

Wilder-Smith, Annelies, Duane J. Gubler, Scott C. Weaver, Thomas P. Monath, David L.

Heymann, and Thomas W. Scott. 2017. "Epidemic Arboviral Diseases: Priorities for

Research and Public Health." *The Lancet Infectious Diseases* 17 (3): e101–6.

https://doi.org/10.1016/S1473-3099(16)30518-7.

Wimberly, Michael C., Justin K. Davis, Michelle V. Evans, Andrea Hess, Philip M. Newberry,

Nicole Solano-Asamoah, and Courtney C. Murdock. 2020. "Land Cover Affects

Microclimate and Temperature Suitability for Arbovirus Transmission in an Urban

Landscape." *PLOS Neglected Tropical Diseases* 14 (9): e0008614.

https://doi.org/10.1371/journal.pntd.0008614.

Yoshioka, Miho, Jannelle Couret, Frances Kim, Joseph McMillan, Thomas R. Burkot, Ellen M.

Dotson, Uriel Kitron, and Gonzalo M. Vazquez-Prokopec. 2012. "Diet and Density

Dependent Competition Affect Larval Performance and Oviposition Site Selection in the

Mosquito Species Aedes Albopictus (Diptera: Culicidae).” *Parasites & Vectors* 5 (1):

225. https://doi.org/10.1186/1756-3305-5-225.

Zhang, Qian, Kaiyuan Sun, Matteo Chinazzi, Ana Pastore y Piontti, Natalie E. Dean, Diana

Patricia Rojas, Stefano Merler, et al. 2017. “Spread of Zika Virus in the Americas.”

Proceedings of the National Academy of Sciences 114 (22): E4334–43.

https://doi.org/10.1073/pnas.1620161114.

Zhou, Bin, Diego Rybski, and Jürgen P. Kropp. 2017. “The Role of City Size and Urban Form in

the Surface Urban Heat Island.” *Scientific Reports* 7 (1): 4791.

https://doi.org/10.1038/s41598-017-04242-2.

Table 1. Daily mean microclimate values (95% confidence intervals) across seasons (summer and fall) and land use type (rural, suburban, urban). Temperature (°C) data were collected from data loggers placed in the larval environment, while relative humidity (%) data were collected from loggers placed <1m above ground.

	Summer			Fall		
	Rural	Suburban	Urban	Rural	Suburban	Urban
Temperature (°C)						
T_{min}	22.5 (22.0, 22.9)[a]	22.7 (22.2, 23.2)[ab]	23.3 (22.8, 23.7)[b]	17.5 (17.1, 17.9)[c]	18.2 (17.8, 18.7)[d]	18.4 (18.0, 18.8)[d]
T_{mean}	25.0 (24.5, 25.6)[a]	25.1 (24.5, 25.6)[a]	25.7 (25.2, 26.3)[a]	21.1 (20.6, 21.6)[b]	21.3 (20.8, 21.8)[b]	21.6 (21.2, 22.1)[b]
T_{max}	28.7 (27.4, 29.9)[a]	28.3 (27.0, 29.5)[a]	29.2 (28.0, 30.5)[a]	25.7 (24.6, 26.8)[b]	26.4 (25.2, 27.6)[b]	25.3 (24.2, 26.4)[b]
T_{range}	6.37 (5.26, 7.47)[a]	5.65 (4.51, 6.80)[a]	6.10 (5.03, 7.18)[a]	8.00 (6.93, 9.08)[a]	8.17 (7.02, 9.32)[a]	6.69 (5.62, 7.77)[b]
Relative Humidity (%)						
R_{min}	80.8 (74.9, 87.1)[a]	75.7 (70.4, 81.5)[ab]	68.0 (63.4, 73.0)[b]	70.1 (65.1, 75.4)[c]	68.9 (63.8, 74.5)[cd]	61.4 (57.0, 66.2)[d]
R_{mean}	96.0 (92.9, 99.3)[a]	93.9 (90.8, 97.0)[a]	88.2 (85.5, 91.1)[b]	92.1 (89.1, 95.2)[c]	90.3 (87.3, 93.5)[c]	84.5 (81.7, 87.3)[d]
R_{max}	100.0 (99.0, 101.0)[a]	100.0 (99.0, 100.9)[a]	98.8 (97.9, 99.7)[a]	99.6 (98.7, 100.6)[a]	99.3 (98.3, 100.3)[a]	97.4 (96.5, 98.3)[b]
R_{range}	19.8 (15.5, 24.2)[a]	24.6 (20.4, 28.9)[ab]	31.0 (27.0, 35.0)[b]	28.8 (24.6, 33.1)[c]	29.4 (24.9, 33.8)[c]	34.6 (30.4, 38.9)[c]

* Letters represent significant differences as measured by Tukey's multiple comparison of means, adjusting for significance with the Holm-Bonferroni method.

859 **Table 2.** Mean demographic trait values [female emergence (%), female development time

860 (days), female wing length (mm), intrinsic growth rate] (95% confidence intervals) across

861 seasons (summer and fall), land use type (rural, suburban, urban), and larval density treatment

862 (30 = low, 60 = medium, 120 = high).

	Female Emergence (%)	Development Time (days)	Wing Length (mm)	Per Capita Growth Rate (days^{-1})
Season				
Summer	22.6 (19.9, 25.4)[a]	16.1 (15.5,16.7)[a]	2.40 (2.35, 2.45)[a]	0.114 (0.109, 0.119)[a]
Fall	6.6 (5.33, 8.05)[b]	18.5 (17.6, 19.4)[b]	2.09 (2.03, 2.15)[b]	0.045 (0.040, 0.051)[b]
Land Use Type				
Rural	12.3 (9.7, 15.6)[a]	17.4 (16.3, 18.6)[a]	2.24 (2.16, 2.33)[a]	0.078 (0.070, 0.086)[a]
Suburban	13.2 (10.5, 16.6)[a]	17.2 (16.2, 18.4)[a]	2.23 (2.15, 2.31)[a]	0.082 (0.074, 0.090)[a]
Urban	12.0 (9.3, 15.3)[a]	17.0 (15.9, 18.2)[a]	2.24 (2.16, 2.32)[a]	0.078 (0.071, 0.087)[a]
Density Treatment				
Low	28.9 (25.2, 32.9)[a]	17.8 (17.1, 18.5)[a]	2.35 (2.29, 2.40)[a]	0.113 (0.108, 0.119)[a]
Medium	12.4 (10.5, 14.6)[b]	17.0 (16.3, 17.7)[b]	2.28 (2.22, 2.33)[b]	0.085 (0.079, 0.090)[b]
High	4.8 (3.7, 6.3)[c]	16.9 (16.0, 18.0)[b]	2.10 (2.03, 2.17)[c]	0.041 (0.035, 0.047)[c]

865 ***** Letters represent significant differences as measured by Tukey's multiple comparison of

866 means, adjusting for significance with the Holm-Bonferroni method.

868   **Table 3.** Generalized Linear Mixed Model results of the interaction between season (summer vs.

869   fall) and larval density treatment (30 = low, 60 = medium, 120 = high) on demographic rates.

870   Cells represent mean values (95% confidence intervals).

	Density Treatment	Female Emergence (%)	Development Time (days)	Wing Length (mm)	Per Capita Growth Rate (days^{-1})
Summer	Low	30.9 (26.1, 36.1)[a]	15.7 (15.0, 16.4)[a]	2.53 (2.46, 2.60)[a]	0.133 (0.126, 0.140)[a]
	Medium	20.6 (17.2, 24.4)[b]	15.9 (15.2, 16.5)[a]	2.49 (2.42, 2.56)[a]	0.115 (0.108, 0.123)[b]
	High	17.6 (14.9, 20.8)[c]	16.7 (16.0, 17.3)[b]	2.20 (2.14, 2.26)[b]	0.093 (0.086, 0.101)[c]
Fall	Low	27.0 (22.5, 32.0)[d]	20.1 (19.3, 21.1)[c]	2.18 (2.12, 2.25)[d]	0.094 (0.087, 0.101)[d]
	Medium	7.2 (5.6, 9.1)[e]	18.2 (17.3, 19.2)[d]	2.09 (2.02, 2.15)[e]	0.054 (0.047, 0.061)[e]
	High	1.2 (0.7, 1.9)[f]	17.2 (15.6, 19.0)[b]	2.00 (1.90, 2.11)[e]	–0.011 (-0.021, -0.002)[f]

873   * Letters represent significant differences as measured by pairwise comparison using Tukey's

874   multiple comparison of means, adjusting for significance with the Holm-Bonferroni method.