JOURNAL NAME: Ecology

MANUSCRIPT TYPE: Article

MANUSCRIPT TITLE: Mosquito population dynamics are shaped by interactions among

larval density, temperature, and humidity

AUTHORS: Nicole Solano[1,2], Emily C. Herring[3], Carl W. Hintz[4], Philip M. Newberry[1,2],

Annakate M. Schatz[1,2], Joseph W. Walker[2*], Gregory R. Jacobs[5], Craig W. Osenberg[1], Courtney

C. Murdock[1,2†]

AFFILIATIONS:

1. Odum School of Ecology, University of Georgia, Athens, GA, USA

2. Center for Ecology of Infectious Diseases, University of Georgia, Athens, GA, USA

3. Department of Population Health, College of Veterinary Medicine, University of

Georgia, Athens, GA, USA

4. Department of Applied Ecology, North Carolina State University, Raleigh, NC, USA

5. Natural Resources and the Environment Section, Ashley School of Global Development

and the Environment, Cornell University, Ithaca, NY, USA

CORRESPONDING AUTHOR: Nicole Solano (nicole.n.solano@gmail.com)

* Current address: Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT, USA
† Current address: Department of Entomology, College of Agriculture and Life Sciences, Cornell University, Ithaca, NY, USA

KEY WORDS

Aedes albopictus; Density-Dependence*;* Fitness*;* Per Capita Growth Rate*;* Seasonality*;* Life

History Traits; Land Use

DATA AVAILABILITY STATEMENT

The data and code that support the findings of this study are openly available on Figshare:

https://figshare.com/s/93b37d27373bb0f48896.

ABSTRACT (English)

Understanding how variation in key abiotic and biotic factors interact at ecologically relevant

spatial scales is crucial for predicting population dynamics, distributions, and abundances. This

is especially true for vectors that transmit human pathogens. However, studies investigating the

effects of environmental variation on vectors have typically investigated environmental factors in

isolation or in laboratory experiments that examine constant environmental conditions that often

do not occur in the field. To address these limitations, we conducted a semi-field experiment in

Athens, Georgia using the invasive Asian tiger mosquito (*Aedes albopictus*). We selected nine

sites that varied in impervious surface and vegetation cover to explore effects of natural variation

in microclimate (specifically, temperature and humidity) on mosquitoes. We manipulated

conspecific larval density at each site and repeated the experiment in the summer and fall to

further increase the variability in temperature and humidity. We then evaluated the original

design features (land cover, larval density, and season), and their interactions, on the mean

proportion of females emerging, juvenile development time, size upon emergence, and estimated

per capita population growth (i.e., fitness). We found significant effects of larval density, land

cover, and season on all response variables, including a non-intuitive decrease in development

time with increasing larval density in the fall. We repeated these analyses using the

hypothesized microclimate drivers of these effects: temperature and humidity. In general, the

model using the microclimate variables outperformed the model using land use and season,

highlighting the roles of temperature and humidity and their interactions with density on

mosquito traits and dynamics. Our study demonstrates that ignoring the interaction between

variation in biotic (e.g., intraspecific competition) and abiotic (e.g., temperature and humidity)

variables could reduce the accuracy and precision of models used to predict mosquito population

and pathogen transmission dynamics, especially those inferring dynamics at finer-spatial scales

across which transmission and control occur.

ABSTRACT (Spanish)

Es crucial comprender cómo los factores abióticos y bióticos interactúan a escalas

ecológicamente relevantes para predecir la dinámica poblacional, la distribución, y la

abundancia. Esto es especialmente cierto para vectores que transmiten patógenos humanos. Sin

embargo, la mayoría de los estudios han considerado factores ambientales de forma aislada o en

condiciones de laboratorio poco realistas. Para superar estas limitaciones, realizamos un

experimento de semi-campo en Athens, Georgia, con el mosquito invasor *Aedes albopictus*.

Manipulamos la densidad larval en nueve sitios que variaban en cobertura vegetal y superficie

impermeable, y repetimos el experimento en el verano y otoño. Luego, evaluamos los efectos de

la cobertura del suelo, la densidad larval, y la estación en la proporción de hembras adultas

emergidas, el tiempo de desarrollo, el tamaño al emerger, y el crecimiento poblacional per cápita.

Encontramos efectos significativos de todas las variables, así como fuertes interacciones entre

estación y densidad intraespecífica de mosquitos, incluyendo una disminución no intuitiva del

tiempo de desarrollo en el otoño con mayor densidad larval. Repetimos los análisis usando

factores microclimáticas hipotetizados como impulsores de estos efectos: temperatura y

humedad. En general, el modelo que incorpora variables microclimáticas superó al modelo

basado en el uso de suelo y estacionalidad, evidenciando la importancia de la temperatura y la

humedad, así como sus interacciones con la densidad larval, en las características biológicas y la

dinámica poblacional de los mosquitos. Nuestros resultados demuestran que la interacción entre

variación biótica (competencia intraespecífica) y abiótica (temperatura y humedad), es

fundamental para mejorar la precisión de los modelos que predicen la dinámica poblacional de

mosquitos y la transmisión de patógenos a escalas locales de transmisión y control.

INTRODUCTION

Three processes generally determine the distribution and abundances of organisms: environmental filtering (abiotic variables), species interactions (biotic variables), and constraints on organismal dispersal (Leibold and Chase, 2018; Menge and Sutherland, 1976; Vellend, 2016). The effects of abiotic and biotic variables on species distribution and abundances have become some of the most important issues in ecology today, especially in the context of rapid biodiversity loss (McKinney and Lockwood, 1999) climate change (Parmesan, 2006) and human-mediated dispersal (Bullock and Pufal, 2020). Multiple environmental factors (both abiotic and biotic) affect the demography, vital rates (e.g., development, growth, and survival), and population dynamics of ectothermic organisms. However, predicting the effects of abiotic and biotic variables on organismal distribution and abundance patterns is difficult because their effects on processes that determine organismal fitness and population dynamics are scale-dependent (Levin, 1992; McGill, 2010; Wiens, 1989).

It has been long hypothesized that environmental filtering is strongest at larger, more regional scales because climate variables (e.g., temperature, relative humidity, precipitation) exhibit little variation at fine-spatial scales (Houghton et al., 2002; Levin, 1992; McGill, 2010; Wiens, 1989). With the emphasis on climate warming in the ecological literature, there has been a strong focus on understanding the effects of temperature on the distribution and abundance of ectothermic organisms, as well as their traits (Bale et al., 2002; Helmuth et al., 2010; Mordecai et al., 2019; Parmesan, 2006; Pörtner and Farrell, 2008). These effects have been leveraged in species distribution models and mechanistic models to predict the current and future geographic ranges of ectothermic organisms (Amarasekare, 2024; Huey et al., 2001; Simon and Amarasekare, 2024). However, biotic variables, like intra- and inter-species competition,

availability of resources, and presence and density of biological enemies (predators and

parasites), also have important effects on organismal fitness and population dynamics (Beale et

al., 2014; Ewing et al., 2019; Golding et al., 2015; Leach et al., 2016). These biotic factors affect

individuals and can vary among local neighborhoods; thus, the effects of these factors on species

distribution and abundance patterns are hypothesized to arise at fine-spatial scales (Levin, 1992;

McGill, 2010) but are often not incorporated into species distribution or mechanistic models

(Cohen et al., 2016). Finally, abiotic and biotic factors can interact to affect species

distributions, which could be especially relevant for organisms with complex life cycles, such as

amphibians or insects, that use very different habitats across their life stages (Govindarajulu and

Anholt, 2006; Indermaur et al., 2010). When this spatial variation is ignored, predictions of a

given species' population dynamics averaged over space or time can lead to considerable bias if

the effects of these factors are nonlinear (Bozinovic et al., 2011; Martin and Huey, 2008; Ruel

and Ayres, 1999). As a result, mathematical models might overlook the influence of hot spots or

ecological traps on a given species' population dynamics.

One of the areas in which these issues are readily apparent is in the field of mosquito

ecology. Climate and land use change are rapidly altering the landscapes across which

mosquitoes occur, altering the ecological relationships mosquitoes share with their environments

and their vertebrate hosts. This has led to biological invasions of mosquito species and their

parasites to new areas of the world (Roiz et al., 2024; Taylor et al., 2024) with consequences for

the emergence of new diseases (Chala and Hamde, 2021; Lounibos, 2002; Yan et al., 2024).

Understanding and predicting how future changes in climate and land use will alter mosquito

dynamics and the spread of vector-borne pathogens is particularly relevant. Thus, there has been

considerable investigation into the effects of temperature on mosquito distributions, population

dynamics, and parasite transmission (reviewed in Kraemer et al., 2019; Liu-Helmersson et al.,

2019; Mordecai et al., 2019; Ryan et al., 2023; Sinka et al., 2020). Recent work also suggests

that relative humidity can have important effects on larval traits and mosquito population

dynamics, possibly through effects of humidity on the surface tension of habitats (see Brown et

al., 2023; Huxley et al. 2025). In addition to abiotic variables, mosquitoes experience intra- and

inter-specific competition during their larval stage, which can affect mosquito survival,

reproduction, and generation time by slowing development, reducing adult size, and lowering

teneral reserves (e.g., lipids and proteins) of adult mosquitoes (Briegel, 2003) with significant

implications for mosquito fitness and vectorial capacity (Alto and Bettinardi, 2013; Alto et al.,

2005; Moller-Jacobs et al., 2014; Shapiro et al., 2016).

Currently, there has been little investigation of the potential interactive effects of abiotic

and biotic factors on mosquito population dynamics. Most studies that have considered the

interaction between abiotic (i.e., temperature, relative humidity) and biotic factors (i.e.,

intraspecific competition) on mosquito life history traits and population dynamics have been

conducted in the lab under constant environmental conditions that may not reflect field

environments (Evans et al., 2021; Miazgowicz et al., 2020; Tesla et al., 2018; Shapiro et al.,

2017). In the field, abiotic variables like temperature vary diurnally, seasonally, and across

space (e.g., due to variation in land use and land cover) (Evans et al., 2018, 2019; Murdock et al.,

2017; Wimberly et al., 2020).

To improve our understanding of how field-relevant variation in key abiotic and biotic

factors interact at spatial scales relevant for mosquito fitness and population dynamics, we

conducted a semi-field experiment using the Asian tiger mosquito (*Aedes albopictus*) in Athens,

Georgia during summer and fall 2017. *Ae. albopictus* is a highly invasive mosquito in the United

States and other areas of the world, as well as a competent vector of arboviruses including

dengue and chikungunya (Benedict et al., 2007; Bonizzoni et al., 2013). We selected nine sites

that varied in impervious surface and vegetation cover, and conducted an experiment across the

sites at two different times to leverage natural spatial and temporal variation in temperature and

relative humidity, and explored their effects on mosquito traits and population dynamics (Brown

et al., 2023; Murdock et al., 2017; Wimberly et al., 2020). During both experiments, we also

manipulated conspecific larval density, a biotic variable known to have large effects on metrics

of mosquito fitness and population growth rates (Juliano, 2009; Teng and Apperson, 2000). We

simultaneously monitored spatial and temporal variation in temperature and humidity. Although

we designed our study based upon density, land use and season, we were specifically interested

in the interactive effects of conspecific density, temperature and relative humidity, each of which

is known to affect mosquito demography in isolation (e.g., Alto and Juliano, 2001; Brown et al.,

2023; Mordecai et al., 2019), while controlling for other factors that might vary with season and

land use (e.g., leaf litter or detrital inputs). We hypothesized that temperature and humidity

would serve as "master variables" and would provide a more general and better predictor of

mosquito dynamics than land use and season. Understanding the drivers of spatial and temporal

heterogeneity on the dynamics of invasive species like *Ae. albopictus* is crucial for predicting

and managing their future invasive potential, as well as for other species with similar ecology

(e.g., *Aedes aegypti)* that play an important role in human arbovirus transmission.

METHODS

Experimental design

164
165

We conducted the study between June 2017 and October 2017 in Athens-Clarke County,

Georgia, USA. We utilized nine previously established field sites that ranged in impervious

surface, vegetation cover, and local microclimate (Appendix S1: Figure S1) (Evans et al., 2019).

Within each of the nine sites, we placed three trays in full shade separated by approximately 6-9

m. Each tray held four wide-mouth mason jars each containing 300 mL of leaf infusion (Evans

et al., 2018; Murdock et al., 2017). Three jars in each tray received a specific larval density

treatment (30, 60, or 120 *Ae. albopictus* first instar larvae from a laboratory colony established

from field collected *Ae. albopictus* in 2015, Athens, GA;; Murdock et al., 2017). We covered the

top of the jars with mesh netting and staked a wire cage with a vinyl roof over the trays (see

Murdock et al., 2017) to prevent changes in habitat volume from rainfall, the addition of

nutrients due to falling vegetation, and the loss of jars from animal disturbance. To understand

how seasonal variation in abiotic factors may alter relationships between conspecific larval

density and traits governing mosquito population dynamics, we conducted this experiment in the

summer (June 11 - July 25) and fall (September 8 - October 18) 2017. Jars were visited daily

and at the same time of day. Any adults that had emerged were collected and placed on ice for

transport back to the lab. Mosquitoes were then stored at -20°C until we determined their sex

(Darsie and Ward, 2005) and wing length.

Microclimate variables

184
To measure variation in water temperature, we deployed one data logger (Monarch

Instruments, Amherst, NH, USA: Radio Frequency Identification (RFID) Temperature Track-It

Logger) to one jar in each tray that had been filled with leaf infusion but contained no larvae.

We also uniformly distributed six data loggers of the same brand throughout each site at

approximately 0.5-1.0 m above the ground and in full shade to measure relative humidity. All

loggers recorded microclimate variables every ten minutes with data downloaded monthly.

Mosquito life history traits

In this study, we only report female data since females determine mosquito population

growth rates and pathogen transmission. To calculate the proportion of mosquito larvae and

pupae that survived to emerge as adults (i.e., emergence),),) we divided the total number of

females that emerged by half the total number of 1^{st} instar larvae placed into the habitat (e.g., 15,

30, or 60 assuming half of the 1^{st} instar larvae were female). Juvenile development time was

defined as the number of days between the start of the experiment and the day an adult female

emerged. One wing was removed from each individual, mounted onto a glass slide using clear

nail polish, and measured with a dissecting microscope and micrometer to index body size

(Packer and Corbet, 1989). To estimate the per capita growth rates of mosquito populations in

response to our experimental treatments we used equation 7 from Livdahl & Sugihara (1984):

203
$$r' = \frac{ln\left(\frac{1}{N_0}\sum_x A_x\ f(w_x)\right)}{D + \frac{\sum_x x A_x\ f(w_x)}{\sum_x A_x\ f(w_x)}} \tag{1}$$

where N_0 is the initial number of females in a cohort (i.e., half of the density treatment), A_x is the

number of females that emerged on day x, and D is the time to reproduction following

emergence (which we assume to be 14 days: Lounibos et al., 2002). The fecundity $f(w_x)$ of

females that emerged on day x, with an average wing length of w_x, was estimated from the linear

relationship between wing length and number of eggs laid from Lounibos et al. (2002):

$$f(w_x) = -121.240 + (78.02 * w_x) \qquad\qquad (2)$$

In 12 cases, no females emerged from a jar (these all occurred in the fall at high density). Since

removing those jars would elevate and bias r' for that treatment, we imputed their per capita

growth rate by setting $r' = -0.05$ day^{-1}, which was just below the most negative r' observed in

the other jars.

Data Analyses

We took three complementary approaches to analyze the results. First, we examined

variation in water temperature and relative humidity across land use and season. To assess how

daily microclimate conditions varied, we used linear mixed effects models to investigate how the

daily mean, minimum, maximum, and range in air temperature (T_{mean}, T_{min}, T_{max}, T_{range}) and

relative humidity (R_{mean}, R_{min}, R_{max}, R_{range}) varied with design variables.. All models included

fixed effects of season, land-use type, and their interaction, and random effects of site, logger

(nested within site), and day.

Second, we investigated how juvenile survival and development rate, adult size, and

estimated per capita population growth (i.e., fitness) of *Ae. albopictus* varied with land use,

season, and larval density. Although our putative interest was microclimate (and not land use and

season per se), our sampling design was based on these categorical factors, so we started with

those analyses before moving on to exploration of the effects of temperature and humidity. We

also used GLMMs to determine how emergence (binomial distribution; *logit* link function),

larval development rate (lognormal distribution; *log* link function), wing length (lognormal

distribution; *log* link function), and per capita growth rate (Gaussian distribution; *identity* link

function) varied across seasons, land-use types, and density treatments. Those models all had

season, land-use type, density treatment, and their interactions as fixed effects, with site and tray

(within site) as nested random effects. Analyses of development rate and wing length also

included jars (within tray) as an additional nested random effect.

Third, we constructed models that replaced land use and season (the design variables)

with temperature and humidity (the putative proximate drivers of variation mosquito dynamics).

To begin our analyses of the effects of microclimate on mosquito traits, we first explored the

correlations among the microclimate variables to avoid multicollinearity. Based on correlation

analyses, we included only T_{min}, T_{max}, and R_{min} in our models since they were of particular

biological importance to us and they were not strongly correlated to each other (Appendix S1:

Figure S2). We analyzed effects of the microclimate variables using GLMMs and the same

distributions and link functions for each response variable as described above, although in this

case the predictors were continuous rather than categorical.

Finally, to compare the use of the microclimate variables vs. season and land use for each

response variable, we ran three models – *null*, *base*, and *microclimate* – and used the corrected

Akaike Information Criterion (AIC_c) to evaluate their relative performance. The *null model* only

included density treatment as a fixed effect (e.g., it lacked land use and season). The *base model*

incorporated the original design factors (season, land-use type, density, and their interactions) as

fixed effects.. The *microclimate model* replaced the effects of land use and season with the

microclimate variables, and therefore included the fixed effects of density, along with the main

effects of T_{min}, T_{max}, R_{min} and their interactions with density. For each response variable, all

three models included the effect of density and an identical random effect structure, but they

varied in how they incorporated environmental variables. Microclimate variables experienced by

larvae in a jar were characterized by the mean environment experienced by those larvae as

determined by their distribution of emergence dates (i.e., for each larvae we calculated the mean

experienced microclimate per larvae from day 0 to its day of emergence and then averaged these

means across all females that emerged from the jar). In 12 cases in the fall at high density, no

females emerged from a jar, so the microclimate variables for those jars were imputed based on

emergence patterns from other jars at that site, season and density. Two sites did not have any

females emerge (one rural site and one suburban site); therefore, we used the emergence patterns

from other jars at the two other sites of the same land-use type, season, and density.

All analyses were conducted in R 4.5.1 (R Core Team, 2025). Code to run analyses and

create figures can be found on Figshare (Solano, 2026). Models were fit using the *glmmTMB*

package (Brooks et al., 2017), and compared by AIC_c using the *bblme* package (Bolker and R

Development Core Team, 2023). Type III analysis of variance (ANOVA) was conducted for all

models using the *car* package (Fox et al., 2024). The *emmeans* package was used to obtain

means and confidence intervals after adjusting for random effects (Lenth et al., 2025). We

assessed model fit using the scaled residuals from the *DHARMa* package (Hartig et al., 2024).

<u>RESULTS</u>

Microclimate

Five of the 54 aquatic temperature loggers failed (three in the summer and two in the

fall), and 36 of the 108 terrestrial loggers failed or were stolen (16 in the summer and 20 in the

fall). Despite these losses, failure rates were roughly equivalent across sites and microclimate

variation across season and land use types was well represented by the remaining loggers. Water

temperature (i.e., mean daily T_{min}, T_{mean}, and T_{max}) and relative humidity (mean daily R_{min}, R_{mean},

and R_{max}) metrics all varied significantly with season, land use, and their interaction (Appendix

S1: Figure S3, Tables S1 and S2), demonstrating that land use and season captured significant

spatiotemporal variation in mosquito-relevant microclimates.

Emergence

282   Of the 5670 first instar *Ae. albopictus* placed into the field during each season, a total of

3156 mosquitoes (56% of the starting population) successfully emerged as adults during the

summer while only 1233 mosquitoes (38%) emerged in the fall (emergence is equivalent to

survival through the larval and pupal stages). Assuming that the initial frequency of females was

50%, there were significant main effects of season and conspecific density on the proportion of

female first instar larvae that survived to emergence, with greater survival in the low-density

treatment relative to medium- and high- density treatments (Figure 1a, Appendix S1: Table S3).

There also were interactions between season and density (Figure 1a) as well as land-use type and

density (Appendix S1: Figure S4, Table S3). For example, there was little difference in the

survival of female juveniles in the low-density treatments in the summer and fall; however, the

negative effects of increasing conspecific density on juvenile survival was much greater in the

fall (Figure 1a). Further, female juveniles at low conspecific densities survived best in suburban

sites, while those at intermediate densities survived best in urban sites and those at high densities

survived best in rural sites (Appendix S1: Figure S4).

296   Some of these effects likely arose through the influence of the microclimate variation

across land use and season. For example, the microclimate model substantially outperformed the

null model ($\Delta AIC_c = 393$; Table 1), with the probability of female emergence (juvenile survival)

decreasing with increasing density and increasing with increasing temperature (T_{min}) and relative

humidity (R_{min}). In addition to these main effects, we observed interactions of both microclimate

variables with density (Figure 2a): as density increased, the positive effect of temperature (i.e.,

T_{min}) and relative humidity (R_{min}) increased in magnitude (Figure 2a). However, the base model

outperformed the microclimate model ($\Delta AIC_c = 6.3$; Table 1), suggesting that the temperature

and humidity variables did not capture as much of the spatiotemporal drivers of juvenile survival

as did the original design variables (land use and season).

Female Development Time

Both season and conspecific larval density significantly affected the average amount of

time required for adult female emergence (Table 1, Appendix S1: Table S3), with females on

average developing more quickly in the summer (day 9 mean emergence date) relative to the fall

(day 13 mean emergence date) and in the low-density treatment (Figure 1b, Appendix S1: Figure

S4b). However, a significant interaction between season and larval density demonstrates that the

main effect of density is being driven by the relationship between mean female development

time (time to emergence) and larval density in the summer (Appendix S1: Figure S4, Table S3).

Females reared in the low-density treatment were the fastest to develop in the summer but were

the slowest to develop in the fall (Figure 1b).

These effects on development time were better predicted using the microclimate variables

($\Delta AIC_c > 30$ for the microclimate model compared to both the null and base models).

Development time decreased with increasing daily T_{min}, and this effect was strongest for

mosquitoes developing in low density environments (Figure 2b). In contrast to the other life

history parameters, there was little evidence to suggest a role of relative humidity on

development time (Table 1).

Wing Length

We observed significant main effects of season and density on female wing length

(Appendix S1: Figure S4, Table S3). Females reared in the summer had longer wings compared

to the fall, and females developing in the high conspecific larval treatment had significantly

shorter wings than mosquitoes from the low-density treatment (Figure 1c, Appendix S1: Figure

S4). The observed variation in wing length was better predicted by the microclimate model

compared to the null or based models (Table 1: $\Delta AIC_c > 18$). Increased temperature (reflected in

both T_{min} and T_{max}) increased (Figure 2c, Table 1). Additionally, we observed a significant

interaction between density and daily R_{min}, with increases in relative humidity resulting in

increased wing size at low and medium densities but not at high density (Figure 2c, Table 1).

Estimated Per Capita Population Growth Rate

Integrating these responses revealed strong effects of season, conspecific larval density,

and the interaction between these two factors on the estimated per capita population growth rate,

r' (Equation 1, Figure 1d, Appendix S1: Table S3). In general, the estimated per capita growth

rate was greater in the summer relative to the fall and was greater at low densities than at high

densities (Figure 1d). A significant interaction between season and density was observed due to

greater declines in the estimated per capita growth rate with increasing conspecific density in the

fall relative to the summer (Figure 1d, Appendix S1: Tables S3). Additionally, we observed

negative per capita growth rates at the highest conspecific density in the fall, while all other

combinations of factors led to positive projected per capita growth (Figure 1d, Appendix S1:

Figure S4). These effects may have arisen through the effects of T_{min} and R_{min}, both of which

were positively associated with increased per capita growth (Figure 2d, Table 1), although both

microclimate variables also exhibited significant interactions with density. For example, the

effect of increasing T_{min} was greater on larvae living at high densities and more muted for those

living at lower densities (Figure 2d).

Effects of microclimate on mosquito performance

In presenting our results, we initially emphasized the effects of our base model because

our study was designed with respect to land use and season. However, for two of the four

response variables (development time and wing length), the model based on microclimate

performed better than the base model by a considerable degree (e.g., $\Delta AIC_c > 18$: Table 1), and

the microclimate and base models performed similarly for per capita population growth rate

($\Delta AIC_c=1.1$). In contrast, the base model performed better than the microclimate model for

emergence ($\Delta AIC_c=6.3$). The null model, which excluded abiotic variables, always performed

the poorest (Table 1; $\Delta AIC_c >80$). In general, the microclimate models showed that mosquito

performance increased with increasing warming temperature (T_{min} or T_{max}) and increasing

minimum relative humidity (R_{min}), although the magnitude of these effects often depended on

density (Table 1, Figure 2). Thus, the microclimate effects captured the observed variation in

mosquito traits as effectively as season and land use and further highlighted the interactions of

biotic (density) and with abiotic (microclimate) variables.

DISCUSSION

To date, the majority of studies investigating the effects of various abiotic and biotic

factors on mosquito fitness, population dynamics, vector competence, and transmission potential

have explored the effect of varying a single environmental variable such as temperature (e.g.,

Agyekum et al., 2022; Liu-Helmersson et al., 2014; Monteiro et al., 2007), conspecific density

(e.g., Hawley, 1985; Reiskind and Lounibos, 2009; Sauers et al., 2022), or food resources (e.g.,

Levi et al., 2014; Telang and Wells, 2004). Further, these studies are often conducted in

laboratory environments under relatively constant conditions. However, mosquito populations in

the field will be subject to the effects of many different environmental variables that fluctuate

diurnally and seasonally and likely interact. Previous work in the *Ae. albopictus* system suggests

that subtle variation in microclimate in the field can translate into significant variation in

mosquito life history traits, population dynamics, and arbovirus transmission potential (Evans et

al., 2018; 2019; Murdock et al., 2017; Wimberly et al., 2020). Our experimental study builds on

these results by showing that season and land use (via their effects on temperature and relative

humidity) can interact with variation in key biotic variables (conspecific density) to affect

metrics of mosquito fitness. The results from this study have broad implications for our

understanding and ability to predict the effects of current and future environmental variation on

mosquito fitness and population dynamics, and the environmental suitability for mosquito-borne

disease transmission.

Land use affects microclimate with subtle effects on metrics of mosquito fitness and

population dynamics

Like previous studies conducted in the same region (Athens, Georgia) (Evans et al., 2018;

Murdock et al., 2017; Wimberly et al., 2020), urban sites had higher minimum temperatures than

rural sites and were drier than non-urban land use types. This is consistent with the Urban Heat

Island (UHI) effect, where nighttime temperatures (analogous to T_{min}) are 1-2.5ºC higher

(Hibbard et al., 2017) than surrounding areas due to the higher percentage of impervious surfaces

in urban areas that absorb, store, and re-emit heat (e.g., asphalt and concrete). However, in non-

urban areas, hydrologically permeable land covers re-emit less heat (e.g., forests, shrublands,

wetlands), have higher evapotranspiration rates, and often provide ample shading resulting in

lower temperatures and higher relative humidity than urban areas (Bowler et al., 2010; Brown et

al., 2023; Huang et al., 1987; Jabbar et al., 2023). Despite the variation in microclimate

observed across land use, this variation was small (Appendix S1: Figure S2) and for the most

part did not significantly affect components of mosquito fitness or estimated per capita growth

rates within a given season (Appendix S1: Figure S4). This is not necessarily surprising as

Athens-Clarke County is a small city in Georgia, with less variation in impervious surface cover,

vegetation cover, and ambient temperatures than seen in larger cities in the U.S. and worldwide

(Oke, 1973; Peng et al., 2011; Zhou et al., 2017). We did observe effects of land use on the

relationship between conspecific density and juvenile mosquito survival (i.e., emergence),

suggesting the effects of intra-specific competition could potentially be modified across an urban

gradient. However, while significant, the effects of the interaction between land use and density

on metrics of mosquito fitness were relatively small (Appendix S1: Figure S4), likely because

variation in microclimate (e.g., T_{min} and R_{min}) was not large across Athens within a given season

(Appendix S1: Figure S3).

Seasonal variation in microclimate affects metrics of mosquito fitness and population

dynamics

Seasonal variation in microclimate was much larger than what we observed across land

use, with temperature and relative humidity being higher during the summer field trial relative to

the fall. As a result, we observed strong seasonal effects on juvenile survival, the juvenile

development time, adult body size, and estimated per capita growth rates. Mosquitoes are

ectothermic, and their trait performance has been shown to have non-linear relationships with

temperature variation across a diversity of mosquito species (reviewed by: Mordecai et al.,

2019). For example, laboratory work conducted under a range of constant temperatures predicts

Ae. albopictus juvenile survival and mosquito development rates are maximized at ~25°C and

~35°C, respectively (Mordecai et al., 2017). Thus, it is unsurprising that we generally saw faster

juvenile development times and increased adult female emergence in the warmer summer field

trial (with average temperatures ranging from 25-25.7°C) than the cooler fall (with temperatures

ranging from 21.1-21.6°C) (Figures 1, 2, Appendix S1: Figure S3).

Seasonal variation in microclimate shapes the relationship between intra-specific

competition and metrics of mosquito fitness and population dynamics

In general, our results support previous research that shows decreasing temperature or increasing

conspecific larval densities result in decreased juvenile survival, smaller adults upon emergence,

and decreased per capita growth rates (Lord, 1998; Sauers et al., 2022; Walker et al., 1991).

However, our results also highlight a critical and novel role of seasonality and land use in

modifying the effects of conspecific larval density on mosquito fitness and population dynamics.

In both the summer and fall, we observed a negative effect of increasing conspecific larval

density on juvenile survival (i.e., the proportion of adult females emerging). However, this trend

was much stronger during the fall trial (Figure 1a). This could be due to a hurricane that

occurred at the beginning of the fall trial (day three) when larvae were in their most vulnerable

stage (i.e., first instar) and/or due to cooler fall temperatures being below the thermal optimum

for *Ae. albopictus* larval survival (~25°C: Mordecai et al., 2017). For example, mean

temperatures in the fall were <15ºC for three consecutive days. These temperatures, which can

result in high *Ae. albopictus* larval and pupal mortality (reviewed in Waldock et al., 2013),

combined with the additional stress of developing under high intra-specific competition, could

synergistically increase juvenile mortality. This interactive effect of temperature and density on

juvenile survival is further supported by our microclimate analyses, which demonstrated that

effects of cooler temperatures were more severe at higher densities (Figure 2a).

Contrary to our initial hypothesis and many prior studies (e.g., Ezeakacha and Yee, 2019;

Sauers et al., 2022), larvae appeared to develop faster with higher conspecific densities in the

cooler fall (Figures 1b, 2b). Several factors could explain this result. First, the timing of

juvenile mortality is known to affect the amount of resources available to the surviving larvae

(Evans et al., 2022; McIntire and Juliano, 2018; Neale and Juliano, 2019). If mortality of early

instar larvae was elevated by cooler fall temperatures and high intra-specific densities, the

surviving larvae would have experienced reduced intra-specific competition and increased access

to food resources, both of which allow larvae to develop faster (Evans et al., 2022; Neale and

Juliano, 2019; Teng and Apperson, 2000). Stage-specific overcompensation and the Hydra

effect have been detected in *Ae. albopictus* when early larval mortality occurred (McIntire and

Juliano, 2018). Another possible explanation may be survival bias or frailty. Very few females

survived in the fall from the medium-density treatment (7.2%) and even fewer from the high-

density treatment (1.2%). If the few individuals that did survive were of higher quality (e.g.,

competitive ability), then their subsequent development time might be expected to be greater

than observed in the low-density treatment. A third explanation could be that larvae facultatively

alter their development rate based on a stressor such as low temperatures and/or low quantity of

nutritional resources due to increased competition. While not demonstrated before in *Ae.*

albopictus, previous work in the *Ae. triseriatus* and *Ae. communis* systems have shown larvae

can speed up development rates in environments subject to evaporation (Chodorowski, 1969;

Juliano and Stoffregen, 1994).

Our results contrast with the extensive literature documenting relationships between

mosquito wing size (i.e. size upon emergence) with variation in conspecific density (Alto et al.,

2005; Evans et al., 2021; Lord, 1998) or temperature (Kingsolver and Huey, 2008; Rueda et al.,

1990). Warmer temperatures have been shown to typically result in smaller wings and body size

– "hotter is smaller"– and cooler temperatures typically result in larger wings and body size –

"cooler is bigger" hypothesis (Alto and Juliano, 2001; Ezeakacha and Yee, 2019; Kingsolver and

Huey, 2008; Rueda et al., 1990). Our results contradict these hypotheses. Instead, we

consistently found that wing length increased with an increase in the minimum temperature,

although the slope of this relationship was less for mosquitoes developing at high conspecific

densities (Figure 2c). Only a few studies have demonstrated smaller wing lengths at cooler

temperatures (Lounibos et al., 2002; Reiskind and Zarrabi, 2012). This result could be attributed

to slower microbial growth in the fall and mosquito anabolism being less efficient (due to cooler

temperatures) (Reiskind and Janairo, 2018). Thus, with females in the warmer summer

experiencing more microbially rich habitats regardless of larval density treatment as well as a

greater capacity to incorporate nutrients into tissue, facilitating more rapid adult female

emergence and larger body sizes in the summer. Finally, like development time, mosquito size

upon emergence converges at high densities regardless of season, suggesting that high amounts

of intra-specific competition could eventually outweigh the influence of variation in abiotic

factors such as temperature and relative humidity.

In general, while growth rates declined as larval conspecific densities increased (see also:

Edgerly and Livdahl, 1992; Lord, 1998; Reiskind and Lounibos, 2009) and increased at higher

temperatures (Alto and Juliano, 2001), we found the estimated per capita growth rates at higher

temperatures (e.g., in the summer) were less affected by intra-specific competition than at lower

temperatures (e.g., in the fall). This intriguing result suggests that mosquito populations might

be less strongly regulated by density-dependent processes as temperatures increase on the

landscape Most studies to date have considered the effects of temperature or conspecific density

on growth rates in isolation, without considering the possible interaction. Those that have

considered the interaction between temperature and intra-specific competition on per capita

growth rates have generally found intra-specific competition to intensify as temperatures warm

(Evans et al., 2021; Teng and Apperson, 2000), which is not what we observed in this study.

However, we did not account for possible effects of temperature and relative humidity on adult

longevity or on size-specific fecundity, which could potentially modify our estimates of the per

capita population growth. Given the interest in vector-pathogen dynamics in response to global

change, interactions between temperature and density dependence warrant further investigation.

CONCLUSION

This study has several important implications. It is generally thought that biotic factors

(e.g., intra-specific and inter-specific competition, quality and quantity of habitat, biological

enemies) will vary more at fine spatial scales, driving local variation in organismal fitness and

population dynamics, whereas abiotic factors will dominate regional effects because they vary at

larger spatial scales (McGill, 2010; Russell et al., 2006; Wiens 1989). Our study demonstrates

that while variation in biotic factors (e.g., intra-specific competition) and abiotic factors (e.g.,

temperature and relative humidity) are both influential, they interact to affect mosquito

population dynamics at local scales. Ignoring the interaction between variation in biotic and

abiotic variables could reduce the accuracy and precision of models (no matter their scale) used

to predict species distributions and population dynamics, as well as the transmission dynamics of

mosquito-borne pathogens. By not accounting for these interactions, mismatches in model

predictions could result in unexpected spatial and temporal variation in species densities and

host-parasite interactions, especially for organisms with complex life cycles like mosquitoes

(Moller-Jacobs et al., 2014; Shapiro et al., 2016). While our results demonstrate the critical role

of abiotic factors like temperature and relative humidity that are linked to variation in land use

and season, future work should incorporate other aspects of land use and seasonality, such as

habitat quality that arises through variation in detrital inputs and precipitation, which will affect

the strength of intraspecific competition and potentially the interaction with abiotic variables.

CONTRIBUTIONS

CCM designed the experiment. NS, ECH, CWH, PMN, AMS, and JWW conducted fieldwork.

NS led, and GRJ assisted with the statistical analyses, with guidance from CCM and CWO. NS

drafted the manuscript with guidance from CCM and CWO. All authors were involved in

reviewing and editing the manuscript. All authors read and approved the final manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank members of the Murdock lab (UGA and Cornell), Osenberg lab (UGA), and Drake lab

(UGA) for their helpful discussions and field support. We particularly thank Sydney Habegger,

Luis Hernandez, and Abigail LeCroy for their assistance in the lab and field, Juliana Hoyos,

Maria Luisa Müller Theissen, and Kimberly Archbold for their feedback on the Spanish abstract,

Ashutosh Pathak for the thoughtful discussions and guidance with the statistical analyses, the

property owners who provided us access to the study sites, and several reviewers and Kathryn

Cottingham for helpful comments on previous submissions. We also appreciate the financial

support from the National Science Foundation Research Experiences for Undergraduates (Grant

no: 1659683), the National Science Foundation Research Traineeship: Interdisciplinary Disease

Ecology Across Scales (Grant no: 1545433), the National Institutes of Health Office of Research

Infrastructure Programs (Grant No: 2T35OD010433-11), and start-up resources provided by the

University of Georgia.

REFERENCES

Agyekum, Thomas P., John Arko-Mensah, Paul K. Botwe, et al. 2022. "Effects of Elevated

Temperatures on the Growth and Development of Adult *Anopheles gambiae* (s.l.)

(Diptera: Culicidae) Mosquitoes." *Journal of Medical Entomology* 59 (4): 1413–20.

https://doi.org/10.1093/jme/tjac046.

Alto, Barry W., and David Bettinardi. 2013. "Temperature and Dengue Virus Infection in

Mosquitoes: Independent Effects on the Immature and Adult Stages." *The American*

Journal of Tropical Medicine and Hygiene 88 (3): 497. https://doi.org/10.4269/ajtmh.12-

0421.

Alto, B. W. & Juliano, S. A. 2001. Precipitation and temperature effects on populations of *Aedes*

albopictus (Diptera: Culicidae): Implications for range expansion. Journal of Medical

Entomology 38:646-656

Alto, Barry W., L. Philip Lounibos, Stephen Higgs, and Steven A. Juliano. 2005. "Larval

Competition Differentially Affects Arbovirus Infection in *Aedes Mosquitoes*." *Ecology*

86 (12): 3279–88. https://doi.org/10.1890/05-0209.

Amarasekare, Priyanga. 2024. "Temperature-Dependent Dispersal and Ectotherm Species'

Distributions in a Warming World." *The Journal of Animal Ecology* 93 (4): 428–46.

https://doi.org/10.1111/1365-2656.14054.

Bale, Jeffery S., Gregory J. Masters, Ian D. Hodkinson, et al. 2002. "Herbivory in Global

Climate Change Research: Direct Effects of Rising Temperature on Insect Herbivores."

Global Change Biology 8 (1): 1–16. https://doi.org/10.1046/j.1365-2486.2002.00451.x.

Beale, Colin M., Mark J. Brewer, and Jack J. Lennon. 2014. "A New Statistical Framework for

the Quantification of Covariate Associations with Species Distributions." *Methods in*

Ecology and Evolution 5 (5): 421–32. https://doi.org/10.1111/2041-210X.12174.

Benedict, Mark Q., Rebecca S. Levine, William A. Hawley, and L. Philip Lounibos. 2007.

"Spread of The Tiger: Global Risk of Invasion by The Mosquito *Aedes albopictus*."

Vector-Borne and Zoonotic Diseases 7 (1): 76–85.

https://doi.org/10.1089/vbz.2006.0562.

Bolker, Ben and R. Development Core Team. 2023. *Bbmle: Tools for General Maximum*

Likelihood Estimation. V. 1.0.25.1. Released December 9. https://cran.r-

project.org/web/packages/bbmle/index.html.

Bonizzoni, Mariangela, Giuliano Gasperi, Xioaguang Chen, and Anthony A. James. 2013. "The

Invasive Mosquito Species *Aedes albopictus*: Current Knowledge and Future

Perspectives." *Trends in Parasitology* 29 (9): 460–68.

https://doi.org/10.1016/j.pt.2013.07.003.

Bowler, Diana E., Lisette Buyung-Ali, Teri M. Knight, and Andrew S. Pullin. 2010. "Urban

Greening to Cool Towns and Cities: A Systematic Review of the Empirical Evidence."

Landscape and Urban Planning 97 (3): 147–55.

https://doi.org/10.1016/j.landurbplan.2010.05.006.

Bozinovic, Francisco, Daniel A. Bastías, Francisca Boher, Sabrina Clavijo-Baquet, Sergio A.

Estay, and Michael J. Angilletta. 2011. "The Mean and Variance of Environmental

Temperature Interact to Determine Physiological Tolerance and Fitness." *Physiological*

and Biochemical Zoology 84 (6): 543–52. https://doi.org/10.1086/662551.

Briegel, Hans. 2003. "Physiological Bases of Mosquito Ecology." *Journal of Vector Ecology:*

Journal of the Society for Vector Ecology 28 (1): 1–11.

Brooks, Mollie E., Kasper Kristensen, Koen J. van Benthem, et al. 2017. "glmmTMB Balances

Speed and Flexibility Among Packages for Zero-Inflated Generalized Linear Mixed

Modeling." *The R Journal* 9 (2): 378–400. https://doi.org/10.32614/RJ-2017-066.

Brown, Joel J., Mercedes Pascual, Michael C. Wimberly, Leah R. Johnson, and Courtney C.

Murdock. 2023. "Humidity – The Overlooked Variable in the Thermal Biology of

Mosquito-Borne Disease." *Ecology Letters* 26 (7): 1029–49.

https://doi.org/10.1111/ele.14228.

Bullock, James M., and Gesine Pufal. 2020. "Human-Mediated Dispersal as a Driver of

Vegetation Dynamics: A Conceptual Synthesis." *Journal of Vegetation Science* 31 (6):

943–53. https://doi.org/10.1111/jvs.12888.

Chala, Bayissa, and Feyissa Hamde. 2021. "Emerging and Re-Emerging Vector-Borne Infectious

Diseases and the Challenges for Control: A Review." *Frontiers in Public Health* 9

(October). https://doi.org/10.3389/fpubh.2021.715759.

Chodorowski, A. 1969. "The Desiccation of Ephemeral Pools and the Rate of Development in

Aedes communis Larvae." *Polish Archive of Hydrobiology* 16: 79–91.

Cohen, Jeremy M., David J. Civitello, Amber J. Brace, et al. 2016. "Spatial Scale Modulates the

Strength of Ecological Processes Driving Disease Distributions." PNAS Plus.

Proceedings of the National Academy of Sciences 113 (24): E3359–64.

https://doi.org/10.1073/pnas.1521657113.

Darsie, Richard F., and Ronald A. Ward. 2005. *Identification and Geographical Distribution of*

the Mosquitos of North America, North of Mexico. University Press of Florida.

Gainesville, FL.

Edgerly, Janice S., and Todd P. Livdahl. 1992. "Density-Dependent Interactions Within a

Complex Life Cycle: The Roles of Cohort Structure and Mode of Recruitment." *Journal*

of Animal Ecology 61 (1): 139–50. https://doi.org/10.2307/5517.

Evans, Katherine G., Zoey R. Neale, Brendan Holly, Cecilia C. Canizela, and Steven A. Juliano.

2022. "Survival-Larval Density Relationships in the Field and Their Implications for

Control of Container-Dwelling *Aedes Mosquitoes.*" *Insects* 14 (1): 17.

https://doi.org/10.3390/insects14010017.

Evans, Michelle V., John M. Drake, Lindsey Jones, and Courtney C. Murdock. 2021. "Assessing

Temperature-Dependent Competition between Two Invasive Mosquito Species."

Ecologtcal Applications 31 (5): e02334. https://doi.org/10.1002/eap.2334.

Evans, Michelle V., Carl W. Hintz, Lindsey Jones, et al. 2019. "Microclimate and Larval Habitat

Density Predict Adult *Aedes albopictus* Abundance in Urban Areas." *The American*

Journal of Tropical Medicine and Hygiene 101 (2): 362–70.

https://doi.org/10.4269/ajtmh.19-0220.

Evans, Michelle V., Justine C. Shiau, Nicole Solano, Melinda A. Brindley, John M. Drake, and

Courtney C. Murdock. 2018. "Carry-over Effects of Urban Larval Environments on the

Transmission Potential of Dengue-2 Virus." *Parasites & Vectors* 11 (1): 426.

https://doi.org/10.1186/s13071-018-3013-3.

Ewing, David A., Bethan V. Purse, Christina A. Cobbold, Stefanie M. Schäfer, and Steven M.

White. 2019. "Uncovering Mechanisms behind Mosquito Seasonality by Integrating

Mathematical Models and Daily Empirical Population Data: Culex Pipiens in the UK."

Parasites & Vectors 12 (1): 1. https://doi.org/10.1186/s13071-019-3321-2.

Ezeakacha, Nnaemeka F., and Donald A. Yee. 2019. "The Role of Temperature in Affecting

Carry-over Effects and Larval Competition in the Globally Invasive Mosquito *Aedes*

albopictus." *Parasites & Vectors* 12 (1): 1. https://doi.org/10.1186/s13071-019-3391-1.

Fox, John, Sanford Weisberg, Brad Price, et al. 2024. *Car: Companion to Applied Regression*. V.

3.1-3. Released September 27. https://cran.r-project.org/web/packages/car/index.html.

Golding, Nick, Miles A. Nunn, and Bethan V. Purse. 2015. "Identifying Biotic Interactions

Which Drive the Spatial Distribution of a Mosquito Community." *Parasites & Vectors* 8

(1): 1. https://doi.org/10.1186/s13071-015-0915-1.

Govindarajulu, Purnima P., and Bradley R. Anholt. 2006. "Interaction between Biotic and

Abiotic Factors Determines Tadpole Survival Rate under Natural Conditions."

Écoscience 13 (3): 413–21. https://doi.org/10.2980/i1195-6860-13-3-413.1.

Hartig, Florian, Lukas Lohse, and Melina de Souza leite. 2024. *DHARMa: Residual Diagnostics*

for Hierarchical (Multi-Level / Mixed) Regression Models. V. 0.4.7. Released October

18. https://cran.r-project.org/web/packages/DHARMa/index.html.

Hawley, W. A. 1985. "The Effect of Larval Density on Adult Longevity of a Mosquito, *Aedes*

sierrensis: Epidemiological Consequences." *Journal of Animal Ecology* 54 (3): 955–64.

JSTOR. https://doi.org/10.2307/4389.

Helmuth, Brian, Bernardo R. Broitman, Lauren Yamane, et al. 2010. "Organismal Climatology:

Analyzing Environmental Variability at Scales Relevant to Physiological Stress." *Journal*

of Experimental Biology 213 (6): 995–1003. https://doi.org/10.1242/jeb.038463.

Hibbard, K. A., F. M. Hoffman, D. Huntzinger, et al. 2017. *Changes in Land Cover and*

Terrestrial Biogeochemistry. Climate Science Special Report: Fourth National Climate

Assessment, Volume I. U.S. Global Change Research Program.

https://doi.org/10.7930/J0416V6X.

Houghton, J. T., James J. McCarthy, and Bert Metz. 2002. "Climate Change 2001: The Scientific

Basis." *Foreign Affairs* 81 (1): 208. https://doi.org/10.2307/20033020.

Huang, Y. J., H. Akbari, H. Taha, and A. H. Rosenfeld. 1987. "The Potential of Vegetation in

Reducing Summer Cooling Loads in Residential Buildings." Journal of Applied

Meteorology and Climatology. *Journal of Applied Meteorology and Climatology* 26 (9):

1103–16.

Huey, Raymond B., David Berrigan, and Donald B. Miles. 2001. "Temperature, Demography,

and Ectotherm Fitness." *American Naturalist* 158 (2): 204. 4784459.

https://doi.org/10.1086/321314.

Huxley, Paul J., Joel J. Brown, Brandy St. Laurent, et al. 2025. Beyond temperature: Relative

humidity systematically shifts the temperature dependence of population growth in a

malaria vector. bioRxiv 2025.05.30.656372. https://doi.org/10.1101/2025.05.30.656372

Indermaur, Lukas, Benedikt R. Schmidt, Klement Tockner, and Michael Schaub. 2010. "Spatial

Variation in Abiotic and Biotic Factors in a Floodplain Determine Anuran Body Size and

Growth Rate at Metamorphosis." *Oecologia* 163 (3): 637–49.

https://doi.org/10.1007/s00442-010-1586-4.

Jabbar, Hajer Khaled, Mustafa N. Hamoodi, and Amjed N. Al-Hameedawi. 2023. "Urban Heat

Islands: A Review of Contributing Factors, Effects and Data." *IOP Conference Series:*

Earth and Environmental Science 1129 (1): 012038. https://doi.org/10.1088/1755-

1315/1129/1/012038.

Juliano, SA, and TL Stoffregen. 1994. "Effects of Habitat Drying on Size at and Time to

Metamorphosis in the Tree Hole Mosquito *Aedes triseriatus*." *Oecologia* 97 (3): 369–76.

28313632, pp. 369–76. PubMed. https://doi.org/10.1007/bf00317327.

Juliano, Steven A. 2009. "Species Interactions Among Larval Mosquitoes: Context Dependence

Across Habitat Gradients." *Annual Review of Entomology* 54 (Volume 54, 2009): 37–56.

https://doi.org/10.1146/annurev.ento.54.110807.090611.

Kingsolver, Joel G., and Raymond B. Huey. 2008. "Size, Temperature, and Fitness: Three

Rules." *Evolutionary Ecology Research* 10: 251–68.

Kraemer, Moritz U. G., Robert C. Reiner, Oliver J. Brady, et al. 2019. "Past and Future Spread

of the Arbovirus Vectors *Aedes aegypti* and *Aedes albopictus*." *Nature Microbiology* 4

(5): 854–63. https://doi.org/10.1038/s41564-019-0376-y.

Leach, Katie, W. Ian Montgomery, and Neil Reid. 2016. "Modelling the Influence of Biotic

Factors on Species Distribution Patterns." *Ecological Modelling* 337 (October): 96–106.

https://doi.org/10.1016/j.ecolmodel.2016.06.008.

Leibold, M., and JM Chase. 2018. *Metacommunity Ecology | Princeton University Press*.

Princeton University Press.

Lenth, Russell V., Balazs Banfai, Ben Bolker, et al. 2025. *Emmeans: Estimated Marginal*

Means, Aka Least-Squares Means. V. 1.11.2-8. Released August 27. https://cran.r-

project.org/web/packages/emmeans/index.html.

Levi, Tal, Eitan Ben-Dov, Preeti Shahi, Dov Borovsky, and Arieh Zaritsky. 2014. "Growth and

Development of *Aedes aegypti* Larvae at Limiting Food Concentrations." *Acta Tropica*

133 (May): 42–44. https://doi.org/10.1016/j.actatropica.2014.02.001.

Levin, Simon A. 1992. "The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur

Award Lecture." *Ecology* 73 (6): 1943–67. https://doi.org/10.2307/1941447.

Liu-Helmersson, Jing, Åke Brännström, Maquins Odhiambo Sewe, Jan C. Semenza, and Joacim

Rocklöv. 2019. "Estimating Past, Present, and Future Trends in the Global Distribution

and Abundance of the Arbovirus Vector *Aedes aegypti* Under Climate Change

Scenarios." *Frontiers in Public Health* 7 (June).

https://doi.org/10.3389/fpubh.2019.00148.

Liu-Helmersson, Jing, Hans Stenlund, Annelies Wilder-Smith, and Joacim Rocklöv. 2014.

"Vectorial Capacity of *Aedes aegypti*: Effects of Temperature and Implications for

707   Global Dengue Epidemic Potential." *PLOS ONE* 9 (3): e89783.

708   https://doi.org/10.1371/journal.pone.0089783.

709 Livdahl, Todd P., and George Sugihara. 1984. "Non-Linear Interactions of Populations and the

710   Importance of Estimating Per Capita Rates of Change." *Journal of Animal Ecology* 53

711   (2): 573–80. JSTOR. https://doi.org/10.2307/4535.

712 Lord, Cynthia C. 1998. "Density Dependence in Larval *Aedes albopictu*s (Diptera: Culicidae)."

713   *Journal of Medical Entomology* 35 (5): 5.

714 Lounibos, L. Philip. 2002. "Invasions by Insect Vectors of Human Disease." *Annual Review of*

715   *Entomology* 47 (Volume 47, 2002): 233–66.

716   https://doi.org/10.1146/annurev.ento.47.091201.145206.

717 Lounibos, L.P., S. Suarez, Z. Menendez, et al. 2002. "Does Temperature Affect the Outcome of

718   Larval Competition between *Aedes aegypti* and *Aedes albopictus*?" *Journal of Vector*

719   *Ecology* 27 (1): 86–95.

720 Martin, Tara Laine, and Raymond B. Huey. 2008. "Why 'Suboptimal' Is Optimal: Jensen's

721   Inequality and Ectotherm Thermal Preferences." *The American Naturalist* 171 (3): E102–

722   18. https://doi.org/10.1086/527502.

723 McGill, Brian J. 2010. "Matters of Scale." *Science* 328 (5978): 575–76.

724   https://doi.org/10.1126/science.1188528.

725 McIntire, Kristina M., and Steven A. Juliano. 2018. "How Can Mortality Increase Population

726   Size? A Test of Two Mechanistic Hypotheses." *Ecology* 99 (7): 1660–70.

727   https://doi.org/10.1002/ecy.2375.

McKinney, Michael L., and Julie L. Lockwood. 1999. "Biotic Homogenization: A Few Winners

Replacing Many Losers in the Next Mass Extinction." *Trends in Ecology & Evolution* 14

(11): 450–53. https://doi.org/10.1016/S0169-5347(99)01679-1.

Menge, Bruce A., and John P. Sutherland. 1976. "Species Diversity Gradients: Synthesis of the

Roles of Predation, Competition, and Temporal Heterogeneity." *The American Naturalist*

110 (973): 351–69.

Miazgowicz, K. L., M. S. Shocket, S. J. Ryan, et al. 2020. "Age Influences the Thermal

Suitability of *Plasmodium falciparum* Transmission in the Asian Malaria Vector

Anopheles stephensi." *Proceedings of the Royal Society B*, ahead of print, July 29. World.

https://doi.org/10.1098/rspb.2020.1093.

Moller-Jacobs, Lillian L., Courtney C. Murdock, and Matthew B. Thomas. 2014. "Capacity of

Mosquitoes to Transmit Malaria Depends on Larval Environment." *Parasites & Vectors*

7 (1): 1. https://doi.org/10.1186/s13071-014-0593-4.

Monteiro, Laura C. C., José R. B. de Souza, and Cleide M. R. de Albuquerque. 2007. "Eclosion

Rate, Development and Survivorship of *Aedes albopictus* (Skuse) (Diptera: Culicidae)

under Different Water Temperatures." *Neotropical Entomology* 36 (6): 966–71.

https://doi.org/10.1590/s1519-566x2007000600021.

Mordecai Erin A., Jeremy M. Cohen, Michelle V. Evans, et al. 2017. "Detecting the Impact of

Temperature on Transmission of Zika, Dengue, and Chikungunya Using Mechanistic

Models." PLoS Negl Trop Dis 11(4): e0005568.

https://doi.org/10.1371/journal.pntd.0005568

Mordecai, Erin A., Jamie M. Caldwell, Marissa K. Grossman, et al. 2019. "Thermal Biology of

Mosquito-Borne Disease." *Ecology Letters* 22 (10): 1690–708.

https://doi.org/10.1111/ele.13335.

Murdock, Courtney C., Michelle V. Evans, Taylor D. McClanahan, Kerri L. Miazgowicz, and

Blanka Tesla. 2017. "Fine-Scale Variation in Microclimate across an Urban Landscape

Shapes Variation in Mosquito Population Dynamics and the Potential of *Aedes albopictus*

to Transmit Arboviral Disease." *PLOS Neglected Tropical Diseases* 11 (5): e0005640.

https://doi.org/10.1371/journal.pntd.0005640.

Neale, Zoey R., and Steven A. Juliano. 2019. "Finding the Sweet Spot: What Levels of Larval

Mortality Lead to Compensation or Overcompensation in Adult Production?" *Ecosphere*

10 (9): e02855. https://doi.org/10.1002/ecs2.2855.

Oke, T. R. 1973. "City Size and the Urban Heat Island." *Atmospheric Environment* 7 (8): 769–

79. https://doi.org/10.1016/0004-6981(73)90140-6.

Packer, M. J., and Philip S. Corbet. 1989. "Size Variation and Reproductive Success of Female

Aedes punctor (Diptera: Culicidae)." *Ecological Entomology* 14 (3): 297–309.

https://doi.org/10.1111/j.1365-2311.1989.tb00960.x.

Parmesan, Camille. 2006. "Ecological and Evolutionary Responses to Recent Climate Change."

Annual Review of Ecology, Evolution, and Systematics 37 (Volume 37, 2006): 637–69.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110100.

Peng, Shushi, Shilong Piao, Philippe Ciais, et al. 2011. "Surface Urban Heat Island Across 419

Global Big Cities." *Environmental Science & Technology* 46 (2): 696-703.

https://doi.org/10.1021/es2030438.

Pörtner, Hans O., and Anthony P. Farrell. 2008. "Physiology and Climate Change." *Science* 322

(5902): 690–92. https://doi.org/10.1126/science.1163156.

R Core Team. 2025. *R: A Language and Environment for Statistical Computing.* R Foundation

for Statistical Computing. Vienna, Austria. https://www.R-project.org/

Reiskind, M. H., and L. P. Lounibos. 2009. "Effects of Intraspecific Larval Competition on

Adult Longevity in the Mosquitoes *Aedes aegypti* and *Aedes albopictus*." *Medical and*

Veterinary Entomology 23 (1): 62–68. https://doi.org/10.1111/j.1365-2915.2008.00782.x.

Reiskind, Michael H., and Ali A. Zarrabi. 2012. "Is Bigger Really Bigger? Differential

Responses to Temperature in Measures of Body Size of the Mosquito, *Aedes albopictus*."

Journal of Insect Physiology 58 (7): 911–17.

https://doi.org/10.1016/j.jinsphys.2012.04.006.

Reiskind, Michael, and M. Shawn Janairo. 2018. "Tracking *Aedes aegypti* (Diptera: Culicidae)

Larval Behavior Across Development: Effects of Temperature and Nutrients on

Individuals' Foraging Behavior." *Journal of Medical Entomology* 55 (5): 1086–92.

https://doi.org/10.1093/jme/tjy073.

Roiz, David, Paulina A. Pontifes, Fréderic Jourdain, et al. 2024. "The Rising Global Economic

Costs of Invasive *Aedes* Mosquitoes and *Aedes*-Borne Diseases." *Science of The Total*

Environment 933 (July): 173054. https://doi.org/10.1016/j.scitotenv.2024.173054.

Rueda, L. M., K. J. Patel, R. C. Axtell, and R. E. Stinner. 1990. "Temperature-Dependent

Development and Survival Rates of *Culex quinquefasciatu*s and *Aedes aegypti* (Diptera:

Culicidae)." *Journal of Medical Entomology* 27 (5): 892–98.

https://doi.org/10.1093/jmedent/27.5.892.

Ruel, Jonathan J., and Matthew P. Ayres. 1999. "Jensen's Inequality Predicts Effects of

Environmental Variation." *Trends in Ecology & Evolution* 14 (9).

https://doi.org/10.1016/s0169-5347(99)01664-x.

Russell, Roly, Spencer A. Wood, Gary Allison, and Bruce A. Menge. 2006. "Scale,

Environment, and Trophic Status: The Context Dependency of Community Saturation in

Rocky Intertidal Communities." *The American Naturalist* 167 (6): E158–70.

https://doi.org/10.1086/504603.

Ryan, Sadie J., Catherine A. Lippi, Oswaldo C. Villena, Aspen Singh, Courtney C. Murdock,

and Leah R. Johnson. 2023. "Mapping Current and Future Thermal Limits to Suitability

for Malaria Transmission by the Invasive Mosquito *Anopheles stephensi*." *Malaria*

Journal 22 (1): 104. https://doi.org/10.1186/s12936-023-04531-4.

Sauers, Logan A., Kelsey E. Hawes, and Steven A. Juliano. 2022. "Non-Linear Relationships

between Density and Demographic Traits in Three *Aedes* Species." *Scientific Reports* 12

(May): 8075. https://doi.org/10.1038/s41598-022-11909-y.

Shapiro, Lillian L. M., Courtney C. Murdock, Gregory R. Jacobs, Rachel J. Thomas, and

Matthew B. Thomas. 2016. "Larval Food Quantity Affects the Capacity of Adult

Mosquitoes to Transmit Human Malaria." *Proceedings of the Royal Society B: Biological*

Sciences, ahead of print, July 13. World. https://doi.org/10.1098/rspb.2016.0298.

Shapiro, Lillian L. M., Shelley A. Whitehead, and Matthew B. Thomas. 2017. "Quantifying the

Effects of Temperature on Mosquito and Parasite Traits That Determine the Transmission

Potential of Human Malaria." *PLOS Biology* 15 (10): e2003489.

https://doi.org/10.1371/journal.pbio.2003489.

Simon, Margaret W., and Priyanga Amarasekare. 2024. "Predicting the Fundamental Thermal

Niche of Ectotherms." *Ecology* 105 (5): e4289. https://doi.org/10.1002/ecy.4289.

Sinka, M. E., S. Pironon, N. C. Massey, et al. 2020. "A New Malaria Vector in Africa: Predicting

the Expansion Range of *Anopheles stephensi* and Identifying the Urban Populations at

Risk." *Proceedings of the National Academy of Sciences* 117 (40): 24900–24908.

https://doi.org/10.1073/pnas.2003976117.

Solano, Nicole. 2026. "Data and Code to Reproduce: Solano et al., 2026 'Mosquito Population

Dynamics Are Shaped by Interactions among Larval Density, Temperature, and

Humidity.'" Figshare. https://doi.org/https://doi.org/10.6084/m9.figshare.25804639.

Taylor, Roz, Louisa A. Messenger, Tarekegn A. Abeku, Sian E. Clarke, Rajpal S. Yadav, and Jo

Lines. 2024. "Invasive *Anopheles stephensi* in Africa: Insights from Asia." *Trends in*

Parasitology 40 (8): 731–43. https://doi.org/10.1016/j.pt.2024.06.008.

Telang, Aparna, and Michael A. Wells. 2004. "The Effect of Larval and Adult Nutrition on

Successful Autogenous Egg Production by a Mosquito." *Journal of Insect Physiology* 50

(7): 677–85. https://doi.org/10.1016/j.jinsphys.2004.05.001.

Teng, Hwa-Jen, and Charles S. Apperson. 2000. "Development and Survival of Immature *Aedes*

albopictus and *Aedes triseriatus* (Diptera: Culicidae) in the Laboratory: Effects of

Density, Food, and Competition on Response to Temperature." *Journal of Medical*

Entomology 37 (1): 40–52. https://doi.org/10.1603/0022-2585-37.1.40.

Tesla, Blanka, Leah R. Demakovsky, Erin A. Mordecai, et al. 2018. "Temperature Drives Zika

Virus Transmission: Evidence from Empirical and Mathematical Models." *Proceedings*

of the Royal Society B 285 (1884): 20180795. https://doi.org/10.1098/rspb.2018.0795.

Vellend, Mark. 2016. *The Theory of Ecological Communities*. Princeton University Press.

Princeton, NJ.

Waldock, Joanna, Nastassya L. Chandra, Jos Lelieveld, et al. 2013. "The Role of Environmental

Variables on *Aedes albopictus* Biology and Chikungunya Epidemiology." *Pathogens and*

Global Health 107 (5): 224–41. https://doi.org/10.1179/2047773213Y.0000000100.

Walker, Edward D., Daniel L. Lawson, Richard W. Merritt, William T. Morgan, and Michael J.

Klug. 1991. "Nutrient Dynamics, Bacterial Populations, and Mosquito Productivity in

Tree Hole Ecosystems and Microcosms." *Ecology* 72 (5): 1529–46.

https://doi.org/10.2307/1940953.

Wiens, J. A. 1989. "Spatial Scaling in Ecology." *Functional Ecology* 3 (4): 385–97.

https://doi.org/10.2307/2389612.

Wimberly, Michael C., Justin K. Davis, Michelle V. Evans, et al. 2020. "Land Cover Affects

Microclimate and Temperature Suitability for Arbovirus Transmission in an Urban

Landscape." *PLOS Neglected Tropical Diseases* 14 (9): e0008614.

https://doi.org/10.1371/journal.pntd.0008614.

Yan, Jiayue, Andrew J. Mackay, and Chris M. Stone. 2024. "Dynamics of Invasive Mosquitoes:

Introduction Pathways, Limiting Factors, and Their Potential Role in Vector-Borne

Pathogen Transmission." *Frontiers in Tropical Diseases* 5 (December).

https://doi.org/10.3389/fitd.2024.1503120.

Zhou, Bin, Diego Rybski, and Jürgen P. Kropp. 2017. "The Role of City Size and Urban Form in

the Surface Urban Heat Island." *Scientific Reports* 7 (1): 4791.

https://doi.org/10.1038/s41598-017-04242-2.

Table 1. Results from Generalized Linear Mixed Effects Models testing the effects of larval density treatment (30 = low, 60 =

medium, 120 = high), land-use type (rural, suburban, urban), and/or microclimate (T_{min}, T_{max}, R_{min}) on mosquito traits across summer

& fall seasons. The mosquito traits were analyzed using three model structures: (1) the Null model – main effect of larval density (but

no other fixed effect); (2) the Base model – main effects of season, land use, and larval density and their interactions; and (3) the

Microclimate model – main effects of larval density, water temperature (T_{min}, T_{max}) and relative humidity (R_{min}) and the interactions

between density and T_{min}, T_{max}, and R_{min}. Experimental site and tray (nested in site) were included as random factors in all models, and

jar (nested in tray) was included for development time and wing length models. ΔAIC_c values, degrees of freedom, and statistically

significant fixed effects are shown. Significance is denoted as *p < 0.05* (*), *p < 0.01* (**), and *p < 0.001* (***). The best model(s)

based on $\Delta AIC_c \leq 2$ is indicated in bold.

Response Variable	Model	$\Delta AICc$	df	Significant Predictors
	Null	399.3	5	Density***
Emergence	**Base**	**0**	**20**	**Density***, Season:Density***, Landuse:Density*****
	Microclimate	6.3	14	Density***, Density:Tmin***, Density:Tmax*, Density:Rmin***
	Null	92	7	Density*
Development Time	Base	30.5	22	Season***, Density**, Season:Density*,
	Microclimate	**0**	**16**	**Density*, Tmin***, Density:Tmin***
	Null	80.4	7	Density***
Wing Length	Base	18.7	22	Season***, Density***
	Microclimate	**0**	**16**	**Density***, Tmin***, Tmax*, Rmin***, Density:Rmin***
	Null	215.4	6	Density***
Intrinsic Growth Rate (r')	**Base**	**0**	**21**	**Season***, Density***, Season:Density*****
	Microclimate	**1.1**	**15**	**Density***, Tmin***, Rmin*, Density:Tmin***, Density:Rmin*****

 FIGURE CAPTIONS

Figure 1. Results of mixed-effects model for *Ae. albopictus:* (**a**) the proportion of female larvae

that emerged (i.e., survival probability), (**b**) female development time (days), (**c**) female wing

length (mm), and (**d**) estimated per capita intrinsic growth rate, r' (day^{-1}), for each density

treatment (low = 30 larvae, medium = 60 larvae, high = 120 larvae) across two seasons (summer

= red, fall = orange). Horizonal and vertical lines give the mean +/- 95% CI, and the dots

indicate raw values [based on females in (b), (c), and jars for (a), (d)] adjusted for random

effects.

Figure 2. Results of mixed-effects model for *Ae. albopictus:* (**a**) the proportion of female larvae

that emerged (i.e., survival probability), (**b**) female development time (days), (**c**) female wing

length (mm), and (**d**) estimated per capita growth rate, r' (day^{-1}), for each of the three density

treatments (low = 30 larvae, medium = 60 larvae, high = 120 larvae), as influenced by minimum

temperature (T_{min}, ºC), and minimum relative humidity (R_{min}, %). Dots indicate raw values

[based on females in (b), (c), and jars for (a), (d)] adjusted for random effects. Black lines give

the predicted effects of T_{min} for each density treatment.

Figure 1

Figure 2