DnaA modulates the gene expression and morphology of the Lyme disease spirochete

Andrew C. Krusenstjerna[1], Nerina Jusufovic[1], Timothy C. Saylor[1], Brian Stevenson[1,2]

1. Department of Microbiology, Immunology, and Molecular Genetics, University of
Kentucky College of Medicine, Lexington, Kentucky, USA
2. Department of Entomology, University of Kentucky, Lexington, Kentucky, USA

Address for correspondence:

Department of Microbiology, Immunology, and Molecular Genetics

University of Kentucky, College of Medicine

Lexington, Ky. 40536-0278

Email: brian.stevenson@uky.edu

Phone: 1-859-257-9358

ABSTRACT.

All bacteria encode a multifunctional DNA-binding protein, DnaA, which initiates

chromosomal replication. Despite having the most complex, segmented bacterial genome, little is

known about *Borrelia burgdorferi* DnaA and its role in maintaining the spirochete's physiology.

In this work we utilized inducible CRISPR-interference and overexpression to modulate cellular

levels of DnaA to better understand this essential protein. Dysregulation of DnaA, either up or

down, increased or decreased cell lengths, respectively, while also significantly slowing

replication rates. Using fluorescent microscopy, we found the DnaA CRISPRi mutants had

increased numbers of chromosomes with irregular spacing patterns. DnaA-depleted spirochetes

also exhibited a significant defect in helical morphology. RNA-seq of the conditional mutants

showed significant changes in the levels of transcripts involved with flagellar synthesis,

elongation, cell division, virulence, and other functions. These findings demonstrate that the

DnaA plays a commanding role in maintaining borrelial growth dynamics and protein

expression, which are essential for the survival of the Lyme disease spirochete.

IMPORTANCE.

Lyme disease is the most prevalent tick-borne infection in the Northern Hemisphere.

Borrelia burgdorferi, the causative spirochete bacteria, has been maintained in nature for

millennia in a consistent enzootic cycle between *Ixodes* ticks and various small vertebrate hosts.

During the tick's blood meal, *B. burgdorferi* substantially increases its replication rate, alters its

repertoire of outer surface proteins, and disseminates into the new vertebrate host. Across

eubacteria, DnaA is the master regulatory protein that initiates chromosomal replication and acts

as a transcription factor to regulate specific pathways. Here, we describe the roles that *B.*

burgdorferi DnaA has on the physiology and gene expression of this medically important

pathogen.

Introduction.

Replication plays a significant role in the enzootic lifecycle of *Borrelia burgdorferi*, the

Lyme disease spirochete. An extracellular bacterial parasite, *B. burgdorferi* exclusively colonizes

Ixodid ticks and assorted vertebrates (1, 2). Naïve tick larvae acquire the Lyme bacteria when

they feed on an infected host. After this single blood meal, the larvae drop off, overwinter, and

molt to their nymphal stage. During this time, *B. burgdorferi* persists within the midgut of the

nutrient-deplete tick. Once the nymph emerges and takes its blood meal, the spirochetes obtain

the requisite materials to replicate, divide, and disseminate from the tick to the new vertebrate

host (3, 4).

The highly conserved AAA+ family protein DnaA initiates the replication of bacterial

chromosomes (5-7). DnaA monomers recognize the *oriC* locus and cooperatively multimerize to

form a helical structure. The DnaA filament promotes the separation of AT-rich DNA elements,

which recruits helicase and replication machinery. DnaA not only initiates chromosomal

replication but also binds elsewhere in the genome to regulate gene expression (8, 9). We

recently found that *B. burgdorferi* DnaA directly regulates the *dnaX-ebfC* operon, which codes

for the Tau (τ) subunit of DNA polymerase III holoenzyme, DnaX, and a regulatory nucleoid-

associated protein, EbfC (10). These genes are highly expressed during periods of rapid

spirochete replication (10-12). At the tick-vertebrate interface, we hypothesize that *B.*

burgdorferi DnaA not only commits to its replication initiation function but coordinates the

expression of genes needed for vertebrate infection (10, 13, 14).

Replication of the *B. burgdorferi* linear chromosome proceeds bidirectionally from the

centrally-located *oriC* (15). Outside of this, little is known about borrelial DNA replication, its

regulation, or coordination with other cellular processes. Many of the genes involved in those

pathways are essential, which has hampered their investigation. Historically, studying essential

loci required the development or isolation of conditional mutants. The first studies into bacterial

DnaA proteins utilized temperature-sensitive mutants (16). In the era of molecular genetics,

many tools have been developed to produce conditional phenotypes more readily, such as

inducible promoters. Overexpression has been a reliable means of elucidating a protein's

function in *B. burgdorferi* and other bacteria (14, 17, 18). Recently, inducible CRISPR

interference (CRISPRi) systems have been developed and added to the repertoire of tools for

studying borrelial biology (19, 20). This work will describe the consequence of CRISPRi-

mediated knockdown of the essential replication initiator protein DnaA in *B. burgdorferi*. Using

this approach, coupled with overexpression, we observed profound consequences of DnaA-

dysregulation on spirochete morphology, chromosome partitioning, and gene expression.

RESULTS.

Construction and validation of plasmids to dysregulate levels of DnaA in *B. burgdorferi*. DnaA is essential for initiating chromosomal replication, so deleting DnaA is lethal to bacteria. Thus, to gain insight into the impacts of DnaA on borrelial physiology, we produced IPTG-inducible constructs to reduce or elevate cellular levels of DnaA. Knockdown was carried out using the newly refined all-in-one CRISPR interference (CRISPRi) shuttle vector, which places the expression of both the sgRNA and a borrelial codon-optimized dCas9 gene under the control of the Lac repressor (20). We designed two CRISPRi constructs with sgRNAs targeting either the template ($dnaA_{T1}$) or non-template ($dnaA_{NT1}$) DNA strands directly 5' or within the *dnaA* gene (**Fig. 1A**). Overexpression was achieved using plasmid pACK121, which contains an inducible *dnaA* with an N-terminal 3xFLAG tag. All constructs were transformed into *B. burgdorferi* strain B31-e2, hereafter referred to as e2. The spirochetes tolerated each construct well, replicating at rates comparable to the parental strain without the inducer (**Fig. 1B**), with one exception. The $dnaA_{T1}$ CRISPRi strain replicated slightly slower than the others, although it reached the same final density. This may suggest leakiness of the hybrid *lac* promoter that controls the sgRNA and dCas9 gene (B. Murphy & W. Zückert, personal communication).

To assess the efficacy of these inducible constructs, spirochetes of each strain were grown to mid-exponential growth phase ($3-5 \times 10^7$ cells/mL), then incubated overnight with 0.5 mM IPTG. The *dnaA*-targeting CRISPRi strains decreased the expression of DnaA, with the construct targeting the template strand, $dnaA_{T1}$, consistently yielding the most effective gene silencing (**Fig. 1C** and **D**). The potency of the $dnaA_{T1}$ construct likely explains the slowed spirochete growth seen in the absence of IPTG (**Fig. 1B**). Overexpression resulted in elevated quantities of DnaA protein and transcript (**Fig. 1C** and **D**). Neither the CRISPRi empty vector

(EV) nor the parental e2 control strains yielded changes in DnaA levels when IPTG was added,

demonstrating that we could specifically and effectively alter cellular DnaA concentrations.

Proper DnaA expression is essential for borrelial growth. Given its essential nature

for chromosomal replication, we first sought to evaluate the consequence of DnaA dysregulation

on *B. burgdorferi* growth and cell division. To achieve this, all strains were grown to mid-

exponential phase and passaged into fresh media with 0.5 mM of IPTG at a density of $1x10^5$

cells/mL. Bacterial numbers of each culture were counted every day for seven days to generate

growth curves.

CRISPRi knockdown of DnaA levels substantially reduced the generation time and

carrying capacity of *B. burgdorferi* (**Fig. 2A** and **B**). The severity of these phenotypes was

consistent with the knockdown efficacy of the two constructs, with $dnaA_{NT1}$ cultures reaching a

maximum density that was one log lower and $dnaA_{T1}$ cultures maxing out two logs lower than

the controls. This suggests that reducing the cellular concentration of DnaA directly limits the

number of division cycles, which is consistent with DnaA's role as the chromosomal replication

initiator.

Overproducing DnaA slowed growth during the first four days post-inoculation (dpi),

growing at a rate similar to the $dnaA_{T1}$ culture (**Fig. 2A** and **B,** early). After that time, however,

cultures consistently resumed logarithmic growth, yielding generation times that were

indistinguishable from the parental strain (**Fig. 2B**, late). We hypothesized this abrupt

resumption of growth was due to mutations in the inducible *dnaA* plasmid that eliminated

overexpression. To test this, we passaged spirochetes that had resumed growth into fresh media

with IPTG to see if growth would still be perturbed. Consistent with our hypothesis, these

"conditioned" bacteria immediately grew at rates that were similar to the control strain (**Fig. 2C**),

and induction of the 3xFLAG tagged DnaA did not occur (**Fig. 2D**). To further test our

hypothesis, we extracted the DnaA overexpression plasmid from the *B. burgdorferi* cultures and

transformed them into *E. coli*. We sequenced purified plasmids from two colonies derived from

the same induced borrelial culture. Each clone had unique mutations within the *dnaA* ORF that

truncated the full-length protein (**Fig. 2E**). Notably, both mutations resulted in the complete loss

of domain IV, the DNA-binding domain of DnaA. Plasmid isolated from transformed *B.*

burgdorferi, which had never been exposed to IPTG, had no mutations within the inducible

dnaA. These escape mutants, taken with the growth dynamics, highlight the stressful nature of

dnaA overexpression, which has consistently been observed in other bacterial organisms (21-23).

Unlike the overexpression strain, we never observed a resumption of growth in the

induced *dnaA* CRISPRi strains. Furthermore, normal growth was attainable from knocked-down

cultures after passaging spirochetes into fresh media without IPTG. These collective data

demonstrate that precisely controlled levels of DnaA are required for optimal borrelial growth.

DnaA affects spirochete cell division/elongation. Chromosomal replication is

intimately tied to cellular elongation and division (24-26). Having found that the conditional

DnaA mutants had replication defects, we also sought to assess the impact of DnaA

dysregulation on *B. burgdorferi* cell length. Over seven days, we imaged and measured

spirochetes. In the parental e2 cultures, we observed a consistent pattern wherein median cell

length peaked during early exponential phase, followed by a steady decrease, then stabilization

once the cultures reached stationary phase (**Fig. 3A**, **3E,** and **3I**).

These findings sharply contrast with what we observed in the CRISPRi knockdown

strains, where the spirochetes steadily maintained median lengths of 30-40 μm (**Fig. 3B-C**, **3F**,

3H, and **3I**); the parental e2 strain had median lengths that spanned from 24-32 μm. The ranges

(maximum-minimum) of the cell lengths within these knockdown strains (147 µm, $dnaA_{T1}$; 134

µm, $dnaA_{NT1}$) were greater than that of the overexpression (77 µm) and parental (101 µm)

bacteria.

149   When DnaA was overexpressed, the median cell length decreased by 2 days post-

induction (**Fig. 3D**, **3H**, and **3K**). As cultures accumulated mutations in the overexpressed *dnaA*,

cell lengths gradually increased to a peak at day 5, then decreased. This oscillation in spirochete

length aligned with the observed growth pattern of this strain, where slowed replication

corresponded to cells of shorter size and rapid replication to longer. This suggests that high

levels of DnaA can diminish borrelial cell length. These results demonstrate that DnaA plays a

role in regulating *B. burgdorferi* cell elongation and/or division and that maintaining proper

DnaA levels is critical to these processes.

157   **Impact of DnaA on ploidy and partitioning.** *B. burgdorferi* is polyploid during

exponential growth *in vitro* (27). As DnaA is the master initiator of chromosomal replication and

affects *B. burgdorferi* morphology, we hypothesized that the ploidy and partitioning of

chromosomes could be altered when DnaA levels are dysregulated. To test this, we transformed

the parental e2 and $dnaA_{T1}$ strains with a construct expressing the chromosomal ParB protein

fused to mCherry (pBSV2G_P_{0826}-mCherry$_{Bb}$-ParB), which has been demonstrated to bind near

oriC (27). We first grew the two strains to mid-exponential phase without IPTG and measured

the length and ParB-*oriC* puncta per cell. The parental bacteria were smaller than those of the

uninduced $dnaA_{T1}$, consistent with our prior observations. Yet, both strains had similar numbers

of ParB-*oriC* (**Fig. 4**). After taking these measurements, the $dnaA_{T1}$ culture was split in half, and

IPTG was added to one of the cultures. The parental, uninduced, and induced strains were then

incubated and examined for three more days.

On the first day post-induction, as the cultures entered stationary phase ($\geq 1 \times 10^8$

cells/mL), all strains experienced a dramatic decline in the total ParB-*oriC* puncta per cell,

consistent with prior observations (27). Over the next two days, the number of these continued to

decrease in the parental and uninduced cultures, whereas the induced culture plateaued and

ended with significantly more ParB-*oriC* puncta. Furthermore, the induced cell lengths, while

longer than the parental, were the same as the uninduced cells, indicating the difference in puncta

wasn't simply due to differences in cell length. With this, we concluded that depleting DnaA

impairs the completion of borrelial chromosomal replication and cell division.

In addition to differences in the number of ParB-*oriC* puncta in the *dnaA*-conditional

mutant, we also noted an apparent difference in their spacing (**Fig. 5A-B**) relative to the parental

strain (**Fig. 5C**). Some cells had wide gaps between the foci, while others had foci that were

close together. We assessed the cultures two- and three-days post-induction and found about

23.7% of parental e2 cells had irregular puncta spacing. The *dnaA*$_{T1}$ CRISPRi strains, in contrast,

had significantly greater proportions of 53.8% and 55.3% of spirochetes with irregular spacing

without and with added IPTG, respectively (**Fig. 5D**). Thus, knocking down DnaA also perturbs

chromosomal partitioning.

Dysregulation of DnaA disrupts flagellar homeostasis. Depletion of DnaA also

resulted in *B. burgdorferi* cells with aberrant helicity. Specifically, we observed spirochetes with

a complete or partial loss of their characteristic corkscrew shape (**Fig. 6A** and **B**, respectively).

Some of these tube-shaped cells were substantially elongated and showed evidence of

incomplete division (**Fig. 6C** and **Supplemental Video**). About 13% of the spirochetes in the

dnaA$_{T1}$-induced cultures and 8.7% of the uninduced exhibited abnormal morphologies (**Fig. 6D**).

The parental and empty vector strains, independent of IPTG, did not exhibit such elevated

proportions of these cells, indicating that the observed helical abnormalities were due to targeted

silencing of the *dnaA* gene. The high level of abnormal cells in the uninduced *dnaA*$_{T1}$ cultures,

relative to the controls, can again be explained by leaky expression of the *dnaA* sgRNA encoded

in the CRISPRi construct.

The characteristic corkscrew appearance of *B. burgdorferi* is due to the 7-11 flagella

found in the periplasm (28). The motors that direct the movement of the endoflagella are

normally localized to the bacterium's poles (29). The observed impacts on growth and

partitioning suggest coordination between replication and flagellar assembly at the sites of

division. This is not unheard of. In *E. coli*, for example, DnaA is known to be involved in

flagellar regulation (30, 31).

DnaA is a global regulator of borrelial gene expression. DnaA is well known to be a

transcription factor in many other bacteria, with regulons driving specific phenotypes (9). In *B.*

burgdorferi, we previously demonstrated that DnaA controls the expression of the *dnaX-ebfC*

operon (10). Considering this and the dramatic impacts of DnaA dysregulation on the Lyme

spirochete, we queried whether the myriad phenotypes of the conditional mutants are due to

DnaA-dependent transcriptional effects. To address this question, we performed RNA-seq on

three strains with different levels of DnaA enrichment: wild-type (e2), DnaA-up (e2 +

pACK121), and DnaA-down (e2 + *dnaA*$_{T1}$). All the strains had the plasmids cp26, lp17, lp54,

cp32-1, cp32-3, and cp32-4, as assessed by PCR and whole genome sequencing. The bacteria for

RNA-seq analysis were grown to mid-exponential phase, induced with 0.5 mM IPTG, and

incubated overnight. We chose this strategy to mitigate the chance of escape-mutant

development from confounding the results.

For the analyses of the RNA-seq data, comparisons were made between the three groups.

We considered a gene to be differentially expressed if the false discovery rate (FDR) was less

than or equal to 0.05 and had a log$_2$FC (fold change) of ≥ 1 or ≤ -1. With these thresholds, 216

genes were noted as being differentially expressed between the wild-type and DnaA-down

cultures (72 down, 144 up; **Fig. 7A**), 84 genes between WT and DnaA-up (40 down, 44 up; **Fig.**

7B), and 259 between DnaA-down and DnaA-up (105 down, 154 up; **Fig. 7C**). Principle

component analysis showed that the replicates of each strain clustered together (**Fig. 7D**). Most

of the genes that were differentially expressed are encoded on the chromosome: DnaA-down vs.

e2: 69.9%; DnaA-up vs. e2: 72.6% (**Fig. 7E**). Of the plasmids, the linear lp54 replicon had the

most impacted genes for the DnaA-down vs. e2 comparison (8.3%), while the circular cp32-1

had the most for the DnaA-up vs e2 comparison (7.1%; **Fig. 7E**).

As expected, the RNA-seq showed that compared to the e2 parent, *dnaA* transcript was

about 2.8-fold less abundant in the CRISPRi strain and 15-fold more abundant in the

overexpression strain. Typically, we normalize transcript levels to *ftsK*, a gene previously

observed to have stable expression across growth phases (32). RNA-seq showed that *ftsK* was

impacted by *dnaA* dysregulation (discussed below). Given this, we performed validations

normalizing to *rpoB*, which did not change under the tested conditions. Using this approach,

dnaA transcript was 80-fold less abundant in the knockdown vs. the WT and 16-fold more

abundant in the overexpression vs. the WT (**Fig. 7E**). DnaA protein levels were about half as

abundant in the knockdown and 13 times more in overexpression than the WT (**Fig. 10B**).

Transcript levels of many genes were most affected when *dnaA* levels were reduced. We

initially hypothesized that transcriptomic profiles between the conditional mutants relative to the

parental strain would reciprocally mirror each other. Only one gene fit this hypothesis, BB_0413

(2.2-fold increase for $dnaA_{T1}$/e2; 2.3-fold decrease for pACK121/e2). Indeed, in many cases,

transcript levels for a gene would experience the same degree of change regardless of whether

DnaA was increased or decreased. Such instances suggested that some of the observed changes

in gene expression were not due to DnaA itself but rather a shared cellular response to the

changes in DnaA levels. Despite some of these complications, the RNA-seq data set offered

insights that could explain the phenotypes observed in the conditional *dnaA*-mutant strains.

The earliest interrogations into borrelial DnaA focused on its function as a transcription

factor (10, 14). Consistent with our prior *in vitro* studies, knocking down *dnaA* significantly

decreased transcription of the *dnaX-ebfC* operon (10). This operon codes for the τ-subunit of the

DNA polymerase III holoenzyme (DnaX) and the nucleoid-associated protein, EbfC. Other

impacted replication genes included the DNA polymerase III β-clamp (*dnaN*), DNA polymerase

I (*polA*), DNA topoisomerase IV (*parEC*), and telomere resolvase (*resT*). Except for *dnaN*, these

genes were downregulated when *dnaA* levels were decreased. The upregulation of *dnaN*, seen

when comparing DnaA-Up to DnaA-down conditions, suggests that DnaA represses the

expression of this gene. This possibility is unsurprising given that the *oriC* is directly 5' of the

dnaN ORF, and we have shown that DnaA binds this region (10).

In the *dnaA*-knockdown *B. burgdorferi*, we observed cells that were considerably

elongated. In our evaluation of the borrelial genome, we identified 22 homologs of genes that

encode components of either the bacterial elongasome or divisome (**Table 1**). Of these, 39.1%

were differentially expressed in the $dnaA_{T1}$ CRISPRi strain (**Fig. 8A**). Among these were *mreB*,

mreC, *mreD,* and *mrdA*, which constitute an apparent operon and encode core components of the

bacterial elongasome (**Fig. 8B**). MreB is an actin-like cytoskeleton protein that polymerizes

across the bacterial inner membrane to regulate the spatiotemporal peptidoglycan synthesis of

growing cells (33-35). Knockdown of MreB in *B. burgdorferi* results in bulging and cell

widening at the division sites (19). MreC and MreD are membrane-embedded proteins that

interact and regulate the peptidoglycan crosslinking activity of PBP2 (penicillin-binding protein

2; MrdA) (36). The peptidoglycan polymerase RodA and PBP1a interact with PBP2, with the

former stimulating its activity. Except for *mreD*, the genes in the apparent *mre* operon were

downregulated in *dnaA*-deficient *B. burgdorferi*. A DNA sequence that is the same as those

found in the *oriC*, which may constitute the borrelial DnaA-box, is located upstream of the *mreB*

ORF, suggesting DnaA may directly regulate the expression of these genes. The decreased

transcript levels for *mreB*, *mreC*, and *mrdA* would suggest decreased elongation, which is the

opposite of what we observed (**Fig. 3**). This apparent contradiction suggests that filamentation

may be due to defects in cell division.

The transcripts of four important divisome proteins were impacted when *dnaA* was

knocked down: FtsA, FtsK, FtsE, and FtsX (**Fig. 8A** and **B**). FtsA is the membrane anchor

protein for the septation marker protein FtsZ (37, 38), and its transcript increased about 2-fold in

the conditional *dnaA* mutant. Surprisingly, DnaA-overexpressed *B. burgdorferi* also experienced

a 2-fold increase in *ftsA*. Transcripts for FtsK, the DNA translocase, declined 2-fold. FtsE and

FtsX, encoded next to each other, had their transcripts decreased 2-fold and increased almost 3-

fold, respectively. Interestingly, a second allele of *ftsX* (*ftsX**), which is truncated, is located

directly upstream of *ftsEX,* and transcript levels of this gene were substantially affected, with a

189-fold increase. Like the *mre* locus, DnaA could regulate these genes directly as potential

DnaA-boxes are adjacent to these loci.

In the conditional *dnaA*-knockdown *B. burgdorferi,* we observed spirochetes that had lost

their helical structure and hypothesized these cells had disrupted flagella. The *B. burgdorferi*

genome encodes nearly three dozen flagellar genes (**Fig. 9A**). When *dnaA* was knocked down,

six of these genes were significantly impacted: *fliQ*, *fliR*, *flgD*, *flaA*, *flgV*, and *flgA* (**Fig. 9B**). All

these genes except for *flaA* and *flgA* are encoded in the *flgB* superoperon. FliQ and FliR are

components of the flagellar export apparatus (39). FlgD is a scaffolding protein that is required

for flagellar hook formation (40, 41). FlaA is the minor borrelial flagellar filament protein (42,

43). FlgV is a protein first characterized as a homolog of the RNA chaperone Hfq, but recent

work has demonstrated the protein localizes with flagellar motors and modulates their assembly

(44, 45). FlgA is a chaperone involved in the formation of the P-ring of the flagellar motor (46).

The transcripts for *fliR* and *flgD* increased the most of these genes, about 15 and 4-fold,

respectively, when DnaA levels were reduced. The changes in the expression of these genes

could alter flagellar assembly homeostasis and explain the observed abnormalities in helicity in

the *dnaA*-deficient spirochetes. It is also possible that defects in septation disrupted the

localization and assembly of flagella.

In addition to affecting genes for essential cellular processes, DnaA significantly

impacted the expression of outer surface proteins that are involved in vertebrate infection and

virulence, such as ErpA, decorin binding protein DbpA, and OspC (**Fig. 7A-C** and **Fig. 10E-F**).

The Erps are multifunctional adhesins that bind vertebrate host factors such as laminin,

plasminogen, and complement proteins (47-52). Transcripts for *erpA* increased in the *dnaA*

knockdown and overexpression conditions (**Fig. 10E-F**). Interestingly, this increase in transcript

didn't correspond to increased ErpA protein (**Fig. 10B-D**).

The *erp* genes are encoded on the cp32 plasmids and have conserved operators that allow

uniform regulation (53). The Erps are turned on when the tick begins to feed, a time of rapid

borrelial replication, and remain on during vertebrate infection (54). Three proteins bind the *erp*

operators and regulate their expression: BpaB (<u>B</u>orrelial cp32 <u>ParB</u> analog), BpuR, and EbfC.

BpaB represses *erp* transcription along with the co-repressor BpuR (**Fig. 10A**) (55-57). EbfC is

the antirepressor that antagonizes BpaB to allow for *erp* transcription (56). DnaA was previously

hypothesized to regulate *erp* transcription by activating EbfC and repressing BpuR (10, 13, 14).

EbfC transcript and protein levels decreased when *dnaA* was knocked down, consistent with that

model (**Fig. 10B** and **E**). BpuR transcript, however, did not significantly change, but its protein

levels appeared to decline with a decrease in cellular DnaA concentrations (**Fig. 10B** and **E**).

BpuR alone does not repress *erp* expression, so this cannot account for the increase in ErpA

transcript. Thus, we looked at the protein levels of the *erp* repressor. Despite the lack of a

change in the transcript, immunoblots showed a decrease in BpaB protein levels in *dnaA*

CRISPRi *B. burgdorferi*, which can explain the increase in *erpA* transcript (**Fig. 10B, C,** and **E**).

BpaB levels were similar to those of the parental e2 strain during *dnaA* overexpression, and

ErpA transcript and protein were unaffected, likely due to normal levels of the antirepressor

EbfC. Overall, these results demonstrate that DnaA not only affects the expression of genes

involved in basic cellular processes but also those involved in maintaining the Lyme spirochete's

infection cycle.

DISCUSSION

In this study, we sought to understand the roles of DnaA in controlling the physiology of

B. burgdorferi. To do this, we utilized inducible CRISPRi and overexpression vectors.

Overexpressing DnaA was toxic to the bacteria and was consistently overcome by mutations in

the inducible *dnaA*. This, along with the difference in DnaA levels from overexpression vs.

knockdown, 13-fold DnaA increase vs. 2-fold decrease, likely explains some of the results we

encountered. Nevertheless, increasing DnaA levels yielded significantly different phenotypes,

such as spirochete size. CRISPRi was the more reliable and effective means of producing

conditional *dnaA* mutants in *B. burgdorferi*. This approach targeting *dnaA* has been successfully

described in *E. coli*, *Lactobacillus plantarum,* and *Streptococcus pyogenes* (58-60). An attempt

was made in *Pseudomonas putida,* but clones couldn't be recovered (61).

We found that lowering the cellular levels of DnaA profoundly reduced cell division,

consistent with the protein's function as the chromosomal replication initiator. The reduction of

DnaA also coincided with decreased expression of many essential replication genes. It is known

that one of these genes, *dnaX*, is directly regulated by DnaA (10). This suggests that DnaA may

directly regulate those other genes, or *B. burgdorferi* may have mechanisms to sense replication

initiation and coordinate gene expression accordingly.

DnaA-deficient spirochetes also increased in length. The bacterial cell cycle is typically

divided into three phases: B, C, and D (62). The B and D phases correspond to the birth and

division of the bacteria, respectively. The C phase corresponds to everything in between, namely,

chromosomal replication, along with elongation/growth. Replication is initiated by DnaA and

then carried out by the replisome. Elongation is facilitated by the elongasome, which synthesizes

peptidoglycan to allow for cell growth. The DnaA-depleted spirochetes, although considerably

longer than the parental strain, had overall decreased expression of elongasome genes, the *mre*

locus in particular. This suggested a division issue, but we cannot rule out the possibility that

DnaA is required for cross-talk between replication and cell growth. Indeed, it is likely that

replication, elongation, and division are all interconnected. Lyme and relapsing *Borreliae*

elongate at three distinct zones along their length: 1/4, 1/2, and 3/4 (63). This localization pattern

at the mid-cell suggests an interplay between elongation and division machinery. How or if

DnaA directly plays into this delicate balance remains to be assessed.

Many bacterial cell division proteins are denoted "Fts" for their filamentous phenotypes

caused by temperature-sensitive mutations (64). Knocking down DnaA affected the expression

of *ftsA*, *ftsEX*, and *ftsK*. The dysregulation of these genes potentially explains both the elongated

phenotype and the irregular spacing of the ParB-*oriC* puncta.

FtsA localizes to the inner leaflet of the inner membrane and binds FtsZ. Increased FtsA,

seen when DnaA was knocked down, could decrease Z-ring assembly and, thus, cytokinesis. In

E. coli, overexpression of FtsA causes cells to filament (65). Overexpressed DnaA spirochetes

had elevated *ftsA* transcript, but no changes were observed in *ftsEX*. The FtsEX complex

regulates the activity of peptidoglycan hydrolases and FtsA (66-69). Disruption of FtsE and

FtsX levels could, therefore, prevent the breakdown of peptidoglycan and Z-ring formation at the

site of division. The combined perturbation of *ftsA* and *ftsEX* expression in the *dnaA* CRISPRi

strain potentially explains the observed elongated phenotype.

In addition to changes in spirochete length in the DnaA-depleted *B. burgdorferi*, we

observed abnormal spacing of ParB-*oriC* puncta. These bacteria had decreased transcripts of the

membrane-embedded DNA translocase FtsK, a core component of the divisome (70-72). FtsK,

through its C-terminal αβγ-domain, resolves chromosome dimers and translocates DNA from the

septum to allow for cytokinesis (73-76). A decrease in FtsK could thus result in the failure to

separate sister chromosomes, leading to cells with abnormal ploidy. This could also explain why

some *oriC* sites were located close together in the *dnaA* knockdown strain (**Fig. 5**). FtsK also

recruits downstream divisome proteins through its N-terminal transmembrane domain (71, 77-

79). Therefore, reducing FtsK may also limit the recruitment of divisome proteins to the septa.

Motility is perhaps the most important "virulence" factor that *B. burgdorferi* employs to

infect and colonize its vertebrate hosts. By knocking down DnaA in the Lyme spirochete, we

observed a substantial impact on the bacteria's helicity. This corkscrew morphology is due to

periplasmic flagella (28). Key genes, such as FlgD and FliR, were overexpressed when levels of

DnaA were lowered. FliR is a component of the flagellar motor's export apparatus, which

transports flagellar filament substrates from the cytoplasm to the periplasm (39). Disrupting the

balance of this subunit would very likely impair the overall function and structure of the flagellar

motor and filament. By increasing FliR, it is possible that the export of the flagellar cargo to the

periplasmic space would be affected, which in turn would prevent flagella formation. The

flagellar motors are normally localized to the poles and developing septa of *B. burgdorferi*. This

suggests that replication, elongation, or division processes may coordinate the formation of

flagellar motors at the dividing mid-cell. In *E. coli*, DnaA regulates genes involved in flagellar

assembly (30, 31, 80). Whether DnaA in *B. burgdorferi* acts through a similar mechanism is

unknown.

In addition to the many morphological changes incurred by dysregulation of DnaA, there

were also significant impacts on the expression of vertebrate infection-related outer surface

proteins. Many of these antigenic proteins, such as OspC, DbpA, and the Erps are turned on

during the tick blood meal, a period defined by rapid replication. Relative to these genes, levels

of transcript for the alternative sigma factor RpoS were not significantly changed by

manipulation of DnaA levels. We previously demonstrated that DnaA regulates the *erp*

antirepressor EbfC and may regulate the *erp* co-repressor BpuR (10, 14). With our CRISPRi

approach, we were able to validate our prior data on *dnaX-ebfC* regulation. While we could not

detect any substantial differences in BpuR with DnaA manipulation, this might have been a

consequence of assessing mid-exponential spirochetes grown at 34 ºC; BpuR expression is

highest when grown at 23 ºC (14, 57). When DnaA levels were reduced, expression of ErpA

increased. We suggest this could be due to decreased levels of *erp* repressor BpaB. While this

was unexpected, it is logical that a partitioning protein be decreased when *dnaA* transcription is

knocked down. This suggests that these extrachromosomal DNAs can sense changes in

chromosomal replication. How this is mediated remains to be investigated.

We observed gene expression level changes in a vast number of regulatory networks.

This led us to examine if *dnaA* knockdown or overexpression impacted any known regulatory

factors. Among the known networks, we observed modest changes in gene expression levels of

the following regulators: Hk1, Rrp1, and SpoVG. These changes did not meet the log fold

change threshold but were significant (FDR $\leq$ 0.05) and had a log fold change of approximately

1.5. When comparing the $dnaA_{T1}$ knockdown strain to the parental e2 strain, these genes were

downregulated by a fold change of 1.52, 1.61, and 1.53, respectively. While not meeting our set

threshold, it is possible the combined modest expression changes in these regulators considerably

impacted the expression levels of the genes they modulate. For example, the nucleic acid binding

protein SpoVG, the response regulator Hk1, and diguanylate cyclase Rrp1 are all known to

regulate the glycerol metabolism (*glpFKD*) operon, which is essential for *B. burgdorferi*

colonization and persistence in ticks (81-85). The downregulation of the *glpFKD* operon (~2.6 to

4-fold) is likely attributable to the decreased expression of *spoVG*, *hk1*, and *rrp1*. These

observations link DnaA and/or DNA replication to crucial borrelial networks, including the c-di-

GMP regulatory network (Hk1/Rrp1) critical to completing the *B. burgdorferi* enzootic life

cycle.

Another regulatory network impacted was the Hk2/Rrp2 pathway, whose expression

levels were increased by 8.2 and 1.5, respectively. The Hk2/Rrp2 two-component system is a

known activator of the RpoN/RpoS alternative sigma factor cascade, which regulates essential

virulence factors like OspC (86-88). Deletion of *rrp2* is lethal, and investigations into *rrp2*

regulation have relied on conditional mutants (18). Although neither *rpoN* nor *rpoS* levels

changed significantly, we note that *rpoS* transcription can be activated by the housekeeping

sigma factor RpoD in addition to RpoN (89). This further provides strong evidence that the

Hk2/Rrp2 pathway has impacts outside of *rpoS* regulation potentially mediated by proper *dnaA*

levels and/or associated with DNA replication. In addition, this data supports our past data

showing that *ospC* regulation, through the repressor Gac, can occur independently of RpoS

regulation (90). Additional data from our lab indicates that DnaA is a positive activator of OspC

(unpublished results). This supports the RNA-seq data in this study, showing that knockdown of

DnaA levels reduces *ospC* transcript.

Organisms have evolved complex and varied molecular circuits to regulate cellular

homeostasis in ever-changing environments. Bacterial pathogens, especially, have developed

such networks in their tussle against host defenses. During the tick blood meal, *B. burgdorferi*

must take advantage of the nutrients to propagate its numbers, reprogram its transcriptome, and

disseminate to the vertebrate. Failure to do so means death, as *B. burgdorferi* cannot be

transmitted vertically to tick offspring (91). In *B. burgdorferi,* DnaA, the master regulator of

replication initiation, appears to form a complex regulatory network to coordinate these essential

processes (**Fig. 11**). Future studies need to be conducted to determine which of the DnaA-

dysregulated effects described here are due to DnaA itself or a consequence of a lack of initiation

of chromosomal replication.

METHODS AND MATERIALS.

Bacterial strains, plasmids, and genetic manipulations. Studies were conducted using

the readily transformable *B. burgdorferi* clonal strain B31-e2 (92). Spirochetes were grown in

liquid Barbour-Stoener-Kelly II (BSK-II) medium supplemented with 6% rabbit serum (v/v) at

35 ºC. For all experiments, *B. burgdorferi* was diluted from mid-exponential phase (3-5 x 10^7

cells/mL) into fresh BSK-II medium. Cultures were only used if they were at least two passages

away and no more than three from the initial -80ºC stock. Borrelial culture densities were

enumerated using a Petroff-Hauser counting chamber and dark-field microscopy.

The *dnaA* overexpression (pACK121) and CRISPRi plasmids (pACK136, *dnaA*$_{T1}$;

pACK138, *dnaA*$_{NT1}$) were generated by GenScript. The pACK121 plasmid is a derivative of

pJSB268 (17). The plasmid was designed to replace the *luc* gene with the *B. burgdorferi dnaA*

gene containing an N-terminal 3xFLAG epitope. The pACK136 (*dnaA*$_{T1}$) and pACK137

(*dnaA*$_{NT1}$) plasmids are derivatives of pJJW101 (20). These plasmids were created by inserting

the desired sgRNA sequence between the BsaI sequences. The sgRNA targets used in this study

were chosen using CRISPy-web (https://crispy.secondarymetabolites.org) and B31 chromosome

contig (NC_001318). Whole-plasmid sequencing was performed by Plasmidsaurus using Oxford

Nanopore Technology with custom analysis and annotation.

Growth curve analysis. Growth curves for each strain were generated from three

independent experiments. Cultures were seeded at the same initial density (1 or 5 x 10^5 cells/mL)

and enumerated over seven days. Generation times were calculated in Microsoft Excel by fitting

an exponential curve to the data points denoting the beginning and end of logarithmic growth.

Statistically significant differences were determined by one-way ANOVA.

Immunoblot analyses. Polyclonal DnaA antibodies were generated by Thermo Scientific

in rabbits using recombinant GST-DnaA generated in-house (10). The serum from the final bleed

was purified by affinity purification using MBP-DnaA on an amylose column. Murine

monoclonal anti-FlaB antibodies were used to assess the even loading of SDS-PAGE gels (93).

Other primary antibodies were generated previously (56, 57, 94). Cells for immunoblot were

pelleted and washed three times with 1x PBS, pH 7.4. Approximately 10^7 spirochetes were

loaded per lane for SDS-PAGE. Goat anti-Rabbit Alexa Fluor 488 (1:10, 000, Thermo Scientific)

and Goat anti-Mouse IRDye800 (1:5,000, Licor) were used as secondary antibodies.

Densitometric analyses were performed using ImageLab software (BioRad). The intensities of

the bands of interest were normalized to the corresponding loading control.

Quantitative reverse transcription-PCR (qRT-PCR). Mid-exponential phase (3-5 x

10^7 cells/mL) bacteria were washed three times with PBS (pH 7.4) before RNA extraction using

Qiagen Mini RNA kits. Genomic DNA was cleared by on-column DNase treatment. RNA was

reverse transcribed using iScript cDNA synthesis kits (Bio-Rad). Quantitative PCR was

conducted using iTaq Universal SYBR Green Supermix (Bio-Rad) and a Bio-Rad CFX96 Touch

Real-time PCR thermocycler. The primer sets used for this study were designed using the IDT

Primer Quest Tool (https://www.idtdna.com/PrimerQuest/Home/Index). Cq values were

normalized to *ftsK* or *rpoB* (ΔCq) and then to the parental strain or uninduced control (ΔΔCq).

Fold expression was determined using the function $2^{-\Delta\Delta Cq}$. If $\Delta\Delta Cq > 0$, then the function $-2^{\Delta\Delta Cq}$

was used.

Cell length analysis. Wet mounts of live bacteria were visualized by dark-field with an

Olympus Bx51 microscope at 400x total magnification. Micrographs were taken using a C-

mounted Accu-scope Excelis HD camera. Cell lengths were determined using Captavision+

software. A conversion factor of 13.80 pixels/μm was used for all measurements. For each

replicate and time point, approximately 50 spirochetes were measured.

Fluorescence microscopy and image analysis. Spirochetes were pelleted by

centrifugation, washed twice with PBS (pH 7.4), and diluted to a final concentration of 1×10^5

cells/mL. Ten microliters of cells were mounted on a glass slide. Cells were visualized using an

Olympus Bx51 microscope with a Cool LED p-E300 illumination system. Micrographs of dark-

field and red fluorescence were captured for each field of view using a C-mounted Accu-scope

Excelis HD camera. Micrographs were merged in ImageJ, and the background was inverted

using EZreverse (https://amsterdamstudygroup.shinyapps.io/ezreverse/) with the HSL color

space option (95).

Whole Genome Sequencing (WGS) and analysis. The parental *B. burgdorferi* B31-e2,

dnaA$_{T1}$ CRISPRi, and *dnaA*-overexpression strains were sequenced to verify plasmid content and

determine if mutations occurred in the *dnaA* sequence of the generated strains. Briefly, the

strains were grown in duplicate to mid-exponential phase ($3\text{-}5 \times 10^7$ cells/mL). Two mLs of cells

were pelleted and washed twice with PBS (pH 7.4) and frozen at -80 ºC. DNA extraction and

whole genome sequencing were performed by SeqCenter. Briefly, the cells were sent to the

SeqCenter facility. DNA was extracted from the pellets using the ZymoBIOMICS^{TM} DNA

Miniprep Kit according to the manufacturer's protocol. DNA concentrations were determined by

Qubit.

Sequencing libraries were prepared via the tagmentation-based and PCR-based Illumina

DNA Prep kit and custom IDT 10 bp unique dual indices (UDI) with a target insert size of 320

bp. The sequencing was performed either in one or more multiplexed flow-cell runs on an

Illumina NovaSeq 6000 sequencer resulting in 2x151 bp paired-end reads. Quality control steps

including demultiplexing and adapter trimming were performed with bcl-convert (v4.1.5). Reads

were aligned to the *B. burgdorferi* B31 genome assembly (GCF_000008685.2_ASM868v2).

Variant calling was performed using BreSeq (v. 0.38.1) under default settings.

RNA sequencing (RNA-seq) and analysis. The same cultures of the parental *B.*

burgdorferi B31-e2, *dnaA*$_{T1}$ CRISPRi, and *dnaA*-overexpression strains used for WGS were used

for RNA-seq. All strains were grown in duplicate to mid-exponential growth phase (3-5 x 10^7

cells/mL) and induced with 0.5 mM IPTG overnight at 35 ºC. Nine mLs of the cells were

pelleted and washed two times with PBS (pH 7.4) and frozen at -80 ºC. The cells were sent to

SeqCenter for RNA extraction and sequencing using their intermediate RNA analysis with

replicates (Prokaryotic) service. Briefly, their method consisted of quality control and adapter

trimming by bclconvert (version 4.1.5) and mapping reads using HISAT2 (version 2.2.0) to the

RefSeq version of the *B. burgdorferi* B31 genome assembly (GCF_000008685.2_ASM868v2).

The read counts were uploaded into R (version 4.0.2) and normalized using the edgeR Trimmed

Mean of M values (TMM) algorithm (1.14.5). Differential gene expression analysis was

performed with edgeR using the normalized TMM values. Only plasmids that were confirmed to

be present were included in the analyses. We further filtered the data set by setting the thresholds

for differential expression at $\log_2 FC \geq 1$ or ≤ -1 and an FDR ≤ 0.05. Differentially expressed

genes (DEGs) were visualized via volcano plots made using VolcaNoseR (96). DEGs belonging

to the elongasome and divisome were visualized by heatmap. The heatmap was generated in R (v

4.4.1) using the pheatmap package, with the log2 transformed normalized counts per million

(CPMs) (97). The clustering method used for the heatmap was the default method of complete

linkage, while the distance measure used was correlation.

To identify flagellar, cell division, and elongation homologs, the genome was directly

searched for annotations, or the sequences of characterized proteins from other bacteria were

aligned by BLAST against the *B. burgdorferi* genome.

536   Acknowledgments

537   This research was funded by NIH grant R21 AI147139. We thank Dr. Wolfram Zückert and

538   Bryan Murphy for their help designing and troubleshooting the CRISPRi shuttle vectors and Dr.

539   Jon Blevins for providing the pJSB268 shuttle vector. BioRender was used to make the

540   schematics in Figures 1, 2, 8, 9, and 10.

Works Cited.

1.   Radolf JD, Caimano MJ, Stevenson B, Hu LT. 2012. Of ticks, mice and men:

     understanding the dual-host lifestyle of Lyme disease spirochaetes. Nat Rev Microbiol

     10:87-99.

2.   Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JW, Li X, Mead PS. 2016.

     Lyme borreliosis. Nat Rev Dis Primers 2:16090.

3.   De Silva AM, Fikrig E. 1995. Growth and migration of *Borrelia burgdorferi* in *Ixodes*

     ticks during blood feeding. Am J Trop Med Hyg 53:397-404.

4.   Piesman J, Oliver JR, Sinsky RJ. 1990. Growth kinetics of the Lyme disease spirochete

     (*Borrelia burgdorferi*) in vector ticks (*Ixodes dammini*). Am J Trop Med Hyg 42:352-7.

5.   Speck C, Messer W. 2001. Mechanism of origin unwinding: sequential binding of DnaA

     to double- and single-stranded DNA. EMBO J 20:1469-76.

6.   Kaur G, Vora MP, Czerwonka CA, Rozgaja TA, Grimwade JE, Leonard AC. 2014.

     Building the bacterial orisome: high-affinity DnaA recognition plays a role in setting the

     conformation of *oriC* DNA. Mol Microbiol 91:1148-63.

7.   Ozaki S, Katayama T. 2012. Highly organized DnaA-*oriC* complexes recruit the single-

     stranded DNA for replication initiation. Nucleic Acids Res 40:1648-65.

8.   Messer W, Weigel C. 1997. DnaA initiator--also a transcription factor. Mol Microbiol

     24:1-6.

9.   Menikpurage IP, Woo K, Mera PE. 2021. Transcriptional Activity of the Bacterial

     Replication Initiator DnaA. Front Microbiol 12:662317.

10.   Krusenstjerna AC, Saylor TC, Arnold WK, Tucker JS, Stevenson B. 2023. *Borrelia*

burgdorferi DnaA and the Nucleoid-Associated Protein EbfC Coordinate Expression of

the *dnaX-ebfC* Operon. J Bacteriol 205:e0039622.

11.   Medrano MS, Ding Y, Wang X-G, Lu P, Coburn J, Hu LT. 2007. Regulators of

expression of the oligopeptide permease A proteins of *Borrelia burgdorferi*. Journal of

bacteriology 189:2653-2659.

12.   Jutras BL, Bowman A, Brissette CA, Adams CA, Verma A, Chenail AM, Stevenson B.

2012. EbfC (YbaB) is a new type of bacterial nucleoid-associated protein and a global

regulator of gene expression in the Lyme disease spirochete. J Bacteriol 194:3395-406.

13.   Stevenson B, Krusenstjerna AC, Castro-Padovani TN, Savage CR, Jutras BL, Saylor TC.

2022. The Consistent Tick-Vertebrate Infectious Cycle of the Lyme Disease Spirochete

Enables Borrelia burgdorferi To Control Protein Expression by Monitoring Its

Physiological Status. J Bacteriol 204:e0060621.

14.   Jutras BL, Savage CR, Arnold WK, Lethbridge KG, Carroll DW, Tilly K, Bestor A, Zhu

H, Seshu J, Zuckert WR, Stewart PE, Rosa PA, Brissette CA, Stevenson B. 2019. The

Lyme disease spirochete's BpuR DNA/RNA-binding protein is differentially expressed

during the mammal-tick infectious cycle, which affects translation of the SodA

superoxide dismutase. Mol Microbiol 112:973-991.

15.   Picardeau M, Lobry JR, Hinnebusch BJ. 1999. Physical mapping of an origin of

bidirectional replication at the centre of the *Borrelia burgdorferi* linear chromosome. Mol

Microbiol 32:437-45.

16.   Hirota Y, Ryter A, Jacob F. 1968. Thermosensitive mutants of *E. coli* affected in the

processes of DNA synthesis and cellular division. Cold Spring Harb Symp Quant Biol

33:677-93.

17.   Blevins JS, Revel AT, Smith AH, Bachlani GN, Norgard MV. 2007. Adaptation of a

luciferase gene reporter and *lac* expression system to *Borrelia burgdorferi*. Appl Environ

Microbiol 73:1501-13.

18.   Groshong AM, Gibbons NE, Yang XF, Blevins JS. 2012. Rrp2, a prokaryotic enhancer-

like binding protein, is essential for viability of *Borrelia burgdorferi*. J Bacteriol

194:3336-42.

19.   Takacs CN, Scott M, Chang Y, Kloos ZA, Irnov I, Rosa PA, Liu J, Jacobs-Wagner C.

2020. A CRISPR interference platform for selective downregulation of gene expression

in *Borrelia burgdorferi*. Appl Environ Microbiol doi:10.1128/AEM.02519-20.

20.   Murphy BT, Wiepen JJ, He H, Pramanik AS, Peters JM, Stevenson B, Zuckert WR.

2023. Inducible CRISPRi-Based Operon Silencing and Selective in Trans Gene

Complementation in *Borrelia burgdorferi*. J Bacteriol 205:e0046822.

21.   Zheng W, Li Z, Skarstad K, Crooke E. 2001. Mutations in DnaA protein suppress the

growth arrest of acidic phospholipid-deficient *Escherichia coli* cells. EMBO J 20:1164-

72.

22.   Jonas K, Chen YE, Laub MT. 2011. Modularity of the bacterial cell cycle enables

independent spatial and temporal control of DNA replication. Curr Biol 21:1092-101.

23.   Anderson ME, Smith JL, Grossman AD. 2022. Multiple mechanisms for overcoming

lethal over-initiation of DNA replication. Mol Microbiol 118:426-442.

24.   Cooper S, Helmstetter CE. 1968. Chromosome replication and the division cycle of

606   *Escherichia coli* Br. Journal of Molecular Biology 31:519-540.

607   25.   Donachie WD. 1968. Relationship between Cell Size and Time of Initiation of DNA

608   Replication. Nature 219:1077-1079.

609   26.   Wallden M, Fange D, Lundius EG, Baltekin Ö, Elf J. 2016. The Synchronization of

610   Replication and Division Cycles in Individual *E. coli* Cells. Cell 166:729-739.

611   27.   Takacs CN, Wachter J, Xiang Y, Ren Z, Karaboja X, Scott M, Stoner MR, Irnov I,

612   Jannetty N, Rosa PA, Wang X, Jacobs-Wagner C. 2022. Polyploidy, regular patterning of

613   genome copies, and unusual control of DNA partitioning in the Lyme disease spirochete.

614   Nat Commun 13:7173.

615   28.   Motaleb MA, Corum L, Bono JL, Elias AF, Rosa P, Samuels DS, Charon NW. 2000.

616   *Borrelia burgdorferi* periplasmic flagella have both skeletal and motility functions. Proc

617   Natl Acad Sci U S A 97:10899-904.

618   29.   Li C, Bakker RG, Motaleb MA, Sartakova ML, Cabello FC, Charon NW. 2002.

619   Asymmetrical flagellar rotation in *Borrelia burgdorferi* nonchemotactic mutants. Proc

620   Natl Acad Sci U S A 99:6169-74.

621   30.   Wu D, Baigalmaa L, Yao Y, Li G, Su M, Fan L, Morigen. 2021. The *Escherichia coli*

622   QseB/QseC signaling is required for correct timing of replication initiation and cell

623   motility. Gene 773:145374.

624   31.   Mizushima T, Tomura A, Shinpuku T, Miki T, Sekimizu K. 1994. Loss of flagellation in

625   *dnaA* mutants of *Escherichia coli*. J Bacteriol 176:5544-6.

626   32.   Arnold WK, Savage CR, Brissette CA, Seshu J, Livny J, Stevenson B. 2016. RNA-Seq of

627   *Borrelia burgdorferi* in Multiple Phases of Growth Reveals Insights into the Dynamics of

Gene Expression, Transcriptome Architecture, and Noncoding RNAs. PLoS One

11:e0164165.

33.   Jones LJ, Carballido-Lopez R, Errington J. 2001. Control of cell shape in bacteria:

helical, actin-like filaments in *Bacillus subtilis*. Cell 104:913-22.

34.   Kruse T, Bork-Jensen J, Gerdes K. 2005. The morphogenetic MreBCD proteins of

Escherichia coli form an essential membrane-bound complex. Mol Microbiol 55:78-89.

35.   Figge RM, Divakaruni AV, Gober JW. 2004. MreB, the cell shape-determining bacterial

actin homologue, co-ordinates cell wall morphogenesis in *Caulobacter crescentus*. Mol

Microbiol 51:1321-32.

36.   Liu X, Biboy J, Consoli E, Vollmer W, den Blaauwen T. 2021. MreC and MreD balance

the interaction between the elongasome proteins PBP2 and RodA. PLOS Genetics

16:e1009276.

37.   Pichoff S, Lutkenhaus J. 2005. Tethering the Z ring to the membrane through a conserved

membrane targeting sequence in FtsA. Mol Microbiol 55:1722-34.

38.   Pichoff S, Lutkenhaus J. 2002. Unique and overlapping roles for ZipA and FtsA in septal

ring assembly in *Escherichia coli*. Embo j 21:685-93.

39.   Minamino T, Macnab RM. 1999. Components of the *Salmonella* flagellar export

apparatus and classification of export substrates. J Bacteriol 181:1388-94.

40.   Kutsukake K, Doi H. 1994. Nucleotide sequence of the *flgD* gene of *Salmonella*

typhimurium which is essential for flagellar hook formation. Biochim Biophys Acta

1218:443-6.

41.   Ohnishi K, Ohto Y, Aizawa S, Macnab RM, Iino T. 1994. FlgD is a scaffolding protein

needed for flagellar hook assembly in *Salmonella typhimurium*. J Bacteriol 176:2272-81.

42.   Ge Y, Li C, Corum L, Slaughter CA, Charon NW. 1998. Structure and expression of the

652   FlaA periplasmic flagellar protein of *Borrelia burgdorferi*. J Bacteriol 180:2418-25.

653   43.   Motaleb MA, Sal MS, Charon NW. 2004. The decrease in FlaA observed in a *flaB*

654   mutant of Borrelia burgdorferi occurs posttranscriptionally. J Bacteriol 186:3703-11.

655   44.   Lybecker MC, Abel CA, Feig AL, Samuels DS. 2010. Identification and function of the

656   RNA chaperone Hfq in the Lyme disease spirochete *Borrelia burgdorferi*. Mol Microbiol

657   78:622-35.

658   45.   Zamba-Campero M, Soliman D, Yu H, Lasseter AG, Chang Y-Y, Liu J, Aravind L,

659   Jewett MW, Storz G, Adams PP. 2024. Broadly conserved FlgV controls flagellar

660   assembly and *Borrelia burgdorferi* dissemination in mice. bioRxiv

661   doi:10.1101/2024.01.09.574855:2024.01.09.574855.

662   46.   Juhas M, Ajioka JW. 2015. Identification and validation of novel chromosomal

663   integration and expression loci in *Escherichia coli* flagellar region 1. PLoS One

664   10:e0123007.

665   47.   Stevenson B, Brissette CA. 2023. Erp and Rev Adhesins of the Lyme Disease

666   Spirochete's Ubiquitous cp32 Prophages Assist the Bacterium during Vertebrate

667   Infection. Infect Immun 91:e0025022.

668   48.   Stevenson B, El-Hage N, Hines MA, Miller JC, Babb K. 2002. Differential binding of

669   host complement inhibitor factor H by *Borrelia burgdorferi* Erp surface proteins: a

670   possible mechanism underlying the expansive host range of Lyme disease spirochetes.

671   Infect Immun 70:491-7.

672   49.   Brissette CA, Haupt K, Barthel D, Cooley AE, Bowman A, Skerka C, Wallich R, Zipfel

673   PF, Kraiczy P, Stevenson B. 2009. *Borrelia burgdorferi* infection-associated surface

674   proteins ErpP, ErpA, and ErpC bind human plasminogen. Infect Immun 77:300-6.

675   50.   Brissette CA, Verma A, Bowman A, Cooley AE, Stevenson B. 2009. The *Borrelia*

676   *burgdorferi* outer-surface protein ErpX binds mammalian laminin. Microbiology

677   (Reading) 155:863-872.

678   51.   Garrigues RJ, Thomas S, Leong JM, Garcia BL. 2022. Outer surface lipoproteins from

679   the Lyme disease spirochete exploit the molecular switch mechanism of the complement

680   protease C1s. J Biol Chem 298:102557.

681   52.   Pereira MJ, Wager B, Garrigues RJ, Gerlach E, Quinn JD, Dowdell AS, Osburne MS,

682   Zückert WR, Kraiczy P, Garcia BL, Leong JM. 2022. Lipoproteome screening of the

683   Lyme disease agent identifies inhibitors of antibody-mediated complement killing. Proc

684   Natl Acad Sci U S A 119:e2117770119.

685   53.   Babb K, McAlister JD, Miller JC, Stevenson B. 2004. Molecular characterization of

686   *Borrelia burgdorferi erp* promoter/operator elements. J Bacteriol 186:2745-56.

687   54.   Miller JC, von Lackum K, Babb K, McAlister JD, Stevenson B. 2003. Temporal analysis

688   of *Borrelia burgdorferi* Erp protein expression throughout the mammal-tick infectious

689   cycle. Infect Immun 71:6943-52.

690   55.   Burns LH, Adams CA, Riley SP, Jutras BL, Bowman A, Chenail AM, Cooley AE,

691   Haselhorst LA, Moore AM, Babb K, Fried MG, Stevenson B. 2010. BpaB, a novel

692   protein encoded by the Lyme disease spirochete's cp32 prophages, binds to *erp* Operator

693   2 DNA. Nucleic Acids Res 38:5443-55.

694   56.   Jutras BL, Verma A, Adams CA, Brissette CA, Burns LH, Whetstine CR, Bowman A,

695   Chenail AM, Zuckert WR, Stevenson B. 2012. BpaB and EbfC DNA-binding proteins

696   regulate production of the Lyme disease spirochete's infection-associated Erp surface

697   proteins. J Bacteriol 194:778-86.

698   57.   Jutras BL, Chenail AM, Carroll DW, Miller MC, Zhu H, Bowman A, Stevenson B. 2013.

699   Bpur, the Lyme disease spirochete's PUR domain protein: identification as a

700   transcriptional modulator and characterization of nucleic acid interactions. J Biol Chem

701   288:26220-34.

702   58.   Si F, Li D, Cox SE, Sauls JT, Azizi O, Sou C, Schwartz AB, Erickstad MJ, Jun Y, Li X,

703   Jun S. 2017. Invariance of Initiation Mass and Predictability of Cell Size in *Escherichia*

704   *coli*. Curr Biol 27:1278-1287.

705   59.   Myrbraten IS, Wiull K, Salehian Z, Havarstein LS, Straume D, Mathiesen G, Kjos M.

706   2019. CRISPR Interference for Rapid Knockdown of Essential Cell Cycle Genes in

707   *Lactobacillus plantarum*. mSphere 4.

708   60.   Bjånes E, Stream A, Gibson PS, Bravo AM, Dahesh S, Baker JL, Varble A, Nizet V,

709   Veening J-W. 2024. An efficient *in vivo*-inducible CRISPR interference system for group

710   A *Streptococcus* genetic analysis and pathogenesis studies. bioRxiv

711   doi:10.1101/2024.02.22.581527:2024.02.22.581527.

712   61.   Fenster JA, Werner AZ, Tay JW, Gillen M, Schirokauer L, Hill NC, Watson A, Ramirez

713   KJ, Johnson CW, Beckham GT, Cameron JC, Eckert CA. 2022. Dynamic and single cell

714   characterization of a CRISPR-interference toolset in *Pseudomonas putida* KT2440 for β-

715   ketoadipate production from p-coumarate. Metab Eng Commun 15:e00204.

62.   Meunier A, Cornet F, Campos M. 2021. Bacterial cell proliferation: from molecules to

cells. FEMS Microbiol Rev 45.

63.   Jutras BL, Scott M, Parry B, Biboy J, Gray J, Vollmer W, Jacobs-Wagner C. 2016. Lyme

disease and relapsing fever *Borrelia* elongate through zones of peptidoglycan synthesis

that mark division sites of daughter cells. Proc Natl Acad Sci U S A 113:9162-70.

64.   Van De Putte P, Van D, Roersch A. 1964. The selection of mutants of *Escherichia coli*

with impaired cells division at elevated temperature. Mutat Res 106:121-8.

65.   Wang H, Henk MC, Gayda RC. 1993. Overexpression of *ftsA* induces large bulges at the

septal regions in *Escherichia coli*. Current Microbiology 26:175-181.

66.   Meisner J, Montero Llopis P, Sham LT, Garner E, Bernhardt TG, Rudner DZ. 2013.

FtsEX is required for CwlO peptidoglycan hydrolase activity during cell wall elongation

in *Bacillus subtilis*. Mol Microbiol 89:1069-83.

67.   Li J, Xu X, Shi J, Hermoso JA, Sham L-T, Luo M. 2023. Regulation of the cell division

hydrolase RipC by the FtsEX system in *Mycobacterium tuberculosis*. Nature

Communications 14:7999.

68.   Xu X, Li J, Chua W-Z, Pages MA, Shi J, Hermoso JA, Bernhardt T, Sham L-T, Luo M.

2023. Mechanistic insights into the regulation of cell wall hydrolysis by FtsEX and EnvC

at the bacterial division site. Proceedings of the National Academy of Sciences

120:e2301897120.

69.   Du S, Pichoff S, Lutkenhaus J. 2016. FtsEX acts on FtsA to regulate divisome assembly

and activity. Proceedings of the National Academy of Sciences 113:E5052-E5061.

70.   Diez AA, Farewell A, Nannmark U, Nyström T. 1997. A mutation in the *ftsK* gene of

738   *Escherichia coli* affects cell-cell separation, stationary-phase survival, stress adaptation,

739   and expression of the gene encoding the stress protein UspA. J Bacteriol 179:5878-83.

740   71.   Liu G, Draper GC, Donachie WD. 1998. FtsK is a bifunctional protein involved in cell

741   division and chromosome localization in *Escherichia coli*. Mol Microbiol 29:893-903.

742   72.   Steiner W, Liu G, Donachie WD, Kuempel P. 1999. The cytoplasmic domain of FtsK

743   protein is required for resolution of chromosome dimers. Mol Microbiol 31:579-83.

744   73.   Crozat E, Rousseau P, Fournes F, Cornet F. 2014. The FtsK family of DNA translocases

745   finds the ends of circles. J Mol Microbiol Biotechnol 24:396-408.

746   74.   Yu XC, Weihe EK, Margolin W. 1998. Role of the C terminus of FtsK in *Escherichia

747   coli* chromosome segregation. J Bacteriol 180:6424-8.

748   75.   Capiaux H, Lesterlin C, Pérals K, Louarn JM, Cornet F. 2002. A dual role for the FtsK

749   protein in *Escherichia coli* chromosome segregation. EMBO Rep 3:532-6.

750   76.   Aussel L, Barre FX, Aroyo M, Stasiak A, Stasiak AZ, Sherratt D. 2002. FtsK Is a DNA

751   motor protein that activates chromosome dimer resolution by switching the catalytic state

752   of the XerC and XerD recombinases. Cell 108:195-205.

753   77.   Begg KJ, Dewar SJ, Donachie WD. 1995. A new *Escherichia coli* cell division gene,

754   *ftsK*. J Bacteriol 177:6211-22.

755   78.   Grenga L, Luzi G, Paolozzi L, Ghelardini P. 2008. The *Escherichia coli* FtsK functional

756   domains involved in its interaction with its divisome protein partners. FEMS

757   microbiology letters 287:163-167.

79.   Chen JC, Beckwith J. 2001. FtsQ, FtsL and FtsI require FtsK, but not FtsN, for co-

localization with FtsZ during *Escherichia coli* cell division. Molecular microbiology

42:395-413.

80.   Sperandio V, Torres AG, Kaper JB. 2002. Quorum sensing *Escherichia coli* regulators B

and C (QseBC): a novel two-component regulatory system involved in the regulation of

flagella and motility by quorum sensing in *E. coli*. Mol Microbiol 43:809-21.

81.   Savage CR, Jutras BL, Bestor A, Tilly K, Rosa PA, Tourand Y, Stewart PE, Brissette

CA, Stevenson B. 2018. *Borrelia burgdorferi* SpoVG DNA- and RNA-Binding Protein

Modulates the Physiology of the Lyme Disease Spirochete. J Bacteriol 200.

82.   Saylor TC, Savage CR, Krusenstjerna AC, Jusufovic N, Zuckert WR, Brissette CA,

Motaleb M, Schlax PJ, Stevenson B. 2023. Quantitative analyses of interactions between

SpoVG and RNA/DNA. Biochem Biophys Res Commun 654:40-46.

83.   Pappas CJ, Iyer R, Petzke MM, Caimano MJ, Radolf JD, Schwartz I. 2011. *Borrelia*

burgdorferi requires glycerol for maximum fitness during the tick phase of the enzootic

cycle. PLoS Pathog 7:e1002102.

84.   He M, Ouyang Z, Troxell B, Xu H, Moh A, Piesman J, Norgard MV, Gomelsky M, Yang

XF. 2011. Cyclic di-GMP is essential for the survival of the Lyme disease spirochete in

ticks. PLoS pathogens 7:e1002133.

85.   Caimano MJ, Dunham-Ems S, Allard AM, Cassera MB, Kenedy M, Radolf JD. 2015.

Cyclic di-GMP modulates gene expression in Lyme disease spirochetes at the tick-

mammal interface to promote spirochete survival during the blood meal and tick-to-

mammal transmission. Infect Immun 83:3043-60.

86.   Blevins JS, Xu H, He M, Norgard MV, Reitzer L, Yang XF. 2009. Rrp2, a sigma54-

781   dependent transcriptional activator of *Borrelia burgdorferi*, activates *rpoS* in an

782   enhancer-independent manner. J Bacteriol 191:2902-5.

783   87.   Xu Q, Shi Y, Dadhwal P, Liang FT. 2012. RpoS regulates essential virulence factors

784   remaining to be identified in Borrelia burgdorferi. PLoS One 7:e53212.

785   88.   Ouyang Z, Zhou J, Norgard MV. 2014. Synthesis of RpoS is dependent on a putative

786   enhancer binding protein Rrp2 in *Borrelia burgdorferi*. PLoS One 9:e96917.

787   89.   Samuels DS. 2011. Gene regulation in *Borrelia burgdorferi*. Annual review of

788   microbiology 65:479-99.

789   90.   Castro-Padovani TN, Saylor TC, Husted OT, Krusenstjerna AC, Jusufovic N, Stevenson

790   B. 2023. Gac Is a Transcriptional Repressor of the Lyme Disease Spirochete's OspC

791   Virulence-Associated Surface Protein. J Bacteriol doi:10.1128/jb.00440-22:e0044022.

792   91.   Rollend L, Fish D, Childs JE. 2013. Transovarial transmission of *Borrelia* spirochetes by

793   *Ixodes scapularis*: a summary of the literature and recent observations. Ticks Tick Borne

794   Dis 4:46-51.

795   92.   Casjens S, van Vugt R, Tilly K, Rosa PA, Stevenson B. 1997. Homology throughout the

796   multiple 32-kilobase circular plasmids present in Lyme disease spirochetes. J Bacteriol

797   179:217-27.

798   93.   Barbour AG, Hayes SF, Heiland RA, Schrumpf ME, Tessier SL. 1986. A *Borrelia*-

799   specific monoclonal antibody binds to a flagellar epitope. Infect Immun 52:549-54.

800   94.   El-Hage N, Babb K, Carroll JA, Lindstrom N, Fischer ER, Miller JC, Gilmore RD, Jr.,

801   Mbow ML, Stevenson B. 2001. Surface exposure and protease insensitivity of *Borrelia*

802   *burgdorferi* Erp (OspEF-related) lipoproteins. Microbiology (Reading) 147:821-830.

803   95.   Song X, Goedhart J. 2024. EzReverse – a Web Application for Background Adjustment

804   of Color Images. bioRxiv doi:10.1101/2024.05.27.594095:2024.05.27.594095.

805   96.   Goedhart J, Luijsterburg MS. 2020. VolcaNoseR is a web app for creating, exploring,

806   labeling and sharing volcano plots. Scientific Reports 10.

807   97.   Kolde R, Kolde MR. 2015. Package 'pheatmap'. R package 1:790.

808   98.   Zhao X, Zhang K, Boquoi T, Hu B, Motaleb MA, Miller KA, James ME, Charon NW,

809   Manson MD, Norris SJ, Li C, Liu J. 2013. Cryoelectron tomography reveals the

810   sequential assembly of bacterial flagella in *Borrelia burgdorferi*. Proc Natl Acad Sci U S

811   A 110:14390-5.

812   99.   Qin Z, Tu J, Lin T, Norris SJ, Li C, Motaleb MA, Liu J. 2018. Cryo-electron tomography

813   of periplasmic flagella in *Borrelia burgdorferi* reveals a distinct cytoplasmic ATPase

814   complex. PLoS Biol 16:e3000050.

815   100.   Chang Y, Moon KH, Zhao X, Norris SJ, Motaleb MA, Liu J. 2019. Structural insights

816   into flagellar stator–rotor interactions. eLife 8:e48979.

Figure Legends.

Figure 1. Validation of conditional dysregulation of DnaA. (A) Two sgRNAs were designed

to knock down borrelial *dnaA* transcription by CRISPRi. One sgRNA targeted the template

strand directly upstream of the ORF ($dnaA_{T1}$), while the other targeted the template strand within

the ORF ($dnaA_{NT1}$). **(B)** The *dnaA* CRISPRi bacteria, CRISPRi empty vector bacteria, and the

strain with the *dnaA* overexpression plasmid grew at the same rate as the parental e2. Adding

IPTG to these strains resulted in appropriate knockdown or overexpression of DnaA protein **(C)**

and transcript **(D)**, as assessed by immunoblot and qRT-PCR. The overexpression shuttle vector

encodes a DnaA with an N-terminal 3xFLAG moiety and thus migrates above the native protein.

Error bars in (B) and (D) represent the standard error of the mean (SEM).

Figure 2. Appropriate expression of DnaA is required for *B. burgdorferi* replication. (A)

Growth curve studies were conducted on different *B. burgdorferi* strains incubated with 0.5 mM

IPTG. Three independent cultures of each strain were measured for growth curve analysis. Error

bars represent the SEM. The dashed line indicates when the overexpression strain resumed

growth. **(B)** The generation time was determined from the growth curve analyses from three

individual cultures of each strain. For the overexpression strain, the growth rate was determined

before (early; 1-4 dpi) and after the resumption of growth (late; 4-7 dpi). The resumption of

replication in that strain was evidently due to mutation of the overexpressed *dnaA* gene. The p-

values are indicated as follows: **, $p \leq 0.01$; ***, $p \leq 0.001$; ns, not significant, $p > 0.05$. **(C)**

IPTG-treated DnaA overexpression spirochetes that resumed growth (conditioned) lost

sensitivity to IPTG when passaged to fresh media and did not overproduce 3xFLAG-DnaA as

assessed by immunoblot **(D)**, supporting the conclusion that accumulated mutations in the

overexpression *dnaA* permitted replication. **(E)** Sequenced plasmid from these conditioned

spirochetes showed unique mutations (S = substitution; I= Insertion; D = deletion) in the

3xFLAG-*dnaA* ORF that caused a frameshift and truncation of the full-length protein. The

location of the truncation is indicated by *.

Figure 3. DnaA dysregulation affects spirochete cell length. (A-D) Cell lengths of the 0.5 mM

IPTG-treated *B. burgdorferi* B31-e2 strains measured over seven days (gray = e2; red = $dnaA_{T1}$;

$dnaA_{NT1}$ = blue; P_{lac}::3xFLAG-*dnaA* = purple). The line represents the median cell length. **(E-H)**

The median cell length (filled-in circle) with error bars representing the 95% CI superimposed

against the corresponding growth curve (empty circle) from Fig. 2. The dashed line indicates

when the overexpression strain resumed growth. **(I-K)** Representative images of observed

phenotypes. **(I)** The e2 cell of approximate median length at 2 dpi. **(J)** DnaA-depleted cell with

the characteristic elongation phenotype. **(K)** Median size of a DnaA overexpression spirochete at

2 dpi.

Figure 4. Knockdown of DnaA increases the number of chromosomes per spirochete. The

B. burgdorferi CRISPRi $dnaA_{T1}$ and parent B31-e2 strains were transformed with a construct

encoding a mCherry-tagged ParB protein, which binds near the *oriC*, to identify the locations of

the chromosomes. The two strains were grown without inducer until mid-log ($t = 0$), and then the

$dnaA_{T1}$ strain was divided into two cultures, and one of those was induced with 0.5 mM IPTG.

Cultures were tracked for three more days. **(A)** Representative merged micrographs (blue = dark

field; red = mCherry-ParB) of spirochetes with the approximate median numbers of ParB-*oriC*

puncta **(B)** and cell length **(C)**. Significant differences in numbers of puncta and cell length at

day 0 were determined by the Mann-Whitney U test. Differences between the three groups (e2, -

IPTG, and + IPTG) on days 1-3 were tested for by the Kruskal-Wallis test and then Dunn's test

for multiple comparisons. Multiplicity-adjusted p-values are reported. **, p ≤ 0.01; ***, p ≤

0.001; ****, p ≤ 0.0001; ns, not significant, p > 0.05. Approximately 50 spirochetes were

imaged for each strain on each day. Data were gathered from two independent experiments.

Figure 5. Irregular spacing of *oriC* in *dnaA* knockdown *B. burgdorferi*. (A-B) Representative

images of spirochetes containing the $dnaA_{T1}$ CRISPRi plasmid with irregular spacing of ParB-

oriC puncta (red). DnaA-depleted cells had large regions where no foci were observed

(brackets). Some cells also had foci that were close together (arrows). **(C)** Parental B31-e2

spirochetes with approximately the same number of ParB-*oriC* puncta as the mutant but exhibit

regular oriC spacing. **(D)** The number of spirochetes with irregular ParB-*oriC* was quantified

(e2, *n* = 184; uninduced $dnaA_{T1}$, *n* = 198 cells; induced $dnaA_{T1}$ *n* = 206). One-way ANOVA with

Tukey's post-hoc test was done to determine the significance and make comparisons. The

multiplicity-adjusted p-values are reported **, $p \leq 0.01$; ns, not significant, $p > 0.05$.

Figure 6. DnaA-depletion affects borrelial helicity. Representative micrographs of observed

phenotypes. When *dnaA* was knocked down, spirochetes with no **(A)** or asymmetrical helicity

(B) were observed. **(C)** Some of the abnormal cells were elongated and showed division defects.

(D) Spirochetes exhibiting similar defects were observed at a low level in the B31-e2 strain

(0.7%, *n* = 1514). The empty vector (EV) control cultures had a similar proportion, independent

of the addition of IPTG (uninduced: 2.1%, *n* = 1439; induced: 1.6%, *n* = 1608). The bacteria

with the $dnaA_{T1}$ CRISPRi plasmid had significantly more cells with perturbed helicity that was

dependent on induction (uninduced: 8.7%, *n* = 1437; induced: 13%, *n* = 1472). Spirochetes were

counted from three independent cultures. Statistical significance was determined by one-way

ANOVA with a Holm-Šídák's post-hoc test, and multiplicity-adjusted p-values are reported. ***,

$p \leq 0.001$, ****, $p \leq 0.0001$; ns, not significant, $p > 0.05$.

Figure 7. RNA-seq of DnaA-dysregulated *B. burgdorferi*. The RNA from three *B. burgdorferi*

strains with different cellular levels of DnaA were sequenced: B31-e2 (parent), CRISPRi $dnaA_{T1}$

(DnaA-down), and overexpression pACK121 (DnaA-up). **(A-C)** Volcano plots plotting the fold

change (log_2FC) against each gene's false discovery rate ($-log_{10}(FDR)$). The threshold for

significance was set at a $log_2FC \geq 1$ or ≤ -1 ($\geq$ 2-fold change) and an FDR of ≤ 0.05. **(D)** PCA

plot showing the clustering of the replicate samples that were sequenced. **(E)** Donut charts

showing the replicons on which the significantly affected genes are encoded. **(F)** qRT-PCR

results of *dnaA* transcripts in the sequenced samples. Cq values were normalized to *ftsK* or *rpoB*

(ΔCq) and then to the parental strain ($\Delta\Delta$Cq). Error bars represent the standard deviation (SD).

Figure 8. Impacts of DnaA on *B. burgdorferi* elongation and division genes. (A) Heat map of

the division and elongation genes that were significantly affected when DnaA was knocked

down. **(B)** Top: Locus maps of the select elongation and division genes to scale. The sites of

potential DnaA-boxes are denoted as purple boxes below the genes. Bottom: Simplified schema

of the hypothesized borrelial elongasome and divisome containing the highlighted genes of

interest. Informed by the work of Liu et al. and Hu et al. (36, 69) **(C)** Bar graph showing the fold

change detected by RNA-seq of the elongation and division genes. The dashed lines indicate the

2-fold threshold for meaningful gene expression changes.

Figure 9. Impacts of DnaA on *B. burgdorferi* flagellar genes. (A) The borrelial flagellar motor

is a complex molecular machine (bottom right) that is comprised of nearly three dozen proteins

encoded in four separate loci on the chromosome (top). The schematic of the *B. burgdorferi*

flagellar motor was informed from data by Zhao et al., Qin et al., and Chang et al. (98-100). **(B)**

RNA-seq showed the transcripts for five flagellar genes increased when DnaA was knocked

down. Two of these genes, FliQ and FliR (purple), are constituents of the flagellar export

apparatus. The dashed lines indicate the 2-fold threshold for meaningful gene expression

changes.

Figure. 10. Impacts of DnaA on the Erp regulatory network. (A) Solid black lines indicate

established interactions. Dashed lines indicate hypothesized interactions. Activation is denoted

by lines with arrowheads, and inhibition by lines without arrowheads. Note that co-repressive or

anti-repressive activities, i.e., BpuR and EbfC, respectively, are indicated by arrows directed at

lines. **(B)** Representative immunoblots from samples of *B. burgdorferi* that were analyzed by

RNA-seq. Quantitation of target proteins when DnaA was knocked down **(C)** or overexpressed

(D). Band intensities for each replicate immunoblot were normalized to FlaB and then the

parental e2 strain. **(C-D)** The dotted lines at 1 represent where the protein abundance is the same

as e2. Bar graphs showing the fold change detected by RNA-seq in the DnaA-down/e2 **(E)** and

DnaA-up comparisons **(F). (E-F)** The dotted lines indicate the 2-fold threshold for meaningful

gene expression changes.

Figure 11. Roles and impacts of DnaA dysregulation on borrelial physiology. The DnaA of

B. burgdorferi functions as the chromosome replication initiator and a transcription factor.

Altering levels of this essential protein modulate aspects of DNA replication and bacterial

physiology. The data presented here point to crosstalk between the systems controlling

replication and bacterial morphology. Future studies will need to be conducted to ascertain the

direct roles DnaA plays in maintaining this delicate balance.

Table 1. Homologous elongation and division genes of *B. burgdorferi*.

Protein	Locus Tag	Elongasome	Divisome	RNA-seq Differential Expression		
				DnaA-down vs e2	DnaA-up vs e2	DnaA-down vs DnaA-up
FtsE	BB_0080	-	+	Down	NA	Down
FtsX	BB_0081	-	+	Up	NA	Up
FtsI/PBP3	BB_0136	+	+	NA	NA	NA
MurE	BB_0201	+	+	NA	NA	NA
FtsK	BB_0257	-	+	Down	NA	Down
FtsZ	BB_0299	-	+	NA	NA	NA
FtsA	BB_0300	-	+	Up	Up	NA
FtsQ/DivlB	BB_0301	-	+	NA	NA	NA
FtsW	BB_0302	-	+	NA	NA	NA
MraY	BB_0303	+	+	NA	NA	NA
MurF	BB_0304	+	+	NA	NA	NA
MurD	BB_0585	+	+	NA	NA	NA
MurB	BB_0598	+	+	NA	NA	NA
Ami	BB_0666	-	+	NA	NA	NA
MreB	BB_0715	+	-	Down	NA	Down
MreC	BB_0716	+	-	Down	NA	Down
MreD	BB_0717	+	-	Up	NA	Up
PBP2	BB_0718	+	-	Down	NA	Down
RodA	BB_0719	+	-	NA	NA	NA
PBP1a	BB_0732	+	-	Down	NA	Down
MurG	BB_0767	+	+	NA	NA	NA
MurJ	BB_0810	+	+	NA	Down	Up
MurC	BB_0817	+	+	NA	NA	NA

941   * Plus sign (+) indicates the protein is part of the complex, minus sign (-) indicates it is not

942   * NA = Not affected

944   **Table 2.** Oligonucleotides used in this study.

Target	Name	Sequence	Purpose	Source
dnaA ORF	qPCR *dnaA* F	CATGTGACCGATCTCCTTCTG	qPCR	(10)
	qPCR *dnaA* R	CGACAATAGCTGCTCTGAGTT		
ftsK ORF	qPCR *ftsK* F	GACCTTCTGATGAGCCAATGT	qPCR	(10)
	qPCR *ftsk* R	GCTGCTCTGTTGTAACCTATCT		
rpoB ORF	qPCR *rpoB* F	GTTCATCTGGGACAAGGAAGAG	qPCR	This study
	qPCR *rpoB* R	CAGACTCAAGGCCCTCATTAAG		

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Dark Field **mCherry-ParB**

Figure 6.

Figure 7.

Figure 8.

Figure 9.

Figure 10.

Figure 11.