The long-chain fatty acid-CoA ligase FadD1 monitors *Pseudomonas aeruginosa* quorum sensing as a receptor of *cis*-2-decenoic acid

Shihao Song[1,2,11], Xia Li[3,11], Jingyun Liu[2,4,11], Bing Wang[1], Yang Si[5], Hongguang Han[1], Xiuyun Sun[1,3], Mingfang Wang[3], Binbin Cui[3,6], Guangliang Wu[6], Yongliang Huo[7], Liangxiong Xu[8], Beile Gao[9], Liang Yang[10], Xiaoxue Wang[9], Lian-Hui Zhang[2], Yinyue Deng[3,*]

[1]*Key Laboratory of Tropical Biological Resources of Ministry of Education, School of Pharmaceutical Sciences, Hainan University, Haikou 570228, China*

[2]*Integrative Microbiology Research Center, South China Agricultural University, Guangzhou 510642, China*

[3]*School of Pharmaceutical Sciences (Shenzhen), Shenzhen Campus of Sun Yat-sen University, Sun Yat-sen University, Shenzhen 518107, China*

[4]*Zhengzhou Shuqing Medical College, Zhengzhou 450064, China*

[5]*Department of Microbiology, College of Life Sciences, Nanjing Agricultural University, Nanjing 210095, China*

[6]*Pharmacy Department, The Affiliated LiHuiLi Hospital of Ningbo University, Ningbo315046, China*

[7]*Experimental Animal Center, Guangzhou Medical University, Guangzhou, Guangdong, 511436, China*

[8]*School of Life Sciences, Huizhou University, Huizhou 516007, China*

[9]*Guangdong Key Laboratory of Marine Materia Medica, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China*

[10]*School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.*

[11]*These authors contributed equally: Shihao Song, Xia Li and Jingyun Liu.*

Corresponding author:

Yinyue Deng: dengyle@mail.sysu.edu.cn

Running title: *Cis*-2-decenoic acid controls *P. aeruginosa* virulence

Abstract

Diffusible signal factor (DSF)-family quorum sensing (QS) signals are widely utilized by many bacterial pathogens to modulate various biological functions and virulence. Previous studies showed that *cis*-2-decenoic acid (*cis*-DA) is involved in modulation of biofilm dispersion in *Pseudomonas aeruginosa*, but its signaling mechanism remains vague. Here, we report that *cis*-DA regulates the physiology and virulence of *P. aeruginosa* through the long-chain fatty acid-CoA ligase FadD1. *Cis*-DA specifically binds to FadD1 with high affinity and enhances the binding of FadD1 to the promoter DNA region of *lasR*. Further analysis showed that the FadD1 is a response regulator of *cis*-DA with a DNA-binding leucine zipper motif to control the transcription of various target genes. Moreover, FadD1 exhibited catalytic activity on *cis*-2-dodecenoic acid (BDSF) of *Burkholderia cenocepacia* and enhanced the competitiveness of *P. aeruginosa*. Together, our work presents a new DSF-type QS signaling system, which is highlighted by its receptor and response regulator evolved from a canonical enzyme of fatty acid metabolism.

Introduction

Quorum sensing (QS) is a cell-to-cell communication mechanism utilized by numerous species of bacteria. The first QS signal discovered in gram-negative bacteria is the acyl homoserine lactone (AHL) produced by *Vibrio fischeri*[1-5]. In addition to AHL-family signals, there are many other types of QS signals[6-16], including the diffusible signal factor (DSF)-family signals. *Pseudomonas aeruginosa* is a major source of opportunistic infections in both immunocompromised individuals and cystic fibrosis patients and has evolved at least three types of QS systems, e.g., *las*, *pqs* and *rhl*[17-19]. The *las* and *rhl* AHL QS systems use N-3-oxo-dodecanoyl-l-homoserine lactone (3-oxo-C12-HSL) and N-butyryl-l-homoserine lactone (C4-HSL), respectively, to regulate biological functions, such as biofilm formation, motility and virulence factor production[20-24]. The *pqs* system employs 2-heptyl-3-hydroxy-4(1H)-quinolone (PQS), to regulate biological functions and virulence[25-29]. These QS systems of *P. aeruginosa* are hierarchically interrelated, and the *las* system was verified to monitor both the *rhl* and *pqs* systems[17, 30]. Moreover, it was also revealed that the fatty acid signaling molecule *cis*-2-decenoic acid (*cis*-DA) exhibited important functions in *P. aeruginosa*[31]. The production of *cis*-DA in *P. aeruginosa* needs an enoyl-CoA synthase encoded by DspI[32]. The inactivation of DspI leads to a significant reduction in biofilm dispersion[33,34]. However, it is not yet clear how *cis*-DA regulates biofilm

dispersion, and its receptor and downstream signaling network remain to be investigated.

Fatty acyl coenzyme A (CoA) synthetases (FACSs; fatty acid-CoA ligases) are a class of enzymes

that activate alkanoic acids to CoA esters[35]. FACSs are widely distributed in both prokaryotic and

eukaryotic organisms and exhibit broad substrate specificity[36]. The long-chain fatty acid-CoA ligase

(FadD) is responsible for the activation of endogenous long-chain fatty acids into acyl-CoAs[37]. RpfB,

which has recently been reported as a long-chain fatty acid-CoA ligase of *Xanthomonas campestris*

pv. campestris (*Xcc*), is involved in the activation of a wide range of fatty acids to their CoA esters *in*

vitro[38]. Intriguingly, RpfB is required for the turnover of both *cis*-11-methyl-dodecenoic acid (DSF) and

cis-2-dodecenoic acid (BDSF) QS signals *in vivo*[39]. The *rpfB* deletion mutant produced high levels of

both DSF and BDSF signals and displayed increased virulence in *Xcc*[40,41]. Moreover, RpfB-dependent

QS signal turnover was also detected in various bacterial species[40].

Transcription factors (TFs) are proteins that bind to specific DNA sequences, thereby regulating the

transcription of genetic information from DNA to messenger RNA[42]. Once bound to DNA, these

proteins can promote or block the recruitment of RNA polymerase to specific genes to control

transcriptional expression[43]. Although there is great diversity in the structure and function of DNA-

binding proteins, some polypeptide motifs enable binding to the major groove of DNA, including helix-

turn-helix (HTH), helix-loop-helix (HLH), zinc fingers, and leucine zippers[44-47].

In this study, we demonstrated that FadD1 of *P. aeruginosa* could convert BDSF to *cis*-2-

dodecenoic acid-CoA and reduce the competitiveness of *B. cenocepacia*. FadD1 also controls the

physiology and virulence of *P. aeruginosa* by sensing *cis*-DA, then monitors the QS hierarchy and

controls the expression of various target genes as a global transcriptional regulator. These findings

indicate that FadD1 is a member of a new class of response regulators of QS signal that controls the

important biological functions and virulence of bacterial pathogens, with an additional role in fatty acid

oxidation to catalyze the formation of fatty acyl-CoA.

Results

FadD1 of *P. aeruginosa* shows enzyme activity on the BDSF signal

BDSF was first found to be involved in the regulation of biofilm formation, motility and virulence of *B.*

cenocepacia[9-12,48,49]. It was subsequently reported that BDSF from *B. cenocepacia* interferes with QS

systems and the type III secretion system (T3SS) of *P. aeruginosa*[50]. Interestingly, we found that

BDSF could also be degraded when it was added to the culture of *P. aeruginosa* PAO1. The amount

of BDSF in the culture of *P. aeruginosa* decreased to 21.78% after 6 h of incubation (Fig. 1a). Previous

work demonstrated that RpfB is a long-chain fatty acid-CoA ligase responsible for BDSF and DSF

signal turnover in *Xcc*[38,39]. To discover the enzyme that degrades BDSF, we then searched RpfB

homologs in the genome of *P. aeruginosa* PAO1 by using the Basic Local Alignment Search Tool

(BLAST) algorithm[51]. We found six RpfB homologs, FadD1-6[52]; among them, FadD1 showed the

highest homology, with 55.35% identity with RpfB. Therefore, FadD1, which contains 562 aa with a

calculated molecular weight of 61.67 kDa, was purified to homogeneity using affinity chromatography

(Fig. 1b). The *in vitro* enzymatic activity assays showed that when FadD1 was mixed with BDSF, the

free BDSF level in the mixture decreased nearly undetectable after 1 h (Fig. 1c). These results

suggested that FadD1 exhibits enzyme activity on BDSF. Consistently, deletion of *fadD1* in *P.*

aeruginosa caused a substantial reduction in BDSF degradation activity (Fig. 1d) but did not affect

the growth rate of the bacterial cells (Supplementary Fig. 1).

We then investigated the *in vivo* activity of FadD1 in *B. cenocepacia* H111 by overexpressing the

fadD1 gene. Overexpression of *fadD1* did not affect the growth rate but markedly reduced the BDSF

production of the bacterial cells (Supplementary Fig. 2a, b). As expected, deletion of *rpfF$_{Bc}$* resulted

in complete abolishment of the production of BDSF in *B. cenocepacia* H111 (Supplementary Fig. 2b).

Interestingly, overexpression of *fadD1* in the *B. cenocepacia* wild-type strain reduced biofilm formation,

swarming motility and protease activity, which could be restored by the addition of exogenous BDSF

at 5 μM (Supplementary Fig. 2c-e). Overexpression of *fadD1* in the wild-type strain H111 also

decreased cytotoxicity (Supplementary Fig. 2f).

FadD1 boosts the competitive capability of *P. aeruginosa* against *B. cenocepacia*

As BDSF mediates the cross-talk between *B. cenocepacia* and *P. aeruginosa* [50] and FadD1 exhibits

catalysis activity on BDSF, we investigated whether FadD1 plays a role in the competitive interactions

between the two bacterial species. To this end, we cocultured the green fluorescent protein (GFP)

fluorescence-labeled *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and complemented strains in the

presence of the *B. cenocepacia* H111 strain labeled with mCherry and measured the BDSF production

of *B. cenocepacia* (Fig. 1d) and the mean fluorescence intensity (MFI) of both *B. cenocepacia* and *P.*

aeruginosa at different time points as indicated (Fig. 1e, f). The results showed that deletion of *fadD1*

caused a drastic reduction in the degradation capability of *P. aeruginosa* toward the BDSF signal. The BDSF signal yield of *B. cenocepacia* cocultured with the *P. aeruginosa* deletion mutant Δ*fadD1* was more than 5 times higher than that when it was cocultured with wild-type PAO1 or the complemented strain Δ*fadD1*(*fadD1*) at 9 h postinoculation (Fig. 1d). On the other hand, we found that the GFP mean fluorescence intensity (MFI) of the *P. aeruginosa* Δ*fadD1* strain was lower than that of the *P. aeruginosa* wild-type and complemented strains when they were cocultured with the *B. cenocepacia* strains (Fig. 1e). The mCherry mean fluorescence intensity (MFI) of the *B. cenocepacia* H111 strain cocultured with the *P. aeruginosa* Δ*fadD1* strain was higher than that of the *B. cenocepacia* H111 strain cocultured with the *P. aeruginosa* wild-type and complemented strains (Fig. 1f). These results indicated that FadD1 plays an important role in the competition between *P. aeruginosa* and *B. cenocepacia*.

Deletion of FadD1 impairs biological functions and pathogenicity in *P. aeruginosa*

It was determined that FadD1 plays an important role in the interaction between *P. aeruginosa* and *B. cenocepacia*. We continued to study whether FadD1plays a role in the regulation of biological functions and pathogenesis in *P. aeruginosa*. We found that deletion of *fadD1* resulted in decreases in biofilm formation, swarming, and pyocyanin production by 30.47%, 36.72% and 62.15%, respectively (Fig. 2a-c). Deletion of *fadD1* also attenuated cytotoxicity by 69.78% when A549 cells were incubated with the *P. aeruginosa* strains at 8 h postinoculation (Fig. 2d). We tested whether there was a change in the expression level of upstream (*PA3298*) and downstream (*fadD2*) genes of the *fadD1* mutant gene. The results showed that there was no change in the expression of upstream and downstream genes in the *fadD1* deletion mutant (Supplementary Fig. 3). To further study the regulatory roles of FadD1 in controlling bacterial physiology, we analyzed and compared the transcriptomes of the wild-type strain and the Δ*fadD1* mutant strain by using RNA sequencing (RNA-Seq). Differential gene expression analysis showed that the expression levels of 23 genes were increased and those of 208 genes were decreased in the Δ*fadD1* mutant strain compared with the expression levels in the wild-type strain ($|Log_2$ fold-change$| \geq 1.0$) (Supplementary Table 1 and Supplementary Fig. 4a). These differentially expressed genes are associated with a range of biological functions (Supplementary Table 1 and Supplementary Fig. 4b). These genes include the *las*, *rhl* and *pqs* QS system genes *PA1432* (*lasI*), *PA1430* (*lasR*), *PA3476* (*rhlI*), *PA3477* (*rhlR*),

PA0996 (*pqsA*), and *PA1003* (*mvfR*), which were shown to be involved in pathogenicity in *P.*
aeruginosa.

FadD1 monitors the QS systems of *P. aeruginosa*

As FadD1 obviously affects the QS-regulated genes and phenotypes of *P. aeruginosa* (Fig. 2a-d,
Supplementary Fig. 4 and Supplementary Table 1), we inferred that FadD1 might affect the QS
systems. Analysis of the expression profiles of *lacZ* under the control of the *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR*
and *pqsA* promoters showed that deletion of *fadD1* resulted in reduced expression levels of *lasR*, *lasI*,
rhlR, *rhlI*, *mvfR* and *pqsA* (Fig. 2e-j), which are signal molecule receptor genes and synthase-
encoding genes of the *las*, *rhl* and *pqs* QS systems, respectively. RT-qPCR analysis and RNA-seq
also showed that the mutation of *fadD1* caused a decrease in the expression levels of *lasR*, *lasI*, *rhlR*,
rhlI, *mvfR* and *pqsA* (Fig. 2k). Therefore, we measured and compared the production of 3-oxo-C12-
HSL, C4-HSL and PQS in the wild-type, *fadD1* mutant, and complemented strains. We found that the
production of 3-oxo-C12-HSL, C4-HSL and PQS was reduced in the *fadD1* mutant strain (Fig. 2l-n).
These results demonstrated that FadD1 positively regulated the *las*, *rhl* and *pqs* QS systems in *P.*
aeruginosa.

FadD1 regulates target gene expression by directly binding to the promoter

FadD1 obviously affects the QS gene expression levels and QS signal production of *P. aeruginosa*.
Therefore, we performed electrophoretic mobility shift assays (EMSAs) to test whether the
transcriptional regulation of the signal molecule synthase-encoding genes and receptor genes of the
las, *rhl* and *pqs* QS systems is achieved by direct binding of FadD1 to their promoters. The 339-bp,
321-bp, 270-bp, 330-bp, 306-bp and 324-bp DNA fragments from the *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and
pqsA promoters were PCR-amplified and used as probes. Surprisingly, as shown in Figure. 3*A*, the
lasR promoter DNA fragment formed stable DNA-protein complexes with FadD1 and migrated slower
than unbound probes. The amount of labeled probe that bound to FadD1 increased with increasing
amounts of FadD1 but decreased in the presence of a 50-fold greater concentration of unlabeled
probe (Fig. 3a). However, FadD1 did not bind to the promoters of the other tested genes, i.e., *rhlR*,
mvfR, *lasI*, *rhlI* and *pqsA* (Supplementary Fig. 5 and 6).

To identify the DNA motif recognized by FadD1 and the genes directly regulated by FadD1, we

performed chromatin immunoprecipitation sequencing (ChIP-Seq) analysis. ChIP-Seq data analysis revealed the potential FadD1 binding site as 5'-AGGACGG-3' in the *lasR* gene promoter probe (Fig. 3b). The results showed that FadD1 could directly bind to multiple target gene promoters to regulate their transcription and expression (Supplementary Table 2). To further examine the binding sites, three target genes, *flgM*, *katA* and *PA0692,* were chosen for EMSA analysis. The amounts of labeled probes of *flgM*, *katA* and *PA0692* bound to FadD1 were also increased with increasing amounts of FadD1 (Supplementary Fig. 7). Then, the potential binding sites 5'-AGGACGG-3', 5'-TGGCCGG-3', 5'-AGGAGGG-3', and 5'-AGGGCGG-3' were deleted from the promoter regions of *lasR*, *flgM*, *katA* and *PA0692*, respectively. EMSA analysis showed that no DNA-protein complex was formed when the binding site was deleted from the probes (Fig. 3c and Supplementary Fig. 8), indicating that this fragment is essential for the binding of FadD1 to the *lasR, flgM*, *katA* and *PA0692* promoters. Together, these results demonstrated that FadD1 regulates target gene expression by directly binding to a specific region in the gene promoter.

The leucine zipper motif of FadD1 is responsible for binding with DNA

Since FadD1 can bind to *lasR* and regulate transcription, and the domain structure analysis of FadD1 by using the HMMER (https://www.ebi.ac.uk/Tools/hmmer/) shows that it has an AMP-binding enzyme domain (AMP-B) and an AMP-binding enzyme C-terminal domain (AMP-B_C) (Fig. 3d). We then purified the AMP-B domain and the AMP-B_C domain for EMSA (Fig. 3e). The results showed that only the AMP-B domain can bind to the promoter region of *lasR* (Fig. 3f, g). To determine the location of the new DNA binding motif in the AMP-B domain, we used the protein motif analysis website (https://www.novopro.cn/tools/motifscan.html). A structure called "leucine zipper" was characterized at 275 to 296 amino acids **L**TGNHNI**L**ITNPRD**L**PSMLKD**L** in AMP-B domain (Fig. 3d and Supplementary Fig. 9). The leucine zipper consists of a periodic repetition of leucine residues at every seventh position over a distance covering eight helical turns. This motif is found in many eukaryotic transcription factors[45]. However, in bacterial transcription factor, this motif is only found in the classical DNA-binding HTH domain of the LysR family regulator MetR[46] and a BDSF response regulator DsfR identified recently[47]. So we speculated that the four leucine residues L275, L282, L289 and L296 might be important for the binding of FadD1 to target gene promoters. we then introduced four site-specific substitutions into FadD1 (named FadD1[LM]), and the results showed that the simultaneous

mutation of L275A, L282A, L289A and L296A abolished the binding of FadD1 to the *lasR* promoter

(Fig. 3i). In addition, *in trans* expression of FadD1^{LM} did not rescue the defective biofilm, swarming

motility and pyocyanin production phenotypes of the Δ*fadD1* strain (Fig. 3j-l). We then used the

HDOCK server to automatically predict the interaction between FadD1 and the DNA-binding site 5'-

AGGACGG-3'. The binding model of DNA and FadD1 and the detailed contacts of the complex are

shown in Supplementary Fig. 10 and Supplementary Table 3, respectively. The binding sites of FadD1

are in the leucine zipper structure. The results are consistent with the conclusion that the DNA-binding

motif of FadD1 is the leucine zipper.

FadD1 is a receptor of *cis*-DA

Since *cis*-DA is a DSF-type QS signal in *P. aeruginosa*, and the FadD1 controls QS-regulated

phenotypes as a global transcriptional regulator in *P. aeruginosa*, we then measured whether there is

an interaction between *cis*-DA and FadD1. To test this hypothesis, we purified FadD1 to performed

microscale thermophoresis (MST) analysis to test whether FadD1 binds *cis*-DA. As shown in Fig. 4a,

FadD1 exhibited binding activity to *cis*-DA, with an estimated dissociation constant (K_D) of 1.06 ± 0.14

μM. To further confirm FadD1 is a receptor of *cis*-DA, we generated the *cis*-DA deficient mutant Δ*dspI*,

the double deletion mutants Δ*dspI*Δ*fadD1* and *in trans* expressed *fadD1* in the Δ*dspI* and measured

the motility activity and pyocyanin of these strains. The results showed that *in trans* expression of

fadD1 rescued the defective motility and pyocyanin phenotypes of the Δ*dspI* strain (Fig. 4b, c).

Interestingly, we found that the addition of exogenous *cis*-DA rescued the impaired motility and

pyocyanin of Δ*dspI* (Fig. 4b, c); however, addition of exogenous *cis*-DA showed no effect on these

phenotypes of the mutant Δ*fadD1* and Δ*dspI*Δ*fadD1* (Fig. 4b, c). We then measured the gene

expression levels of *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and *pqsA* in the wild-type strain and the Δ*dspI* strain.

The results indicated that deletion of *dspI* caused a significant decrease in the gene expression levels

of *las*, *rhl* and *pqs* QS system genes (Fig. 4d-i). Therefore, we measured and compared the production

of 3-oxo-C12-HSL, C4-HSL and PQS in the wild-type, *dspI* mutant, and complemented strains. The

result showed that the production of 3-oxo-C12-HSL, C4-HSL and PQS was reduced in the *dspI*

mutant strain (Fig. 4j-l). These results demonstrated that both *cis*-DA and FadD1 positively regulates

the *las*, *rhl* and *pqs* QS systems in *P. aeruginosa*.

cis-DA enhances the binding of FadD1 to target gene promoters

To determine how the binding of *cis*-DA to FadD1 might affect the activity of FadD1, we examined the effects of *cis*-DA on the binding of FadD1 to the *lasR* promoter by EMSA. As shown in Fig. 5a, the binding of FadD1 to the promoter probe of *lasR* was enhanced when *cis*-DA was present in the reaction mixtures, and the amount of probe bound to FadD1 increased as the amount of *cis*-DA increased (Fig. 5a).

Then, we attempted to identify the *cis*-DA-binding sites in FadD1. Autodocking analysis revealed 2 amino acid residues, Gly488 (G488) and Arg552 (R552), that might be critical for the interaction between FadD1and *cis*-DA (Fig. 5b). To test whether the roles of these acid residues are related to the binding to *cis*-DA, we then generated two single point mutants (FadD1^{G488A} and FadD1^{R552A}) (Fig. 5c). MST analysis showed that the mutant FadD1^{G488A} and FadD1^{R552A} did not bind to *cis*-DA (Fig. 5d, e). Consistent with this, EMSA analysis showed that mutation of the *cis*-DA-binding site (Gly488 and Arg552) completely eliminated the effect of *cis*-DA on FadD1 (Fig. 5f, g). In addition, *in trans* expression of *fadD1*^{G488A} and *fadD1*^{R552A} only partially rescued motility and pyocyanin (Fig. 5h, i), and the addition of exogenous 20 μM *cis*-DA showed no effect on these phenotypes of the mutant Δ*dspl*Δ*fadD1*(*fadD1*^{G488A}) and Δ*dspl*Δ*fadD1*(*fadD1*^{R552A}) (Fig. 5h, i). Thus, we concluded that FadD1 is a specific response regulator of *cis*-DA in *P. aeruginosa*.

Reconstruction of FadD2 with the leucine zipper motif

Our study found that the DNA binding motif of FadD1 is a leucine zipper motif. Therefore, we used the protein motif analysis website (https://www.novopro.cn/tools/motifscan.html) to analyze whether the other FadD proteins, FadD2-6, contain a leucine zipper structure. The results showed that the other FadD proteins did not contain a leucine zipper structure, and EMSA analysis showed that FadD2, which shares about 59.82% amino acid identity with FadD1, did not bind to the *lasR* gene promoter (Supplementary Fig. 11a, b). To further investigate the role of the leucine zipper motif of FadD1, we reconstructed FadD2 by adding the leucine zipper structure of FadD1 to the same part of FadD2 (named FadD2$_{LZ}$) and tested whether the reconstructed FadD2 has regulatory activity similar to that of FadD1. Therefore, FadD2$_{LZ}$ was purified to homogeneity using affinity chromatography (Supplementary Fig. 11c), and assayed whether it could bind to the *lasR* promoter. Intriguingly, we found that the *lasR* promoter DNA fragment formed stable DNA-protein complexes with FadD2$_{LZ}$ and

migrated more slowly than unbound probes. The amount of labeled probe that bound to FadD2$_{LZ}$ increased with increasing amounts of FadD2$_{LZ}$ but decreased in the presence of a 50-fold greater concentration of unlabeled probe (Supplementary Fig. 11d). Interestingly, addition of exogenous *cis*-DA also enhanced the binding of FadD2$_{LZ}$ to the *lasR* promoter probe at a final concentration of 20 μM (Supplementary Fig. 11e). In addition, *in trans* expression of the *fadD2$_{LZ}$* gene rescued the defective swarming motility and pyocyanin production phenotypes of the Δ*fadD1* strain (Supplementary Fig. 11f, g). And we also found that the addition of exogenous *cis*-DA rescued the impaired motility and pyocyanin of Δ*dspI*Δ*fadD1*(*fadD2$_{LZ}$*) (Supplementary Fig. 11f, g). These results showed that the leucine zipper motif plays a critical role in the regulatory activity of FadD1.

Analysis of the catalytic motif of FadD1

Previous studies reported that fatty acyl-CoA ligase plays a crucial role in catalyzing the formation of fatty acyl-CoA through hydrolysis of pyrophosphate[35,36]. To test whether FadD1 can function in the uptake and activation steps of fatty acid degradation, we evaluated its enzymatic activity *in vitro*. The results showed that FadD1 had high activity on lauric acid (C12:0), while it displayed moderate activity on BDSF but no activity on *cis*-DA (Supplementary Fig. 12a). In addition, the previous study found that mutation of Trp433, Thr436 and Arg453, the substrate binding sties, results in complete inactivation of fatty acyl-CoA synthetase in *E. coli* (37). We continued to generated three single point mutants (FadD1^{W434A}, FadD1^{T437A} and FadD1^{R454A}) to explore whether the mutation affects the enzyme activity of FadD1 (Supplementary Fig. 12b). Similarly, it was shown that the FadD1^{W434A}, FadD1^{T437A} and FadD1^{R454A} exerted no enzyme activity on *cis*-DA, BDSF and lauric acid (C12:0) (Supplementary Fig. 12c-e). To further characterize the relationship between enzyme activity of FadD1 and perception of *cis*-DA, we chose to test the regulatory activity of the mutant FadD1^{R454A}. We *in trans* expressed the *fadD1^{R454A}* in the Δ*dspI*Δ*fadD1* mutant strain. Interestingly, it was found that addition of exogenous *cis*-DA rescued the impaired motility of Δ*dspI*Δ*fadD1*(*fadD1^{R454A}*) (Supplementary Fig. 12f). Consistent with this, EMSA analysis showed that mutation of the substrate binding stie (Arg454) would not affect the perception of *cis*-DA (Supplementary Fig. 12g).

Homologs of FadD1 are widespread in bacteria

To determine whether FadD1 is widely distributed, homologs of FadD1 were sought in the genome
10

database by using the BLAST program. It was indicated that homologs of FadD1 are distributed across many different bacterial species, including in the genera *Pseudomonas*, *Azotobacter*, *Azomonas*, and *Halopseudomonas* (Supplementary Table 4). Furthermore, the homologs of FadD1 in the genera *Pseudomonas, Azotobacter* and *Azomonas* contain a leucine zipper structure (Supplementary Table 4). We then selected *P. fluorescens* Migula ATCC17518, which contains both a FadD1$_{Pf}$ protein with a leucine zipper structure and a *lasR$_{Pf}$* gene, and the FadD1$_{Pf}$ protein was purified by affinity chromatography and tested for its interaction with the *lasR$_{Pf}$* gene promoter. The EMSA results showed that the FadD1$_{Pf}$ protein from *P. fluorescens* could also bind to the *lasR$_{Pf}$* gene promoter (Supplementary Fig. 13), indicating that the transcriptional regulatory mechanism of FadD1 might be widely conserved in bacteria.

Discussion

Fatty acid-CoA ligase plays a vital role in fatty acid metabolism, as it can convert fatty acids to fatty acyl-CoA in bacteria[37]. In this study, we found that the fatty acid-CoA ligase FadD1 of *P. aeruginosa* not only exhibits high activity for the catalysis of lauric acid (C12:0) to lauric acid-CoA (Supplementary Figure. 12a), and employs a key role in the bacterial competition with *B. cenocepacia* by degrading its QS signal BDSF (Fig. 1), but also plays a critical role in the regulation of *P. aeruginosa* physiology and pathogenesis by acting as the receptor and response regulator of its QS signal *cis*-DA (Fig. 2 and Supplementary Table 1). We demonstrated that FadD1 can directly bind to the promoter of the target genes, thereby regulate the transcription levels of the target genes (Fig. 3a and Supplementary Fig. 7). It was further identified that mutation of the leucine zipper structure of FadD1 abolished the binding of FadD1 to the target gene promoters (Fig. 3i). Interestingly, we found that FadD2 does not contain a leucine zipper structure, and EMSA analysis showed that FadD2 did not bind to the *lasR* gene promoter (Supplementary Fig. 11b). However, the reconstructed FadD2 with addition of the leucine zipper structure of FadD1 to the same part of FadD2 (named FadD2$_{LZ}$) obtained the regulatory activity similar to that of FadD1. It could bind to the *lasR* gene promoter and restore the *fadD1* mutant phenotypes (Supplementary Fig. 11). Furthermore, our findings indicate that FadD1 is widely conserved in many other bacterial species, including *Pseudomonas*, *Azotobacter*, and *Azomonas* species (Supplementary Table 4), indicating that FadD1 with a leucine zipper structure might be a new family of transcriptional regulatory proteins widely conserved in bacteria.

The fatty acid molecule *cis*-DA was first identified in *P. aeruginosa* to act as the autoinducer of biofilm dispersion[31]. It has also been shown to induce biofilm dispersion in a range of Gram-negative and Gram-positive bacteria and in the fungal pathogen *Candida albicans*[31]. Although the previous study reported that the two-component sensor/response regulator hybrid DspS (PA4112, PA14_10770) is required and essential for native biofilm dispersion in response to the cell-to-cell signaling molecule *cis*-DA[34], there was no exploration of other functions of *cis*-DA. In this study, we found that deletion of *dspI*, which is the synthase of *cis*-DA, caused an impairment in motility and pyocyanin production (Fig. 4b, c). Interestingly, in *trans* expression of *fadD1* rescued the defective motility and pyocyanin phenotypes of the Δ*dspI* strain (Fig. 4b, c). However, the addition of exogenous *cis*-DA showed no effect on these phenotypes of the mutant Δ*fadD1* and Δ*dspI*Δ*fadD1* (Fig. 4b, c). Moreover, *cis*-DA is shown to bind to FadD1 with high affinity and enhanced the binding of FadD1 to the target gene promoter probes (Fig. 4a, 5a). We also identified that two amino acid residues at positions Gly488 (G488) and Arg552 (R552) are critical for the interaction between FadD1 and *cis*-DA (Fig. 5). Mutations at Gly488 and Arg552 abolished the binding between FadD1 and *cis*-DA (Fig. 5d, e). Another interesting observation in this study was that mutation at the substrate binding stie Arg454 (R454) completely abolished the enzyme activity of FadD1 on both BDSF and lauric acid (C12:0) (Supplementary Fig. 12c-e), but could not affect the effect of *cis*-DA on FadD1 (Supplementary Fig. 12g), suggesting that the *cis*-DA-binding site is different from the substrate-binding stie in FadD1. Together, FadD1 was identified to be a new response regulator of *cis*-DA QS signal.

BDSF has been identified to be a DSF-family QS signal in *B. cenocepacia*[11]. As *cis*-DA shares a similar chemical structure with that of BDSF, we then continued to explore the interaction between BDSF and FadD1. Although FadD1 showed a degradation activity on BDSF, we found that FadD1 tightly binds to BDSF with an estimated dissociation constant (K_D) of 1.65 ± 0.21 μM (Supplementary Fig. 14a). Interestingly, the addition of exogenous BDSF also enhanced the binding of FadD1 to the *lasR* promoter probe at a final concentration of 10 μM (Supplementary Fig. 14b). To distinguish the role of BDSF to be a "substrate" or "ligand" of FadD1, we then used EMSA analysis and the result showed that mutation of the *cis*-DA-binding sites abolished the perception of *cis*-DA (Fig. 5), but will not affect the effect of BDSF on FadD1 (Supplementary Fig. 14c). However, EMSA analysis showed that mutation of the substrate-binding stie (Arg454) completely abolished the effect of BDSF on the binding of FadD1 to target gene promoter (Supplementary Fig. 14d), while not affect the effect of *cis*-

DA on FadD1 (Supplementary Fig. 12g). These results suggested that BDSF also interferes with the binding of FadD1 to target gene promoters *in vitro* as a substrate of the canonical enzyme FadD1.

In *P. aeruginosa,* there are at least three QS systems, namely, *las*, *rhl* and *pqs*, which are involved in the regulation of a broad range of genes important for metabolism and virulence[18,19]. These QS systems are hierarchically interrelated, and the *las* system was verified to monitor both the *rhl* and *pqs* systems[17,30]. In this study, we identified that FadD1 affects the QS-regulated phenotypes, including the biofilm formation, swarming, and pyocyanin in *P. aeruginosa* (Fig. 2a-c). Surprisingly, we also identified that FadD1 is a *cis*-DA receptor protein that positively regulates the *las*, *rhl* and *pqs* QS systems in *P. aeruginosa* (Fig. 2, 6). These results indicated that FadD1 is not only a response regulator of *cis*-DA but also an important upstream regulator of the QS signaling network in *P. aeruginosa*. FadD1 regulates numerous target genes and physiological functions by monitoring the QS signaling systems, which broadens our understanding of the QS signaling network of *P. aeruginosa*.

In summary, our findings suggest that FadD1, which contains a leucine zipper structure in the AMP-B domain, is not only an enzyme of fatty oxidation but also a receptor of *cis*-DA that directly controls target gene expression and controls the QS network in *P. aeruginosa*. In addition, the BLAST search revealed that homologs of FadD1 containing the leucine zipper structure are highly conserved in many bacteria (Supplementary Table 4), suggesting that the regulatory mechanism of FadD1 might be present in various bacterial species. Consistently, our recent study revealed that the fatty acyl-CoA ligase DsfR (BCAM2136), which contains a leucine zipper motif, acts as a global transcriptional regulator to control *B. cenocepacia* virulence by sensing BDSF[47]. Different from FadD1, DsfR has a specific BDSF-sensing site and showed no enzyme activity on BDSF, while FadD1 only binds with BDSF on the substrate-binding sites. These results indicate that FadD1 and DsfR share some common characteristics and are distinguished from each other. In conclusion, our work presents a unique and widely conserved signal receptor of *cis*-DA, which might be an important new type of QS signal receptor in bacteria.

Methods

Bacterial strains and growth conditions.

The bacterial strains and plasmids used in this work are listed in Supplementary Table 5. *P. aeruginosa*

PAO1, *P. fluorescens* Migula ATCC17518 and *B. cenocepacia* H111 strains obtained from the

American Type Culture Collection (ATCC) were maintained in Luria-Bertani broth (LB; 10 g/L tryptone,

5 g/L yeast extract (#PL0021, oxoid), 5 g/L NaCl (#S805275, Macklin)) at 37°C. The following

antibiotics were added when necessary: ampicillin (#A830931, Macklin) and kanamycin (#K885955,

Macklin) at 100 µg mL^{-1}; tetracycline (#T829835, Macklin), 10 µg mL^{-1}; and gentamicin (#G810322,

Macklin), 10 µg mL^{-1}. BDSF (#C957977, Macklin) and *cis*-DA (#Z911073, Macklin) (HPLC ≥ 99%)

were dissolved in methanol at a final concentration of 100 mM, and this solution was added to the

medium in the experiments. Bacterial growth was determined by measuring the optical density at 600

nm.

BSDF extraction

401 The extraction of BDSF from culture supernatants was conducted following previously described

402 methods[11,53]. To quantify BDSF production in the culture of *B. cenocepacia* strain or *P. aeruginosa*

403 strain, 100 mL of the supernatant was collected. Its crude ethyl acetate (#E809178, Macklin) extract

404 was passed through a 0.22 µm Minisart filter unit (#16541-K, Sartorius) and was then condensed to

405 1 mL for LC-MS analysis.

Quantification of BDSF and *cis*-DA

Quantification of BDSF and *cis*-DA was performed by using an LC-MS system, which consisted of an

ACQUITY UPLC system and a Waters Q-Tof Premier high-resolution mass spectrometer[53]. An

ACQUITY UPLC BEH C18 column 1.7 µm (2.1 × 50 mm) was used for the chromatography analysis

of BDSF and *cis*-DA, which was eluted with a CH$_3$OH (methyl alcohol, #A452, ThermoFisher) gradient

in water at 65-100% supplemented with 0.01% formic acid at a flow rate of 0.4 mL·min^{-1} for 10 min.

Next, 65% CH$_3$OH was used for 3 min. The entire elution column was introduced into the Q-TOF mass

spectrometer according to the manufacturer's instructions. The BDSF and *cis*-DA levels in the culture

supernatant were measured using the peak area in the extracted ion chromatogram.

Protein expression and purification

418 The coding regions of proteins were amplified with the primers listed in Supplementary Table 6, and

419 were attached and fused to the expression vectors pET-21a, pET-28a and PDBHT2. The fusion gene

constructs were transformed into *E. coli* strain BL21. Affinity purification of the 6×His fusion proteins was performed using Ni-NTA resin (#SA035010, Smart-lifesciences). The His-tag was removed using TEV protease (#P2307, Beyotime), and the cleaved fusion protein was analyzed by SDS-PAGE[54,55].

Enzyme activity assays

The activity of FadD1 was determined following the methods[56]. *cis*-DA, BDSF or lauric acid (C12:0) were dissolved in TME buffer (Tris-HCl (#T917657, Macklin), 60 mM; $MgCl_2$ (#M813763, Macklin), 10 mM; EDTA (#E885215, Macklin), 1 mM; coenzyme A (CoASH) (#C824132, Macklin), 0.5 mM; and Triton X-100 (#T6328, Macklin), 0.1%; pH 7.5) and added to the mixture at a final concentration of 200 μM, while the final concentrations of FadD1 and its derivatives were 20 μM. The reaction mixture was incubated at 37°C, and the reaction was stopped by placing sample tubes in boiling water for 5, 15, 30 and 60 min. *cis*-DA, BDSF or lauric acid (C12:0) levels were measured by LC-MS spectrometry.

Construction of in-frame deletion mutants and complementation

P. aeruginosa PAO1 was used as the parental strain for the generation of an in-frame deletion mutant of *fadD1* and *dspI* by the following method described previously[14]. The primers used to generate upstream and downstream regions flanking *fadD1* and *dspI* are listed in Supplementary Table 7. For complementation analysis, the coding regions of *fadD1* and *dspI* were amplified by PCR and cloned into the plasmid pBBR1-MCS5, pBBR1-MCS2 and pLAFR3 using the primers listed in Supplementary Table 6. The resulting constructs were conjugated into the *P. aeruginosa* PAO1 mutant, *B. cenocepacia* H111 strain by electroporation[14,54].

Bacterial growth analysis

An overnight bacterial culture in LB medium was inoculated into fresh media to an OD_{600} of 0.01. A 10 mL cell suspension was grown in 50 mL conical centrifuge tubes at 37°C with shaking at 220 rpm. Bacterial growth was determined by measuring the optical density at 600 nm every 4 hours. LB medium was used as the negative control[47].

Competition assays in the mixed culture

The green fluorescent protein expression vector was used and transformed into the *P. aeruginosa*

PAO1 WT, ΔfadD1 and ΔfadD1(fadD1) strains. The mCherry fluorescent protein expression vector

was introduced into the *B. cenocepacia* H111 strain by electroporation. Then, the *P. aeruginosa* PAO1

WT, ΔfadD1 and ΔfadD1(fadD1) strains were cocultured with the *B. cenocepacia* H111 strain at an

initial ratio of 1:4 (vol/vol) at OD_{600} = 0.1 at 37°C with shaking at 220 rpm for 12 h. Then, the

quantification of BDSF was performed by using an LC-MS system, and the mixed cultures were

analyzed by a Spectra Max i3x multifunctional enzyme labeling instrument (Molecular Devices, CA,

USA)[57].

***P. aeruginosa* phenotype assays**

For the analysis of biofilm formation[58], bacterial cells were grown overnight at 37°C and inoculated

into LB media (150 μL per well) in 96-well polypropylene microtiter plates. Microtiter plates with fitted

lids were incubated at 37°C with shaking at 200 rpm for 12 h. The plates were then washed to remove

planktonic cells and stained for 20 min with 1% (weight/vol) crystal violet (#C6158, Sigma-Aldrich).

After washing with water, 200 μL of 95% (vol/vol) ethanol was added to the wells to release the stain.

Biofilm was determined by measuring the absorbance of the resulting solution at 570 nm.

Motility was determined on 0.3% semisolid agar (#BD214010, Becton, Dickinson and Company).

Bacteria were inoculated into the center of plates containing 0.8% tryptone, 0.5% glucose, and 0.3%

agar. The plates were incubated at 30°C for 18 h before the diameter of the colony was measured[54].

The pyocyanin assay was performed with the method described previously[49]. Briefly, bacteria were

cultured at 37°C for approximately 12 hours and centrifuged at 13,000 rpm for 1 min, and 1.5 mL

supernatants were collected and extracted with double volume chloroform with vigorous shaking at

room temperature for 30 min. The solvent phase was transferred to a new tube containing 1 mL of 1

N HCl (#H399545, Sigma-Aldrich). The mixture was shaken gently to transfer pyocyanin to the

aqueous phase. The quantity of pyocyanin was determined by measurement of absorbance at 520

nm and normalization against the cell density.

***B. cenocepacia* phenotype assays**

Biofilm formation in 96-well polypropylene microtiter dishes was assayed essentially as described

previously by *P. aeruginosa*. Swarming motility was determined on semisolid agar (0.3%). Bacteria

were inoculated into the center of plates containing 0.8% tryptone, 0.5% glucose, and 0.3% agar. The

plates were incubated at 30°C for 18 h before the diameter of the colony was measured.

Protease assays were performed following a previously described method[14]. Briefly, bacteria were

cultured at 37°C for approximately 12 hours. Cultures were centrifuged at 13,000 rpm for 5 min, and

the supernatants were removed and filtered through a 0.22 μm membrane. One hundred microliters

of supernatant was incubated at 30°C with an equal volume of azocasein (#A875599, Macklin)

dissolved in proteolytic buffer (5 mg/mL) for 30 min. The reaction was stopped by the addition of 400

μL of 10% (weight/vol) TCA (#T818877, Macklin) buffer. After incubation for 2 min at room temperature,

the mixture was centrifuged at 13,000 rpm for 1 min to remove the remaining azocasein. Supernatants

were removed and mixed with 700 μL of 525 mM NaOH (#S817977, Macklin). The absorbance of the

azopeptide supernatant was measured at a wavelength of 442 nm. Protease activity was obtained

after normalization of the absorbance against the corresponding cell density.

Cytotoxicity assays

Cytotoxicity was assessed by measuring the release of LDH from human A549 cells[54]. The A549 cell

line was cultured in Dulbecco's modified Eagle's medium (#D6429, Sigma-Aldrich) at 37°C with 5%

CO_2. A549 cells were seeded in 96-well plates at a density of 5×10^4 cells per well and grown until

they reached ~90% confluence. Culture supernatants were removed, and the cell monolayer was

washed once with phosphate-buffered saline (PBS). Fresh bacterial cells cultured to an OD_{600} of 1.0

were washed and diluted in DMEM. The diluted bacterial cells were then added to the A549 cell

monolayers at a multiplicity of infection (MOI) of approximately 1000. After incubation for 8 h, the

cytotoxicity was determined by measuring the released lactate dehydrogenase (LDH) in the

supernatants using a cytotoxicity detection kit (#4744926001, Roche).

Construction of reporter strains and measurement of β-galactosidase activity

The promoters of *lasR, lasI, rhlR*, *rhlI, mvfR* and *pqsA* were amplified using the primer pairs listed in

Supplementary Table 6. The resulting products were inserted upstream of the promoter less *lacZ* gene

in the vector pME2-*lacZ*. Transconjugants were then selected on LB agar plates supplemented with

tetracycline and X-gal (#X8050, Solarbio). Measurement of β-galactosidase activities was performed

following previously described methods[14,54,55].

Quantitative reverse transcription PCR assays

Bacterial cells were cultured at 37°C in LB broth to the logarithmic-growth phase (OD_{600}=1.0). Cells were collected by centrifugation, and total RNA was extracted using an Eastep Super Total RNA Extraction Kit (#LS1040, Promega). cDNA synthesis was performed using a HiScript III 1st Strand cDNA Synthesis Kit (#11141ES60, Yeasen). Primers for RT-qPCR are listed in Supplementary Table 6. The RT-qPCR experiments were performed on a QuantStudio™ 7 RT-qPCR System (Thermo Fisher) using SYBR qPCR Green Master Mix (#11201ES08, Yeasen). The expression of the target genes was normalized to the expression level of 16S RNA. The relative transcript abundance was calculated by the $2^{-\Delta\Delta Ct}$ method[55]. At least three biological replicates were performed per sample.

Quantitative analysis of QS signal production in *P. aeruginosa*

Bacterial cells were grown in LB medium overnight with agitation at 37°C. One liter of culture supernatant was collected by centrifugation and extracted with an equal volume of ethyl acetate. The crude extract (organic phase) was dried using a rotary evaporator and dissolved in methanol. All of the above samples were kept at 4°C until analysis. Ultrahigh-performance liquid chromatography-electrospray ionization tandem mass spectrometry (UHPLC-ESI-MS/MS) was performed in a Shimadzu LC-30A UHPLC system with a Waters C18 column (1.8 µm, 150 × 2.1 mm) and a Shimadzu 8060 QQQ-MS mass spectrometer with an ESI source interface. The mass spectrometer was operated in positive-ion mode. The mobile phase was prepared as 0.1% formic acid/water and acetonitrile[57].

RNA-Seq analysis

RNA was isolated from *P. aeruginosa* strains grown to OD_{600}=1.0 by using the Eastep Super Total RNA Extraction Kit (Promega, Madison, USA). The concentration was measured using a Qubit® RNA Assay Kit in a Qubit® 2.0 Fluorometer (Life Technologies, CA, USA). The purity and integrity were checked using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA) and the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA), respectively. Sequencing libraries were generated using the NEBNext® Ultra™ Directional RNA Library Prep Kit for Illumina® (NEB, USA) and then purified by the AMPure XP system. Finally, library quality was assessed by using the Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples

was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS

(Illumina). After cluster generation, the library preparations were sequenced on an Illumina HiSeq

platform, and paired-end reads were generated. Trimmed sequence reads were aligned to the *P.*

aeruginosa PAO1 genome sequence using Bowtie2-2.2.3, and normalized read counts were

compared using HTSeq v0.6.1 as described previously[59,60]. Then, the FPKM of each gene was

calculated based on the 26 lengths of this gene, and the read count was mapped to the gene.

Electrophoretic gel mobility shift assay

Electrophoretic mobility shift assay (EMSA) experiments were performed as described

previously[53,54,61]. The DNA probes used for EMSA were prepared by PCR amplification using the

primer pairs listed in Supplementary Table 6. The purified PCR products were 3-end-labeled with

biotin following the manufacturer's instructions (#20148, Thermo Fisher). The DNA-protein binding

reactions were performed according to the manufacturer's instructions (#20148, Thermo Fisher). A 5%

polyacrylamide gel was used to separate the DNA-protein complexes. After UV cross-linking, the

biotin-labeled probes were detected in the membrane using a Biotin luminescent detection kit (#20148,

Thermo Fisher).

Chromatin immunoprecipitation sequencing analyses

Chromatin immunoprecipitation sequencing (ChIP-Seq) was performed according to previously

published methods with some modifications[62]. ChIP-Seq analysis was conducted by the Wuhan

IGENEBOOK Biotechnology Co. Ltd. (Wuhan, China). The anti-HIS antibody (ab9108, 1:1000, Abcam)

was used for immunoprecipitation, and DNA fragments enriched by ChIP were sequenced on an

Illumina HiSeq 2000. After removing sequencing adaptors and low-quality bases, the clean reads

were mapped to the *P. aeruginosa* PAO1 genome using BWA software (version 0.7.15-r1140). The

peak caller MACS (version, 2.1.1.20160309; *q* value, 0.05) was used to localize the potential binding

sites of FadD1[63].

Molecular docking FadD1

The 3D structure of the double-stranded DNA was built in the MOE1 DNA/RNA Builder. Protein-DNA

docking in the HDOCK2 server was used for molecular docking simulation of FadD1 and DNA[64,65].

The 3D structure of FadD1 (Q9HYU4) predicted by AlphaFold2 was set as the receptor, and the 3D structure of the DNA was set as the ligand[66]. The HDOCK server automatically predicts their interaction through a hybrid algorithm of template-based and template-free docking. First, the server performs template-based modeling to identify possible homologous DNA templates. Second, the intrinsic scoring function for protein-RNA interactions is integrated in template-free docking. Finally, the top 100 predicted complex structures are provided to users for download, of which the top 10 models will be visualized on the result web page. The intermolecular contacts from the most likely poses were further evaluated. The models of the complex and interface residues were analyzed using MOE. Molecular graphics were generated by PyMOL.

AutoDock and PyMOL were used for docking and binding site analysis. The crystal structure of FadD1 from the AlphaFold2 was used for docking with *cis*-DA by AutoDock software. To screen the potential binding sites, a grid box was prepared to wrap the entire macromolecule. The receptor grid was centered at center_x = -1.038, center_y = 1.49, center_z = 11.31, size_x = 126, size_y = 126, and size_z = 126. The exhaustiveness was set at 50. The top scoring ligand conformation was chosen, and the potential binding sites were visualized in 3D by PyMOL. The potential binding sites were visualized in 2D by Protein Plus (https://proteins.plus/).

Microscale thermophoresis assay

Protein-binding experiments were carried out with a Nano Temper 16 Monolith NT.115 instrument (NanoTemper Technologies; www.nanotemper-technologies.com)[55]. In brief, the protein was labeled with the L014 Monolith NT.115 Protein Labelling Kit (#MO-L018, Nano Temper). The labeled protein and different titers of unlabeled *cis*-DA were mixed and loaded onto standard treated silicon capillaries (#MO-L022, Nano Temper), and fluorescence was measured. The measurements were carried out at 20% LED power and 40% MST power.

Statistical analysis

No statistical method was used to predetermine the sample size. No data were excluded from the analyses. The data are presented as the means ± standard deviations. Statistical analyses were performed with Prism 8 software (GraphPad). Statistical significance is indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ns = no significance (one-way ANOVA or two-way ANOVA). Biological

replicates and numbers of independent experiments were stated in the legends. All experiments presented as representative LC-MS graphs or gels were repeated at least 3 times with similar results.

Reporting summary

Further information on the research design is available in the Nature Research Reporting Summary linked to this article.

Data Availability

RNA-seq and ChIP-seq data generated in this study have been deposited in the National Center for Biotechnology Information Sequence Read Archive (SRA) database under accession code PRJNA1121197 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1121197) and PRJNA1120965 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1120965), respectively. The data supporting the findings of this study are available within the article and Supplementary Information file.

Reference

1. Nealson, K.H. & Hastings, J.W. Bacterial bioluminescence: its control and ecological significance. *Microbiol. Rev.* **43**, 496-518 (1979). doi: 10.1128/mr.43.4.496-518.1979.
2. Fuqua, C. & Greenberg, E.P. Listening in on bacteria: acyl-homoserine lactone signalling. *Nat. Rev. Mol. Cell Biol.* 3, 685-695 (2002). doi: 10.1038/nrm907.
3. Papenfort, K. & Bassler, B.L. Quorum sensing signal-response systems in Gram-negative bacteria. *Nat. Rev. Microbiol.* **14**, 576-588 (2016). doi: 10.1038/nrmicro.2016.89.
4. Whiteley, M., Diggle, S.P. & Greenberg, E.P. Progress in and promise of bacterial quorum sensing research. *Nature* **551**, 313-320 (2017). doi: 10.1038/nature24624.
5. Chadha, J., Harjai, K. & Chhibber, S. Revisiting the virulence hallmarks of *Pseudomonas aeruginosa*: a chronicle through the perspective of quorum sensing. *Environ. Microbiol.* **24**, 2630-2656 (2022). doi: 10.1111/1462-2920.15784.
6. Flavier, A.B., Clough, S.J., Schell, M.A. & Denny, T.P. Identification of 3-hydroxypalmitic acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*. *Mol. Microbiol.* **26**, 251-259 (1997). doi: 10.1046/j.1365-2958.1997.5661945.x.
7. Barber, C.E. *et al.*, A novel regulatory system required for pathogenicity of *Xanthomonas campestris* is mediated by a small diffusible signal molecule. *Mol. Microbiol.* **24**, 555-566 (1997). doi: 10.1046/j.1365-2958.1997.3721736.x.
8. Déziel, E. *et al.*, Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *Proc. Natl Acad. Sci. U.S.A.* **101**, 1339-1344 (2004). doi: 10.1073/pnas.0307694100.
9. Zhang, L.H. & Dong, Y.H. Quorum sensing and signal interference: diverse implications. *Mol. Microbiol.* **53**, 1563-1571 (2004). doi: 10.1111/j.1365-2958.2004.04234.x.

10. Diggle, S.P. *et al.*, The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS
play multifunctional roles in quorum sensing and iron entrapment. *Chem. Biol.* **14**, 87-96 (2007).
doi: 10.1016/j.chembiol.2006.11.014.

11. Boon, C. *et al.*, A novel DSF-like signal from *Burkholderia cenocepacia* interferes with *Candida*
albicans morphological transition. *ISME J.* **2**, 27-36 (2008). doi: 10.1038/ismej.2007.76.

12. He, Y.W. & Zhang, L.H. Quorum sensing and virulence regulation in *Xanthomonas campestris*.
FEMS Microbiol. Rev. **32**, 842-57 (2008). doi: 10.1111/j.1574-6976.2008.00120.x.

13. Deng, Y., Wu, J., Tao, F. & Zhang, L.H. Listening to a new language: DSF-based quorum sensing
in Gram-negative bacteria. *Chem. Rev.* **111**, 160-173 (2011). doi: 10.1021/cr100354f.

14. Deng, Y. *et al.*, *Cis*-2-dodecenoic acid receptor RpfR links quorum-sensing signal perception with
regulation of virulence through cyclic dimeric guanosine monophosphate turnover. *Proc. Natl Acad.*
Sci. U.S.A. **109**, 15479-15484 (2012). doi: 10.1073/pnas.1205037109.

15. Dulcey, C.E. *et al.*, The end of an old hypothesis: the *pseudomonas* signaling molecules 4-
hydroxy-2-alkylquinolines derive from fatty acids, not 3-ketofatty acids. *Chem. Biol.* **20**, 1481-1491
(2013). doi: 10.1016/j.chembiol.2013.09.021.

16. Wang, M., Li, X., Song, S., Cui, C., Zhang, L.H. & Deng, Y. The *cis*-2-Dodecenoic Acid (BDSF)
Quorum Sensing System in *Burkholderia cenocepacia*. *Appl. Environ. Microbiol.* **88**, e0234221
(2022). doi: 10.1128/aem.02342-21.

17. de Kievit, T.R., Kakai, Y., Register, J.K., Pesci, E.C. & Iglewski, B.H. Role of the *Pseudomonas*
aeruginosa las and *rhl* quorum-sensing systems in *rhlI* regulation. *FEMS Microbiol. Lett.* **212**, 101-
106 (2002). doi: 10.1016/s0378-1097(02)00735-8.

18. Lee, J. & Zhang, L.H. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*.
Protein Cell **6**, 26-41 (2015). doi: 10.1007/s13238-014-0100-x.

19. Mukherjee, S., Moustafa, D., Smith, C.D., Goldberg, J.B. & Bassler, B.L. The RhlR quorum-
sensing receptor controls *Pseudomonas aeruginosa* pathogenesis and biofilm development
independently of its canonical homoserine lactone autoinducer. *PLoS Pathog.* **13**, e1006504
(2017). doi: 10.1371/journal.ppat.1006504.

20. Fuqua, W.C., Winans, S.C. & Greenberg, E.P. Quorum sensing in bacteria: the LuxR-LuxI family
of cell density-responsive transcriptional regulators. *J. Bacteriol.* **176**, 269-275 (1994). doi:
10.1128/jb.176.2.269-275.1994.

21. Pesci, E.C., Pearson, J.P., Seed, P.C. & Iglewski, B.H. Regulation of *las* and *rhl* quorum sensing
in *Pseudomonas aeruginosa*. *J. Bacteriol.* **179**, 3127-3132 (1997). doi: 10.1128/jb.179.10.3127-
3132.

22. Withers, H., Swift, S. & Williams, P. Quorum sensing as an integral component of gene regulatory
networks in Gram-negative bacteria. *Curr. Opin. Microbiol.* **4**, 186-193 (2001). doi: 10.1016/s1369-
5274(00)00187-9.

23. Ng, W.L. & Bassler, B.L. Bacterial quorum-sensing network architectures. *Annu. Rev. Genet.* **43**,
197-222 (2009). doi: 10.1146/annurev-genet-102108-134304.

24. Rutherford, S.T. & Bassler, B.L. Bacterial quorum sensing: its role in virulence and possibilities for
its control. *C.S.H. Perspect. Med.* **2**, a012427 (2012). doi: 10.1101/cshperspect.a012427.

25. Jander, G., Rahme, L.G. & Ausubel, F.M. Positive correlation between virulence of *Pseudomonas*
aeruginosa mutants in mice and insects. *J. Bacteriol.* **182**, 3843-3845 (2000). doi:
10.1128/JB.182.13.3843-3845.

26. Cao, H., Krishnan, G., Goumnerov, B., Tsongalis, J., Tompkins, R. & Rahme, L.G. A quorum

sensing-associated virulence gene of *Pseudomonas aeruginosa* encodes a LysR-like
transcription regulator with a unique self-regulatory mechanism. *Proc. Natl Acad. Sci. U.S.A.* **98**,
14613-14618 (2001). doi: 10.1073/pnas.251465298.

27. Wade, D.S. *et al.*, Regulation of *Pseudomonas* quinolone signal synthesis in *Pseudomonas aeruginosa*. *J. Bacteriol.* **187**, 4372-4380 (2005). doi: 10.1128/JB.187.13.4372-4380.

28. Déziel, E. *et al.*, The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation: multiple quorum sensing-regulated genes are modulated without affecting *lasRI*, *rhlRI* or the production of N-acyl-L-homoserine lactones. *Mol. Microbiol.* **55**, 998-1014 (2005). doi: 10.1111/j.1365-2958.2004.04448.x.

29. García-Reyes, S., Soberón-Chávez, G. & Cocotl-Yanez, M. The third quorum-sensing system of *Pseudomonas aeruginosa*: *Pseudomonas* quinolone signal and the enigmatic PqsE protein. *J. Med. Microbiol.* **69**, 25-34 (2020). doi: 10.1099/jmm.0.001116.

30. Choi, Y. *et al.*, Growth phase-differential quorum sensing regulation of anthranilate metabolism in *Pseudomonas aeruginosa*. *Mol. Cells* **32**, 57-65 (2011). doi: 10.1007/s10059-011-2322-6.

31. Davies, D.G. & Marques, C.N. A fatty acid messenger is responsible for inducing dispersion in microbial biofilms. *J. Bacteriol.* **191**, 1393-1403 (2009). doi: 10.1128/JB.01214-08.

32. Amari, D.T., Marques, C.N. & Davies, D.G. The putative enoyl-coenzyme A hydratase DspI is required for production of the *Pseudomonas aeruginosa* biofilm dispersion autoinducer *cis*-2-decenoic acid. *J. Bacteriol.* **195**, 4600-4610 (2013). doi: 10.1128/JB.00707-13.

33. Rahmani-Badi, A., Sepehr, S., Fallahi, H. & Heidari-Keshel, S. Dissection of the *cis*-2-decenoic acid signaling network in *Pseudomonas aeruginosa* using microarray technique. *Front. Microbiol.* **6**, 383 (2015). doi: 10.3389/fmicb.2015.00383.

34. Kalia, M., Amari, D., Davies, D.G. & Sauer, K. *cis*-DA-dependent dispersion by *Pseudomonas aeruginosa* biofilm and identification of *cis*-DA-sensory protein DspS. *mBio*. **14**, e0257023 (2023). doi: 10.1128/mbio.02570-23.

35. Hume, A.R., Nikodinovic-Runic, J. & O'Connor, K.E. FadD from *Pseudomonas putida* CA-3 is a true long-chain fatty acyl coenzyme A synthetase that activates phenylalkanoic and alkanoic acids. *J. Bacteriol.* **191**, 7554-7565 (2009). doi: 10.1128/JB.01016-09.

36. Ruth, K., de Roo, G., Egli, T. & Ren, Q. Identification of two acyl-CoA synthetases from *Pseudomonas putida* GPo1: one is located at the surface of polyhydroxyalkanoates granules. *Biomacromolecules* **9**, 1652-1659 (2008). doi: 10.1021/bm8001655.

37. Black, P.N., Zhang, Q., Weimar, J.D. & DiRusso, C.C. Mutational analysis of a fatty acyl-coenzyme A synthetase signature motif identifies seven amino acid residues that modulate fatty acid substrate specificity. *J. Biol. Chem.* **272**, 4896-4903 (1997). doi: 10.1074/jbc.272.8.4896.

38. Bi, H., Yu, Y., Dong, H., Wang, H. & Cronan, J.E. *Xanthomonas campestris* RpfB is a fatty Acyl-CoA ligase required to counteract the thioesterase activity of the RpfF diffusible signal factor (DSF) synthase. *Mol. Microbiol.* **93**, 262-275 (2014). doi: 10.1111/mmi.12657.

39. Zhou, L., Wang, X.Y., Sun, S., Yang, L.C., Jiang, B.L. & He, Y.W. Identification and characterization of naturally occurring DSF-family quorum sensing signal turnover system in the phytopathogen *Xanthomonas. Environ. Microbiol.* **17**, 4646-4658 (2015). doi: 10.1111/1462-2920.12999.

40. Wang, X.Y., Zhou, L., Yang, J., Ji, G.H. & He, Y.W. The RpfB-Dependent Quorum Sensing Signal Turnover System Is Required for Adaptation and Virulence in Rice Bacterial Blight Pathogen *Xanthomonas oryzae* pv. oryzae. *Mol. Plant Microbe. Interact.* **29**, 220-230 (2016). doi: 10.1094/MPMI-09-15-0206-R.

41. Tian, X.Q., Wu, Y., Cai, Z. & Qian, W. BDSF is a degradation-prone quorum-sensing signal
detected by the histidine kinase RpfC of *Xanthomonas campestris* pv. *campestris*. *Appl. Environ.*
Microbiol. **88**, e0003122 (2022). doi: 10.1128/aem.00031-22.

42. Maston, G.A., Evans, S.K. & Green, M.R. Transcriptional regulatory elements in the human
genome. *Annu. Rev. Genomics. Hum. Genet.* **7**, 29-59 (2006). doi:
10.1146/annurev.genom.7.080505.115623.

43. Wang, G., Wang, F., Huang, Q., Li, Y., Liu, Y. & Wang, Y. Understanding Transcription Factor
Regulation by Integrating Gene Expression and DNase I Hypersensitive Sites. *Biomed. Res. Int.*
2015, 757530 (2015). doi: 10.1155/2015/757530.

44. Clark, D.P. & Pazdernik, N.J. Chapter e14 - Protein Structure and Function, *Mol. Biol. (Second*
Edition), 2013, Pages e280-e284, https://doi.org/10.1016/B978-0-12-378594-7.00048-2.

45. Miller, M. The importance of being flexible: the case of basic region leucine zipper transcriptional
regulators. *Curr. Protein Pept. Sci.* **10**, 244-269 (2009). doi: 10.2174/138920309788452164.

46. Maxon, M.E., Wigboldus, J., Brot, N. & Weissbach, H. Structure-function studies on *Escherichia*
coli MetR protein, a putative prokaryotic leucine zipper protein. *Proc. Natl Acad. Sci. U.S.A.* **87**,
7076-7079 (1990). doi: 10.1073/pnas.87.18.7076.

47. Li, X. *et al.*, Regulation of *Burkholderia cenocepacia* virulence by the fatty acyl-CoA ligase DsfR
as a response regulator of quorum sensing signal. *Cell Rep.* **43**, 114223 (2024). doi:
10.1016/j.celrep.2024.114223.

48. Deng, Y., Boon, C., Eberl, L. & Zhang, L.H. Differential modulation of *Burkholderia cenocepacia*
virulence and energy metabolism by the quorum-sensing signal BDSF and its synthase. *J.*
Bacteriol. **191**, 7270-7278 (2009). doi: 10.1128/JB.00681-09.

49. Deng, Y., Wu, J., Eberl, L. & Zhang, L.H. Structural and functional characterization of diffusible
signal factor family quorum-sensing signals produced by members of the *Burkholderia cepacia*
complex. *Appl. Environ. Microbiol.* **76**, 4675-4683 (2010). doi: 10.1128/AEM.00480-10.

50. Deng, Y., Boon, C., Chen, S., Lim, A. & Zhang, L.H. *Cis*-2-dodecenoic acid signal modulates
virulence of *Pseudomonas aeruginosa* through interference with quorum sensing systems and
T3SS. *BMC Microbiol.* **13**, 231 (2013). doi: 10.1186/1471-2180-13-231.

51. Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. Basic local alignment search tool.
J. Mol. Biol. **215**, 403-410 (1990). doi: 10.1016/S0022-2836(05)80360-2.

52. Zarzycki-Siek, J. *et al.*, Elucidating the *Pseudomonas aeruginosa* fatty acid degradation pathway:
identification of additional fatty acyl-CoA synthetase homologues. *PLoS One* **8**, e64554 (2013).
doi: 10.1371/journal.pone.0064554.

53. Cui, C. *et al.*, A novel two-component system modulates quorum sensing and pathogenicity in
Burkholderia cenocepacia. *Mol. Microbiol.* **108**, 32-44 (2018). doi: 10.1111/mmi.13915.

54. Yang, C. *et al.*, *Burkholderia cenocepacia* integrates *cis*-2-dodecenoic acid and cyclic dimeric
guanosine monophosphate signals to control virulence. *Proc. Natl Acad. Sci. U.S.A.* **114**, 13006-
13011 (2017). doi: 10.1073/pnas.1709048114.

55. Guo, Q. *et al.*, Elongation factor P modulates *Acinetobacter baumannii* physiology and virulence
as a cyclic dimeric guanosine monophosphate effector. *Proc. Natl Acad. Sci. U.S.A.* **119**,
e2209838119 (2022). doi: 10.1073/pnas.2209838119.

56. Kang, Y., Zarzycki-Siek, J., Walton, C.B., Norris, M.H. & Hoang, T.T. Multiple FadD acyl-CoA
synthetases contribute to differential fatty acid degradation and virulence in *Pseudomonas*
aeruginosa. *PLoS One* **5**, e13557 (2010). doi: 10.1371/journal.pone.0013557.

57. Cui, B. *et al.*, The cell-cell communication signal indole controls the physiology and interspecies communication of *Acinetobacter baumannii*. *Microbiol. Spectr.* **10**, e0102722 (2022). doi: 10.1128/spectrum.01027-22.

58. Song, S. *et al.*, Anthranilic acid from *Ralstonia solanacearum* plays dual roles in intraspecies signalling and inter-kingdom communication. *ISME J.* **14**, 2248-2260 (2020). doi: 10.1038/s41396-020-0682-7.

59. Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357-359 (2012). doi: 10.1038/nmeth.1923.

60. Trapnell, C. *et al.*, Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**, 511-515 (2010). doi: 10.1038/nbt.1621.

61. Song, S. *et al.*, An anthranilic acid-responsive transcriptional regulator controls the physiology and pathogenicity of *Ralstonia solanacearum*. *PLoS Pathog.* **18**, e1010562 (2022). doi: 10.1371/journal.ppat.1010562.

62. Landt, S.G. *et al.*, ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Res.* **22**, 1813-1831 (2012). doi: 10.1101/gr.136184.111.

63. Kleinman, C.L., Sycz, G., Bonomi, H.R., Rodríguez, R.M., Zorreguieta, A. & Sieira, R. ChIP-seq analysis of the LuxR-type regulator VjbR reveals novel insights into the *Brucella* virulence gene expression network. *Nucleic Acids Res.* **45**, 5757-5769 (2017). doi: 10.1093/nar/gkx165.

64. Molecular Operating Environment (MOE), Chemical Computing Group Inc., 1010 Sherbooke St. West, Suite #910, Montreal, QC, Canada, H3A 2R7 (2018).

65. Yan, Y., Tao, H., He, J. & Huang, S.Y. The HDOCK server for integrated protein-protein docking. *Nat. Protoc.* **15**, 1829-1852 (2020). doi: 10.1038/s41596-020-0312-x.

66. Jumper, J. *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021). doi: 10.1038/s41586-021-03819-2.

Acknowledgments

796 This work was financially supported by the National Key Research and Development Program of China (2021YFA0717003 to Y.D.), the National Natural Science Foundation of China (32300033 to S.S.) and the Scientific Research Foundation of Hainan University (KYQD(ZR)-23006 to S.S.).

Author contributions

801 Y.D. and S.S. conceived the project. S.S., X.L., J.L., and Y.D. designed the research. S.S., X.L., J.L., B.W., Y.S., H.H., X.S., M.W., and B.C. performed the research. S.S., X.L., J.L., G.W., Y.H., L.X., B.G., L.Y., X.W., L-H.Z., and Y.D. analyzed the data. S.S. and Y.D. wrote the paper.

Competing interests

806 The authors declare no competing interests.

Figure 1. Effect of *fadD1* on the competitive capability of *P. aeruginosa* against *B. cenocepacia*.

a Analysis of the BDSF signal quenching ability of *P. aeruginosa* PAO1 (n = 3 biological replicates). **b** SDS-PAGE of the purified FadD1 protein. **c** Analysis of the enzymatic quenching activity of FadD1 on BDSF signal (*n* = 3 biological replicates). **d** The BDSF signal production of the *B. cenocepacia* H111 strain cocultured with *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains (*n* = 3 biological replicates). **p=0.0016, ***p<0.001, ***p<0.001. **e** Green mean fluorescence intensity of the *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains carrying the green fluorescent protein expression vector (*n* = 3 biological replicates). *p=0.0306, **p=0.0021, **p=0.0050, ***p<0.001, **p=0.0017, ***p<0.001, ***p<0.001. **f** Red mean fluorescence intensity of the *B. cenocepacia* H111 strain carrying the mCherry fluorescent protein expression vector (*n* = 3 biological replicates). ***p<0.001, **p=0.0016, ***p<0.001, **p=0.0011, ***p<0.001, **p=0.0020. The data are the means ± standard deviations of three independent experiments. The *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains carried the green fluorescent protein expression vector. The mCherry fluorescent protein expression vector was carried by the *B. cenocepacia* H111 strain. The *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains were cocultured with *the B. cenocepacia* H111 strain at a ratio of 1:4 (v/v) at OD_{600} = 0.1. The statistical comparisons were

824 performed using two-way ANOVA. In **b** experiment was performed three times and representative

825 images from one experiment are shown.

Figure 2. Effects of *fadD1* on the QS-regulated phenotypes and QS of *P. aeruginosa*. Analysis

of biofilm formation **a**, swarming motility **b**, pyocyanin production **c**, and virulence **d** in *P. aeruginosa*

PAO1(**a**, *n* = 5 biological replicates; **b**, **c**, **d**, *n* = 3 biological replicates). **a** ***p<0.001, ***p<0.001. **b**

p=0.0025, **p=0.0081. **c ***p<0.001, ***p<0.001. **d** ***p<0.001, ***p<0.001. Effects of *fadD1* on the

gene expression levels of *lasR* **e**, *lasI* **f**, *rhlR* **g**, *rhlI* **h**, *mvfR* **i** and *pqsA* **j** by assessing the β-

galactosidase activity of the promoter-*lacZ* transcriptional fusions and by RT-qPCR **k**, which are the

signal molecule synthase-encoding genes and receptor genes of the *rhl*, *las* and *pqs* QS systems in

P. aeruginosa PAO1, respectively (*n* = 3 biological replicates). **e** ***p<0.001, ***p<0.001, ***p<0.001.

f ***p<0.001, ***p<0.001, ***p<0.001, ***p<0.001. **g** ***p<0.001, ***p<0.001, ***p<0.001. **h** *p=0.025,

***p<0.001, ***p<0.001, ***p<0.001. **i** ***p<0.001, ***p<0.001, ***p<0.001. **j** **p=0.0015, ***p<0.001,

***p<0.001. The production of 3-oxo-C12-HSL **l**, C4-HSL **m** and PQS **n** in the *P. aeruginosa* PAO1

wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains (*n* = 3 biological replicates). The production of QS

850 signals in the *P. aeruginosa* PAO1 wild-type strain was arbitrarily defined as 100%. **l** ***p<0.001,

851 ***p<0.001. **m** ***p<0.001, **p=0.0015. **n** ***p<0.001, ***p<0.001. The data are means ± standard

852 deviations of three independent experiments. The statistical comparisons were performed using one-

853 way ANOVA or two-way ANOVA (**a**, **b**, **c**, **d**, **l**, **m**, **n**, one-way ANOVA; **e**, **f**, **g**, **h**, **i**, **j**, two-way ANOVA).

Figure 3. Analysis of the binding of FadD1 to target gene promoters. a EMSA analysis of *in vitro*

binding of FadD1 to the promoter of *lasR*, in which the biotin-labeled 339-bp *lasR* promoter DNA probe

was used for the protein binding assay. A protein-DNA complex, represented by a band shift, was

formed when different concentrations of FadD1 protein were incubated with the probe at room

temperature for 30 min. **b** A potential FadD1 binding region was identified by ChIP-seq. The

conserved sequence is shown in orange. **c** Analysis of the binding between FadD1 and the mutated

lasR promoter with deletion of the FadD1 binding sequence AGGACGG. EMSA analysis was

performed *in vitro*. **d** Domain structure analysis of FadD1 (https://www.ebi.ac.uk/Tools/hmmer/). The

red box represents the leucine zipper motif (https://www.novopro.cn/tools/motifscan.html) in AMP-B

domain. **e** SDS-PAGE of the purified AMP-B domain and the AMP-B_C domain of the FadD1 protein.

EMSA analysis of *in vitro* binding of the AMP-B domain **f** and the AMP-B_C domain **g** to the promoter

of *lasR*. **h** SDS-PAGE of the purified mutated FadD1 protein (FadD1^{LM}: L275A L282A L289A L296A).

i EMSA analysis of the *in vitro* binding of the mutated FadD1 protein to the promoter of *lasR*. *In trans*

expression of the mutated *fadD1* could not restore biofilm formation **j**, swarming motility **k**, and

pyocyanin production **l** in the *P. aeruginosa* Δ*fadD1* strain (**j**, *n* = 6 biological replicates; **k**, **l**, *n* = 3

biological replicates). **j** ***p<0.001, ***p<0.001, ^{ns}*p*=0.9362. **k** **p=0.0029, **p=0.0026, ^{ns}*p*=0.9919. **l**

***p<0.001, ***p<0.001, ^{ns}*p*>0.9999. The EMSA experiments were performed three times, and

representative images from one experiment are shown. The data are the means ± standard deviations

of three independent experiments. The statistical comparisons were performed using one-way

ANOVA.

Figure 4. Effects of FadD1 on *cis*-DA-regulated phenotypes of *P. aeruginosa*. a MST analysis of the binding of *cis*-DA to the FadD1. "Fnorm (‰)" indicates the fluorescence time trace changes in the MST response. Complementation of the *dspI* mutant with *fadD1*. *In trans* expression of *fadD1* complemented swarming motility **b** and pyocyanin production **c** in the *dspI*-deficient mutant. Effects of *dspI* on the gene expression levels of *lasR* **d**, *lasI* **e**, *rhlR* **f**, *rhlI* **g**, *mvfR* **h**, *pqsA* **i** and the production of 3-oxo-C12-HSL **j**, C4-HSL **k** and PQS **l** of the *P. aeruginosa* PAO1 strains. The production of QS signals in the *P. aeruginosa* PAO1 wild-type strain was arbitrarily defined as 100%. n = 3 biological replicates. **b** ***$p<0.001$, ***$p<0.001$, *$p=0.0181$, **$p=0.0047$, $^{ns}p>0.9999$, $^{ns}p>0.9999$. **c** ***$p<0.001$, ***$p<0.001$, $^{ns}p=0.9984$, $^{ns}p>0.9999$. **d, e, f, g, h, l, j, k, l** ***$p<0.001$. The data are means ± standard deviations of three independent experiments. The statistical comparisons were performed using one-way ANOVA.

Figure 5. Effects of *cis*-DA on the binding of FadD1 to the promoters of the target genes. a
EMSA analysis of the *in vitro* binding of FadD1 to the promoter of *lasR* with the addition of different
amounts of *cis*-DA. **b** The potential residues (which are shown as stick models and are labeled) of
FadD1 involved in *cis*-DA binding analyzed by AutoDocking software. The hydrogen bonds were
represented by yellow dashed lines. **c** SDS–PAGE gel electrophoresis of purified FadD1 variants.
MST analysis of the binding of BDSF to the **d** FadD1^{G488A} and **e** FadD1^{R552A}. EMSA analysis of the *in*
vitro binding of **f** FadD1^{G488A} and **g** FadD1^{R552A} to the promoters of *lasR* with the addition of different
amounts of *cis*-DA. The EMSA experiments were performed three times, and representative images
from one experiment are shown. Analysis of the perception of *cis*-DA by FadD1^{G488A} and FadD1^{R552A},
cis-DA-regulated phenotypes: **h** swarming motility and **i** pyocyanin production in *P. aeruginosa* (n = 3
biological replicates). **h** ***p<0.001, ***p<0.001, ^{ns}*p*>0.9999, ^{ns}*p*>0.9999. **i** ***p<0.001, ***p<0.001,
^{ns}*p*=0.9996, ^{ns}*p*>0.9999. The data are means ± standard deviations of three independent experiments.
The statistical comparisons were performed using one-way ANOVA.

Figure 6. Schematic representation of *cis*-DA-regulated QS signaling in *P. aeruginosa*. DspI
synthesized *cis*-DA signal molecule. FadD1, the signal receptor of *cis*-DA, perceives *cis*-DA signal
and regulates the transcription of *lasR*, which positively regulates the *las*, *rhl* and *pqs* QS systems in
P. aeruginosa. DspS is also the sensor of *cis*-DA involved in biofilm dispersion in response to *cis*-DA.
The individual QS circuits are highly interconnected.