The long-chain fatty acid-CoA ligase FadD1 monitors *Pseudomonas aeruginosa* quorum sensing as a receptor of *cis*-2-decenoic acid

Shihao Song[1,2,11], Jingyun Liu[2,3,11], Bing Wang[1], Yang Si[4], Hongguang Han[1], Xiuyun Sun[1,5], Mingfang Wang[5], Binbin Cui[5,6], Guangliang Wu[6], Yongliang Huo[7], Liangxiong Xu[8], Beile Gao[9], Liang Yang[10], Xiaoxue Wang[9], Lian-Hui Zhang[2], Yinyue Deng[5,*]

[1]Key Laboratory of Tropical Biological Resources of Ministry of Education, School of Pharmaceutical Sciences, Hainan University, Haikou 570228, China

[2]Integrative Microbiology Research Center, South China Agricultural University, Guangzhou 510642, China

[3]Zhengzhou Shuqing Medical College, Zhengzhou 450064, China

[4]Department of Microbiology, College of Life Sciences, Nanjing Agricultural University, Nanjing 210095, China

[5]School of Pharmaceutical Sciences (Shenzhen), Shenzhen Campus of Sun Yat-sen University, Sun Yat-sen University, Shenzhen 518107, China

[6]Pharmacy Department, The Affiliated LiHuiLi Hospital of Ningbo University, Ningbo315046, China

[7]Experimental Animal Center, Guangzhou Medical University, Guangzhou, Guangdong, 511436, China

[8]School of Life Sciences, Huizhou University, Huizhou 516007, China

[9]Guangdong Key Laboratory of Marine Materia Medica, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China

[10]School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.

[11]These authors contributed equally: Shihao Song and Jingyun Liu.

Corresponding author:

Yinyue Deng: dengyle@mail.sysu.edu.cn

Running title: *Cis*-2-decenoic acid controls *P. aeruginosa* virulence

Highlights

The long-chain fatty acid-CoA ligase FadD1 acts as a global transcriptional regulator in *P. aeruginosa*.

FadD1 is a receptor of QS signal *cis*-DA and functions at the top of the QS hierarchy in *P. aeruginosa*.

The homolog of FadD1 in *P. fluorescens* could bind to the target gene promoter by responding to *cis*-DA.

In brief

Pseudomonas aeruginosa is a human pathogen with antibiotic resistance and a wide range of dynamic defenses makes it an extremely challenging organism to treat in modern-day medicine. It employs *cis*-2-decenoic acid (*cis*-DA) quorum sensing to regulate the physiology and pathogenesis. Song *et al.* demonstrate that the long-chain fatty acid-CoA ligase FadD1 controls the important biological functions and virulence in *P. aeruginosa* as a novel response regulator of *cis*-DA signal.

SUMMARY

Diffusible signal factor (DSF)-family quorum sensing (QS) signals are widely utilized by many bacterial pathogens to modulate various biological functions and virulence. Previous studies showed that *cis-2*-decenoic acid (*cis*-DA) is involved in modulation of biofilm dispersion in *Pseudomonas aeruginosa*, but its signaling mechanism remains vague. Here, we report that *cis*-DA regulates the physiology and virulence of *P. aeruginosa* through the long-chain fatty acid-CoA ligase FadD1. *Cis*-DA specifically binds to FadD1 with high affinity and enhances the binding of FadD1 to the promoter DNA region of *lasR*. Further analysis showed that the FadD1 is a response regulator of *cis*-DA with a DNA-binding leucine zipper motif to control the transcription of various target genes. Moreover, FadD1 exhibited catalytic activity on *cis*-2-dodecenoic acid (BDSF) of *Burkholderia cenocepacia* and enhanced the competitiveness of *P. aeruginosa*. Together, our work presents a new DSF-type QS signaling system, which is highlighted by its receptor and response regulator evolved from a canonical enzyme of fatty acid metabolism.

INTRODUCTION

Quorum sensing (QS) is a cell-to-cell communication mechanism utilized by numerous species of bacteria. The first QS signal discovered in gram-negative bacteria is the acyl homoserine lactone (AHL) produced by *Vibrio fischeri*[1-5]. In addition to AHL-family signals, there are many other types of QS signals[6-16], including the diffusible signal factor (DSF)-family signals. *Pseudomonas aeruginosa* is a major source of opportunistic infections in both immunocompromised individuals and cystic fibrosis patients and has evolved at least three types of QS systems, e.g., *las*, *pqs* and *rhl*[17-19]. The *las* and *rhl* AHL QS systems use N-3-oxo-dodecanoyl-l-homoserine lactone (3-oxo-C12-HSL) and N-butyryl-l-homoserine lactone (C4-HSL), respectively, to regulate biological functions, such as biofilm formation, motility and virulence factor production[20-24]. The *pqs* system employs 2-heptyl-3-hydroxy-4(1H)-quinolone (PQS), to regulate biological functions and virulence[25-29]. These QS systems of *P. aeruginosa* are hierarchically interrelated, and the *las* system was verified to monitor both the *rhl* and *pqs* systems[17, 30]. Moreover, it was also revealed that the fatty acid signaling molecule *cis*-2-decenoic acid (*cis*-DA) exhibited important functions in *P. aeruginosa*[31]. The production of *cis*-DA in *P.*

aeruginosa needs an enoyl-CoA synthase encoded by DspI[32]. The inactivation of DspI leads to a

significant reduction in biofilm dispersion[33,34]. However, it is not yet clear how *cis*-DA regulates biofilm

dispersion, and its receptor and downstream signaling network remain to be investigated.

Fatty acyl coenzyme A (CoA) synthetases (FACSs; fatty acid-CoA ligases) are a class of enzymes

that activate alkanoic acids to CoA esters[35]. FACSs are widely distributed in both prokaryotic and

eukaryotic organisms and exhibit broad substrate specificity[36]. The long-chain fatty acid-CoA ligase

(FadD) is responsible for the activation of endogenous long-chain fatty acids into acyl-CoAs[37]. RpfB,

which has recently been reported as a long-chain fatty acid-CoA ligase of *Xanthomonas campestris*

pv. campestris (*Xcc*), is involved in the activation of a wide range of fatty acids to their CoA esters *in*

vitro[38]. Intriguingly, RpfB is required for the turnover of both *cis*-11-methyl-dodecenoic acid (DSF) and

cis-2-dodecenoic acid (BDSF) QS signals *in vivo*[39]. The *rpfB* deletion mutant produced high levels of

both DSF and BDSF signals and displayed increased virulence in *Xcc*[40,41]. Moreover, RpfB-dependent

QS signal turnover was also detected in various bacterial species[40].

Transcription factors (TFs) are proteins that bind to specific DNA sequences, thereby regulating the

transcription of genetic information from DNA to messenger RNA[42]. Once bound to DNA, these

proteins can promote or block the recruitment of RNA polymerase to specific genes to control

transcriptional expression[43]. Although there is great diversity in the structure and function of DNA-

binding proteins, some polypeptide motifs enable binding to the major groove of DNA, including helix-

turn-helix (HTH), helix-loop-helix (HLH), zinc fingers, and leucine zippers[44-47].

In this study, we demonstrated that FadD1 of *P. aeruginosa* could convert BDSF to *cis*-2-dodecenoic

acid-CoA and reduce the competitiveness of *B. cenocepacia*. FadD1 also controls the physiology and

virulence of *P. aeruginosa* by sensing *cis*-DA, then monitors the QS hierarchy and controls the

expression of various target genes as a global transcriptional regulator. These findings indicate that

FadD1 is a member of a new class of response regulators of QS signal that controls the important

biological functions and virulence of bacterial pathogens, with an additional role in fatty acid oxidation

to catalyze the formation of fatty acyl-CoA.

RESULTS

FadD1 of *P. aeruginosa* shows enzyme activity on the BDSF signal

BDSF was first found to be involved in the regulation of biofilm formation, motility and virulence of *B. cenocepacia*[9-12,48,49]. It was subsequently reported that BDSF from *B. cenocepacia* interferes with QS systems and the type III secretion system (T3SS) of *P. aeruginosa*[50]. Interestingly, we found that BDSF could also be degraded when it was added to the culture of *P. aeruginosa* PAO1. The amount of BDSF in the culture of *P. aeruginosa* decreased to 21.78% after 6 h of incubation (Figure 1A). Previous work demonstrated that RpfB is a long-chain fatty acid-CoA ligase responsible for BDSF and DSF signal turnover in *Xcc*[38,39]. To discover the enzyme that degrades BDSF, we then searched RpfB homologs in the genome of *P. aeruginosa* PAO1 by using the Basic Local Alignment Search Tool (BLAST) algorithm[51]. We found six RpfB homologs, FadD1-6[52]; among them, FadD1 showed the highest homology, with 55.35% identity with RpfB. Therefore, FadD1, which contains 562 aa with a calculated molecular weight of 61.67 kDa, was purified to homogeneity using affinity chromatography (Figure 1B). The *in vitro* enzymatic activity assays showed that when FadD1 was mixed with BDSF, the free BDSF level in the mixture decreased nearly undetectable after 1 h (Figure 1C). These results suggested that FadD1 exhibits enzyme activity on BDSF. Consistently, deletion of *fadD1* in *P. aeruginosa* caused a substantial reduction in BDSF degradation activity (Figure 1D) but did not affect the growth rate of the bacterial cells (Figure S1).

We then investigated the *in vivo* activity of FadD1 in *B. cenocepacia* H111 by overexpressing the *fadD1* gene. Overexpression of *fadD1* did not affect the growth rate but markedly reduced the BDSF production of the bacterial cells (Figure S2A, B). As expected, deletion of *rpfF_{Bc}* resulted in complete abolishment of the production of BDSF in *B. cenocepacia* H111 (Figure S2B). Interestingly, overexpression of *fadD1* in the *B. cenocepacia* wild-type strain reduced biofilm formation, swarming motility and protease activity, which could be restored by the addition of exogenous BDSF at 5 μM (Figure S2C-E). Overexpression of *fadD1* in the wild-type strain H111 also decreased cytotoxicity (Figure S2F).

FadD1 boosts the competitive capability of *P. aeruginosa* against *B. cenocepacia*

As BDSF mediates the cross-talk between *B. cenocepacia* and *P. aeruginosa* [50] and FadD1 exhibits

catalysis activity on BDSF, we investigated whether FadD1 plays a role in the competitive interactions

between the two bacterial species. To this end, we cocultured the green fluorescent protein (GFP)

fluorescence-labeled *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and complemented strains in the

presence of the *B. cenocepacia* H111 strain labeled with mCherry and measured the BDSF

production of *B. cenocepacia* (Figure 1D) and the mean fluorescence intensity (MFI) of both *B.*

cenocepacia and *P. aeruginosa* at different time points as indicated (Figure 1E, F). The results

showed that deletion of *fadD1* caused a drastic reduction in the degradation capability of *P.*

aeruginosa toward the BDSF signal. The BDSF signal yield of *B. cenocepacia* cocultured with the *P.*

aeruginosa deletion mutant Δ*fadD1* was more than 5 times higher than that when it was cocultured

with wild-type PAO1 or the complemented strain Δ*fadD1*(*fadD1*) at 9 h postinoculation (Figure 1D).

On the other hand, we found that the GFP mean fluorescence intensity (MFI) of the *P. aeruginosa*

Δ*fadD1* strain was lower than that of the *P. aeruginosa* wild-type and complemented strains when

they were cocultured with the *B. cenocepacia* strains (Figure 1E). The mCherry mean fluorescence

intensity (MFI) of the *B. cenocepacia* H111 strain cocultured with the *P. aeruginosa* Δ*fadD1* strain

was higher than that of the *B. cenocepacia* H111 strain cocultured with the *P. aeruginosa* wild-type

and complemented strains (Figure 1F). These results indicated that FadD1 plays an important role in

the competition between *P. aeruginosa* and *B. cenocepacia*.

Deletion of FadD1 impairs biological functions and pathogenicity in *P. aeruginosa*

It was determined that FadD1 plays an important role in the interaction between *P. aeruginosa* and

B. cenocepacia. We continued to study whether FadD1plays a role in the regulation of biological

functions and pathogenesis in *P. aeruginosa*. We found that deletion of *fadD1* resulted in decreases

in biofilm formation, swarming, and pyocyanin production by 30.47%, 36.72% and 62.15%,

respectively (Figure 2A-C). Deletion of *fadD1* also attenuated cytotoxicity by 69.78% when A549 cells

were incubated with the *P. aeruginosa* strains at 8 h postinoculation (Figure 2D). We tested whether

there was a change in the expression level of upstream (*PA3298*) and downstream (*fadD2*) genes of

the *fadD1* mutant gene. The results showed that there was no change in the expression of upstream

and downstream genes in the *fadD1* deletion mutant (Figure S3). To further study the regulatory roles

of FadD1 in controlling bacterial physiology, we analyzed and compared the transcriptomes of the

wild-type strain and the Δ*fadD1* mutant strain by using RNA sequencing (RNA-Seq). Differential gene

expression analysis showed that the expression levels of 23 genes were increased and those of 208

genes were decreased in the Δ*fadD1* mutant strain compared with the expression levels in the wild-

type strain (|Log$_2$ fold-change| ≥ 1.0) (Table S1 and Figure S4A). These differentially expressed genes

are associated with a range of biological functions (Table S1 and Figure S4B). These genes include

the *las*, *rhl* and *pqs* QS system genes *PA1432* (*lasI*), *PA1430* (*lasR*), *PA3476* (*rhlI*), *PA3477* (*rhlR*),

PA0996 (*pqsA*), and *PA1003* (*mvfR*), which were shown to be involved in pathogenicity in *P.*

aeruginosa.

FadD1 monitors the QS systems of *P. aeruginosa*

As FadD1 obviously affects the QS-regulated genes and phenotypes of *P. aeruginosa* Figure 2A-D,

Figure S4 and Table S1), we inferred that FadD1 might affect the QS systems. Analysis of the

expression profiles of *lacZ* under the control of the *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and *pqsA* promoters

showed that deletion of *fadD1* resulted in reduced expression levels of *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and

pqsA (Figure 2E-J), which are signal molecule receptor genes and synthase-encoding genes of the

las, *rhl* and *pqs* QS systems, respectively. RT-qPCR analysis and RNA-seq also showed that the

mutation of *fadD1* caused a decrease in the expression levels of *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and *pqsA*

(Figure 2K). Therefore, we measured and compared the production of 3-oxo-C12-HSL, C4-HSL and

PQS in the wild-type, *fadD1* mutant, and complemented strains. We found that the production of 3-

oxo-C12-HSL, C4-HSL and PQS was reduced in the *fadD1* mutant strain (Figure 2L-N). These results

demonstrated that FadD1 positively regulated the *las*, *rhl* and *pqs* QS systems in *P. aeruginosa*.

FadD1 regulates target gene expression by directly binding to the promoter

FadD1 obviously affects the QS gene expression levels and QS signal production of *P. aeruginosa*.

Therefore, we performed electrophoretic mobility shift assays (EMSAs) to test whether the

transcriptional regulation of the signal molecule synthase-encoding genes and receptor genes of the

las, *rhl* and *pqs* QS systems is achieved by direct binding of FadD1 to their promoters. The 339-bp,

321-bp, 270-bp, 330-bp, 306-bp and 324-bp DNA fragments from the *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and

pqsA promoters were PCR-amplified and used as probes. Surprisingly, as shown in Figure. 3A, the

lasR promoter DNA fragment formed stable DNA-protein complexes with FadD1 and migrated slower

than unbound probes. The amount of labeled probe that bound to FadD1 increased with increasing

amounts of FadD1 but decreased in the presence of a 50-fold greater concentration of unlabeled

probe (Figure 3A). However, FadD1 did not bind to the promoters of the other tested genes, i.e., *rhlR*,

mvfR, *lasI*, *rhlI* and *pqsA* (Figure S5 and Figure S6).

To identify the DNA motif recognized by FadD1 and the genes directly regulated by FadD1, we

performed chromatin immunoprecipitation sequencing (ChIP-Seq) analysis. ChIP-Seq data analysis

revealed the potential FadD1 binding site as 5'-AGGACGG-3' in the *lasR* gene promoter probe (Figure

3B). The results showed that FadD1 could directly bind to multiple target gene promoters to regulate

their transcription and expression (Table S2). To further examine the binding sites, three target genes,

flgM, *katA* and *PA0692,* were chosen for EMSA analysis. The amounts of labeled probes of *flgM*, *katA*

and *PA0692* bound to FadD1 were also increased with increasing amounts of FadD1 (Figure S7).

Then, the potential binding sites 5'-AGGACGG-3', 5'-TGGCCGG-3', 5'-AGGAGGG-3', and 5'-

AGGGCGG-3' were deleted from the promoter regions of *lasR*, *flgM*, *katA* and *PA0692*, respectively.

EMSA analysis showed that no DNA-protein complex was formed when the binding site was deleted

from the probes (Figure 3C and Figure S8), indicating that this fragment is essential for the binding of

FadD1 to the *lasR, flgM*, *katA* and *PA0692* promoters. Together, these results demonstrated that

FadD1 regulates target gene expression by directly binding to a specific region in the gene promoter.

The leucine zipper motif of FadD1 is responsible for binding with DNA

Since FadD1 can bind to *lasR* and regulate transcription, and the domain structure analysis of FadD1

by using the HMMER (https://www.ebi.ac.uk/Tools/hmmer/) shows that it has an AMP-binding

enzyme domain (AMP-B) and an AMP-binding enzyme C-terminal domain (AMP-B_C) (Figure 3D).

We then purified the AMP-B domain and the AMP-B_C domain for EMSA (Figure 3E). The results

showed that only the AMP-B domain can bind to the promoter region of *lasR* (Figure 3F, G). To

determine the location of the new DNA binding motif in the AMP-B domain, we used the protein motif

analysis website（https://www.novopro.cn/tools/motifscan.html). A structure called "leucine zipper"

was characterized at 275 to 296 amino acids **L**TGNHNI**L**ITNPRD**L**PSMLKD**L** in AMP-B domain

(Figure 3D and Figure S9). The leucine zipper consists of a periodic repetition of leucine residues at

every seventh position over a distance covering eight helical turns. This motif is found in many

eukaryotic transcription factors[45]. However, in bacterial transcription factor, this motif is only found in

the classical DNA-binding HTH domain of the LysR family regulator MetR[46] and a BDSF response

regulator DsfR identified recently[47]. So we speculated that the four leucine residues L275, L282, L289

and L296 might be important for the binding of FadD1 to target gene promoters. We then introduced

four site-specific substitutions into FadD1 (named FadD1[LM]), and the results showed that the

simultaneous mutation of L275A, L282A, L289A and L296A abolished the binding of FadD1 to the

lasR promoter (Figure 3I). In addition, *in trans* expression of FadD1[LM] did not rescue the defective

biofilm, swarming motility and pyocyanin production phenotypes of the Δ*fadD1* strain (Figure 3J-L).

We then used the HDOCK server to automatically predict the interaction between FadD1 and the

DNA-binding site 5'-AGGACGG-3'. The binding model of DNA and FadD1 and the detailed contacts

of the complex are shown in Figure S10 and Table S3, respectively. The binding sites of FadD1 are

in the leucine zipper structure. The results are consistent with the conclusion that the DNA-binding

motif of FadD1 is the leucine zipper.

FadD1 is a receptor of *cis*-DA

Since *cis*-DA is a DSF-type QS signal in *P. aeruginosa*, and the FadD1 controls QS-regulated

phenotypes as a global transcriptional regulator in *P. aeruginosa*, we then measured whether there

is an interaction between *cis*-DA and FadD1. To test this hypothesis, we purified FadD1 to performed

microscale thermophoresis (MST) analysis to test whether FadD1 binds *cis*-DA. As shown in Figure

4A, FadD1 exhibited binding activity to *cis*-DA, with an estimated dissociation constant (K_D) of 1.06 ±

0.14 µM. To further confirm FadD1 is a receptor of *cis*-DA, we generated the *cis*-DA deficient mutant

Δ*dspI*, the double deletion mutants Δ*dspI*Δ*fadD1* and *in trans* expressed *fadD1* in the Δ*dspI* and

measured the motility activity and pyocyanin of these strains. The results showed that *in trans*

expression of *fadD1* rescued the defective motility and pyocyanin phenotypes of the Δ*dspI* strain

(Figure 4B, C). Interestingly, we found that the addition of exogenous *cis*-DA rescued the impaired

motility and pyocyanin of Δ*dspI* (Figure 4B, C); however, addition of exogenous *cis*-DA showed no

effect on these phenotypes of the mutant Δ*fadD1* and Δ*dspI*Δ*fadD1* (Figure 4B, C). We then measured

the gene expression levels of *lasR*, *lasI*, *rhlR*, *rhlI*, *mvfR* and *pqsA* in the wild-type strain and the

Δ*dspI* strain. The results indicated that deletion of *dspI* caused a significant decrease in the gene

expression levels of *las*, *rhl* and *pqs* QS system genes (Figure 4D-I). Therefore, we measured and

compared the production of 3-oxo-C12-HSL, C4-HSL and PQS in the wild-type, *dspI* mutant, and

complemented strains. The result showed that the production of 3-oxo-C12-HSL, C4-HSL and PQS

was reduced in the *dspI* mutant strain (Figure 4J-L). These results demonstrated that both *cis*-DA and

FadD1 positively regulates the *las*, *rhl* and *pqs* QS systems in *P. aeruginosa*.

cis-DA enhances the binding of FadD1 to target gene promoters

To determine how the binding of *cis*-DA to FadD1 might affect the activity of FadD1, we examined the

effects of *cis*-DA on the binding of FadD1 to the *lasR* promoter by EMSA. As shown in Figure 5A, the

binding of FadD1 to the promoter probe of *lasR* was enhanced when *cis*-DA was present in the

reaction mixtures, and the amount of probe bound to FadD1 increased as the amount of *cis*-DA

increased (Figure 5A).

Then, we attempted to identify the *cis*-DA-binding sites in FadD1. Autodocking analysis revealed 2

amino acid residues, Gly488 (G488) and Arg552 (R552), that might be critical for the interaction

between FadD1and *cis*-DA (Figure 5B). To test whether the roles of these acid residues are related

to the binding to *cis*-DA, we then generated two single point mutants (FadD1^{G488A} and FadD1^{R552A})

(Figure 5C). MST analysis showed that the mutant FadD1^{G488A} and FadD1^{R552A} did not bind to *cis*-DA

(Figure 5D, E). Consistent with this, EMSA analysis showed that mutation of the *cis*-DA-binding site

(Gly488 and Arg552) completely eliminated the effect of *cis*-DA on FadD1 (Figure 5F, G). In addition,

in trans expression of *fadD1*^{G488A} and *fadD1*^{R552A} only partially rescued motility and pyocyanin (Figure

5H, I), and the addition of exogenous 20 µM *cis*-DA showed no effect on these phenotypes of the

mutant Δ*dspI*Δ*fadD1*(*fadD1^{G488A}*) and Δ*dspI*Δ*fadD1*(*fadD1^{R552A}*) (Figure 5H, I). Thus, we concluded

that FadD1 is a specific response regulator of *cis*-DA in *P. aeruginosa*.

Reconstruction of FadD2 with the leucine zipper motif

Our study found that the DNA binding motif of FadD1 is a leucine zipper motif. Therefore, we used

the protein motif analysis website (https://www.novopro.cn/tools/motifscan.html) to analyze whether

the other FadD proteins, FadD2-6, contain a leucine zipper structure. The results showed that the

other FadD proteins did not contain a leucine zipper structure, and EMSA analysis showed that FadD2,

which shares about 59.82% amino acid identity with FadD1, did not bind to the *lasR* gene promoter

(Figure S11A, B). To further investigate the role of the leucine zipper motif of FadD1, we

reconstructed FadD2 by adding the leucine zipper structure of FadD1 to the same part of FadD2

(named FadD2$_{LZ}$) and tested whether the reconstructed FadD2 has regulatory activity similar to that

of FadD1. Therefore, FadD2$_{LZ}$ was purified to homogeneity using affinity chromatography (Figure

S11C), and assayed whether it could bind to the *lasR* promoter. Intriguingly, we found that the *lasR*

promoter DNA fragment formed stable DNA-protein complexes with FadD2$_{LZ}$ and migrated more

slowly than unbound probes. The amount of labeled probe that bound to FadD2$_{LZ}$ increased with

increasing amounts of FadD2$_{LZ}$ but decreased in the presence of a 50-fold greater concentration of

unlabeled probe (Figure S11D). Interestingly, addition of exogenous *cis*-DA also enhanced the

binding of FadD2$_{LZ}$ to the *lasR* promoter probe at a final concentration of 20 μM (Figure S11E). In

addition, *in trans* expression of the *fadD2$_{LZ}$* gene rescued the defective swarming motility and

pyocyanin production phenotypes of the Δ*fadD1* strain (Figure 11F, G). And we also found that the

addition of exogenous *cis*-DA rescued the impaired motility and pyocyanin of Δ*dspI*Δ*fadD1*(*fadD2$_{LZ}$*)

(Figure S11F, G). These results showed that the leucine zipper motif plays a critical role in the

regulatory activity of FadD1.

Analysis of the catalytic motif of FadD1

Previous studies reported that fatty acyl-CoA ligase plays a crucial role in catalyzing the formation of

fatty acyl-CoA through hydrolysis of pyrophosphate[35,36]. To test whether FadD1 can function in the

uptake and activation steps of fatty acid degradation, we evaluated its enzymatic activity *in vitro*. The results showed that FadD1 had high activity on lauric acid (C12:0), while it displayed moderate activity on BDSF but no activity on *cis*-DA (Figure S12A). In addition, the previous study found that mutation of Trp433, Thr436 and Arg453, the substrate binding sties, results in complete inactivation of fatty acyl-CoA synthetase in *E. coli* (37). We continued to generated three single point mutants (FadD1^{W434A}, FadD1^{T437A} and FadD1^{R454A}) to explore whether the mutation affects the enzyme activity of FadD1 (Figure S12B). Similarly, it was shown that the FadD1^{W434A}, FadD1^{T437A} and FadD1^{R454A} exerted no enzyme activity on *cis*-DA, BDSF and lauric acid (C12:0) (Figure S12C-E). To further characterize the relationship between enzyme activity of FadD1 and perception of *cis*-DA, we chose to test the regulatory activity of the mutant FadD1^{R454A}. We *in trans* expressed the *fadD1^{R454A}* in the Δ*dspI*Δ*fadD1* mutant strain. Interestingly, it was found that addition of exogenous *cis*-DA rescued the impaired motility of Δ*dspI*Δ*fadD1*(*fadD1^{R454A}*) (Figure S12F). Consistent with this, EMSA analysis showed that mutation of the substrate binding stie (Arg454) would not affect the perception of *cis*-DA (Figure S12G).

Homologs of FadD1 are widespread in bacteria

To determine whether FadD1 is widely distributed, homologs of FadD1 were sought in the genome database by using the BLAST program. It was indicated that homologs of FadD1 are distributed across many different bacterial species, including in the genera *Pseudomonas*, *Azotobacter*, *Azomonas*, and *Halopseudomonas* (Table S4). Furthermore, the homologs of FadD1 in the genera *Pseudomonas, Azotobacter* and *Azomonas* contain a leucine zipper structure (Table S4). We then selected *P. fluorescens* Migula ATCC17518, which contains both a FadD1$_{Pf}$ protein with a leucine zipper structure and a *lasR$_{Pf}$* gene, and the FadD1$_{Pf}$ protein was purified by affinity chromatography and tested for its interaction with the *lasR$_{Pf}$* gene promoter. The EMSA results showed that the FadD1$_{Pf}$ protein from *P. fluorescens* could also bind to the *lasR$_{Pf}$* gene promoter (Figure S13), indicating that the transcriptional regulatory mechanism of FadD1 might be widely conserved in bacteria.

DISCUSSION

Fatty acid-CoA ligase plays a vital role in fatty acid metabolism, as it can convert fatty acids to fatty

acyl-CoA in bacteria[37]. In this study, we found that the fatty acid-CoA ligase FadD1 of *P. aeruginosa*

not only exhibits high activity for the catalysis of lauric acid (C12:0) to lauric acid-CoA (Figure. S12A),

and employs a key role in the bacterial competition with *B. cenocepacia* by degrading its QS signal

BDSF (Figure 1), but also plays a critical role in the regulation of *P. aeruginosa* physiology and

pathogenesis by acting as the receptor and response regulator of its QS signal *cis*-DA (Figure 2 and

Table S1). We demonstrated that FadD1 can directly bind to the promoter of the target genes, thereby

regulate the transcription levels of the target genes (Figure 3A and Figure S7). It was further identified

that mutation of the leucine zipper structure of FadD1 abolished the binding of FadD1 to the target

gene promoters (Figure 3I). Interestingly, we found that FadD2 does not contain a leucine zipper

structure, and EMSA analysis showed that FadD2 did not bind to the *lasR* gene promoter (Figure

S11B). However, the reconstructed FadD2 with addition of the leucine zipper structure of FadD1 to

the same part of FadD2 (named FadD2$_{LZ}$) obtained the regulatory activity similar to that of FadD1. It

could bind to the *lasR* gene promoter and restore the *fadD1* mutant phenotypes (Figure S11).

Furthermore, our findings indicate that FadD1 is widely conserved in many other bacterial species,

including *Pseudomonas*, *Azotobacter*, and *Azomonas* species (Table S4), indicating that FadD1 with

a leucine zipper structure might be a new family of transcriptional regulatory proteins widely conserved

in bacteria.

The fatty acid molecule *cis*-DA was first identified in *P. aeruginosa* to act as the autoinducer of

biofilm dispersion[31]. It has also been shown to induce biofilm dispersion in a range of Gram-negative

and Gram-positive bacteria and in the fungal pathogen *Candida albicans*[31]. Although the previous

study reported that the two-component sensor/response regulator hybrid DspS (PA4112,

PA14_10770) is required and essential for native biofilm dispersion in response to the cell-to-cell

signaling molecule *cis*-DA[34], there was no exploration of other functions of *cis*-DA. In this study, we

found that deletion of *dspl*, which is the synthase of *cis*-DA, caused an impairment in motility and

pyocyanin production (Figure 4B, C). Interestingly, in *trans* expression of *fadD1* rescued the defective

motility and pyocyanin phenotypes of the Δ*dspl* strain (Figure 4B, C). However, the addition of

exogenous *cis*-DA showed no effect on these phenotypes of the mutant Δ*fadD1* and Δ*dspl*Δ*fadD1*

(Figure 4B, C). Moreover, *cis*-DA is shown to bind to FadD1 with high affinity and enhanced the

binding of FadD1 to the target gene promoter probes (Figure 4A and Figure 5A). We also identified

that two amino acid residues at positions Gly488 (G488) and Arg552 (R552) are critical for the

interaction between FadD1 and *cis*-DA (Figure 5). Mutations at Gly488 and Arg552 abolished the

binding between FadD1 and *cis*-DA (Figure 5D, E). Another interesting observation in this study was

that mutation at the substrate binding stie Arg454 (R454) completely abolished the enzyme activity

of FadD1 on both BDSF and lauric acid (C12:0) (Figure S12C-E), but could not affect the effect of

cis-DA on FadD1 (Figure S12G), suggesting that the *cis*-DA-binding site is different from the

substrate-binding stie in FadD1. Together, FadD1 was identified to be a new response regulator of

cis-DA QS signal.

BDSF has been identified to be a DSF-family QS signal in *B. cenocepacia*[11]. As *cis*-DA shares a

similar chemical structure with that of BDSF, we then continued to explore the interaction between

BDSF and FadD1. Although FadD1 showed a degradation activity on BDSF, we found that FadD1

tightly binds to BDSF with an estimated dissociation constant (K_D) of 1.65 ± 0.21 µM (Figure S14A).

Interestingly, the addition of exogenous BDSF also enhanced the binding of FadD1 to the *lasR*

promoter probe at a final concentration of 10 µM (Figure S14B). To distinguish the role of BDSF to be

a "substrate" or "ligand" of FadD1, we then used EMSA analysis and the result showed that mutation

of the *cis*-DA-binding sites abolished the perception of *cis*-DA (Figure 5), but will not affect the effect

of BDSF on FadD1 (Figure S14C). However, EMSA analysis showed that mutation of the substrate-

binding stie (Arg454) completely abolished the effect of BDSF on the binding of FadD1 to target gene

promoter (Figure S14D), while not affect the effect of *cis*-DA on FadD1 (Figure S12G). These results

suggested that BDSF also interferes with the binding of FadD1 to target gene promoters *in vitro* as a

substrate of the canonical enzyme FadD1.

In *P. aeruginosa,* there are at least three QS systems, namely, *las*, *rhl* and *pqs*, which are involved

in the regulation of a broad range of genes important for metabolism and virulence[18,19]. These QS

systems are hierarchically interrelated, and the *las* system was verified to monitor both the *rhl* and

pqs systems[17,30]. In this study, we identified that FadD1 affects the QS-regulated phenotypes,

including the biofilm formation, swarming, and pyocyanin in *P. aeruginosa* (Figure 2A-C). Surprisingly, we also identified that FadD1 is a *cis*-DA receptor protein that positively regulates the *las, rhl* and *pqs* QS systems in *P. aeruginosa* (Figure 2 and Figure 6). These results indicated that FadD1 is not only a response regulator of *cis*-DA but also an important upstream regulator of the QS signaling network in *P. aeruginosa*. FadD1 regulates numerous target genes and physiological functions by monitoring the QS signaling systems, which broadens our understanding of the QS signaling network of *P. aeruginosa*.

In summary, our findings suggest that FadD1, which contains a leucine zipper structure in the AMP-B domain, is not only an enzyme of fatty oxidation but also a receptor of *cis*-DA that directly controls target gene expression and controls the QS network in *P. aeruginosa*. In addition, the BLAST search revealed that homologs of FadD1 containing the leucine zipper structure are highly conserved in many bacteria (Table S4), suggesting that the regulatory mechanism of FadD1 might be present in various bacterial species. Consistently, our recent study revealed that the fatty acyl-CoA ligase DsfR (BCAM2136), which contains a leucine zipper motif, acts as a global transcriptional regulator to control *B. cenocepacia* virulence by sensing BDSF[47]. Different from FadD1, DsfR has a specific BDSF-sensing site and showed no enzyme activity on BDSF, while FadD1 only binds with BDSF on the substrate-binding sites. These results indicate that FadD1 and DsfR share some common characteristics and are distinguished from each other. In conclusion, our work presents a unique and widely conserved signal receptor of *cis*-DA, which might be an important new type of QS signal receptor in bacteria.

Limitations of the study

* Limitations of the Study: As it reads right now, the section mostly focuses on future directions of the work to answer outstanding open questions. While future directions can belong to this section, it is mostly the place to openly discuss potential caveats of the work (technical, conceptual, or other limitations) with the intent of promoting clarity and transparency. I would like to suggest that you further develop and edit this section in this direction.

Our findings demonstrated that FadD1 is a specific receptor protein of *cis*-DA QS signal. It was

revealed that *cis*-DA binds to FadD1 to enhance the binding of FadD1 to the promoter DNA of target

genes. However, it remains unknown how does the binding to *cis*-DA induce the regulatory activity of

FadD1 on the transcription of target genes.

STAR METHODS

Detailed methods are provided in the online version of this paper and include the following:

KEY RESOURCES TABLES

RESOURCE AVAILABILITY

Lead contact

Materials availability

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Bacterial strains and culture conditions

Cell culture

METHOD DETAILS

BSDF extraction

Quantification of BDSF and cis-DA

Protein expression and purification

Enzyme activity assays

Construction of in-frame deletion mutants and complementation

Bacterial growth analysis

Competition assays in the mixed culture

P. aeruginosa phenotype assays

B. cenocepacia phenotype assays

Cytotoxicity assays

Construction of reporter strains and measurement of β-galactosidase activity

Quantitative reverse transcription PCR assays

Quantitative analysis of QS signal production in P. aeruginosa

RNA-Seq analysis

Electrophoretic gel mobility shift assay

Chromatin immunoprecipitation sequencing analyses

Molecular docking FadD1

Microscale thermophoresis assay

STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

448   Supplemental information can be found online at

ACKNOWLEDGMENTS

451   This work was financially supported by the National Key Research and Development Program of

452   China (2021YFA0717003 to Y.D.), the National Natural Science Foundation of China (32300033 to

453   S.S.) and the Scientific Research Foundation of Hainan University (KYQD(ZR)-23006 to S.S.).

AUTHOR CONTRIBUTIONS

456   Y.D. and S.S. conceived the project. S.S., J.L., and Y.D. designed the research. S.S., J.L., B.W., Y.S.,

457   H.H., X.S., M.W., and B.C. performed the research. S.S., J.L., G.W., Y.H., L.X., B.G., L.Y., X.W., L-

458   H.Z., and Y.D. analyzed the data. S.S. and Y.D. wrote the paper.

DECLARATION OF INTERESTS

461   The authors declare that they have no conflict of interest.

INCLUSION AND DIVERSITY

464   We support inclusive, diverse, and equitable conduct of research.

REFERENCES

1. Nealson, K.H. and Hastings, J.W. (1979). Bacterial bioluminescence: its control and ecological

significance. Microbiol Rev *43*, 496-518. doi: 10.1128/mr.43.4.496-518.1979.

2. Fuqua, C. and Greenberg, E.P. (2002). Listening in on bacteria: acyl-homoserine lactone

signalling. Nat Rev Mol Cell Biol *3*, 685-695. doi: 10.1038/nrm907.

3. Papenfort, K. and Bassler, B.L. (2016). Quorum sensing signal-response systems in Gram-

negative bacteria. Nat Rev Microbiol *14*, 576-588. doi: 10.1038/nrmicro.2016.89.

4. Whiteley, M., Diggle, S.P. and Greenberg, E.P. (2017). Progress in and promise of bacterial
quorum sensing research. Nature *551*, 313-320. doi: 10.1038/nature24624.

5. Chadha, J., Harjai, K. and Chhibber, S. (2022). Revisiting the virulence hallmarks of
Pseudomonas aeruginosa: a chronicle through the perspective of quorum sensing. Environ
Microbiol *24*, 2630-2656. doi: 10.1111/1462-2920.15784.

6. Flavier, A.B., Clough, S.J., Schell, M.A. and Denny, T.P. (1997). Identification of 3-hydroxypalmitic
acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*. Mol
Microbiol *26*, 251-259. doi: 10.1046/j.1365-2958.1997.5661945.x.

7. Barber, C.E. *et al.*, (1997). A novel regulatory system required for pathogenicity of *Xanthomonas
campestris* is mediated by a small diffusible signal molecule. Mol Microbiol *24*, 555-566. doi:
10.1046/j.1365-2958.1997.3721736.x.

8. Déziel, E. *et al.*, (2004). Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs)
reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. Proc Natl Acad Sci U
S A *101*, 1339-1344. doi: 10.1073/pnas.0307694100.

9. Zhang, L.H. and Dong, Y.H. (2004). Quorum sensing and signal interference: diverse implications.
Mol Microbiol *53*, 1563-1571. doi: 10.1111/j.1365-2958.2004.04234.x.

10. Diggle, S.P. *et al.*, (2007). The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and
PQS play multifunctional roles in quorum sensing and iron entrapment. Chem Biol *14*, 87-96. doi:
10.1016/j.chembiol.2006.11.014.

11. Boon, C. *et al.*, (2008). A novel DSF-like signal from *Burkholderia cenocepacia* interferes with
Candida albicans morphological transition. ISME J *2*, 27-36. doi: 10.1038/ismej.2007.76.

12. He, Y.W. and Zhang, L.H. (2008). Quorum sensing and virulence regulation in *Xanthomonas
campestris*. FEMS Microbiol Rev *32*, 842-57. doi: 10.1111/j.1574-6976.2008.00120.x.

13. Deng, Y., Wu, J., Tao, F. and Zhang, L.H. (2011). Listening to a new language: DSF-based
quorum sensing in Gram-negative bacteria. Chem Rev *111*, 160-173. doi: 10.1021/cr100354f.

14. Deng, Y. *et al.*, (2012). *Cis*-2-dodecenoic acid receptor RpfR links quorum-sensing signal
perception with regulation of virulence through cyclic dimeric guanosine monophosphate turnover.
Proc Natl Acad Sci U S A *109*, 15479-15484. doi: 10.1073/pnas.1205037109.

15. Dulcey, C.E. *et al.*, (2013). The end of an old hypothesis: the *pseudomonas* signaling molecules
4-hydroxy-2-alkylquinolines derive from fatty acids, not 3-ketofatty acids. Chem Biol *20*, 1481-
1491. doi: 10.1016/j.chembiol.2013.09.021.

16. Wang, M., Li, X., Song, S., Cui, C., Zhang, L.H. and Deng, Y. (2022). The *cis*-2-Dodecenoic Acid
(BDSF) Quorum Sensing System in *Burkholderia cenocepacia*. Appl Environ Microbiol *88*,
e0234221. doi: 10.1128/aem.02342-21.

17. de Kievit, T.R., Kakai, Y., Register, J.K., Pesci, E.C. and Iglewski, B.H. (2002). Role of the
Pseudomonas aeruginosa las and *rhl* quorum-sensing systems in *rhlI* regulation. FEMS Microbiol
Lett *212*, 101-106. doi: 10.1016/s0378-1097(02)00735-8.

18. Lee, J. and Zhang, L.H. (2015). The hierarchy quorum sensing network in *Pseudomonas
aeruginosa*. Protein Cell *6*, 26-41. doi: 10.1007/s13238-014-0100-x.

19. Mukherjee, S., Moustafa, D., Smith, C.D., Goldberg, J.B. and Bassler, B.L. (2017). The RhlR
quorum-sensing receptor controls *Pseudomonas aeruginosa* pathogenesis and biofilm

development independently of its canonical homoserine lactone autoinducer. PLoS Pathog *13*,
e1006504. doi: 10.1371/journal.ppat.1006504.

20. Fuqua, W.C., Winans, S.C. and Greenberg, E.P. (1994). Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. J Bacteriol *176*, 269-275. doi: 10.1128/jb.176.2.269-275.1994.

21. Pesci, E.C., Pearson, J.P., Seed, P.C. and Iglewski, B.H. (1997). Regulation of *las* and *rhl* quorum sensing in *Pseudomonas aeruginosa*. J Bacteriol *179*, 3127-3132. doi: 10.1128/jb.179.10.3127-3132.

22. Withers, H., Swift, S. and Williams, P. (2001). Quorum sensing as an integral component of gene regulatory networks in Gram-negative bacteria. Curr Opin Microbiol *4*, 186-193. doi: 10.1016/s1369-5274(00)00187-9.

23. Ng, W.L. and Bassler, B.L. (2009). Bacterial quorum-sensing network architectures. Annu Rev Genet *43*, 197-222. doi: 10.1146/annurev-genet-102108-134304.

24. Rutherford, S.T. and Bassler, B.L. (2012). Bacterial quorum sensing: its role in virulence and possibilities for its control. C S H Perspect Med *2*, a012427. doi: 10.1101/cshperspect.a012427.

25. Jander, G., Rahme, L.G. and Ausubel, F.M. (2000). Positive correlation between virulence of *Pseudomonas aeruginosa* mutants in mice and insects. J Bacteriol *182*, 3843-3845. doi: 10.1128/JB.182.13.3843-3845.

26. Cao, H., Krishnan, G., Goumnerov, B., Tsongalis, J., Tompkins, R. and Rahme, L.G. (2001). A quorum sensing-associated virulence gene of *Pseudomonas aeruginosa* encodes a LysR-like transcription regulator with a unique self-regulatory mechanism. Proc Natl Acad Sci U S A *98*, 14613-14618. doi: 10.1073/pnas.251465298.

27. Wade, D.S. *et al.*, (2005). Regulation of *Pseudomonas* quinolone signal synthesis in *Pseudomonas aeruginosa*. J Bacteriol *187*, 4372-4380. doi: 10.1128/JB.187.13.4372-4380.

28. Déziel, E. *et al.*, (2005). The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation: multiple quorum sensing-regulated genes are modulated without affecting *lasRI*, *rhlRI* or the production of N-acyl-L-homoserine lactones. Mol Microbiol *55*, 998-1014. doi: 10.1111/j.1365-2958.2004.04448.x.

29. García-Reyes, S., Soberón-Chávez, G. and Cocotl-Yanez, M. (2020). The third quorum-sensing system of *Pseudomonas aeruginosa*: *Pseudomonas* quinolone signal and the enigmatic PqsE protein. J Med Microbiol *69*, 25-34. doi: 10.1099/jmm.0.001116.

30. Choi, Y. *et al.*, (2011). Growth phase-differential quorum sensing regulation of anthranilate metabolism in *Pseudomonas aeruginosa*. Mol Cells *32*, 57-65. doi: 10.1007/s10059-011-2322-6.

31. Davies, D.G. and Marques, C.N. (2009). A fatty acid messenger is responsible for inducing dispersion in microbial biofilms. J Bacteriol *191*, 1393-1403. doi: 10.1128/JB.01214-08.

32. Amari, D.T., Marques, C.N. and Davies, D.G. (2013). The putative enoyl-coenzyme A hydratase DspI is required for production of the *Pseudomonas aeruginosa* biofilm dispersion autoinducer *cis*-2-decenoic acid. J Bacteriol *195*, 4600-4610. doi: 10.1128/JB.00707-13.

33. Rahmani-Badi, A., Sepehr, S., Fallahi, H. and Heidari-Keshel, S. (2015). Dissection of the *cis*-2-decenoic acid signaling network in *Pseudomonas aeruginosa* using microarray technique. Front Microbiol *6*, 383. doi: 10.3389/fmicb.2015.00383.

34. Kalia, M., Amari, D., Davies, D.G. and Sauer, K. (2023). *cis*-DA-dependent dispersion by

Pseudomonas aeruginosa biofilm and identification of *cis*-DA-sensory protein DspS. mBio *14*, e0257023. doi: 10.1128/mbio.02570-23.

35. Hume, A.R., Nikodinovic-Runic, J. and O'Connor, K.E. (2009). FadD from *Pseudomonas putida* CA-3 is a true long-chain fatty acyl coenzyme A synthetase that activates phenylalkanoic and alkanoic acids. J Bacteriol *191*, 7554-7565. doi: 10.1128/JB.01016-09.

36. Ruth, K., de Roo, G., Egli, T. and Ren, Q. (2008). Identification of two acyl-CoA synthetases from *Pseudomonas putida* GPo1: one is located at the surface of polyhydroxyalkanoates granules. Biomacromolecules *9*, 1652-1659. doi: 10.1021/bm8001655.

37. Black, P.N., Zhang, Q., Weimar, J.D. and DiRusso, C.C. (1997). Mutational analysis of a fatty acyl-coenzyme A synthetase signature motif identifies seven amino acid residues that modulate fatty acid substrate specificity. J. Biol. Chem *272*, 4896-4903. doi: 10.1074/jbc.272.8.4896.

38. Bi, H., Yu, Y., Dong, H., Wang, H. and Cronan, J.E. (2014). *Xanthomonas campestris* RpfB is a fatty Acyl-CoA ligase required to counteract the thioesterase activity of the RpfF diffusible signal factor (DSF) synthase. Mol Microbiol *93*, 262-275. doi: 10.1111/mmi.12657.

39. Zhou, L., Wang, X.Y., Sun, S., Yang, L.C., Jiang, B.L. and He, Y.W. (2015). Identification and characterization of naturally occurring DSF-family quorum sensing signal turnover system in the phytopathogen *Xanthomonas*. Environ Microbiol *17*, 4646-4658. doi: 10.1111/1462-2920.12999.

40. Wang, X.Y., Zhou, L., Yang, J., Ji, G.H. and He, Y.W. (2016). The RpfB-Dependent Quorum Sensing Signal Turnover System Is Required for Adaptation and Virulence in Rice Bacterial Blight Pathogen *Xanthomonas oryzae* pv. oryzae. Mol Plant Microbe Interact *29*, 220-230. doi: 10.1094/MPMI-09-15-0206-R.

41. Tian, X.Q., Wu, Y., Cai, Z. and Qian, W. (2022). BDSF is a degradation-prone quorum-sensing signal detected by the histidine kinase RpfC of *Xanthomonas campestris* pv. *campestris*. Appl Environ Microbiol *88*, e0003122. doi: 10.1128/aem.00031-22.

42. Maston, G.A., Evans, S.K. and Green, M.R. (2006). Transcriptional regulatory elements in the human genome. Annu Rev Genomics Hum Genet *7*, 29-59. doi: 10.1146/annurev.genom.7.080505.115623.

43. Wang, G., Wang, F., Huang, Q., Li, Y., Liu, Y. and Wang, Y. (2015). Understanding Transcription Factor Regulation by Integrating Gene Expression and DNase I Hypersensitive Sites. Biomed Res Int *2015*, 757530. doi: 10.1155/2015/757530.

44. Clark, D.P. and Pazdernik, N.J. (2013). Chapter e14 - Protein Structure and Function, Mol. Biol. (Second Edition), Pages e280-e284. https://doi.org/10.1016/B978-0-12-378594-7.00048-2.

45. Miller, M. (2009). The importance of being flexible: the case of basic region leucine zipper transcriptional regulators. Curr Protein Pept Sci *10*, 244-269. doi: 10.2174/138920309788452164.

46. Maxon, M.E., Wigboldus, J., Brot, N. and Weissbach, H. (1990). Structure-function studies on *Escherichia coli* MetR protein, a putative prokaryotic leucine zipper protein. Proc Natl Acad Sci U S A *87*, 7076-7079. doi: 10.1073/pnas.87.18.7076.

47. Li, X. *et al.*, (2024). Regulation of *Burkholderia cenocepacia* virulence by the fatty acyl-CoA ligase DsfR as a response regulator of quorum sensing signal. Cell Rep *43*, 114223. doi: 10.1016/j.celrep.2024.114223.

48. Deng, Y., Boon, C., Eberl, L. and Zhang, L.H. (2009). Differential modulation of *Burkholderia cenocepacia* virulence and energy metabolism by the quorum-sensing signal BDSF and its

synthase. J Bacteriol *191*, 7270-7278. doi: 10.1128/JB.00681-09.

49. Deng, Y., Wu, J., Eberl, L. and Zhang, L.H. (2010). Structural and functional characterization of diffusible signal factor family quorum-sensing signals produced by members of the *Burkholderia cepacia* complex. Appl Environ Microbiol *76*, 4675-4683. doi: 10.1128/AEM.00480-10.

50. Deng, Y., Boon, C., Chen, S., Lim, A. and Zhang, L.H. (2013). *Cis*-2-dodecenoic acid signal modulates virulence of *Pseudomonas aeruginosa* through interference with quorum sensing systems and T3SS. BMC Microbiol *13*, 231. doi: 10.1186/1471-2180-13-231.

51. Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990). Basic local alignment search tool. J Mol Biol *215*, 403-410. doi: 10.1016/S0022-2836(05)80360-2.

52. Zarzycki-Siek, J. *et al.*, (2013). Elucidating the *Pseudomonas aeruginosa* fatty acid degradation pathway: identification of additional fatty acyl-CoA synthetase homologues. PLoS One *8*, e64554. doi: 10.1371/journal.pone.0064554.

53. Cui, C. *et al.*, (2018). A novel two-component system modulates quorum sensing and pathogenicity in *Burkholderia cenocepacia*. Mol Microbiol *108*, 32-44. doi: 10.1111/mmi.13915.

54. Yang, C. *et al.*, (2017). *Burkholderia cenocepacia* integrates *cis*-2-dodecenoic acid and cyclic dimeric guanosine monophosphate signals to control virulence. Proc Natl Acad Sci U S A *114*, 13006-13011. doi: 10.1073/pnas.1709048114.

55. Guo, Q. *et al.*, (2022). Elongation factor P modulates *Acinetobacter baumannii* physiology and virulence as a cyclic dimeric guanosine monophosphate effector. Proc Natl Acad Sci U S A *119*, e2209838119. doi: 10.1073/pnas.2209838119.

56. Kang, Y., Zarzycki-Siek, J., Walton, C.B., Norris, M.H. and Hoang, T.T. (2010). Multiple FadD acyl-CoA synthetases contribute to differential fatty acid degradation and virulence in *Pseudomonas aeruginosa*. PLoS One *5*, e13557. doi: 10.1371/journal.pone.0013557.

57. Cui, B. *et al.*, (2022). The cell-cell communication signal indole controls the physiology and interspecies communication of *Acinetobacter baumannii*. Microbiol Spectr *10*, e0102722. doi: 10.1128/spectrum.01027-22.

58. Song, S. *et al.*, (2020). Anthranilic acid from *Ralstonia solanacearum* plays dual roles in intraspecies signalling and inter-kingdom communication. ISME J *14*, 2248-2260. doi: 10.1038/s41396-020-0682-7.

59. Langmead, B. and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*, 357-359. doi: 10.1038/nmeth.1923.

60. Trapnell, C. *et al.*, (2010). Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol *28*, 511-515. doi: 10.1038/nbt.1621.

61. Song, S. *et al.*, (2022). An anthranilic acid-responsive transcriptional regulator controls the physiology and pathogenicity of *Ralstonia solanacearum*. PLoS Pathog *18*, e1010562. doi: 10.1371/journal.ppat.1010562.

62. Landt, S.G. *et al.*, (2012). ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. Genome Res *22*, 1813-1831. doi: 10.1101/gr.136184.111.

63. Kleinman, C.L., Sycz, G., Bonomi, H.R., Rodríguez, R.M., Zorreguieta, A. and Sieira, R. (2017). ChIP-seq analysis of the LuxR-type regulator VjbR reveals novel insights into the *Brucella* virulence gene expression network. Nucleic Acids Res *45*, 5757-5769. doi: 10.1093/nar/gkx165.

64. Molecular Operating Environment (MOE), Chemical Computing Group Inc., 1010 Sherbooke St.
West, Suite #910, Montreal, QC, Canada, H3A 2R7 (2018).
65. Yan, Y., Tao, H., He, J. and Huang, S.Y. The HDOCK server for integrated protein-protein docking.
Nat Protoc *15*, 1829-1852 (2020). doi: 10.1038/s41596-020-0312-x.
66. Jumper, J. *et al.*, (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*,
583-589. doi: 10.1038/s41586-021-03819-2.

STAR METHODS

KEY RESOURCES TABLES

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-His	Abcam	ab9108
Bacterial		
Bacterial strains	see Table S5	N/A
Oligonucleotides		
For deletion	see Table S6	N/A
For EMSA	see Table S6	N/A
For *in trans* expression	see Table S6	N/A
For recombinant protein	see Table S6	N/A
For qRT-PCR	see Table S6	N/A
Recombinant DNA		
Plasmids	see Table S5	N/A
Chemicals, peptides, and recombinant proteins		
Tryptone	oxoid	LP0042B
Yeast extract	oxoid	PL0021
NaCl	Macklin	S805275
Agar	Macklin	A800728
Gentamicin	Macklin	G810322
Tetracycline	Macklin	T829835
Kanamycin	Macklin	K885955
Ampicillin	Macklin	A830931
2×T5 Super PCR Mix	Tsinekg	TSE005
ClonExpress II One Step Cloning Kit	Vazyme	C112-02
X-gal	Solarbio	X8050
HisTrap affinity columns	Smart-lifesciences	SA035010

TEV protease	Beyotime	P2307
LightShift Chemiluminescent EMSA Kit	Thermo Fisher	20148
Crystal violet	Sigma-Aldrich	C6158
Monolith NT.115 Protein Labeling Kit	Nano Temper	MO-L014
Silicon capillaries	Nano Temper	MO-L022
PBS buffer	Macklin	917808
Tris	Beyotime	ST760
Eastep Super Total RNA Extraction Kit	Promega	LS1040
cDNA synthesis	Yeasen	11141ES60
SYBR qPCR Green Master Mix	Yeasen	11201ES08
Formaldehyde	Macklin	F809902
Glycine	Macklin	810676
Complete proteinase inhibitor cocktail	Roche	4693116001
SDS	Lablead	L5751
EDTA	Solarbio	E8040
Mini-protease inhibitor cocktail	Roche	11836170001
LiCl	ThermoFisher	AM9480
NP40	ThermoFisher	85124
Deoxycholic acid	Solarbio	D8460
$NaHCO_3$	Macklin	S837271
RNase A	ThermoFisher	EN0531
Proteinase K	Macklin	39450-01-6
IPTG	Macklin	I811719
Glucose	Sigma-Aldrich	G6152
Sucrose	Aladdin	S112228
CytoTox 96 Kit	Promega	G1780
Dulbecco's Modified Eagle's Medium-high glucose (DMEM)	Sigma-Aldrich	D6046
Fetal Bovine Serum	Sigma	F4135
Trypsin EDTA	GIBCO	25200072
DMSO	Sigma-Aldrich	D5879
BDSF	Abcam	ab145341
C4-HSL	Sigma-Aldrich	67605-85-0
3-oxo-C12-HSL	Sigma-Aldrich	168982-69-2

cis-DA	Sigma-Aldrich	15790-91-7
Lauric acid	Sigma-Aldrich	143-07-7
Deposited data		
Raw and Analyzed RNA-Seq Data	This study	PRJNA1121197
Raw and Analyzed ChIP-Seq Data	This study	PRJNA1120965
Software and algorithms		
Prism	GraphPad	https://graphpad.com:443/ scientific-software/prism/
AutoDock	AutoDock Tools	https://autodock.scripps.edu/
Pymol	PyMOL 3.0	https://pymol.org/
Protein Plus	Protein Plus	https://proteins.plus/
HDOCK	HDOCK sever	http://hdock.phys.hust.edu.cn/

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact Yinyue Deng (dengyle@mail.sysu.edu.cn).

Materials availability

Constructs and reagents in this study will be made available upon request, but a completed

Materials Transfer Agreement may be required if there is potential for commercial application.

Data and code availability

Unprocessed data related to figures in this manuscript are available from the lead contact upon

request.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this work paper is available

from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Bacterial strains and culture conditions

The bacterial strains and plasmids used in this work are listed in Table S5. *P. aeruginosa* PAO1, *P.*

fluorescens Migula ATCC17518 and *B. cenocepacia* H111 strains obtained from the American Type

Culture Collection (ATCC) were maintained in Luria-Bertani broth (LB; 10 g/L tryptone, 5 g/L yeast

extract, 5 g/L NaCl) at 37°C. The following antibiotics were added when necessary: ampicillin and

kanamycin at 100 µg mL^{-1}; tetracycline, 10 µg mL^{-1}; and gentamicin, 10 µg mL^{-1}. BDSF and *cis*-DA

(HPLC ≥ 99%) were dissolved in methanol at a final concentration of 100 mM, and this solution was

added to the medium in the experiments. Bacterial growth was determined by measuring the optical

density at 600 nm.

Cell culture

A549 cell were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10%

fetal bovine serum (FBS) and 1% penicillin–streptomycin, cells were cultivated in a humidified

atmosphere containing 5% CO_2 at 37 °C.

METHOD DETAILS

BSDF extraction

The extraction of BDSF from culture supernatants was conducted following previously described

methods[11,53]. To quantify BDSF production in the culture of *B. cenocepacia* strain or *P. aeruginosa*

strain, 100 mL of the supernatant was collected. Its crude ethyl acetate extract was passed through

a 0.22 µm Minisart filter unit and was then condensed to 1 mL for LC-MS analysis.

Quantification of BDSF and *cis*-DA

Quantification of BDSF and *cis*-DA was performed by using an LC-MS system, which consisted of an

ACQUITY UPLC system and a Waters Q-Tof Premier high-resolution mass spectrometer[53]. An

ACQUITY UPLC BEH C18 column 1.7 µm (2.1 × 50 mm) was used for the chromatography analysis

of BDSF and *cis*-DA, which was eluted with a CH_3OH gradient in water at 65-100% supplemented

with 0.01% formic acid at a flow rate of 0.4 mL·min^{-1} for 10 min. Next, 65% CH_3OH was used for 3

min. The entire elution column was introduced into the Q-TOF mass spectrometer according to the

manufacturer's instructions. The BDSF and *cis*-DA levels in the culture supernatant were measured

using the peak area in the extracted ion chromatogram.

Protein expression and purification

700 The coding regions of proteins were amplified with the primers listed in Table S6, and were attached

701 and fused to the expression vectors pET-21a, pET-28a and PDBHT2. The fusion gene constructs

702 were transformed into *E. coli* strain BL21. Affinity purification of the 6×His fusion proteins was

703 performed using Ni-NTA resin (Smart-lifesciences). The His-tag was removed using TEV protease

704 (Beyotime), and the cleaved fusion protein was analyzed by SDS-PAGE[54,55].

Enzyme activity assays

The activity of FadD1 was determined following the methods[56]. *cis*-DA, BDSF or lauric acid (C12:0)

were dissolved in TME buffer (Tris-HCl, 60 mM; $MgCl_2$, 10 mM; EDTA, 1 mM; coenzyme A (CoASH),

0.5 mM; and Triton X-100, 0.1%; pH 7.5) and added to the mixture at a final concentration of 200 μM,

while the final concentrations of FadD1 and its derivatives were 20 μM. The reaction mixture was

incubated at 37°C, and the reaction was stopped by placing sample tubes in boiling water for 5, 15,

30 and 60 min. *cis*-DA, BDSF or lauric acid (C12:0) levels were measured by LC-MS spectrometry.

Construction of in-frame deletion mutants and complementation

P. aeruginosa PAO1 was used as the parental strain for the generation of an in-frame deletion mutant

of *fadD1* and *dspI* by the following method described previously[14]. The primers used to generate

upstream and downstream regions flanking *fadD1* and *dspI* are listed in Table S7. For

complementation analysis, the coding regions of *fadD1* and *dspI* were amplified by PCR and cloned

into the plasmid pBBR1-MCS5, pBBR1-MCS2 and pLAFR3 using the primers listed in Table S6. The

resulting constructs were conjugated into the *P. aeruginosa* PAO1 mutant, *B. cenocepacia* H111

strain by electroporation[14,54].

Bacterial growth analysis

An overnight bacterial culture in LB medium was inoculated into fresh media to an OD_{600} of 0.01. A

10 mL cell suspension was grown in 50 mL conical centrifuge tubes at 37°C with shaking at 220 rpm.

Bacterial growth was determined by measuring the optical density at 600 nm every 4 hours. LB

medium was used as the negative control[47].

Competition assays in the mixed culture

The green fluorescent protein expression vector was used and transformed into the *P. aeruginosa*

PAO1 WT, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains. The mCherry fluorescent protein expression vector

was introduced into the *B. cenocepacia* H111 strain by electroporation. Then, the *P. aeruginosa* PAO1

WT, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains were cocultured with the *B. cenocepacia* H111 strain at an

initial ratio of 1:4 (vol/vol) at OD_{600} = 0.1 at 37°C with shaking at 220 rpm for 12 h. Then, the

quantification of BDSF was performed by using an LC-MS system, and the mixed cultures were

analyzed by a Spectra Max i3x multifunctional enzyme labeling instrument (Molecular Devices, CA,

USA)[57].

P. aeruginosa **phenotype assays**

For the analysis of biofilm formation[58], bacterial cells were grown overnight at 37°C and inoculated

into LB media (150 μL per well) in 96-well polypropylene microtiter plates. Microtiter plates with fitted

lids were incubated at 37°C with shaking at 200 rpm for 12 h. The plates were then washed to remove

planktonic cells and stained for 20 min with 1% (weight/vol) crystal violet (Sigma-Aldrich). After

washing with water, 200 μL of 95% (vol/vol) ethanol was added to the wells to release the stain.

Biofilm was determined by measuring the absorbance of the resulting solution at 570 nm.

Motility was determined on 0.3% semisolid agar (Becton, Dickinson and Company). Bacteria were

inoculated into the center of plates containing 0.8% tryptone, 0.5% glucose, and 0.3% agar. The

plates were incubated at 30°C for 18 h before the diameter of the colony was measured[54].

The pyocyanin assay was performed with the method described previously[49]. Briefly, bacteria were

cultured at 37°C for approximately 12 hours and centrifuged at 13,000 rpm for 1 min, and 1.5 mL

supernatants were collected and extracted with double volume chloroform with vigorous shaking at

room temperature for 30 min. The solvent phase was transferred to a new tube containing 1 mL of 1

N HCl (Sigma-Aldrich). The mixture was shaken gently to transfer pyocyanin to the aqueous phase.

The quantity of pyocyanin was determined by measurement of absorbance at 520 nm and

normalization against the cell density.

***B. cenocepacia* phenotype assays**

Biofilm formation in 96-well polypropylene microtiter dishes was assayed essentially as described

previously by *P. aeruginosa*. Swarming motility was determined on semisolid agar (0.3%). Bacteria

were inoculated into the center of plates containing 0.8% tryptone, 0.5% glucose, and 0.3% agar. The

plates were incubated at 30°C for 18 h before the diameter of the colony was measured.

Protease assays were performed following a previously described method[14]. Briefly, bacteria were

cultured at 37°C for approximately 12 hours. Cultures were centrifuged at 13,000 rpm for 5 min, and

the supernatants were removed and filtered through a 0.22 μm membrane. One hundred microliters

of supernatant was incubated at 30°C with an equal volume of azocasein (Macklin) dissolved in

proteolytic buffer (5 mg/mL) for 30 min. The reaction was stopped by the addition of 400 μL of 10%

(weight/vol) TCA (Macklin) buffer. After incubation for 2 min at room temperature, the mixture was

centrifuged at 13,000 rpm for 1 min to remove the remaining azocasein. Supernatants were removed

and mixed with 700 μL of 525 mM NaOH (Macklin). The absorbance of the azopeptide supernatant

was measured at a wavelength of 442 nm. Protease activity was obtained after normalization of the

absorbance against the corresponding cell density.

Cytotoxicity assays

Cytotoxicity was assessed by measuring the release of LDH from human A549 cells[54]. The A549 cell

line was cultured in Dulbecco's modified Eagle's medium (Sigma-Aldrich) at 37°C with 5% CO_2. A549

cells were seeded in 96-well plates at a density of 5×10^4 cells per well and grown until they reached

~90% confluence. Culture supernatants were removed, and the cell monolayer was washed once

with phosphate-buffered saline (PBS). Fresh bacterial cells cultured to an OD_{600} of 1.0 were washed

and diluted in DMEM. The diluted bacterial cells were then added to the A549 cell monolayers at a

multiplicity of infection (MOI) of approximately 1000. After incubation for 8 h, the cytotoxicity was

determined by measuring the released lactate dehydrogenase (LDH) in the supernatants using a

cytotoxicity detection kit (Roche).

Construction of reporter strains and measurement of β-galactosidase activity

The promoters of *lasR, lasI, rhlR, rhlI, mvfR* and *pqsA* were amplified using the primer pairs listed in

Table S6. The resulting products were inserted upstream of the promoter less *lacZ* gene in the vector

pME2-*lacZ*. Transconjugants were then selected on LB agar plates supplemented with tetracycline

and X-gal (Solarbio). Measurement of β-galactosidase activities was performed following previously

described methods[14,54,55].

Quantitative reverse transcription PCR assays

Bacterial cells were cultured at 37°C in LB broth to the logarithmic-growth phase (OD_{600}=1.0). Cells

were collected by centrifugation, and total RNA was extracted using an Eastep Super Total RNA

Extraction Kit (Promega). cDNA synthesis was performed using a HiScript III 1st Strand cDNA

Synthesis Kit (Yeasen). Primers for RT-qPCR are listed in Table S6. The RT-qPCR experiments were

performed on a QuantStudio™ 7 RT-qPCR System (Thermo Fisher) using SYBR qPCR Green Master

Mix (Yeasen). The expression of the target genes was normalized to the expression level of 16S RNA.

The relative transcript abundance was calculated by the $2^{-\Delta\Delta Ct}$ method[55]. At least three biological

replicates were performed per sample.

Quantitative analysis of QS signal production in *P. aeruginosa*

Bacterial cells were grown in LB medium overnight with agitation at 37°C. One liter of culture

supernatant was collected by centrifugation and extracted with an equal volume of ethyl acetate. The

crude extract (organic phase) was dried using a rotary evaporator and dissolved in methanol. All of

the above samples were kept at 4°C until analysis. Ultrahigh-performance liquid chromatography-

electrospray ionization tandem mass spectrometry (UHPLC-ESI-MS/MS) was performed in a

Shimadzu LC-30A UHPLC system with a Waters C18 column (1.8 μm, 150 × 2.1 mm) and a Shimadzu

8060 QQQ-MS mass spectrometer with an ESI source interface. The mass spectrometer was

operated in positive-ion mode. The mobile phase was prepared as 0.1% formic acid/water and

acetonitrile[57].

RNA-Seq analysis

RNA was isolated from *P. aeruginosa* strains grown to OD_{600}=1.0 by using the Eastep Super Total

RNA Extraction Kit (Promega, Madison, USA). The concentration was measured using a Qubit® RNA

Assay Kit in a Qubit® 2.0 Fluorometer (Life Technologies, CA, USA). The purity and integrity were

checked using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA) and the RNA Nano

6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA), respectively.

Sequencing libraries were generated using the NEBNext® Ultra™ Directional RNA Library Prep Kit

for Illumina® (NEB, USA) and then purified by the AMPure XP system. Finally, library quality was

assessed by using the Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples

was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS

(Illumina). After cluster generation, the library preparations were sequenced on an Illumina HiSeq

platform, and paired-end reads were generated. Trimmed sequence reads were aligned to the *P.*

aeruginosa PAO1 genome sequence using Bowtie2-2.2.3, and normalized read counts were

compared using HTSeq v0.6.1 as described previously[59,60]. Then, the FPKM of each gene was

calculated based on the 26 lengths of this gene, and the read count was mapped to the gene.

Electrophoretic gel mobility shift assay

Electrophoretic mobility shift assay (EMSA) experiments were performed as described

previously[53,54,61]. The DNA probes used for EMSA were prepared by PCR amplification using the

primer pairs listed in Table S6. The purified PCR products were 3-end-labeled with biotin following

the manufacturer's instructions (Thermo Fisher). The DNA-protein binding reactions were performed

according to the manufacturer's instructions (Thermo Fisher). A 5% polyacrylamide gel was used to

separate the DNA-protein complexes. After UV cross-linking, the biotin-labeled probes were detected

in the membrane using a Biotin luminescent detection kit (Thermo Fisher).

Chromatin immunoprecipitation sequencing analyses

Chromatin immunoprecipitation sequencing (ChIP-Seq) was performed according to previously published methods with some modifications[62]. ChIP-Seq analysis was conducted by the Wuhan IGENEBOOK Biotechnology Co. Ltd. (Wuhan, China). The anti-HIS antibody (ab9108, 1:1000, Abcam) was used for immunoprecipitation, and DNA fragments enriched by ChIP were sequenced on an Illumina HiSeq 2000. After removing sequencing adaptors and low-quality bases, the clean reads were mapped to the *P. aeruginosa* PAO1 genome using BWA software (version 0.7.15-r1140). The peak caller MACS (version, 2.1.1.20160309; *q* value, 0.05) was used to localize the potential binding sites of FadD1[63].

Molecular docking FadD1

The 3D structure of the double-stranded DNA was built in the MOE1 DNA/RNA Builder. Protein-DNA docking in the HDOCK2 server was used for molecular docking simulation of FadD1 and DNA[64,65]. The 3D structure of FadD1 (Q9HYU4) predicted by AlphaFold2 was set as the receptor, and the 3D structure of the DNA was set as the ligand[66]. The HDOCK server automatically predicts their interaction through a hybrid algorithm of template-based and template-free docking. First, the server performs template-based modeling to identify possible homologous DNA templates. Second, the intrinsic scoring function for protein-RNA interactions is integrated in template-free docking. Finally, the top 100 predicted complex structures are provided to users for download, of which the top 10 models will be visualized on the result web page. The intermolecular contacts from the most likely poses were further evaluated. The models of the complex and interface residues were analyzed using MOE. Molecular graphics were generated by PyMOL.

AutoDock and PyMOL were used for docking and binding site analysis. The crystal structure of FadD1 from the AlphaFold2 was used for docking with *cis*-DA by AutoDock software. To screen the potential binding sites, a grid box was prepared to wrap the entire macromolecule. The receptor grid was centered at center_x = -1.038, center_y = 1.49, center_z = 11.31, size_x = 126, size_y = 126, and size_z = 126. The exhaustiveness was set at 50. The top scoring ligand conformation was chosen,

and the potential binding sites were visualized in 3D by PyMOL. The potential binding sites were

visualized in 2D by Protein Plus (https://proteins.plus/).

Microscale thermophoresis assay

Protein-binding experiments were carried out with a Nano Temper 16 Monolith NT.115 instrument

(NanoTemper Technologies; www.nanotemper-technologies.com)[55]. In brief, the protein was labeled

with the L014 Monolith NT.115 Protein Labelling Kit (Nano Temper). The labeled protein and different

titers of unlabeled *cis*-DA were mixed and loaded onto standard treated silicon capillaries (Nano

Temper), and fluorescence was measured. The measurements were carried out at 20% LED power

and 40% MST power.

STATISTICAL ANALYSIS

No statistical method was used to predetermine the sample size. No data were excluded from the

analyses. The data are presented as the means ± standard deviations. Statistical analyses were

performed with Prism 8 software (GraphPad). Statistical significance is indicated as follows: *$p < 0.05$;

**$p < 0.01$; ***$p < 0.001$; ns = no significance (one-way ANOVA or two-way ANOVA). Biological

replicates and numbers of independent experiments were stated in the legends. All experiments

presented as representative LC-MS graphs or gels were repeated at least 3 times with similar results.

FIGURE LEGENDS

Figure 1. Effect of *fadD1* on the competitive capability of *P. aeruginosa* against *B. cenocepacia*.

(A) Analysis of the BDSF signal quenching ability of *P. aeruginosa* PAO1 (n = 3 biological replicates).

(B) SDS-PAGE of the purified FadD1 protein. (C) Analysis of the enzymatic quenching activity of

FadD1 on BDSF signal (*n* = 3 biological replicates). (D) The BDSF signal production of the *B.*

cenocepacia H111 strain cocultured with *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*)

strains (*n* = 3 biological replicates). (E) Green mean fluorescence intensity of the *P. aeruginosa* PAO1

wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains carrying the green fluorescent protein expression vector

(*n* = 3 biological replicates). (F) Red mean fluorescence intensity of the *B. cenocepacia* H111 strain

carrying the mCherry fluorescent protein expression vector (n = 3 biological replicates). The data are

the means ± standard deviations of three independent experiments. The *P. aeruginosa* PAO1 wild-

type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains carried the green fluorescent protein expression vector. The

mCherry fluorescent protein expression vector was carried by the *B. cenocepacia* H111 strain. The

P. aeruginosa PAO1 wild-type*,* Δ*fadD1* and Δ*fadD1*(*fadD1*) strains were cocultured with *the B.*

cenocepacia H111 strain at a ratio of 1:4 (v/v) at OD_{600} = 0.1. The statistical comparisons were

performed using two-way ANOVA (*p < 0.05; **p < 0.01; ***p < 0.001).

Figure 2. Effects of *fadD1* on the QS-regulated phenotypes and QS of *P. aeruginosa*. Analysis

of biofilm formation (A), swarming motility (B), pyocyanin production (C), and virulence (D) in *P.*

aeruginosa PAO1((A), n = 5 biological replicates; (B), (C), (D), n = 3 biological replicates). Effects of

fadD1 on the gene expression levels of *lasR* (E), *lasI* (F), *rhlR* (G), *rhlI* (H), *mvfR* (I) and *pqsA* (J) by

assessing the β-galactosidase activity of the promoter-*lacZ* transcriptional fusions and by RT-qPCR

(K), which are the signal molecule synthase-encoding genes and receptor genes of the *rhl*, *las* and

pqs QS systems in *P. aeruginosa* PAO1, respectively ((E), (F), (G), (H), (I), (J), n = 6 biological

replicates; (K), n = 3 biological replicates). The production of 3-oxo-C12-HSL (L), C4-HSL (M) and

PQS (N) in the *P. aeruginosa* PAO1 wild-type, Δ*fadD1* and Δ*fadD1*(*fadD1*) strains (n = 3 biological

replicates). The production of QS signals in the *P. aeruginosa* PAO1 wild-type strain was arbitrarily

defined as 100%. The data are means ± standard deviations of three independent experiments. The

statistical comparisons were performed using one-way ANOVA or two-way ANOVA ((A), (B), (C), (D),

(L), (M), (M), one-way ANOVA; (E), (F), (G), (H), (I), (J), two-way ANOVA; *p < 0.05; **p < 0.01; ***p

< 0.001).

Figure 3. Analysis of the binding of FadD1 to target gene promoters. (A) EMSA analysis of *in*

vitro binding of FadD1 to the promoter of *lasR,* in which the biotin-labeled 339-bp *lasR* promoter DNA

probe was used for the protein binding assay. A protein-DNA complex, represented by a band shift,

was formed when different concentrations of FadD1 protein were incubated with the probe at room

temperature for 30 min. (B) A potential FadD1 binding region was identified by ChIP-seq. The

conserved sequence is shown in orange. (C) Analysis of the binding between FadD1 and the mutated

lasR promoter with deletion of the FadD1 binding sequence AGGACGG. EMSA analysis was

performed *in vitro*. (D) Domain structure analysis of FadD1 (https://www.ebi.ac.uk/Tools/hmmer/). The

red box represents the leucine zipper motif (https://www.novopro.cn/tools/motifscan.html) in AMP-B

domain. (E) SDS-PAGE of the purified AMP-B domain and the AMP-B_C domain of the FadD1 protein.

EMSA analysis of *in vitro* binding of the AMP-B domain (F) and the AMP-B_C domain (G) to the

promoter of *lasR*. (H) SDS-PAGE of the purified mutated FadD1 protein (FadD1^{LM}: L275A L282A

L289A L296A). (I) EMSA analysis of the *in vitro* binding of the mutated FadD1 protein to the promoter

of *lasR*. *In trans* expression of the mutated *fadD1* could not restore biofilm formation (J), swarming

motility (K), and pyocyanin production (L) in the *P. aeruginosa* Δ*fadD1* strain ((J), $n = 6$ biological

replicates; (K), (L), $n = 3$ biological replicates). The EMSA experiments were performed three times,

and representative images from one experiment are shown. The data are the means ± standard

deviations of three independent experiments. The statistical comparisons were performed using one-

way ANOVA (**$p < 0.01$; ***$p < 0.001$; ns = no significance).

Figure 4. Effects of FadD1 on *cis*-DA-regulated phenotypes of *P. aeruginosa*. (A) MST analysis

of the binding of *cis*-DA to the FadD1. "Fnorm (‰)" indicates the fluorescence time trace changes in

the MST response. Complementation of the *dspI* mutant with *fadD1*. *In trans* expression of *fadD1*

complemented swarming motility (B) and pyocyanin production (C) in the *dspI*-deficient mutant.

Effects of *dspI* on the gene expression levels of *lasR* (D), *lasI* (E), *rhlR* (F), *rhlI* (G), *mvfR* (H), *pqsA*

(I) and the production of 3-oxo-C12-HSL (J), C4-HSL (K) and PQS (L) of the *P. aeruginosa* PAO1

strains. The production of QS signals in the *P. aeruginosa* PAO1 wild-type strain was arbitrarily

defined as 100%. $n = 3$ biological replicates. The data are means ± standard deviations of three

independent experiments. The statistical comparisons were performed using one-way ANOVA (*$p <$

0.05; **$p < 0.01$; ***$p < 0.001$).

Figure 5. Effects of *cis*-DA on the binding of FadD1 to the promoters of the target genes. (A)

EMSA analysis of the *in vitro* binding of FadD1 to the promoter of *lasR* with the addition of different

amounts of *cis*-DA. (B) The potential residues (which are shown as stick models and are labeled) of

FadD1 involved in *cis*-DA binding analyzed by AutoDocking software. The hydrogen bonds were

represented by yellow dashed lines. (C) SDS–PAGE gel electrophoresis of purified FadD1 variants.

MST analysis of the binding of BDSF to the (D) FadD1^{G488A} and (E) FadD1^{R552A}. EMSA analysis of

the *in vitro* binding of (F) FadD1^{G488A} and (G) FadD1^{R552A} to the promoters of *lasR* with the addition of

different amounts of *cis*-DA. The EMSA experiments were performed three times, and representative

images from one experiment are shown. Analysis of the perception of *cis*-DA by FadD1^{G488A} and

FadD1^{R552A}, *cis*-DA-regulated phenotypes: (H) swarming motility and (I) pyocyanin production in *P.*

aeruginosa (*n* = 3 biological replicates). The data are means ± standard deviations of three

independent experiments. The statistical comparisons were performed using one-way ANOVA (***p

< 0.001; ns = no significance).

Figure 6. Schematic representation of *cis*-DA-regulated QS signaling in *P. aeruginosa*. DspI

synthesized *cis*-DA signal molecule. FadD1, the signal receptor of *cis*-DA, perceives *cis*-DA signal

and regulates the transcription of *lasR*, which positively regulates the *las*, *rhl* and *pqs* QS systems in

P. aeruginosa. DspS is also the sensor of *cis*-DA involved in biofilm dispersion in response to *cis*-DA.

The individual QS circuits are highly interconnected.