Distribution, Prevalence, and Antibiotic Resistance of Non‑Tuberculous Mycobacteria on Hainan Island

Xinru Xu[1#], Jingjing Huang[2#], Hongkun Wang[1], Tianchen Xiang[1], Jun Liu[1], Li Zheng[1,3,*]

1. Department of Immunology, Key Laboratory of Tropical Translational Medicine of Ministry of Education, Hainan Medical University, Haikou, Hainan, PR China.
2. Institute of Tuberculosis Control and Research, Hainan Provincial Center of Disease Control and Prevention, Haikou, Hainan, PR China.
3. Hainan Medical University-The University of Hong Kong Joint Laboratory of Tropical Infectious Diseases, Haikou, Hainan, PR China.

*Correspondence:
Li Zheng (culturelily@sina.com)
#These authors contributed equally to this work.

ABSTRACT

Background: Non-tuberculous mycobacteria (NTM) represent a diverse group of environmental mycobacteria, with certain species implicated in infections affecting multiple tissues, including the lungs. The incidence of NTM-related diseases has been on the rise globally, particularly in developing countries. This study aimed to systematically investigate the ecological distribution, molecular identification, and antibiotic susceptibility profiles of NTM isolates in Hainan Island, furnishing a scientific basis for local control and prevention strategies. Methods: From 2015 to 2018, 158 suspected NTM samples were collected across 14 cities on Hainan Island. Following standardized sample processing and DNA extraction, molecular identification was conducted through PCR amplification of the rpoB and hsp65 genes, and sequencing, with species confirmation via GenBank database comparison. Phylogenetic analysis was performed to elucidate interspecies relationships and drug susceptibility testing was carried out for five antibiotics on M. abscessus isolates. Results: 118 NTM isolates were successfully identified, encompassing 24 distinct species. M. abscessus and M. intracellulare, accounting for 23.73% and 22.03% respectively, were the most prevalent. Phylogenetic analysis revealed clusters indicative of pathogenic and potentially pathogenic traits among different NTM species, with M. abscessus and M. fortuitum clustering as rapidly growing, pathogenic mycobacteria. Drug sensitivity testing indicated universal resistance of M. abscessus to doxycycline, yet high sensitivity to amikacin (71.4%) and linezolid (50.0%). Additionally, non-NTM species such as Nocardia thailandica, and Tsukamurella paurometabola were detected, suggesting potential misdiagnosis in clinical settings. Conclusion: This study provides a comprehensive dataset on the ecological distribution, molecular characterization, and antibiotic resistance of NTM in Hainan Island, highlighting the dominance of M. abscessus and M. intracellulare in local NTM infections.

Keywords: Non-tuberculous Mycobacteria (NTM); Species Identification; RNA Polymerase Beta Subunit (rpoB); Heat Shock Protein 65 (Hsp65)

INTRODUCTION

Non-tuberculous mycobacteria (NTM), belonging to the Mycobacteriaceae family, ubiquitously inhabit natural environments such as soil, water, and even the human body. With over 200 species documented, approximately 60 of which are known to be pathogenic, these organisms can affect a wide array of tissues including skin, soft tissue, lymphatic system, synovial fluid, joints, bones, and lungs[1]。 In immunocompromised individuals, pathogenic NTM can lead to opportunistic NTM pulmonary disease (NTM-PD) and extrapulmonary infections[2]. The global incidence of NTM infections has been increasing, particularly in developing countries [3]. In China, the prevalence of NTM infections rose from 4.9% in 1990 to 22.9% in 2010, with the proportion of NTM-related lung infections increasing from 4.24% to 12.68% by 2021 [4-5], Misdiagnosis of NTM as multidrug-resistant tuberculosis or co-infections with other chronic diseases contributes to escalating mortality rates[6-7]. The AIDS pandemic has further heightened recognition of NTM's clinical significance [8]; however, due to the absence of mandatory reporting systems for NTM infections in many nations, current epidemiological data is limited and potentially underestimates the actual burden[9-10] . An accurate understanding of NTM epidemiology and the implementation of effective surveillance systems are thus crucial for guiding control strategies.

Most regions in China have established comprehensive monitoring systems and research strategies for NTM, especially in developed areas like Shanghai, Beijing, and Guangdong, which feature extensive temporal coverage, diverse sample collection, and a wealth of NTM subspecies, leading to well-developed theoretical frameworks and even rare identifications [11-15]. NTM research of Hainan Island initiated in 2019[16], encountered challenges of geographical limitations and inadequate sample diversity, indicating a significant lag in research compared to other provinces. This study aims to enhance the breadth and depth of NTM investigation in Hainan, providing a robust foundation for more precise and targeted intervention strategies, ensuring their scientific rigor and effectiveness, and advancing the scientific frontiers in NTM prevention and control.

Mycobacterium avium complex (MAC), the most common slowly growing mycobacteria (SGM), is the predominant cause of NTM pulmonary disease globally[17], with strains like M. avium, M. intracellulare, M. chimaera, M. marseillense, M. bouchedurhonense, M. timonense, yongonense, and M. paraintracellulare recognized as pathogens [18]. The M. abscessus complex (MABC), dominated by M. abscessus subspecies, accounts for 65%-80% of rapidly growing mycobacteria (RGM) pulmonary infections[19] [20], with three key subspecies identified. M. kansasii ranks as the second most frequent NTM pathogen in some European countries[21], whereas in China, MAC and MABC prevail, specifically M. intracellulare and M. abscessus, along with other pathogens such as M. fortuitum, M. gordonae, and M. kansasii. In China, NTM distributions exhibit a southern, coastal, and warm-humid regional predominance over northern, inland, and cold-dry areas[2,5].

Drug susceptibility testing of numerous NTM isolates indicates clarithromycin (CLR) and amikacin (AMK) as highly effective against MAC, serving as first-line agents . MAC shows resistance to linezolid (LZD; 42.86%) and moxifloxacin (MFX; 25.32%), suggesting their use as second-line options. MABC exhibits high resistance to moxifloxacin, ciprofloxacin, and tobramycin (>95%), but retains susceptibility to tigecycline (98.53%) and amikacin (64.71%) [22], highlighting their potential as first-line therapies. M. fortuitum complex displays sensitivity to amikacin and moxifloxacin but is generally resistant to clarithromycin, doxycycline, and trimethoprim/sulfamethoxazole[23]. Given the suboptimal outcomes of current NTM treatments, the exploration of alternative antibiotics such as clofazimine, bedaquiline, delamanid, and tedizolid is warranted. These drugs have shown in vitro activity against specific NTM species, including M. intracellulare, M. avium (clofazimine-sensitive), M. abscessus, M. fortuitum, and M. chelonae (tedizolid-responsive), highlighting their therapeutic potential [24]. This research delves into the distribution and antibiotic resistance profiles of NTM in Hainan, elucidating the current NTM prevalence scenario and augmenting the province's NTM research corpus.

METHODS

Ethics Statement:
This study was conducted with the approval of the Ethics Committee of Hainan Medical University, under the ethics number HYLL-2021-170.

Study Population and Sample Collection:

The experimental samples were sourced from the Hainan Tuberculosis Prevention and Control Institution, covering the period between 2015-2018. A total of 158 sputum and bronchoalveolar lavage fluid specimens, clinically suspected of NTM infection, were collected from 14 cities and counties in Hainan Island, including Baisha, Baoting, Changjiang, Chengmai, Danzhou, Ding'an, Dongfang, Haikou, Ledong, Lingao, Lingshui, Qionghai, Sanya, and Wenchang.

Materials and Equipment:

Bacterial genomic DNA extraction kit (Tiangen Biotech, Beijing), Template DNA, buffer, dNTPs, TaqDNA polymerase (Tiangen Biotech, Beijing), Hsp65 and rpoB gene amplification primers (Tsingke Biotech, Beijing), PCR thermal cycler (Bio-Rad Laboratories, Shanghai), DNA recovery kit (Tiangen Biotech, Beijing). High-speed benchtop centrifuge (Eppendorf, Germany).

Clinical Sample Collection:

Patients are instructed to fast for at least 1-2 hours before sample collection. Before expectoration, patients perform a mouthwash with 0.1% spirulina solution three times, followed by rinsing with clean water, to minimize oral flora contamination. Sputum Production: Deep coughing is encouraged to ensure productive sputum collection into sterile plastic containers with a Spiral mouth, provided by the hospital, emphasizing the importance of collecting deep sputum such as purulent, mucoid, or blood-streaked specimens. Saliva samples may also be accepted if no productive sputum can be produced. Each patient provides three types of sputum samples (morning, night, and on-site) over a day. Morning and night sputa are self-collected at home, while the on-site sample is obtained at the hospital. Samples are transported to the tuberculosis laboratory within 4 hours of collection. If immediate transportation is not feasible, samples are refrigerated at 4℃ until delivery. In the absence of a provincial tuberculosis control institute, samples are sent to designated tuberculosis laboratories in each city for initial testing. Subsequently, positive cultures are forwarded to the Hainan Provincial Center for Disease Control and Prevention for confirmation and further analysis.

PCR-based Species Identification:

DNA was extracted from each sample to serve as a template for subsequent PCR reactions using primers targeting rpoB and hsp65 genes. Primer sequences were as follows: hsp65-F: 5'-ACC AAC GAT GGT GTG TCC AT-3', hsp65-R: 5'-CTT GTC GAA CCG CAT ACC CT-3', rpoB-F: 5'- GAC GAC ATC GAC CAC TTC GG -3', rpoB-R: 5'-GGG GTC TCG ATC GGG CAC AT-3'. Amplification was conducted in20 µL reaction mixtures containing 2 µL 10×PCR buffer, 200 µM dNTPs, 0.25 µM of each primer, and 1U TransTaqHiFiDNA polymerase (Tiangen Biotech, Beijing). Thermal cycling conditions comprised initial denaturation at 94°C for 5 minutes; 30 cycles of 94°C denaturation for 30 seconds, annealing at 60°C for 30 seconds, extension at 72°C for 1 minute; and a final extension at 72°C for 10 minutes. Products were analyzed by agarose gel electrophoresis, and DNA fragments were recovered using a gel extraction kit, and sent to Hainan Nanshan Biotechnology Co., Ltd. for sequencing. Ultimately, nucleotide sequences were aligned with GenBank references via the NCBI BLAST website on 04/22/2024 (https://blast.ncbi.nlm.nih.gov/Blast.cg).

Phylogenetic Tree Construction:

To explore evolutionary relationships between hsp65 and rpoB genes, multiple sequence alignment was performed using the Muscle algorithm in MEGA software (v11.0)

151 with default parameters. Neighbor-joining (NJ) method was applied in MEGA to construct
152 phylogenetic trees. To enhance tree topology reliability, a bootstrapping process involving
153 1000 iterations was executed. For intuitive comprehension, the phylogenetic tree was
154 refined and annotated using the Tree Visualization by One Table (TvBOT,
155 https://www.chiplot.online/tvbot.html, version 2.6) online tool.
156
NTM Susceptibility Testing:
According to Clinical and Laboratory Standards Institute (CLSI) guidelines, broth
microdilution assays were employed for in vitro antimicrobial susceptibility testing of
selected NTM isolates. Minimum inhibitory concentration (MIC) values require clinical
context interpretation and adjustment. MIC was defined as the lowest concentration
inhibiting visible growth of the tested strain. The MicroDST™ system (Yinked AUTOBIO
Diagnostics Co., Ltd., Zhuhai, China) was utilized for resistance testing of NTM isolates.
MIC breakpoints and sensitivity/resistance determination were interpreted per
manufacturer instructions. Five antibiotics were tested against M.abscessus isolates:
moxifloxacin, amikacin, linezolid, azithromycin, and doxycycline.

RESULTS

NTM Subspecies Characterization Findings
From 158 clinical samples suspected of NTM collected between 2015 and 2018,157
isolates were successfully species-identified, comprising 118 NTM (across 24 species) and
39 non-NTM organisms. Identification primarily relied on rpoB gene sequencing, with
hsp65 sequencing serving as a supplementary tool for conclusive determination. The
species distribution, as shown in Figure 1, was led by M. abscessus (28/118, 23.73%), M.
intracellulare (26/118, 22.03%), M. msimiae (17/118, 14.41%), M. paraintracellare (16/118,
13.56%), and M. colombiense (5/118, 4.24%). MTB(23), Nocardia
thailandica(3/16,18.75%), Tsukamurella paurometabola(2/16,12.5%), and Gordonia
jinghuaiqii(2/16,12.5%), were also identified. Detailed composition ratios are illustrated in
Supplemental Table 1.

179
180   **Figure 1** Distribution of non-tuberculous Mycobacteria isolated from patients with
181 suspected pulmonary NTM in Hainan Island, China. Abbreviation： NTM: non-tuberculous
182 Mycobacteria. RGM: rapidly growing mycobacteria; SGM: slowly growing mycobacteria.

183
rpoB Gene Phylogeny of NTM

A neighbor-joining (NJ) phylogenetic tree was constructed based on NTM rpoB
sequences, revealing clear speciation with pathogenic and potentially pathogenic species
clustering characteristics. M. fortuitum and M. abscessus grouped into a distinct cluster of
rapidly growing, pathogenic mycobacteria. Strong bootstrap support validated the tree's
reliability, with M. sherrisii and M. simiae closely related per hsp65 and rpoB analysis,
reflecting proximity to MAC, MABC, and M. simiae complexes(Figure 2). HSP 65 Gene
Phylogeny of NTM was shown in Supplemental Figure 1.

192
193   **Figure 2** Phylogenetic tree of rpoB gene from 118 NTM using the neighbor-joining
194 method with MEGA11.0 software. Tree-scale 0.007 represents the evolutionary distance
195 unit, and Bootstrap values represents the self-expanding value, which is used to test the
196 calculated evolutionary Tree branch confidence.
197
Spatial Distribution of NTM in Hainan Island

The 118 NTM isolates were distributed across 14 cities in Hainan, with distinct
patterns. Northeastern regions reported higher NTM presence compared to southwestern,
and SGMs outnumbered RGMs. MAC and MABC were dominant NTM groups, featuring
M. intracellulare and M. abscessus respectively. SGMs were detected in all except
Baoting, while RGMs were absent in Chengmai and Baisha. Key distributions included M.
intracellulare in Qionghai, Lingshui, Dongfang, and Danzhou, and M. abscessus in Dingan,

Wenchang, and Haikou. Species like M. algericus, M. sinensis, M. bacteremicum, and
others showed unique distributions. Refer to Figure 3 for comprehensive mapping.

207
208 **Figure 3** Distribution map of NTM in 14 cities and counties in Hainan Province (Baisha,
209 Baoting, Changjiang, Chengmai, Danzhou, Ding 'an, Dongfang, Haikou, Ledong, Lingao,
210 Lingshui, Qionghai, Sanya, Wenchang).
211
212
213 Susceptibility Test Results of M. abscessus

214   The MICs and breakpoints for seven antibiotics against 28 M. abscessus isolates are
215 detailed in Table 1. The isolates displayed high levels of antibiotic resistance, with
216 universal resistance to doxycycline but retained sensitivity to amikacin (71.4%) and
217 linezolid (50.0%). Complete resistance and sensitivity profiles are elaborated in Table 1.

Table 1 Drug susceptibility test of M.abscessus（n=28）

Antimicrobial agent	MIC （μg/mL）			Number （%）		
	MIC range	MIC 50	MIC 90	S	I	R
Amikacin	1-64	4	64	(20/28)71.4%	(3/28)10.7%	(5/28)17.9%
Linezolid	2-32	8	32	(14/28)50.0%	(8/28)28.6%	(6/28)21.4%
Moxifloxacin	0.125-16	4	16	(8/28)28.6%	(6/28)21.4%	(14/28)50.0%
Azithromycin	1-32	16	32	(9/28)32.1%	(5/28)17.9%	(14/28)50.0%
Doxycycline	16->128	>128	>128	0	0	(28/28)100%

Abbreviation：S, susceptible; I, intermediary; R, resistant

219
DISCUSSION

This study identified five major groups of NTM: MAC, MABC, M. simiae complex, M. fortuitum
complex, and other NTM species. MAC, a common causative agent of human NTM diseases

affecting various organs such as lungs, soft tissues, lymph nodes, and bone marrow, with MAC
pulmonary disease being the most prevalent[22]。, was also the primary NTM pathogen detected in
our study from pulmonary disease cases in Hainan, with a total of 51 isolates (comprising M.
intracellulare, M. paraintracellare, M. colombiense, M. timonense, M. avium). This aligns with
previous reports by Zhong [16] and Wang [22] who identified 149 and 154 MAC isolates respectively,
reinforcing the widespread distribution of MAC in Hainan. MABC, another dominant NTM group,
predominantly consisted of M. abscessus and its subspecies, differing slightly from Zhong and
Wang's findings where M. intracellulare was more prevalent. M. simiae complex emerged as the
third most frequent NTM in Hainan, further supporting its presence as a significant species in the
region, contrasting its less frequent detection in other provinces[2]. The NTM profile in Hainan
generally aligns with previous studies by Zhong [16,25],Wang[22],and Wu[26], showing a higher
prevalence of SGMs compared to RGMs. Huang[27] in Beijing also found M. intracellulare as the
dominant NTM, followed by M. abscessus and M. avium, mirroring our results where M. abscessus
and M. intracellulare dominate. M. abscessus is most prevalent in Hainan, likely due to the region's
hot and humid climate, which favors its survival over M. intracellulare, which thrives better in the
drier climate of Beijing.
Comparatively, Northern regions' literature reported M. intracellulare as the predominant
species with SGMs more common than RGMs, consistent with our Hainan findings. In central
China, M. intracellulare, M. avium, and M. kansasii were the main SGMs, while RGMs included M.
abscessus, M. chelonae, and M. fortuitum, similar to Hainan's leading species. However, only one
M. avium was isolated in Hainan, without M. kansasii, which is more frequently found in northern
regions. Southwest China's NTM composition also showed SGM dominance, led by M. avium, M.
intracellulare, and M. kansasii, again echoing Hainan's trend of more SGMs than RGMs.
Notably, M. kansasii and M. smegmatis were not detected in Hainan, despite their prevalence
in other parts of China. For instance, M. avium was the dominant NTM in Sichuan [28], unlike in
Hainan where M. abscessus dominates. Coastal southeastern regions, such as Fujian[29-30]、
Zhejiang[31-34], and Jiangsu[35-36] reported higher M. avium frequencies, contrasting with Hainan's
single M. avium isolate, highlighting geographic variation.
Considering Hainan and Taiwan's insular geography and similar latitudes, comparing NTM
distributions reveals insights. Huang et al. [37] found M. abscessus and MAC to be the most
common in Taiwan's NTM-pulmonary infections, mirroring Hainan's MAC dominance and M.
abscessus prevalence. However, the contrast in NTM distribution between northern and southern
Taiwan (where M. abscessus and MAC predominate respectively) differs from Hainan's northeast
and southwest (both MAC-dominated). Hainan's northeastern NTM abundance contrasts with
Taiwan's southern dominance, possibly influenced by its latitude, tropical/subtropical climates, and
topography. Hainan's central mountainous terrain might influence coastal NTM prevalence,
reflecting the island's central-high and peripheral-low geography. Overall, these findings
underscore the impact of geographical and environmental factors on NTM distribution patterns
across regions.
The distribution of NTM within Hainan Island exhibits a characteristic pattern, with fewer
occurrences in the central mountainous regions compared to the surrounding coastal areas.
Factors beyond climate, such as the mountainous terrain of the island's center, should also be
considered when examining this phenomenon. Remarkable disparities exist among different
regions in Hainan regarding dominant NTM species, with Danshan County dominated by M.
simiae, Qionghai, Changjiang, Lingao, Chengmai, and Chengmai featuring M. intracellulare
prominently, Dingan and Wenchang showing M. abscessus as the leading species, while Haikou
and Ledong highlighted M. paraintracellulare. The top three municipalities in terms of NTM
isolation frequency were Danshan, Qionghai, and Dingan, differing from previous research on
Hainan's NTM distribution from 2015-2018 [16], which listed Haikou, Dongfang, and Wenchang as
the top locations, and the incidence data from2018-2022 [26] placing Haikou, Dongfang, and
Wanning as the top three. These changes in regional distribution over time likely relate to
economic development, living conditions, lifestyle improvements, and public health awareness. As
opportunistic pathogens, NTM typically invades organs such as the lungs, skin, lymph nodes, and
bones when host immunity is compromised, with lung infections being most common[38-39].
Although NTM exhibits lower virulence and pathogenicity than MTB, their non-specific clinical
symptoms and radiological similarities to MTB lead to diagnostic challenges. Their natural and

acquired drug resistance further complicates clinical management[39] underscoring the importance
of NTM species identification and epidemiological investigations.
In Hainan, the hot and humid environment fosters NTM survival and dissemination, making
epidemiological studies in this high NTM-endemic region crucial for local control efforts. Given M.
abscessus's natural and acquired resistance to many common antibiotics, resistance testing to
identify effective treatments is essential; clinical practice often employs anti-tubercular drugs[40].
Guidelines[41] recommend clarithromycin, azithromycin, amikacin, and cefoxitin for M. abscessus.
Our susceptibility tests revealed rates of 71.4% for amikacin, 50.0% for linezolid, 32.1% for
azithromycin, and 28.6% for moxifloxacin. The high resistance to doxycycline (100%) highlights the
need for caution when using this drug clinically. The intrinsic resistance mechanisms of M.
abscessus include its waxy, opaque cell wall, drug transport systems, antibiotic modification, and
target gene polymorphisms[42]. Subspecies differences in resistance rates [43], particularly with
higher clarithromycin resistance in M. abscessus subsp. abscessus than subsp. massiliense,
indicating a need for further study to guide treatment.
Accurate and rapid NTM species identification is vital for disease management, reducing
iatrogenic harm, and preventing misdiagnosis [25]. Our study used rpoB sequencing to identify 24
NTM species, complementing Hsp65 sequencing that identified 18 species. While Hsp65 is a
highly expressed antigen and a powerful genotypic identifier [44], incorporating additional methods
like rpoB sequencing provides more precision. Compared to Hsp65, rpoB offers several
advantages [45] including sufficient information to distinguish the most recognized mycobacteria and
a greater variability for species differentiation. The rpoB-based phylogenetic tree, for example,
separates M. intracellulare from M. paraintracellare more accurately.
The study unexpectedly identified numerous non-NTM species, including previously
undetected MTB and species of Corynebacterium, Actinomadura, Gordonia, Tsukamurella,
Streptomyces, and Nocardia, all belonging to the actinobacteria phylum. Streptomyces
tubbatahanensis, not previously reported in China, suggests new perspectives in streptomycete
pathogenesis. Tsukamurella paurometabola, rarely pathogenic but previously isolated in Hainan [22]
indicates its opportunistic nature. Nocardia species, ubiquitous yet some highly pathogenic
especially in immunocompromised individuals, with Nocardia nova known as the second most
prevalent [46].　Gordonia species, isolated from various environments, can cause opportunistic
infections in immunocompromised hosts. Corynebacterium tuberculostearicum is omnipresent on
human skin, yet its health impacts require further investigation. [47]
As an expansion and deepening of previous work in Hainan, this study leverages the province-
wide TB control network for multi-dimensional NTM testing, achieving notable progress in
geographic coverage and species diversity. By involving hospitals at all levels and standardizing
specimen submission, a comprehensive and representative NTM repository was established,
surpassing single-center biases and enhancing sample representativeness. This collaborative,
cross-regional approach lays a firm foundation for elucidating regional NTM patterns and potential
epidemiological rules. It also underscores the direct influence of public health policies on
monitoring activities, emphasizing the need for flexibility and adaptability in future surveillance
systems to accurately track NTM epidemiology. Collectively, this study enriches Hainan's NTM
epidemiological data, informing targeted control strategies, and highlighting the value of
interdisciplinary and inter-institutional cooperation in addressing NTM public health challenges.

Conclusion

This study has provided an in-depth exploration into the ecological distribution and drug
resistance patterns of NTM in Hainan Island, highlighting M. abscessus and MAC as principal
pathogens driving NTM infections in the region, predominantly in Danshan, Qionghai, Dingan,
Dongfang, and Wenchang. Novel findings include the documentation of previously unreported
NTM species in Hainan, specifically M. novum, M. algericus, M. conceptionense, M. vanbaalenii,
M. eburneum, M. nebraskense, M. sinensis, M. bacteremicum, M. frederiksbergense, M. phlei.
Moreover, the study underscores the therapeutic potential of amikacin and linezolid against M.
abscessus.

FUNDING

This study was supported by the Natural Science Foundation of Hainan Province
(No:821QN0893, No: 820QN274) and the Natural Science Project of Hainan Provincial
Department of Education (No: Hnky2022-38, No: Hnky2020-38). XX, HW, TX, and YW
acknowledge support from the Innovation and Entrepreneurship Training Program for College
Students of Hainan Medical University (No:S202211810034, X202211810063).

Author Contribution

338   XX, HW, and TX conducted the experiments; JH was responsible for sample collection; XX,
339 JL, and JH performed the data analysis; XX and LZ wrote and revised the manuscript; LZ designed
340 the study.

REFERENCES

[1]Mortaz e Moloudizargari-M-Varahram-M. What Immunological Defects Predispose to Non-
tuberculosis Mycobacterial Infections?.[J], 2018.
[2]Xia Yu,Pengnian Liu,Guan Liu, et al. The prevalence of non-tuberculous mycobacterial infections
in mainland China: Systematic review and meta-analysis[J]. Journal of Infection, 2016, 73: 558-567.
[3]Jason E. Stout,Won-Jung Koh,Wing Wai Yew. Update on pulmonary disease due to non-
tuberculous mycobacteria[J]. International Journal of Infectious Diseases, 2016, 45: 123-134.
[4]Lixia Wang,Hui Zhang,Yunzhou Ruan, et al. Tuberculosis prevalence in China, 1990–2010; a
longitudinal analysis of national survey data[J]. The Lancet, 2014, 383: 2057-2064.
[5]Sun Qing,Yan Jun,Liao Xinlei, et al. Trends and Species Diversity of Non-tuberculous
Mycobacteria Isolated From Respiratory Samples in Northern China, 2014–2021[J]. Frontiers in
Public Health, 2022, 10.
[6]Nicolas Veziris,Claire Andréjak,Stéphane Bouée, et al. Non-tuberculous mycobacterial pulmonary
diseases in France: an 8 years nationwide study[J]. BMC Infectious Diseases, 2021, 21.
[7]Radha Gopalaswamy,Sivakumar Shanmugam,Rajesh Mondal, et al. Of tuberculosis and non-
tuberculous mycobacterial infections – a comparative analysis of epidemiology, diagnosis and
treatment[J]. Journal of Biomedical Science, 2020, 27.
[8]Robert Steinbrook. Tuberculosis and HIV in India[J]. New England Journal of Medicine, 2007, 356:
1198-1199.
[9]P López-Roa,E Aznar,J Cacho, et al. Epidemiology of Non-Tuberculous Mycobacteria isolated
from clinical specimens in Madrid, Spain, from 2013 to 2017[J]. European Journal of Clinical
Microbiology & Infectious Diseases, 2020, 39: 1089-1094.
[10]Huizheng Zhang,Ming Luo,Ke Zhang, et al. Species identification and antimicrobial susceptibility
testing of non-tuberculous mycobacteria isolated in Chongqing, Southwest China[J]. Epidemiology
and Infection, 2020, 149.
[11]LiuLin Luo,Li Liang,RanRan Zhang, et al. Using Vitek MS v3.0 To Identify Nontuberculous
Mycobacteria in Liquid Media in a Clinical Microbiology Laboratory[J]. Microbiology Spectrum, 2022,
10.
[12]Shuai Chen Wei Piao Chuang Yue Hong Jin Li Li Qi Jiang Le Cai Ji. Increasing Trends and
Species Diversity of Nontuberculous Mycobacteria in A Coastal Migrant City-Shenzhen, China.[J].
Biomedical and Environmental Sciences : Bes, 2022, 35(2): 146-150.
[13]Qi Liu,Jingli Du,Huiru An, et al. Clinical characteristics of patients with non-tuberculous
mycobacterial pulmonary disease: a seven-year follow-up study conducted in a certain tertiary
hospital in Beijing[J]. Frontiers in Cellular and Infection Microbiology, 2023, 13: 1205225.
[14]Qing Zhang,Heping Xiao,Liping Yan. PCR-reverse blot hybridization assay in respiratory
specimens for rapid detection and differentiation of mycobacteria in HIV-negative population[J]. BMC
Infectious Diseases, 2021, 21.
[15]Jing-jing Huang,Ying-xing Li,Ying Zhao, et al. Prevalence of nontuberculous mycobacteria in a
tertiary hospital in Beijing, China, January 2013 to December 2018[J]. BMC Microbiology, 2020, 20.
[16] Yeteng Zhong, Mingguan Lin, Chong Lin, et al. [Characteristics of non-tuberculous
Mycobacterial infection in suspected tuberculosis patients in Hainan area] [J]. Chinese Journal of
Infection Control, 2019, 18(08): 701-707.

[17]D. Rebecca Prevots,Theodore K. Marras. Epidemiology of Human Pulmonary Infection with
Nontuberculous Mycobacteria[J]. Clinics in Chest Medicine, 2015, 36: 13-34.
[18] Yuanchun Li. Analysis of the Subspecies Identification, Drug Resistant Profiles and the
Correlation between the Drug-resistant Phenotype and Genotype in Mycobacterium Avium Complex
[D]. Beijing Hospital of the Ministry of Health, 2021
[19]David E. Griffith,Timothy Aksamit,Barbara A. Brown-Elliott, et al. An Official ATS/IDSA Statement:
Diagnosis, Treatment, and Prevention of Nontuberculous Mycobacterial Diseases[J]. American
Journal of Respiratory and Critical Care Medicine, 2007, 175: 367-416.
[20]Qi Guo,Haiqing Chu,Meiping Ye, et al. The Clarithromycin Susceptibility Genotype Affects the
Treatment Outcome of Patients with Mycobacterium abscessus Lung Disease[J]. Antimicrobial
Agents and Chemotherapy, 2018, 62.
[21]Wouter Hoefsloot,Jakko van Ingen,Claire Andrejak, et al. The geographic diversity of
nontuberculous mycobacteria isolated from pulmonary samples: an NTM-NET collaborative study[J].
European Respiratory Journal, 2013, 42: 1604-1613.
[22]Jieying Wang,Zhuoling Chen,Yuni Xu, et al. Screening and Drug Resistance Analysis of Non-
Tuberculous Mycobacteria in Patients with Suspected Pulmonary Tuberculosis on the Hainan Island,
China[J]. Infection and Drug Resistance, 2023, Volume 16: 463-476.
[23]Anna Borek,Anna Zabost,Agnieszka Głogowska, et al. New RAPMYCOI SensititreTM
Antimicrobial Susceptibility Test for Atypical Rapidly Growing Mycobacteria (RGM)[J]. Diagnostics,
2022, 12: 1976.
[24]Wei-Chang Huang,Ming-Chih Yu,Yi-Wen Huang. Identification and drug susceptibility testing for
nontuberculous mycobacteria[J]. Journal of the Formosan Medical Association, 2020, 119: S32-S41.
[25] Yeteng Zhong, Zhuolin Chen, Jieying Wang, et al. [Species identification of non-tuberculous
Mycobacteria from patients with pulmonary infection in Hainan] [J]. Disease Surveillance, 2022,
37(08): 1053-1058.
[26] Haifeng Wu, Minhua Weng, Rui Liu, et al. [Epidemiological status and clinical characteristics of
225 cases of non-tuberculous mycobacterial pulmonary disease in Hainan Province] [J]. China
Tropical Medicine, 2024, 24(04): 463-467.
[27]Prevalence of nontuberculous mycobacteria in a tertiary hospital in Beijing, China, January
2013 to December 2018.[J].
[28] Jiao Yang, Tianping Li, Chao Yang, et al. [Distribution and drug resistance of non-tuberculous
mycobacteria in the skin and soft tissue infections] [J]. Chinese Journal of Laboratory Diagnosis,
2020, 24(08): 1266-1268.
[29]Huang Mingxiang,Lin Yinxia,Chen Xinchao, et al. The value of gene chip detection of
bronchoalveolar lavage fluid in the diagnosis of nontuberculous mycobacterial lung disease[J].
Annals of Palliative Medicine, 2021, 10: 6438-6445.
[30]Evaluation of the MeltPro Myco Assay for the Identification of Non-Tuberculous Mycobacteria[J].
[31]Jintian Xu,Ping Li,Shengchao Zheng, et al. Prevalence and risk factors of pulmonary
nontuberculous mycobacterial infections in the Zhejiang Province of China[J]. Epidemiology and
Infection, 2019, 147.
[32]Hong Fang,Yanwan Shangguan,Hao Wang, et al. Multicenter evaluation of the biochip assay for
rapid detection of mycobacterial isolates in smear-positive specimens[J]. International Journal of
Infectious Diseases, 2019, 81: 46-51.
[33]Yelei Zhu,Wenya Hua,Zhengwei Liu, et al. Identification and characterization of nontuberculous
mycobacteria isolated from suspected pulmonary tuberculosis patients in eastern china from 2009
to 2019 using an identification array system[J]. The Brazilian Journal of Infectious Diseases, 2022,
26: 102346.
[34] Dandan Xia, Lianpeng Wu. [Analysis of identification results of 967 strains of non-tuberculosis
Mycobacteria in Wenzhou from 2016 to 2021] [J]. Chinese Journal of Health Laboratory Technology,
2022, 32(09): 1065-1068.
[35]Chunmei Hu,Lili Huang,Min Cai, et al. Characterization of non-tuberculous mycobacterial
pulmonary disease in Nanjing district of China[J]. Bmc Infectious Diseases, 2019, 19.
[36] Haixia Zhang, Jing Huang, Yuanyuan Xiao, et al. Distribution and drug resistance analysis of
Mycobacterium tuberculosis and nontuberculosis mycobacteria in Nanjing from 2017 to 2020 [J].
Infectious Disease Information, 2022, 35(03): 259-263. 263.

[37]Hung-Ling Huang,Meng-Hsuan Cheng,Po-Liang Lu, et al. Epidemiology and Predictors of NTM
Pulmonary Infection in Taiwan - a Retrospective, Five-Year Multicenter Study[J]. Scientific Reports,
2017, 7.
[38] Mahsa Dastranj,Abbas Farahani,Abdolrazagh Hashemi Shahraki, et al. Molecular identification
and distribution of non-tuberculous mycobacteria isolated from clinical specimens by PCR-
sequencing method in the West of Iran[J]. The Clinical Respiratory Journal, 2017, 12: 996-1002.
[39] Xinyu Yang, Yanfeng Zhao, Junli Yi, et al. [Analysis of risk factors and drug resistance of
clinical isolates of non-tuberculous mycobacteria in Beijing prevention and control institutions in
2019] [J]. Journal of Clinical Pulmonary Medicine, 2022, 27(08): 1148-1153.
[40]Dongping Wang,Wenhong Lin,Hongyan Cheng, et al. Clinical Characteristics and Antimicrobial
Susceptibility of Mycobacterium intracellulare and Mycobacterium abscessus Pulmonary Diseases:
A Retrospective Study[J]. Canadian Journal of Infectious Diseases and Medical Microbiology, 2022,
2022: 1-9.
[41]Charles L Daley,Jonathan M Iaccarino,Christoph Lange, et al. Treatment of Nontuberculous
Mycobacterial Pulmonary Disease: An Official ATS/ERS/ESCMID/IDSA Clinical Practice Guideline[J].
Clinical Infectious Diseases, 2020, 71: 905-913.
[42]Rose Lopeman,James Harrison,Maya Desai, et al. Mycobacterium abscessus: Environmental
Bacterium Turned Clinical Nightmare[J]. Microorganisms, 2019, 7: 90.
[43] Xinyu Wang. Subspecies distribution and drug resistance characteristics of Mycobacterium
abscessus complex clinical isolates [D]. Guangzhou Medical University, 2023
[44]Dyshelly Nurkartika Pascapurnama,Nabilla Zavitri,Raspati Koesoemadinata, et al. Identification
of Significant Pathogenic Nontuberculous Mycobacteria Species from Presumptive TB Patients
Using Partial hsp65 Gene Sequencing[J]. Infection and Drug Resistance, 2023, Volume 16: 6923-
6930.
[45] Bum-Joon Kim,Seung-Hyun Lee,Mi-Ae Lyu, et al. Identification of Mycobacterial Species by
Comparative Sequence Analysis of the RNA Polymerase Gene (rpoB)[J]. Journal of Clinical
Microbiology, 1999, 37: 1714-1720.
[46]Carleigh N. Bruce,Neda Esmaili. Periocular Infection With Nocardia nova Following
Dacryocystorhinostomy With Silicone Stent Intubation[J]. Ophthalmic Plastic & Reconstructive
Surgery, 2023, 39: e202-e204.
[47] PayalJoglekar ClaytonDeming Katherine P.Lemon Heidi H.Kong Julia A.Segre SeanConlan
Nashwa M.Ahmed. Genomic characterization of the C. tuberculostearicum species complex, a
ubiquitous member of the human skin microbiome[Z]: Cold Spring Harbor Laboratory, 2023.
472