Genetic and cellular basis of impaired phagocytosis and photoreceptor degeneration in

CLN3 disease.

Jimin Han[1-3], Sueanne Chear[4], Jana Talbot[4], Vicki Swier[6], Clarissa Booth[6], Cheyenne Reuben-

Thomas[1-3], Sonal Dalvi[1-3], Jill M. Weimer[6,7], Alex W. Hewitt[5], Anthony L. Cook[4,] Ruchira

Singh[1-3*]

1. Department of Ophthalmology, University of Rochester, Rochester, NY, 14620, USA

2. Department of Biomedical Genetics, University of Rochester, Rochester, NY, 14620, USA

3. Center for Visual Science, University of Rochester, Rochester, NY, 14620, USA

4. Wicking Dementia Research and Education Centre, University of Tasmania, Tasmania,

Australia

5. Menzies Institute for Medical Research, University of Tasmania, Tasmania, Australia

6. Pediatrics & Rare Diseases Group; Sanford Research, Sioux Falls, South Dakota, USA

7. Department of Pediatrics; Sanford School of Medicine, University of South Dakota, Sioux

Falls, South Dakota, USA

*__Lead Contact and Corresponding Author:__

Ruchira Singh, University of Rochester,

601 Elmwood Ave, Rochester, NY 14642

(585) 276-7338, ruchira_singh@urmc.rochester.edu

Abstract

CLN3 Batten disease (also referred to as Juvenile Neuronal Ceroid Lipofuscinosis; JNCL) is a

lysosomal storage disorder that typically initiates with retinal degeneration but is followed by

seizure onset, motor decline and premature death. Patient-derived iPSC-RPE cells show

defective phagocytosis of photoreceptor outer segments (POSs). Because modifier genes are

implicated in CLN3 disease, our goal here was to investigate a direct link between *CLN3*

mutation and POS phagocytosis defect by diseased RPE cells. Isogenic control and *CLN3* mutant

stem cell lines were generated by CRISPR-Cas9-mediated biallelic deletion of exons 7 and 8. A

transgenic Yucatan miniswine model of CLN3 disease ($CLN3^{\Delta 7\text{-}8/\Delta 7\text{-}8}$ pigs) was also used to study

the impact of $CLN3^{\Delta 7\text{-}8/\Delta 7\text{-}8}$ mutation on POS phagocytosis. POS phagocytosis by cultured RPE

cells was analyzed by Western blotting and immunohistochemistry. Electroretinogram, optical

coherence tomography and histological analysis of $CLN3^{\Delta 7\text{-}8}$ and wild-type miniswine eyes were

carried out at 6-, 36-, or 48-month age. $CLN3^{\Delta 7\text{-}8/\Delta 7\text{-}8}$ RPE (*CLN3* RPE) displayed reduced POS

binding and consequently decreased uptake of POS compared to isogenic control RPE cells .

Furthermore, wild-type miniswine RPE cells phagocytosed POS isolated from $CLN3^{\Delta 7\text{-}8/\Delta 7\text{-}8}$

(*CLN3*) Yucatan miniswine less efficiently than POS isolated from wild-type miniswines.

Consistent with decreased POS phagocytosis, lipofuscin/autofluorescence was decreased in

CLN3 miniswine RPE at 36 months-of-age and was followed by almost complete loss of

photoreceptors at 48 months of age. These results show that $CLN3^{\Delta 7\text{-}8/\Delta 7\text{-}8}$ mutation (that affects

up to 85% patients) is independently sufficient for promoting both impaired POS phagocytosis

and photoreceptor cell loss in CLN3 disease.

Keywords: CLN3 disease, juvenile neuronal ceroid lipofuscinosis, Batten disease, retinal

pigment epithelium, phagocytosis, photoreceptor outer segment, gene-editing.

Introduction

CLN3 disease is the most common form of Batten disease (neuronal ceroid lipofuscinoses or NCLs) that develops in early childhood. Symptoms include progressive loss of vision and neurodegeneration of the brain, ultimately leading to premature death[1,2]. Most affected CLN3 disease patients (~75%) have a 966 bp deletion that entirely removes exon 7 and 8 from the *CLN3* transcript[3]. Despite this, there is significant variability in the disease phenotype of CLN3 disease patients, including those harboring the same common *CLN3* mutation[4-6]. The lack of genotype-phenotype relationship in CLN3 disease has suggested a contribution of modifier genes to the disease development and progression[4].

Murine models have been widely used as an *in vivo* model of CLN3 disease[7-11]. However, despite some evidence of molecular and pathological alterations, there is poor recapitulation of retinal degeneration phenotype in these models[12-14]. To overcome the lack of visual degeneration of the existing CLN3 models, a transgene miniswine model was developed that carries the orthologous common 966 bp deletion found in most human patients[96]. Importantly, *CLN3[Δ7-8/Δ7-8]* Yucatan miniswine (*CLN3* miniswine) shows both progressive neuronal loss, motor dysfunction, and vision impairment phenotypes similar to those seen in individuals witch CLN3 disease[15].

Although retina degeneration is a prominent pathology in CLN3 disease, there is limited knowledge of the disease mechanism(s) underlying retinal degeneration in CLN3 disease. Clinical and histopathologic studies on CLN3 disease retina have shown autofluorescent lipopigment buildup in neurons with degeneration of several retinal cell type(s), including the retinal pigment epithelium (RPE)[16-18]. Furthermore, histopathologic analyses of CLN3 disease

donor eyes showed decreased autofluorescent material, lipofuscin, in the RPE and in contrast

elevated autofluorescence in the photoreceptor cell layer[16,19-21]. We have recently replicated

decreased RPE autofluorescence in the CLN3 disease donor eyes in patient-derived human

induced pluripotent stem cell (hiPSC)- RPE cells (harboring the common 966 bp deletion) post-

chronic daily exposure to a physiological load of photoreceptor outer segments (POSs)[20].

However, because this study compared CLN3 disease patient-derived hiPSC-RPE cells to non-

isogenic control hiPSC-RPE cells, it is unclear if the observed POS phagocytosis defect in CLN3

disease is a direct consequence of *CLN3* mutation. Note that, as previously mentioned modifier

genes have strongly implicated in molecular alterations observed in CLN3 disease[4].

CRISPR-based gene editing has been routinely used for *in vitro* correction of genetic mutations

in human pluripotent stem cells, including hiPSCs and human embryonic stem cells (hESCs)[22-24].

In fact, CRISPR-based gene editing of hiPSCs and hESCs have been routinely utilized to

decipher genotype-phenotype relationship in several retinal diseases[25-28]. For example, we have

previously used gene-corrected hiPSC lines from patients with Doyne honeycomb retinal

dystrophy (DHRD) to show that R345W mutation in EFEMP1 is causal for drusen accumulation

in DHRD[29]. Similarly, CRISPR-Cas9 mediated repair *of CLCN2* mutation rescued dysfunctional

POS phagocytosis by iPSC-RPE cells from patients with *CLCN2*-associated retina

degeneration[30]. Also, Neural retina-specific leucine zipper protein (NRL)-deficient hESC-retina

organoids engineered via CRISPR-Cas9 gene editing showed a direct role of NRL in rod

photoreceptor specification[31]. Overall, gene-editing strategies including both correction of

genetic defect in patient hiPSCs and introduction of disease-associated mutation in control

hESC/hiPSCs provides a suitable approach to definitively link the independent contribution of

genetic variants to molecular and pathological defects in a human-based cellular model.

Therefore, to link disease-causing *CLN3* mutation to the observed impaired phagocytosis in

CLN3 disease, here, we used CRISPR/Cas9 gene editing to generate isogenic H9 hESC lines

carrying homozygous deletion of exons 7 and 8 in CLN3 (referred to here in as $CLN3^{\Delta7\text{-}8/\Delta7\text{-}8}$).

Additionally, we utilized a large animal model of CLN3 disease to study the impact of common

966 bp *CLN3* deletion mutation on POS phagocytosis and autofluorescence/lipofuscin

accumulation.

Materials and Methods

Availability of data and materials

Request for original data files and reagents and materials used in this study will be fulfilled by contacting the corresponding author, Dr. Ruchira Singh.

Procurement and use of hESC line.

H9 hESCs were obtained from WiCell and used under approval from the Tasmania Health and Medical Human Research Ethics Committee (#13502) and Institutional Regulatory Board (IRB) (RSRB00056538)at the University of Rochester and conformed with the ethical norms and the declaration of Helsinki.

Animals.

WT and transgenic ($CLN3^{Δ7-8/Δ7-8}$) miniswine[8] used in these studies were obtained from Exemplar Genetics and all animal studies were conducted following the approval of the Institutional Animal Care and Use Committees at Exemplar Genetics (Protocol # MRP2018-004).

CRISPR-Cas9-mediated editing of *CLN3*

Editing of *CLN3* to delete exons 7 and 8 was done as previously described[32,33]. Briefly, 8×10^5 H9 hESCs in single-cell suspension were resuspended in 100 µL electroporation buffer from the Human Stem Cell Nucleofector Solution 2 kit (Lonza, VPH-5022). Alt-R Cas9 Electroporation enhancer (1.1 µM) (IDT, #1075915) and assembled CRISPR/Cas ribonucleoprotein consisting of 1.4 µM dual crRNA (*Table 1*) (IDT) and 1.2 µM Alt-R S.p. Hifi Cas9 Nuclease V3 (IDT,

#1081060) were mixed with cells and transferred to a nucleofection cuvette. Electroporation was

conducted using Amaxa Nucleofector 2b Device (Lonza) with Nucleofector program B-016.

Electroporated cells were seeded at low density (2,000 cells/well) in a Matrigel (Corning,

FAL354277)-coated 6 well-plates with CloneR (STEMCELL Technologies, #05889)/

mTeSR^{TM}1 media (STEMCELL Technologies, #85850). When single-cell colonies were

approximately 200 µm in diameter, each colony was picked and seeded into 2 wells of duplicate

Matrigel-coated 96 well-plates with one plate for cryopreservation and the other for genotyping.

PCR screening and genotyping

When cells from the genotyping 96-well plate were 50-60% confluent, DNA extraction was done

using QuickExtract DNA Extraction Solution (Epicentre, #E09050) according to manufacturer's

instructions. Subsequently, 50 ng of DNA extract was amplified through PCR reaction using

KAPA HotStart PCR Kit, with dNTPs (Millenium Science, ROC-07958897001), and CLN3 F1

and CLN3 R1 primers that span the targeted region (*Table 1*).

Characterization of pluripotency

Embryonic stem cell colonies at 80% confluency were dissociated with ReLeSR (STEMCELL

Technologies, #05872). Cell aggregates were transferred into ultra-low attachment 6-well plates

in Complete KSR EB medium (KnockOut™ SR (Life Technologies, #10828-028) 20%,

DMEM/F-12, GlutaMAX™ supplement (Life Technologies, #10565018) 79%, MEM Non-

Essential Amino Acids solution 1% (Life Technologies, #11140050), 2-mercaptoethanol 0.1 mM

(Sigma, M6250) for embryoid body (EB) formation. Medium was refreshed every 2 days. On

day 4, EBs were plated onto Matrigel-coated wells for further differentiation in Complete KSR

EB medium for 16 days. Pluripotency markers were analyzed by quantitative real-time PCR

(qRT-PCR) and immunocytochemistry.

For qRT-PCR analyses, total RNA was isolated from positive clones with the RNeasy Plus Mini

Kit (Qiagen, #74134) according to the manufacturer's protocol. Reverse transcription of 1 μg

RNA was performed using Omniscript RT kit (Qiagen, #205111). CLN3 transcript was

amplified, and PCR products were sequenced with primers, CLN3 F2 and CLN3 R2 that span

exons 6-9 (*Table 1*).

For immunocytochemical analyses, hESC colonies were passaged and cultured on Matrigel-

coated coverslips in mTeSR1 medium until 60% confluency. Cells were then fixed with 4%

paraformaldehyde in PBS for 20 minutes at room temperature. Fixed cells were incubated for

one hour in blocking solution (5% fetal bovine serum (FBS) in PBS and 5% FBS with 0.1%

Triton X-100 in PBS for extracellular and intracellular markers respectively) followed by

incubation with the primary antibodies for NANOG, OCT4, TRA-1-60 and SSEA4 overnight at

4°C. This was followed by incubation in host-specific secondary antibody for one hour at room

temperature and thereafter mounting and visualization by fluorescent microscopy. Further details

of primary antibody concentration and source can be found in *Table 2*.

Virtual karyotyping

Genomic DNA (200 ng) from isogenic H9 *CLN3*$^{\Delta7-8/\Delta7-8}$cell lines were analyzed for copy number

variation using Illumina HumanCytoSNP-12 beadchip array. B allele frequency (BAF) and log R

ratio (LRR) of each single nucleotide polymorphism (SNP) marker were collected from

GenomeStudio (Illumina, USA). These were used for CNV analyses which was performed using

PennCNV[34] and QuantiSNP[35] with default parameter settings. Genomic regions having at least

20 contiguous single nucleotide polymorphisms (SNPs) or genomic regions with SNPs spanning

at least 1 MB were designated as chromosomal copy number variation[36].

RPE differentiation and culture from hESCs

Differentiation of hESC to RPE was performed as previously described[20,37]. Briefly, embryoid

bodies (EBs) were generated from colonies of hESCs cultured in either mTeSR™ or mTeSR™

(Stemcell technologies). On day 6 of EB culture, EBs were plated onto laminin-coated tissue

culture plates and fed with neural induction medium or NIM whose composition is described

in[20,37]. The cell culture medium was switched to retinal differentiation medium (RDM) on day

14. RDM composition is described in[20,37]. hiPSC-RPE cells were dissected in patches from

adherent cultures around day 60-90 when characteristic RPE morphology could be observed. The

RPE patches at passage P0 were seeded and passaged onto either again a non-permeable plastic

24-well plate or onto transwell (Corning) insert (0.4 µm pore size). RPE cells at < P4 were used

in all experiments.

Primary miniswine RPE culture

Primary miniswine RPE cultures were performed as previously described[39,40]. Briefly, after

enucleation, eyes were placed in 0.2 % povidone-iodine solution for 10 minutes on ice followed

by 5 rinses in 1000U/mL penicillin-streptomycin solution. The anterior portion of the eye and the

retina was next removed (note the retina was used for POS isolation) and eyecups were

transferred to 6-well plates filled with 1ml of prewarmed 0.5% trypsin with 5.3 mM EDTA in

HBSS without calcium and magnesium. Subsequently, the 6 well plates were placed in a 37°C

incubator for 30 minutes. At the end of the incubation, RPE cells were collected in a 15mL tube

containing prewarmed media (1X DMEM with 4.5 g/L glucose, L-glutamine, and sodium

pyruvate and 10% FBS) and gently centrifuged (300 rcf) to collect the RPE cell pellet. The RPE

cell pellet was resuspended and plated in a T25 flask in culture media (1X DMEM with 4.5 g/L

glucose, L-glutamine, and sodium pyruvate, 1% NEAA, 1% penicillin/streptomycin) containing

10 %FBS. Once the RPE cells have reached confluence, the FBS concentration in the culture

media was reduced to 1% and subsequently RPE cells were passaged and replated onto laminin-

coated 24-wells or transwells.

POS phagocytosis assay

POS phagocytosis assay was performed as previously described20. Bovine POSs were obtained

commercially from InVision BioResources (Cat. #98740, Seattle, WA) and porcine POSs

prepared from wild-type and CLN3 miniswine as previously described[38-40]. Note that POS was

isolated eyes from three WT and four CLN3 miniswine eyes of ~30-36 months of age.

For evaluating POS uptake, as described earlier[20], mature monolayer of RPE cell in culture were

fed unlabeled POS or FITC-labeled POS (~10-40 POS/RPE cell)[2] for two hour at 37°C with 10%

FBS media supplementation. Thereafter, to remove POS (unbound) on the RPE cell surface,

RPE cells were thoroughly washed with 1X PBS. Subsequently, RPE cells were either harvested

for Western blotting or processed for immunocytochemistry experiments. Quantitative analysis

of the number of FITC-POS phagocytosed included a consideration of FITC-particle size.

Specifically, a threshold of <5µm allowed us to evaluate the amount bound, ingested POS but

eliminate aggregated POS from the analyses.

For evaluating POS binding and POS internalization, we used a previously published protocol[41]

that was also utilized to evaluate POS binding versus internalization in CLN3 disease hiPSC-

RPE[20]. Briefly, the position of FITC-particles relative to ZO1 was used to evaluate the amount of

bound versus internalized FITC-POS particles[42]. Specifically, to analyze the relative position of

the FITC-POS within the cell, FITC-POS fed RPE cells post-immunocytochemical labeling with

ZO1 antibody were imaged as confocal z-stacks spanning the entire height of the RPE cells. The

images were subsequently analyzed with ImageJ software to determine the number of apical

versus basal FITC-POS particles relative to ZO1.

RPE Autofluorescence measurements

As previously described[20,43], for *in vitro* experiments evaluating autofluorescence accumulation,

RPE cells were fed POS daily (~20-40/RPE cell/day) for 14 days. At day 14, to remove any

unbound POS, RPE monolayer was thoroughly washed with 1X PBS. RPE cells were then

immediately fixed and processed by immunocytochemistry where the red channel (ex: 546 nm

and em: 560-615 nm) was exclusively used for evaluation of RPE autofluorescence without any

antibody staining. Quantitative analyses of autofluorescent particles were performed per DAPI-

stained cells for count and per viewing area for area covered by the autofluorescent particles

using ImageJ/FIJI and Microsoft Excel software.

For analysis of autofluorescence in miniswine eye samples, miniswine RPE whole mounts were

utilized. Miniswine eyes were injected with 4% PFA following enucleation for 1h prior to

dissection to remove the anterior portion and vitreous. The remaining retina/RPE eyecup was

submerged in 4% PFA for a fixation of at least 4h at 4°C. Following two 1X PBS washes,

samples were prepared for wholemount staining by carefully removing the RPE/choriocapillaris

from the sclera. The RPE wholemount was then permeabilized/blocked in blocking buffer (1%

bovine serum albumin (ImmunoReagents, Inc.), 0.2% Triton-X-100, 0.2% Tween-20) for 1h

room temperature. Subsequently RPE Samples were washed and incubated with DAPI dye

(Hoechst 33342; Life Technologies) for 15-30min at room temperature. After one last wash with

1X PBS, RPE samples were mounted onto slides with Prolong Gold (Life Technologies P36930)

and cover slipped and imaged by confocal microscopy in the red channel (ex: 546 nm and em:

560-615 nm).

Western blot

Western blotting was performed as described[20]. Briefly, RPE cells were lysed in Radio-

Immunoprecipitation Assay (RIPA) buffer containing protease inhibitor cocktail (Sigma-

Aldrich). Subsequently, protein quantification was performed with Pierce BCA assay and equal

amount of protein was loaded for each sample and resolved on 4-20% Tris-HCl gradient gels,

followed by transfer onto Polyvinylidene Fluoride (PVDF) membranes. The PVDF membrane

was subsequently blocked with 5% dry milk in 1X PBS or commercially bought blocking buffer

(LI-COR) for 1h at room temperature followed by primary antibody incubation overnight at 4°C.

After primary antibody incubation, PVDF membrane was washed with 0.1% Tween-PBS and

incubated in host-specific HRP or IRDye-conjugated secondary antibody (1:10,000) in 5% dry

milk 0.1% Tween-PBS for 1h at room temperature. Azure C500 (Azure Biosystem) and Image

Studio were used for analysis and quantification of the Western blot images.

Note that all primary antibodies used for Western blotting analyses are listed in *Table 2*.

Immunocytochemistry

Immunocytochemical analysis was performed as previously described[20,43]. Briefly, RPE on

transwells were permeabilized and blocked in blocking buffer (10% normal donkey serum, 0.1%

Triton-X-100) for 1h at room temperature. This was followed by overnight incubation in primary

antibody in 0.5X blocker buffer at 4°C. Next day, RPE in transwells were washed and

subsequently incubated with host-specific Alexa-conjugated secondary antibody (1:500) in 0.5X

blocking buffer for 1h at room temperature. The RPE in transwells were next incubated with

Hoechst 33342 (Life Technologies) for 15 min to stain the nuclei, then washed in 1X PBS for 5

min. Subsequently, transwell membranes were cut out and mounted onto slides with Prolong

Gold (Life Technologies) and imaged with a confocal microscope (LSM 510 META, Zeiss, or

Eclipse Ti2, Nikon). Confocal microscopy images were analyzed with ImageJ/FIJI software. All

primary antibodies used can be found in *Table 2*.

Histological analyses of miniswine retina

Whole eyes were fixed for ~3 weeks in 10% neutral buffered formalin. Subsequently, the retina

was dissected and the midperiphery region of the retina was used for processing and analyses.

Specifically, the dissected retina tissue was further fixed at room temperature in 4% PFA

for 1h. After 1X PBS washes, the retina was processed for plastic embedding (Technovit 7100

kit; Electron Microscopy Sciences) as previously described[15]. Retina sections were cut at 3 μm

thick and dried on a slide warmer. Multiple Stain Solution (Polysciences, Inc.) was used to stain

the retina sections on slides, which were then mounted with permount (Fisher Scientific) and

imaged by light microscopy.

Transmission electron microscopy analyses

Transmission electron microscopy (TEM) was performed as previously described[20]. Briefly, RPE grown on transwells were fixed overnight in a solution containing 0.1 M sodium cacodylate, 2.5% glutaraldehyde and 4% paraformaldehyde. Next, RPE samples on transwell membrane were embedded in epoxy resin and processed to obtain 60 nm thick RPE section, then imaged at the Electron Microscopy Core at the University of Rochester using a Hitachi H-7650 Transmission Electron Microscope.

TER measurements

EVOM2 volt-ohm meter (World Precision Instruments) was used to measure TER of RPE grown in transwell inserts as described[44] using an). A TER measurements were reported as resistance per area or $\Omega.cm^{-2}$ after blank subtraction,

ERG and OCT measurements

ERG was performed using RET*eval*® device (LKC technologies) under anesthesia as previously described [15]. Briefly for scotopic ERG measurements, animals were dark adapter for 20 mins. Photopic ERG measurements were conducted under standard illumination with exposure to room light for over 10 minutes. Scotopic and photopic a-wave and b-wave amplitudes data from each eye for each analysis was measured separately but graphical representation of an individual measurement (e.g., scotopic a wave) shows averaged data for both eyes together.

Retinal layer thickness of miniswine via OCT was assessed using the Leica instrument spectral-domain OCT system (Bioptigen Envisu R2200; Leica, Danaher, Washington, DC). Prior to

imaging, eyes were dilated to increase the field of view using topical tropicamide 1% and

ophthalmic 2.5% phenylephrine eye drops. For each eye, high density scans consisted of

1000x100x1 (A scans x B scans x repeated B scans) for averaging, with a 12mm x 12mm

volume scan centered upon the retina superior to the optic nerve. Multiple images were obtained

from each eye for analysis. OCT analysis to access retinal layer thickness was performed on 3

scans per eye using Bioptigen InVivoVue and ImageJ software

Experimental set-up, data analyses and statistical testing.

For all experiments comparing isogenic control and *CLN3* RPE (with the exception of

immunocytochemistry for RPE signature proteins), parallel age-matched cultures of RPE

monolayer in transwell inserts that displayed TER > 150 $\Omega.\text{cm}^{-2}$ were utilized. Data is

presented as mean ± sem. Two-tailed unpaired student's t-test was used to compute significance

and Microsoft Excel was used to plot the graphs. For quantitative analysis of confocal

microscopy data, at least 5 distinct images (different viewing areas) per sample were used in

each experiment. In each individual figure n refers to the biological replicate sample size for data

from either isogenic control versus *CLN3$^{Δ7-8/Δ7-8}$* hESC-RPE cells derived from distinct EB

differentiations, or *ex vivo* or *in vivo* analyses of age-and sex matched wild-type and *CLN3*

miniswine retina and RPE.

Results

Generation and characterization of isogenic control and CLN3$^{Δ7-8/Δ7-8}$ (*CLN3*) hRPE cells .

To generate *CLN3$^{Δ7-8/Δ7-8}$* isogenic cell lines, we utilized a well-characterized hESC line, H9[45].

CLN3$^{Δ7-8/Δ7-8}$ intron 6 and the other within intron 8 which are spaced 968 bp apart in the *CLN3*

gene (*Figure 1a*). Our first attempt of CRISPR editing yielded a H9 *CLN3$^{+/Δ7-8}$* cell line with

monoallelic deletion of exons 7-8 in CLN3 gene (*Figure. 1b*). To obtain a biallelic deletion of

exons 7-8 in the *CLN3* gene, we repeated CRISPR editing on the H9 *CLN3$^{+/Δ7-8}$* cell line.

Subsequent PCR revealed two truncated DNA bands (*Figure. 1b*), which despite having

different deletion sizes had complete deletion of exons 7-8 from both CLN3 alleles. The

successful biallelic deletion of exons 7-8 was confirmed through sequencing of the *CLN3*

transcript using cDNA and primers targeting regions outside exons 7-8 in the homozygous H9

CLN3$^{Δ7-8/Δ7-8}$ cell line (*Figure. 1c*).

Genome-wide copy number variation analysis demonstrated that the H9 *CLN3$^{Δ7-8/Δ7-8}$* isogenic

cells have normal diploid karyotypes (46,XX). No chromosomal alterations were detected in the

edited cells when compared to the unedited control cells based on analysis of the B allele

frequency and Log R ratio (*Figure. S1*).

Pluripotency of isogenic H9 and H9 *CLN3$^{Δ7-8/Δ7-8}$* cell lines was confirmed by positive reactivity

for pluripotency markers OCT4, NANOG, TRA-1-60, and SSEA4 (*Figure. S2*). Quantitative

real-time polymerase chase reaction (qRT-PCR) analysis of embryoid bodies also demonstrated

expression of genes specific for all three germ layer markers (endoderm, ectoderm, and

mesoderm) (***Figure. S2***).

Isogenic control and *CLN3*$^{Δ7-8/Δ7-8}$ H9 ESC lines successfully differentiated to RPE cells using

our previously described protocols[20,29,37,46]. Light microscopy and transmission electron

microscopy (TEM) of control RPE and *CLN3*$^{Δ7-8/Δ7-8}$ RPE (referred henceforth to as *CLN3*

hRPE) showed the expected cell morphology including presence of elongated RPE microvilli in

both control and *CLN3* hRPE cultures (***Figure 1d***). Confocal microscopy imaging post-

immunocytochemical analyses also showed similar and expected localization of tight junction

protein, ZO1, in both control and *CLN3* hRPE cultures (***Figure 1e***). Similarly, consistent with

polarized expression of specific proteins in RPE cells, orthogonal view of confocal microscopy

images of control and *CLN3* hRPE showed apical localization of microvilli protein, EZR, and

basolateral localization of BEST1 (***Figure 1e***). Qualitative Western blotting analyses showed

robust presence of RPE signature proteins, BEST1, CRALBP, EZR, MERTK, RPE65, as well as

loading control (ACTN) in control and *CLN3* hRPE cells. In addition, in agreement with

transepithelial fluid movement and formation of a polarized epithelial monolayer[20,47], light

microscopy images showed presence of fluid domes in both control and *CLN3* hRPE cultures

(***Figure 1f***). Lastly, transepithelial epithelial resistance (TER) recordings showed formation of a

tight epithelial barrier with both control and *CLN3* hRPE developing physiological TER of ~150

$Ω$*cm^{-2} expected for RPE monolayer *in vivo*[48].

Overall, CRISPR-Cas9 editing, we were able to generate pluripotent isogenic control and

CLN3$^{Δ7-8/Δ7-8}$ H9 hESC lines. Notably, isogenic control versus *CLN3* hRPE cells showed similar

morphology with presence of several RPE signature proteins and the expected RPE cell polarity

and tight junction characteristics akin to native human RPE *in vivo*.

Disease causing *CLN3* mutation is independently sufficient to promote impair POS binding

and consequently impair POS phagocytosis by RPE cells.

We have previously shown reduced POS binding and consequently impaired POS phagocytosis

by patient-derived CLN3 disease hiPSC-RPE cells compared to control hiPSC-RPE cells[1]. To

link CLN3 function to POS phagocytosis by RPE cells, we evaluated uptake of non-diseased

POS by isogenic control versus *CLN3* hRPE cells.

Confocal microscopy imaging post-immunocytochemical analyses of control and *CLN3* mutant

RPE cells that were fed FITC-labeled POS (~20-40 POS/RPE cell) for 2h showed fewer number

of phagocytosed POSs in *CLN3* hRPE cells compared to parallel cultures of control RPE cells

(***Figure 2a-2c***). Autofluorescence accumulation (lipofuscin) in RPE cells is a consequence of

POS phagocytosis and we have previously been able to show disease-associated reduction in

lipofuscin (autofluorescence) in CLN3 disease patient-derived hiPSC-RPE cells after chronic (14

day) daily supplementation of a physiological dose of POS by measuring autofluorescence levels

in the red channel (ex: 546 nm and em: 560-615 nm). Consistently, confocal microscopy

analyses of autofluorescence levels in the red channel in POS-fed RPE cells (~40 POS per day

for 14 days) showed decreased accumulation of autofluorescent material (count, area) in the

CLN3 hRPE cells compared to control RPE cells (***Figure 2d-2f***).

To confirm that reduced POS phagocytosis by *CLN3* mutant RPE cells is a direct consequence

of defective POS binding by *CLN3* mutant RPE cells, we next utilized a previously published

protocol[20,41] to analyze bound versus internalized FITC-labeled POS (~20-40 POS/RPE cell)

post-2h POS feeding in isogenic control versus *CLN3* hRPE cells (***Figure 3***). Compared to

parallel cultures of control RPE cells, *CLN3* mutant hRPE that displayed decrease in total POS

uptake (***Figure 2b, 2c and 3a***) also showed decrease in both the amount of apically localized

(bound) and basally localized (internalized) FITC-POS particles relative to ZO1 (***Figure 3b-3e***).

However, no difference in the internalized FITC-POS relative to bound FITC-POS was seen in

control versus *CLN3* hRPE (***Figure 3f***).

Overall, these data confirm that *CLN3* mutation is independently sufficient to promote impaired

POS binding and consequently defective POS phagocytosis by CLN3 disease RPE cells.

***CLN3* mutant POS are phagocytosed less efficiently by wild-type RPE cells.**

Our published paper[20] and results presented in this study (***Figure 2, 3***) show that disease-causing

CLN3 mutation leads to impaired POS phagocytosis by RPE cells. However, these data lack the

consideration of diseased POS to the phagocytosis process. Note that POS disorganization is an

early pathological hallmark of CLN3 disease retina in living human eye[2,16,49,50]. Furthermore,

several studies have implicated inflammation and oxidative stress in CLN3 disease[51-56] and it has

previously been shown that oxidized POS and POS exposed to free radicals are phagocytosed

less efficiently by RPE cells[57,58]. Therefore, we next investigated the phagocytosis of control

versus *CLN3* mutant POS.

We chose to utilize *CLN3* mutant POS isolated from *CLN3$^{Δ7-8/Δ7-8}$* Yucatan miniswine in these

experiments as it is well-established that POS in stem cell-derived retina organoids lack

reproducibility and consistency potentially due to the lack of RPE[59,60]. Parallel cultures of wild-

type miniswine RPE primary were fed either unlabeled wild-type miniswine POS or *CLN3*

miniswine POS (~10 POS/RPE cell) for 2 h. Subsequently, phagocytosis of POSs was measured

by quantitative Western blotting analyses measuring the levels of RHO, a POS-specific protein,

within the wild-type miniswine RPE cells (***Figure 4a, 4b***).

CLN3 miniswine POS were phagocytosed less efficiently by wild-type miniswine RPE cultures

compared to wild-type miniswine POS (***Figure 4c, 4d***). In fact, wild-type miniswine POSs were

phagocytosed over 5-fold more efficiently than the *CLN3* miniswine POS (***Figure 4c, 4d***). These

data suggests that CLN3 POS contribute to defective POS phagocytosis by RPE cells in CLN3

disease.

<u>Disease-associated reduction in autofluorescence/lipofuscin levels precedes loss of outer</u>

<u>nuclear layer in *CLN3* miniswine model</u>.

Lipofuscin, the autofluorescence accumulation that accumulates in RPE cells because of

incomplete digestion of POS by RPE cells, is reduced in CLN3 disease donor RPE cells[16,21].

CLN3 disease and CLN3 hRPE cells phagocytose (uptake) less POS and consequently have

reduced RPE autofluorescence/lipofuscin compared to control RPE cells[20] (***Figure 2e, 2f***). Thus,

we utilized autofluorescence levels in the spectral wavelength consistent with lipofuscin, as a

surrogate for efficacy of POS phagocytosis by RPE cells in *CLN3* miniswine retina.

Longitudinal analyses of autofluorescence levels (EX: EM) in the RPE cells of age- and sex-

matched wild-type versus *CLN3* miniswine retina showed similar RPEautofluorescence in wild-

type and *CLN3* miniswine retina at 6 months of age (***Figure 5a-5c***). In contrast, there was

reduced lipofuscin in the *CLN3* miniswine RPE compared to wild-type miniswine RPE at 36

months of age (***Figure 5a-5c***). Note that in our previous study[15], *CLN3* miniswine had shown

reduced photopic and scotopic a wave at ~30-36 months of age. ERG analyses of *CLN3*

miniswine retina at 3 and 6 months of age, when there is no reduction in RPE

lipofuscin/autofluorescence (***Figure 5a-5c***), showed similar photopic and scotopic a-wave

amplitudes in wild-type and *CLN3* miniswine retina (***Figure S3***). Similarly, photopic and

scotopic b-wave recordings were similar in wild-type and *CLN3* miniswine retina at 3 and 6

months of age (***Figure S3***).

Longitudinal histological analyses of *CLN3* miniswine retina at the 36- and 48-month timepoint

showed the presence of POSs and multiple layers of outer nuclear layer (ONL) nuclei in the 36

month *CLN3* miniswine retina. In contrast, POSs were almost entirely gone with a near-complete

loss of ONL in *CLN3* miniswine retina at 48 months of age. Note that optical coherence

tomography (OCT) analyses of wild-type versus *CLN3* miniswine retina showed no significant

changes in any of the retina cell layer including the ONL at 6 months of age, although the nerve

fiber layer (NFL) was slightly reduced when measurements were taken 1 mm from the optic

nerve (***Figure S4***).

Overall, these data show that reduction in RPE lipofuscin/autofluorescence coincides with

photopic and scotopic visual deficits and precedes overt lack of POSs and ONL loss in *CLN3*

miniswine retina.

Discussion

CLN3 disease is the most common form of neuronal ceroid lipofuscinoses (NCLs). Apart from neurological symptoms, vision loss due to retinal degeneration is an early clinical hallmark of CLN3 disease. In a previous study, we used a CLN3 disease patient-derived hiPSC-RPE to show impaired POS phagocytosis[20]. Because gene modifiers have been strongly implicated in CLN3 disease[4], we generated isogenic control and *CLN3* H9 hESC lines (hESCs; H9), as well as using *CLN3* miniswine model, to show that impaired POS phagocytosis is a direct consequence of disease-causing *CLN3* mutation ($CLN3^{\Delta7-8/\Delta7-8}$), which contributes to reduced RPE lipofuscin/autofluorescence in CLN3 disease[20,49].

Histopathological and clinical characterizations of CLN3 disease have suggested that retina degeneration starts at the POS[2,61]. Furthermore, CLN3 disease donor eyes show decreased RPE lipofuscin/autofluorescence[49]. Despite this, there have been limited investigation of POS phagocytosis in CLN3 disease[12,13,20]. This is presumably due to the postulation that photoreceptor cell death and the consequent lack of POS lead to reduced lipofuscin accumulation in RPE cells[49]. Here, we provide evidence that POS phagocytosis dysfunction and subsequently reduced RPE lipofuscin in CLN3 disease is a consequence of both primary RPE dysfunction (***Figure 2, 3***) and POS alterations (***Figure 4***). Specifically, by comparing the uptake of non-diseased POS by parallel cultures of isogenic control and *CLN3* hRPE cells (***Figure 2, 3***), we show that cell autonomous RPE dysfunction due to *CLN3* mutation ($CLN3^{\Delta7-8/\Delta7-8-8}$) is sufficient for reduced POS phagocytosis in CLN3 disease. Similarly, by comparing the uptake of wild-type versus *CLN3* POS by wild-type RPE cells (***Figure 4)***, we show that POS alterations in CLN3

disease also independently contribute to decreased POS phagocytosis by RPE cells in CLN3

disease.

A limitation of our study is that we did not validate the role of POS alterations in impaired POS

phagocytosis by RPE cells in the human-relevant hESC model. Similarly, we did not validate the

POS phagocytosis defect *in vivo* in the *CLN3* miniswine model. This was due to consideration of

the most optimal experimental approach and reduction in the use of tissue from a large animal

model of the disease. The hESC-RPE, which have been consistently shown to possess properties

of human RPE cells *in vivo*[62,63], provided a suitable model system to investigate the ability of

control versus *CLN3* RPE cells to phagocytose shed POSs. Similarly, the availability of *in vitro*

protocols to assess POS binding versus internalization[20,41] allowed us to interrogate the precise

role of POS binding versus POS internalization in impaired POS phagocytosis by *CLN3* hRPE

cells (***Figure 3***).

In contrast to RPE cells, the reproducibility of POS differentiation and maturation in the stem

cell-derived retina organoid is variable[59,60]. Therefore, to avoid confounding variability due to

experimental model system, we relied on primary POSs isolated from wild-type versus *CLN3*

miniswine in experiments evaluating the role of diseased POS on the efficiency of POS

phagocytosis by RPE cells in CLN3 disease. Note that recent advances in protocol for POS

isolation from native retina allowed us to obtain sufficient POS from 3-4 miniswines.

Furthermore, we were able to use RPE from the animals sacrificed for POS isolation to compare

autofluorescence accumulation between wild-type versus *CLN3* miniswine RPE cells.

Reduced lipofuscin/autofluorescence accumulation preceded the loss of POSs and

photoreceptors in the *CLN3* miniswine model and coincided with the earliest timepoint of

scotopic and photopic visual deficit in *CLN3* miniswine (***Figure 5, S3***)[15]. This is consistent with

pathological studies of human eyes that have shown neuronal depletion in CLN3 disease retina

starting at the outer segments of the photoreceptors and proceeding inwards to the ganglion cell

layer[2,61]. Apart from highlighting a central role of photoreceptor-RPE interaction in CLN3

disease (that needs further investigation), these data further highlight the value of *CLN3*

miniswine model for studying the retina pathobiology of CLN3 disease.

Conclusions

Using both *in vivo* (*CLN3* miniswine) and *in vitro* (isogenic hESC control and *CLN3* lines), we show that the most common disease-causing mutation in CLN3 disease ($CLN3^{\Delta ex7/\Delta 8/\Delta ex7/\Delta 8}$) leads to POS alterations and primary RPE dysfunction that contribute to defective POS phagocytosis and reduced RPE autofluorescence/lipofuscin in CLN3 disease. Furthermore, reduced RPE autofluorescence/lipofuscin coincides with the earliest timepoint of scotopic and photopic visual deficit and precedes the loss of POSs and photoreceptors in the in a large animal model of CLN3 disease.

Funding

This work was supported by R01EY028167 (R.S.), R01EY030183 (R.S.), R01EY033192 (R.S.), funding from ForeBatten Foundation to both Singh and Weimer laboratory, Research to Prevent Blindness Unrestricted Challenge Grant to Department of Ophthalmology at University of Rochester and University Research Award given to R.S. by the University of Rochester. We would also like to acknowledge the Batten Disease Support and Research Association Australia (RGP204), assistance and Royal Hobart Hospital Research Foundation (17-205) to A.L.C. and A.W.H.

Author's contribution

Conceptualization and methodology: A.L.C., A.W.H., J.M.W.; R.S.

Experiment and analyses: A.L.C., A.W.H., C.B., C.R.T., J.H., J.M.W.; J.T., S.C., S.D., R.S.

Writing: Initial draft: R.S.

Writing – review & editing: A.L.C., A.W.H., C.B., C.R.T., J.H., J.M.W.; J.T., S.C., S.D., R.S.

Resources, A.L.C., A.W.H., J.M.W.; R.S.

Acknowledgments

We would like to acknowledge support of Exemplar Genetics staff in conducting studies (ERG,

OCT, tissue recovery) that was performed at the Exemplar Genetics facilities.

References

1.   Haltia M, Goebel HH. The neuronal ceroid-lipofuscinoses: a historical introduction.

Biochim Biophys Acta. Nov 2013;1832(11):1795-800. doi:10.1016/j.bbadis.2012.08.012

2.   Anderson GW, Goebel HH, Simonati A. Human pathology in NCL. *Biochim Biophys*

Acta. Nov 2013;1832(11):1807-26. doi:10.1016/j.bbadis.2012.11.014

3.   Mole SE, Williams RE, Goebel HH. Correlations between genotype, ultrastructural

morphology and clinical phenotype in the neuronal ceroid lipofuscinoses. *Neurogenetics*. Sep

2005;6(3):107-26. doi:10.1007/s10048-005-0218-3

4.   Lebrun AH, Moll-Khosrawi P, Pohl S, et al. Analysis of potential biomarkers and

modifier genes affecting the clinical course of CLN3 disease. *Mol Med*. 2011;17(11-12):1253-

61. doi:10.2119/molmed.2010.00241

5.   Sakti DH, Cornish EE, Fraser CL, et al. Early recognition of CLN3 disease facilitated by

visual electrophysiology and multimodal imaging. *Doc Ophthalmol*. Jun 2023;146(3):241-256.

doi:10.1007/s10633-023-09930-1

6.   Schulz A, Kohlschütter A, Mink J, Simonati A, Williams R. NCL diseases - clinical

perspectives.   *Biochim   Biophys   Acta*.   Nov   2013;1832(11):1801-6.

doi:10.1016/j.bbadis.2013.04.008

7.   Mitchison HM, Bernard DJ, Greene NDE, et al. Targeted Disruption of the Cln3 Gene

Provides a Mouse Model for Batten Disease. *Neurobiology of Disease*. 1999/10/01/

1999;6(5):321-334. doi:https://doi.org/10.1006/nbdi.1999.0267

8.   Finn R, Kovács AD, Pearce DA. Altered sensitivity of cerebellar granule cells to

glutamate receptor overactivation in the Cln3(Δex7/8)-knock-in mouse model of juvenile

neuronal ceroid lipofuscinosis. *Neurochem Int*. May 2011;58(6):648-55.

doi:10.1016/j.neuint.2011.02.003

9.   Cotman SL, Vrbanac V, Lebel LA, et al. Cln3(Deltaex7/8) knock-in mice with the

common JNCL mutation exhibit progressive neurologic disease that begins before birth. *Hum

Mol Genet*. Oct 15 2002;11(22):2709-21. doi:10.1093/hmg/11.22.2709

10.   Weimer JM, Benedict JW, Getty AL, et al. Cerebellar defects in a mouse model of

juvenile neuronal ceroid lipofuscinosis. *Brain Res*. Apr 17 2009;1266:93-107.

doi:10.1016/j.brainres.2009.02.009

11.   Pontikis CC, Cella CV, Parihar N, et al. Late onset neurodegeneration in the Cln3-/-

mouse model of juvenile neuronal ceroid lipofuscinosis is preceded by low level glial activation.

Brain Res. Oct 15 2004;1023(2):231-42. doi:10.1016/j.brainres.2004.07.030

12.   Zhong Y, Mohan K, Liu J, et al. Loss of CLN3, the gene mutated in juvenile neuronal

ceroid lipofuscinosis, leads to metabolic impairment and autophagy induction in retinal pigment

epithelium. *Biochim Biophys Acta Mol Basis Dis*. 10 01 2020;1866(10):165883.

doi:10.1016/j.bbadis.2020.165883

13.   Wavre-Shapton ST, Calvi AA, Turmaine M, et al. Photoreceptor phagosome processing

defects and disturbed autophagy in retinal pigment epithelium of Cln3Δex1-6 mice modelling

juvenile neuronal ceroid lipofuscinosis (Batten disease). *Hum Mol Genet*. Dec 15

2015;24(24):7060-74. doi:10.1093/hmg/ddv406

14.   Seigel GM, Lotery A, Kummer A, et al. Retinal pathology and function in a Cln3

knockout mouse model of juvenile Neuronal Ceroid Lipofuscinosis (batten disease). *Mol Cell

Neurosci*. Apr 2002;19(4):515-27. doi:10.1006/mcne.2001.1099

15.   Swier VJ, White KA, Johnson TB, et al. A novel porcine model of CLN3 Batten disease

recapitulates clinical phenotypes. *Dis Model Mech*. Jun 12 2023;doi:10.1242/dmm.050038

16.   Bensaoula T, Shibuya H, Katz ML, et al. Histopathologic and immunocytochemical

analysis of the retina and ocular tissues in Batten disease. *Ophthalmology*. Sep

2000;107(9):1746-53.

17.   Mole SE, Williams RE. Neuronal ceroid-lipfuscinosis. Gene Reviews; 1993-2014.

18.   Katz ML, Johnson GS, Tullis GE, Lei B. Phenotypic characterization of a mouse model

of juvenile neuronal ceroid lipofuscinosis. *Neurobiol Dis*. Feb 2008;29(2):242-53.

doi:10.1016/j.nbd.2007.08.017

19.   Katz ML, Rodrigues M. Juvenile ceroid lipofuscinosis. Evidence for methylated lysine in

neural storage body protein. *Am J Pathol*. Feb 1991;138(2):323-32.

20.   Tang C, Han J, Dalvi S, et al. A human model of Batten disease shows role of CLN3 in

phagocytosis at the photoreceptor-RPE interface. *Commun Biol*. 02 05 2021;4(1):161.

doi:10.1038/s42003-021-01682-5

21.   Bozorg S, Ramirez-Montealegre D, Chung M, Pearce DA. Juvenile neuronal ceroid

lipofuscinosis (JNCL) and the eye. *Surv Ophthalmol*. Jul-Aug 2009;54(4):463-71.

doi:10.1016/j.survophthal.2009.04.007

22.   Bassett AR. Editing the genome of hiPSC with CRISPR/Cas9: disease models. *Mamm*

Genome. Aug 2017;28(7-8):348-364. doi:10.1007/s00335-017-9684-9

23.   Findlay GM. Linking genome variants to disease: scalable approaches to test the

functional impact of human mutations. *Hum Mol Genet*. Oct 01 2021;30(R2):R187-R197.

doi:10.1093/hmg/ddab219

24.   Kim HS, Kweon J, Kim Y. Recent advances in CRISPR-based functional genomics for

the study of disease-associated genetic variants. *Exp Mol Med*. Apr 2024;56(4):861-869.

doi:10.1038/s12276-024-01212-3

25.   Kurzawa-Akanbi M, Tzoumas N, Corral-Serrano JC, et al. Pluripotent stem cell-derived

models of retinal disease: Elucidating pathogenesis, evaluating novel treatments, and estimating

toxicity. *Prog Retin Eye Res*. May 2024;100:101248. doi:10.1016/j.preteyeres.2024.101248

26.   Afanasyeva TAV, Athanasiou D, Perdigao PRL, et al. CRISPR-Cas9 correction of a

nonsense   mutation   in.   *Mol   Ther   Methods   Clin   Dev*.   Jun   08   2023;29:522-531.

doi:10.1016/j.omtm.2023.05.012

27.   Nolan ND, Cui X, Robbings BM, et al. CRISPR editing of anti-anemia drug target

rescues   independent   preclinical   models   of   retinitis   pigmentosa.   *Cell   Rep   Med*.   Apr   16

2024;5(4):101459. doi:10.1016/j.xcrm.2024.101459

28.   Lim RR, Shirali S, Rowlan J, et al. CFH Haploinsufficiency and Complement Alterations

in Early-Onset Macular Degeneration. *Invest Ophthalmol Vis Sci*. Apr 01 2024;65(4):43.

doi:10.1167/iovs.65.4.43

29.   Galloway CA, Dalvi S, Hung SSC, et al. Drusen in patient-derived hiPSC-RPE models of

macular   dystrophies.   *Proc   Natl   Acad   Sci   U   S   A*.   Sep   2017;114(39):E8214-E8223.

doi:10.1073/pnas.1710430114

30.   Xu P, Chen Z, Ma J, et al. Biallelic CLCN2 mutations cause retinal degeneration by

impairing retinal pigment epithelium phagocytosis and chloride channel function. *Hum Genet*.

Apr 2023;142(4):577-593. doi:10.1007/s00439-023-02531-7

31.   Cuevas E, Holder DL, Alshehri AH, Tréguier J, Lakowski J, Sowden JC. NRL. *Stem*

Cells. Apr 2021;39(4):414-428. doi:10.1002/stem.3325

32.   Wang Q, Chear S, Wing K, et al. Use of CRISPR/Cas ribonucleoproteins for high

throughput gene editing of induced pluripotent stem cells. *Methods*. Oct 2021;194:18-29.

doi:10.1016/j.ymeth.2021.02.009

33.   Chear S, Perry S, Wilson R, et al. Lysosomal alterations and decreased

electrophysiological activity in CLN3 disease patient-derived cortical neurons. *Dis Model Mech*.

Dec 01 2022;15(12)doi:10.1242/dmm.049651

34.   Wang K, Li M, Hadley D, et al. PennCNV: an integrated hidden Markov model designed

for high-resolution copy number variation detection in whole-genome SNP genotyping data.

Genome Res. Nov 2007;17(11):1665-74. doi:10.1101/gr.6861907

35.   Colella S, Yau C, Taylor JM, et al. QuantiSNP: an Objective Bayes Hidden-Markov

Model to detect and accurately map copy number variation using SNP genotyping data. *Nucleic*

Acids Res. 2007;35(6):2013-25. doi:10.1093/nar/gkm076

36.   Neavin D, Nguyen Q, Daniszewski MS, et al. Single cell eQTL analysis identifies cell

type-specific genetic control of gene expression in fibroblasts and reprogrammed induced

pluripotent stem cells. *Genome Biol*. Mar 05 2021;22(1):76. doi:10.1186/s13059-021-02293-3

37.   Manian KV, Galloway CA, Dalvi S, et al. 3D iPSC modeling of the retinal pigment

epithelium-choriocapillaris complex identifies factors involved in the pathology of macular

degeneration. *Cell Stem Cell*. 05 06 2021;28(5):846-862.e8. doi:10.1016/j.stem.2021.02.006

38.   Singh R, Shen W, Kuai D, et al. iPS cell modeling of Best disease: insights into the

pathophysiology of an inherited macular degeneration. *Hum Mol Genet*. Feb 2013;22(3):593-

607. doi:10.1093/hmg/dds469

39.   Parinot C, Rieu Q, Chatagnon J, Finnemann SC, Nandrot EF. Large-scale purification of

porcine or bovine photoreceptor outer segments for phagocytosis assays on retinal pigment

epithelial cells. *J Vis Exp*. Dec 12 2014;(94)doi:10.3791/52100

40.   Toops KA, Tan LX, Lakkaraju A. A detailed three-step protocol for live imaging of

intracellular traffic in polarized primary porcine RPE monolayers. *Exp Eye Res*. Jul 2014;124:74-

85. doi:10.1016/j.exer.2014.05.003

41.   Müller C, Charniga C, Temple S, Finnemann SC. Quantified F-Actin Morphology Is

Predictive of Phagocytic Capacity of Stem Cell-Derived Retinal Pigment Epithelium. *Stem Cell*

Reports. 03 13 2018;10(3):1075-1087. doi:10.1016/j.stemcr.2018.01.017

42.   Muller C, Charniga C, Temple S, Finnemann SC. Quantified F-Actin Morphology Is

Predictive of Phagocytic Capacity of Stem Cell-Derived Retinal Pigment Epithelium. *Stem Cell*

Reports. Mar 13 2018;10(3):1075-1087. doi:10.1016/j.stemcr.2018.01.017

43.   Dalvi S, Galloway CA, Winschel L, et al. Environmental stress impairs photoreceptor

outer segment (POS) phagocytosis and degradation and induces autofluorescent material

accumulation in hiPSC-RPE cells. *Cell Death Discov*. 2019;5:96. doi:10.1038/s41420-019-0171-

9

44.   Kennedy BP, Payette P, Mudgett J, et al. A natural disruption of the secretory group II

phospholipase A2 gene in inbred mouse strains. *J Biol Chem*. Sep 22 1995;270(38):22378-85.

doi:10.1074/jbc.270.38.22378

45.   Amit M, Carpenter MK, Inokuma MS, et al. Clonally derived human embryonic stem cell

lines maintain pluripotency and proliferative potential for prolonged periods of culture. *Dev Biol*.

Nov 15 2000;227(2):271-8. doi:10.1006/dbio.2000.9912

46.   Galloway CA, Dalvi S, Shadforth AMA, et al. Characterization of Human iPSC-RPE on

a Prosthetic Bruch's Membrane Manufactured From Silk Fibroin. *Invest Ophthalmol Vis Sci*. Jun

2018;59(7):2792-2800. doi:10.1167/iovs.17-23157

47.   Mircheff AK, Miller SS, Farber DB, Bradley ME, O'Day WT, Bok D. Isolation and

provisional identification of plasma membrane populations from cultured human retinal pigment

epithelium. *Invest Ophthalmol Vis Sci*. May 1990;31(5):863-78.

48.   Sonoda S, Spee C, Barron E, Ryan SJ, Kannan R, Hinton DR. A protocol for the culture

and differentiation of highly polarized human retinal pigment epithelial cells. *Nat Protoc*.

2009;4(5):662-73. doi:10.1038/nprot.2009.33

49.   Ouseph MM, Kleinman ME, Wang QJ. Vision loss in juvenile neuronal ceroid

lipofuscinosis (CLN3 disease). *Ann N Y Acad Sci*. May 2016;1371(1):55-67.

doi:10.1111/nyas.12990

50.   Radke J, Stenzel W, Goebel HH. Human NCL Neuropathology. *Biochim Biophys Acta*.

May 2015;doi:10.1016/j.bbadis.2015.05.007

51.   Vidal-Donet JM, Cárcel-Trullols J, Casanova B, Aguado C, Knecht E. Alterations in

ROS activity and lysosomal pH account for distinct patterns of macroautophagy in LINCL and

JNCL fibroblasts. *PLoS One*. 2013;8(2):e55526. doi:10.1371/journal.pone.0055526

52.   Benedict JW, Sommers CA, Pearce DA. Progressive oxidative damage in the central

nervous system of a murine model for juvenile Batten disease. *J Neurosci Res*. Oct

2007;85(13):2882-91. doi:10.1002/jnr.21416

53.   Tuxworth RI, Chen H, Vivancos V, Carvajal N, Huang X, Tear G. The Batten disease

gene CLN3 is required for the response to oxidative stress. *Hum Mol Genet*. May 15

2011;20(10):2037-47. doi:10.1093/hmg/ddr088

54.   Dannhausen K, Möhle C, Langmann T. Immunomodulation with minocycline rescues

retinal degeneration in juvenile neuronal ceroid lipofuscinosis mice highly susceptible to light

damage. *Dis Model Mech*. Sep 05 2018;11(9)doi:10.1242/dmm.033597

55.   Groh J, Berve K, Martini R. Fingolimod and Teriflunomide Attenuate Neurodegeneration

in Mouse Models of Neuronal Ceroid Lipofuscinosis. *Mol Ther*. Aug 02 2017;25(8):1889-1899.

doi:10.1016/j.ymthe.2017.04.021

56.   Brooks AI, Chattopadhyay S, Mitchison HM, Nussbaum RL, Pearce DA. Functional

categorization of gene expression changes in the cerebellum of a Cln3-knockout mouse model

for Batten disease. *Mol Genet Metab*. Jan 2003;78(1):17-30. doi:10.1016/s1096-7192(02)00201-

9

57.   Gupta P, Tirgan N, Kalariya NM, Godley BF, Motamedi M. Kinetics of Oxidized Rod

Outer Segment Phagocytosis Under Chronic Oxidative Stress in an in-vitro Model of Age-

Related   Macular   Degeneration.   *Investigative   Ophthalmology   &   Visual   Science*.

2010;51(13):494-494.

58.   Becquet F, Goureau O, Soubrane G, Coscas G, Courtois Y, Hicks D. Superoxide inhibits

proliferation and phagocytic internalization of photoreceptor outer segments by bovine retinal

pigment epithelium in vitro. *Exp Cell Res*. Jun 1994;212(2):374-82. doi:10.1006/excr.1994.1157

59.   Singh RK, Nasonkin IO. Limitations and Promise of Retinal Tissue From Human

Pluripotent Stem Cells for Developing Therapies of Blindness. *Front Cell Neurosci*.

2020;14:179. doi:10.3389/fncel.2020.00179

60.   Capowski EE, Samimi K, Mayerl SJ, et al. Reproducibility and staging of 3D human

retinal organoids across multiple pluripotent stem cell lines. *Development*. Dec

2018;doi:10.1242/dev.171686

61.   Radke J, Stenzel W, Goebel HH. Human NCL Neuropathology. *Biochim Biophys Acta*.

Oct 2015;1852(10 Pt B):2262-6. doi:10.1016/j.bbadis.2015.05.007

62.   Rowland TJ, Blaschke AJ, Buchholz DE, Hikita ST, Johnson LV, Clegg DO.

Differentiation of human pluripotent stem cells to retinal pigmented epithelium in defined

conditions using purified extracellular matrix proteins. *J Tissue Eng Regen Med*. Aug

2013;7(8):642-53. doi:10.1002/term.1458

63.   Dalvi S, Galloway CA, Singh R. Pluripotent Stem Cells to Model Degenerative Retinal

Diseases: The RPE Perspective. *Adv Exp Med Biol*. 2019;1186:1-31. doi:10.1007/978-3-030-

28471-8_1

    Figure Legends

Figure 1. Generation and characterization of isogenic control and CLN3 hESC RPE.

a) Schematic depicting CRISPR-Cas9 deletion of exons 7 and 8 of CLN3 in H9 hESC**.** Note the

CLN3 locus and the location of dual guides to excise exons 7-8 from the CLN3 gene.

b) Identification of exons 7-8 deletion in CLN3 gene in positive clones. Genomic PCR of clones

with homozygous deletion (lane 1 and 2) and heterozygous deletion (lane 3). RT-PCR analysis

of homozygous 1kb deletion clone (lane 4) and control (lane 5).

c) Sequencing alignment of cDNA for control (top panel) and *CLN3$^{Δ7\text{-}8/Δ7\text{-}8}$* (bottom panel) cell

lines shows successful biallelic deletion of exons 7-8 reflected as splicing of exon 6 to exon 9

(indicated by an arrow).

d) Representative light microscopy images (top panel) and electron microscopy images (bottom

panel) showing the expected RPE morphology and apical microvilli in isogenic control and

CLN3$^{Δ7\text{-}8/Δ7\text{-}8}$ hESC RPE (*CLN3* hRPE) (scale bar = 100µm).

e) Representative confocal microscopy images showing similar and expected localization of tight

junction protein, ZO1 (red, top panel), RPE microvilli protein, EZR (red, middle panel), and

basolaterally expressed RPE protein, BEST1 (green, bottom panel) in control and *CLN3* hRPE

cells. Nuclei were stained with DAPI (blue) (scale bar = 10µm).

f) Representative Western blot images showing expected presence of RPE signature proteins in

both control and CLN3 hRPE cells. ACTN served as the loading control in these experiments.

g) Representative light microscopy images showing evidence of transepithelial fluid flux with

presence of fluid domes in polarized monolayers of control and *CLN3* hRPE cells. Note the

presence of RPE both within the fluid dome (focal plane 1, top panel) and outside the fluid dome

(focal plane 2, bottom panel) (scale bar = 50 μm).

h) Transepithelial resistance (TER) measurements showing evidence of a tight epithelial

monolayer in both control and *CLN3* hRPE cultures. Note the dotted line represent the known

TER threshold for RPE cells in *vivo*[48].

n ≥ 3 for all experiments in figure 1

Figure 2. Evaluation of POS phagocytosis and RPE autofluorescence by control versus

***CLN3* hRPE cells.**

a) Schematic of experimental assay utilized to measure POS uptake by control and *CLN3* hRPE

cells after feeding a physiologic dose of POS (~20-40 POS/RPE cell) for 2h.

b, c) Representative confocal microscopy images (b, scale bar 10 µm) and quantitative analyses

(c) showing decreased phagocytosis of FITC-POS particles (green, panel b) but similar

localization of tight junction protein (ZO1) in *CLN3* hRPE cells compared to control hRPE cells.

Nuclei were stained with DAPI (blue) (scale bar = 10μm). ***p ≤ 0.001.

d) Schematic of experimental assay utilized to measure RPE autofluorescence after daily chronic

POS feeding (~20-40 POS/RPE cell) for a duration of 14 days by control and *CLN3* hRPE cells

after feeding a physiologic dose of POS (~20-40 POS/RPE cell) for 2h.

e, f) Representative confocal microscopy images (e, scale bar 10μm) and quantitative analyses (f,

f') showing decreased RPE autofluorescence (count, area) in the red channel (ex: 546 nm and

em: 560-615 nm) but similar localization of tight junction protein (ZO1) in *CLN3* hRPE cells fed

daily with POS compared to control hRPE cells fed daily with POS. Nuclei were stained with

DAPI (blue) (scale bar = 10μm). ***p ≤ 0.001.

n ≥ 3 for all experiments in figure 2

Figure 3. Evaluation of POS binding versus internalization by control versus *CLN3* hRPE

cells.

a-c) Representative confocal images of FITC-POS fed (2h) RPE monolayers showing decreased

number of total FITC-POS particles (a), apical FITC-POS compared to ZO1 (consistent with

reduced FITC-POS binding (b), and basal FITC-POS compared to ZO1 (consistent with reduced

FITC-POS internalization) (c) in *CLN3* hRPE cells compared to control hRPE cells. Note that a

published protocol that utilized position of ZO-1 relative to FITC-POS to estimate bound versus

internalized POS was used in these experiments (scale bar = 10 μm).

d-f) Quantitative analyses showing decreased number of both bound FITC-POS (d) and

internalized FITC-POS (e) in *CLN3* hRPE cells compared to control hRPE cells. Consistent with

a POS binding defect, the number of internalized FITC-POS relative to number of bound FITC-

POS was unchanged between control and *CLN3* hRPE cells (**f**). *p ≤ 0.05, ***p ≤ 0.001.

n ≥ 3 for all experiments in figure 3

Figure 4. Evaluation of POS uptake of wild-type (WT) versus *CLN3* POS by RPE cells.

a, b) Schematic of experimental assay utilized to measure POS phagocytosis by WT primary

miniswine RPE post feeding of WT miniswine POS versus *CLN3* pig POS for 2h.

c, d) Representative Western blot image (c) and quantification (d) showing decreased

phagocytosis of *CLN3* miniswine POS by WT miniswine RPE compared to WT miniswine POS

by WT miniswine RPE. ***p ≤ 0.001.

n ≥3 for figure 4 experiment.

Figure 5. Longitudinal comparison of RPE autofluorescence levels and retina histology in

CLN3 **miniswine eye.**

a-c) Representative confocal images (a, b) and quantitative analyses (c, c') showing similar

levels of autofluorescence accumulation in RPE of WT and *CLN3* miniswine at 6-month of age.

In contrast, decreased autofluorescence accumulation was observed in *CLN3* miniswine RPE

compared to WT miniswine RPE at the 36-months of age (scale bar 10 µm). $**p \leq 0.01$.

d) Representative images of *CLN3* miniswine retina sections at 36 and 48-months age showing

presence of POSs at 36 months of age but absence of POSs with significant loss of the

photoreceptor cell nuclei/ outer nuclear layer (ONL) at 48-months of age.

n≥3 for all experiments in figure 5. Scale bar = 5 µm.

a

5'   3'

Exon:
1   7 8   15

966 bp *CLN3* gene deletion spanning exons 7 & 8

Control hESC line

| Exon 6 | sg1 | Exon 7 | Exon 8 | sg2 | Exon 9 |

Exon 6 | Exon 9

CLN3$^{Δ7-8/Δ7-8}$ hESC line

b

1   2   3   4   5

746 bp
613 bp

c

Exon 6 — Exon 7

Control

Exon 6 — Exon 9

CLN3$^{Δex7/8/Δex7/8}$

d

Control   *CLN3*

e

Control   *CLN3*

ZO-1/DAPI

EZR/DAPI

BEST1/DAPI

f

	Control	*CLN3*
BEST1		
CRALBP		
EZR		
MERTK		
RPE65		
ACTN		

g

Control   *CLN3*

Focal Plane 1

Focal Plane 2

h

TER (Ω*cm^{-2})

Control   CLN3

a

Wash and Measure FITC

Feed FITC-POS (~20-40 POS/RPE cell)

0h   2h

b   Control   *CLN3*

DAPI/FITC/ZO1

c

# of Total FITC-POS Particles/ 100 Cells (Normalized to Control)

***

Control   CLN3

d

Measure autofluorescence

Feed POS (~20-40 POS/RPE cell)

Day: 0   14

e   Control   *CLN3*

DAPI/ZO1/autofluorescence

f

# of Autofluorescence Per DAPI-stained cell (Normalized to Control)

***

Control   CLN3

f'

Autofluorescence Per viewing area (Normalized to Control)

***

Control   CLN3

a

Wash and Measure RHO

Feed unlabeled POS (10 POS/RPE cell)

0h   2h

b

WT RPE   WT RPE

+ WT POS   + *CLN3* POS

c

WT   *CLN3*

RHO

ACTN

d

RHO/ACTN (Normalized to WT)

***