# **Title:** Sex-biased gene expression across tissues reveals unexpected differentiation in the gills of the threespine stickleback

**Authors**: Florent Sylvestre[1,*], Nadia Aubin-Horth[1], Louis Bernatchez[1,†]

[1]: Département de Biologie, Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada

†: Deceased

*: Corresponding author

Correspondence: Florent.sylvestre.1@ulaval.ca

**Orcid**:

Florent Sylvestre: 0000-0002-8769-9242

Nadia Aubin-Horth: 0000-0002-9030-634X

Louis Bernatchez: 0000-0002-8085-9709

**Abstract:**

Sexual dimorphism can evolve through sex-specific regulation of the same gene set. However, sex chromosomes can also facilitate this by directly linking gene expression to sex. Moreover, heteromorphic sex chromosomes often exhibit different gene content, which contributes to sexual dimorphism. Understanding patterns of sex-biased gene expression across organisms is important for gaining insight about the evolution of sexual dimorphism and sex chromosomes. Moreover, studying gene expression in species with recently established sex chromosomes can help understand the evolutionary dynamics of gene loss and dosage compensation. The threespine stickleback is known for its strong sexual dimorphism, especially during the reproductive period. Sex is determined by a young XY sex chromosome pair with three non-recombining regions that have started to degenerate. Using the high multiplexing capability of 3' QuantSeq to sequence the sex-biased transcriptome of liver, gills and brain, we provide the first characterization of sex-specific transcriptomes from ~80 stickleback (40 males and 40 females) collected from a natural population during the reproductive period. We find that the liver is extremely differentiated (36% of autosomal genes) and reflects ongoing reproduction, while the brain shows very low levels of differentiation (0.78%) with no particular functional enrichment. Finally, the gills exhibit high levels of differentiation (5%), suggesting that sex should be considered in physiological and ecotoxicological studies of gill responses in fishes. We also find that sex-biased gene expression in X-linked genes is mainly driven by a lack of dosage compensation. However, sex-biased expression of genes that have conserved copies on both sex chromosomes is likely driven by the degeneration of Y allele expression and a down-regulation of male-beneficial mutations on the X chromosome.

*Keywords*: sex, Liver, Gills, Brain, Stickleback, gene expression, dosage compensation

## Introduction

Species with sexual reproduction often exhibit sexual dimorphism (Lande, 1980). Because sexes share most of their genetic material except for potential sex chromosomes, sexual dimorphism can arise from the different regulation of the same set of genes (Ellegren & Parsch, 2007; Tosto et al., 2023). Sex chromosomes thus represent a key step in understanding the evolution of sexes. Sex-biased gene expression has been described in many systems as dependent on both life stages (Djordjevic et al., 2022) and tissues (Rodríguez-Montes et al., 2023). Compiled data for five tissues in five mammals and a bird species show that sex-biased gene expression varies in intensity across tissues, species and sexual maturity, with the contribution of sex chromosomes being also variable (Rodríguez-Montes et al., 2023).

The variety of sex determinism systems and reproductive behaviours present in teleost fish make them an interesting vertebrate group to study sex-biased gene expression (Thresher, 1984; Devlin & Nagahama, 2002; Kobayashi et al., 2013). Patterns of sex-biased gene expression have been described in fish somatic tissues, such as the brain or liver, which are known to play a role in reproduction. Sex-biased gene differentiation in the brain seems highly species dependent, with about a thousand genes identified in salmonids (Hale et al., 2018) but only a handful in the Gulp pipefish *syngnathus scovelli* (Beal et al., 2018) or the zebrafish *Danio renio* (Yuan et al., 2019). In cichlids, gene expression levels in the brain are associated with social status and gonadic sex (Renn et al., 2008; Schumer et al., 2011). Liver is sexually dimorphic, specifically in oviparous species (Qiao et al., 2016; Darolti & Mank, 2023) and many genes have been identified as sexually biased in salmonids (Sutherland et al., 2019) and across cichlid taxa (Lichilín et al., 2021). On the contrary, other tissues such as the gills have received strong attention in the context of adaptation to salinity, pollution or hypoxia (Scott et al., 2004; Van Der Meer et al., 2005; Gonzalez et al., 2006) but studies rarely account for potential sex dimorphism in the response of this tissue (but see (Lichilín et al., 2021). Given the dynamic nature of sex-biased gene expression, we need a better characterization of sex-biased gene expression in natural populations if we want to understand the processes underlying the evolution of sexual dimorphism.

Sex chromosomes play an important role in sex dimorphism (Rice, 1984). However, understanding patterns of sex-biased gene expression on sex chromosomes is particularly complex in species with non-recombining sex chromosomes, as they tend to degenerate and may involve dosage compensation (Bachtrog, 2013). The lowered effective size of the Y (or W) chromosome, present in only one sex, leads to lowered efficiency of natural selection and the degeneration of Y chromosomes (Charlesworth & Charlesworth, 2000). Genes that have lost their Y or W copy (hemizygous genes) are expected to exhibit a reduced expression level that does not come from sex-specific gene regulation. However, lowered gene activity on sex-chromosome can have widespread effects on autosomal genes (Wijchers et al., 2010), and mutation reestablishing the ancestral level of expression in the heterogametic sex is expected to be advantageous, leading to the evolution of dosage compensation. Global dosage compensation, where the X (or Z) chromosome is overexpressed to compensate for the loss of Y (or W) genes was long thought to be necessary in systems with degenerated sex chromosomes, but accumulating literature outside

*Drosophila* and mammals suggest that it is not necessarily the case (Mank, 2013), with many groups
exhibiting no or partial dosage compensation. Moreover, many genes are dose insensitive (i.e.,
their expression level does not affect the phenotypic outcome), therefore they do not need to be
compensated. In fish, global dosage compensation has rarely been found (Darolti et al., 2019) and
we still lack knowledge about the extent of the evolution of dosage compensation in this highly
diverse group.
The three-spined stickleback is a model fish species with a young XY sex-chromosome system,
at most 13–26 million years-old (Peichel et al., 2020), and strong sexual dimorphism. The sex
chromosome consists of a pseudoautosomal region which is still recombining, and three
evolutionary strata which have evolved through successive inversion (Peichel et al., 2020). Studies
have found sex-specific splicing and protein expression (Viitaniemi & Leder, 2011; Naftaly et al.,
2021), but few have described sex-specific patterns of gene expression in that species. The brain is
the most studied tissue (Primmer et al., 2013; Kaitetzidou et al., 2022) and shows limited sex-biased
gene expression outside the sex chromosomes (but see (Kitano et al., 2020). Similar results have
been observed in the liver (Leder et al., 2010). Accumulation of sex-biased genes on sexual
chromosomes seems to be associated with a lack of global dosage compensation (Leder et al.,
2010; Schultheiß et al., 2015; White et al., 2015) in that species, coupled with a potential partial
dosage compensation in one of the evolutionary strata of the Y chromosome. However, these
studies used laboratory-raised individuals outside the reproductive period when sticklebacks
exhibit the strongest sexual dimorphism, and thus more studies are needed to understand sex-
biased gene expression in natural populations during the reproductive period.
In this study, we took advantage of the high sample multiplexing capability offered by QuantSeq
3′-UTR sequencing (Moll et al., 2014) to profile the transcriptome of ~40 samples per sex in three
tissues (liver, brain, and gills) in adults from a natural population of threespine stickleback from
eastern Canada. We aimed at 1) describing the sex-specific transcriptome of this species in each
tissue and 2) making use of the recently sequenced Y chromosome data to understand the
dynamics of dosage compensation. According to work in other species, we expected the liver to be
the most differentiated somatic tissue between sexes, as it plays an important role during
reproduction in teleost fish (Meng et al., 2016). The brain is known to be sexually dimorphic during
the reproduction of the threespine stickleback (Kotrschal et al., 2012) and we expected substantial
sex-biased gene expression for that tissue. Finally, sex is not a factor usually accounted for when
studying gills, an extremely important tissue for physiological regulation, and when it is, there is
almost no sex-biased expression (Lichilín et al., 2021). We therefore predicted that this would be a
neutral tissue with little to no sex-biased expression. We also expected to find that most genes on
sex-chromosome exhibit sex-biased gene expression caused by the lack of global dosage
compensation in that species.

# Material and methods

**Ethics statement**

This study was approved by the Comité de Protection des Animaux de l'Université Laval (CPAUL, approval number SIRUL 109096). A fish permit was issue by the Ministère des Forêts, de la Faune et des Parcs du Québec (permit number 2018 04 11 005 01 S P) for sampling.

**Sampling and Sequencing:**

We collected adult anadromous three-spined sticklebacks from tide pools of the St Lawrence River at Baie de l'Isle verte (48.009961, −69.407070). Brain, liver, and gills were dissected (under five, seven and ten minutes respectively) and preserved in RNAlater at -20 C. In 2022, we performed RNA extractions using the RNeasy mini Kit (Qiagen). We disrupted samples is 700 µL (brain) or 1400 µL (Gills, Liver) of trizol using a mixermix 400 at 30 Hz for 3 minutes or until complete tissue disruption. After three minutes, we added 140 µL (or 280 µL) of chloroform, homogenized the solution and waited five minutes before centrifuging for 15 min at 12,000g. We collected the upper phase and added 550 µL (or 1100 µL) of ethanol before transferring 700 µL into a RNeasy Mini column (Qiagen). We then centrifuged for 15s at 11,000g, discarded the flow-through and repeated the operation until all the collected phase has been used. We then proceeded with the extraction protocol following manufacturer's instruction, including a DNase step, but replacing buffer RW1 by buffer RWT from miRNeasy Mini Kit (Qiagen), as it yielded better results for the brain. We then generated QuantSeq libraries using 3′ mRNA-Seq Library Prep Kit (Lexogen) with dual indexing to identify each individual and 18 PCR cycles for library amplification. After quality check on a 2100 Bioanalyzer (Agilent Technologies) and concentration estimation using Quant-iT PicoGreen (Invitrogen), 227 libraries were pooled to equimolarity and sent to the Centre d'Expertise et de Services Génome Québec (Montréal, QC, Canada) for 50 bp single-end sequencing on 2 lanes of an Illumina Novaseq X. Tissue disruption and quality check were performed at the Plateforme d'Analyse Genomique of Laval university (Québec, QC, Canada).

**Alignment and Expression Counts**

After quality check using FASTQC v0.11.8, we used fastp v 0.15.0 (Chen et al., 2018) to trim poly-A tails, Illumina adapters, and the first 12bp of each read, as they showed biased composition. We discarded reads shorter than 20bp long and proceeded with the alignment. We used STAR v2.7.2b (Dobin et al., 2013) two-pass mode to align reads to the stickleback reference genome V5 (Peichel et al., 2020; Nath et al., 2021), accessible on NCBI as RefSeq assembly GCF_016920845.1), excluding the Y reference sequence for the females and its pseudo-autosomal region (PAR) for males. We used STAR two-pass mode to discover reads junctions and improve mapping accuracy in the second pass, then quantified gene expression using HTSeq v0.11.3. (Anders et al., 2015) in union mode after filtering out multi-mapping reads. Counts from each sample were merged using custom R scripts, leading to four datasets: the total datasets comprising all samples, and one dataset for each tissue (Liver, Gills and Brain).

## Filtering, normalization, and quality check

Unless stated otherwise, analyses were performed in R v 4.3.2 (R Core Team, 2021) and python 3.9.12 (Rossum & Drake, 2010). For each dataset, we applied the same filtering procedure. First, samples with fewer than 5,000,000 raw reads counts were excluded, as preliminary principal component analysis (PCA) showed that they clustered together (result not shown). Then, we kept only genes with one count per million (cpm) in at least 10 samples. To identify potential errors in the dataset (in particular, samples where tissue was misidentified), we first performed a PCA on autosomal genes across all samples, using blind Vst normalization as implemented in DESeq2 v1.40.2 (Love et al., 2014). For all other analyses, read counts normalization was carried out independently for each tissue using the average of ratio normalization factor implemented in DESeq2. Given the specificity of genes on sex chromosomes, only autosomal genes were used for the calculation of the scaling factors in each tissue, which were then used to normalize sex-chromosome genes independently. Note that gene length was not accounted for in the normalization process, as it could generate a bias in our dataset as QuantSeq only sequences the poly-A tail of each transcript, not the full transcript.

## Differential Expression Analysis of Autosomal Genes

Within each tissue, we used Wilcoxon rank-sum tests to identify differentially expressed genes (DEG), using the Benjamini-Hochberg procedure to control the false discovery rate (Benjamini & Hochberg, 1995). This approach was selected as a recent study suggested that the classically used negative binomial models implemented in Deseq2 are subject to increased false positive rates in large sample size datasets (Li, Ge, et al., 2022) We also calculated the log-fold change (LFC) of gene expression between sexes as, for each gene:

$$LFC_g = \frac{\sum_{m=1}^{N_m} log2(g_m + 0.5)}{N_m} - \frac{\sum_{f=1}^{N_f} log2(g_f + 0.5)}{N_f}$$

with gm and gf being the normalized read count for each male and female, and Nm and Nf the number of individuals from each sex.

## Functional Characterization of DEG

To explore common functions among sex-biased genes, we performed Gene Ontology (GO) enrichment analysis. To do so, we used blastX (Camacho et al., 2009) to gather the Swiss-Prot (Schneider et al., 2009) annotation for each sequence of the three-spined stickleback transcriptome available on NCBI (RefSeq assembly GCF_016920845.1) and gathered gene ontology information from the associated UniProt entry (The UniProt Consortium, 2023). We then summarized transcript-level GO at the gene level using information from the NCBI annotation of our reference genome with a custom python script (lien?). We used goatools (Klopfenstein et al. 2018) to perform Fisher's exact test for enrichment at a q-value threshold of 0.05 using the Benjamini-Hochberg procedure. We also combined Zfin (zfin.atlassian.net), genecards (www.genecards.org) and Uniprot (www.uniprot.org) databases to provide a more precise

functional characterization of 1) the 10 genes with the lowest q-value and 2) the 10 genes with the
highest LFC for each tissue, as well as all DEG shared by all tissues. Functional annotations
described in the results section are based on information from these databases unless otherwise
noted.
To explore the genomic distribution of DEGs, we performed a Fisher test for enrichment analysis
for each chromosome to test whether 1) some chromosomes are enriched in DEG considering their
number of genes and 2) DEG within a chromosome are enriched toward male-biased or female-
biased genes considering the global distribution of sex bias toward each sex. We used a Benjamini-
Hochberg procedure to control the false discovery rate independently for the two hypotheses
tested.

**Identification of Shared Genes Between X and Y**

The study of sex-biased gene expression on sex chromosomes is complexified by independent
gene gain or loss on their non-recombining region. To identify gene loss or gain, we first extracted
transcript sequences from the reference genome using gffread (Pertea and Pertea 2020) and the
NCBI genome annotation for our genome version, using -C option to remove transcripts with no
CDS. We did the same using the available ninespine stickleback reference genome (Varadharajan
et al., 2019) and annotation (GenBank accession GCA_949316345.1), then used Orthofinder v2.5.5
(Emms & Kelly, 2019) to identify orthogroups with default parameters. We used a custon python
script to identify transcripts likely to be orthologs between X and Y chromosomes and summarize
the information at the gene level, as each gene can have several transcripts. We accepted an
orthologous relationship if 1) an orthogroup was composed of one transcript from X, one from the
Y and other members originated from the nine spined stickleback. To account for the fact that one
gene can have several transcripts, we accepted an orthogroup with many X or Y transcripts if they
belonged to the same X or Y gene, but filtered out genes for which transcripts did not all belong to
the same orthogroup (except if the transcript was not associated with any other orthogroup).
Transcripts that fell in an orthogroup with only Y or X transcripts were categorized as hemiploid Y
(no gene copy on X) or hemiploid X (no gene copy on Y). Orthogroups that contained autosomal
genes, likely reflecting gene gain on the sex chromosome, were removed from analysis as well as
genes with multiple copies on either sex chromosome. Finally, genes for which transcripts couldn't
be assigned to any orthogroup were considered as hemiploid X or Y. Gene counts for genes still
shared between X and Y chromosomes were pooled in males.  We assigned each gene to one of the
three known evolutionary strata of sex chromosomes using its central position, and breakpoints
defined in Peichel et al. (2020). Given that the pseudoautosomal region of the Y chromosome was
excluded from the read mapping step, read counts for that region were already correct and genes
were considered as still sharinga copy ignoring Orthofinder results.

**Sex-Biased Gene Expression on the Sex Chromosomes**

We used the merged counts to test for sex-biased gene expression on sex chromosome using
the same method as for the autosome. To test for potential dosage compensation, we estimated
the autosome to sex chromosome median expression ratio for all genes, and independently for

hemiploid X and Y genes for males and females. The autosomal median was calculated as the
median across all genes of the mean log2 normalized read counts across all samples. Median
expression for genes on the sex chromosome was calculated similarly but separating individuals
by sex. We estimated one median for all genes, genes that still have a copy on both chromosomes
and genes specific to each sex chromosome. This calculation was done for the whole chromosome
and for each stratum independently (PAR, strata I, strata II and strata III). Confidence intervals (CI)
were calculated using bias correction bootstraping for both autosomal and sex chromosome genes
and significance assessed using overlap of the CI with 0. Finally, to understand the processes
underlying the evolution of sex-biased gene expression for genes with a copy conserved on each
chromosome, we estimated X and Y allele expression in males. We then compared the expression
level of female, male and X and Y alleles in sex-biased genes to their expression levels in unbiased
genes using Wilcoxon rank-sum test.

# Results and discussion

### Sequencing, Mapping and Filtering

The two lines combined rendered 4,247,758,804 reads with an average of 20,319,408 [CI $_{95\%}$:
18,047,379 – 22,591,436] for the gills, 18,847,630 [17,937,603 – 19,757,656] for the brain and
17,073,329 [14,606,317 – 19,540,341] for the liver. The average read length after quality trimming
was 38 bp. After removing samples with fewer than 5,000,000 reads (ten for the liver, one for the
brain), we kept 67 samples in the liver (35 Females; 32 Males), 77 for the brain (38; 39) and 72 in gills
(36; 36). Dataset include 59 samples sequenced in all tissue, but 10 were specific to gills and brain,
five to brain and liver and three to gills and liver. Three samples were sequenced only in brain. We
had a percent mapping of uniquely mapped reads to the reference genome ranging from 44.3% to
88.61% (median 76.76%). After filtering, we identified five overexpressed genes in the liver but none
in the gills or the brain. Liver was the tissue with the lowest number of expressed autosomal genes
(14,402) compared to gills (17624) and brain (17930), with 13957 autosomal genes expressed in all
three tissues. Using PCA to screen our dataset for potential cross tissue contamination or
mislabelling revealed no evident issue (Fig1 A) as each tissue formed a distinct group, confirming
the quality of the dataset.

### Tissues Differ in the Magnitude of the Sex-Biased Expression

Patterns of sex-biased gene expression varied greatly between liver, gills and brain, both in
terms of the number of genes differentially expressed and their function (figure 1B). Liver
transcriptomes showed strong sex specificity, with 5303 sex-biased genes (hereafter SBG, "sex-
biased gene") with a q-value ≤ 5% after a Benjamini-Hocheberg correction. In comparison, using
the same threshold, we found 973 SBG in gills and only 141 in the brain (Figure 1B). Genes also
exhibited a wider range of differential expression in the liver compared to gills and brain (Figure
1 C-D-E). Across all tissues, no chromosome showed enrichment for SBG (Fig. 2, Fisher's test q-
value> 0.4).

**Sex-biased genes found in all tissues are implicated in cell physiology, cell-cell signalling and gene expression modulation**

Most SBG are unique to a tissue, with only 42 genes significant in all three tissues (figure 1B, table S1). Eight of them are related to basic cell physiology such as growth, cytoskeleton, and differentiation. Five genes are involved in cell-cell signalling or adhesion and are mainly known to play a role in neuron communication and development. Seven genes are involved in gene expression modulation and affect either DNA methylation, transcription, or alternative splicing. If fish, sex-specific methylation is known for its role in modulating the expression level of reproduction-related genes (Laing et al., 2018; Li, Chen, et al., 2022) and in sex determinism

**Figure 1:** Patterns of autosomal sex-biased gene expression in threespine stickleback. A) PCA of gene expression on all tissues. B) Overlap of sex-biased genes between tissues. Inset barplot represents tissue specific repartition of male (blue) and female-biased genes. Volcano plots of sex-biased gene expression in C) liver, D) gills and E) brain. Coloured dots correspond to genes significant at a 5% false discovery rate based on a Wilcoxon-rank-sum test. Dotted lines represent a log-fold change of 1 (doubled expression).

(Gemmell et al. 2019). Similarly, sex-specific transcript usage can provide an alternative route from gene regulation to generate sexual dimorphism (Telonis-Scott et al., 2009; Naftaly et al., 2021). Hence, those genes could play an important role in regulating sex-biased gene expression and

dimorphism across tissues. Other functions found in shared SBG among tissues involve the
immune system, testosterone response (one gene) and two nuclear genes with mitochondrial
function.
In most cases, genes showed the same directionality in all tissues (Fig. 3), except for two genes:
slc16a13, a monocarboxylic acid transporter, and esr2b, an estrogen receptor. Both are female-
biased in liver but male-biased in the brain and gills. While the function of slc16a13 is hard to
interpret in our context, as the substrate of this member of a large family of solute transporter is
unknown (Halestrap, 2012), the gene esr2b code for an estrogen receptor. In teleosts, estrogen
receptors are involved in several biological processes, including reproductive processes (Nelson &
Habibi, 2013). In the brain, esr2b is thought to play a role in reproduction through the
291

Figure 2: Proportion of sex biased genes across chromosomes and sex in each tissue.
Chromosome VII and XVIII show enrichment toward male-biased function in liver (q-value
≤0.05), as well as the mitochondria with lower confidence (q-value = 0.12. No chromosomes
are significantly enriched in sex-biased genes.

regulation of gonadotropin production, which is involved in gametogenesis (Muriach et al., 2008)
in both sexes and have been found to have a higher expression in males in the pituitary gland of
the fathead minnow (Pimephales promelas) (Filby & Tyler, 2005). In the liver, esr2b showed strong
expression levels and was female biased (LFC of -0.35), and could be associated with vitellogenesis
process, although it seems to vary across species (Dominguez et al., 2014; Chen et al., 2019). We
still lack knowledge on the effect of sex hormones in the gills.
In the case of a pattern of directionality in gene expression shared by only two tissues, we only
found discordance when the comparison included the liver (Fig. S1). While part of this is explained

by the liver having both more SBG and more shared SBG with other tissues in general, it suggests
that this tissue might have a different usage of the same gene set.
306

307 **Figure 3:** Heatmap of log-fold change in gene expression between sexes for 42 genes
308 significantly differentially expressed in all tissues. Bar on top shows concordance in the
309 direction of expression bias.

**Sex-biased genes in the brain are few and not enriched for particular functions**

The brain showed the lowest number of SBG with 141 genes differentially expressed between
sexes on the autosomes (0.78% of expressed genes), equally distributed between males and
females and across chromosomes (Figure 2). We found no enrichment for any gene ontology term
using the 5% threshold (supplementary tables). These results are in accordance with a similar
study in the same species in which they found a higher number of expressed genes but included
sex chromosomes in their analyses (Kaitetzidou et al., 2022). Looking into the most significant
genes (10 with highest p-value and 10 with highest LFC), we found that only 12 of them are specific
to the brain (Table S2). We were unable to annotate three of them (LOC120812970, LOC120817963
and LOC120833148). The remaining 9 genes were associated with various biological functions. The
growth hormone-releasing hormone (ghrh), more highly expressed in males, is the first hormone
secreted in the growth hormone axis, which in fish is not only involved in growth but also
reproduction, metabolism and immune function (Chang & Wong, 2009). TTC29 is involved in cilium
movement and is mainly described in sperm flagellum (Bereketoğlu et al., 2022). Other genes
included ihhb (LOC120819658), which is involved in neural and chondrocye development (Wu et
al., 2001; Chung et al., 2013), hcn3 (si:dkey-197j19.6), an ion channel which is essential for neuronal
function, MMP13 or 18 (LOC120822795), which modulates angiogenesis in the brain (Ma et al.,
2016), and ecm1a (LOC120810788), which codes for an extracellular matrix protein involved in
signal transduction.

**Sex-biased genes in gills are associated with ion-related functions and immune defense**

We identified 973 SBG in the gills (5.5% of expressed genes), of which 60.8% are male-biased.
Significant genes are distributed homogeneously in the genome, with no chromosome significantly
enriched in SBG of biased toward a sex (Fig. 2). Synaptic signaling and organisation represent 48%
of significantly enriched GO terms (Table S3), most of them (48% of total significant terms) related
to synaptic signalling and organization. However, looking at descriptions of genes within those GO
categories on Zfin or Genbank revealed that many genes code for ion channels or pumps, which

are a core function of the gill tissue (Perry et al., 2003), yet the associated GO functions have only
been described in the brain (Table S4). This suggests that gene ontology analysis in gills suffer from
the lack of gill-specific information. Other GOs include various biosynthetic and metabolic
processes, as well as cell adhesion, communication, and development. Most strongly differentiated
genes (Table S5) in gills involve two genes with potential role in pathogen resistance, hhipl1 and
CLEC4M (LOC120817010) and genes with basic cellular functions (LOC120828377, si:dkeyp-92c9.4
and LOC120810538). Asic2 codes for an ion channel, and this gene family is involved in Na+ intake
in rainbow trout gills (Dymowska et al., 2014). Three genes are involved in neuronal function,
including one ion channel, rem2, and a galanin receptor, galr2b.  Dpysl3, is also male biased in the
liver is supposed to have an effect of peripheric axon growth. As for the gene ontology, the function
of these genes in the gills is unclear. Finally, three genes (LOC120816929, LOC120817829 and
LOC120821053) could not be annotated.
Most studies on gills transcriptomes focused on their role in osmoregulation and respiratory
processes in responses to anoxia, salinity, or various contaminants (Scott et al., 2004; Van Der Meer
et al., 2005; Gonzalez et al., 2006). Works have also been interested in gills' function in defence
against pathogens, as they represent a direct entry for infection and parasites (Mitchell & Rodger,
2011). However, a survey of the physiology literature illustrates that these studies do not usually
account for sex. When they do, few SBG are identified, for example, in African cichlids (Lichilín et
al., 2021). However, our results identify numerous SBG in animals all found in the same
environmental conditions. This unexpected result highlights the importance of accounting for sex
when studying gills, as the extent to which this tissue might respond differently to various
challenges between sexes is also poorly understood.

**The Liver is a Hotspot of Sex-Biased Gene Expression**

We identified 5303 SBG in the liver (36.8% of total expressed genes), 60.3% of which are male-
biased. SBG are uniformly distributed among chromosomes (Fisher's exact test qvalue ≥ 0.45,
Figure 2), except for chromosomes V and XVIII, which show enrichment in male-biased genes, and
the mitochondria which shows marginal enrichment for male-biased function (FDR of 0.12). While
comparing the number of genes between studies is complex as life stages, condition and filtering
have a deep impact on detected SBG, widespread sex-biased gene expression has been found in
the liver of other fish species, such as *Salvelinus fontinalis*, in which SBG represents 16.1% of the
total gene expression using more stringent filtering (LFC>= 1.5) than we applied (Sutherland et al.,
2019). Similarly, very low levels of SBG are identified in cichlids (Lichilín et al., 2021) using a LFC of
2 as a cutoff but these observations vary across species. In our study, only 396 (2.6%) in the liver
passed this cutoff (644 for the 1.5 cutoff), suggesting that sex-specific regulation of gene expression
in the liver mostly occurs through subtle regulatory changes, with a median absolute LFC for
significant genes at 0.63. Enriched gene ontology terms in the liver are mainly related to metabolic
and biosynthetic processes. The immune system also seems differentiated between the sexes, with
enriched processes such as humoral immune response and response to external stimulus. We also
identified an enrichment in hemostasis and coagulation regulation (Table S6). Among the most
significant genes in terms of p-value and fold change (Table S7), numerous genes were involved in

response to estrogen and estradiol (fam20cl, vtg3, LOC100190882, LOC100190880, LOC120823934
blasting to vtg1, and two vtg2). Other genes showed functions related to gene expression
regulation (e.g. lbx2, st18), response to pathogens (LOC120810467, LOC120820940, blasting to the
fucolectin-1 and CHIT1), and cellular differentiation (LOC120824638, blasting to srda3a).
We identify five genes representing more than 15% of reads in at least one sample: apoa2,
lect2.1, LOC120808851, LOC120810467 and LOC120823934, hereafter referred to as overexpressed
genes (Figure 4). All five overexpressed genes showed sex-biased gene expression, according to
Wilcoxon rank-sum tests (p.value <10-6). Apoa2 (apolipoprotein A-II) and lect2.1 (leukocyte cell
derived chemotaxin 2.1), which were more highly expressed in males, have functions related to
lipid transport and immune system. Apolipoproteins are involved in lipid transports in vertebrates
but have also been found to have antimicrobial activity in teleost fish (Concha et al., 2003), among
other functions. Leukocyte cell derived chemotaxin2 have been known to have chemotactic
activity in humans, but also have antibacterial activity in other vertebrates and in teleost fish.
BlastX results for females-biased genes (LOC120808851, LOC120810467 and LOC120823934, figure
4) are indicative of ongoing preparation for reproduction. LOC120808851 is located on the sex
chromosome and shows similarity to the ZP3 (Zona pellucida sperm-binding protein 3) Uniprot
entry, a protein that mediates sperm-binding during fertilization. According to Orthofinder results,
it is part of a cluster of duplicated genes on the chrXIX (LOC120808849, LOC120809240,
LOC120808850, LOC120808851) with a single copy conserved on the Y chromosome, which
supports the importance of this function for females. LOC120810467 shows similarity to the
Fucolectin-1 entry, which belongs to a family of genes involved in innate immunity (Honda et al.,
2000) that has been found to be accumulating in European seabass' (Dicentrachus labras) eggs
(Parisi et al., 2010). Finally, LOC120823934 shows similarity to Vitellogenin-2, which is a precursor
to several egg-yolk proteins (Tata, 1976). This result is concordant with the observation of high
levels of estrogen receptors in liver as described before, and further confirmed by the presence of
vtg3, another vitellogenin coding gene, and vtg-2 among the list of most significant genes in liver
(LFC = -5.69). Moreover, as described before, other genes related to response to estrogen also are
among the most significant genes. Moreover, both ZP3 and vitellogenin are genes known to be
expressed in liver (Sano et al., 2017), at least in recent teleosts, further confirming the quality of the
dataset
The liver is the only tissue in which we observed sex-biased expression of mitochondrial genes.
We identified 13 genes with sex-biased gene expression (~56% of expressed genes in the liver's
mitochondria), all male-biased. These genes include ATP6 and 8, COX2, ND1,2,3,5 and two transfer
RNA. In sticklebacks, parental care from the male during the reproductive period, coupled with
gonadal development, is associated with a strong depletion of energy reserves (Chellappa et al.,
1989; Huntingford et al., 2001). While the development of eggs is also costly for females, the strong
involvement in nest building, defence, and parental care by males could generate a high energetic
need in males associated with the metabolic processing of energy reserves in the liver.
414

**Sex-biased Gene Expression on Sexual Chromosome Mostly Reflects Gene Loss in Non-Recombining Regions**

Most genes located on sex chromosomes exhibited sex-biased gene expression (499 in liver, 755 in gills and 813 in brain). Contrasting with the autosomal pattern of sex-biased gene expression, gills and brain exhibited the strongest pattern of SBG, and the higher number of genes expressed in those tissues is not sufficient to explain it (respectively 71%, 83% and 87% of expressed genes in liver, gills and brain). Most SBG are caused by genes having lost their Y copy (83.9%, 71.6% and 70.2% of significant genes in liver, gills and brain), suggesting a lack of global dosage compensation in all studied tissues. Genes orthology relationships for sex chromosomes are available in Table S8. When looking at the ratio of gene expression between sex chromosomes and autosomes, we found that gene expression was greatly lowered uniquely in males for genes having lost their Y copy (95% confidence interval: [-1.77; -1.29] in brain, [-1.5;-1.00] in gills and [-1.50;-1.01] in liver for males; [-0.71; -0.28], [-0.41; -0.03] and [-0.70; -0.07] in females), but not for genes still having a Y orthologue ([-0.48; 0.24], [-0.59,0.22] and [-0.26, 0.18] in males; [-0.23; 0.39], [-0.27;0.5] and [-0.34, 0.71] in females). This pattern holds across all evolutionary strata and tissue (Fig. 5, Table S9). Estimated ratios for the pseudo-autosomal regions or genes with coding sequences on both chromosomes in non-recombining strata did not statistically differ from 0 (Table S9), except for female-biased expression in stratum II in the brain ([0.10, 1.35], and stratum I in the gills ([0.39, 1.61]). Genes having lost their X copy exhibhit a similar pattern but with overall higher median sex-chromosome to autosome expression ratio. In stratum I, it is not statistically different from 0 ([-1.31, 0.02], [-1.44, 0.10] and [-0.92, 0.69]).

Figure 5: Sex chromosomes to autosome expression ratio across sex-chromosome
evolutionary strata. Hemizygous X and hemizygous Y genes respectively lost their Y- and X-
coding sequence, while diploid genes are still have a copy on both chromosome. For diploid
genes, strata was defined using the Y copy position Confidence intervals from 1000
bootstraps are shown. Sample sizes for each median are indicated in Table S8.

Lack of dosage compensation has already been described in the brain (Schultheiß et al., 2015;
White et al., 2015). and our results extend this conclusion to liver in and gills, and to genes with a
lost X copy which had not been included in previous studies. Dosage compensation is expected to
evolve when reduced expression in the heterogametic sex affects phenotype, i.e., affects the
protein level and its interaction network within the organism. In sticklebacks, hemizigous genes
tend to be dosage insensitive, meaning that protein quantities are independent from its expression
level (Peichel et al., 2020). This suggests that there is no selective pressure to evolve dosage
compensation and is corroborated by the fact that conserved genes are dosage-sensitive and
evolving under purifying selection (White et al., 2015).

Apart from chromosome degeneration, gene expression on sex chromosomes is expected to
evolve through several processes. First, as the X chromosome is more often transmitted to females,
it is expected to accumulate dominant female-beneficial mutations that could lead to an increase
in expression of the X copy (Bachtrog et al., 2011). Sex chromosomes are also expected to be
enriched in both active and resolved sexual conflicts, in which case gene expression should
increase depending on the sex in which they are beneficial (Vicoso & Charlesworth, 2006; Bachtrog
et al., 2011). Finally, the lack of recombination and lowered sample size of the Y chromosome can
lead to the accumulation of loss-of-function mutations, leading to the  progressive loss of Y-copy
expression (Charlesworth & Charlesworth, 2000; Bachtrog, 2013). We observed a feminization of

stratum I in gills and II in brain, which had previously been described for stratum II (Leder et al.,
2010; White et al., 2015), suggesting a role for female-beneficial mutations in the evolution of gene
expression on the X chromosome. We did not observe feminization of the pseudo-autosomal region
as previously reported by White et al (2015). This could be caused by the use of autosomal
expression level instead of a closely related species to estimate ancestral expression rate.

To better understand the drivers of sex-biased gene expression of sex chromosomes, we
compared expression of X and Y alleles in sex-biased genes with conserved copies on both
chromosomes (excluding the still recombining PAR) to the allelic expression of unbiased genes (Fig.
6). In all tissues, we found a lowered expression of the Y allele of female-biased genes compared to
unbiased genes (wilcoxon rank-sum test p-value: $1\times10^{-4}$ in brain, $8\times10^{-4}$ in gills and $2\times10^{-2}$ in liver)
while X expression remained similar (all p-value > 0.7), which also resulted in lowered expression
in males in the brain (p-value$1\times10^{-2}$). This suggests that the degeneration of the Y chromosome
coupled with the absence of dosage compensation in this species is the main driver of female-
biased gene evolution, as suggested by previous work (White et al., 2015). Note that we also found
that in the liver overexpression in females also occurred in female-biased genes ($2\times10^{-2}$).

Figure 6: Genotype and allele-specific gene expression across male-biased, unbiased and
female-biased genes. Male-biased and female-biased genes were defined as genes with log2
fold change in expression under -0.5 and above 0.5. We assessed significance using Wilcoxon
rank-sum test comparing within genotype expression levels of male-biased and female-
biased genes to unbiased genes. *: p-value <0.05, ** p-value <0.01, *** p-value <0.001

Similarly, we observed that male-biased gene expression in females ($6\times10^{-5}$, $1\times10^{-2}$ and $7\times10^{-2}$)
but not males (all p-value > $2\times10^{\wedge-1}$) is reduced compared to unbiased genes. We also observed
reduced expression of the X allele in males in the brain and gills ($1\times10^{-4}$, $7\times10^{-3}$), suggesting that this
could be the result of a systematic down-regulation of some genes on the X-chromosome, which

could be a signal of ongoing demasculinization, i.e. the loss of male-advantageous gene on the X
chromosome (Gurbich & Bachtrog, 2008) Concordant with that hypothesis, genes identified as
specific to the Y chromosome tended to have higher expression than genes specific to the X
chromosome in males, suggesting that they are associated with male-beneficial functions.

**Conclusion**

Our study characterizes the sex-specific transcriptome of brain, gills and liver for the threespine
stickleback during its reproductive period. We find low levels of differentiation between sexes in
the brain compared to the level of dimorphism shown in behavioural and morphological studies.
On the opposite, the gills exhibit pronounced sexual dimorphism that is usually not reported or
accounted for in the literature, suggesting that the importance of sex as a cofactor in gill studies
has been underestimated. The liver appeared to be strongly differentiated between the sexes, as
expected for teleost fish. Sex chromosomes are a hotspot of intersex differentiation in all tissues,
with ~70% of genes being differentially expressed. This pattern seems to be caused both by an
ongoing degeneration of the non-recombining region of this sex chromosome coupled with the
absence of dosage compensation mechanism, and a potential repression of male-advantageous
mutations on the X chromosome, although further investigation of gene sequence evolution would
be necessary.

**Acknowledgement:**

Authors are thankful to I. Caza-Allard E. Reni-Nolin for their help during fieldwork and tissue dissection. We thank S. Delaive and F-A Deschênes-Picard for a second year of sampling, although not used in this manuscript. We also thank S. Bernatchez, R. Bouchard, A. Perreault-Payette and C. Berger for occasional assistance and advice during fieldwork. We thank B. Bougas for her help during mRNA extraction and library construction, and Y. Dorant and E. Normandeau for their help during data processing. We also thank C. Venney for english-proofing the manuscript. This project is part of the Ressources Aquatiques Québec (RAQ) research program. L. Bernatchez, who co-conceptualized and co-supervised this project as the PhD advisor of the first author, passed away before the end of it. We are extremely grateful for his involvement, support and motivation until the end.

**Data accessibility:**

Raw sequencing reads and unfiltered read counts are available through NCBI GEO, accession GSE269432. All analysis in the manuscript and related code  are available at https://doi.org/10.5281/zenodo.11477976

**Authors contributions:**

L.B and N.A.H conceptualized and supervised the project. F.S did the fieldwork, generated and filtered the dataset, and performed analyses. N.A.H helped with interpretation. F.S wrote the draft of the manuscript, and N.A.H reviewed and approved it.

**Funding:**

This research was funded by a Fonds de recherche du Québec - Nature et technologies (https://frqnet.frq.gouv.qc.ca/) team grant to L.B. and N.A.-H. (grant number: 254,537). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Conflict of interest**

The authors have no conflict of interest to declare.

**Bibliography:**

Anders S, Pyl PT, Huber W (2015) HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*, **31**, 166–169. https://doi.org/10.1093/bioinformatics/btu638

Bachtrog D (2013) Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nature Reviews Genetics*, **14**, 113–124. https://doi.org/10.1038/nrg3366

Bachtrog D, Kirkpatrick M, Mank JE, McDaniel SF, Pires JC, Rice W, Valenzuela N (2011) Are all sex chromosomes created equal? *Trends in Genetics*, **27**, 350–357. https://doi.org/10.1016/j.tig.2011.05.005

Beal AP, Martin FD, Hale MC (2018) Using RNA-seq to determine patterns of sex-bias in gene expression in the brain of the sex-role reversed Gulf Pipefish (Syngnathus scovelli). *Marine Genomics*, **37**, 120–127. https://doi.org/10.1016/j.margen.2017.09.005

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, **57**, 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bereketoğlu MB, Abdullayev R, Tuğ Bozdoğan S (2022) Current Approach to Genetic Causes of Male Infertility and Genetic Counseling. *Düzce Tıp Fakültesi Dergisi*, **24**, 7–16. https://doi.org/10.18678/dtfd.1183283

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421. https://doi.org/10.1186/1471-2105-10-421

Chang JP, Wong AOL (2009) Chapter 4 Growth Hormone Regulation in Fish. In: *Fish Physiology* , pp. 151–195. Elsevier. https://doi.org/10.1016/S1546-5098(09)28004-6

Charlesworth B, Charlesworth D (2000) The degeneration of Y chromosomes (B Charlesworth, PH Harvey, Eds,). *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, **355**, 1563–1572. https://doi.org/10.1098/rstb.2000.0717

Chellappa S, Huntingford FA, Strang RHC, Thomson RY (1989) Annual variation in energy reserves in male three-spined stickleback, Gasterosteus aculeatm L. (Pisces, Gasterosteidae). *Journal of Fish Biology*, **35**, 275–286. https://doi.org/10.1111/j.1095-8649.1989.tb02976.x

Chen Y, Tang H, He J, Wu X, Wang L, Liu X, Lin H (2019) Interaction of nuclear ERs and GPER in vitellogenesis in zebrafish. *The Journal of Steroid Biochemistry and Molecular Biology*, **189**, 10–18. https://doi.org/10.1016/j.jsbmb.2019.01.013

Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, **34**, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Chung A-Y, Kim S, Kim E, Kim D, Jeong I, Cha YR, Bae Y, Park SW, Lee J, Park H-C (2013) Indian Hedgehog b Function Is Required for the Specification of Oligodendrocyte Progenitor Cells in the Zebrafish CNS. *The Journal of Neuroscience*, **33**, 1728–1733. https://doi.org/10.1523/JNEUROSCI.3369-12.2013

Concha MI, Molina S, Oyarzún C, Villanueva J, Amthauer R (2003) Local expression of apolipoprotein A-I gene and a possible role for HDL in primary defence in the carp skin. *Fish & Shellfish Immunology*, **14**, 259–273. https://doi.org/10.1006/fsim.2002.0435

Darolti I, Mank JE (2023) Sex-biased gene expression at single-cell resolution: cause and consequence of sexual dimorphism. *Evolution Letters*, **7**, 148–156. https://doi.org/10.1093/evlett/qrad013

Darolti I, Wright AE, Sandkam BA, Morris J, Bloch NI, Farré M, Fuller RC, Bourne GR, Larkin DM, Breden F, Mank JE (2019) Extreme heterogeneity in sex chromosome differentiation and dosage compensation in livebearers. *Proceedings of the National Academy of Sciences*, **116**, 19031–19036. https://doi.org/10.1073/pnas.1905298116

Devlin RH, Nagahama Y (2002) Sex determination and sex differentiation in fish: an overview of genetic, physiological, and environmental influences. *Aquaculture*, **208**, 191–364. https://doi.org/10.1016/S0044-8486(02)00057-1

Djordjevic J, Dumas Z, Robinson-Rechavi M, Schwander T, Parker DJ (2022) Dynamics of sex-biased gene expression during development in the stick insect Timema californicum. *Heredity*, **129**, 113–122. https://doi.org/10.1038/s41437-022-00536-y

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15–21. https://doi.org/10.1093/bioinformatics/bts635

Dominguez GA, Bisesi JH, Kroll KJ, Denslow ND, Sabo-Attwood T (2014) Control of Transcriptional Repression of the Vitellogenin Receptor Gene in Largemouth Bass (Micropterus Salmoides) by Select Estrogen Receptors Isotypes. *Toxicological Sciences*, **141**, 423–431. https://doi.org/10.1093/toxsci/kfu145

Dymowska AK, Schultz AG, Blair SD, Chamot D, Goss GG (2014) Acid-sensing ion channels are involved in epithelial Na$^+$ uptake in the rainbow trout *Oncorhynchus mykiss*. *American Journal of Physiology-Cell Physiology*, **307**, C255–C265. https://doi.org/10.1152/ajpcell.00398.2013

Ellegren H, Parsch J (2007) The evolution of sex-biased genes and sex-biased gene expression. *Nature Reviews Genetics*, **8**, 689–698. https://doi.org/10.1038/nrg2167

Emms DM, Kelly S (2019) OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology*, **20**, 238. https://doi.org/10.1186/s13059-019-1832-y

Filby AL, Tyler CR (2005) Molecular Characterization of Estrogen Receptors 1, 2a, and 2b and Their Tissue and Ontogenic Expression Profiles in Fathead Minnow (Pimephales promelas)1. *Biology of Reproduction*, **73**, 648–662. https://doi.org/10.1095/biolreprod.105.039701

Gonzalez P, Baudrimont M, Boudou A, Bourdineaud J-P (2006) Comparative Effects of Direct Cadmium Contamination on Gene Expression in Gills, Liver, Skeletal Muscles and Brain of the Zebrafish (Danio rerio). *BioMetals*, **19**, 225–235. https://doi.org/10.1007/s10534-005-5670-x

Gurbich TA, Bachtrog D (2008) Gene content evolution on the X chromosome. *Current Opinion in Genetics & Development*, **18**, 493–498. https://doi.org/10.1016/j.gde.2008.09.006

Hale MC, McKinney GJ, Thrower FP, Nichols KM (2018) Evidence of sex-bias in gene expression in the brain transcriptome of two populations of rainbow trout (Oncorhynchus mykiss) with divergent life histories (CS Rosenfeld, Ed,). *PLOS ONE*, **13**, e0193009. https://doi.org/10.1371/journal.pone.0193009

Halestrap AP (2012) The monocarboxylate transporter family—Structure and functional characterization. *IUBMB Life*, **64**, 1–9. https://doi.org/10.1002/iub.573

Honda S, Kashiwagi M, Miyamoto K, Takei Y, Hirose S (2000) Multiplicity, Structures, and Endocrine
and Exocrine Natures of Eel Fucose-binding Lectins. *Journal of Biological Chemistry*, **275**, 33151–
33157. https://doi.org/10.1074/jbc.M002337200

Huntingford FA, Chellappa S, Taylor AC, Strang RHC (2001) Energy reserves and reproductive
investment in male three-spined sticklebacks, *Gasterosteus aculeatus*. *Ecology of Freshwater*
*Fish*, **10**, 111–117. https://doi.org/10.1034/j.1600-0633.2001.100206.x

Kaitetzidou E, Gilfillan GD, Antonopoulou E, Sarropoulou E (2022) Sex-biased dynamics of three-
spined stickleback (Gasterosteus aculeatus) gene expression patterns. *Genomics*, **114**, 266–277.
https://doi.org/10.1016/j.ygeno.2021.12.010

Kitano J, Kakioka R, Ishikawa A, Toyoda A, Kusakabe M (2020) Differences in the contributions of
sex linkage and androgen regulation to sex-biased gene expression in juvenile and adult
sticklebacks. *Journal of Evolutionary Biology*, **33**, 1129–1138. https://doi.org/10.1111/jeb.13662

Kobayashi Y, Nagahama Y, Nakamura M (2013) Diversity and Plasticity of Sex Determination and
Differentiation in Fishes. *Sexual Development*, **7**, 115–125. https://doi.org/10.1159/000342009

Kotrschal A, Räsänen K, Kristjánsson BK, Senn M, Kolm N (2012) Extreme Sexual Brain Size
Dimorphism in Sticklebacks: A Consequence of the Cognitive Challenges of Sex and Parenting?
(A Iwaniuk, Ed,). *PLoS ONE*, **7**, e30055. https://doi.org/10.1371/journal.pone.0030055

Laing LV, Viana J, Dempster EL, Uren Webster TM, Van Aerle R, Mill J, Santos EM (2018) Sex-specific
transcription and DNA methylation profiles of reproductive and epigenetic associated genes in
the gonads and livers of breeding zebrafish. *Comparative Biochemistry and Physiology Part A:*
*Molecular & Integrative Physiology*, **222**, 16–25. https://doi.org/10.1016/j.cbpa.2018.04.004

Lande R (1980) Sexual dimorphism, sexual selection, and adaptation in polygenic characters.
*Evolution*, **34**, 292–305. https://doi.org/10.1111/j.1558-5646.1980.tb04817.x

Leder EH, Cano JM, Leinonen T, O'Hara RB, Nikinmaa M, Primmer CR, Merilä J (2010) Female-Biased
Expression on the X Chromosome as a Key Step in Sex Chromosome Evolution in Threespine
Sticklebacks. *Molecular Biology and Evolution*, **27**, 1495–1503.
https://doi.org/10.1093/molbev/msq031

Li P, Chen J, Zhu C, Pan Z, Li Q, Wei H, Wang G, Cheng W, Fu B, Sun Y (2022) DNA Methylation
Difference between Female and Male Ussuri Catfish (Pseudobagrus ussuriensis) in Brain and
Gonad Tissues. *Life*, **12**, 874. https://doi.org/10.3390/life12060874

Li Y, Ge X, Peng F, Li W, Li JJ (2022) Exaggerated false positives by popular differential expression
methods when analyzing human population samples. *Genome Biology*, **23**, 79.
https://doi.org/10.1186/s13059-022-02648-4

Lichilín N, El Taher A, Böhne A (2021) Sex-biased gene expression and recent sex chromosome
turnover. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **376**, 20200107.
https://doi.org/10.1098/rstb.2020.0107

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biology*, **15**, 550. https://doi.org/10.1186/s13059-014-0550-8

Ma F, Martínez-San Segundo P, Barceló V, Morancho A, Gabriel-Salazar M, Giralt D, Montaner J,
Rosell A (2016) Matrix metalloproteinase-13 participates in neuroprotection and neurorepair

after cerebral ischemia in mice. *Neurobiology of Disease*, **91**, 236–246. https://doi.org/10.1016/j.nbd.2016.03.016

Mank JE (2013) Sex chromosome dosage compensation: definitely not for everyone. *Trends in Genetics*, **29**, 677–683. https://doi.org/10.1016/j.tig.2013.07.005

Meng S, Qiu L, Hu G, Fan L, Song C, Zheng Y, Wu W, Qu J, Li D, Chen J, Xu P (2016) Effects of methomyl on steroidogenic gene transcription of the hypothalamic-pituitary-gonad-liver axis in male tilapia. *Chemosphere*, **165**, 152–162. https://doi.org/10.1016/j.chemosphere.2016.09.024

Mitchell SO, Rodger HD (2011) A review of infectious gill disease in marine salmonid fish: Infectious gill disease in salmonids. *Journal of Fish Diseases*, **34**, 411–432. https://doi.org/10.1111/j.1365-2761.2011.01251.x

Moll P, Ante M, Seitz A, Reda T (2014) QuantSeq 3′ mRNA sequencing for RNA quantification. *Nature Methods*, **11**, i–iii. https://doi.org/10.1038/nmeth.f.376

Muriach B, Carrillo M, Zanuy S, Cerdá-Reverter JM (2008) Distribution of estrogen receptor 2 mRNAs (Esr2a and Esr2b) in the brain and pituitary of the sea bass (Dicentrarchus labrax). *Brain Research*, **1210**, 126–141. https://doi.org/10.1016/j.brainres.2008.02.053

Naftaly AS, Pau S, White MA (2021) Long-read RNA sequencing reveals widespread sex-specific alternative splicing in threespine stickleback fish. *Genome Research*, **31**, 1486–1497. https://doi.org/10.1101/gr.274282.120

Nath S, Shaw DE, White MA (2021) Improved contiguity of the threespine stickleback genome using long-read sequencing. *G3 Genes|Genomes|Genetics*, **11**, jkab007. https://doi.org/10.1093/g3journal/jkab007

Nelson ER, Habibi HR (2013) Estrogen receptor function and regulation in fish and other vertebrates. *General and Comparative Endocrinology*, **192**, 15–24. https://doi.org/10.1016/j.ygcen.2013.03.032

Parisi MG, Cammarata M, Benenati G, Salerno G, Mangano V, Vizzini A, Parrinello N (2010) A serum fucose-binding lectin (DlFBL) from adult Dicentrarchus labrax is expressed in larva and juvenile tissues and contained in eggs. *Cell and Tissue Research*, **341**, 279–288. https://doi.org/10.1007/s00441-010-1004-6

Peichel CL, McCann SR, Ross JA, Naftaly AFS, Urton JR, Cech JN, Grimwood J, Schmutz J, Myers RM, Kingsley DM, White MA (2020) Assembly of the threespine stickleback Y chromosome reveals convergent signatures of sex chromosome evolution. *Genome Biology*, **21**, 177. https://doi.org/10.1186/s13059-020-02097-x

Perry SF, Shahsavarani A, Georgalis T, Bayaa M, Furimsky M, Thomas SLY (2003) Channels, pumps, and exchangers in the gill and kidney of freshwater fishes: Their role in ionic and acid-base regulation. *Journal of Experimental Zoology Part A: Comparative Experimental Biology*, **300A**, 53–62. https://doi.org/10.1002/jez.a.10309

Primmer CR, Papakostas S, Leder EH, Davis MJ, Ragan MA (2013) Annotated genes and nonannotated genomes: cross-species use of Gene Ontology in ecology and evolution research. *Molecular Ecology*, **22**, 3216–3241. https://doi.org/10.1111/mec.12309

Qiao Q, Le Manach S, Sotton B, Huet H, Duvernois-Berthet E, Paris A, Duval C, Ponger L, Marie A, Blond A, Mathéron L, Vinh J, Bolbach G, Djediat C, Bernard C, Edery M, Marie B (2016) Deep sexual

dimorphism in adult medaka fish liver highlighted by multi-omic approach. *Scientific Reports*, **6**, 32459. https://doi.org/10.1038/srep32459

R Core Team (2021) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Renn SCP, Aubin-Horth N, Hofmann HA (2008) Fish and chips: functional genomics of social plasticity in an African cichlid fish. *Journal of Experimental Biology*, **211**, 3041–3056. https://doi.org/10.1242/jeb.018242

Rice WR (1984) Sex chromosomes and the evolution of sexual dimorphism. *Evolution*, **38**, 735–742. https://doi.org/10.2307/2408385

Rodríguez-Montes L, Ovchinnikova S, Yuan X, Studer T, Sarropoulos I, Anders S, Kaessmann H, Cardoso-Moreira M (2023) Sex-biased gene expression across mammalian organ development and evolution. *Science*, **382**, eadf1046. https://doi.org/10.1126/science.adf1046

Rossum G van, Drake FL (2010) *The Python language reference*. Python Software Foundation, Hampton, NH.

Sano K, Kawaguchi M, Katano K, Tomita K, Inokuchi M, Nagasawa T, Hiroi J, Kaneko T, Kitagawa T, Fujimoto T, Arai K, Tanaka M, Yasumasu S (2017) Comparison of Egg Envelope Thickness in Teleosts and its Relationship to the Sites of ZP Protein Synthesis. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, **328**, 240–258. https://doi.org/10.1002/jez.b.22729

Schneider M, Lane L, Boutet E, Lieberherr D, Tognolli M, Bougueleret L, Bairoch A (2009) The UniProtKB/Swiss-Prot knowledgebase and its Plant Proteome Annotation Program. *Journal of Proteomics*, **72**, 567–573. https://doi.org/10.1016/j.jprot.2008.11.010

Schultheiß R, Viitaniemi HM, Leder EH (2015) Spatial Dynamics of Evolving Dosage Compensation in a Young Sex Chromosome System. *Genome Biology and Evolution*, **7**, 581–590. https://doi.org/10.1093/gbe/evv013

Schumer M, Krishnakant K, Renn SCP (2011) Comparative gene expression profiles for highly similar aggressive phenotypes in male and female cichlid fishes ( *Julidochromis* ). *Journal of Experimental Biology*, **214**, 3269–3278. https://doi.org/10.1242/jeb.055467

Scott GR, Richards JG, Forbush B, Isenring P, Schulte PM (2004) Changes in gene expression in gills of the euryhaline killifish *Fundulus heteroclitus* after abrupt salinity transfer. *American Journal of Physiology-Cell Physiology*, **287**, C300–C309. https://doi.org/10.1152/ajpcell.00054.2004

Sutherland BJG, Prokkola JM, Audet C, Bernatchez L (2019) Sex-Specific Co-expression Networks and Sex-Biased Gene Expression in the Salmonid Brook Charr *Salvelinus fontinalis*. *G3 Genes|Genomes|Genetics*, **9**, 955–968. https://doi.org/10.1534/g3.118.200910

Tata JR (1976) The expression of the vitellogenin gene. *Cell*, **9**, 1–14. https://doi.org/10.1016/0092-8674(76)90047-7

Telonis-Scott M, Kopp A, Wayne ML, Nuzhdin SV, McIntyre LM (2009) Sex-Specific Splicing in Drosophila: Widespread Occurrence, Tissue Specificity and Evolutionary Conservation. *Genetics*, **181**, 421–434. https://doi.org/10.1534/genetics.108.096743

The UniProt Consortium (2023) UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Research*, **51**, D523–D531. https://doi.org/10.1093/nar/gkac1052

Thresher RE (1984) *Reproduction in reef fishes*. T.F.H. Publications; Distributed in the U.S. by T.F.H.
Publications, British Crown Colony of Hong Kong: Neptune City, NJ.

Tosto NM, Beasley ER, Wong BBM, Mank JE, Flanagan SP (2023) The roles of sexual selection and
sexual conflict in shaping patterns of genome and transcriptome variation. *Nature Ecology &*
*Evolution*, **7**, 981–993. https://doi.org/10.1038/s41559-023-02019-7

Van Der Meer DLM, Van Den Thillart GEEJM, Witte F, De Bakker MAG, Besser J, Richardson MK,
Spaink HP, Leito JTD, Bagowski CP (2005) Gene expression profiling of the long-term adaptive
response to hypoxia in the gills of adult zebrafish. *American Journal of Physiology-Regulatory,*
*Integrative and Comparative Physiology*, **289**, R1512–R1519.
https://doi.org/10.1152/ajpregu.00089.2005

Varadharajan S, Rastas P, Löytynoja A, Matschiner M, Calboli FCF, Guo B, Nederbragt AJ, Jakobsen
KS, Merilä J (2019) *Genome sequencing of the nine-spined stickleback (* Pungitius pungitius *)*
*provides insights into chromosome evolution*. Genomics. https://doi.org/10.1101/741751

Vicoso B, Charlesworth B (2006) Evolution on the X chromosome: unusual patterns and processes.
*Nature Reviews Genetics*, **7**, 645–653. https://doi.org/10.1038/nrg1914

Viitaniemi HM, Leder EH (2011) Sex-Biased Protein Expression in Threespine Stickleback,
*Gasterosteus aculeatus*. *Journal of Proteome Research*, **10**, 4033–4040.
https://doi.org/10.1021/pr200234a

White MA, Kitano J, Peichel CL (2015) Purifying Selection Maintains Dosage-Sensitive Genes during
Degeneration of the Threespine Stickleback Y Chromosome. *Molecular Biology and Evolution*,
**32**, 1981–1995. https://doi.org/10.1093/molbev/msv078

Wijchers PJ, Yandim C, Panousopoulou E, Ahmad M, Harker N, Saveliev A, Burgoyne PS, Festenstein
R (2010) Sexual Dimorphism in Mammalian Autosomal Gene Regulation Is Determined Not Only
by Sry but by Sex Chromosome Complement As Well. *Developmental Cell*, **19**, 477–484.
https://doi.org/10.1016/j.devcel.2010.08.005

Wu Q, Zhang Y, Chen Q (2001) Indian hedgehog Is an Essential Component of Mechanotransduction
Complex to Stimulate Chondrocyte Proliferation. *Journal of Biological Chemistry*, **276**, 35290–
35296. https://doi.org/10.1074/jbc.M101055200

Yuan W, Jiang S, Sun D, Wu Z, Wei C, Dai C, Jiang L, Peng S (2019) Transcriptome profiling analysis
of sex-based differentially expressed mRNAs and lncRNAs in the brains of mature zebrafish
(Danio rerio). *BMC Genomics*, **20**, 830. https://doi.org/10.1186/s12864-019-6197-9

767