The MICOS Complex Regulates Mitochondrial Structure and Oxidative Stress During Age-Dependent Structural Deficits in the Kidney

Zer Vue[1*], Praveena Prasad[2*], Han Le[1*], Kit Neikirk[1], Chanel Harris[1], Edgar Garza-Lopez[3], Eric Wang[4], Alexandria Murphy[2], Brenita Jenkins[2], Larry Vang[1], Estevão Scudese[1], Bryanna Shao[1], Ashlesha Kadam[5], Jianqiang Shao[6], Andrea G. Marshall[1], Amber Crabtree[1], Benjamin Kirk[3], Alice Koh[1], Genesis Wilson[1], Ashton Oliver[1], Taylor Rodman[1], Kinuthia Kabugi[1], Ho-Jin Koh[7], Quinton Smith[4], Elma Zaganjor[1], Celestine N. Wanjalla[8], Chandravanu Dash[9], Chantell Evans[10], Mark A. Phillips[11], David Hubert[11], Olujimi Ajijola[12], Aaron Whiteside[13], Young Do Koo[3,14], André Kinder[15], Mert Demirci[8], Claude F. Albritton[8,16], Nelson Wandira[8], Sydney Jamison[8], Taseer Ahmed[8], Mohammad Saleem[8], Dhanendra Tomar[5], Clintoria R. Williams[13], Mariya T. Sweetwyne[17], Sandra A. Murray[18], Anthonya Cooper[18], Annet Kirabo[8,19,20,21], Pooja Jadiya[22], Anita Quintana[23], Prasanna Katti[24,25], Dao☐Fu Dai[26], Melanie R. McReynolds[#2], Antentor Hinton, Jr[#1]

Affiliations:
1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA.
2. Department of Biochemistry and Molecular Biology, The Huck Institute of the Life Sciences, Pennsylvania State University, State College, PA 16801
3. Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA.
4. Department of Chemical and Biomolecular Engineering, University of California, Irvine, CA, 92697, USA.
5. Department of Internal Medicine, Section of Cardiovascular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157 USA
6. Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA.
7. Department of Biological Sciences, Tennessee State University, Nashville, TN 37209, USA
8. Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232
9. Department of Biochemistry, Cancer Biology, Pharmacology and Neuroscience, Meharry Medical College, Nashville, TN, United States
10. Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27708, USA
11. Department of Integrative Biology, Oregon State University, Corvallis, OR, 97331, USA.
12. UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA, USA.
13. Department of Neuroscience, Cell Biology and Physiology, Wright State University, Dayton, OH 45435 USA
14. Fraternal Order of Eagles Diabetes Research Center, Iowa City, Iowa, USA
15. Artur Sá Earp Neto University Center - UNIFASE-FMP, Petrópolis Medical School, Brazil
16. Department of Biomedical Sciences, School of Graduate Studies, Meharry Medical College, Nashville, TN 37208-3501, USA.
17. Department of Laboratory Medicine and Pathology, University of Washington, Seattle, WA, 98195, USA
18. Department of Cell Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA, 15260, USA
19. Vanderbilt Center for Immunobiology, Vanderbilt University, Nashville, TN, 37232, USA.

20. Vanderbilt Institute for Infection, Immunology and Inflammation, Vanderbilt University,
Nashville, TN, 37232, USA.
21. Vanderbilt Institute for Global Health, Vanderbilt University, Nashville, TN, 37232, USA.
22. Department of Internal Medicine, Section of Gerontology and Geriatric Medicine, Sticht
Center for Healthy Aging and Alzheimer's Prevention, Wake Forest University School of
Medicine, Winston-Salem, NC
23. Department of Biological Sciences, Border Biomedical Research Center, The University of
Texas at El Paso, El Paso, Texas, USA
24. National Heart, Lung and Blood Institute, National Institutes of Health, 9000 Rockville Pike,
Bethesda, MD 20892, USA
25. Department of Biology, Indian Institute of Science Education and Research (IISER) Tirupati,
AP, 517619, India
26. Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD,
USA.
61
62   *These authors share co-first authorship.
63   #These authors share co-senior authorship.
64
65   Corresponding Author:
66   Antentor Hinton
67   Department of Molecular Physiology and Biophysics
68   Vanderbilt University
69   antentor.o.hinton.jr@Vanderbilt.Edu 319-383-3095
70
71

72   **ABSTRACT**:
73

The kidney filters nutrient waste and bodily fluids from the bloodstream, in addition to
secondary functions of metabolism and hormone secretion, requiring an astonishing amount of
energy to maintain its functions. In kidney cells, mitochondria produce adenosine triphosphate
(ATP) and help maintain kidney function. Due to aging, the efficiency of kidney functions
begins to decrease. Dysfunction in mitochondria and cristae, the inner folds of mitochondria, is a
hallmark of aging. Therefore, age-related kidney function decline could be due to changes in
mitochondrial ultrastructure, increased reactive oxygen species (ROS), and subsequent
alterations in metabolism and lipid composition. We sought to understand if there is altered
mitochondrial ultrastructure, as marked by 3D morphological changes, across time in tubular
kidney cells. Serial block facing-scanning electron microscope (SBF-SEM) and manual
segmentation using the Amira software were used to visualize murine kidney samples during the
aging process at 3 months (young) and 2 years (old). We found that 2-year mitochondria are
more fragmented, compared to the 3-month, with many uniquely shaped mitochondria observed
across aging, concomitant with shifts in ROS, metabolomics, and lipid homeostasis.
Furthermore, we show that the mitochondrial contact site and cristae organizing system
(MICOS) complex is impaired in the kidney due to aging. Disruption of the MICOS complex
shows altered mitochondrial calcium uptake and calcium retention capacity, as well as
generation of oxidative stress. We found significant, detrimental structural changes to aged
kidney tubule mitochondria suggesting a potential mechanism underlying why kidney diseases

occur more readily with age. We hypothesize that disruption in the MICOS complex further
exacerbates mitochondrial dysfunction, creating a vicious cycle of mitochondrial degradation
and oxidative stress, thus impacting kidney health.
96
97 **Keywords**: Mitochondria; Metabolism; Kidney; 3DEM; ROS; MICOS complex
98

INTRODUCTION:

Kidneys are principally known for their role in the excretion of waste products, though their full functions go far beyond, including hormonal signaling, which makes the kidney critical for many other functions, such as blood pressure regulation [1]. However, dysfunction may occur in various states, such as the sudden loss of kidney function, acute kidney injury (AKI), and chronic kidney diseases (CKD) [2] of various etiologies. An estimated 90% of individuals are not aware of their AKI and CKD, thus the exact magnitude of kidney diseases remains difficult to measure [3]. However, poor treatment outcomes for kidney diseases and associations with other conditions, such as cardiovascular disease, emphasize the importance of developing new effective treatments [4]. Kidneys are among the most mitochondrial-rich tissues in the body[5], therefore one approach to expand our understanding of kidney pathological processes is the study of mitochondria's roles in kidney functions [5].

It is well established that mitochondria provide numerous critical cellular functions beyond oxidative phosphorylation[6]. Mitochondria play a role in cell signaling, calcium regulation, apoptosis, and general homeostasis. Mitochondrial genes also encode the pathways responsible for ATP generation[7]. Mitochondria dysfunction has been implicated in kidney diseases [8–11] as well as other diseases impacting mitochondrial-rich organs including muscle diseases [12,13], neurological diseases [14,15], and obesity-related diabetes [16,17]. Notably, specific mechanisms of mitochondrial dysfunction that may govern each disease state, and ways to rescue mitochondria function, remain nascent research topics. Therefore, it is important to explore the therapeutic relevance of mitochondria.

Studies have shown that mitochondrial dysfunction is one of the hallmarks of AKI pathogenesis, making mitochondria a critical target to restore kidney function to pre-disease states [2,18]. Other focus areas for kidney research include autosomal dominant polycystic kidney disease, which has been shown to shift mitochondrial function to anaerobic respiration[19], limiting oxidative capacity, through numerous mechanisms, such as calcium signaling. Mitochondrial primary roles, frequency, and connections in different tissues may vary significantly. For mitochondrial calcium regulation, the mitochondrial calcium uniporter (MCU), which regulates calcium influx in mitochondria, is more active in the kidney than in other mitochondrial-rich organs like the liver and the heart [20]. Still, the mechanisms that mediate kidney mitochondrial dynamics remain unclear.

One link between mitochondria and kidney diseases is aging. Aging is the greatest risk factor for both CKD and AKI. While mitochondrial dysfunction is a hallmark of aging [21], an increased risk of AKI and CKD is believed to be due to aging [22,23]. It is well established that mitochondria across organ systems lose peak function with aging [24–26]. Past research has implicated that in mouse kidneys, mitochondrial bioenergetics is lost due to proton leak, reducing electron transfer [27]. Furthermore, clearance of damaged mitochondria via mitophagy responses are also blunted across aging in kidney proximal tubules [28]. However, it is poorly understood how mitochondria function declines with age and the relationship between mitochondrial dysfunction and mitochondrial structure in kidneys. It is possible that similar to other models, there is decreased fusion in the mitochondria, resulting in damaged mitochondria and cristae breakdown. This was shown by previous studies looking at kidneys with age via transmission electron microscopy (TEM) [29]. However, investigating mitochondria at 2D does not provide sufficient details and subcellular structures of mitochondria [30,31]. Thus, we performed 3D reconstruction through serial block face-scanning electron microscopy, which allows for a broader analysis range [32,33], to determine how mitochondrial networking and broad structures are

altered across aging in mouse kidneys. We studied mice at two ages: 3-month mice, representing
a young adult phenotype, and 2-year mice, representing a geriatric model [34].
Here, we investigated changes in the structure and physical appearance of kidneys in
young and old groups. We then studied mitochondria and cristae morphological changes in both
2D and 3D, and mitochondrial reactive oxygen species (ROS) production in young and aged
mice. Mitochondrial ROS synthesis mainly occurs on the electron transport chain located in the
inner mitochondrial membrane during the process of oxidative phosphorylation [35,36]. Damaged or
dysfunctional mitochondria are harmful to the cells because they release substances that promote
cell death and create ROS which causes TEC apoptosis [36]. The relationships between
mitochondrial oxidative stress, ROS production, and mitophagy are closely intertwined, and
these processes are all involved in pathological conditions of AKI [36].
Past studies have suggested that dysfunction of the mitochondrial contact site and cristae
organizing system (MICOS) complex, a group of proteins regulating cristae morphology, can
cause oxidative stress [37]. Given that we have previously found age-dependent losses in the
MICOS complex correlating with mitochondrial structural defects in skeletal muscle and cardiac
tissue [38,39], we hypothesized that a similar phenotype may be observed in the kidney, which
could confer oxidative damage and impaired calcium homeostasis characteristic of age-related
AKI and CKD [40,41]. We found that deletion of the MICOS genes resulted in mitochondrial
structural changes and impairments in mitochondrial calcium regulation in the kidney. Because
we showed that aging affects both the kidneys and mitochondria structure and MICOS, we
investigated the metabolomic and lipidomic changes in aging kidneys to understand further the
pathways that may mediate mitochondrial and MICOS roles in aging kidney vulnerability to
injury and disease.
168
Results:
Human Aging Causes Minimal Changes in Kidney Size
Previous studies have utilized magnetic resonance imaging of solid renal masses as a proxy for
pathologic classification and defining kidney structure [42,43]. Generally, it has been found that
following the age of 60, there is a reduction in kidney volume at a rate of approximately 16 cubic
centimeters per decade [44]. Thus, we utilized magnetic resonance imaging to determine how the
kidney is remodeled during the aging process. By enrolling female and male participants
(Figures 1A–D), we created a "young" cohort (n = 14) consisting of individuals under 50 years
old and an "old" cohort of individuals older than 60 years old (n = 20) (Supplemental File 1). For
both sexes, the kidney area did not show a significant change (Figures 1E–F). In-phase, which
refers to aligned fat and water molecules, and out-of-phase, or opposed phase, intensity were
similarly minimally changed in both females and males across aging (Figures 1G-J). We did
observe that the old cohort of males had a significantly reduced in-phase intensity (Figure 1H).
From there, we proceeded to calculate a ratio of in-phase to out-of-phase intensity, which
showed no significant differences (Figures 1K-L). When male and female subjects were
combined, kidney from male and female was not significantly differentiated across the aging
process (Supplemental Figure 1A). We generally observed minimal sex-dependent differences
during the aging process (Supplemental Figures 1B–E). Interestingly, old females did have a
significantly lower cross-sectional area than aged males, increasing a potential increased
susceptibility to aging with sex-dependent differences in kidney aging (Figure 1B). While we
could not confirm subjects had kidney disease, these results support a slight age-related decline

in kidney structure. However, while gross morphological changes may be minimal, we sought to
further elucidate tissue changes with aging.
192
Murine in Aging Results in Interstitial Fibrosis and Oxidative Stress
Previous studies have shown that interstitial fibrosis on kidney biopsy is regarded as a
prognostic indicator, although its effectiveness as a diagnostic marker can be mixed, generally it
can represent a nephropathy [45,46]. We looked at young (4–5 months old; Figure 2A) and old (21–
23 months old; Figure 2B) C57BL/6J mice with trichrome blue to stain connective tissue blue.
Concurrent with previous studies [47], we found that the trichrome area percentage increased in
comparing young (5.8%) and old mice (11.0%), indicating a higher degree of interstitial fibrosis
(Figure 2C). From there, we used immunohistochemistry to look at nitrotyrosin, as stained in
brown areas. Studies have shown that increased nitrotyrosine levels correlate with renal
dysfunction and inflammatory processes, serving as a biomarker for kidney diseases such as AKI
and CKD, as well as overall mortality in these disease states [48–50]. Our results showed that a
significant increase in nitrotyrosine in tubular epithelial cells and podocytes of old mice, as
compared to their young counterparts (Figures 2D-F). This indicates that oxidative stress occurs
with aging, so we sought to understand how mitochondrial structure also changes.
207
208
Ultrastructural Analysis of TEM shows Larger Mitochondria with Poor Cristae
Morphology
Previous research has indicated that aging can influence the dynamics and morphological
structure of the mitochondria in the kidney [51]. To see mitochondrial and cristae structure changes
across aging, we first looked at the mitochondrial and cristae TEM images in 3-month and 2-
year-old mice (Figures 3A-B) tubules. Interestingly, we noticed that the area of the mitochondria
increased when comparing 3-month [mean of 1.57 μm^2 ± 2.09 μm^2 standard deviation (SD)] and
2-year-old (2.54 μm^2 ± 3.33 μm^2 SD) mice (Figure 3C). Similarly, the circularity index of
mitochondria increased with age between 3-month (0.746 ± 0.174 SD) and 2-year (0.813 ± 0.116
SD) mice (Figure 3D). However, the mitochondrial number did not significantly change (Figure
3E). This suggests that mitochondria have higher normalcy with aging, with more relative area
for oxidative phosphorylation, but these increases may also indicate pathological swelling of
mitochondria [52]. Thus, we turned our attention to cristae, the inner folds of mitochondria that
regulate bioenergetics [53], which showed significant losses in structural integrity across aging in
the kidney. Specifically, the number of cristae decreased consistently across aging (Figures 3A-
B). We looked at cristae score, which is a qualitative ordinal scale that assigns a whole-number
score from "0" to "4" representing little-to-no cristae and normal well-formed cristae,
respectively [54–56]. We found that cristae underwent significant age-related defects, with 3-month
samples having mostly well-formed cristae (mean cristae score of 3.44 ± 0.794 SD) while 2-year
samples showing few-well-formed examples of cristae (1.71 ± 0.693 SD) (Figure 3F). To verify
these findings, we performed a qualitative assessment in SBF-SEM of cristae. We noted the 3D
morphology of cristae in the 3-month sample (Figure 3G) was well formed and plentiful,
concurrent with our TEM findings; in contrast, 2-year samples (Figure 3H) qualitatively showed
many more areas lacking cristae, and the cristae present was often scattered or limited in area.
Together, based on these findings, aging in murine kidneys results in disrupted cristae
and larger mitochondria, which could be due to swelling. However, it should be noted that TEM
can be limited in analyzing mitochondrial changes, beyond those of cristae structure, across the

aging process, especially in the context of capturing mitochondrial changes in volume. Thus, we
shifted to using SBF-SEM to rigorously analyze mitochondrial volumetric changes.
238
239

SBF-SEM Reveals Aging Results in Reduced Mitochondrial Volume in Kidney Tissue
Based on our observations of the lack of cristae folding in aging kidney tissue, we used 3-
D techniques to image kidney tissue biopsies from young (3-month-old) and aged (2-year-old)
mice with SBF-SEM. With ranges of 10 µm for the x- and y-planes and 50 µm for the z-plane,
SBF-SEM enables 3-D reconstruction of mitochondria with an accurate spatial resolution that
cannot be seen in 2-D [32,33,57]. To elucidate the changes in intermyofibrillar mitochondria in
relation to aging, we surveyed approximately 250 intermyofibrillar mitochondria from each of
the four male mice (n = 4) sampled (Figure 4A) at each age time point with SBF-SEM 3-D
reconstruction methods, for a total of 1702 mitochondria surveyed cumulatively across age-
points. At a 10 µm by 10 µm image stack resolution, ∼300 serial section orthogonal (ortho)
slices with a total imaging depth of 50 µM (Figure 4B) were manually traced at transverse
intervals (Figure 4C). This allowed for 3-D reconstructions of mitochondria to be created (Figure
4D) and visualized in 3-D from various viewpoints (Figure 4E; Videos 1-2).
253
With these observations, we sought to determine if mitochondrial networks changed in
response to aging. In Figure 5, we show representative orthoslice images of the kidney tissue at
each aging point (Figures 5A-A'), the overlay of the 3-D reconstruction on orthoslice (Figures
5B-B'), and the isolated 3-D reconstruction (Figures 5C-C'), with each color representing an
independent mitochondrion. We found that the median mitochondrial area did not significantly
change between 3-month (mean 8.29 μm^2 ± 10.1 μm^2 SD) versus 2-year (6.46 µm ± 5.31 µm
SD) cohorts, unlike our TEM findings, despite great interindividual heterogeneity (Figures 5D-
D'). However, the perimeter decreased between 3-month (mean 14,328 µm ± 17,723 µm) versus
2-year (10,241 µm ± 8,273 µm SD) cohorts, which also showed less intra-individual
heterogeneity, generally (Figures 5E-E'). This trend towards smaller mitochondria was reflected
when we compared volume between 3-month (0.920 μm^3 ± 1.06 μm^3 SD) versus 2-year (0.741
920 μm^3 ± 0.695 μm^3 SD) cohorts (Figures 5F-F'). Compared with our previous results in
cardiac and skeletal muscle, both unaged and aged kidney mitochondria exhibit a relatively
larger size [39]. This also shows that while TEM was able to capture some dynamics of cristae, 3D
images showed that TEM age-related increase in area was incorrect, as kidney mitochondria
become smaller in mice moving from adolescence to adulthood. When mitochondrial
quantifications from each mouse were compared, they showed intergroup heterogeneity,
particularly with one 3-month mice consistently having larger mitochondria, along with
persistent intraindividual variability (Figures 5D'-F').
273
To further elucidate age-related changes and characterize the mitochondrial types in each
age cohort, we used mito-otyping, a method similar to karyotyping, to organize mitochondria
based on their volumes to better visualize the overall mitochondrial diversity (Figure 5G).
Critically, this approach revealed that there were few significant changes in morphology with
only branching showing reductions. In combination, the aged kidney mitochondrial morphology
resembled healthy mitochondria with a reduced size that lacks a phenotype or fragmentation.
Since mitochondria showed a variety of structural changes due to aging, we turned our attention
to the MICOS complex as a potential mechanistic regulator of these age-related changes.

Age-Related Loss of the MICOS Complex Causes Loss of Mitochondrial Structure in HEK293 cells.

Although it is established that the MICOS complex is critical for cristae dynamics [58,59], we have also found that it can modulate overall mitochondrial structure in aging [60], yet it is unclear how aging affects the MICOS complex in aging kidney. Thus, we looked for age-related changes in three core components of the MICOS complex: *Mitofilin* (MIC60), *Chchd3* (MIC19), and *Chchd6* (MIC25), each of which are fundamental to the formation of the MICOS complex and cristae integrity [61–64]. We also looked for changes in *Opa1*, which is epistatic to a *Mitofilin*, a component of the MICOS complex, and is also known to serve as an age-dependent regulator of mitochondrial dynamics [65]. We found that the MICOS complex mRNA expression also decreased in the kidney with age (Figures 6A-D). We found, as expected, that *Opa1* mRNA decreased by over 50% between 3 months and 2 years (Figure 6A). *Mitofilin* also decreased significantly with the largest drop in fold change of any MICOS complex component (Figure 6B). Likewise, *Chchd3* and *Chchd6* mRNA transcripts also decreased with age but slightly less than *Mitofilin* (Figure 6D). Notably, while Opa1 can be epistatic to the MICOS complex, the MICOS complex modulates cristae integrity independently of it, suggesting these changes in qPCR mRNA transcripts represent non-interconnected pathways [66]. To further understand the role of mitochondrial dynamics upon the loss of these MICOS genes, we also studied the impact of losing the MICOS complex.

Since the loss of *Opa1* triggers changes in morphology [65,67], we used it as a positive control for morphological changes. We performed siRNA deletions of *Chchd3*, *Mitofilin*, *Chchd6*, and Opa1 in immortalized human embryonic kidney cells (HEK293 cells). From there, we performed TEM in each of these conditions as compared to a control (Figure 7A-E). As expected, *Opa1* deletion led to significant decreases in mitochondria area, perimeter, and length with an inverse increase in circularity index, which was expected as a result of impaired fusion dynamics (Figures 7F-I). *Chchd3* deletion shows an even more drastic phenotype of reduced mitochondrial area, while *Chchd6* deletion shows a small decrease compared to the control and *Mitofilin* deletion demonstrates no significant differences (Figure 7F). Interestingly, *Chchd3* deletion HEK293 shows a higher perimeter and length despite its decreases in area and circularity index (Figures 7F-I). *Chchd3* KO cells had nearly completely elongated mitochondria, unlike those in *Opa1* deletion. Together this shows that while the phenotype of the MICOS complex KO is distinct from the loss of OPA1, beyond only its canonic roles in cristae integrity [59], it also can modulate mitochondrial structure. Since cristae and mitochondrial dysfunction were a hallmark change in aging kidneys, we sought to understand further functional implications of the age-dependent loss of the MICOS complex.

Knockdown of MIC60 and CHCHD6 Generates Oxidative Stress in HEK293 cells.

As previously reviewed [68], oxidative stress generated across aging can confer susceptibility to kidney pathologies with characteristics that mimic aspects of kidney aging or CKD. While oxidative stress can arise through pleiotropic mechanisms including inflammation and reduced antioxidant activity [36,68,69], free radicals can be generated through the NADPH oxidases and mitochondrial electron transport chain. Thus, we examined if loss of the MICOS complex, conferring alterations in cristae integrity, also results in dysfunction of cellular respiration processes to lead to oxidative stress. To explore this paradigm, we knocked down

Chchd6 and *Mitofilin* since they both have central roles in interacting with the SAM complex to modulate cristae integrity [70]. Intracellular total and mitochondrial ROS levels were assessed by different fluorescent dyes: MitoPY1 with high specificity for H_2O_2 (Figure 7J), MitoBright, for mitochondrial ROS, and DCFDA for generalized intracellular ROS and mitochondrial superoxide production (Figure 7K). Using microscopy-based ROS quantification, we measured increases in mitochondrial ROS levels in both *Mitofilin* and *Chchd6* deletion (Figure 7L). Antimycin A treatment was used as a positive control to validate ROS quantification. Mitofilin knockdown in 293 HEK cells showed significantly increased ROS by all fluorescent dyes, whereas only MitoBright indicating mitochondrial ROS and DCFDA, indicating general intracellular ROS as well as mitochondrial superoxide production, were significantly elevated in CHCHD6 silenced cells (Figures 7N-O). In summary, the suppression of both CHCHD6 and MIC60 resulted in the elevation of ROS, implying the MICOS complex has a role in mitochondrial ROS homeostasis.

Knockdown of MIC60 and CHCHD6 impairs mCa^{2+} handling in HEK293 cells.

Ca^{2+} influences mitochondrial cristae structure [71,72]. To elucidate MICOS core components' (MIC60 and CHCHD6) role in mitochondrial calcium (mCa^{2+}) regulation, we examined mCa^{2+} uptake and retention in HEK293 cells. Both *MITOFILIN* and *CHCHD6* deletion cells show reduced mCa^{2+} uptake (Figure 8A). Furthermore, reduced mCa^{2+} uptake correlated with early permeability transition pore opening (Figure 8B). *MITOFILIN* and *CHCHD6* deletion cells showed a significant reduction in mCa^{2+} retention capacity compared to controls (Figures 7C-D). Furthermore, we confirmed for all of these experiments that siRNA successfully reduced the protein expression of MIC60 and CHCHD6 (Figures 8E-F). These findings indicate MICOS' crucial role in maintaining physiological mCa^{2+} homeostasis, with altered MICOS complex and cristae structure rendering mitochondria susceptible to Ca^{2+} flux dysregulation and mCa^{2+}-induced cell death. Since MERCs are well understood to be modulators of calcium homeostasis dependent on tethering distances [73,74], we qualitatively examined MERCs in 3D structure in aging (Figures 8G-J; Videos 3-4). We observed in aging tissue there was generally a decrease in MERCs with a phenotype of wrappER, a shape that is commonly reported in liver [75]. Since aging confers a loss of MERCs, this suggests impaired Ca^{2+}, paralleling the dysfunction that arises with the loss of the MICOS complex. Together, these results suggested the roles of mitochondria and MICOS in aging kidneys. To further investigate the pathways that regulate mitochondrial changes, metabolomics, and lipidomic profiling was studied in the kidney.

Global metabolic and lipidomic profiling highlights dynamic changes in the aged kidney.

Following our observations of dysregulation of mitochondrial structure in aging, we sought to better understand other age-related metabolic regulators that may be associated with MICOS-mediated changes in ROS production and Ca^{2+} flux. In aging samples, we conducted comprehensive metabolomic and lipidomic profiling to understand the consequences of these morphological imbalances. The metabolomics analysis unveiled disruptions in amino acid biosynthesis, altered nucleotide metabolism, and dysregulated redox signaling in aging kidney tissues (Figures 9A-B; D-H; SFigure 2). Amino acids play a pivotal role in kidney mitochondrial function, contributing to energy production, gluconeogenesis, nitrogen metabolism, protein

synthesis, antioxidant defense, and specialized metabolic pathways within mitochondria [76].
Consequently, the altered shape and size of mitochondria with age accompany disruption to
normal amino acid metabolism, prompting an exploration into whether these changes are
causative or consequential in the aging process. A notable finding was the significant impact on
glycine, serine, and threonine metabolism in aged kidneys (Figures 9B; D-E). These amino acids
are intricately linked to mitochondrial function, contributing to one-carbon metabolism,
providing substrates for energy production, and maintaining cellular homeostasis. Our findings
suggest disruptions to the mitochondrial glycine cleavage system (GCS), influencing the
synthesis of purines, pyrimidines, and other small molecules (Figure 9E). Additionally, threonine
catabolism, contributing to acetyl-CoA production for energy through oxidative phosphorylation
in the mitochondria, showed significant decreases, suggesting that altered mitochondrial
morphology may disrupt energy metabolism and cellular homeostasis (Figure 9D). A noteworthy
decrease in valine, leucine, and isoleucine in aged kidney tissues (Figures 9F-H) raised interest.
Although the biosynthesis of these branched-chain amino acids occurs in the cytoplasm, they
play crucial roles in regulating mitochondrial biogenesis, autophagy, and cellular signaling.
Significant depletions in tissue NAD^+, NADP, NAM, and an increase in NADH pools were also
detected, indicating an age-related imbalance of cofactors (Figures 9I-L). Together, this
underscores disruptions to these metabolites could be key contributors to declining kidney health
and function with age due to morphological changes.
393
Lipidomic profiling of young and aged kidney tissues unveiled age-related changes in
both lipid classes and lipid chain lengths (Figures 10A-C). In aging kidneys, significant
alterations were observed in the triglycerides oligomers (TGO), triglycerides (TG), sterols (ST),
N-acylethanolamines (NAE), lyso-phosphatidyl-inositol (LPI), dihexoylceramides (Hex2Cer),
dilysocariolipin (DLCL), and cardiolipin (CL) lipid classes when compared to other lipid groups
(Figures 10C). We also noted significant differences in lipid chain lengths with age in the
kidney, which impacts membrane integrity, fluidity, and functionality (Figures 10C). Thus, our
lipidomic profiling revealed disruptions in lipid classes that contribute to both energy
metabolism and the maintenance of mitochondrial membrane structure, integrity, and function.
This study also sheds light on novel roles for lipid classes such as NAEs and LPIs in the context
of kidney aging.
405
406
407 **Discussion:**
408
Structural Analysis
In the past, numerous studies have looked at kidneys across aging or disease states via
TEM, which provides high-resolution 2D images [29,77,78]. While TEM is a useful technique to
understand changes in cristae integrity, it cannot accurately capture many structural details of
mitochondria, such as diverse structures mitochondria may adapt to depending on cellular
conditions [79], and we found that TEM area findings directly contradict our findings of decreased
mitochondrial volume with aging. Here we utilized SBF-SEM to perform 3D reconstruction in
aged mouse kidney, which showed many novel phenotypes and losses in mitochondrial volume
that were otherwise not captured by MRI or TEM methods.
We have previously used 3D reconstruction to observe aged skeletal muscle in mouse [39].
We previously observed increased fragmentation which was counteracted by increased

complexity. In aging murine kidneys, mitochondrial complexity did not undergo significant changes with aging. We did note a high rate of diverse or unique mitochondrial shapes, which may in turn confer functional implications [31,79]. In this study, we found a large amount of variation in mitochondrial shape in both 3-month and 2-year aged cohorts, which show a mix of elongated, compact, big volume, small volume, nanotunnels, donut-shaped, and branching structural phenotypes. While our previous studies in skeletal and cardiac tissue showed a dominant phenotype across aging, typically of fragmentation, mitochondria in the kidney undergo different shapes, and 3-month and 2-year samples do not present vastly different phenotypes.

One notable phenotype we observed is mitochondria donuts. We found that the 3-month samples displayed many branched forms of mitochondria with high complexity and formed within them donut-like structures. Past research utilizing 3D reconstruction in the aged brain in monkeys found a high rate of the donut mitochondria phenotype in the aged cohort, which had impaired memory function [80]. Even beyond age, samples with mitochondrial donuts, which resulted in smaller synapses, had worse memory than older cohorts that had normal mitochondria [80]. While it is established that mitochondria donuts are a hallmark of mitochondrial dysfunction [7], interestingly it seems they are differentially expressed in tissue types and correlates to each tissue's functions. It has been suggested that their increased surface area relative to volume allows for them to maintain more organelle contacts at the cost of lower ATP production [79]. It has also been found that, unlike swollen mitochondria, donut mitochondria maintain more of their internal structure and, potentially as a result, are not the target of mitophagy [81]. Since our analysis shows that more donuts occur in the younger samples, these may represent a positive phenotype in some cases. Therefore, the exact roles of donut mitochondria still remain unclear and may extend beyond what has been previously hypothesized.

Beyond changes in relative bioenergetics between mitochondrial shapes, their roles in calcium homeostasis and other biomolecular pathways deserve further research. For instance, past studies have also suggested decreasing activity of the Akt pathway, which is upstream from the mTORC pathway, as a mechanism to restore autophagy, clear out defective mitochondria, and restore biogenetics [82]. However, given that other studies have shown that donut mitochondria may not be subject to autophagic clearing of mitochondria [81], suggesting a potential reason for different relative rates of mitochondrial donuts. Additionally, research of mitochondrial aging in the kidney has found that mtDNA is more error-prone across aging, with up to a 5-fold increase in the number of point mutations and deletions [82], which may also be responsible for the alterations in the mitochondria structure we observe. Thus, the exact molecular underpinnings of these various shapes are deserving of further research.

Finally, we also found differences in the structure of mitochondria, depending on their location. Through 3D reconstruction, we found that in all the aging samples of the kidney, next to the nucleus, the mitochondria are round. The further away from the nucleus, the more unique mitochondria structures, such as large volume (increased mitochondrial function capacity), small volume, elongated (relatively greater surface area facilitates interaction with the surrounding environment), compact, nano tunnels, and donut-shaped (increased surface area for interaction) are present [31]. The change in shapes likely arises as a result of mitochondrial stress [79,83]. Whereas, the areas that are not undergoing stress, likely present typical and elongated mitochondria. Therefore, it is possible that different areas of kidney undergo stress, potentially linked to stress from filtration, while others are not as susceptible to stress as aging goes. In a previous study, we have found that mitochondria in heart retain their morphology [38]. Therefore,

there may be a similar mechanism that helps retain morphology for intracellular regions, such as
perinuclear kidney mitochondria. Notably, the kidney houses at least 16 types of epithelial cells
[84], and the kidney also has distinct regions including the cortex, medulla, and renal sections,
which serve differing functions [85]. However, our study did not permit the differentiation of these
separate regions. Thus, future studies may consider using methods such as SDS-PAGE to further
differentiate kidney samples [86]. Thus, future studies may further explore this by developing ways
to better separate epithelial and globular areas of kidney for SBF-SEM and seeing if there are
region- or area-dependent differences across aging kidney mitochondria.

The MICOS Complex as a Master Regulator

Importantly, in translating the impact of this study to AKI and CKD, mitochondria are
known to be impactors in the pathophysiology of these diseases [87]. Mitochondrial dynamics are
complex and observing key regulators of mitochondrial form, and thus function, may explain
changes that occur in kidney disease states. Key regulators of mitochondria include OPA1
(regulating mitochondrial fusion) and DRP1 (regulating mitochondrial fission), and they may be
responsible for changes observed in the kidney. Past research has shown that in AKI there is a
decrease of OPA1 expression and increased DRP1 expression, suggesting the likely
fragmentation of mitochondria [2]. However, beyond models with decreased expression of DRP1
not being viable, mitochondrial fission is also important to maintain various roles including
microtubule trafficking [88]. Therefore, this study sought to find other targets and changes in
mitochondrial structure beyond simple alterations in fusion and fission, which is often the extent
of what TEM may be able to survey, and the MICOS complex is one such compelling target.
Aging in kidneys is well-established by us and others to cause interstitial fibrosis and
oxidative stress [47,89–91]. Our results suggested age-related loss of the MICOS complex leads to
mitochondrial structural loss, generating oxidative stress and dysregulating calcium homeostasis.
Since the MICOS complex forms across cristae junctions, understanding of the interdependency
of different MICOS complex proteins is still evolving, but currently, it is understood that some
integral proteins such as MIC60 (*Mitofilin*) regulate the expression of other proteins including
MIC10 and MIC19 [92]. Similarly, MIC60/MIC19 (*Mitofilin/Chchd3*), unlike other MICOS
complex proteins, assemble independently of cardiolipin, with MIC19 being responsible for the
regulation of subcomplex distribution [59]. Past studies of the MICOS complex in the kidney have
been limited, but they generally show that mitochondria-rich regions including the kidney have a
high rate of MIC60 and its isoforms, with a deletion of *Mitofilin* resulting in lethal disruption of
the overall complex [92]. This underscores the central role of *Mitofilin*, relative to other
components of the MICOS complex, with functions that extend beyond cristae and
mitochondrial dynamics to nucleoid distribution, suggesting roles in mtDNA synthesis [93]. This
has been recapitulated by other studies showing that *Mitofilin* depletion decreases mtDNA
transcription, resulting in impaired bioenergetics in the kidney, as previously review [61]. Notably,
we saw a most marked age-related decrease in *Mitofilin*, compared with other components
(Figure 6), yet *Mitofilin* also showed a less drastic mitochondrial phenotype when knocked out
compared with other MICOS complex proteins (Figure 7). While structural analysis of MICOS
complex knockouts is limited by only being in TEM, this underscores the importance of
consideration other roles of the MICOS complex beyond its extensively reviewed and well-
understood role in cristae dynamics and biogenesis [58,94].
The role of the MICOS complex in disease states remains more controversial. Generally,
loss of the MICOS complex has been shown to reduce cardiac ATP levels, thus impairing tissue

integrity [95]. Studies within other tissue types, such as the liver have shown that *Chchd3* depletion results in impaired MERCs to induce fatty liver disease with SLC25A46 involvement [96]. As previously reviewed [97], the MICOS complex has thus been involved in neurodegenerative disorders, metabolic syndromes, cardiac dysfunctions, and muscle pathologies. In the kidney, as previously reviewed, impairment of *Mitofilin* has specifically been implicated in the pathophysiology of mtDNA-renal diseases, diabetic kidney disease, kidney failure, and reperfusion [61]. Interestingly, other studies have suggested a protective mechanism by the loss of the MICOS complex. It is possible that loss of the MICOS complex, despite aberrant cristae structure, is a protective factor across aging by having an unexpected, pronounced lifespan extension in *Podospora anserina* [98]. Specifically, it has been suggested that Miro-MIC60 interactions impair cellular respiration and cause oxidative stress to prevent mitophagy, thus lending increased susceptibility to Parkinson's disease and Friedreich's ataxia [99]. This underscores the need to better understand the impact of MICOS complex loss.

Notably, contrary to other studies showing that Miro-MIC60 interactions cause oxidative stress, we found that deletion of the MICOS complex components, *CHCHD6* and *MITOFILIN*, resulted in mitochondrial and cell oxidative stress. As previously reviewed [61], oxidative stress has occurred upon *MITOFILIN* deletion in some tissue types such as cardiac, but the interlinking of MICOS complex and oxidative stress remains poorly elucidated. Of relevance, within the kidney, oxidative stress mediates age-associated renal cell death and has been linked to numerous pathological conditions, as previously reviewed [68]. Since the loss of the MICOS complex is well-understood to impair bioenergetics and ATP production [60,62], our findings suggest that the closely linked process of free radical generation is also bolstered. MICOS-generated ROS may have various effects; for example, they can reduce NAD+, which we observed in our aged tissue, to cause alterations in glycolysis, TCA cycle, and oxidative phosphorylation, as previously reviewed [100]. As previously reviewed, changes in fuel availability lead to alterations in TCA metabolite levels, with downstream effects in reducing mitochondrial calcium uptake and lowering matrix Ca^{2+} levels, which in turn decreases Ca^{2+}-dependent TCA cycle enzyme activity, including pyruvate dehydrogenase and α-ketoglutarate dehydrogenase, and in some cases induce autophagy as a compensatory mechanism for changes in substrate availability [101]. Since we saw concomitantly mCa^{2+} uptake is also decreased with silencing of the MICOS complex, this suggests a vicious cycle through which ROS-dependent NAD+ and calcium-dependent TCA metabolites are lost due to the MICOS complex; however, this pathway needs to be more thoroughly explored. Alternatively, oxidative stress can cause mitochondrial permeability transition pore (mPTP) openings, which adaptively release excess ROS to maintain mitochondrial homeostasis but in pathological permeant conditions can engage in destructive ROS-dependent ROS release [102,103]. While mPTP openings can be transient, calcium-dependent lowering of membrane potential can also cause permanent openings which confer increased risk to apoptotic pathways [104], suggesting an alternative pathway through which a feedback loop may arise due to ROS generation and calcium dysregulation following silencing of proteins involved in the MICOS complex.

In murine renal tubular epithelial cells, an MCU-dependent increase in mitochondrial calcium accumulation results in oxidative stress and, ultimately, senescence [105]. This study underscores the importance of further explicating the role of MICOS in senescence and the therapies that target senescence. For example, a recent study elucidated that diminished expression of GLIS1 in age-related kidney aging models correlates with impaired mitochondrial quality control mechanisms, while increased GLIS1 interaction with PGC1-α helps maintain

mitochondrial stability, thereby suggesting GLIS1 as a potential therapeutic target for mitigating
cell senescence and age-related renal fibrosis [106]. Furthermore, the roles of the MICOS complex
in affecting calcium highlights the importance of investigation of other regulators of
mitochondrial Ca2+ uniporter (e.g., MICU1, MCU, EMRE), some of which have recently been
identified to have roles in cristae morphology [72]. While MICU1 has increasingly been shown to
have a role in cristae morphology [71,72], the interconnectedness of these proteins has not yet been
studied in the context of downstream effectors of the MICOS complex.

Notably, past studies have shown that nicotinamide adenine dinucleotide phosphate
reduced oxidase (NOX) and its oxidative stress contributes to ER stress-induced apoptosis,
contributing to renal dysfunction [107]. Similarly, NOXs have been indicated to play a role in acute
kidney injury by promoting oxidative stress [35,108]. Beyond underscoring the therapeutic potential
of NOXs, their interdependence with ER stress also highlights the importance of studying
MERCs further. MERCs, contact sites under 50 nm which can be caused by ER stress, have
previously been associated with calcium signaling and lipid metabolism, but recent research has
further suggested a potential role in senescence [109]. Here, we did not comprehensively study
MERCs, which are known to be implicated with the calcium homeostasis with which
mitochondria engage in [110]. However, a qualitative analysis did show that wrappER forms
principally in young samples (Figures 8G-H). Past studies show that rough endoplasmic
reticulum may curve to closely wrapped around the mitochondria and maintain lipid
homeostasis, which has been termed wrappER [111]. Thus, the lipidomic shifts we observed with
aging may be possibly caused by deficient lipid flux and impaired cristae structure without
wrappER. This compartment, which has numerous functions including fatty acid secretion, may
be an organelle linking mitochondria and peroxisomes for overall lipid balance regulation [112].
Given calcium homeostasis dysfunction is a potential avenue of kidney disease [2], it remains
important to consider in the future how calcium homeostasis is impacted across aging through
MERC modulation, especially as qualitatively, less MERCs appear to exist in an aged sample.

Metabolic and Lipidomic Changes
Notably, across aging we saw changes in metabolism, which have been known to arise in
part due to oxidative stress generated by the MICOS complex [refs]. Oxidative stress triggers the
activation of poly(ADP-ribose) polymerase (PARP) as a DNA repair mechanism, which in turn
consumes nicotinamide adenine dinucleotide (NAD+), thus NAD+ levels decreased with age in
both males and females [113]. Of relevance, extracellular NAD+ triggers a pathway involving
cAMP to cause an influx of influx of extracellular Ca^{2+} and subsequent superoxide and nitric
oxide generation [114]. A decrease in NAD+ levels has been implicated with disruption of
mitochondrial homeostasis and function to lead to diabetic kidney injury [115]. Studies using
nicotinamide riboside to boost NAD+ levels have shown that it may actually impair inter-
organelle communication and not restore cristae dynamics [116]. This suggests that loss of the
MICOS complex may partially result in a loss of NAD+ levels which may not be easily restored.

Our metabolomic data shows impairment of Pentose Phosphate Pathway (PPP); impaired
glutamine conversion to glutamate, which is required for alpha-ketoglutarate (TCA) and
glutathione (GSH) biosynthesis. Elevated glutamine also activates the mTOR pathway, which
has been suggested to be activated to modulate mitochondria biogenesis in the kidney and is
known to be activated in the aging process[117,118]. PPP is essential to supply NADPH, which is
required for the endogenous glutathione antioxidant system. NAPDH is a central co-factor of
lipid and redox homeostasis [119]. PPP is also essential to supply nucleosides and nucleotides;

Depletion of nucleotides, such as Xanthine and Ribose, promotes genomic instability[120].
Additionally, our results align with existing literature on dysregulated NAD^+ metabolism in
aging kidneys[121]. Significant depletions in tissue NAD^+, NADP, NAM, and an increase in
NADH pools were detected, indicating an age-related imbalance of cofactors (Figures 8I-L). The
decline in NAD+ is linked to a redox shift hypothesis, where more NAD+ is converted to NADH
without adequate reduction back to NAD^{+}[122]. Our data supports this hypothesis, highlighting the
importance of maintaining an adequate pool of mitochondrial NAD^+ for optimal function.
Further supporting a disruption in redox balance was the detection of decreased FAD (flavin
adenine dinucleotide) with age in mouse kidneys (Figure 8M). Similar to NAD(H), FAD is
involved in various energy production pathways, including oxidative phosphorylation, the TCA
cycle, beta-oxidation of fatty acids, and the electron transport chain. In summary, our
metabolomic profiling provides evidence of disruptions to energy metabolism with age,
potentially acting as either a consequence or precursor to the observed morphological phenotypes
we observed in aged kidney tissues.
Our lipidomics profiling showed profound changes in lipid classes. These lipid classes
play diverse and intricate roles in kidney mitochondria, participating in key cellular processes.
TGOs and TGs serve as energy substrates and act as a storage form of fatty acids in kidney
mitochondria [123,124]. Disruptions to these lipids can impact mitochondrial beta-oxidation,
providing acetyl-CoA for energy production. Sterols contribute to the structural integrity and
fluidity of mitochondrial membranes, aligning with our data and observations [125,126]. Maintaining
proper membrane structure is crucial for mitochondrial function, a feature lost with age in kidney
tissues. The role of NAEs in endocannabinoid signaling is recognized [127,128], but their specific
function in kidney mitochondria remains unclear. Notably, changes in the NAE lipid class with
age in the kidney are of special interest. Additionally, while LPIs are known signaling molecules,
their exact impact on mitochondrial function in the kidney requires further investigation. Here
we present two novel cases of lipid classes playing a role with age in the kidney. Hex2Cer
contributes to the composition of mitochondrial membranes [129,130], supporting our observation
that changes in shape and size contribute to aging in the kidney. Lastly, cardiolipins (DLCL and
CL), unique phospholipids predominantly found in the inner mitochondrial membrane, are
crucial for maintaining mitochondrial membrane integrity, cristae structure, and the function of
respiratory chain complexes [131,132]. Changes in CL content with age support the observed decline
in mitochondrial stability and function of our samples.
636
Conclusion

Together, our results underscore that the aging of murine kidney tissue causes cristae
disarray and impaired mitochondrial structure with smaller volumes. This happens alongside
wide-spread metabolic and lipidomic shifts, as well as increased fibrosis and oxidative stress,
which cumulatively confer decreased oxidative capacity and increased risk of age-related disease
states including CKD and AKI [40]. We further found that the MICOS complex is lost with kidney
aging, absent of changes of other common regulators of mitochondria, and cristae morphology.
While the age-dependent loss of the MICOS complex likely accounts for the loss of cristae
architecture, silencing of some components of the MICOS complex in HEK cells confers a
similar structure to that of aged tissue. The MICOS complex silencing further causes both
oxidative stress, reflective of aged states, and reduced mitochondrial calcium uptake. It is
plausible that these changes together create a vicious cycle: MICOS loss drives oxidative stress,
leading to calcium-dependent TCA dysregulation and NAD^+ dysregulation, driving more

oxidative stress and mtDNA loss, leading to a reduction in MICOS complex transcripts, causing
dysfunctional mitochondrial, producing more oxidative stress byproducts, leading to age-
dependent disease states.

EXPERIMENTAL PROCEDURES
Animal Care and Maintenance
Per protocols previously described [38,60,133], the care and maintenance of the male C57BL/6J mice
conformed to the National Institute of Health's guidelines for the use of laboratory animals. The
University of Iowa's Institutional Animal Care and Use Committee (IACUC) or University of
Washington IACUC approved the housing and feeding of these mice. Anesthesia was achieved
using a mixture of 5% isoflurane and 95% oxygen.

Human Sample Cohort:

All human samples were obtained from Brazilian cohorts according to the CAEE (Ethics
Appreciation Presentation Certificate) guidelines. Samples from young individuals were
collected, and experiments were performed under CAEE number 61743916.9.0000.5281;
samples from older individuals were collected under CAEE number 10429819.6.0000.5285.

Immunohistochemistry

Young (4–5 months old) and old (21–23 months old) C57BL/6J were maintained on a chow diet.
Kidney slices were embedded in OCT, with section processing as previously described [47].
Nitrotyrosine (MilliporeSigma, #06□284,1:1000), and anti-mouse IgG-HRP (Abcam; ab97046)
staining was performed to measure oxidative stress. Masson trichrome staining was quantified
with representative images at low power from each kidney section deconvoluted and thresholded
to calculate the blue area relative to the total tissue area in ImageJ.

RNA Extraction and RT-qPCR
Using TRIzol reagent (Invitrogen), total RNA was isolated from tissues and further purified with
the RNeasy kit (Qiagen Inc). RNA concentration was determined by measuring absorbance at
260 nm and 280 nm using a NanoDrop 1000 spectrophotometer (NanoDrop products,
Wilmington, DE, USA). Approximately 1 µg of RNA was reverse-transcribed using a High-
Capacity cDNA Reverse Transcription Kit (Applied Biosciences, Carlsbad CA). Quantitative
PCR (qPCR) was then performed using SYBR Green (Life Technologies, Carlsbad, CA) [16]. For
qPCR, 50 ng of cDNA was loaded into each well of a 384-well plate, with the reaction carried
out on an ABI Prism 7900HT system (Applied Biosystems) with the following cycle: 1 cycle at
95°C for 10 min; 40 cycles of 95°C for 15 s; 59°C for 15 s, 72°C for 30 s, and 78°C for 10 s; 1
cycle of 95°C for 15 s; 1 cycle of 60°C for 15 s; and one cycle of 95°C for 15 s. GAPDH
normalization was used to present the data as fold changes. qPCR primers used were from
previously published sequences [133], as detailed in Table 1.

Table 1: qPCR Primers Used

Gene	Primers	
Opa-1	Forward	5'-ACCAGGAGACTGTGTCAA-3'
	Reverse	5'-TCTTCAAATAAACGCAGAGGTG-3'
Chchd3	Forward	5'-GAAAAGAATCCAGGCCCTTCCACGCGC-3'
	Reverse	5'-CAGTGCCTAGCACTTGGCACAACCAGGAA-3'
Chchd6	Forward	5'-CTCAGCATGGACCTGGTAGGCACTGGGC-3'
	Reverse	5'-GCCTCAATTCCCACATGGAGAAAGTGGC-3'
Mitofilin	Forward	5'-CCTCCGGCAGTGTTCACCTAGTAACCCCTT-3'
	Reverse	5'-TCGCCCGTCGACCTTCAGCACTGAAAACCTAT-3'

CRISPR-Cas9 Knockouts

All cell types were infected with the following adenoviruses for gene knockouts: control CRISPR/Cas9 (sc-418922), CHCHD6 CRISPR (sc-425817), CHCHD3 CRISPR (sc-425804), and mitofilin CRISPR (sc-429376) (Santa Cruz Biotechnology, California, US), alongside appropriate guide RNAs (Table 2). The CRISPR mixture, prepared with 2.5% CRISPR/Cas9 and 2.5% RNAiMax (ThermoFisher Scientific; 13778075) in Opti-MEM (Gibco; 31985070), was incubated for 20 minutes at room temperature. Post-incubation, cells were treated with the CRISPR mixture and incubated at 37°C. Medium changes and cell washes were performed, and experiments were conducted 3 and 7 days post-infection.

705   Table 2: Guide RNA and Plasmids Used

Gene Name	Type of Plasmid	CAS Number
Mitofilin	CRISPR/Cas9 KO (m)	sc-429376
CHCHD6	CRISPR/Cas9 KO (m)	sc-425817
CHCHD3	CRISPR/Cas9 KO (m)	sc-425804
Control	CRISPR/Cas9 KO (m)	sc-418922

706
707

Serial Block-Face Scanning Electron Microscope (SBF-SEM) Processing of Mouse Muscle Fibers

SBF-SEM was performed according to previously defined protocols [134–136]. Anesthesia was induced in male mice using 5% isoflurane. Post skin and hair removal, the liver was treated with 2% glutaraldehyde in 100 mM phosphate buffer for 30 minutes, dissected into 1-mm³ cubes, and further fixed in a solution containing 2.5% glutaraldehyde, 1% paraformaldehyde, and 120 mM sodium cacodylate for 1 hour.

Fixation and subsequent steps collected onto formvar-coated slot grids (Pella, Redding CA), stained and imaged as previously described [134–136]. This includes tissue washing with 100 mM cacodylate buffer, incubation in a mixture of 3% potassium ferrocyanide and 2% osmium tetroxide, followed by dehydration in an ascending series of acetone concentrations. The tissues were then embedded in Epoxy Taab 812 hard resin. Sectioning and imaging of sample was performed using a VolumeScope 2 SEM (Thermo Fisher Scientific, Waltham, MA). Conventional TEM analysis was performed on 300–400 serial sections from each sample, following staining and imaging protocols. Subsequently, analyzed, via imaging was performed

under low vacuum/water vapor conditions with a starting energy of 3.0 keV and beam current of
0.10 nA. Sections of 50 nm thickness were cut allowing for imaging at 10 nm × 10 nm × 50 nm
spatial resolution.

LCMS Methods for Metabolomics
Frozen kidney tissues were weighed and ground with liquid nitrogen in a cryomill (Retsch) at 25
Hz for 45 seconds, before extracting tissues 40:40:20 acetonitrile: methanol: water +0.5% FA
+15% NH4HCO3[137] with a volume of 40μL solvent per 1mg of tissue, vortexed for 15 seconds,
and incubated on dry ice for 10 minutes. Kidney tissue samples were then centrifuged at 16,000
g for 30 minutes. The supernatants were transferred to new Eppendorf tubes and then centrifuged
again at 16,000 g for 25 minutes to remove and residual debris before analysis. Extracts were
analyzed within 24 hours by liquid chromatography coupled to a mass spectrometer (LC-MS).
The LC–MS method was based on hydrophilic interaction chromatography (HILIC) coupled to
the Orbitrap Exploris 240 mass spectrometer (Thermo Scientific) [138]. The LC separation was
performed on a XBridge BEH Amide column (2.1 x 150 mm, 3.5 μm particle size, Waters,
Milford, MA). Solvent A is 95%: 5% H2O: acetonitrile with 20 mM ammonium acetate and
20mM ammonium hydroxide, and solvent B is 90%: 10% acetonitrile: H2O with 20 mM
ammonium acetate and 20mM ammonium hydroxide. The gradient was 0 min, 90% B; 2 min,
90% B; 3 min, 75% B; 5 min, 75% B; 6 min, 75% B; 7 min, 75% B; 8 min, 70% B; 9 min, 70%
B; 10 min, 50% B; 12 min, 50% B; 13 min, 25% B; 14min, 25% B; 16 min, 0% B; 18 min, 0%
B; 20 min, 0% B; 21 min, 90% B; 25 min, 90% B. The following parameters were maintained
during the LC analysis: flow rate 150 mL/min, column temperature 25 °C, injection volume 5 µL
and autosampler temperature was 5 °C. For the detection of metabolites, the mass spectrometer
was operated in both negative and positive ion mode. The following parameters were maintained
during the MS analysis: resolution of 180,000 at m/z 200, automatic gain control (AGC) target at
3e6, maximum injection time of 30 ms and scan range of m/z 70-1000. Raw LC/MS data were
converted to mzXML format using the command line "msconvert" utility [139]. Data were
analyzed via the EL-MAVEN software version 12.
751
LCMS Methods for Lipidomic Profiling
Kidney tissues were homogenized using a Retsch CryoMill. The homogenate was mixed with 1
mL of Extraction Buffer containing IPA/H2O/Ethyl Acetate (30:10:60, v/v/v) and Avanti
Lipidomix Internal Standard (diluted 1:1000) (Avanti Polar Lipids, Inc. Alabaster, AL). Samples
were vortexed and transferred to bead mill tubes for homogenization using a VWR Bead Mill at
6000 g for 30 seconds, repeated twice. The samples were then sonicated for 5 minutes and
centrifuged at 15,000 g for 5 minutes at 4°C. The upper phase was transferred to a new tube and
kept at 4°C. To re-extract the tissues, another 1 mL of Extraction Buffer (30:10:60, v/v/v) was
added to the tissue pellet-containing tube. The samples were vortexed, homogenized, sonicated,
and centrifuged as described earlier. The supernatants from both extractions were combined, and
the organic phase was dried under liquid nitrogen gas. The dried samples were reconstituted in
300 µL of Solvent A (IPA/ACN/H2O, 45:35:20, v/v/v). After brief vortexing, the samples were
sonicated for 7 minutes and centrifuged at 15,000 g for 10 minutes at 4°C. The supernatants were
transferred to clean tubes and centrifuged again for 5 minutes at 15,000 g at 4°C to remove any
remaining particulates. For LC-MS lipidomic analysis, 60 µL of the sample extracts were
transferred to mass spectrometry vials. Sample analysis was performed within 36 hours after
extraction using a Vanquish UHPLC system coupled with an Orbitrap Exploris 240™ mass
spectrometer equipped with a H-ESI™ ion source (all Thermo Fisher Scientific). A Waters

(Milford, MA) CSH C18 column (1.0 × 150 mm × 1.7 µm particle size) was used. Solvent A
consisted of ACN:H2O (60:40; v/v) with 10 mM Ammonium formate and 0.1% formic acid,
while solvent B contained IPA:ACN (95:5; v/v) with 10 mM Ammonium formate and 0.1%
formic acid. The mobile phase flow rate was set at 0.11 mL/min, and the column temperature
was maintained at 65 °C. The gradient for solvent B was as follows: 0 min 15% (B), 0–2 min
30% (B), 2–2.5 min 48% (B), 2.5–11 min 82% (B), 11–11.01 min 99% (B), 11.01–12.95 min
99% (B), 12.95–13 min 15% (B), and 13–15 min 15% (B). Ion source spray voltages were set at
4,000 V and 3,000 V in positive and negative mode, respectively. Full scan mass spectrometry
was conducted with a scan range from 200 to 1000 m/z, and AcquireX mode was utilized with a
stepped collision energy of 30% with a 5% spread for fragment ion MS/MS scan.
780
781
Quantification of TEM Micrographs and Parameters Using ImageJ
Samples were fixed in a manner to avoid any bias, per established protocols [136]. Following
preparation, tissue was embedded in 100% Embed 812/Araldite resin with polymerization at 60
°C overnight. After ultrathin sections (90–100 nm) were collected, they were post-stained with
lead citrate and imaged (JEOL 1400+ at 80 kV, equipped with a GatanOrius 832 camera). The
National Institutes of Health (NIH) *ImageJ* software was used for quantification of TEM images,
as described previously [55,140].
789
Segmentation and Quantification of 3D SBF-SEM Images Using Amira
SBF-SEM images were manually segmented in Amira to perform 3D reconstruction, as
described previously [38,39,134,140]. 300-400 slices were used and analyzed by a blind individual.
250 total mitochondria across from 3 mice were collected for each quantification. For 3D
reconstruction of cardiomyocytes, 10 cells and a total of about 200 mitochondria. Quantification
of 3D structures was performed using the Amira software with built-in parameters or previously
described measurements [134].
797
Assessment of ROS levels
HEK293 WT cells (0.2 millions) were plated in 35 mm dishes. The next day, MIC60
(ThermoFisher, 136128) and CHCHD6 (ThermoFisher, 34035) siRNAs were transfected into
HEK293 cells using Lipofectamine RNAiMax (Invitrogen) according to the manufacturing
instructions. After incubation for 30 hrs, cells were co-stained for 30 min at 37°C with two
different dyes for ROS detection: MitoBright ROS Deep Red (10 µM, Dojindo Laboratories) for
mitochondrial superoxide detection, and DCFDA (10 µM, Invitrogen) for intracellular total ROS
detection. Following the incubation with staining dyes, cells were washed thrice with 1X HBSS
and ROS imaging was done using a confocal microscope (FV4000, Olympus Life Science).
807
808 For mitochondrial H_2O_2 imaging, cells were incubated with MitoPY1 (5 µM, Bio-Techne) for 45
809 min at 37°C. Cells were then washed with 1x HBSS and imaged using a confocal microscope
810 (FV4000, Olympus Life Science). ImageJ was used for the quantification of fluorescence
811 intensities.
812
813 *Mitochondrial Intracellular Calcium*
814 Knockdown of MIC60 and CHCHD6 in HEK293 cells.

815 The MIC60 and CHCHD6 siRNAs along with scramble siRNA control were transfected into
816 HEK293 cells using Lipofectamine RNAiMax (Invitrogen) according to the manufacturer's
817 instructions. After incubation of 48 hrs, cells were used for Calcium measurements.
818
Measurement of mitochondrial Ca^{2+} uptake in HEK293 cells.
Mitochondrial Ca^{2+} uptake was assessed using a multi-wavelength excitation dual-wavelength
emission fluorimeter (Delta RAM, Photon Technology Int.) with slight modifications following
the protocol outlined in Tomar et al., 2016 (PMID: 27184846). An equal number of cells
($2.5x10^6$ cells) were uniformly cleansed with Ca^{2+}/Mg^{2+}-free DPBS (GIBCO) and subsequently
permeabilized in 1 mL of intracellular medium (ICM- 120 mM KCl, 10 mM NaCl, 1 mM
KH_2PO_4, 20 mM HEPES-Tris, pH 7.2) containing 20 µg/ml digitonin, 1.5 µM thapsigargin to
inhibit the SERCA pump and 2.5 mM succinate to energize the mitochondria. The loading of
Fura-FF (1 µM) at the 0 s time point facilitated the measurement of mitochondrial Ca^{2+} uptake.
Fluorescence was recorded at the 340- and 380-nm ex/510-nm em, with continuous stirring at
37°C, and at specified time points a bolus of 5 µM Ca^{2+} and the mitochondrial uncoupler FCCP
(10 µM) were introduced in the cell suspension.
831
Assessment of mCa^{2+} retention capacity (CRC).
To assess mCa2+ retention capacity (CRC), 2×106 cells were resuspended in an
intracellular-like medium containing (120 mM KCl, 10 mM NaCl, 1 mM KH2PO4, 20 mM
HEPES-Tris, pH 7.2), 1.5 µM thapsigargin to inhibit SERCA so that the movement of
Ca2+ was solely influenced by mitochondrial uptake, 20-µg/ml digitonin, supplemented with
2.5 µM succinate. All solutions underwent Chelex 100 treatment to eliminate traces of
Ca^{2+} (Sigma). Digitonin-permeabilized cells were loaded with the ratiometric reporters FuraFF
at a concentration of 1 µM. Fluorescence was recorded using spectrofluorometer (Delta RAM,
Photon Technology Int.) at 340- and 380-nm ex/510-nm em. Following baseline recordings, a
repetitive series of Ca^{2+} boluses (5 µM) were introduced at indicated time points. Upon
reaching a steady state recording, a protonophore, 10 µM FCCP, was added to collapse the
$\Delta\psi m$ and release matrix free-Ca^{2+}. The number of Ca^{2+} boluses taken up by cells was counted to
calculate mitochondrial CRC.
845
Data Analysis
GraphPad Prism (La Jolla, CA, USA), was used for all statistical analysis. All experiments
involving SBF-SEM and TEM data had at least three independent experiments. Statistics were
not handled by those conducting the experiments. The black bars represent the standard error of
the mean. For all analysis, one-way ANOVA was performed with tests against each independent
group and significance was assessed using Fisher's protected least significant difference (LSD)
test. *, **, ***, **** were set to show significant difference, denoting $p < 0.05$, $p < 0.01$, $p <$
0.001, and $p < 0.0001$, respectively.
854
855

856 **FUNDING**
857 All authors have no competing interests.
858 This project was funded by the National Institute of Health (NIH) NIDDK T-32, number
859 DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; National
860 Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training

in Molecular Endocrinology to A.C.; NSF MCB #2011577I to S.A.M.; The UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award) from Burroughs Welcome Fund (BWF) ID # 1021868.01, BWF Ad-hoc Award, NIH Small Research Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation to A.H.J.; and National Institutes of Health grant HD090061 and the Department of Veterans Affairs Office of Research Award I01 BX005352 to J.G. Howard Hughes Medical Institute Hanna H. Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to M.R.M); and Burroughs Wellcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to M.R.M). National Institutes of Health Grants: R21DK119879 (to C.R.W.) and R01DK-133698 (to C.R.W.), American Heart Association Grant 16SDG27080009 (to C.R.W.) and by an American Society of Nephrology KidneyCure Transition to Independence Grant (to C.R.W.). Doris Duke Clinical Scientist Development Award grant 2021193, Burroughs Wellcome Fund grant 1021480, K23 HL156759, and R01 DK112262 (CNW). NIH Grants R01HL147818, R03HL155041, and R01HL144941 (A. Kirabo). NIH Grant R00DK120876 (D.T.), Harold S. Geneen Charitable Trust Awards Program (D.T.), Alzheimer's Association AARG-NTF-23-1144888 (D.T.). NIH Grant R00AG065445 (P.J.), Alzheimer's Association 24AARG-D-1191292 (P.J.), Wake ADRC REC and Development grant P30AG072947 (P.J.). American Heart Association Grant 23POST1020344 (A.K.). NIH K01AG062757 to (M.T.S.) Its contents are solely the responsibility of the authors and do not necessarily represent the official view of the NIH. The contents are solely the responsibility of the authors and do not necessarily represent the official view of the NIH. The funders had no role in study design, data collection, and analysis, decision to publish, or preparation of the manuscript. We would also like to acknowledge the Huck Institutes' Metabolomics Core Facility (RRID:SCR_023864) for use of the OE 240 LCMS and Drs. Imhoi Koo, Ashley Shay, and Sergei Koshkin for helpful discussions on sample preparation and analysis.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

References:

1. Ferguson, M. A. & Waikar, S. S. Established and Emerging Markers of Kidney Function. *Clinical Chemistry* **58**, 680–689 (2012).

2. Ishimoto, Y. & Inagi, R. Mitochondria: a therapeutic target in acute kidney injury. *Nephrology Dialysis Transplantation* **31**, 1062–1069 (2016).

3. Chronic Kidney Disease in the United States, 2019. *Fluoride Action Network* https://fluoridealert.org/studytracker/38332/ (2020).

4. Duann, P. & Lin, P.-H. Mitochondria Damage and Kidney Disease. in *Mitochondrial*

Dynamics in Cardiovascular Medicine (ed. Santulli, G.) 529–551 (Springer International

Publishing, Cham, 2017). doi:10.1007/978-3-319-55330-6_27.

5. Bhargava, P. & Schnellmann, R. G. Mitochondrial energetics in the kidney. *Nat Rev Nephrol*

13, 629–646 (2017).

6. Glancy, B. Visualizing mitochondrial form and function within the cell. *Trends in molecular*

medicine **26**, 58–70 (2020).

7. Picard, M. & McEwen, B. S. Mitochondria impact brain function and cognition. *Proceedings*

of the National Academy of Sciences **111**, 7–8 (2014).

8. Duchen, M. R. & Szabadkai, G. Roles of mitochondria in human disease. *Essays in*

Biochemistry **47**, 115–137 (2010).

9. Mao, J. *et al.* The relationship between kidney disease and mitochondria: a bibliometric

study. *Renal Failure* **46**, 2302963 (2024).

10. Parasyri, M. *et al.* Renal Phenotype in Mitochondrial Diseases: A Multicenter Study. *Kidney*

Diseases **8**, 148–159 (2022).

11. Forbes, J. M. & Thorburn, D. R. Mitochondrial dysfunction in diabetic kidney disease. *Nat*

Rev Nephrol **14**, 291–312 (2018).

12. Alway, S. E., Mohamed, J. S. & Myers, M. J. Mitochondria Initiate and Regulate Sarcopenia.

Exerc Sport Sci Rev **45**, 58–69 (2017).

13. Hepple, R. T. Mitochondrial Involvement and Impact in Aging Skeletal Muscle. *Frontiers in*

Aging Neuroscience **6**, (2014).

14. Flannery, P. J. & Trushina, E. Mitochondrial dynamics and transport in Alzheimer's disease.

Molecular and Cellular Neuroscience **98**, 109–120 (2019).

15. Zhang, L. *et al.* Altered brain energetics induces mitochondrial fission arrest in Alzheimer's

Disease. *Scientific reports* **6**, 1–12 (2016).

16. Boudina, S. *et al.* Mitochondrial energetics in the heart in obesity-related diabetes: direct

evidence for increased uncoupled respiration and activation of uncoupling proteins. *Diabetes*

56, 2457–2466 (2007).

17. Friederich, M., Hansell, P. & Palm, F. Diabetes, oxidative stress, nitric oxide and

mitochondria function. *Current diabetes reviews* **5**, 120–144 (2009).

18. Venkatachalam, M. A. & Weinberg, J. M. The tubule pathology of septic acute kidney

injury: a neglected area of research comes of age. *Kidney International* **81**, 338–340 (2012).

19. Padovano, V., Podrini, C., Boletta, A. & Caplan, M. J. Metabolism and mitochondria in

polycystic kidney disease research and therapy. *Nat Rev Nephrol* **14**, 678–687 (2018).

20. Fieni, F., Bae Lee, S., Jan, Y. N. & Kirichok, Y. Activity of the mitochondrial calcium

uniporter varies greatly between tissues. *Nat Commun* **3**, 1317 (2012).

21. Bratic, A. & Larsson, N.-G. The role of mitochondria in aging. *The Journal of clinical*

investigation **123**, 951–957 (2013).

22. Stenvinkel, P. & Larsson, T. E. Chronic Kidney Disease: A Clinical Model of Premature

Aging. *American Journal of Kidney Diseases* **62**, 339–351 (2013).

23. Perico, N., Remuzzi, G. & Benigni, A. Aging and the kidney. *Current Opinion in*

Nephrology and Hypertension **20**, 312–317 (2011).

24. Figueiredo, P. A., Mota, M. P., Appell, H. J. & Duarte, J. A. The role of mitochondria in

aging of skeletal muscle. *Biogerontology* **9**, 67–84 (2008).

25. Lesnefsky, E. J., Chen, Q. & Hoppel, C. L. Mitochondrial Metabolism in Aging Heart. *Circ*

Res **118**, 1593–1611 (2016).

26. Zhang, J. *et al.* Alterations in mitochondrial dynamics with age-related Sirtuin1/Sirtuin3

deficiency impair cardiomyocyte contractility. *Aging Cell* **20**, e13419 (2021).

27. Serviddio, G. *et al.* Bioenergetics in aging: mitochondrial proton leak in aging rat liver,

kidney and heart. *Redox Report* **12**, 91–95 (2007).

28. Yamamoto, T. *et al.* Time-dependent dysregulation of autophagy: Implications in aging and

mitochondrial homeostasis in the kidney proximal tubule. *Autophagy* **12**, 801–813 (2016).

29. Cui, J. *et al.* Age-related changes in the function of autophagy in rat kidneys. *AGE* **34**, 329–

339 (2012).

30. Neikirk, K. *et al.* Call to Action to Properly Utilize Electron Microscopy to Measure

Organelles to Monitor Disease. *European Journal of Cell Biology* 151365 (2023)

doi:10.1016/j.ejcb.2023.151365.

31. Jenkins, B. C. *et al.* Mitochondria in disease: changes in shapes and dynamics. *Trends*

Biochem Sci **49**, 346–360 (2024).

32. Marshall, A. G. *et al.* Serial Block Face-Scanning Electron Microscopy as a Burgeoning

Technology. *Adv Biol (Weinh)* e2300139 (2023) doi:10.1002/adbi.202300139.

33. Courson, J. A. *et al.* Serial Block-Face Scanning Electron Microscopy (SBF-SEM) of

Biological Tissue Samples. *J Vis Exp* 10.3791/62045 (2021) doi:10.3791/62045.

34. Dutta, S. & Sengupta, P. Men and mice: Relating their ages. *Life Sci* **152**, 244–248 (2016).

35. Li, M. S. *et al.* NADPH oxidase-2 mediates zinc deficiency-induced oxidative stress and

kidney damage. *American Journal of Physiology-Cell Physiology* **312**, C47–C55 (2017).

36. Tomsa, A. M., Alexa, A. L., Junie, M. L., Rachisan, A. L. & Ciumarnean, L. Oxidative stress

as a potential target in acute kidney injury. *PeerJ* **7**, e8046 (2019).

37. Genin, E. C. *et al.* CHCHD 10 mutations promote loss of mitochondrial cristae junctions

with impaired mitochondrial genome maintenance and inhibition of apoptosis. *EMBO*

molecular medicine **8**, 58–72 (2016).

38. Vue, Z. *et al.* Three-Dimensional Mitochondria Reconstructions of Murine Cardiac Muscle

Changes in Size Across Aging. *American Journal of Physiology-Heart and Circulatory*

Physiology (2023) doi:10.1152/ajpheart.00202.2023.

39. Vue, Z. *et al.* Mouse Skeletal Muscle Decrease in the MICOS Complex and Altered

Mitochondrial Networks with age. *bioRxiv* 2022–03 (2022).

40. Gyurászová, M., Gurecká, R., Bábíčková, J. & Tóthová, Ľ. Oxidative Stress in the

Pathophysiology of Kidney Disease: Implications for Noninvasive Monitoring and

Identification of Biomarkers. *Oxid Med Cell Longev* **2020**, 5478708 (2020).

41. Hill Gallant, K. M. & Spiegel, D. M. Calcium Balance in Chronic Kidney Disease. *Curr*

Osteoporos Rep **15**, 214–221 (2017).

42. Childs, D. D. *et al.* In-phase signal intensity loss in solid renal masses on dual-echo gradient-

echo MRI: association with malignancy and pathologic classification. *AJR Am J Roentgenol*

203, W421-428 (2014).

43. Outwater, E. K., Bhatia, M., Siegelman, E. S., Burke, M. A. & Mitchell, D. G. Lipid in renal

clear cell carcinoma: detection on opposed-phase gradient-echo MR images. *Radiology* **205**,

103–107 (1997).

44. Fang, Y. *et al.* The ageing kidney: Molecular mechanisms and clinical implications. *Ageing*

Research Reviews **63**, 101151 (2020).

45. Wang, K., Liao, Q. & Chen, X. Research progress on the mechanism of renal interstitial

fibrosis in obstructive nephropathy. *Heliyon* **9**, e18723 (2023).

46. Menn-Josephy, H. *et al.* Renal interstitial fibrosis: an imperfect predictor of kidney disease

progression in some patient cohorts. *Am J Nephrol* **44**, 289–299 (2016).

47. Chu, Y. *et al.* Glutathione peroxidase☐1 overexpression reduces oxidative stress, and

improves pathology and proteome remodeling in the kidneys of old mice. *Aging Cell* **19**,

e13154 (2020).

48. Lv, W., Booz, G. W., Fan, F., Wang, Y. & Roman, R. J. Oxidative Stress and Renal Fibrosis:

Recent Insights for the Development of Novel Therapeutic Strategies. *Front Physiol* **9**, 105

(2018).

49. Walker, L. M. *et al.* Oxidative Stress and Reactive Nitrogen Species Generation during

Renal Ischemia. *Toxicological Sciences* **63**, 143–148 (2001).

50. Qian, J. *et al.* Nitrotyrosine Level Was Associated with Mortality in Patients with Acute

Kidney Injury. *PLOS ONE* **8**, e79962 (2013).

51. Brandt, T. *et al.* Changes of mitochondrial ultrastructure and function during ageing in mice

and Drosophila. *eLife* **6**, e24662.

52. L, G., Nn, van der W., Ij, O., Pj, P. & Am, van der B. Loss of the intermembrane space

protein Mgm1/OPA1 induces swelling and localized constrictions along the lengths of

mitochondria. *The Journal of biological chemistry* **279**, (2004).

53. Cogliati, S., Enriquez, J. A. & Scorrano, L. Mitochondrial cristae: where beauty meets

functionality. *Trends in biochemical sciences* **41**, 261–273 (2016).

54. Eisner, V. *et al.* Mitochondrial fusion dynamics is robust in the heart and depends on calcium

oscillations and contractile activity. *Proc Natl Acad Sci U S A* **114**, E859–E868 (2017).

55. Lam, J. *et al.* A Universal Approach to Analyzing Transmission Electron Microscopy with

ImageJ. *Cells* **10**, 2177 (2021).

56. Crabtree, A. *et al.* Defining Mitochondrial Cristae Morphology Changes Induced by Aging

in Brown Adipose Tissue. *Adv Biol (Weinh)* e2300186 (2023) doi:10.1002/adbi.202300186.

57. McBride, E. L. *et al.* Comparison of 3D cellular imaging techniques based on scanned

electron probes: Serial block face SEM vs. Axial bright-field STEM tomography. *Journal of*

Structural Biology **202**, 216–228 (2018).

58. Anand, R., Reichert, A. S. & Kondadi, A. K. Emerging Roles of the MICOS Complex in

Cristae Dynamics and Biogenesis. *Biology (Basel)* **10**, 600 (2021).

59. Friedman, J. R., Mourier, A., Yamada, J., McCaffery, J. M. & Nunnari, J. MICOS

coordinates with respiratory complexes and lipids to establish mitochondrial inner membrane

architecture. *Elife* **4**, e07739 (2015).

60. Vue, Z. *et al.* 3D reconstruction of murine mitochondria reveals changes in structure during

aging linked to the MICOS complex. *Aging Cell* **22**, e14009 (2023).

61. Feng, Y., Madungwe, N. B. & Bopassa, J. C. Mitochondrial inner membrane protein,

Mic60/mitofilin in mammalian organ protection. *J Cell Physiol* **234**, 3383–3393 (2019).

62. An, J. *et al.* CHCM1/CHCHD6, Novel Mitochondrial Protein Linked to Regulation of

Mitofilin and Mitochondrial Cristae Morphology *. *Journal of Biological Chemistry* **287**,

7411–7426 (2012).

63. Darshi, M. *et al.* ChChd3, an Inner Mitochondrial Membrane Protein, Is Essential for

Maintaining Crista Integrity and Mitochondrial Function *. *Journal of Biological Chemistry*

286, 2918–2932 (2011).

64. Darshi, M. & Taylor, S. S. Mitochondrial ChChD3 acts as a Scaffold for Mitofilin, Sam50

and PKA. (2008).

65. Barrera, M., Koob, S., Dikov, D., Vogel, F. & Reichert, A. S. OPA1 functionally interacts

with MIC60 but is dispensable for crista junction formation. *FEBS Letters* **590**, 3309–3322

(2016).

66. Hu, C. *et al.* OPA1 and MICOS Regulate mitochondrial crista dynamics and formation. *Cell*

Death Dis **11**, 1–17 (2020).

67. Gilkerson, R., De La Torre, P. & St. Vallier, S. Mitochondrial OMA1 and OPA1 as

Gatekeepers of Organellar Structure/Function and Cellular Stress Response. *Frontiers in*

Cell and Developmental Biology **9**, (2021).

68. Percy, C., Pat, B., Poronnik, P. & Gobe, G. Role of oxidative stress in age-associated chronic

kidney pathologies. *Advances in Chronic Kidney Disease* **12**, 78–83 (2005).

69. Stenvinkel, P. *et al.* Chronic Inflammation in Chronic Kidney Disease Progression: Role of

Nrf2. *Kidney Int Rep* **6**, 1775–1787 (2021).

70. Ding, C. *et al.* Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae

structure. *Scientific reports* **5**, 1–11 (2015).

71. Tomar, D. *et al.* MICU1 regulates mitochondrial cristae structure and function independently

of the mitochondrial Ca2+ uniporter channel. *Sci Signal* **16**, eabi8948 (2023).

72. Gottschalk, B. *et al.* MICU1 controls cristae junction and spatially anchors mitochondrial

Ca2+ uniporter complex. *Nat Commun* **10**, 3732 (2019).

73. Hinton, A. *et al.* ATF4 Dependent Increase in Mitochondrial-Endoplasmic Reticulum

Tethering Following OPA1 Deletion in Skeletal Muscle. *Journal of Cellular Physiology*

(2024) doi:10.1101/2022.09.12.507669.

74. Csordás, G. & Hajnóczky, G. SR/ER–mitochondrial local communication: Calcium and

ROS. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1787**, 1352–1362 (2009).

75. Ilacqua, N., Anastasia, I. & Pellegrini, L. Isolation and analysis of fractions enriched in

WrappER-associated mitochondria from mouse liver. *STAR Protocols* **2**, 100752 (2021).

76. Li, Q. & Hoppe, T. Role of amino acid metabolism in mitochondrial homeostasis. *Front Cell*

Dev Biol **11**, 1127618 (2023).

77. Bárcena, C., Martínez, M. A., Ortega, M. P., Muñoz, H. G. & Sárraga, G. U. Mitochondria

with Tubulovesicular Cristae in Renal Oncocytomas. *Ultrastructural Pathology* **34**, 315–320

(2010).

78. O'Toole, J. F., Patel, H. V., Naples, C. J., Fujioka, H. & Hoppel, C. L. Decreased

cytochrome c mediates an age-related decline of oxidative phosphorylation in rat kidney

mitochondria. *Biochemical Journal* **427**, 105–112 (2010).

79. Glancy, B., Kim, Y., Katti, P. & Willingham, T. B. The Functional Impact of Mitochondrial

Structure Across Subcellular Scales. *Frontiers in Physiology* **11**, (2020).

80. Hara, Y. *et al.* Presynaptic mitochondrial morphology in monkey prefrontal cortex correlates

with working memory and is improved with estrogen treatment. *Proceedings of the National*

Academy of Sciences **111**, 486–491 (2014).

81. Zhou, Y. *et al.* Topology-dependent, bifurcated mitochondrial quality control under

starvation. *Autophagy* **16**, 562–574 (2020).

82. Jankauskas, S. S. *et al.* Aged kidney: can we protect it? Autophagy, mitochondria and

mechanisms of ischemic preconditioning. *Cell Cycle* **17**, 1291–1309 (2018).

83. Long, Q. *et al.* Modeling of Mitochondrial Donut Formation. *Biophysical journal* **109**, 892–9

(2015).

84. Balzer, M. S., Rohacs, T. & Susztak, K. How Many Cell Types Are in the Kidney and What

Do They Do? *Annu Rev Physiol* **84**, 507–531 (2022).

85. Agarwal, S. K., Sethi, S. & Dinda, A. K. Basics of kidney biopsy: A nephrologist's

perspective. *Indian J Nephrol* **23**, 243–252 (2013).

86. Samuel, C. S. Determination of Collagen Content, Concentration, and Sub-types in Kidney

Tissue. in *Kidney Research: Experimental Protocols* (eds. Becker, G. J. & Hewitson, T. D.)

223–235 (Humana Press, Totowa, NJ, 2009). doi:10.1007/978-1-59745-352-3_16.

87. Jiang, M. *et al.* Mitochondrial dysfunction and the AKI-to-CKD transition. *American*

Journal of Physiology-Renal Physiology **319**, F1105–F1116 (2020).

88. Bowes, T. & Gupta, R. S. Novel mitochondrial extensions provide evidence for a link

between microtubule-directed movement and mitochondrial fission. *Biochemical and*

Biophysical Research Communications **376**, 40–45 (2008).

89. Gomes, P. *et al.* Aging increases oxidative stress and renal expression of oxidant and

antioxidant enzymes that are associated with an increased trend in systolic blood pressure.

Oxid Med Cell Longev **2**, 138–145 (2009).

90. Zhou, X. J. *et al.* The aging kidney. *Kidney International* **74**, 710–720 (2008).

91. O'Sullivan, E. D., Hughes, J. & Ferenbach, D. A. Renal Aging: Causes and Consequences. *J*

Am Soc Nephrol **28**, 407–420 (2017).

92. Rockfield, S. M. *et al.* Genetic ablation of Immt induces a lethal disruption of the MICOS

complex. *Life Sci Alliance* **7**, e202302329 (2024).

93. Li, H. *et al.* Mic60/Mitofilin determines MICOS assembly essential for mitochondrial

dynamics and mtDNA nucleoid organization. *Cell Death & Differentiation* **23**, 380–392

(2016).

94. Viana, M. P., Levytskyy, R. M., Anand, R., Reichert, A. S. & Khalimonchuk, O. Protease

OMA1 modulates mitochondrial bioenergetics and ultrastructure through dynamic

association with MICOS complex. *iScience* **24**, 102119 (2021).

95. Birker, K. *et al.* Mitochondrial MICOS complex genes, implicated in hypoplastic left heart

syndrome, maintain cardiac contractility and actomyosin integrity. *eLife* **12**, e83385 (2023).

96. Dong, J. *et al.* Mic19 depletion impairs endoplasmic reticulum-mitochondrial contacts and

mitochondrial lipid metabolism and triggers liver disease. *Nat Commun* **15**, 168 (2024).

97. Eramo, M. J., Lisnyak, V., Formosa, L. E. & Ryan, M. T. The 'mitochondrial contact site

and cristae organising system' (MICOS) in health and human disease. *The Journal of*

Biochemistry **167**, 243–255 (2020).

98. Warnsmann, V. *et al.* Disruption of the MICOS complex leads to an aberrant cristae structure

and an unexpected, pronounced lifespan extension in Podospora anserina. *Journal of*

Cellular Biochemistry **123**, 1306–1326 (2022).

99. Li, L. *et al.* A mitochondrial membrane-bridging machinery mediates signal transduction of

intramitochondrial oxidation. *Nat Metab* **3**, 1242–1258 (2021).

100. Yang, Y. & Sauve, A. A. NAD+ metabolism: Bioenergetics, signaling and manipulation

for therapy. *Biochim Biophys Acta* **1864**, 1787–1800 (2016).

101. Tomar, D. & Elrod, J. W. Metabolite regulation of the mitochondrial calcium uniporter

channel. *Cell Calcium* **92**, 102288 (2020).

102. Zorov, D. B., Juhaszova, M. & Sollott, S. J. Mitochondrial Reactive Oxygen Species

(ROS) and ROS-Induced ROS Release. *Physiol Rev* **94**, 909–950 (2014).

103.   Batandier, C., Leverve, X. & Fontaine, E. Opening of the mitochondrial permeability

transition pore induces reactive oxygen species production at the level of the respiratory

chain complex I. *J Biol Chem* **279**, 17197–17204 (2004).

104.   Vianello, A. *et al.* The mitochondrial permeability transition pore (PTP) — An example

of multiple molecular exaptation? *Biochimica et Biophysica Acta (BBA) - Bioenergetics*

1817, 2072–2086 (2012).

105.   Xiong, Y. *et al.* Mitochondrial calcium uniporter promotes kidney aging in mice through

inducing mitochondrial calcium-mediated renal tubular cell senescence. *Acta Pharmacol Sin*

1–14 (2024) doi:10.1038/s41401-024-01298-5.

106.   Xu, L. *et al.* GLIS1 alleviates cell senescence and renal fibrosis through PGC1-α

mediated mitochondrial quality control in kidney aging. *Free Radical Biology and Medicine*

209, 171–184 (2023).

107.   Li, G., Scull, C., Ozcan, L. & Tabas, I. NADPH oxidase links endoplasmic reticulum

stress, oxidative stress, and PKR activation to induce apoptosis. *Journal of Cell Biology* **191**,

1113–1125 (2010).

108.   Jeong, B. Y. *et al.* Oxidative stress caused by activation of NADPH oxidase 4 promotes

contrast-induced acute kidney injury. *PLOS ONE* **13**, e0191034 (2018).

109.   Ziegler, D. V., Martin, N. & Bernard, D. Cellular senescence links mitochondria-ER

contacts and aging. *Commun Biol* **4**, 1–14 (2021).

110.   Carreras-Sureda, A. *et al.* Non-canonical function of IRE1α determines mitochondria-

associated endoplasmic reticulum composition to control calcium transfer and bioenergetics.

Nat Cell Biol **21**, 755–767 (2019).

111.   Anastasia, I. *et al.* Mitochondria-rough-ER contacts in the liver regulate systemic lipid

homeostasis. *Cell Reports* **34**, 108873 (2021).

112.   Ilacqua, N. *et al.* A three-organelle complex made by wrappER contacts with

peroxisomes and mitochondria responds to liver lipid flux changes. *Journal of Cell Science*

135, jcs259091 (2021).

113.   Massudi, H. *et al.* Age-Associated Changes In Oxidative Stress and NAD+ Metabolism

In Human Tissue. *PLOS ONE* **7**, e42357 (2012).

114.   Bruzzone, S. *et al.* Extracellular NAD+ regulates intracellular calcium levels and induces

activation of human granulocytes. *Biochem J* **393**, 697–704 (2006).

115.   Yan, L.-J. NADH/NAD+ Redox Imbalance and Diabetic Kidney Disease. *Biomolecules*

11, 730 (2021).

116.   Lauritzen, K. H. *et al.* Impaired dynamics and function of mitochondria caused by

mtDNA toxicity leads to heart failure. *American Journal of Physiology-Heart and*

Circulatory Physiology **309**, H434–H449 (2015).

117.   Khan, N. A. *et al.* mTORC1 regulates mitochondrial integrated stress response and

mitochondrial myopathy progression. *Cell metabolism* **26**, 419–428 (2017).

118.   Papadopoli, D. *et al.* mTOR as a central regulator of lifespan and aging. *F1000Res* **8**,

F1000 Faculty Rev-998 (2019).

119.   Ilter, D. *et al.* NADK-mediated *de novo* NADP(H) synthesis is a metabolic adaptation

essential for breast cancer metastasis. *Redox Biology* **61**, 102627 (2023).

120.   Bester, A. C. *et al.* Nucleotide Deficiency Promotes Genomic Instability in Early Stages

of Cancer Development. *Cell* **145**, 435–446 (2011).

121.   McReynolds, M. R. *et al.* NAD+ flux is maintained in aged mice despite lower tissue

1169   concentrations. *Cell Syst* **12**, 1160-1172.e4 (2021).

1170   122.   McReynolds, M. R., Chellappa, K. & Baur, J. A. Age-related NAD+ decline. *Exp*

1171   *Gerontol* **134**, 110888 (2020).

1172   123.   Mapuskar, K. A. *et al.* Mitochondrial Oxidative Metabolism: An Emerging Therapeutic

1173   Target to Improve CKD Outcomes. *Biomedicines* **11**, 1573 (2023).

1174   124.   Console, L. *et al.* The Link Between the Mitochondrial Fatty Acid Oxidation

1175   Derangement and Kidney Injury. *Front Physiol* **11**, 794 (2020).

1176   125.   Zheng Koh, D. H. & Saheki, Y. Regulation of Plasma Membrane Sterol Homeostasis by

1177   Nonvesicular Lipid Transport. *Contact (Thousand Oaks)* **4**, 25152564211042451 (2021).

1178   126.   Tian, S., Ohta, A., Horiuchi, H. & Fukuda, R. Oxysterol-binding protein homologs

1179   mediate sterol transport from the endoplasmic reticulum to mitochondria in yeast. *J Biol*

1180   *Chem* **293**, 5636–5648 (2018).

1181   127.   Tsuboi, K., Uyama, T., Okamoto, Y. & Ueda, N. Endocannabinoids and related N-

1182   acylethanolamines: biological activities and metabolism. *Inflamm Regen* **38**, 28 (2018).

1183   128.   Boachie, N. *et al.* Circulating Endocannabinoids and N-Acylethanolamines in Individuals

1184   with Cannabis Use Disorder-Preliminary Findings. *Brain Sci* **13**, 1375 (2023).

1185   129.   Peng, K.-Y. *et al.* Mitochondrial dysfunction-related lipid changes occur in nonalcoholic

1186   fatty liver disease progression. *J Lipid Res* **59**, 1977–1986 (2018).

1187   130.   Schenkel, L. C. & Bakovic, M. Formation and regulation of mitochondrial membranes.

1188   *Int J Cell Biol* **2014**, 709828 (2014).

131. Paradies, G., Paradies, V., Ruggiero, F. M. & Petrosillo, G. Role of Cardiolipin in

Mitochondrial Function and Dynamics in Health and Disease: Molecular and

Pharmacological Aspects. *Cells* **8**, 728 (2019).

132. Chen, W.-W., Chao, Y.-J., Chang, W.-H., Chan, J.-F. & Hsu, Y.-H. H.

Phosphatidylglycerol Incorporates into Cardiolipin to Improve Mitochondrial Activity and

Inhibits Inflammation. *Sci Rep* **8**, 4919 (2018).

133. Pereira, R. O. *et al.* OPA 1 deficiency promotes secretion of FGF 21 from muscle that

prevents obesity and insulin resistance. *The EMBO journal* **36**, 2126–2145 (2017).

134. Garza-Lopez, E. *et al.* Protocols for Generating Surfaces and Measuring 3D Organelle

Morphology Using Amira. *Cells* **11**, 65 (2022).

135. Neikirk, K. *et al.* Systematic Transmission Electron Microscopy-Based Identification and

3D Reconstruction of Cellular Degradation Machinery. *Advanced Biology* **7**, 2200221

(2023).

136. Hinton, A. *et al.* A Comprehensive Approach to Sample Preparation for Electron

Microscopy and the Assessment of Mitochondrial Morphology in Tissue and Cultured Cells.

Adv Biol (Weinh) e2200202 (2023) doi:10.1002/adbi.202200202.

137. Lu, W., Wang, L., Chen, L., Hui, S. & Rabinowitz, J. D. Extraction and Quantitation of

Nicotinamide Adenine Dinucleotide Redox Cofactors. *Antioxidants & Redox Signaling* **28**,

167–179 (2018).

138. Wang, L. *et al.* Peak Annotation and Verification Engine for Untargeted LC–MS

Metabolomics. *Anal. Chem.* **91**, 1838–1846 (2019).

139.   Adusumilli, R. & Mallick, P. Data Conversion with ProteoWizard msConvert. in

Proteomics: Methods and Protocols (eds. Comai, L., Katz, J. E. & Mallick, P.) 339–368

(Springer, New York, NY, 2017). doi:10.1007/978-1-4939-6747-6_23.

140.   Hinton, A. *et al.* A comprehensive approach for artifact-free sample preparation and

assessment of mitochondrial morphology in tissue and cultured cells. *bioRxiv* (2021).

1215
1216

Figure Legend:

Main Figures:
Graphical Abstract: Kidney aging causes a decline in the MICOS complex, concomitant with metabolic, lipidomic, and mitochondrial structural alterations.

Figure 1: Comparative analyses of kidney characteristics in young and old subjects differentiated   by   sex.
Cross-sectional imaging of kidney anatomy data from (A) females under 50 years old (aged 14–42 years old; n = 8), (B) males under 50 years old (aged 14–48 years old; n = 6), (C) females over 60 years old (aged 65–90 years old; n = 9), and (D) males over 60 years old (aged 60–88 years old; n = 11). (E) Measurements of left kidney cross-sectional area, calculated as the product of length and width, in females and (F) males. (G) Measurements of left kidney in-phase, representing when magnetic fields of fat and water molecules are in alignment, in females and (H) males. (I) Measurements of left kidney out-of-phase, representing when magnetic fields of fat and water molecules are not in alignment, in females and (J) males. (K) The ratio of the in-phase measurements to out-of-phase, in females and (L) males. For all panels, error bars indicate SEM, Mann–Whitney tests were used for statistical analysis, and statistical significance is denoted as ns (not significant), $*p < 0.05$, and $**p < 0.01$.

Figure 2: Changes in fibrosis and oxidative damage in kidney of young and old murine samples.
(A) Representative images of Masson trichrome staining of the kidney cortex in young (4–5 months old) and (B) old (21–23 months old) samples. (C) Quantification of blue areas. (D) Immunohistochemistry for nitrotyrosine (brown color) in young (4–5 months old) and (E) old (21–23 months old) samples. (F) Quantification of the staining density of nitrotyrosine. For all panels, dots represent independent samples, error bars indicate SEM, unpaired t-tests were used for statistical analysis, and statistical significance is denoted as ns (not significant), $*p < 0.05$, and $**p < 0.01$.

Figure 3: Changes in kidney tissue mitochondria and cristae across aging revealed in transmission electron microscopy (TEM). (A) Representative transmission electron micrographs for kidney tissue at 3 month and (B) two-year in male mice. Boxes show cristae magnified to enhance the details of cristae. (C) Quantification of key mitochondrial characteristics included mitochondrial area and (D) circularity index, which measures mitochondrial shape. (E) Total mitochondria in regions of interest quantification. (F) For cristae, cristae score (D), a measurement of the quality observed cristae, is shown. (G) Using serial block face scanning electron microscopy, representative images of cristae morphology in three-dimensions are shown in three-month and (H) 2-year samples. Each dot represents one mitochondrion, with a variable number in each condition with mitochondria measurements in 3 months having n of approximately 1050 and in 2 years of having n of approximately 1450. Cristae score has n of 1093. For all panels, error bars indicate SEM, mann–Whitney tests were used for statistical analysis, and significance values indicate $****P \leq 0.0001$ and ns, not significant.

Figure 4: Schematic of reconstructing kidney tissue mitochondria using serial block facing-scanning electron microscopy (SBF-SEM). (A) Schematic depicting removal of the kidney. (B) Following embedded fixation, SBF-SEM allows for ortho-slice alignment. (C) Manual segmentation of ortho slices was performed to ultimately yield (D) 3-dimensional (3-D) reconstructions of mitochondria (E) 3-D reconstruction of individually colored mitochondria from a transverse view and longitudinal view in kidney tissue of different ages.

Figure 5: Changes in kidney tissue mitochondria morphology across aging revealed in serial block facing-scanning electron microscopy (SBF-SEM). (A) Representative SBF-SEM orthogonal (ortho) slice for kidney tissue in three-month and (A') 2-year samples. (B) 3-D reconstructions of mitochondria in male kidney tissues of three-months and 2-years (B') overlaid on ortho slices. (C) 3-D reconstructed and isolated mitochondria for clear visualization from three-months and (C') 2-years. (D) 3-D reconstructions were then quantified by 3-D area of the average mitochondria (E) perimeter of the average mitochondria, and (F), mitochondrial volume in kidney tissue from 3-month and 2-year samples. (D'-F') Individual values of the mitochondrion in each mouse to show intra-animal heterogeneity. (G) Mito-otyping to arrange mitochondria on the basis of their volume to view qualitative differences in morphology. Each dot represents the average of a single mouse with a varied number of mitochondria surveyed within each mouse (n=4, with each sample having 83-251 mitochondria). In total, 3-month samples included 740 mitochondria, while 2-year samples included 962 mitochondria, which were used for statistical analysis. For all panels, error bars indicate SEM, mann–Whitney tests were used for statistical analysis, and significance values indicate $**P \leq 0.01$, $***P \leq 0.001$, and ns, not significant.

Figure 6: Transcription and Western Blotting of *Opa1* and mitochondrial contact site and cristae organizing system (MICOS) genes in aging kidney tissue. (A–D) Quantitative polymerase chain reaction (qPCR) analyzing the gene transcript fold changes of Opa-1 and MICOS across aging: (A) *Opa1* transcripts, (B) *Mitofilin* transcripts, (C) *Chchd3* transcript, and (D) *Chchd6* transcripts. (E) Western Blot of OPA1, mitochondrial dynamic proteins, and MICOS protein expression. For all panels, error bars indicate SEM, and Mann–Whitney tests were used for statistical analysis. Each dot represents an individual qPCR run (n=4). Significance values indicate $***P \leq 0.001$ and $****P \leq 0.0001$. For all western blotting experiments, n = 4.

Figure 7: Loss of mitochondrial contact site and cristae organizing system (MICOS) genes result in mitochondrial structure changes and oxidative stress in the kidney. (A-E) Individual knockout (KO) of *Opa1*, *Mitofilin*, *Chchd3*, and *Chchd6* and representative transmission electron micrographs. (F–H) quantification upon KO state of each MICOS gene and Opa1 (n = 10 cells) was performed in 3-D reconstruction: (F) average single mitochondrion area, (G) average single mitochondrion perimeter, (H) average single mitochondrion circularity index, and (I) average single mitochondrion length across individual MICOS KO. (J) 4′,6-diamidino-2-phenylindole (DAPI) staining, MitoPY1 (5 uM, 45 min at 370 c magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA (*MITOFILIN* KD), and CHCHD6-siRNA (*CHCHD6* KD) transfected permeabilized HEK293 cells. (K) 4′,6-diamidino-2-phenylindole (DAPI) staining, MitoBright Deep Red (10 uM, 30 min at 37^0 c), DCFDA (10 uM, 30 min at 37^0 c, magnification of 60x), and merge channels in scramble-siRNA (control),

MIC60-siRNA (*MITOFILIN* KD), and CHCHD6-siRNA (*CHCHD6* KD) transfected
permeabilized HEK293 cells. (L) Plate reader-based reactive oxygen species (ROS)
quantification. (M) Microscopy-based ROS quantification of MitoPY1 orange, (N) DCFDA, and
(O) MitoSox Deep Red. For all statistical tests, a one-way ANOVA statistical test was performed
with Dunnett's multiple comparisons test. For 3D microscopy, each dot represents a
mitochondrion, with their number varied between control (n=81), *Opa1* KO (n=153), *Chchd3*
KO (n=139), *Chchd6* KO (n=180), and *Mitofilin* KO (n=156). Significance values indicate *P ≤
0.05, **P ≤ 0.01, ***P ≤ 0.001 and ****P ≤ 0.0001; ns, not significant.

1317

Figure 8: mCa^{2+} uptake and calcium retention capacity are reduced in *MITOFILIN*
(MIC60) and *CHCHD6* knockdown HEK293 cells. (A) Representative traces of mitochondrial
calcium uptake in scramble-siRNA (control), MIC60-siRNA (*MITOFILIN* KD), and CHCHD6-
siRNA (*CHCHD6* KD) transfected permeabilized HEK293 cells. (B) Percentage of mCa2+
uptake rate calculated from (C) representative traces of mitochondrial calcium retention capacity
in control, *MITOFILIN* KD, and *CHCHD6* KD HEK293 cells. The number of boluses of calcium
taken up by cells is shown in circles. (D) Percentage change in mitochondrial calcium retention
capacity calculated from representative traces of mitochondrial calcium retention capacity. (E)
Western blot showing siRNA-mediated KD of *CHCHD6*/CHCHD6 in HEK293 cells. (F)
Western blot showing siRNA-mediated KD of *MITOFILIN*/MIC60 in HEK293 cells. (G) Serial
block face scanning electron microscopy obtained representative images of mitochondria
endoplasmic reticulum contact site morphology overlaid on orthoslice and (H) isolated in three
dimensions in three-month and (I-J) 2-year samples. For all statistical tests, one-way ANOVA
statistical test was performed with Dunnett's multiple comparisons test. N=3-5 for all calcium
experiments, as run in triplicates. Significance values indicate **P ≤ 0.01.

1333
1334

Figure 9. Global metabolomic profiling uncovers dysregulated metabolic pathways and lipid
classes with age in kidney tissues. (A) Metabolomics heatmap showing the relative abundance
of the top 25 metabolite hits with age. (B) Metabolic pathway analysis revealing cluster of
metabolites related to signaling networks that are disrupted in the aging kidney. (D-M)
Metabolite pools illustrating the metabolic pathways that are altered with age in the kidney–
Redox/NAD$^+$ Metabolism and Amino Acid Metabolism/Biosynthesis. For each tissue and
metabolite in the heatmaps, the aged samples were normalized to the median of the young
samples and then log2 transformed. Young, n= 6; aged, n= 6. For all panels, error bars indicate
SEM, ** indicates P≤ 0.01; and *P≤ 0.05, calculated with Student's t-test.

1344

Figure 10: Global lipidomic profiling uncovers dysregulated metabolic pathways and lipid
classes with age in kidney tissues. (A) Heatmap showing disrupted and enriched lipid classes
based on comparisons between young and old kidney tissues. (B) Lipid class enrichment (C) and
lipid chain length enrichment based on comparisons between young and old kidney.
Significantly different lipid classes represented in the figures are those with adjusted p-values <
0.05 (note: p-values were adjusted to correct for multiple comparisons using an FDR procedure)
and log fold changes greater than 1 or less than –1. Young, n= 6; aged, n= 6. For all panels, error
bars indicate SEM, ** indicates P≤ 0.01; and *P≤ 0.05, calculated with Student's t-test.

1353
1354 **Supplement:**

Supplementary Figure 1: Comparative analyses of kidney characteristics in young and old participants. (A) Metrics defined in Figure 1 grouped to look at overall age-related differences, regardless of sex. (B) Kidney cross-sectional area (CSA), (C) in-phase measurement, (D) out-of-phase measurement, and (E) in-phase to out-of-phase ratio, differentiated by sex. One-way ANOVA statistical test performed with post hoc Tukey's test. Each dot represents an individual participant (n=6-11); full patient information is available in File 1. For all panels, error bars indicate SEM, significance values indicate *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, and ****$P \leq 0.0001$; ns, not significant.

Supplementary Figure 2: Global metabolomic and lipidomic profiling uncovers dysregulated metabolic pathways and lipid classes and chain lengths with age in kidney tissues. (SA) Metabolomics heatmap showing the relative abundance of metabolites.

Supplementary File 1: Full kidney characteristics, including age, of young and old participants analyzed in Figure 1.

Video 1: 3D reconstruction of mitochondria from various angles in 3-month kidney samples.

Video 2: 3D reconstruction of mitochondria from various angles in 2-year kidney samples.

Video 3: 3D reconstruction of wrappER (ER represented in green) from 3-month kidney samples.

Video 4: 3D reconstruction of MERCs (ER represented in blue) from 2-year kidney samples.

Graphical Abstract:

The MICOS complex regulates cristae morphology

Kidney aging causes a decline in the MICOS complex, concomitant with metabolic, lipidomic, and mitochondrial structural alterations

MICOS Complex

Aging

HEK

MICOS
siRNA

ROS

Ca²⁺

Knockout of the MICOS complex contributes to oxidative stress and calcium dysregulation

Figure 1

Figure 2

Figure 3

Figure 4

A Mouse Kidney

B SBF-SEM 3D imaging
10μm 10μm 50μm
3μm

C
1 2 3 4 5
6 7 8 9 10
11 12 13 14 15
16 17 18 19 20
21 22 23 24 25
0.5μm

D
90°
0.5μm

E

	3-Month	2-Year
Transverse		2μm
Longitudinal		

Figure 5

Figure 6

Figure 7

Figure 8

A

Dg+Tg

Ca²⁺ (5µM)

FCCP

FuraFF ΔR/R₀

Time (sec)

— scr-siRNA
— Mitofilin-siRNA
— Chchd6-siRNA6

B

%ₘCa²⁺ Uptake Rate

Control   CHCHD6 KD   MITOFILIN KD

C

Dg+Tg

Scr-siRNA ⑯

Ca²⁺ (5µM)

FCCP

FuraFF ΔR/R₀

Time (sec)

Dg+Tg

Chchd6-siRNA ⑨

Ca²⁺ (5µM)

FCCP

FuraFF ΔR/R₀

Time (sec)

Dg+Tg

Mitofilin-siRNA ⑩

Ca²⁺ (5µM)

FCCP

FuraFF ΔR/R₀

Time (sec)

D

% Mitochondrial CRC

Control   CHCHD6 KD   MITOFILIN KD

E

Scr-siRNA   CHCHD6-siRNA   kDa

CHCHD6   23

ATP5A   50

F

Scr-siRNA   MIC60-siRNA   kDa

MIC60   90

ATP5A   50

G

H

I

J

Figure 9

Figure 10

A

B

C