Title: The spatial organization of ascending auditory pathway microstructural maturation from infancy through adolescence using a novel fiber tracking approach

Authors: Kirsten M. Lynch[1] Stefanie C. Bodison[2], Ryan P. Cabeen[1,] Arthur W. Toga[1], Courtney C.J. Voelker[3]

Affiliations:
[1]Laboratory of Neuro Imaging (LONI), USC Mark and Mary Stevens Institute for Neuroimaging and Informatics, USC Keck School of Medicine, Los Angeles, CA, USA
[2]Department of Occupational Therapy, College of Public Health and Health Professions, University of Florida, Gainesville, FL, USA
[3]Otology/Neurotology, Pacific Neuroscience Institute, Los Angeles, CA

Corresponding author:
Kirsten M. Lynch
kirsten.lynch@loni.usc.edu
USC Stevens Neuroimaging and Informatics Institute
Keck School of Medicine of USC
University of Southern California
2025 Zonal Ave.
Los Angeles, CA 90033

Abstract:

Auditory perception is established through experience-dependent stimuli exposure during sensitive developmental periods; however, little is known regarding the structural development of the central auditory pathway in humans. The present study characterized the regional developmental trajectories of the ascending auditory pathway from the brainstem to the auditory cortex from infancy through adolescence using a novel diffusion MRI-based tractography approach and along-tract analyses. We used diffusion tensor imaging (DTI) and neurite orientation dispersion and density imaging (NODDI) to quantify the magnitude and timing of auditory pathway microstructural maturation. We found spatially varying patterns of white matter maturation along the length of the tract, with inferior brainstem regions developing earlier than thalamocortical projections and left hemisphere tracts developing earlier than the right. These results help to characterize the processes that give rise to functional auditory processing and may provide a baseline for detecting abnormal development.

Keywords: Auditory pathway, child development, microstructure, tractography, NODDI, white matter

1. Introduction

The auditory system is a fundamental sensory perceptual system that enables the identification and location of sounds in the external environment. Auditory perception is established through exposure to experience-dependent stimuli during critical developmental periods of heightened plasticity in early life (Gabard-Durnam and McLaughlin, 2020) and plays a critical role in cognitive development (Ortiz-Mantilla et al., 2019). Infants born with congenital sensorineural hearing loss face many obstacles in the acquisition of speech later in life (Kral et al., 2019); however, cochlear implantation early in life during sensitive developmental periods can restore auditory function and enable normal acquisition of spoken language (Nicholas and Geers, 2013; Niparko et al., 2010; Robbins et al., 2004). Given the important role of myelinating processes on the closure of critical periods of auditory plasticity (Kalish et al., 2020; Mcgee et al., 2005), characterization of auditory white matter maturation is an important step towards understanding typical and atypical child auditory development, and possibly guiding interventional treatment planning. However, few studies have investigated the neural microstructure of the pathways that facilitate transmission of auditory information to the brain during typical development using neuroimaging techniques.

The ascending auditory pathway conveys spectrotemporal auditory information from the receptors in the inner ear to the primary auditory cortex (Peterson et al., 2021). The neural pathways that support this information transfer are complex and involve polysynaptic relays of decussating fibers through the brainstem, midbrain, thalamus and subcortical white matter (**Figure 1A**). In summary, the vestibulocochlear nerve (cranial nerve VIII) carries auditory information from the organ of Corti within the cochlea to the cochlear nucleus (CN) in the brainstem through the internal auditory canal. Fibers from the CN then bifurcate and synapse in the ispilateral and contralateral superior olivary nuclei (SON) in the trapezoid body of the brainstem where binaural cues essential for auditory localization are processed (Yin and Chan, 1990). Auditory information is then conveyed from the bilateral SON via the lateral lemniscus to the inferior colliculus (IC), which has a tonotopic map with a spatial gradient of preferential frequencies organized from superficial to deep tissue (De Martino et al., 2013). Following segregation of auditory features in the IC, fibers then project to the moderating body of the medial geniculate nucleus (MGN) of the thalamus. Fibers from the MGN then pass through the acoustic radiation and terminate in the primary auditory cortex (A1) located within the superior temporal gyrus, where reciprocal connections with adjacent and contralateral auditory regions are believed to mediate refinement of the incoming auditory information (Westerhausen et al., 2009).

Much of what we know about the development of the human auditory system comes from histological studies in postmortem tissue. Cellular development of the auditory pathway occurs in a step-wise fashion and follows an inferior-to-superior pattern of maturation, where brainstem structures mature earlier than cortical projections (Moore and Linthicum, 2007). Rapid maturation of the cochlea and vestibulocochlear nerve occur prenatally (Moore et al., 1995; Moore and Linthicum, 2001), whereas myelination of the lateral lemniscus pathway in the brainstem and projections from the IC to the MGN are achieved in the perinatal period (Moore et al., 1995; Yakovlev and Lecours, 1967). Myelination of thalamocortical projections to the auditory cortex reach adult-like levels in early childhood, while axonal maturation of intracortical connections in the superficial white matter of the auditory cortex extend past the first decade of life (Moore and Guan, 2001). The auditory pathway undergoes a protracted developmental process that differs markedly from the early maturation observed in other sensory systems (Kinney et al., 1988; Pujol et al., 2006; Yakovlev and Lecours, 1967), presumably to accommodate language acquisition and cognitive development in children. This extended

developmental time course therefore renders the auditory system particularly susceptible to auditory deficits during sensitive windows of enhanced plasticity, as auditory deprivation in infants and young children have more devastating effects on hearing and language acquisition compared to older children and adults (Gabard-Durnam and McLaughlin, 2020). An understanding of the developmental timing of the auditory pathway *in vivo* can therefore provide researchers and clinicians insight into critical shifts in maturational processes to better aid in the diagnosis and treatment of children with acquired or congenital hearing loss.

Diffusion MRI (dMRI) is a non-invasive neuroimaging technique that is sensitive to the displacement patterns of diffusing water molecules and provides unique insight into characteristics of *in vivo* white matter structure (Beaulieu, 2009). Quantitative models derived from dMRI contrasts, such as diffusion tensor imaging (DTI) and neurite orientation dispersion and density imaging (NODDI), provide unique insight into the integrity of white matter microstructure. DTI is a dMRI model sensitive to white matter cellular features that influence diffusion anisotropy, such as axonal orientation dispersion, density, myelination and membrane permeability. While the metrics derived from the DTI model are sensitive to developmental processes, they are not specific and the relative contribution of different cellular features to changes in DTI metrics is unclear. NODDI is a multi-compartment diffusion model that estimates tissue compartments based on the restrictive nature of water diffusion patterns using multi-shell diffusion data (Zhang et al., 2012). The neurite density index, which models the fraction of a voxel occupied by axons, provides a more specific measure of cell density and accounts for more age-related variance during child development compared to the DTI metric fractional anisotropy (FA) (Genc et al., 2017). Therefore, utilization of metrics derived from both DTI and NODDI can provide complementary insight into different developmental processes within the auditory pathway.

The developmental time course of major white matter tract microstructure using DTI and NODDI metrics have been well studied in pediatric populations, which broadly show dynamic and spatially-varying patterns of maturation with age (reviewed in: Lebel and Deoni 2018; Lebel et al. 2019). However, few studies have assessed the microstructural development of the auditory pathway in vivo. In fact, the majority of research on the development of the auditory pathways and their associated cortical functions have been conducted in children with neurodevelopmental conditions, such as autism spectrum disorders (Berman et al., 2016), and varying degrees of hearing loss. Previous studies have shown auditory pathway FA is reduced in young children with congenital sensorineural hearing loss (Wu et al., 2016) and may be predictive of auditory recovery following cochlear implantation (Huang et al., 2015; Wang et al., 2019). However, many of these studies are restricted to tractography of the auditory pathway thalamocortical projections (Berman et al., 2016; Wu et al., 2016) or utilize diffusion characteristics within delineated regions to explore select nuclei of the auditory pathway (Huang et al., 2015; Wang et al., 2019). Given the potential clinical utility of auditory pathway microstructure for neurodevelopmental diagnoses, it is therefore important to understand how these processes emerge in the normative population. However, no studies have yet to use tractography of the auditory pathway from the brainstem to the cortex to characterize white matter microstructural maturation in typical development.

Research into the structural connectivity of the auditory pathway *in vivo* with dMRI, particularly within the brainstem, is complicated by a number of factors. Brainstem anatomy is characterized by decussating fibers and small, thin pathways that run in close proximity to neighboring structures. The high spatial resolution (Ford et al., 2013) and contrast (Deoni et al., 2005; Ford et al., 2013; Nagae-Poetscher et al., 2004) required to resolve the complex organization of auditory brainstem structures are rarely afforded with conventional dMRI.

Furthermore, the brainstem is characterized by a mixture of white matter bundles and subcortical nuclei, and dMRI tractography performance is inhibited by the polysynaptic nature of the auditory pathway (Mori and Tournier, 2013; Wakana et al., 2004). In order to mitigate the inherent difficulties of delineating the auditory pathway from the brainstem to the auditory cortex, we utilize state-of-the-art diffusion analysis techniques, including a high angular resolution diffusion imaging (HARDI) acquisition with a higher order modeling approach to enhance detail to auditory microstructure (Zanin et al., 2019). In particular, we employed multi-compartment modeling techniques including DTI with free-water elimination and NODDI, which are made possible with a multi-shell (or multi-b-value) acquisition, and this provides a way to characterize tissue microstructure properties and reduce partial volume effects that arise from the mixing of distinct tissue compartments. We also use multi-tensor modeling of white matter fibers to improve the accuracy of fiber tracking in areas with crossing fibers, which are especially challenging in the brainstem. Further, to explore developmental differences that correspond to relevant auditory pathway anatomy, we examined microstructure parameters along the auditory pathway by matching pathway trajectories across individuals. While the complexity and fine structure of some parts of the auditory pathway still remain beyond what may be resolved by MRI, these diffusion derived metrics may nevertheless provide a coarse-grain representation of the major features of an individual's auditory pathway, which are useful for establishing population-level growth models across neurodevelopment. To illustrate this point, a recent study by Vos et al., 2015 showed DTI was sufficient to reconstruct the auditory nerve projections in a sample of adults with unilateral hearing loss; however, multi-shell dMRI tractography provided enhanced sensitivity to morphological alterations in the affected auditory nerve over single-shell methods (Cabeen et al., 2018a). Such models can help expand our understanding of the basic neuroscience of the auditory system, and also provide a resource for potential clinical applications in diagnosis, treatment planning, and evaluation of hearing disorders.

The primary goal of this study was to elucidate the magnitude and timing of auditory pathway maturation using advanced computational neuroimaging methods applied in a large neuroimaging dataset of typically developing children (N=105) from infancy through 18 years old. We developed a novel tractography analysis pipeline to characterize the white matter bundles that constitute the ascending auditory pathway from the vestibulocochlear nerve in the brainstem to the auditory cortex. Growth models were applied to quantitative dMRI metrics derived from DTI and NODDI to characterize the magnitude and timing of microstructural features that contribute to white matter maturation of the bilateral brainstem and subcortical auditory pathways. In order to test the hypothesis that auditory pathway maturation proceeds along the inferior-to-superior axis, we use an along-tract analytical approach to characterize spatial microstructural patterns of white matter maturation along the length of each fiber bundle.

2. Methods

2.1 Subjects

Cross-sectional neuroimaging data from 105 unrelated typically developing infants, children and adolescents between 0.1 and 18.8 years of age ($M \pm SD$ = 7.8 ± 4.9 years, 56 female) were included in the present study. Children were scanned at Cincinnati Children's Hospital Medical Center (CCHMC) as a part of the Cincinnati MR Imaging of NeuroDevelopment (C-MIND) study (Holland et al., 2015), which is publicly available at https://nda.nih.gov/edit_collection.html?id=2329. The goals of C-MIND are to characterize normative brain-behavior development and develop standardized pediatric neuroimaging protocols for typically developing children. Typically developing children were recruited through

the CCHMC and were representative of the racial, ethnic and gender composition of the US population. Participants were excluded if they had a history of head trauma, received special education, maintained an average grade below a C- in school, a premature (<37 gestational weeks) or late term (>42 gestational weeks) birth, a birth weight less than the 10th percentile, a chronic illness or serious medical problem, a head circumference less than the 5th or greater than the 95th percentile for age, a body mass index less than the 5th or greater than the 95th percentile for age and sex, a personal or family history of neurologic or psychiatric disease, a full scale IQ less than 75, abnormal MRI findings, or MRI contraindications, such as orthodontic braces or metallic implants. Parent/guardian consent was obtained for all subjects, subject consent was obtained for individuals over 17 years of age, and subject assent was obtained for individuals less than 17 years of age and procedures were approved by the Institutional Review Board of CCHMC.

2.2 Image acquisition

Neuroimaging data was acquired on a 3T Philips Achieva scanner with a 32-channel head coil. For each subject, multi-shell diffusion weighted images (DWI) were collected across 2 scans with the following acquisition parameters: voxel size=2 mm isotropic, acquisition matrix=112x109, field of view (FOV)=224x224x120, flip angle (FA)=90° and 61 noncollinear gradient-encoding directions with 7 non-diffusion weighted (b0) images interspersed throughout each scan and averaged into a single volume. Differences in the acquisition between the 2 scans include: *Scan 1*: b=1000 s/mm^2; repetition time (TR)/echo time (TE)=6614/81 ms; *Scan 2*: b=3000 s/mm^2; TR/TE=8112/104 ms. T1-weighted (T1w) images were used as an anatomical reference and were acquired using a turbo-field echo magnetization-prepared rapid acquisition with gradient echo (MP-RAGE) protocol with the following parameters: voxel size=1 mm isotropic, FOV=256x224x160, FA=8°, TR/TE=8.1/3.7 ms, and inversion time (TI)=939 ms.

2.3 Data preprocessing

Each scan was normalized by its respective b=0 volume to correct for differences in TR/TE. We corrected for motion and eddy-current induced artifacts using affine registration with FSL FLIRT (Jenkinson et al., 2012) to align each DWI to the average baseline and rotate the associated b-vector. Brain extraction was performed using FSL BET (Smith, 2002).

2.4 Diffusion modeling

Diffusion modeling was performed using the Quantitative Imaging Toolkit (QIT) (Cabeen et al., 2018b). The parameters fractional anisotropy (FA), axial diffusivity (AD), radial diffusivity (RD) and mean diffusivity (MD) were obtained from free-water eliminated diffusion tensor imaging (fwe-DTI) (Pasternak et al., 2009) using iterative least squares optimization (Hoy et al., 2014). fwe-DTI has been shown to improve tissue specificity by reducing partial volume effects introduced by extracellular free water (Metzler-Baddeley et al., 2012). The neurite orientation dispersion and density imaging (NODDI) parameter neurite density index (NDI) (Zhang et al., 2012) was calculated using the spherical mean technique (Cabeen et al., 2019). Intra-voxel multi-tensor models to be used for tractography were calculated using FSL BEDPOSTX (Behrens et al., 2007; Jenkinson et al., 2012).

2.5 Tractography of the auditory pathway

We used an atlas-based streamline tractography approach to model the auditory pathway (**Figure 1**). Our modeling approach separated the auditory pathway into two distinct

segments, (1) a lower brainstem pathway that was constrained to connect fibers from the internal auditory canal (IAC) to the inferior colliculus (IC), and (2) an upper pathway constrained to connect the IC to auditory cortex (AC) in Heschl's gyrus of the superior gyrus of the temporal lobe (**Figure 1B**). We chose the split at the IC as it provided an unambiguous point of termination at each sub-pathway, and the while there are indeed intermediate synaptic connections along the auditory pathway (at the cochlear nucleus, the superior olivary complex, and the medial geniculate nucleus), the region boundaries of those intermediate nuclei do not have sufficient T1w or T2w contrast to be clearly distinguished. However, in reconstructing connections between the IAC and IC, we found the pathway trajectory traverses these intermediate structures, as indicated by 7T fMRI atlas data provided by (García-Gomar et al., 2019); hence, we adopted an approach where these intermediate points are accordingly included in the bundle model in a data-driven manner through tractography (**Figure 1C, D**). We manually labeled IAC, IC, and AC regions-of-interest (ROIs) in the IIT template. The AC was defined using the atlas in (Desikan et al., 2006). The deformable registration algorithm in DTI-TK was used to align these ROIs to native space in each individual, and we applied a hybrid tractography algorithm (Cabeen and Toga, 2020) with the multi-tensor models to reconstruct each bundle with the following parameters: a minimum volume fraction of 0.01, maximum turning angle of 65 degrees, five seeds per voxel, a step size of 1.0 mm, and trilinear interpolation of fiber orientations. The hybrid tractography algorithm is an approach inspired by reinforcement learning to better resolve crossing fiber regions. First, probabilistic tractography is performed using the above tracking parameters, then a single dominant fiber orientation is inferred in each voxel from the most likely probabilistic bundle fiber orientation, with spatial regularization using a Markov Random Field. Subsequently deterministic streamline tractography is applied to the inferred dominant fiber orientation field. This process is described in more detail by Cabeen and Toga, 2020, including an evaluation study of the acoustic radiation that shows this provides increased reliability and topographic regularity.

Next, we characterized the microstructure properties of the upper and lower pathways. The average diffusion metric for the lower and upper auditory pathways of the left and right hemispheres was calculated using a core-averaging approach, to avoid partial volume effects due to neighboring tissue types within the brainstem. This was a simple technique similar to skeletonization, in which a centroid curve was identified as the one with minimal total the Hausdorff distance to the other bundle curves. We then computed the average fwe-DTI and NODDI microstructure parameters on this centroid curve.

2.6 Along-tract processing

We further characterized the bundles using an along tract approach, which can reveal more anatomically-specific features than available from whole bundle analysis. This builds on a previous study by Lynch et al., 2020 that the reader may consult for additional detail. Briefly, a single template prototypical curve was extracted from a population-averaged template bundle with distinct equidistant segments spaced every 4 mm. The template prototype curve was deformed to subject space and the cross-sectional bundle vertices were labelled according to its optimal corresponding template segment label. We then summarized each along-tract vertex by the NODDI and DTI metrics using the core-averaging approach described in the previous section. To account for potential misalignment in the bundle matching step, we applied along-tract smoothing using a Laplacian filter with $\lambda = 0.3$ and five iterations.

2.7 Statistical analysis

Statistical analyses were carried out with R version 3.1.2. The relationship between auditory tract diffusion parameters and age were assessed using the stats package (v3.6.2) with the following models: (a) a univariate linear model of the form: metric = B_0 + B_1*age, where B_0 is the y-intercept and B_1 is the aging coefficient or (b) a Brody growth curve implemented with nonlinear least squares regression using for the form: metric = $α$ - ($α$ - $β$)e^{-k*age}, where $α$ is the asymptote, $β$ is the y-intercept, and k is the growth rate. The best fit model for each diffusion parameter with each auditory tract and along-tract vertex was assessed using Bayesian information criteria (BIC). For models best fit with a Brody growth curve, the age at terminal maturation was defined as the age at 90% of the estimated asymptotic value and bootstrap resampling was performed with 10,000 iterations to obtain the standard error around the estimated coefficients (boot v1.3-28). Lateralization in the developmental timing of auditory pathway maturation with diffusion parameters was assessed by computing the laterality index (LI) at each point along the auditory pathway: LI = (X_I - X_C)/(X_I + X_C) where X_I and X_C reflect the terminal age estimates at corresponding points on the ipsilateral and contralateral tract, respectively. The LI ranges from -1 to 1, where positive values denote regions where the terminal age estimate is older in the ipsilateral compared to the contralateral side, while negative values denote regions where the terminal age estimate is younger in the ipsilateral compared to the contralateral side.

3. Results

3.1. Mean fwe-DTI and NODDI metric differences with age

The fwe-DTI metrics AD, RD and MD showed non-linear age-related differences in the bilateral upper and lower auditory pathways that were best expressed using a Brody growth curve (**Table 1**). The left upper auditory pathway also showed an exponential relationship with age using the fwe-DTI parameter FA and NODDI parameter NDI. Age-related variance in FA and NDI were best explained with a linear model for the left lower (FA: B=.004, t(103)=4.10, p<.001; NDI: B=.009, t(103)=6.37, p<.001;), right lower (FA: B=.003, t(103)=3.05, p=.003; NDI: B=.008, t(103)=4.98, p<.001;) and right upper (FA: B=.008, t(103)=10.20, p<.001; NDI: B=.014, t(103)=18.50, p<.001;) auditory pathways. In these pathways, FA and NDI were significantly and positively associated with age. **Figure 2** shows the age-related changes to fwe-DTI and NODDI parameter values averaged across the whole bilateral upper and lower auditory pathways. AD reaches terminal maturation earliest within the right lower (3.15 years), left upper (5.72 years) and right upper (6.54 years) auditory pathways. RD reaches terminal maturation latest within the right lower (5.14 years) and upper (10.89 years) tracts, while NDI reaches terminal maturation latest within the left upper tract (18.28 years). The estimated age at terminal maturation within the left lower tract is approximately equal across parameters with exponential fits (RD: 4.40 years, MD: 4.58 years, AD: 4.83 years). Overall, the left and right lower tracts reach peak parameter values earlier (left: 4.60 years, right: 4.90 years) than the left and right upper tracts (left: 6.18, right: 8.93 years). For MD and RD, the whole left and right lower tracts reached terminal maturation at approximately the same age (RD: LI=-.08, MD: LI=.07), while the right upper tract reached terminal maturation later (RD: LI=-.25, MD: LI=-.20). Conversely, the age at terminal AD maturation was older in the left compared to right lower tract (LI=.21), while no appreciable interhemispheric differences in AD maturational rate were observed in the upper auditory pathway (LI=-.07) (**Table 1**).

3.2. Developmental rate and timing of tract-specific diffusion changes

Along-tract analyses demonstrate the maturational timing for each metric was not uniform (**Figure 3**). In accordance with our whole tract findings, the Brody growth curve provided the

best fit model for all points along the bilateral auditory pathway for AD, RD and MD and the left upper auditory pathway for NDI. While the Brody growth curve provided the best fit model to describe age-related changes to whole left upper auditory tract FA, age-related FA changes along the most superior fibers in the superficial white matter of the left upper auditory tract were best explained with a linear model (**Figure 4**). The regions that reached terminal maturation latest across diffusion parameters include the brainstem corresponding to the inferior colliculus and superficial white matter underlying the auditory cortex (**Figure 3**). For diffusion parameters within the auditory pathway that showed nonlinear age-related changes, the standard error of the growth rate was highest in the most inferior fibers that correspond to the vestibulocochlear projections into the brainstem (**Figure 5**). Similarly, this region also showed the largest standard error for the estimated slope for parameters that were best fit with a linear model (**Figure 6**). **Figure 7** demonstrates the regional lateralization of diffusion parameter maturation within the auditory pathway. Along-tract analyses reveal increased rightward lateralization of terminal age estimates for RD and MD at the level of the inferior colliculus and the superficial white matter near the superior temporal gyrus.

4. Discussion

This paper is the first to characterize the development of the auditory pathway of typically developing children from infancy through adolescence using diffusion neuroimaging. We have identified spatially varying patterns of white matter microstructural development from the brainstem to the subcortical white matter of the auditory cortex that differed across NODDI and fwe-DTI metrics. Our whole- and along-tract findings support previous histological studies that show a general inferior-to-superior pattern of auditory pathway maturation. Furthermore, we found the left auditory pathway reaches peak microstructural maturity earlier than the right hemisphere, which may be an important precursor to the development of left-lateralized language acquisition in development. Together, our findings contribute to a growing body of work that shows white matter development of auditory structures is a dynamic and heterogeneous process.

In the present study, we found AD, RD and MD are associated with rapid and nonlinear age-related decreases from infancy through adolescence, while FA and NDI increase with age and follow a more protracted developmental timeline. While evidence of auditory pathway diffusion changes across development are limited, our findings are in line with the vast majority of cross-sectional and longitudinal studies of global white matter development that show FA increases, while RD and MD decrease with age (Krogsrud et al., 2016; Lebel and Beaulieu, 2011; Lynch et al., 2020; Moura et al., 2016; Pohl et al., 2016; Scherf et al., 2014). Our findings also agree with recent studies that demonstrate positive associations between NDI and age in white matter across childhood (Genc et al., 2017; Lynch et al., 2020; Mah et al., 2017). Previous studies have demonstrated that increased FA accompanied by decreased MD and RD may reflect increased axon caliber, myelination, and axonal density (Barazany et al., 2009; Beaulieu, 2002; Song et al., 2002). While some studies have shown AD decreases with age in white matter (Krogsrud et al., 2016; Lynch et al., 2020; Pohl et al., 2016), the association between AD and age has been shown to be more variable (Moura et al., 2016; Scherf et al., 2014). While the precise mechanisms that contribute to reduced AD in development are not well understood, work in animal models and surgical transections have shown that reduced AD is associated with axonal injury and degradation (Concha et al., 2006; Song et al., 2003), but not myelination (Song et al., 2002). Together, these results suggest that the early age-related decreases of AD, RD and MD may reflect increased axon diameter, as this process reaches adult-like levels by the age of 5 years (Schröder et al., 1988) while the protracted increase in FA and NDI likely reflect myelination that extends well past adolescence (Bartzokis et al., 2010). This is

corroborated by previous lifespan and longitudinal dMRI studies that show white matter development is a nonlinear and regionally varying process that continues into the third decade of life (Lebel et al., 2012; Lebel and Beaulieu, 2011; Yeatman et al., 2012).

There was converging evidence across all diffusion metrics that show an inferior-to-superior spatial maturation pattern within the auditory pathway, where brainstem regions developed earlier than the thalamocortical projections to the primary auditory cortex. Within the lower auditory pathway from the vestibulocochlear nerve to the IC, AD, RD and MD reached adult levels by 5 years of age, with the earliest age estimates observed in the cranial nerves. Furthermore, the lack of a significant relationship between FA and age in the bilateral cochlear nerve suggests that developmental alterations to cochlear nerve FA was complete by the early postnatal period. Histological studies have similarly shown that rapid embryonic myelination of the cochlear nerve occurs prenatally during the second and third trimesters (Moore et al., 1997; Moore and Linthicum, 2001), while maturation of the brainstem pathways continue through the early postnatal period (Moore and Linthicum, 2007). The lower brainstem, which receives primary sensory information from the cochlear nerve, is important for the establishment of auditory perception and is highly conserved across vertebrate species (Lipovsek and Wingate, 2018). Dendritic arborization and myelination of brainstem axons within the lateral lemniscus and trapezoid body commence during the third trimester and reach adult-like levels by 6 to 12 months of age (Moore et al., 1995), which coincides with the first physiological and behavioral responses to external acoustic stimuli (Birnholz and Benacerraf, 1983; Kuhlman et al., 1988). Fibers that project from the IC to the MGN undergo the most rapid maturation during the postnatal period (Yakovlev and Lecours, 1967), which corresponds to the time when phonemic discrimination is observed in infants (Trehub, 1973). In opposition to the available evidence, we found across metrics using dMRI that the auditory pathway that projects through the IC continued to develop well past adolescence. It is unclear what the source of the discrepancy is between our findings and those found with histology, but it is possible that our findings may have been contaminated by the microstructure of neighboring structures given the small size of the IC.

Contrary to the early maturation of the brainstem auditory pathways, we found the upper auditory pathway reached peak maturation between 5 and 11 years of age for AD, RD and MD and between 11 and 18 years for FA and NDI in the left hemisphere. For all diffusion parameters, the late maturation was driven predominantly by portions of the superficial white matter of the acoustic radiation adjacent to the auditory cortex that, for some metrics, has yet to reach adult levels by 19 years of age. Immature thalamocortical projection axons that have restricted terminations in the auditory cortical layers become apparent by 12 months of age (Moore and Guan, 2001) and proceed to myelinate into early childhood. In line with our findings, the most protracted developmental time course for the auditory pathway occurs in the intracortical projections to adjacent auditory regions within superficial white matter, where they reach adult-like levels by late childhood (Moore and Guan, 2001). Due to the increased computational capacity of intra- and inter-cortical connectivity in the auditory cortex, experience-dependent refinement of advanced auditory processing abilities also extend into adolescence (Litovsky, 2015). For example, age-dependent improvements to sound localization (Kaga, 1992), speech-in-noise discrimination (Elliott, 1979) and the identification degraded sounds (Eisenberg et al., 2000; H. et al., 2000) are observed in late childhood. Notably, the protracted structural and functional maturation of the auditory system in concert with multiple sensory and cognitive systems appear to play a crucial role in the development of speech articulation and perception, which continue to improve well into adolescence (Ross et al., 2011; Sutherland et al., 2012).

Our findings demonstrate interhemispheric developmental differences within the auditory pathway, where diffusion microstructure of the left auditory pathway matured earlier than the right. Our whole tract analyses demonstrated age-related NDI and FA changes within the left auditory pathway taper off by late adolescence, while NDI and FA in the right hemisphere show linear growth that continues past the teenage years. Furthermore, the fwe-DTI metrics RD and MD reach mature levels earlier in portions of the left auditory pathway compared to the right, including the IC and the superficial white matter of the auditory radiation. While few neuroimaging studies have explored the lateralization of the auditory pathway development across childhood, a recent study in infants suggests the leftward developmental asymmetry of thalamocortical microstructure may be present as early as 6 months of age (Adibpour et al., 2020). The leftward asymmetry of white matter processes responsible for auditory perception may also confer more efficient and adaptive functional outcomes in the left hemisphere. The auditory cortex preferentially responds to sounds from the contralateral auditory space and dichotic listening paradigms show a predominantly right-ear preference for auditory stimuli, which coincides with more robust functional responsivity in the left auditory cortex (Westerhausen et al., 2014) and this leftward functional asymmetry in the cortex has been observed as early as infancy (Adibpour et al., 2020). Structural and functional lateralization is a phenomenon commonly observed in systems that support the acquisition of language. Numerous studies have also shown hemispheric preference for language functions and the associated microstructural asymmetries of the perisylvian language networks are also present in early life (Budisavljevic et al., 2015; Dubois et al., 2009; Olulade et al., 2020; Reynolds et al., 2019). Because auditory perception is an important module for functional speech perception, it is possible that these structures develop hemispheric specialization in concert with one another.

While the present study represents an important step towards characterizing normative white matter maturation within the auditory pathway, several limitations should be considered. The dMRI acquisition was collected with multiple gradient strengths enable multi-compartment modeling; however, each shell was acquired with a different TR and TE, which may introduce a modeling bias. To overcome this issue, we normalized each diffusion scan with the non-diffusion-weighted volume of the corresponding scan to mitigate the issues introduced with differing acquisition parameters, as used in previous research (Genc et al., 2017; Lynch et al., 2020). Additionally, several diffusion metrics did not reach asymptotic values and appeared to continue maturing past the age range sampled. a broader age range should be considered To improve our estimates of the maturational timing of auditory pathway microstructure, especially because it is known that myelination is decades-long process (Bartzokis et al., 2010). The data available for this study was cross-sectional in nature and therefore age-related trends do not reflect within-subject changes over time. Furthermore, we were unable to compare our results of auditory pathway microstructural development with functional improvements to hearing ability. Therefore, future studies should consider a longitudinal study design with functional auditory tasks to better understand the relationship between the structural and functional components of auditory development. Although, the children are considered typically developing without any clinically noted language acquisition difficulties or documented hearing loss, there were no formal audiograms obtained in the study subjects. Future studies would be enhanced would be to prospectively obtain audiograms and tests of auditory processing to confirm typical developmental processes.

To our knowledge, the present study is the first to characterize the maturational trajectory of the ascending auditory pathway in vivo from the brainstem to the auditory cortex from infancy through adolescence in a large, typically developing cohort. Utilizing along-tract analyses to gain granular insight into spatial maturational patterns of diffusion microstructure, we show brainstem regions develop than subcortical regions, and left hemisphere structures develop earlier than

right. The results from study can be used as a comparative template to distinguish normative and atypical auditory pathway development. Furthermore, the proposed tractography technique will enable other researchers to delineate the fine auditory structures of the brainstem can be used broadly to study children born with abnormal auditory patterns, such as congenital or acquired sensorineural hearing loss.

CRediT Author Statement

Kirsten M. Lynch: Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review and editing, Visualization. **Stefanie C. Bodison**: Conceptualization, Methodology, Writing – original draft preparation, Writing – review & editing. **Ryan P. Cabeen**: Conceptualization, Methodology, Software, Writing – review and editing, Visualization. **Arthur W. Toga:** Conceptualization, Resources, Supervision, Funding acquisition. **Courtney C. J. Voelker**: Conceptualization, Methodology, Writing – original draft preparation, Writing – review & editing, Visualization, Funding acquisition.

Acknowledgements

We acknowledge the contributions of Dr. Laurel Fisher, who brought valuable insight, enthusiasm, and kindness to the project and team, and Caroline O'Driscoll for her assistance with illustrations used in the present study.

Data availability

All data used in this study is available through the Cincinnati MR Imaging of Neurodevelopment (C-MIND) study in the NIMH Data Archive (NDA) (https://nda.nih.gov/edit_collection.html?id=2329). Data processing modules used in the present study with the Quantitative Imaging Toolkit (QIT) can be downloaded at https://cabeen.io/qitwiki/. The tractography workflow used to extract the ascending auditory pathway, including region of interest masks and bundle definitions, can be found here: https://github.com/cabeen/auditory-pathway

Declaration of Interests

Declarations of interest: None

Funding

The image computing resources provided by the Laboratory of Neuro Imaging Resource (LONIR) at USC are supported in part by National Institutes of Health (NIH) National Institute of Biomedical Imaging and Bioengineering (NIBIB) grant P41EB015922. Author RPC is supported in part by grant number 2020-225670 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation. Author KML is supported by the NIH National Institute on Aging (NIA) Institutional Training Grant T32AG058507. Author CCJV was supported by an American Neurotology Society Research Grant (GR1053368). Data collection and sharing for this project was funded by the Cincinnati MR Imaging of Neurodevelopment study (C-MIND), supported by the NIH National Institute of Child Health and Human Development Grant HHSN275200900018C.

References

Adibpour, P., Lebenberg, J., Kabdebon, C., Dehaene-Lambertz, G., Dubois, J., 2020. Anatomo-functional correlates of auditory development in infancy. Dev. Cogn. Neurosci. 42, 100752. doi:10.1016/j.dcn.2019.100752

Barazany, D., Basser, P.J., Assaf, Y., 2009. In vivo measurement of axon diameter distribution in the corpus callosum of rat brain. Brain 132, 1210–1220. doi:10.1093/brain/awp042

Bartzokis, G., Lu, P.H., Tingus, K., Mendez, M.F., Richard, A., Peters, D.G., Oluwadara, B., Barrall, K.A., Finn, J.P., Villablanca, P., Thompson, P.M., Mintz, J., 2010. Lifespan trajectory of myelin integrity and maximum motor speed. Neurobiol. Aging 31, 1554–1562. doi:10.1016/j.neurobiolaging.2008.08.015

Beaulieu, C., 2009. The biological basis of diffusion anisotropy, in: In: Johansen-Berg, H., Behrens, T.E.J. (Eds. . (Ed.), Diffusion MRI: From Quantitative Measurement to In-Vivo Neuroanatomy. Academic Press, pp. 105–129. doi:10.1016/B978-0-12-396460-1.00008-1

Beaulieu, C., 2002. The basis of anisotropic water diffusion in the nervous system - A technical review. NMR Biomed. 15, 435–455. doi:10.1002/nbm.782

Behrens, T.E.J., Berg, H.J., Jbabdi, S., Rushworth, M.F.S., Woolrich, M.W., 2007. Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? Neuroimage 34, 144–155. doi:10.1016/j.neuroimage.2006.09.018

Berman, J.I., Edgar, J.C., Blaskey, L., Kuschner, E.S., Levy, S.E., Ku, M., Dell, J., Roberts, T.P.L., 2016. Multimodal diffusion-MRI and MEG assessment of auditory and language system development in autism spectrum disorder. Front. Neuroanat. 10. doi:10.3389/fnana.2016.00030

Birnholz, J.C., Benacerraf, B.R., 1983. The Development of Human Fetal Hearing. Science (80-.). 222, 516–518.

Budisavljevic, S., Acqua, F.D., Rijsdijk, V., Kane, F., Picchioni, X., Mcguire, P., Toulopoulou, T., Georgiades, A., Kalidindi, S., Kravariti, E., Murray, R.M., Murphy, D.G., Craig, M.C., Catani, M., 2015. Age-Related Differences and Heritability of the Perisylvian Language Networks. J. Neurosci. 35, 12625–12634. doi:10.1523/JNEUROSCI.1255-14.2015

Cabeen, R.P., Kari, E., Volker, C., Marta, F., Kulich, L.M., Shi, Y., Law, M., Toga, A.W., 2018a. Feasibility of Quantitative Diffusion MR Tractography of the Vestibulocochlear Nerve in Children with Unilateral Profound Sensorineural Hearing Loss, in: Proceedings of the International Society for Magnetic Resonance in Medicine (ISMRM).

Cabeen, R.P., Laidlaw, D.H., Toga, A.W., 2018b. Quantitative Imaging Toolkit⃞: Software for Interactive 3D Visualization , Data Exploration , and Computational Analysis of Neuroimaging Datasets, in: ISMRM-ESMRMB Abstracts. pp. 12–14.

Cabeen, R.P., Sepehrband, F., Toga, A.W., 2019. Rapid and accurate NODDI parameter estimation with the spherical mean technique, in: Proc International Society for Magnetic Resonance in Medicine (ISMRM). p. 3363.

Cabeen, R.P., Toga, A.W., 2020. Reinforcement tractography: A hybrid approach for robust

segmentation of complex fiber bundles, in: IEEE 17th International Symposium on Biomedical Imaging (ISBI). pp. 999–1003.

Concha, L., Gross, D.W., Wheatley, B.M., Beaulieu, C., 2006. Diffusion tensor imaging of time-dependent axonal and myelin degradation after corpus callosotomy in epilepsy patients. Neuroimage 32, 1090–1099. doi:10.1016/j.neuroimage.2006.04.187

De Martino, F., Moerel, M., Van De Moortele, P.F., Ugurbil, K., Goebel, R., Yacoub, E., Formisano, E., 2013. Spatial organization of frequency preference and selectivity in the human inferior colliculus. Nat. Commun. 4, 1–8. doi:10.1038/ncomms2379

Deoni, S.C.L., Peters, T.M., Rutt, B.K., 2005. High-resolution T1 and T2 mapping of the brain in a clinically acceptable time with DESPOT1 and DESPOT2. Magn. Reson. Med. 53, 237–241. doi:10.1002/mrm.20314

Desikan, R.S., Ségonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L., Dale, A.M., Maguire, R.P., Hyman, B.T., Albert, M.S., Killiany, R.J., 2006. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage 31, 968–980. doi:10.1016/j.neuroimage.2006.01.021

Dubois, J., Hertz-Pannier, L., Cachia, A., Mangin, J.F., Le Bihan, D., Dehaene-Lambertz, G., 2009. Structural asymmetries in the infant language and sensori-motor networks. Cereb. Cortex 19, 414–423. doi:10.1093/cercor/bhn097

Eisenberg, L.S., Shannon, R. V., Schaefer Martinez, A., Wygonski, J., Boothroyd, A., 2000. Speech recognition with reduced spectral cues as a function of age. J. Acoust. Soc. Am. 107, 2704–2710. doi:10.1121/1.428656

Elliott, L.L., 1979. Performance of children aged 9 to 17 years on a test of speech intelligibility in noise using sentence material with controlled word predictability. J. Acoust. Soc. Am. 66, 651–653. doi:10.1121/1.383691

Ford, A., Colon-Perez, L., Triplett, W.T., Gullett, J.M., Mareci, T.H., FitzGerald, D.B., 2013. Imaging white matter in human brainstem. Front. Hum. Neurosci. 7, 1–11. doi:10.3389/fnhum.2013.00400

Gabard-Durnam, L., McLaughlin, K.A., 2020. Sensitive periods in human development: charting a course for the future. Curr. Opin. Behav. Sci. 36, 120–128. doi:10.1016/j.cobeha.2020.09.003

García-Gomar, M.G., Strong, C., Toschi, N., Singh, K., Rosen, B.R., Wald, L.L., Bianciardi, M., 2019. In vivo probabilistic structural atlas of the inferior and superior colliculi, medial and lateral geniculate nuclei and superior olivary complex in humans based on 7 tesla MRI. Front. Genet. 10, 1–15. doi:10.3389/fnins.2019.00764

Genc, S., Malpas, C.B., Holland, S.K., Beare, R., Silk, T.J., 2017. Neurite density index is sensitive to age related differences in the developing brain. Neuroimage 148, 373–380. doi:10.1016/j.neuroimage.2017.01.023

H., H.D.E., A., W.B., C., H.S., R., M.D., 2000. Age-Related Improvements in Auditory Backward and Simultaneous Masking in 6- to 10-Year-Old Children. J. Speech, Lang. Hear. Res. 43,

1402–1415. doi:10.1044/jslhr.4306.1402

Holland, S., Schmithorst, V.J., Wagner, M., Lee, G., Rajagopal, A., Sroka, M., Felicelli, N., Rupert, A., Clark, K., Toga, A., Freund, L., 2015. The C-MIND project: normative MRI and behavioral data from children from birth to 18 years, in: Proceedings of the 21st Annumal Meeting of the Organization for Human Brain Mapping (OHBM), Honolulu, HA.

Hoy, A.R., Koay, C.G., Kecskemeti, S.R., Alexander, A.L., 2014. Optimization of a free water elimination two-compartment model for diffusion tensor imaging. Neuroimaging 103, 323–333. doi:10.1016/j.neuroimage.2014.09.053.Optimization

Huang, L., Zheng, W., Wu, C., Wei, X., Wu, X., Wang, Y., Zheng, H., 2015. Diffusion tensor imaging of the auditory neural pathway for clinical outcome of cochlear implantation in pediatric congenital sensorineural hearing loss patients. PLoS One 10, 1–9. doi:10.1371/journal.pone.0140643

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL. Neuroimage 62, 782–790. doi:10.1016/j.neuroimage.2011.09.015

Kaga, M., 1992. Development of sound localization. Acta Paediatr Jpn 34, 134–8.

Kalish, B.T., Barkat, T.R., Diel, E.E., Zhang, E.J., Greenberg, M.E., Hensch, T.K., 2020. Single-nucleus RNA sequencing of mouse auditory cortex reveals critical period triggers and brakes. Proc. Natl. Acad. Sci. U. S. A. 117, 11744–11752. doi:10.1073/pnas.2014145117

Kinney, H.C., Brody, B.A., Kloman, A.S., Gilles, F.H., 1988. Sequence of Central Nervous System Myelination in Human Infancy: II. Patterns of Myelination in Autopsied Infants. J. Neuropathol. Exp. Neurol. 47, 217–234. doi:10.1097/00005072-198805000-00003

Kral, A., Dorman, M.F., Wilson, B.S., 2019. Neuronal Development of Hearing and Language: Cochlear Implants and Critical Periods. Annu. Rev. Neurosci. 42, 47–65. doi:10.1146/annurev-neuro-080317-061513

Krogsrud, S.K., Fjell, A.M., Tamnes, C.K., Grydeland, H., Mork, L., Due-Tønnessen, P., Bjørnerud, A., Sampaio-Baptista, C., Andersson, J., Johansen-Berg, H., Walhovd, K.B., 2016. Changes in white matter microstructure in the developing brain-A longitudinal diffusion tensor imaging study of children from 4 to 11years of age. Neuroimage 124, 473–486. doi:10.1016/j.neuroimage.2015.09.017

Kuhlman, K.A., Burns, K.A., Depp, R., Sabbagha, R.E., 1988. Ultrasonic imaging of normal fetal response to external vibratory acoustic stimulation. Am. J. Obstet. Gynecol. 158, 47–51. doi:10.1016/0002-9378(88)90773-9

Lebel, C., Beaulieu, C., 2011. Longitudinal development of human brain wiring continues from childhood into adulthood. J. Neurosci. 31, 10937–47. doi:10.1523/JNEUROSCI.5302-10.2011

Lebel, C., Deoni, S., 2018. The development of brain white matter microstructure. Neuroimage 182, 207–218. doi:10.1016/j.neuroimage.2017.12.097

Lebel, C., Gee, M., Camicioli, R., Wieler, M., Martin, W., Beaulieu, C., 2012. Diffusion tensor

imaging of white matter tract evolution over the lifespan. Neuroimage 60, 340–52. doi:10.1016/j.neuroimage.2011.11.094

Lebel, C., Treit, S., Beaulieu, C., 2019. A review of diffusion MRI of typical white matter development from early childhood to young adulthood. NMR Biomed 32, e3778. doi:10.1002/nbm.3778

Lipovsek, M., Wingate, R.J.T., 2018. Conserved and divergent development of brainstem vestibular and auditory nuclei. Elife 7, 1–25. doi:10.7554/eLife.40232

Litovsky, R., 2015. Development of the auditory system, Handb Clin Neurol. doi:10.1097/00003446-199902000-00008

Lynch, K.M., Cabeen, R.P., Toga, A.W., Clark, K.A., 2020. Magnitude and timing of major white matter tract maturation from infancy through adolescence with NODDI. Neuroimage 212, 116672. doi:10.1016/j.neuroimage.2020.116672

Mah, A., Geeraert, B., Lebel, C., 2017. Detailing neuroanatomical development in late childhood and early adolescence using NODDI. PLoS One 12, 1–16. doi:10.1371/journal.pone.0182340

Mcgee, A.W., Yang, Y., Fischer, Q.S., Daw, N.W., Stephen, M., 2005. Experience-Driven Plasticity of Visual Cortex Limited by Myelin and Nogo Receptor. Science (80-.). 309, 2222–2226.

Metzler-Baddeley, C., O'Sullivan, M.J., Bells, S., Pasternak, O., Jones, D.K., 2012. How and how not to correct for CSF-contamination in diffusion MRI. Neuroimage 59, 1394–1403. doi:10.1016/j.neuroimage.2011.08.043

Moore, J.K., Guan, Y.L., 2001. Cytoarchitectural and axonal maturation in human auditory cortex. JARO - J. Assoc. Res. Otolaryngol. 2, 297–311. doi:10.1007/s101620010052

Moore, J.K., Guan, Y.L., Shi, S.R., 1997. Axogenesis in the human fetal auditory system, demonstrated by neurofilament immunohistochemistry. Anat. Embryol. (Berl). 195, 15–30. doi:10.1007/s004290050021

Moore, J.K., Linthicum, F.H., 2007. The human auditory system: A timeline of development. Int. J. Audiol. 46, 460–478. doi:10.1080/14992020701383019

Moore, J.K., Linthicum, F.H., 2001. Myelination of the human auditory nerve⬜: different time. Ann. Otol. Rhinol. Laryngol. 110, 655–661.

Moore, J.K., Perazzo, L.M., Braun, A., 1995. Time course of axonal myelination in the human brainstem auditory pathway. Hear. Res. 91, 208–209. doi:10.1016/0378-5955(95)00218-9

Mori, S., Tournier, J.-D., 2013. Introduction to Diffusion Tensor Imaging: and Higher Order Models. Academic Press, Oxford, UK.

Moura, L.M., Kempton, M., Barker, G., Salum, G., Gadelha, A., Pan, P.M., Hoexter, M., Del Aquilla, M.A.G., Picon, F.A., Anés, M., Otaduy, M.C.G., Amaro, E., Rohde, L.A., McGuire, P., Bressan, R.A., Sato, J.R., Jackowski, A.P., 2016. Age-effects in white matter using

associated diffusion tensor imaging and magnetization transfer ratio during late childhood and early adolescence. Magn. Reson. Imaging 34, 529–534. doi:10.1016/j.mri.2015.12.021

Nagae-Poetscher, L.M., Jiang, H., Wakana, S., Golay, X., Van Zijl, P.C.M., Mori, S., 2004. High-resolution diffusion tensor imaging of the brain stem at 3 T. Am. J. Neuroradiol. 25, 1325–1330. doi:10.1016/s0513-5117(08)70337-5

Nicholas, J.G., Geers, A.E., 2013. Spoken language benefits of extending cochlear implant candidacy below 12 months of age. Otol. Neurotol. 34, 532–538. doi:10.1097/MAO.0b013e318281e215

Niparko, J.K., Tobey, E.A., Thal, D.J., Eisenberg, L.S., Wang, N.Y., Quittner, A.L., Fink, N.E., 2010. Spoken language development in children following cochlear implantation. JAMA - J. Am. Med. Assoc. 303, 1498–1506. doi:10.1001/jama.2010.451

Olulade, O.A., Seydell-Greenwald, A., Chambers, C.E., Turkeltaub, P.E., Dromerick, A.W., Berl, M.M., Gaillard, W.D., Newport, E.L., 2020. The neural basis of language development: Changes in lateralization over age. Proc. Natl. Acad. Sci. U. S. A. 117, 23477–23483. doi:10.1073/pnas.1905590117

Ortiz-Mantilla, S., Realpe-Bonilla, T., Benasich, A.A., 2019. Early Interactive Acoustic Experience with Non-speech Generalizes to Speech and Confers a Syllabic Processing Advantage at 9 Months. Cereb. Cortex 29, 1789–1801. doi:10.1093/cercor/bhz001

Pasternak, O., Sochen, N., Gur, Y., Intrator, N., Assaf, Y., 2009. Free water elimination and mapping from diffusion MRI. Magn. Reson. Med. 62, 717–730. doi:10.1002/mrm.22055

Peterson, D.C., Reddy, V., Hamel, R.N., 2021. Neuroanatomy, Auditory Pathway. StatPearls Publishing, Treasure Island, FL.

Pohl, K.M., Sullivan, E. V., Rohlfing, T., Chu, W., Kwon, D., Nichols, B.N., Zhang, Y., Brown, S.A., Tapert, S.F., Cummins, K., Thompson, W.K., Brumback, T., Colrain, I.M., Baker, F.C., Prouty, D., De Bellis, M.D., Voyvodic, J.T., Clark, D.B., Schirda, C., Nagel, B.J., Pfefferbaum, A., 2016. Harmonizing DTI measurements across scanners to examine the development of white matter microstructure in 803 adolescents of the NCANDA study. Neuroimage 130, 194–213. doi:10.1016/j.neuroimage.2016.01.061

Pujol, J., Soriano-Mas, C., Ortiz, H., Sebastián-Gallés, N., Losilla, J.M., Deus, J., 2006. Myelination of language-related areas in the developing brain. Neurology 66, 339–43. doi:10.1212/01.wnl.0000201049.66073.8d

Reynolds, J.E., Long, X., Grohs, M.N., Dewey, D., Lebel, C., 2019. Structural and functional asymmetry of the language network emerge in early childhood. Dev. Cogn. Neurosci. 39, 100682. doi:10.1016/j.dcn.2019.100682

Robbins, A.M.C., Koch, D.B., Osberger, M.J., Zimmerman-Phillips, S., Kishon-Rabin, L., 2004. Effect of Age at Cochlear Implantation on Auditory Skill Development in Infants and Toddlers. Arch. Otolaryngol. - Head Neck Surg. 130, 570–574. doi:10.1001/archotol.130.5.570

Ross, L.A., Molholm, S., Blanco, D., Gomez0Rairez, M., Saint-Amour, D., Foxe, J.J., 2011. The

development of multisensory speech perception continues into the late childhood years. Eir J Neurosci 33, 2329–2337. doi:10.1111/j.1460-9568.2011.07685.x.The

Scherf, K.S., Thomas, C., Doyle, J., Behrmann, M., 2014. Emerging structure-function relations in the developing face processing system. Cereb. Cortex 24, 2964–2980. doi:10.1093/cercor/bht152

Schröder, J.M., Bohl, J., von Bardeleben, U., 1988. Changes of the ratio between myelin thickness and axon diameter in human developing sural, femoral, ulnar, facial, and trochlear nerves. Acta Neuropathol. 76, 471–483. doi:10.1007/BF00686386

Smith, S.M., 2002. Fast robust automated brain extraction. Hum. Brain Mapp. 17, 143–155. doi:10.1002/hbm.10062

Song, S.-K., Sun, S.-W., Ramsbottom, M.J., Chang, C., Russell, J., Cross, A.H., 2002. Dysmyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. Neuroimage 17, 1429–1436. doi:10.1006/nimg.2002.1267

Song, S.K., Sun, S.W., Ju, W.K., Lin, S.J., Cross, A.H., Neufeld, A.H., 2003. Diffusion tensor imaging detects and differentiates axon and myelin degeneration in mouse optic nerve after retinal ischemia. Neuroimage 20, 1714–1722. doi:10.1016/j.neuroimage.2003.07.005

Sutherland, M.E., Zatorre, R.J., Watkins, K.E., Hervé, P.Y., Leonard, G., Pike, B.G., Witton, C., Paus, T., 2012. Anatomical correlates of dynamic auditory processing: Relationship to literacy during early adolescence. Neuroimage 60, 1287–1295. doi:10.1016/j.neuroimage.2012.01.051

Trehub, S.E., 1973. Infants' sensitivity to vowel and tonal contrasts. Dev. Psychol. 9, 91–96. doi:10.1037/h0034999

Vos, S.B., Haakma, W., Versnel, H., Froeling, M., Speleman, L., Dik, P., Viergever, M.A., Leemans, A., Grolman, W., 2015. Diffusion tensor imaging of the auditory nerve in patients with long-term single-sided deafness. Hear. Res. 323, 1–8. doi:10.1016/j.heares.2015.01.010

Wakana, S., Jiang, H., Zijl, P.C.M. Van, Mori, S., 2004. Riber tract-based atlas of human white matter anatomy. Radiology 230, 77–87.

Wang, H., Liang, Y., Fan, W., Zhou, X., Huang, M., Shi, G., Yu, H., Shen, G., 2019. DTI study on rehabilitation of the congenital deafness auditory pathway and speech center by cochlear implantation. Eur. Arch. Oto-Rhino-Laryngology 276, 2411–2417. doi:10.1007/s00405-019-05477-7

Westerhausen, R., Grüner, R., Specht, K., Hugdahl, K., 2009. Functional relevance of interindividual differences in temporal lobe callosal pathways: A DTI tractography study. Cereb. Cortex 19, 1322–1329. doi:10.1093/cercor/bhn173

Westerhausen, R., Kompus, K., Hugdahl, K., 2014. Mapping hemispheric symmetries, relative asymmetries, and absolute asymmetries underlying the auditory laterality effect. Neuroimage 84, 962–970. doi:10.1016/j.neuroimage.2013.09.074

Wu, C., Huang, L., Tan, H., Wang, Y., Zheng, H., Kong, L., Zheng, W., 2016. Diffusion tensor imaging and MR spectroscopy of microstructural alterations and metabolite concentration changes in the auditory neural pathway of pediatric congenital sensorineural hearing loss patients. Brain Res. 1639, 228–234. doi:10.1016/j.brainres.2014.12.025

Yakovlev, P., Lecours, A., 1967. The myelogenetic cycles of regional maturation of the brain, in: A. Minkowski (Ed.), Regional Development of the Brain in Early Life. Blackwell Scientific Publications, Oxford, pp. 3–70.

Yeatman, J.D., Wandell, B.A., Mezer, A.A., 2012. Lifespan maturation and degeneration of human brain white matter. Nat Commun 29, 997–1003. doi:10.1016/j.biotechadv.2011.08.021.Secreted

Yin, T.C.T., Chan, J.C.K., 1990. Interaural time sensitivity in medial superior olive of cat. J. Neurophysiol. 64, 465–488. doi:10.1152/jn.1990.64.2.465

Zanin, J., Dhollander, T., Farquharson, S., Rance, G., Connelly, A., Nayagam, B.A., 2019. Review: Using diffusion-weighted magnetic resonance imaging techniques to explore the microstructure and connectivity of subcortical white matter tracts in the human auditory system. Hear. Res. 377, 1–11. doi:10.1016/j.heares.2019.02.014

Zhang, H., Schneider, T., Wheeler-Kingshott, C.A., Alexander, D.C., 2012. NODDI: practical in vivo neurite orientation dispersion and density imaging of the human brain. Neuroimage 61, 1000–16. doi:10.1016/j.neuroimage.2012.03.072

Tables and Figures

Hemisphere	Tract	Metric	α		k		t_0		Age (years)
			coef	SE	coef	SE	coef	SE	
Left	Lower	FA	--	--	--	--	--	--	--
		AD	4.50×10^{-4}	9.67×10^{-6}	0.477	0.112	7.22×10^{-4}	3.72×10^{-5}	4.83
		RD	2.46×10^{-4}	6.71×10^{-6}	0.524	0.104	4.80×10^{-4}	2.75×10^{-5}	4.40
		MD	3.14×10^{-4}	7.26×10^{-6}	0.503	0.101	5.60×10^{-4}	2.93×10^{-5}	4.58
		NDI	--	--	--	--	--	--	--
	Upper	FA	0.444	0.014	0.205	0.053	0.277	0.018	11.22
		AD	6.22×10^{-4}	6.01×10^{-6}	0.402	0.048	9.29×10^{-4}	2.37×10^{-5}	5.72
		RD	3.20×10^{-4}	6.20×10^{-6}	0.352	0.042	6.04×10^{-4}	2.06×10^{-5}	6.54
		MD	4.21×10^{-4}	5.62×10^{-6}	0.367	0.040	7.11×10^{-4}	1.96×10^{-5}	6.27
		NDI	0.646	0.034	0.126	0.028	0.321	0.017	18.28
Right	Lower	FA	--	--	--	--	--	--	--
		AD	4.56×10^{-4}	8.41×10^{-6}	0.732	0.236	7.16×10^{-4}	4.64×10^{-5}	3.15
		RD	2.54×10^{-4}	8.25×10^{-6}	0.448	0.148	4.15×10^{-4}	2.88×10^{-5}	5.14
		MD	3.23×10^{-4}	7.35×10^{-6}	0.577	0.175	5.18×10^{-4}	3.38×10^{-5}	3.99
		NDI	--	--	--	--	--	--	--
	Upper	FA	--	--	--	--	--	--	--
		AD	6.29×10^{-4}	7.72×10^{-6}	0.352	0.054	9.03×10^{-4}	2.51×10^{-5}	6.54
		RD	3.14×10^{-4}	1.12×10^{-5}	0.211	0.035	5.55×10^{-4}	1.73×10^{-5}	10.89
		MD	4.20×10^{-4}	9.06×10^{-6}	0.246	0.037	6.66×10^{-4}	1.82×10^{-5}	9.37
		NDI	--	--	--	--	--	--	--

Table 1: Growth model parameters for auditory pathway microstructure. Fitted Brody growth curve model coefficients (coef) and standard error (SE) for the asymptote (α), growth rate (k) and intercept (t_0) are shown for each metric across each auditory pathway. Bilateral lower tract and right upper tract FA and NDI were best fit with a linear model and growth coefficients are not provided. The age at 90% of α were calculated for tracts fit with a Brody growth curve.

Figure 1: Auditory pathway anatomy. (A) Schematic of the ascending auditory pathway. (B) Tractography of the lower (blue) and upper (red) auditory pathways are shown on a coronal slice of a subject FA map overlaid with intra-voxel multi-tensor models. (C) An axial slice shows the upper auditory pathway projects through the medial geniculate nucleus of the thalamus. (D) A sagittal slice shows the auditory pathway from the cochlear nerve to the inferior colliculus (white sphere). (E) An axial slice shows the complex fiber orientations at the level of the pons.

Figure 2: Age-related changes in auditory pathway microstructure are shown for the left and right upper and lower auditory pathways for DTI parameters (FA, AD, RD and MD) and NODDI NDI. The fitted line for each parameter and tract reflects the best fit model, where blue denotes linear regression and black denotes exponential changes. For models that demonstrate exponential age-related changes, the age at 90% of the asymptotic value is shown with a dashed red line.

Figure 3: Regional variation in the developmental timing of diffusion microstructure within the auditory pathway. The age at 90% of the asymptotic value of the fitted Brody growth curve for each point along the auditory pathway are shown for (A) AD, (B) MD, (C) NDI, (D) RD and (E) FA. Regions shaded in gray denote points where a growth curve did not provide the best fit model.

Figure 4: Regional linear maturation of microstructural parameters within the auditory pathway. The estimated slope (β) is provided at points along the auditory pathway for (A) FA and (B) NDI where the linear model provided the best fit model. Regions shaded in gray denote points where the Brody growth model provided the best fit.

Figure 5: Exponential growth curve error estimates for regional auditory pathway microstructural maturation. The standard error (SE) of the exponential growth rate (k) is provided for parameters along the auditory pathway obtained through bootstrapping with replacement. Regions shaded in gray denote points where the growth curve did not provide the best fit model.

Figure 6: Standard error (SE) for the slope (β) estimated with a linear model for diffusion parameters at points along the auditory pathway. Regions in gray denote points where the exponential growth curve provided the best fit model.

Figure 7: Regional lateralization of microstructural maturation within the auditory pathway. The laterality index (LI) for the age at 90% of α estimated using a Brody growth curve is provided for (A) AD, (B) MD and (C) RD. Red denotes regions that develop *later* than the contralateral side, while blue denotes regions that mature *earlier* than the contralateral side.