Retinoid X Receptor as a Therapeutic Target to Treat Neurological Disorders Associated with α-Synucleinopathy

Assylbek Zhylkibayev [1], Christopher R. Starr [2], M. Iqbal Hossain [3], Sandeep K Barodia [1], Shaida A. Andrabi [3,4], Maria B. Grant [2], Venkatram R. Atigadda [5], Marina S. Gorbatyuk [1]*, and Oleg S. Gorbatyuk [6]*

[1] Department of Biochemistry, Wake Forest University School of Medicine; Winston-Salem, NC, USA, azhylkib@wakehealth.edu (A.Z.); mgorbaty@wakehealth.edu (M.S.G.)

[2] Department of Ophthalmology, University of Alabama at Birmingham School of Medicine; Birmingham, AL, USA, crstarr@uab.edu (C.R.S.); mariagrant@uabmc.edu (M.B.G.)

[3] Department of Pharmacology and Toxicology, University of Alabama at Birmingham School of Medicine, Birmingham, AL, USA, iqbal7@uab.edu (M.I.H.); sandrabi@uab.edu (S.A.A.)

[4] Department of Neurology, University of Alabama at Birmingham School of Medicine, Birmingham, AL, USA, sandrabi@uab.edu (S.A.A.)

[5] Department of Dermatology, University of Alabama at Birmingham Heersink School of Medicine, AL, USA, venkatra@uab.edu

[6] Department of Translational Neuroscience, Wake Forest University School of Medicine, Winston-Salem, NC, USA, ogorbaty@wakehealth.edu

* Correspondence: ogorbaty@wakehealth.edu (O.S.G.); mgorbaty@wakehealth.edu (M.S.G.)

Abstract: This study investigated the therapeutic potential of the nuclear retinoid X receptor (RXR) in mitigating the progression of alpha-synucleinopathies (αSNPs), particularly in Parkinson's disease (PD). PD-like pathology in mice was successfully induced through the co-delivery of AAV expressing human α-synuclein (αS) and αS preformed fibrils (PFFs) into the substantia nigra pars compacta (SNpc). Significant increases in Lewy body (LB)-like inclusions, loss of tyrosine hydroxylase-positive (TH+) neurons, and reductions in dopamine (DA) levels in the striatum were observed. Additionally, diminished levels of PPARα and NURR1, along with elevated GFAP and Iba1, markers of neuroinflammation, microglial activation, and astrocytic gliosis were associated with PD pathogenesis. AAV-mediated overexpression of human RXRα demonstrated preservation of TH+ neurons, prevention of DA decline and attenuation of αS accumulation. Furthermore, RXR-treated PD brains showed a reduced number of GFAP+ and Iba1+ cells, decreased GFAP+ and Iba1+ immunoreactivity, and fewer and less widespread LB-like aggregates. RXR overexpression also enhanced the production of PPARα and NURR1, proteins critical for neuronal survival. These findings suggest that RXRα activation promotes neuroprotection by mitigating αSNPs and chronic neuroinflammation, a major contributor to PD progression. This research underscores the therapeutic potential of targeting nuclear receptors, such as RXR, in neurodegenerative diseases like PD

Keywords: α-Synucleinopathy; Parkinson disease; Retinoid X Receptor ; α-Synuclein; Lewy body, dopamine, inflammation, NURR1, PPARα

1. Introduction

Alpha-synucleinopathies (αSNPs) including Parkinson's disease (PD) and dementia with Lewy bodies (LB), rank as the most prevalent neurodegenerative condition after Alzheimer's disease (AD) [1]. These conditions are characterized by abnormal intraneuronal inclusions of α-synuclein (αS) [2]. While the pathology of PD is multifaceted, involving several interconnected processes, including the degeneration of dopamine (DA) nigral neurons, LB formation, mitochondrial dysfunction, protein aggregation, impaired clearance, and oxidative stress, chronic neuroinflammation stands out as a hallmark of PD pathophysiology [2,3]. The glial cell activation and heightened

levels of pro-inflammatory factors constitute prevalent features of the PD-afflicted brain [4]. To date, only a few treatments exist that interfere with inflammation and immune deficiency, including minocycline, dexamethasone, and several non-steroid anti-inflammatory drugs. While these approaches protect DA neurons from degeneration and glial cell activation, a breakthrough therapy based on innovative molecular mechanisms and targets is an unmet need.

The modulation of nuclear receptors (NRs) has been proposed as one of the most attractive therapeutic strategies for correcting PD pathobiology [5-11]. The nuclear retinoid X receptor (RXR) is of particular interest for therapeutic intervention, due to its binding and activation of both permissive partners, NURR1 and PPARs, whose dysfunctions have been observed in AD, multiple-system atrophy, and PD [12-14] and which are known to regulate expression of inflammation-associated genes [15,16]. Despite the importance of pharmacological RXR activation against PD-causing toxins [5,17,18], effective RXR-based treatments for PD patients are still in development.

Cumulative evidence indicates that abnormal RXR signaling triggers neuronal stress and neuroinflammation in PD pathobiology [6,8]. Activation of RXR binding partners, either NURR1 or PPAR is beneficial for reversing the PD pathology seen in animal and cellular models of PD [5,7,11,15,19-21]. Furthermore, it has been shown that activation of RXR by treatment with the RXR ligand LG100268 protects cultured DA neurons from stress and PD-like pathology induced by toxin 6-hydroxy dopamine (6-OHDA) [6]. The activation of PPAR by the application of fenofibrate and GW501516 reduces neuroinflammation and PD neurodegeneration [15]. Activation of the NURR1/RXRα heterodimer by NURR1 agonists can stand as a monotherapy for PD to increase striatal DA levels and up-regulate NURR1 target genes in PD-like brain conditions [22]. Collectively, these studies lead us to ask whether RXR/PPAR and RXR/NURR1 complexes are destabilized in the experimental PD model and whether delivery of exogenous RXR to the PD brain activates PPARs and/or NURR1, providing an effective interventional treatment aimed at controlling inflammatory response and reversing PD pathobiology.

The absence of evidence for the direct effects of RXR up-regulation on αS induced neurodegeneration in animal models of PD and the disease related αSNPs fueled our interest in conducting the current study. Using recombinant adeno-associate virus (AAV) to overexpress human RXRα (AAV-RXR) in the mouse substantia nigra pars compacta (SNpc), we examined the role of RXR in PD pathogenesis. Specifically, we investigated the impact of RXRα on the inflammatory response and the changes in viability of nigral tyrosine hydroxylase positive (TH+) cells in response to αS-induced neurotoxicity.

2. Materials and Methods

2.1 Primary mouse cortical neuron culture.

All the experiments involving the use of animals were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Alabama at Birmingham. Primary mouse embryonic cortical cells were isolated at embryonic day 15 (E15) as described previously [23]. Briefly, a day before isolation 12-well plates and glass slides were treated with attachment factor – poly-l-Ornithine (#P4957, Sigma-Aldrich, St. Louis, MO, USA). Mouse brains were dissected and placed in DMEM (#11-965-084, Fisher Scientific, Waltham, MA, USA) supplemented with 20% fetal bovine serum (#A4736401, Fisher Scientific, Waltham, MA, USA) were used as dissection medium. Cortical tissue was trypsinized using TypLE (#12-605-010, Fisher Scientific, Waltham, MA, USA) for 10 min at 37^0C. Digested tissue was homogenized by pipetting in DMEM+10% fetal bovine serum and centrifuged at 1200g for 5 min. Supernatant was discarded and replaced with Neurobasal medium (#A1371001, Fisher Scientific, Waltham, MA, USA) containing B27

supplements (#A3582801, Fisher Scientific, Waltham, MA, USA). Tissue pellets were 98
filtered through 40μm cell strainer (#50-146-1426, Fisher Scientific, Waltham, MA, USA) 99
and transferred to plates and slides. On the second in vitro day (DIV 2), growth of glial 100
cells was inhibited by adding 50 μM 5-fluoro-2-deoxyuridine (#f0503, Sigma-Aldrich, St. 101
Louis, MO, USA) for 3 days. AAV-GFP or AAV-RXR were added at DIV-2 as well, while 102
recombinant alpha synuclein pre-formed fibrils (PFFs) from Stress Marq Biosciences 103
(#SPR-322, Victoria, BC, Canada) were added starting DIV 5 at a concentration of 5μg/ml 104
and alpha synuclein protein from Stress Marq Biosciences (#SPR-322E, Victoria, BC, 105
Canada) at a concentration of 5μg/ml. Cells were grown until DIV 14 and further 106
harvested for analysis. 107

2.2 AAV vectors. 108

All vectors were packaged in AAV2/5 capsid and purified as described previously 109
[24]. Titers for all AAV constructs were equalized to 6 x10^{11} vg/ml. 110

2.3 Intracerebral injection of PFFs and AAV vectors 111

All surgical procedures were performed using aseptic techniques and isoflurane gas 112
anesthesia, as previously described [25]. Stereotaxic coordinates for intranigral injections 113
in C57BL6 mice (three-month-old males, 28±3 g) were AP -2.9 mm, ML +1.2 mm, and DV 114
-4.1 mm, relative to the bregma. The total injection volume was 2.0 μl at a speed of 0.5 115
μl/min. For these injections, we used a pulled glass pipet with an outer diameter not 116
exceeding 40–45 μm to avoid needle-related damage and reduce the injury-mediated 117
inflammatory response. 118

2.4 Behavioral test 119

The cylinder test was employed to measure spontaneous forelimb use [26]. A mouse was 120
placed in a glass cylinder and the number of times it rears up and touches the cylinder 121
wall was measured. The wall touches were subsequently scored for the left and right 122
paws. The data is expressed as the percentage of contralateral (left) to injected side paw 123
usage relative to the total number of touches by both paws. 124

2.5 Isolation and Processing of Tissues. 125

Animals were deeply anesthetized by ketamine (50mg/kg) and xylazine (10mg/kg) 126
cocktail intraperitoneally. The brains were removed and divided into two parts. The 127
caudal part containing the SNpc was fixed in ice-cold 4% paraformaldehyde in 0.1 M 128
phosphate buffer, pH 7.4. The fixed parts of brains were stored overnight at 4°C and then 129
transferred into 30% sucrose in 0.1 M PB for cryoprotection. Thirty μm thick coronal 130
sections were cut and further processed for immunohistochemistry. The rostral piece of 131
brain tissue was immediately used to dissect the right and left striatum. Tissue samples 132
were frozen separately on dry ice and kept at -80° C until assayed. In this study, the mice 133
from the 4-week time point of post-surgery were used to obtain SNpc tissue samples for 134
Western blot analysis. Frozen brains were sectioned on a Leica CM1510S cryostat (Leica, 135
Buffalo Grove, IL, USA) into 150 μm slices with SNpc tissue subsequently dissected out 136
under a microscope, as described previously [24]. 137

2.6 Immunohistochemistry. 138

For the bright-field microscopy analysis, 30μm floating sections were preincubated 139
with 1% H2O2–10% methanol for 15 min and then with 5% normal goat serum for 1 h. 140
Sections were incubated overnight at room temperature with anti-TH (1:2000; #MAB318, 141
mouse; Millipore) antibody. Next, incubation with biotinylated secondary anti-mouse 142
antibody was followed by incubation with avidin–biotin–peroxidase complex 143

(VECTASTAIN® Elite® ABC-HRP Kit, Peroxidase (Standard) (PK-6100) Vector Laboratories, Burlingame, CA, USA). Reactions were visualized using NovaRED Peroxidase (HRP) Substrate Kit (Vector NovaRED® Substrate Kit, Peroxidase (SK-4800), Vector Laboratories, Burlingame, CA, USA).

For confocal microscopy, sections were incubated with primary antibodies against human αS (1:1000; 32-8100, mouse; Invitrogen), pS129 (alpha-synuclein (phospho S129); 1:500; [EP1536Y] (ab51253); rabbit monoclonal; Abcam), and TH (anti-tyrosine hydroxylase antibody; 1:1000; AB1542, sheep; Millipore), human RXRα (RXRA monoclonal antibody (K8508); 1:200; 43-390-0; mouse , Invitrogen), and secondary fluorescent antibodies labeled with Alexa Fluor 488, 555, and 647 (1:500 for all; PIA32773-mouse; A21202-mouse; PIA32731TR-rabbit; PIA32794-rabbit; A21448-sheep, Invitrogen). The sections were examined using an AX-R Confocal Microscope coupled to a fully automated Ti2-E system (Nikon Instruments inc., Melville, NY, USA). NIS-Elements Software package was used for post-capture image analysis. Sequential scanning was used to suppress optical crosstalk between the fluorophores in stationary-structure colocalization assays. All manipulations of contrast and illumination on color images as well as color replacement were made using Adobe Photoshop CS software (Version 23.5.5).

2.7 Quantitation of Phospho-Synuclein Aggregates.

To quantify phospho-synuclein aggregates, 30μm floating brain sections were processed for immunofluorescent labeling with pS129 antibody. To count pS129-positive inclusions, an ImageJ macro was developed by modifying a previously described macro designed for automated cell counting.27 Briefly, the parameters of the counter were adjusted to reliably count pS129-positive inclusions. There was strong agreement between macro-counted sections and those counted manually. To count the aggregates in the macro, an area of interest was drawn on a micrograph within ImageJ, and then the macro was run to count the total number of aggregates in the outlined area. The area occupied by aggregates was manually quantified in ImageJ using the freehand tool and the measure function.

2.8 Unbiased Stereology.

The unbiased stereological estimation of the total number of the TH+ neurons in SNpc was performed using the optical fractionator method as described previously [28]. The sampling of cells to be counted was performed using the Micro Brightfield Stereo Investigator System. The software was used to delineate the transduction area at 4x on 30 μm sections and generate counting areas of 100 x 100 μm. The estimate of the total number of neurons and coefficient of error due to the estimation was calculated according to the optical fractionator formula [28].

2.9 DA measurements.

Frozen striatal tissue samples were shipped on dry ice to the Neurochemistry Core Lab at Vanderbilt University Medical Center, Nashville, TN for HPLC analysis of DA, 3,4-dihydroxyphenylacetic acid (DOPAC), and homovanillic acid (HVA) in the striatum. A total of 5-6 animals per group were analyzed for striatal DA metabolites.

2.10 Western blot Analysis.

Brain tissues were collected and lysed with RIPA buffer (Cell signaling, 9806, USA) supplemented with 1% Halt Protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific, 87786, USA). Samples were homogenized using a pestle (Fisherbrand™ RNase-Free Disposable Pellet Pestles; 12-141-368; Fisher, USA) and rotated at 4°C for 30 min, and

centrifuged at 4°C by 12000g speed for 15 min. Based on the Bradford method, the supernatant was used for protein quantitation using a protein assay (Bio-Rad Protein Assay Kit I; 5000001; Hercules, California, USA). 80-100 μg of protein were loaded onto 4–20% Mini-PROTEAN® TGX™ Precast Gel (Bio-Rad, 4561093EDU, USA) and transferred to polyvinylidene difluoride membrane (Biorad,1704272, USA) using the Trans-Blot Turbo Transfer System (BioRad, 1704150, USA). Membranes were incubated for 1hr in 5% skim milk (Bio-Rad, 1706404, USA) prepared with 1X Tris buffered saline (Bio-Rad, 1706435, USA) with 0.1% Tween 20 (Sigma-Aldrich, P1379, USA). Primary antibodies were diluted in 5% Bovine serum albumin (Fisher, BP9703-100, USA) dissolved in 1X TBS. The following primary antibodies were used in the study: Mouse anti-αS (1:2000; BD Transduction Laboratories, Zymed Laboratories #610787), mouse anti-TH (1:2000; Millipore #MAB318), anti-hRXRα mouse (1:200; Abcam #ab50546). Secondary antibodies (1:10000 for all) were purchased from LI-COR: HRP goat anti-Rabbit IgG (926-80011, USA); HRP goat anti-Mouse IgG (926-80010, USA). Images of membranes were captured and analyzed using the Odyssey XF system (LI-COR, USA).

2.11 Statistical Analysis.

Data were analyzed using one-way analysis of variance with Tukey post-test using Prism (GraphPad Software 10.0.2, Inc.). Data are presented as mean ± SE.

3. Results

3.1. Development of PD-like mouse model associated with αS-induced neurodegeneration.

In this study, we employed an animal model of PD-like pathology, as previously described by Thakur et al and Bjorklund et al [29,30]. In order to mimic the PD pathogenesis in C57BL6 adult male mice, we used combined injections of AAV-αS and PFFs (AAV/PFF) into the SNpc. The PFFs were purchased from Stressmarq (SPR-322) and used at a concentration of 5 mg/ml. The AAV containing a blank expression cassette, which is an αS vector with an early stop codon, served as a control virus (BV) [24,31]. Monomeric αS (Stressmarq; SPR-321) was used as a control substance to PFFs. A mixture of BV and monomeric αS (BV/αS) served as a control for combined AAV/PFF administration. Mice were injected with AAVs and/or PFFs on one side of the brain; the other side was kept as a non-transduced internal control (Figure 1).

Figure 1. Fluorescent microscopy of the mouse SNpc at 8 weeks after AAV-GFP and AAV/PFF injections. The upper image shows the expression of the GFP transgene on the injected side of mouse brain. GFP (green) was expressed in most TH+ neurons (red) in SNpc and also can be seen in the SN pars reticulata, as well as surrounding neuronal structures of the mesencephalon. The bottom image illustrates pS129-positive inclusions (white) in SNpc neurons and nearby structures. Scale bar: 0.5 mm.

The study involved several groups of animals: uninjected controls, and those injected with single AAV-αS, BV, AAV-GFP, AAV-RXR, and a combination with AAV/PFF or BV/αS. We established injection conditions to secure αS expression level and consistent with the reported αS levels in postmortem human PD brains[32], we used the AAV-αS titer that induced moderate levels of αS overexpression, which was three-fold higher relative to endogenous levels. To confirm this, we first validated a mouse anti-αS antibody capable of recognizing an epitope shared by both endogenous mouse αS and exogenous human αS (Figure 2A). In these experiments, the ratio of αS in the injected versus uninjected mouse SNpc was measured at 4 weeks post-injection, a time point when neurodegeneration was not yet significant. The αS ratio varied between 2.7 and 3.1, with an average increase of 2.9-fold in total αS expression (Figure 2B).

Figure 2. Comparative analysis of αS mouse models of PD. All AAV vectors have the same volume and titer (6 x1012 vg/ml). The total injection volume (2 μl) of AAV-GFP and AAV-αS was adjusted using PBS to equalize with BV/αS or AAV/PFF. (A) The comparative affinity test of anti-αS antibody used in the study to bind recombinant human and mouse αS proteins of known concentration at 4 weeks post-injection. (B) Measurement of αS expression in animals that had been injected with human αS at 4 weeks postinjection. Western blot (WB) showing nigral extracts taken from the uninjected and injected sides. Thirty micrograms of tissue extract was analyzed by WB using an antibody that recognize

both human and mouse αS. The amount of αS was normalized to β-actin and the ratio of the injected versus uninjected sides was calculated (n=6). (C) Combined AAV/PFF administration into SNpc resulted in more severe loss of nigral TH+ cells and striatal DA compared with individual injections of AAV-αS or PFF at 8 weeks post injection. Upper graph demonstrates unbiased estimation of nigral TH+ cells in SNpc of mice injected with AAV-αS or PFF and a combination of AAVs/PFF or BV/αS. Tissue slices were labeled with antibody to TH as described in Materials and Methods and the percentage of surviving cells was calculated by comparison with the uninjected side in the same animal. Lower graph shows measurement of striatal DA. The amount of DA in striatal tissue was measured on the injected and uninjected sides of individual animals and displayed as the mean percentage of DA remaining on the injected side compared with the uninjected side plus standard error. Both graphs demonstrate the significant loss of nigral TH+ cells and striatal DA in AAV/PFF injected mice. p*< 0.05; **<0.01; ***<0.001; ***,0.0001 (n=4 for each group except of TH+ cell count in uninjected control, n=14).

We observed significant striatal dopamine (DA) and TH+ neuronal loss at 8 weeks post-injection. These effects were most dramatic in mice injected with AAV/PFF as compared to those receiving either rAAV-αS or PFFs alone (Figure 2C), highlighting the advantages of the combined approach in accelerating the development of LB-like features. Notably, no differences were observed between uninjected mice and BV/αS injected controls, nor between control groups and PFFs injections alone. However, substantial differences were detected between control groups and AAV-αS as well as between control groups or AAV-αS and AAV/PFF (Figure 2C). As a result of AAV/PFF injections, we observed the formation of pS129-positive inclusions (Figure 1). These inclusions were associated with increased neuronal vulnerability seen as dramatic reduction in number of TH+ cells in the SNpc accompanied by a decline in striatal DA levels (Figure 2C).

3.2. Combined AAV/PFF injection induce severe neurodegeneration accompanied by inflammatory response and gliosis.

To investigate a link between the observed aggregates, TH+ cell loss, and inflammation, we next analyze the level of GFAP. GFAP is upregulated in response to CNS injury or disease and serves as a marker of astrogliosis. Elevated GFAP levels are observed in conditions such as AD, PD, HD, and ALS and contribute to neurodegeneration via pro-inflammatory signaling [33-35]. During activation, accompanied by upregulated GFAP expression, astrocytes release pro-inflammatory cytokines and chemokines (e.g., IL-1β, TNF-α), which promote immune cell recruitment. Another protein, IBA1, serves as a marker for microglia and their activation, reflecting microglial dynamics in CNS health and disease. Its upregulation in activated microglia makes it a critical tool to study neuroinflammation. Activated microglia release cytokines such as IL-1β, TNF-α, and IL-6, which can exacerbate neuroinflammation. Therefore, we analyzed these markers in the PD brains.

Figure 3 demonstrates a significant increase in both GFAP and IBA1 fluorescent signals ranging from 2- to 3- fold in AAV/PFF injected brains previously identified with dramatic TH+ cell loss compared to BV/αS injections. Results of this experiment indicated a notable inflammatory response associated with the combined AAV/PFF approach.

Figure 3. Combined AAV/PFF injection led to significant inflammation response in mouse brain evident by increase in GFAP and IBA1, markers of astrocytes and microglia. Of note, we did not find the difference between uninjected and injected SNpc of any control groups. p< ***,0.0001 (n=5-6). Scale bar: 1mm.

3.3. RXR diminishes LB-like formation in primary mouse cortical neurons.

Preliminary testing of AAV-RXR vector in primary mouse neuronal cells (Figure 4) demonstrates a powerful RXR effect on the formation of 129-positive LB-like inclusion. Despite diffuse pS129 immunostaining was observed in both GFP and RXR experimental groups, LB-like αS inclusions (denoted by arrows) localized to cell bodies were identified only in GFP-positive cells after nine days of exposure to PFFs and monomeric αS. None of these inclusions were found in hRXRα-expressing neurons, suggesting the therapeutic potential of treatment with exogenous RXR.

Figure 4. Confocal images illustrate AAV-mediated expression of native GFP (a-c) and human RXRα (d-f), both shown in green, in cortical neurons at DIV14 after nine days of exposure to PFFs and monomeric αS. Both experimental groups demonstrate immunofluorescent staining with anti-pS129 antibody (shown in red). However, LB-like

αS inclusions localized to cell bodies denoted by arrows were identified only in GFP-positive cells and none of them were found in hRXRα expressing neurons. DAPI staining (in blue) added in (c). Scale bars: 25 μm.

3.4. RXR overexpression significantly ameliorate PD-associated pathology in mouse PD model.

Previous studies indicated that reduced RXR signaling triggers neuronal stress and neuroinflammation in PD pathobiology [6,8]. To explore the therapeutic potential of RXR activation for PD treatment, we utilized AAV-mediated transduction to deliver either human RXRα or control GFP, alongside AAV/PFF, into the mouse SNpc. Various control and experimental groups were included in the study, incorporating combinations of GFP, human RXRα viral vectors, and control BV/αS. The mice were analyzed at 4- and 8-weeks post-injection (Figure 5 and Figure 6). An unbiased stereology count of nigral TH+ cells in the injected SNpc was compared with the uninjected SNpc for each mouse. No significant differences were observed between animals injected with the GFP, human RXRα vectors, and/or control BV/αS, and the uninjected side at both the 4- and 8-week time points (p = 0.963; data not shown). Therefore, the BV/αS control group at both time points was used for further comparisons.

Figure 5. The SNpc of mice at 4-week after PFFs and AAV vector injections. Confocal imaging reveals pS129-positive aggregates in TH+ neurons in SNpc of mice treated with AAV/PFF. WB analyses show no difference in nigral TH protein levels between control groups and AAV/PFF + AAV-GFP injected brains, which correlates with stereology counts of TH+ cells in this time point. However, TH protein level was increased in AAV/PFF + AAV-RXR treated nigras. The levels of PPAR and NURR1 were significantly downregulated in AAV/PFF injected brains compared to BV/αS and AAV/PFF + AAV-RXR injected brains. No difference was observed between BV/αS and uninjected brains. The level of pS129 protein adjusted to total αS was significantly downregulated in AAV/PFF+AAV-RXR compared to AAV-PFF+AAV-GFP mice. p*< 0.05; **<0.01 (n=6). Scale bar: 25 μm.

Figure 5 presents the study results of the mouse SNpc terminated at 4 weeks post-injection with AAV/PFF. Despite confocal imaging revealing pS129-positive inclusions in TH+ neurons in the SNpc and surrounding nervous structures of mice treated with AAV/PFF, no significant nigral TH+ cell loss or marked reduction in striatal DA were detected in the mouse brains at this point. However, a notable reduction in PPARα levels was observed in the nigral tissue of AAV/PFF-injected mice at 4 weeks post-injection. At the same time, co-injection of AAV/PFF with AAV-RXR led to over 2-fold increase in PPARα level compared with AAV/PFF + AAV-GFP injected mice. In addition to PPARα, RXR overexpression in AAV/PFF-injected mice enhanced NURR1 production in the SNpc at 4 weeks post-injection. This implies that RXRα overexpression in the PD-like brain activates NURR1, which has been proposed as a standalone therapeutic target for PD [22] Consistent with the previous study demonstrating NURR1-induced TH production [17], this NURR1 upregulation was accompanied with significant increase in TH protein. Moreover, the level of pS129 protein was significantly downregulated ($p<0.001$) in AAV/PFF+AAV-RXR compared to AAV-PFF+AAV-GFP mice at 4 weeks postinjection. This finding indicates that NURR1 is a downstream target of RXR activation and underscores the potential of RXR-mediated NURR1 therapy.

Figure 6. AAV-mediated overexpression of human (h) RXRα ameliorates PD-like pathology induced by AAV/PFF intra-nigral injection. Confocal imaging demonstrates expressions of transgenic GFP or hRXR (in green) within TH+ neurons (in red) with pS129 deposits (in white). Unbiased stereology counts of TH+ cells and measurement levels of striatal DA are shown. p*< 0.05; **<0.01; ***<0.001; ****<0.0001.(n=6).

At 8 weeks post-injection, while AAV/PFF treatment induced a significant loss of nigral TH+ cells and striatal DA, the simultaneous delivery of human RXRα and AAV-PFF diminished TH+ cell loss and increased DA level (Figure 6). Additionally, RXRα

overexpression led to a decrease in GFAP immunoreactivity and GFAP+ punctata, 372
indicating reduced recruitment of astrocytes to the AAV/PFF-affected area at 8 weeks 373
post-injection (Figure 7). It is also noteworthy to mention that the microglial marker IBA1 374
was diminished in RXR-overexpressing brains with pS129-positive inclusions, further 375
supporting the anti-inflammatory effects of RXRα overexpression. 376

Motor behavior impairment was assessed using the cylinder test to measure 378
asymmetry in forelimb use. At 4 weeks post-injection, no significant differences were 379
observed between experimental groups (Figure S1). However, by 8 weeks post-injection, 380
the AAV/PF + AAV-GFP group exhibited significant motor impairment compared to the 381
control BV/αS group. In contrast, mice injected with AAV/PFF + AAV-RXR showed a 382
trend toward improvement relative to the AAV/PF + AAV-GFP group, though this 383
difference did not reach statistical significance (Figure S1). 384

385
Figure 7. Confocal images illustrate reduced expression of inflammatory markers, GFAP 386
(in red) and IBA1 (in green), in AAV/PFF brains upon AAV-RXR delivery. TH+ cells are 387
shown in blue. The graphs depict the results of immunofluorescent detection of GFAP 388
and IBA1 expression shown in arbitrary units. p*< 0.05; **<0.01; ***<0.001; 389
****<0.0001.(n=6). 390

Finally, we analyzed the distribution area of pS129-positive inclusions affected by 392
RXR overexpression by comparing injection sites of AAV/PFF + AAV-GFP and AAV/PFF 393
+ AAV-RXR mice. We counted the pS129-positive inclusions in outlined areas (Figure 8). 394

Remarkably, the number of aggregates and the size of the area containing pS129+- LB-like 395
aggregates, measured in μm², significantly decreased with the distance from the point of 396
injection in AAV/PFF + AAV-RXR injected brains compared to AAV/PFF + AAV-GFP 397
injections (p<0.0001). 398

Figure 8. The pS129-positive inclusions (white) at injection sites of mouse brains treated 400
with AAV/PFF + AAV-GFP and AAV/PFF + AAV-RXR. TH-positive cells stained in red. 401
Graphs represent comparative count and distribution area of pS129-positive aggregates. 402
Student t-test, p****<0.0001. 403

4. Discussion 404

In the current study, we present a reliable mouse model that mimics PD-like 405
pathology, including the accumulation of LB-like aggregates, TH neuronal loss, DA 406
deficit, and activation of astrocytes and microglia. Our data indicate that a combination 407
approach delivering AAV-αS and commercial PFFs to the SNpc results in the activation 408
of a severe inflammatory response. This approach significantly accelerates, and mimics 409
PD progression compared to previously reported models based on individual AAV-αS or 410
PFFs injections. This confirms the observation by Bjorklund A et al [30] that the combined 411
AAV/PFF delivery offers several advantages over the standard PFF model due to the 412
enhanced and accelerated αS pathology and glial response induced by the PFFs and 413
elevated αS levels, as shown in Figures 2C and 3. An advantage of the current study is 414
that we demonstrated that the AAV/PFF-mediated neurodegeneration provides a 415
window of opportunity for drug delivery according to our data. A limitation of the current 416
study is that the RXR-based intervention in the PD pathogenesis should be tested at later 417
stages with marked disease progression thus mimicking more of the real-life situation in 418
individuals with PD. 419

The AAV/PFF mouse model provides benefits by triggering an early inflammatory response observed at 3-4 weeks post-injection before severe aggregate formation is observed, as previously reported [30]. Notably, the significant striatal DA and nigral TH+ cell loss was not yet detected at this time point; however, a level of protein expression induced by AAV-mediated gene transfer reaches a maximum [25,36,37]. This point allows the study pathogenic mechanisms triggered by AAV/PFF.

Our data indicates that overexpression of human RXRα in the SNpc ameliorates αS-associated neurodegeneration. This was evident through a diminished loss of striatal DA and nigral TH+ cells, and reduced deposition of pS129-positive aggregates. Furthermore, we revealed reduced GFAP and IBA1-positive fluorescent signaling and number of positive cells. Given the limitation of the current study, including lack of the detailed analysis of activated astrocytes and microglia phenotypes, this reduction suggests fewer recruited astrocytes and IBA1-positive microglial cells, overall indicating a mitigation of neuroinflammation. In agreement with diminished neuroinflammation, the reduction in αS deposits in the PD brain also suggests that RXR upregulation prohibits the aggregation of an insoluble form of the protein. The results of TH+ cell counting, and striatal DA assay align, indicating the neuroprotective effects of RXR upregulation.

Based on our data, we anticipate that PPAR is necessary for RXR-based therapies to enhance TH cell survival and DA levels in the PD brain. We revealed that AAV/PFF treatment downregulates PPARα production. In turn, RXR overexpression induces significant upregulation of PPARα. It is well known that PPAR α, β, and γ are the permissive RXR binding partners; when activated in the central nervous system, RXR plays a critical role [9]. Recent studies have revealed reduced PPARα expression in AD brains [38] and the genetic linkage to the disease [39,40]. The neuroprotective effect of PPAR agonists have recently been proposed using neurological murine models and include anti-inflammatory effects, APP degradation, and Aβ inhibitory functions [41-45] The pharmacological modulation of PPARs by pioglitazone, rosiglitazone, ibuprofen, piroxicam, ciglitazone, and GW1929 was shown to be neuroprotective in AD, ALS, MS, and PD models [46-51]. In particular, in MPTP and 6-OHDA models of PD, preclinical studies have shown that different PPARγ agonists reduce neuronal cell death, microglial activation, and DA level in the hippocampus and the SNpc. [52] [19,53] [15,54]. A conformational change in the RXRa/PPARγ complex allows the heterodimer to bind PPAR-response elements (PPRE) in the nucleus to activate gene transcription. Therefore, it is not surprising that our results have demonstrated that the AAV-mediated overexpression of human RXRα increases the expression of PPARα in mouse brains with PD-like pathology.

Our data also indicate that in addition to PPARα, RXR overexpression in mice with experimental α-synucleinopathy enhances NURR1 production in SNpc. This finding is in accordance with the multiple studies that highlighted the significance of changes in NURR1 expression during the pathogenesis of PD and related disorders. Thus, it has been shown that NURR1 is significantly decreased in the PD nigral neurons containing αS-immunoreactive inclusions and in the DA nigral neurons with neurofibrillary tangles, correlating with the loss of TH and diminished intracellular pathology in both αSNP and tauopathies [55]. Furthermore, downregulation of NURR1 promotes upregulation of the NF-κB/ NLRP3 inflammasome axis [56-58], while NURR1 overexpression and activation by means of HX600 agonist treatment have significant anti-inflammatory effects decreasing the expression of genes involved in inflammation control [59] and reducing microglia-expressed proinflammatory mediators, preventing neuroinflammation and neuronal cell death [17]. It is also worth mentioning that the TH upregulation and anti-inflammatory therapeutic effect from the application of HX600 agonist is associated with the activation of the NURR1/RXR complex, suggesting their dual role in PD development.

In summary, the current study not only validated a PD-like mouse model using commercially available PFFs, making it convenient to study the role of genetic modifiers, but also provided evidence that mice with progressive αSNPs present a model for druggable intervention in PD progression. Our study demonstrated the power of RXRα-based therapy to mitigate PD progression by targeting major pathological events, such as the formation of LB-like structures, loss of TH+ neurons, and neuroinflammatory response. The mechanism of RXR-based intervention requires future investigation. At this point, it is challenging to anticipate whether enhanced RXR will demonstrate a preference for binding to one nuclear receptor over another. Moreover, the RXR-based therapy may enhance the pool of RXR homodimers available to partner with both NURR1 and PPARs to regulate the expression of genes responsible for anti-inflammatory action.

Funding: The study has been partly supported by M. J. Fox Foundation (O.S.G.) and the National Eye Institute, grant number R01 EY035539 (M.S.G.).

Institutional Review Board Statement: The animal study protocol was approved by the University of Alabama at Birmingham - Institutional animal care and use committee (protocol no. 10085).

Data Availability Statement: The research data is available upon request. The raw data are also available as a supplemental material.

Acknowledgments: We would like to thank Dr. David S. Standaert for his critical evaluation of the manuscript and for providing valuable insights that enhanced its quality.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Bostrom, F., Jonsson, L., Minthon, L., and Londos, E. (2007). Patients with Lewy body dementia use more resources than those with Alzheimer's disease. Int J Geriatr Psychiatry 22, 713-719. 10.1002/gps.1738.

2. Lamotte, G., and Singer, W. (2023). Synucleinopathies. Handb Clin Neurol 196, 175-202. 10.1016/B978-0-323-98817-9.00032-6.

3. Goedert, M., Jakes, R., and Spillantini, M.G. (2017). The Synucleinopathies: Twenty Years On. J Parkinsons Dis 7, S51-S69. 10.3233/JPD-179005.

4. Wang, Q., Liu, Y., and Zhou, J. (2015). Neuroinflammation in Parkinson's disease and its potential as therapeutic target. Transl Neurodegener 4, 19. 10.1186/s40035-015-0042-0.

5. Spathis, A.D., Asvos, X., Ziavra, D., Karampelas, T., Topouzis, S., Cournia, Z., Qing, X., Alexakos, P., Smits, L.M., Dalla, C., et al. (2017). Nurr1:RXRalpha heterodimer activation as monotherapy for Parkinson's disease. Proc Natl Acad Sci U S A 114, 3999-4004. 10.1073/pnas.1616874114.

6. Friling, S., Bergsland, M., and Kjellander, S. (2009). Activation of Retinoid X Receptor increases dopamine cell survival in models for Parkinson's disease. BMC Neurosci 10, 146. 10.1186/1471-2202-10-146.

7. Al-Nusaif, M., Yang, Y., Li, S., Cheng, C., and Le, W. (2022). The role of NURR1 in metabolic abnormalities of Parkinson's disease. Mol Neurodegener 17, 46. 10.1186/s13024-022-00544-w.

8. Sharma, S., Shen, T., Chitranshi, N., Gupta, V., Basavarajappa, D., Sarkar, S., Mirzaei, M., You, Y., Krezel, W., Graham, S.L., and Gupta, V. (2022). Correction to: Retinoid X Receptor: Cellular and Biochemical Roles of Nuclear Receptor with a Focus on Neuropathological Involvement. Mol Neurobiol 59, 2051. 10.1007/s12035-022-02767-w.

9. van Neerven, S., Kampmann, E., and Mey, J. (2008). RAR/RXR and PPAR/RXR signaling in neurological and psychiatric diseases. Prog Neurobiol 85, 433-451. 10.1016/j.pneurobio.2008.04.006.

10. Corona, J.C., and Duchen, M.R. (2015). PPARgamma and PGC-1alpha as therapeutic targets in Parkinson's. Neurochem Res 40, 308-316. 10.1007/s11064-014-1377-0.

11. Corona, J.C., and Duchen, M.R. (2016). PPARgamma as a therapeutic target to rescue mitochondrial function in neurological disease. Free Radic Biol Med 100, 153-163. 10.1016/j.freeradbiomed.2016.06.023.

12. Rahimi, A., Faizi, M., Talebi, F., Noorbakhsh, F., Kahrizi, F., and Naderi, N. (2015). Interaction between the protective effects of cannabidiol and palmitoylethanolamide in experimental model of multiple sclerosis in C57BL/6 mice. Neuroscience 290, 279-287. 10.1016/j.neuroscience.2015.01.030.

13. Ferret-Sena, V., Capela, C., and Sena, A. (2018). Metabolic Dysfunction and Peroxisome Proliferator-Activated Receptors (PPAR) in Multiple Sclerosis. Int J Mol Sci 19. 10.3390/ijms19061639.

14. Le, W.D., Xu, P., Jankovic, J., Jiang, H., Appel, S.H., Smith, R.G., and Vassilatis, D.K. (2003). Mutations in NR4A2 associated with familial Parkinson disease. Nat Genet 33, 85-89. 10.1038/ng1066.

15. Perez-Segura, I., Santiago-Balmaseda, A., Rodriguez-Hernandez, L.D., Morales-Martinez, A., Martinez-Becerril, H.A., Martinez-Gomez, P.A., Delgado-Minjares, K.M., Salinas-Lara, C., Martinez-Davila, I.A., Guerra-Crespo, M., et al. (2023). PPARs and Their Neuroprotective Effects in Parkinson's Disease: A Novel Therapeutic Approach in alpha-Synucleinopathy? Int J Mol Sci 24. 10.3390/ijms24043264.

16. Al-Nusaif, M., Lin, Y., Li, T., Cheng, C., and Le, W. (2022). Advances in NURR1-Regulated Neuroinflammation Associated with Parkinson's Disease. Int J Mol Sci 23. 10.3390/ijms232416184.

17. Loppi, S., Kolosowska, N., Karkkainen, O., Korhonen, P., Huuskonen, M., Grubman, A., Dhungana, H., Wojciechowski, S., Pomeshchik, Y., Giordano, M., et al. (2018). HX600, a synthetic agonist for RXR-Nurr1 heterodimer complex, prevents ischemia-induced neuronal damage. Brain Behav Immun 73, 670-681. 10.1016/j.bbi.2018.07.021.

18. Wang, J., Bi, W., Zhao, W., Varghese, M., Koch, R.J., Walker, R.H., Chandraratna, R.A., Sanders, M.E., Janesick, A., Blumberg, B., et al. (2016). Selective brain penetrable Nurr1 transactivator for treating Parkinson's disease. Oncotarget 7, 7469-7479. 10.18632/oncotarget.7191.

19. Machado, M.M.F., Bassani, T.B., Coppola-Segovia, V., Moura, E.L.R., Zanata, S.M., Andreatini, R., and Vital, M. (2019). PPAR-gamma agonist pioglitazone reduces microglial proliferation and NF-kappaB activation in the substantia nigra in the 6-hydroxydopamine model of Parkinson's disease. Pharmacol Rep 71, 556-564. 10.1016/j.pharep.2018.11.005.

20. Argyrofthalmidou, M., Spathis, A.D., Maniati, M., Poula, A., Katsianou, M.A., Sotiriou, E., Manousaki, M., Perier, C., Papapanagiotou, I., Papadopoulou-Daifoti, Z., et al. (2021). Nurr1 repression mediates cardinal features of Parkinson's disease in alpha-synuclein transgenic mice. Hum Mol Genet 30, 1469-1483. 10.1093/hmg/ddab118.

21. Kim, W., Tripathi, M., Kim, C., Vardhineni, S., Cha, Y., Kandi, S.K., Feitosa, M., Kholiya, R., Sah, E., Thakur, A., et al. (2023). An optimized Nurr1 agonist provides disease-modifying effects in Parkinson's disease models. Nat Commun 14, 4283. 10.1038/s41467-023-39970-9.

22. Saijo, K., Winner, B., Carson, C.T., Collier, J.G., Boyer, L., Rosenfeld, M.G., Gage, F.H., and Glass, C.K. (2009). A Nurr1/CoREST pathway in microglia and astrocytes protects dopaminergic neurons from inflammation-induced death. Cell 137, 47-59. 10.1016/j.cell.2009.01.038.

23. Hossain, M.I., Marcus, J.M., Lee, J.H., Garcia, P.L., Singh, V., Shacka, J.J., Zhang, J., Gropen, T.I., Falany, C.N., and Andrabi, S.A. (2021). Restoration of CTSD (cathepsin D) and lysosomal function in stroke is neuroprotective. Autophagy 17, 1330-1348. 10.1080/15548627.2020.1761219.

24. Salganik, M., Sergeyev, V.G., Shinde, V., Meyers, C.A., Gorbatyuk, M.S., Lin, J.H., Zolotukhin, S., and Gorbatyuk, O.S. (2015). The loss of glucose-regulated protein 78 (GRP78) during normal aging or from siRNA knockdown augments human alpha-synuclein (alpha-syn) toxicity to rat nigral neurons. Neurobiol Aging 36, 2213-2223. 10.1016/j.neurobiolaging.2015.02.018.

25. Gorbatyuk, M.S., Shabashvili, A., Chen, W., Meyers, C., Sullivan, L.F., Salganik, M., Lin, J.H., Lewin, A.S., Muzyczka, N., and Gorbatyuk, O.S. (2012). Glucose regulated protein 78 diminishes alpha-synuclein neurotoxicity in a rat model of Parkinson disease. Mol Ther 20, 1327-1337. 10.1038/mt.2012.28.

26. Magno, L.A.V., Collodetti, M., Tenza-Ferrer, H., and Romano-Silva, M.A. (2019). Cylinder Test to Assess Sensory-motor Function in a Mouse Model of Parkinson's Disease. Bio Protoc 9, e3337. 10.21769/BioProtoc.3337.

27. Maidana, D.E., Tsoka, P., Tian, B., Dib, B., Matsumoto, H., Kataoka, K., Lin, H., Miller, J.W., and Vavvas, D.G. (2015). A Novel ImageJ Macro for Automated Cell Death Quantitation in the Retina. Invest Ophthalmol Vis Sci 56, 6701-6708. 10.1167/iovs.15-17599.

28. Kirik, D., Rosenblad, C., Burger, C., Lundberg, C., Johansen, T.E., Muzyczka, N., Mandel, R.J., and Bjorklund, A. (2002). Parkinson-like neurodegeneration induced by targeted overexpression of alpha-synuclein in the nigrostriatal system. J Neurosci 22, 2780-2791. 10.1523/JNEUROSCI.22-07-02780.2002.

29. Thakur, P., Breger, L.S., Lundblad, M., Wan, O.W., Mattsson, B., Luk, K.C., Lee, V.M.Y., Trojanowski, J.Q., and Bjorklund, A. (2017). Modeling Parkinson's disease pathology by combination of fibril seeds and alpha-synuclein overexpression in the rat brain. Proc Natl Acad Sci U S A 114, E8284-E8293. 10.1073/pnas.1710442114.

30. Bjorklund, A., Nilsson, F., Mattsson, B., Hoban, D.B., and Parmar, M. (2022). A Combined alpha-Synuclein/Fibril (SynFib) Model of Parkinson-Like Synucleinopathy Targeting the Nigrostriatal Dopamine System. J Parkinsons Dis 12, 2307-2320. 10.3233/JPD-223452.

31. Gully, J.C., Sergeyev, V.G., Bhootada, Y., Mendez-Gomez, H., Meyers, C.A., Zolotukhin, S., Gorbatyuk, M.S., and Gorbatyuk, O.S. (2016). Up-regulation of activating transcription factor 4 induces severe loss of dopamine nigral neurons in a rat model of Parkinson's disease. Neurosci Lett 627, 36-41. 10.1016/j.neulet.2016.05.039.

32. Bobela, W., Aebischer, P., and Schneider, B.L. (2015). Alphalpha-Synuclein as a Mediator in the Interplay between Aging and Parkinson's Disease. Biomolecules 5, 2675-2700. 10.3390/biom5042675.

33. Zhang, N., Yan, Z., Xin, H., Shao, S., Xue, S., Cespuglio, R., and Wang, S. (2024). Relationship among alpha-synuclein, aging and inflammation in Parkinson's disease (Review). Exp Ther Med 27, 23. 10.3892/etm.2023.12311.

34. Eo, H., Kim, S., Jung, U.J., and Kim, S.R. (2024). Alpha-Synuclein and Microglia in Parkinson's Disease: From Pathogenesis to Therapeutic Prospects. J Clin Med 13. 10.3390/jcm13237243.

35. Harackiewicz, O., and Grembecka, B. (2024). The Role of Microglia and Astrocytes in the Pathomechanism of Neuroinflammation in Parkinson's Disease-Focus on Alpha-Synuclein. J Integr Neurosci 23, 203. 10.31083/j.jin2311203.

36. Burger, C., Gorbatyuk, O.S., Velardo, M.J., Peden, C.S., Williams, P., Zolotukhin, S., Reier, P.J., Mandel, R.J., and Muzyczka, N. (2004). Recombinant AAV viral vectors pseudotyped with viral capsids from serotypes 1, 2, and 5 display differential efficiency and cell tropism after delivery to different regions of the central nervous system. Mol Ther 10, 302-317. 10.1016/j.ymthe.2004.05.024.

37. Gorbatyuk, O.S., Li, S., Nguyen, F.N., Manfredsson, F.P., Kondrikova, G., Sullivan, L.F., Meyers, C., Chen, W., Mandel, R.J., and Muzyczka, N. (2010). alpha-Synuclein expression in rat substantia nigra suppresses phospholipase D2 toxicity and nigral neurodegeneration. Mol Ther 18, 1758-1768. 10.1038/mt.2010.137.

38. Wojtowicz, S., Strosznajder, A.K., Jezyna, M., and Strosznajder, J.B. (2020). The Novel Role of PPAR Alpha in the Brain: Promising Target in Therapy of Alzheimer's Disease and Other Neurodegenerative Disorders. Neurochem Res 45, 972-988. 10.1007/s11064-020-02993-5.

39. Koivisto, A.M., Helisalmi, S., Pihlajamaki, J., Hiltunen, M., Koivisto, K., Moilanen, L., Kuusisto, J., Helkala, E.L., Hanninen, T., Kervinen, K., et al. (2006). Association analysis of peroxisome proliferator-activated receptor gamma polymorphisms and late onset Alzheimer's disease in the Finnish population. Dement Geriatr Cogn Disord 22, 449-453. 10.1159/000095857.

40. Hamilton, G., Proitsi, P., Jehu, L., Morgan, A., Williams, J., O'Donovan, M.C., Owen, M.J., Powell, J.F., and Lovestone, S. (2007). Candidate gene association study of insulin signaling genes and Alzheimer's disease: evidence for SOS2, PCK1, and PPARgamma as susceptibility loci. Am J Med Genet B Neuropsychiatr Genet 144B, 508-516. 10.1002/ajmg.b.30503.

41. Heneka, M.T., Sastre, M., Dumitrescu-Ozimek, L., Hanke, A., Dewachter, I., Kuiperi, C., O'Banion, K., Klockgether, T., Van Leuven, F., and Landreth, G.E. (2005). Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and Abeta1-42 levels in APPV717I transgenic mice. Brain 128, 1442-1453. 10.1093/brain/awh452.

42. Sung, B., Park, S., Yu, B.P., and Chung, H.Y. (2006). Amelioration of age-related inflammation and oxidative stress by PPARgamma activator: suppression of NF-kappaB by 2,4-thiazolidinedione. Exp Gerontol 41, 590-599. 10.1016/j.exger.2006.04.005.

43. Sastre, M., Dewachter, I., Landreth, G.E., Willson, T.M., Klockgether, T., van Leuven, F., and Heneka, M.T. (2003). Nonsteroidal anti-inflammatory drugs and peroxisome proliferator-activated receptor-gamma agonists modulate immunostimulated processing of amyloid precursor protein through regulation of beta-secretase. J Neurosci 23, 9796-9804. 10.1523/JNEUROSCI.23-30-09796.2003.

44. Sastre, M., Dewachter, I., Rossner, S., Bogdanovic, N., Rosen, E., Borghgraef, P., Evert, B.O., Dumitrescu-Ozimek, L., Thal, D.R., Landreth, G., et al. (2006). Nonsteroidal anti-inflammatory drugs repress beta-secretase gene promoter activity by the activation of PPARgamma. Proc Natl Acad Sci U S A 103, 443-448. 10.1073/pnas.0503839103.

45. Camacho, I.E., Serneels, L., Spittaels, K., Merchiers, P., Dominguez, D., and De Strooper, B. (2004). Peroxisome-proliferator-activated receptor gamma induces a clearance mechanism for the amyloid-beta peptide. J Neurosci 24, 10908-10917. 10.1523/JNEUROSCI.3987-04.2004.

46. Chao, X., Xiong, C., Dong, W., Qu, Y., Ning, W., Liu, W., Han, F., Ma, Y., Wang, R., Fei, Z., and Han, H. (2014). Activation of peroxisome proliferator-activated receptor beta/delta attenuates acute ischemic stroke on middle cerebral ischemia occlusion in rats. J Stroke Cerebrovasc Dis 23, 1396-1402. 10.1016/j.jstrokecerebrovasdis.2013.11.021.

47. Kalra, P., Khan, H., Kaur, A., and Singh, T.G. (2022). Mechanistic Insight on Autophagy Modulated Molecular Pathways in Cerebral Ischemic Injury: From Preclinical to Clinical Perspective. Neurochem Res 47, 825-843. 10.1007/s11064-021-03500-0.

48. Prashantha Kumar, B.R., Kumar, A.P., Jose, J.A., Prabitha, P., Yuvaraj, S., Chipurupalli, S., Jeyarani, V., Manisha, C., Banerjee, S., Jeyabalan, J.B., et al. (2020). Minutes of PPAR-gamma agonism and neuroprotection. Neurochem Int 140, 104814. 10.1016/j.neuint.2020.104814.

49. Tong, Q., Wu, L., Gao, Q., Ou, Z., Zhu, D., and Zhang, Y. (2016). PPARbeta/delta Agonist Provides Neuroprotection by Suppression of IRE1alpha-Caspase-12-Mediated Endoplasmic Reticulum Stress Pathway in the Rotenone Rat Model of Parkinson's Disease. Mol Neurobiol 53, 3822-3831. 10.1007/s12035-015-9309-9.

50. Falcone, R., Florio, T.M., Di Giacomo, E., Benedetti, E., Cristiano, L., Antonosante, A., Fidoamore, A., Massimi, M., Alecci, M., Ippoliti, R., et al. (2015). PPARbeta/delta and gamma in a rat model of Parkinson's disease: possible involvement in PD symptoms. J Cell Biochem 116, 844-855. 10.1002/jcb.25041.

51. Lee, Y., Cho, J.H., Lee, S., Lee, W., Chang, S.C., Chung, H.Y., Moon, H.R., and Lee, J. (2019). Neuroprotective effects of MHY908, a PPAR alpha/gamma dual agonist, in a MPTP-induced Parkinson's disease model. Brain Res 1704, 47-58. 10.1016/j.brainres.2018.09.036.

52. Bonato, J.M., Bassani, T.B., Milani, H., Vital, M., and de Oliveira, R.M.W. (2018). Pioglitazone reduces mortality, prevents depressive-like behavior, and impacts hippocampal neurogenesis in the 6-OHDA model of Parkinson's disease in rats. Exp Neurol 300, 188-200. 10.1016/j.expneurol.2017.11.009.

53. Barbiero, J.K., Santiago, R.M., Persike, D.S., da Silva Fernandes, M.J., Tonin, F.S., da Cunha, C., Lucio Boschen, S., Lima, M.M., and Vital, M.A. (2014). Neuroprotective effects of peroxisome proliferator-activated receptor alpha and gamma agonists in model of parkinsonism induced by intranigral 1-methyl-4-phenyl-1,2,3,6-tetrahyropyridine. Behav Brain Res 274, 390-399. 10.1016/j.bbr.2014.08.014.

54. Hassanzadeh, K., Rahimmi, A., Moloudi, M.R., Maccarone, R., Corbo, M., Izadpanah, E., and Feligioni, M. (2021). Effect of lobeglitazone on motor function in rat model of Parkinson's disease with diabetes co-morbidity. Brain Res Bull 173, 184-192. 10.1016/j.brainresbull.2021.05.011.

55. Chu, Y., Le, W., Kompoliti, K., Jankovic, J., Mufson, E.J., and Kordower, J.H. (2006). Nurr1 in Parkinson's disease and related disorders. J Comp Neurol 494, 495-514. 10.1002/cne.20828.

56. Li, T., Tan, X., Tian, L., Jia, C., Cheng, C., Chen, X., Wei, M., Wang, Y., Hu, Y., Jia, Q., et al. (2023). The role of Nurr1-miR-30e-5p-NLRP3 axis in inflammation-mediated neurodegeneration: insights from mouse models and patients' studies in Parkinson's disease. J Neuroinflammation 20, 274. 10.1186/s12974-023-02956-x.

57. Li, W., Liu, X., Tu, Y., Ding, D., Yi, Q., Sun, X., Wang, Y., Wang, K., Zhu, M., and Mao, J. (2020). Dysfunctional Nurr1 promotes high glucose-induced Muller cell activation by up-regulating the NF-kappaB/NLRP3 inflammasome axis. Neuropeptides 82, 102057. 10.1016/j.npep.2020.102057.

58. Paik, S., Kim, J.K., Silwal, P., Sasakawa, C., and Jo, E.K. (2021). An update on the regulatory mechanisms of NLRP3 inflammasome activation. Cell Mol Immunol 18, 1141-1160. 10.1038/s41423-021-00670-3.

59. Briand, O., Helleboid-Chapman, A., Ploton, M., Hennuyer, N., Carpentier, R., Pattou, F., Vandewalle, B., Moerman, E., Gmyr, V., Kerr-Conte, J., et al. (2012). The nuclear orphan receptor Nur77 is a lipotoxicity sensor regulating glucose-induced insulin secretion in pancreatic beta-cells. Mol Endocrinol 26, 399-413. 10.1210/me.2011-1317.

Figure S1. Cylinder test.