Cross-modality representation and multi-sample integration of spatially resolved omics data

Zhen Li[1], Xuejian Cui[1], Xiaoyang Chen[1], Zijing Gao[1], Yuyao Liu[1], Yan Pan[1], Shengquan Chen[2], and Rui Jiang[1,*]

[1] Ministry of Education Key Laboratory of Bioinformatics, Bioinformatics Division at the Beijing National Research Center for Information Science and Technology, Center for Synthetic and Systems Biology, Department of Automation, Tsinghua University, Beijing 100084, China

[2] School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, China

[*] Corresponding author: ruijiang@tsinghua.edu.cn

Abstract

Spatially resolved sequencing technologies have revolutionized the characterization of biological regulatory processes within microenvironment by simultaneously accessing the states of genomic regions, genes and proteins, along with the spatial coordinates of cells, necessitating advanced computational methods for the cross-modality and multi-sample integrated analysis of spatial omics datasets. To address this gap, we propose PRESENT, an effective and scalable contrastive learning framework, for the cross-modality representation of spatially resolved omics data. Through comprehensive experiments on massive spatially resolved datasets, PRESENT achieves superior performance across various species, tissues, and sequencing technologies, including spatial epigenomics, transcriptomics, and multi-omics. Specifically, PRESENT empowers the incorporation of spatial dependency and complementary omics information simultaneously, facilitating the detection of spatial domains and uncovering biological regulatory mechanisms within microenvironment. Furthermore, PRESENT can be extended to the integrative analysis of horizontal and vertical samples across different dissected regions or developmental stages, thereby promoting the identification of hierarchical structures from a spatiotemporal perspective.

Introduction

Single-cell sequencing technologies have enabled the characterization of states and activities within individual cells from specific or even multi-omics perspectives, promoting molecular cell biology and disease research in areas such as cell atlas production, tumor immunology, and cancer genetics[1, 2]. However, an indispensable step of single-cell sequencing techniques involves the dissociation of cells from the tissue of interest, resulting in the loss of positional information[3]. Complex biological regulatory processes and cellular activities within tissue microenvironment are driven by sophisticated cell-cell interactions mediated by ligand-receptor partners, underscoring the importance of spatial distance between cells[4-6]. Recently, spatially resolved omics technologies have achieved simultaneous profiling of spatial locations and various omics layers, encompassing spatially resolved transcriptomics (SRT) [7, 8], spatial assay for transposase-accessible chromatin sequencing (ATAC-seq)[9, 10] and spatial multi-omics[11] (such as spatial CITE-seq[12], SPOTS[13] and MISAR-seq[14]), and accumulated substantial spatial omics sequencing data in repositories and databases[15, 16], thereby granting us a new perspective to decipher gene regulatory mechanisms in the context of tissue microenvironment.

However, spatial coordinate data and omics data exhibit significant modality differences, posing substantial challenges for the analysis of spatial omics data. Specifically, spatial coordinate data anchors the relative positions of cells within the tissue microenvironment, while omics data depicts the states of chromatin regions, genes and proteins involved in the gene regulatory processes within cells. Intended for this problem, cross-modality representation integrates the multi-modal data from spatially resolved datasets, including spatial coordinates, epigenomics, transcriptomics and proteomics, into a low-dimensional matrix. As a crucial step for spatial omics data analysis, cross-modality representation provides efficient and integrative features for various downstream applications and exploratory analyses[17-19], particularly the identification of spatial domains with distinct anatomical structures, cell-

type compositions, omics heterogeneity, and cell-cell interactions[20, 21], thereby facilitating the

comprehension of tissue functions achieved through the coordination of diverse spatial domains.

Due to technical limitations such as size restrictions of captured area during data acquisition, every

sample generated by spatial sequencing technologies typically focuses on a specific dissected region

under a certain condition, while cross-modality representation methods could only be applied to a single

sample because of spatial coordinate discrepancies between samples[22], impeding the exploration of

spatially resolved samples across multiple dissected areas or developmental stages[23]. Therefore,

integrative analysis of multiple samples across spatiotemporal scales, denoted as the multi-sample

integration, is also critical for spatial omics data analysis by providing a comprehensive and systematic

perspective for deciphering hierarchical and complex structures of cellular organizations[22-25]. Moreover,

recent studies have highlighted that integrating disease and normal samples facilitates the identification of

disease-specific domains and promotes the comprehensive exploration of disease onset and progression

mechanisms[25, 26], further emphasizing the significance of incorporating multiple spatially resolved

samples across various conditions.

Several computational methods have been proposed to address the exploration of SRT data[27, 28],

encompassing cross-modality representation methods, e.g., SpaGCN[29], STAGATE[30] and PAST[17], along

with some multi-sample integration methods such as STAligner[25], Stitch3D[31] and GraphST[23].

Nevertheless, compared with the rapid advancement in SRT data, the development of analytical methods

for new spatial sequencing data, namely the spatial ATAC-seq[9, 10] and spatial multi-omics[11, 13, 14] data, is

still stuck in the preliminary stage. Recently, SpatialGlue[32] was proposed as the first attempt for the cross-

modality representation of spatial multi-omics data by utilizing an attentional feature fusion mechanism.

Tian et al. proposed a dependency-aware generative variational autoencoder framework for the cross-

modality representation of SRT, spatial ATAC-seq and spatial CITE-seq data[33]. However, there are still no

methods specially designed for the multi-sample integration of spatial ATAC-seq and spatial multi-omics

data, forcing researchers to resort to integrative algorithms intended for single-cell or SRT data, which

overlook the critical spatial information[3] and the significant gaps between different omics layers[34-37],

respectively. Additionally, most current algorithms are only capable of accomplishing either cross-modal

representation or multi-sample integration for specific spatial omics data, without addressing the common

key issues underlying these two tasks. Therefore, for the widespread application of spatial omics data,

there is an urgent need for a unified, advanced, and scalable framework for the cross-modality and multi-

sample integrated analysis of various spatial omics data.

Both cross-modality representation and multi-sample integration of spatially resolved omics data face

two common key issues. Firstly, different omics layers of spatial data, encompassing transcriptomics,

epigenomics and proteomics, exhibit distinct patterns and distributions[34-37]. For example, compared with

RNA-seq data, ATAC-seq datasets suffer from larger dimensionality, more severe sparsity and higher

dropout rates, while those of proteomics data sequenced by antibody-derived tags (ADT) are opposite,

requiring targeted modeling of each omics layer. Secondly, while spatial multi-omics technologies

provide a systematic landscape of gene regulatory processes within cellular organizations[14] using cross-

modal spatial coordinate and omics data, they also require analytical algorithms with enhanced

capabilities to simultaneously fuse complementary multi-omics signals and spatial information.

Furthermore, there are additional challenges for the integrated analysis of multiple spatial samples. On the

one hand, due to discrepancies in the dissection and placement, the spatial coordinate data between

different samples cannot be directly compared[22], necessitating effective mechanisms to model the cross-

sample relationships. On the other hand, a major challenge in multi-sample integration lies in the intricate

coupling between technical noise and biological variations[38, 39], requiring methods to eliminate batch

effects while preserving important biological signals.

To this end, we propose PRESENT, an effective and scalable contrastive learning framework developed for the cross-modality representation of sPatially **RES**olved omics data, encompassing **E**pigenomics, tra**N**scriptomics, and pro**T**eomics. Through comprehensive experiments on massive real-world datasets generated by diverse spatial sequencing technologies, including but not limited to spatial transcriptomics[7, 8], spatial epigenomics[9, 10] and spatial multi-omics[11, 13, 14], we demonstrated the superior performance of PRESENT in cross-modality representation. Specifically, PRESENT can simultaneously capture spatial dependency and complementary multi-omics information, obtaining interpretable cross-modality representations for various downstream analyses, particularly the spatial domain identification. PRESENT also offers the potential to incorporate various reference data to address issues related to the low sequencing depth and signal-to-noise ratio in spatial omics data. Furthermore, PRESENT can also be extended to the integrative analysis of multiple samples across different dissection regions and developmental stages, effectively eliminating redundant batch effects while maintaining shared and sample-specific biological variations as well as detecting hierarchical functional structures from a spatiotemporal perspective. In summary, as a scalable and integrative tool in the field of spatial omics data analysis, PRESENT is anticipated to provide great potential for systematic and comprehensive exploration of gene regulatory processes and cellular activities in the context of tissue microenvironment.

Results

Overview of PRESENT

Taking a spatial coordinate matrix and an arbitrary combination of epigenomic, transcriptomic, or proteomic count matrices as input (Methods, Fig. 1a), PRESENT is built on a multi-view autoencoder and extracts spatially coherent biological variations contained in each omics layer via an omics-specific encoder consisting of a graph attention neural network[40] (GAT) and a Bayesian neural network[41] (BNN).

The GAT module aims to incorporates spatial dependency and corresponding omics information, while

the BNN module, proved powerful in non-linear regression problems in original studies[41], brings the

potential to leverage reference data from existing sequencing data, according to our previous study[17].

Afterwards, the omics-specific embeddings obtained by encoders are fused by a cross-omics alignment

module which implements a multi-modal contrastive learning loss function, i.e., inter-omics alignment

(IOA) loss[42] (Methods). Additionally, to capture the distinct patterns of various omics layers, PRESENT

includes omics-specific decoders which model the counts of transcriptomic and epigenomic data via a

zero-inflation negative binomial (ZINB) distribution[43] and a zero-inflation Poisson (ZIP) distribution[44]

(Methods), respectively. Due to significantly lower dropout rates, dimensionality, and sparsity, ADT

expression matrix is reconstructed using a multi-layer perceptron (MLP) module.

Compared to cross-modality representation on a single spatial sample, integrative analysis of multiple

samples faces the common challenges such as the characterization of different omics layers and the cross-

modality integration of spatial and omics information. There, we extend the cross-modality representation

framework PRESENT intended for a single sample to the multi-sample integration scenario, thereby

obtaining PRESENT-BC (Methods, Fig. 1b). The two additional critical issues of multi-sample

integration are overcoming the spatial coordinate discrepancies between samples to capture cross-sample

relationships[22], as well as distinguishing between complexly coupled biological variations and technical

batch effects[23-25, 31]. PRESENT-BC adopts a two-stage strategy to address these issues. In the first stage,

PRESENT-BC has almost the same architecture as PRESENT and is pretrained using a cross-sample

graph and omics data from multiple samples. The intra-sample edges of the cross-sample graph are built

using the spatial coordinates, while the inter-sample connections are established using omics information

(Methods). Additionally, the input of decoders of PRESENT-BC concatenates the one-hot encoding of

batch indices of spots, facilitating the extraction of batch effect-free representations. In the second stage,

we introduce an inter-batch alignment loss[45] and a batch-adversarial learning strategy[46] to align spots

from multiple samples and eliminate redundant batch effects. We further propose an intra-batch

preserving loss to maintain biological signals within each batch while integrating multiple samples.

Additionally, we implement a cyclic iterative optimization process to refine the cross-sample graph based

on the joint representations newly obtained from the trained model in each epoch, as the graph not only

enables the characterization of cross-sample relationships, but also empowers the construction of positive

neighbor sets for spots in the inter-batch alignment loss (Methods).

The low-dimensional cross-modality representations obtained by PRESENT facilitate various

downstream applications (Fig. 1c), including visualization and spatial domain detection, and further

reveal biological implications in the identified domains. When extended to multi-sample integration,

PRESENT-BC not only promotes the integrated of horizontal slices from different dissection regions, but

also enables vertical integration of samples from different developmental stages (Fig. 1c), paving the way

to explore the hierarchical cellular activities from a spatiotemporal perspective.

PRESENT facilitates accurate spatial domain identification in spatial ATAC-seq and SRT data

We first accessed the performance of PRESENT on the spatial ATAC mouse embryo datasets which

consist of three developmental stages[10], including E15.5, E13.5 and E12.5. The datasets include six

samples, with each developmental stage containing two replicated samples, denoted as *-S1 and *-S2. We

evaluated the data quality of the six samples using logarithmic total fragment counts, denoted as

Log10(nFrags), and transcription starting site (TSS) enrichment scores. The former metric represents the

cellular sequencing depth of samples[47], while the latter reflected the signal-to-noise ratio of datasets[48]. As

shown in Fig. 2a, b, the samples of E15.5 stage, i.e., E15.5-S1 and E15.5-S2, achieved the highest

sequencing depth and signal-to-noise ratio, while the E13.5-S1 sample and E12.5-S2 sample suffered

from the lowest sequencing depth and signal-to-noise ratio, respectively.

Spatial domain identification, which aims to decipher regions with distinct anatomical structures,

cell-type compositions, and cell-cell interactions[20, 21], is a fundamental step in revealing gene regulatory

processes and tissue functions in spatial omics data[29, 30]. To evaluate the performance of spatial domain

identification based on latent representations, we applied PRESENT and other six baseline methods,

including three methods (SCALE[49], cisTopic[50] and SnapATAC2[51]) for single-cell ATAC-seq data and

three methods (SpaGCN[29], STAGATE[30] and PAST[17]) for SRT data, to the six mouse embryo samples.

Based on the latent representations from these methods, we implemented the Leiden algorithm[52] to

acquire the same number of clusters as the ground truth domains (Methods). Four commonly used

metrics[53, 54], including adjusted rand index (ARI), adjusted mutual information (AMI), normalized mutual

information (NMI) and homogeneity (Homo), were utilized to quantitatively evaluate the performance of

spatial domain detection. As shown in Fig. 2c, PRESENT achieved the best performance, with

improvements of more than 8.60%, 26.13%, 25.91%, and 22.94% compared to the second-best baseline

method across the four metrics, respectively. As a qualitative evaluation and intuitive illustration of the

spatial domain identification, we also demonstrated the uniform manifold approximation and projection

(UMAP) visualization[55] and spatial visualization of E15.5-S1 sample, colored by ground truth domains

(Fig. 2d, e) and the spatial clusters identified based on different methods (Fig. 2f). As shown in Fig. 2e,

baseline methods like cisTopic and STAGATE could only identify the forebrain domain from others.

Although SCALE, SnapATAC2, SpaGCN and PAST roughly recognized the subdomains of central

nervous system, i.e., forebrain, hindbrain and midbrain, the low-dimensional representations acquired by

them are decentralized and scattered, impeding their performance for spatial clustering (Fig. 2c, f). In

contrast, PRESENT effectively distinguished domains including the forebrain, hindbrain, midbrain, facial

& bone, muscle & heart, and meningeal & peripheral nervous system (PNS) 2. Similar analytical results

were also observed on the other five samples (Supplementary Figs. 1-6), exhibiting the potential of

PRESENT in cross-modality representation and downstream visualization and domain identification.

Despite the superiority of PRESENT across the six mouse embryo samples, low sequencing depth

and signal-to-noise ratio stills significantly constrained the effectiveness of the model, illustrated in Fig.

2a, b and Supplementary Fig. 7. Our previous study has systematically demonstrated that integrating

external reference data can effectively address such issues in SRT data[17]. Inspired by this, we collected a

single-nucleus ATAC-seq mouse embryo dataset involving cells from three developmental stages,

including E12, E13 and E15, which was generated in conjunction with the spatial ATAC mouse embryo

data[10]. We utilized the single-nucleus ATAC-seq dataset (denoted as PRESENT-SN) and the other five

spatial ATAC samples (denoted as PRESENT-SP) as the reference data for PRESENT when analyzing

each mouse embryo sample (Methods). As shown in Fig. 2g and Supplementary Fig. 8, the introduction

of prior information contained in reference data enabled enhanced performance of PRESENT, especially

in samples with lower sequencing depth and signal-to-noise ratio, such as the E13.5-S1 and E12.5-S2

samples. This finding was further validated by the UMAP visualization of the two low-quality samples

(Fig. 2h and Supplementary Figs. 3 and 6). Specifically, PRESENT-SN & SP successfully identified

spatial domains including liver, limb, forebrain, hindbrain and midbrain, which other baseline methods or

even PRESENT failed to distinguish. For systematic evaluation, we further increased the nose level of

datasets by converting lower fragment counts to zero with growing dropout rates, according to a study

that models the dropout events in single-cell data[56]. As shown in Fig. 2i, PRESENT-SN & SP gained

greater advantages as the noise level increased, highlighting the improved benefits of integrating prior

information from reference data with the growing dropout rates in spatial ATAC-seq datasets.

Furthermore, we also carried out comprehensive experiments on SRT axolotl brain datasets generated

by Stereo-seq technique, consisting of six samples from six developmental stages, including stage 44,

stage 54, stage 57, juvenile, adult and metamorphosis[7]. The performance of PRESENT was benchmarked

with several recently developed SRT methods, involving SpaGCN[29], STAGATE[30], PAST[17] and

SpatialPCA[18]. As shown in Supplementary Figs. 9-15, experimental results demonstrated the consistent

superiority of PRESENT in representation, visualization and spatial domain identification. In addition,

utilizing an SRT regenerated axolotl brain data at 20 days post-injury as reference data[7], PRESENT-SP

again gained promoted performance through leveraging prior information contained in reference data

(Supplementary Fig. 9). In summary, PRESENT not only enables effective cross-modality representation

and spatial domain identification in both spatial ATAC-seq and SRT data, but also empowers the

integration of reference data from various sources to address the low sequencing depth and signal-to-

noise ratio of spatially resolved datasets.

PRESENT leverages complementary omics information from paired spatial RNA-ADT data

Next, we collected a spatial RNA-ADT human lymph node dataset generated by 10x Genomics Visium

technology, an in-house data from SpatialGlue[32]. This dataset simultaneously accessed the spatial

locations, transcriptomic and proteomic expression of spots and was manually annotated by the clinician

group according to the histological image[32] (Fig. 3a). We benchmarked PRESENT with SpatialGlue[32], a

representation learning method specially designed for spatial multi-omics data, and several methods

developed for single-cell RNA-ADT paired sequencing data, including totalVI[57], CiteFuse[58] and Seurat-

v4[59]. As shown in Fig. 3b, PRESENT again achieved the best performance on all the four metrics for

spatial domain identification. The UMAP and spatial visualization results were consistent with the

quantitative results (Fig. 3d, e, f). Specifically, CiteFuse and Seurat-v4 obtained unsatisfactory

visualization results, where domains such as medulla cords and sinuses scattered. Although SpatialGlue

and totalVI achieved relatively better results, they failed to recognize capsule from pericapsular adipose

tissue. In contrast, PRESENT successfully distinguished the pericapsular adipose tissue (cluster 8),

capsule (cluster 1), follicle (cluster 3) and cortex (cluster2 and cluster 10). We also directly applied

PRESENT to a single omics layer of the human lymph node dataset, thus leveraging expression

information from only a single omics layer, denoted as RNA-only and ADT-only. As shown in Fig. 3c,

PRESENT, which utilized multiple omics layers (RNA & ADT), achieved improved performance

compared to the model that only leveraged RNA or ADT data, highlighting the capabilities of PRESENT

in integrating complementary omics information.

As the largest secondary lymphoid organ in the body, the spleen plays a significant role in

hematopoiesis and the clearance of erythrocytes, contributing significantly to immune functions[60]. We

further applied PRESENT to a SPOTS mouse spleen dataset[13], which measured spatial transcriptomic and

proteomic profiles simultaneously, to decipher the functionality of spleen in the context of cellular

organizations. Through cross-modality representation and spatial clustering, PRESENT identified four

main spatial domains (Fig. 3g), denoted as the B cell-enriched domain, the T cell-enriched domain and

two macrophage-enriched subdomains, i.e., Mac1-enriched and Mac2-enriched domains, according to our

differentially expressed analysis (Fig. 3h, i). The differentially expressed analysis compared the protein

expression level of each domain with those of the rest domains based on the Wilcoxon rank-sum test[61]

(Fig. 3h), identifying protein markers distinguishing the B cell-enriched domain (CD3, CD4, CD8), the T

cell-enriched domain (CD19, B220, lgD) and the macrophage-enriched domain (F4-80, CD163, CD68),

consistent with the original study[13]. Specifically, the Mac1-enriched and Mac2-enriched domains had

nearly identical protein markers (Fig. 3h), which corresponded to the macrophage markers identified in

the original study[13], thereby making it hard to distinguish the two subdomains purely by ADT data.

Therefore, we further conducted differentially expressed analyses on the transcriptomic data between

different domains (Fig. 3i). Analytical results demonstrated that Mac1-enriched domain could be marked

by genes including *Hbb-bt, Hbb-a1, Rsad2, Hbb-bs, Car2* and *Slc4a1*, while Mac-2 enriched domain

could be recognized with *Igkc, Jchain, Igha, Ighm, Gm2a* and *Cox8a.* The RNA expression levels

between the Mac1 and Mac2-enriched domains were also directly compared and numerous significantly

differentially expressed genes (DEGs) were found (Fig. 3j). In conclusion, despite the similarity of the

two macrophage-enriched subdomains in ADT expression levels, the distinction of transcriptomic

patterns between them could still be identified by PRESENT, emphasizing the ability of PRESENT in

leveraging complementary proteomic and transcriptomic information for effective domain identification.

To further decipher the functional distinction between the two macrophage-enriched subdomains, we

conducted Gene Ontology (GO) enrichment analysis based on the identified DEGs (Methods).

Specifically, for the Mac1-enriched domain, we enriched biological processes including erythrocyte

development, erythrocyte differentiation, erythrocyte homeostasis, myeloid cell development, myeloid

cell differentiation and myeloid cell homeostasis with false discovery rates (FDRs) of 1.29e-8, 5.48e-6,

6.70e-6, 3.59e-7, 8.71e-5 and 2.06e-5, respectively (Fig. 3k and Supplementary Table 1). Macrophages

are myeloid lineage cells which derive from bone marrow[62], reasonably explaining the enrichment of

development, differentiation and homeostasis of myeloid cell in the Mac1-enriched domain. In addition,

previous studies have stressed the importance of macrophages in the entire life of erythrocytes by

providing necessary factors for the survival, proliferation and development of erythrocytes[63, 64], agreeing

with the development, differentiation and homeostasis of erythrocyte found in the Mac1-enriched domain.

In comparison, the Mac2-enriched domain found biological processes including phagocytosis &

engulfment, plasma membrane invagination, membrane invagination, phagocytosis, humoral immune

response and lymphocyte mediated immunity with FDRs of 7.11e-5, 7.11e-5, 7.11e-5, 4.34e-4, 5.58e-4

and 5.94e-4, respectively (Fig. 3k and Supplementary Table 2). Phagocytosis and other related processes,

also called the process of 'cell eating', are commonly considered as classical functions of macrophages[65,

[66], validating the enrichment of macrophages in the Mac2-enriched domain. On the other hand, compared

with the Mac1-enriched domain, the Mac2-enriched domain was spatially close to the B cell-enriched and

T cell-enriched domains (Fig. 3g), thereby explaining the enrichment of lymphocyte mediated immunity

process[67]. In summary, PRESENT effectively integrates complementary omics information in spatial

RNA-ADT data, thereby facilitating accurate spatial domain identification and systematic exploration of

biological processes in the identified domains.

PRESENT extracts cross-modality representations from parallel spatial RNA-ATAC datasets

We further collected the MISAR-seq mouse brain datasets[14], which simultaneously measured the

epigenomics and transcriptomics, i.e., paired spatial RNA-ATAC profiles, of four mouse brain samples

from four developmental stages, involving E11.0, E13.5, E15.5 and E18.5. Compared with spatial-RNA-

ADT data, spatial RNA-ATAC datasets suffer from higher dimension and sparsity. We firstly compared

the performance of PRESENT in spatial domain identification with the spatial multi-omics method

SpatialGlue[32] and six methods designed for the representation of single-cell multi-omics data, including

MOFA+[68], scAI[69], scMDC[70], scMVP[19], Seurat-v4[59] and SnapATAC2[51]. As shown in Fig. 4a, PRESENT

achieved overall the best spatial clustering performance, illustrating the superiority of PRESENT in

extracting spatial multi-omics information into effective cross-modality embeddings for accurate spatial

domain identification. We also used PRESENT to obtain latent representations based on only ATAC data

or RNA data, denoted as ATAC-only and RNA-only respectively, which resulted in decreased

performance, highlighting the importance of integrating complementary multi-omics information in

deciphering biological variations in the context of cellular organizations (Fig. 4b).

Taking the E18.5 mouse brain sample with the most complex spatial structures as an example (Fig.

4c), we intuitively visualized the latent embeddings learned by PRESENT using heatmap (Fig. 4d). The

spots from different spatial domains could be successfully distinguished via the combination of several

latent features or even a single feature in the latent representation space, including large domains such as

cartilage 1-4, mantle zone of dorsal pallium (DPallm), ventricular zone of dorsal pallium (DPallv),

diencephalon & hindbrain and subpallium 1, and small domains like midbrain and thalamus. We further

demonstrated the UMAP visualization of latent representations obtained by different methods on the

E18.5 mouse brain sample (Fig. 4e). Baseline methods, including MOFA+, scAI, scMDC, scMVP and

Seurat-v4, could only roughly separate large domains such as DPallm, DPallv, diencephalon & hindbrain

and subpallium 1. SnapATAC2 could additionally recognize the cartilage 1 but still failed to distinguish

others. In comparison, SpatialGlue and PRESENT not only identified large spatial domains such as

diencephalon & hindbrain, subpallium 1, DPallm, DPallv, cartilage 1-4, but also accurately distinguished

two rare spatial domains, i.e., midbrain and thalamus. In addition, the visualization of spatial clusters

identified based on different methods consistently demonstrated the excellent domain identification

performance of PRESENT, shown in Fig. 4f, g. For example, PRESENT also successfully deciphered

most large domains, including DPallm (cluster 5), DPallv (cluster 6), subpallium 1 (cluster 11), cartilage

1 (cluster 1) and diencephalon & hindbrain (cluster 7), and the two smaller domains, namely midbrain

(cluster 9) and thalamus (cluster 13).

The UMAP and spatial visualization of other three samples of the MISAR-seq mouse brain dataset[14]

achieved similar results (Supplementary Figs. 16-19) and further highlighted the capabilities of

PRESENT in integrating complementary transcriptomic and epigenomic omics information for effective

cross-modality representation. Specifically, PRESENT utilizing only the transcriptomic data of E11.0

brain sample, denoted as RNA-only, successfully captured the difference between the muscle and the

cartilage 2 domain, while the model based solely on epigenomic data, depicted as ATAC-only, failed to do

so (red circles in Fig. 4h). On the other hand, compared with the failure of PRESENT using only

transcriptomic data (RNA-only) in the E15.5 brain sample, the model based on the epigenomic data

(ATAC-only) contributed more to the differentiation between cartilage 2, cartilage 3 and mesenchyme

domains (blue circles in Fig. 4i). Therefore, purely utilizing a single omics layer resulted in biased

interpretation of biological signals in spatial domains. In comparison, when taking both omics layers as

input (RNA & ATAC), PRESENT successfully captured variations among those domains in both E11.0

and E15.5 mouse brain samples (Fig. 4h, i), highlighting the capabilities of PRESENT in leveraging

spatial and complementary transcriptomic & epigenomic information for cross-modality representation

and further facilitating the identification of spatial domains from a systematic perspective.

PRESENT-BC enables effective integration of horizontal and vertical spatially resolved samples

Compared with cross-modality representation on a single spatial sample, integrated analysis of multiple

samples enables the identification of hierarchical structures of cellular organizations across different

dissected areas (horizontally adjacent tissue samples) or developmental stages (vertically serial tissue

samples)[23]. Therefore, building upon the cross-modality representation framework PRESENT, we further

extended PRESENT-BC for the multi-sample integration of horizontal and vertical samples generated by

spatially resolved technologies (Methods, Fig. 1b).

We firstly collected two horizontal SRT mouse brain samples generated by the 10x Genomics Visium

platform, which involves a sagittal anterior section and a sagittal posterior section. We applied two SRT

integration methods, namely STAligner[25] and GraphST[23], and PRESENT-BC to carry out integrated

analysis of these two adjacent brain samples and utilized the anatomic annotation of the sagittal region in

postnatal day 56 (P56) mouse brain provided by Allen Reference Atlas[71] to intuitively compare the

performance of these methods (Fig. 5a). As shown in Fig. 5b-d, analytical results illustrated significant

improvements of PRESENT-BC in shared and sample-specific domain identification compared with

STAligner and GraphST. For example, PRESENT-BC captured the classical hierarchical structures, i.e.,

from outer layer 1 to inner layer 6, of dorsal pallium in both sagittal anterior and posterior sections, as

illustrated in the red rectangle of Fig. 5d. In addition, PRESENT-BC successfully distinguished the

typical cortex Ammonis (CA, cluster 18) and dentate gyrus (DG, cluster 25) structures simultaneously in

the hippocampus (black frames in Fig. 5d), which have been proved to be associated with mood, anxiety

and personality disorders caused by various factors[72]. We further conducted differentially expressed

analysis for domain CA and domain DG, acquiring marker genes including *C1ql2* and *Hpca*, respectively

(Supplementary Figs. 20 and 21). According to previous studies, *C1ql2* serves as a marker gene for

dentate gyrus granule cells (DGCs) enriched in the DG structure[73], while *Hpca* is the encoding gene for a

high-affinity calcium-binding protein restricted to the central nervous system, abundant in pyramidal cells

of the hippocampus CA region[74].

Furthermore, PRESENT-BC also exhibited great potential in capturing sample-specific functional

domains. For instance, PRESENT-BC successfully revealed the sophisticated substructures of the

cerebellar vermis, which only exists in the sagittal posterior section of mouse brain and plays an

important role in coordinating movements of the central body such as trunk, head, and proximal limbs.

Specifically, PRESENT-BC successfully detected the molecular layer (cluster 3), Purkinje cell layer

(cluster 22) and internal granular layer (cluster 17) of cerebellar vermis cortex and the white matter

(cluster 20) of cerebellar vermis. The differentially expressed genes in the corresponding domains

(Supplementary Figs. 22-25), including *S100b*[75], *Car8*[76], *Atp1a1*[77] and *Mbp*[78], have also been reported in

previous studies, further illustrating the effectiveness of PRESENT-BC in uncovering biological

implications in identified spatial domains. For example, *Car8* and *Atp1a1*, the mutation of which cause

the motor coordination defect and dominant Charcot-Marie-Tooth type 2, were proved to be highly

expressed in the granular layers and Purkinje cells in recent studies[76, 77], respectively.

On the other hand, we also quantitatively evaluated the performance of PRESENT-BC in the

integrative analysis of vertically serial tissue samples involving mouse embryo slices from three

developmental stages (E15.5, E13.5 and E12.5) generated by spatial ATAC technique[10]. Due to the lack of

methods specially designed for spatial ATAC data, we compared PRESENT-BC with six integration

methods developed for single-cell data (MNN[79], Scanorama[39], Harmony[38], SCALEX[80], scBasset-BC[81]

and SnapATAC2[51]) and two recently proposed SRT methods (STAligner[25] and GraphST[23]). According to

a recent benchmarking framework for single-cell omics data integration[82], we simultaneously accessed

the batch effect removal and biological variance conservation capabilities of these methods, with seven

metrics for each aspect (Methods). As shown in Fig. 5e, PRESENT-BC achieved the best overall score for

multi-sample integration, ranking first in preserving biological variance and second in removing batch

effects among all the integration methods. The UMAP visualization of spot embeddings across different

developmental stages also demonstrated that compared with other methods, PRESENT-BC gained better

performance in aggregating spots from the same domain but in different batches (Supplementary Fig. 26).

Furthermore, we performed joint spatial clustering based on the joint embeddings obtained by different

methods. As shown in Fig. 5f, g, Scanorama and STAligner could hardly distinguish different spatial

domains across different developmental stages. MNN, Harmony, SCALEX and SnapATAC2 provided

improved results by roughly recognizing forebrain and liver in E12.5 and E13.5 stages, but they still

failed in other domains. In comparison, PRESENT-BC not only identified forebrain (cluster 1) and liver

(cluster 5) in all the three stages, but also gained superior results in domains including hindbrain (cluster

2) and midbrain (cluster 8). In summary, these experimental results have highlighted the superior

performance of PRESENT-BC in integrative analysis of both horizontally adjacent and vertically serial

spatially resolved samples generated by various spatial omics technologies.

PRESENT-BC captures biological signals while eliminating batch effects in spatial RNA-ATAC data

The MISAR-seq mouse brain dataset simultaneously measured the transcriptomic and epigenomic

profiles of developmental brain samples from four stages, including E11.0, E13.5, E15.5 and E18.5,

providing a multi-omics landscape of the developmental mouse brain in the context of cellular

organizations[14]. During the development of mouse brain, some functional structures may be specific to

certain developmental stages (e.g., primary brain), while others may persist throughout the entire

developmental process (e.g., midbrain and muscle). Therefore, the key challenges of integrating the

MISAR-seq mouse brain samples included not only the incorporation of multi-omics information across

multiple samples, but also the conservation of shared and stage-specific biological variations.

Due to the lack of multi-sample integration methods specially designed for spatial multi-omics data,

we compared the performance of PRESENT-BC with six methods developed for single-cell data (MNN[79],

Scanorama[39], Harmony[38], SCALEX[80], scBasset-BC[81] and SnapATAC2[51]) and three recently proposed

SRT methods (STAligner[25], Stitch3D[31] and GraphST[23]). MNN, Scanorama, Harmony and SCALEX could

be applied to batch integration of both RNA-seq and ATAC-seq data, denoted as RNA-* and ATAC-*,

respectively, where * represents the different methods. scBasset-BC was featured as a sequence-based

method specially designed for single-cell ATAC-seq data, while STAligner, Stitch3D and GraphST were

intended for integrative analysis of SRT data. SnapATAC2 has provided guidance for incorporating both

transcriptomic and epigenomic information for latent representation and integrated analysis. These

methods were systematically benchmarked using 14 metrics divided into two aspects, i.e., batch effect

removal and biological variance conservation. As shown in Fig. 6a, scBasset-BC and SnapATAC2

provided satisfactory performance through incorporating sequence and multi-omics information,

respectively. PRESENT-BC, which simultaneously integrated spatial dependency and multi-omics

information of all the samples, achieved the best performance in both batch effect removal and biological

variance conservation among all the 14 methods, thereby illustrating the superiority in multi-sample

integration for spatial RNA-ATAC data.

The UMAP visualization of joint embeddings colored by batch indices (Fig. 6b and Supplementary

Fig. 27a) and ground truth domain labels (Fig. 6c and Supplementary Fig. 27b) also validated the

quantitative results. Specifically, PRESENT-BC successfully fused spots from different developmental

stages (red boxes in Fig. 6b) while maintaining the biologically distinct patterns of DPallm, DPallv,

subpallium 1 & 2, thalamus and muscle (red boxes in Fig. 6c). Specifically, spots of muscle domain

persisted throughout the entire life of mouse brain development, while those of subpallium 1&2, DPallm,

DPallv and thalamus only existed in specific stages. The success of PRESENT-BC in distinguishing these

domains illustrated its capabilities in capturing shared and stage-specific biological variance while

eliminating batch effects. Furthermore, we conducted joint spatial clustering across the four stages based

on the integrated representations obtained by different methods, as shown in Fig. 6d and Supplementary

Fig. 28. Except for the scBasset-BC and SnapATAC2, other baseline methods provide unreliable

clustering results. In contrast, PRESENT-BC successfully detected not only large domains like DPallm

(cluster 7), but also small domains including thalamus (cluster 16) and muscle (cluster 11), which

PRESENT failed to identify through individual analysis on a single sample from E18.5 stage

(Supplementary Fig. 19), highlighting the advantages of integrative analysis of multiple samples. In

summary, these results emphasize that PRESENT-BC enables the conservation of biological signals while

removing batch effects in the integrated analysis of spatial RNA-ATAC samples.

Discussion

Recent breakthroughs in spatial omics sequencing technologies have enabled the simultaneous acquisition

of spatial coordinate information and the regulatory status of the biomolecules within cells, such as

chromatin regions, genes, and proteins, accumulating massive spatial omics data in repositories and

databases[15, 16]. In this article, we proposed PRESENT, an effective and scalable contrastive learning

framework, to address the cross-modality representation of spatially resolved omics data, including

spatial epigenomic, transcriptomic and multi-omics data. To model the distinct patterns and distributions

of different omics layers, we designed omics-specific encoders and decoders for different omics data,

where the transcriptomic and epigenomic decoders model the counts via a zero-inflation negative

binomial (ZINB) distribution[43] and a zero-inflation Poisson (ZIP) distribution[44], respectively (Methods).

In addition, a cross-omics alignment module built on a contrastive learning loss function[42] and graph

attention networks[40] (GAT) and were also introduced to capture the spatial dependency and

complementary multi-omics information (Methods). We further extended the cross-modality

representation framework to the multi-sample integration of spatial omics data, addressing the key

challenges including the inconsistency of spatial coordinates between samples as well as the conservation

of biological variance while eliminating batch effects. Specifically, we adopted an inter-batch alignment

loss[45] and a batch-adversarial learning strategy[46] to eliminate redundant batch effects as well as an intra-

batch preserving loss to maintain biological signals simultaneously. As part of the validation of the model

design, we conducted a comprehensive ablation study to illustrate the effectiveness of these modules, as

shown in Supplementary Figs. 29 and 30.

Throughout comprehensive experiments on different species involving human, mouse and axolotl,

we highlight the superior performance of PRESENT in cross-modality representation of spatial omics

data generated by various spatial sequencing technologies, encompassing spatial ATAC-seq[10], spatial

transcriptomics[7], as well as the spatial RNA-Protein[11, 12] and RNA-ATAC[14] co-profiling techniques.

Specifically, PRESENT effectively integrates the spatial dependency and diverse omics information

contained in spatial data generated by spatial ATAC-seq[10] or spatial transcriptomics[7] techniques (Fig. 2a-

f). It also shows great potential for leveraging various reference data to address issues related to low

sequencing depth and signal-to-noise ratio in spatial omics data (Fig. 2g-i). Additionally, PRESENT

exhibits powerful capabilities in incorporating spatial and complementary multi-omics information,

thereby facilitating explorative analysis in spatial multi-omics data (Fig. 3, 4). More importantly, the

extended multi-sample integration version, namely PRESENT-BC, facilitates the integrative analysis of

multiple spatially resolved samples across distinct dissection areas (Fig. 5a-d) and developmental stages

(Fig. 5e-g, 6), thereby promoting the detection of hierarchical structures and biological functions within

cellular organizations from a spatiotemporal perspective.

Nevertheless, several avenues remain for further improvement of PRESENT. First, although

PRESENT can anchor on spots to analyze a single sample with paired omics sequencing data and

PRESENT-BC is capable of anchoring on shared genomic features to integrate multiple samples, the

framework still struggles with the integration of unpaired samples with distinct features[83], such as the

integration of two samples generated by spatial ATAC-seq and SRT techniques, respectively. Due to

technological and cost limitations, the number of unpaired spatial samples significantly outweighs the

number of paired spatial samples[83], highlighting the importance of unpaired integration. Second, it is also

inspiring to extend the application scenarios of PRESENT to the integration of single-cell and spatial

omics data, which lack crucial spatial information and single-cell resolution, respectively. Third, in

addition to the integrative analysis of multiple samples in the latent space, the alignment of spatial

coordinates in physical space also plays an important role in joint analysis of multiple samples. Moreover,

advanced spatial sequencing technologies like 10x Genomics Visium additionally provide high-resolution

histology image of dissected samples, providing additional morphological features which could be further

leveraged by PRESENT for enhanced analysis of spatially resolved samples.

Methods

Cross-modality representation with PRESENT

PRESENT is constructed upon a multi-view autoencoder framework for representation learning of spots

sequenced by spatial omics technologies, adeptly incorporating cross-modality information,

encompassing spatial coordinate data, epigenomic data generated by the assay for transposase-accessible

chromatin with sequencing (ATAC), transcriptomic data measured by RNA sequencing (RNA), and

proteomic data sequenced by antibody-derived tags (ADT), into a low-dimensional feature matrix. Note

that while the spatial coordinate matrix is a necessary input, the three omics layers are not all mandatory.

In other words, PRESENT can take an arbitrary combination of count matrices from the three omics

layers, including ATAC, RNA and ADT data, as input. In the following sections, taking all the three omics

layers and spatial coordinates as input, we elaborate the architecture of the PRESENT, which consists of

omics-specific encoders, a cross-omics alignment module, and omics-specific decoders (Fig. 1a).

Multi-modality input of PRESENT. The input of PRESENT is a spatial coordinate matrix $M \in R^{n \times 2}$

and three optional fragment count matrices, namely the ATAC data $X^{ATAC} \in R^{n \times r}$, RNA data $X^{RNA} \in$

$R^{n \times g}$ and ADT data $X^{ADT} \in R^{n \times p}$. Here n denotes the number of spots, r the number of input regions,

g the number of highly variable genes and p the number of input proteins. Before feeding the spatial and

omics input into the omics-specific encoders, we transform the spatial coordinate matrix M into an

undirected neighborhood graph G through adding edge connections between all spots and their k nearest

neighbors, where k is set to 6 by default. Afterwards, we apply term frequency-inverse document

frequency (TF-IDF) normalization to the ATAC data[84-87], library size normalization followed by log-

transformation to RNA data[17, 29, 30] and the centered log ratio (CLR) transformation to ADT data [36, 88, 89],

obtaining normalized omics data $\widetilde{X}^{ATAC} \in R^{n \times r}$, $\widetilde{X}^{RNA} \in R^{n \times g}$ and $\widetilde{X}^{ADT} \in R^{n \times p}$, respectively.

Omics-specific encoder. Graph neural networks, such as graph attention networks[40] (GAT), have

showcased successful application in integrating spatial and transcriptomic information for spatial

transcriptomics[17, 23, 29, 30]. On the other hand, research has proved that introducing weight uncertainty into

models through Bayesian neural networks[41] (BNN) improves generalization in nonlinear regression

problems[41] and offers an extension for leveraging prior information from existing sequencing data[17].

Inspired by these studies, we design an omics-specific encoder, consisting of a GAT module and a BNN

module, to capture spatially organized and biologically regulatory signals embedded in each omics layer.

In the following two subsections, taking the ATAC omics-specific encoder as an example, we delineate

the detailed design of GAT and BNN module.

Graph attention network. Given the input undirected neighborhood graph $\boldsymbol{G}$ and the normalized

ATAC matrix $\widetilde{\boldsymbol{X}}^{ATAC} \in R^{n \times r}$, we denote the normalized accessibility vector of spot i in $\widetilde{\boldsymbol{X}}^{ATAC}$ as

$\widetilde{x}_i^{ATAC} \in R^r$ and the spatial neighborhood of spot i as $\mathcal{N}_i$. Initially, we acquire the embedding from the

first hidden layer of spot i, i.e., $\boldsymbol{h}_i^1 \in R^{d_1}$ using a fully connected network (FCN):

$$\boldsymbol{h}_i^1 = \sigma(\boldsymbol{W_1}\widetilde{x}_i^{ATAC} + \boldsymbol{b_1})$$

523   where d_1 represents dimension of the first hidden layer, σ indicates the combination of layer

524   normalization and exponential linear unit (ELU) activation function, and $\boldsymbol{W_1} \in R^{d_1 \times r}$ and $\boldsymbol{b_1} \in R^{d_1}$

525   indicate the weight and bias parameters of the FCN, respectively. Afterwards, The second hidden-layer

526   embedding of spot i, namely $\boldsymbol{h}_i^2 \in R^{d_2}$, is obtained based on a graph attention layer[40], depicted as:

$$\boldsymbol{h}_i^2 = \sigma(\sum_{j \in \mathcal{N}_i} \alpha_{ij}\boldsymbol{W_2}\boldsymbol{h}_j^1)$$

$$\alpha_{ij} = \frac{\sigma_{LeakyReLU}\left(\boldsymbol{a}^T\left(\boldsymbol{W_2}\boldsymbol{h}_i^1 \oplus \boldsymbol{W_2}\boldsymbol{h}_j^1\right)\right)}{\sum_{k \in \mathcal{N}_i} \sigma_{LeakyReLU}\left(\boldsymbol{a}^T\left(\boldsymbol{W_2}\boldsymbol{h}_i^1 \oplus \boldsymbol{W_2}\boldsymbol{h}_k^1\right)\right)}$$

where α_{ij} is the attention score between spot i and j, d_2 denotes the dimension of the second hidden

layer, $\boldsymbol{W_2} \in R^{d_1 \times d_2}$ and $\boldsymbol{a} \in R^{2d_2}$ are learnable parameters of the graph attention layer, $\sigma_{LeakyReLU}$

denotes the LeakyReLU activation function with a negative slope of 0.2, and $\oplus$ refers to the

concatenation function. Finally, h_i^2 undergoes transformation by another graph attention layer to obtain

$h_i^{gat} \in R^{d_{gat}}$ as the output, where d_{gat} denotes the output dimension of GAT module.

Bayesian neural network. According to our previous work, BNN module provides an opportunity to

leverage biological prior information from various sequencing data to improve the analytical performance

in spatial transcriptomic data[17]. Inspired by this, PRESENT includes the BNN module, which assumes

that the parameters of weight matrix, denoted as $W \in R^{d_{bnn} \times r}$, follow Gaussian distributions.

Specifically, each element of W is governed by a Gaussian distribution parameterized by the

corresponding elements in the mean matrix μ^b and variance matrix σ^{b^2}, depicted as:

$$W_{ij} \sim \mathcal{N}\left(\mu_{ij}^b, \sigma_{ij}^{b\,2}\right)$$

In addition, W_{ij} is also governed by a Gaussian prior parameterized by mean μ_{ij}^p and variance $\sigma_{ij}^{p\,2}$.

Inheriting our previous work[17], we calculate the PCA projection weight matrix of normalized ATAC

matrix $\widetilde{X}^{ATAC}$ as the mean matrix μ^p and utilize the standard deviation of all the elements in mean

matrix μ^p to construct the standard deviation matrix σ^p. The distance between the prior and the

distributions of the parameters in BNN module is restricted with Kullback-Leibler divergence (KLD):

$$\mathcal{L}_{BNN} = KLD(\mathcal{N}\left(\mu^b, \sigma^{b^2}\right), \mathcal{N}(\mu^p, \sigma^{p^2}))$$

It is notable that the parameters of prior Gaussian distributions, i.e., the mean matrix μ^p and the variance

matrix $\sigma_{ij}^{p\,2}$, can also be constructed with reference ATAC data, enabling PRESENT to incorporate prior

information from existing ATAC-seq data generated by various technologies, including single-cell ATAC-

seq or spatial ATAC-seq. The unification of peak sets between reference ATAC-seq data and target ATAC-

seq data is conducted according to the merging objects vignette provided by the Signac package[90].

Therefore, the output embeddings of spot i from BNN module, i.e., $h_i^{bnn} \in R^{d_{bnn}}$, is denoted as:

$$h_i^{bnn} = W\widetilde{x}_i^{ATAC} = (\mu^b + \eta \odot \sigma^b)\widetilde{x}_i^{ATAC}$$

where d_{bnn} denotes the output dimension of BNN module, $\boldsymbol{\eta}$ is the standard Gaussian noise matrix and

$\odot$ denotes element-wise product.

Finally, the output embedding of spot i from the ATAC omics-specific encoder, denoted as $\boldsymbol{h}_i^{ATAC} \in$

$R^{d_{ATAC}}$, is the concatenation of output embeddings from the GAT and BNN module:

$$\boldsymbol{h}_i^{ATAC} = \boldsymbol{h}_i^{gat} \oplus \boldsymbol{h}_i^{bnn}$$

where $d_{ATAC} = d_{gat} + d_{bnn}$ is the output dimension of omics-specific encoder for ATAC data.

Cross-omics alignment module. Based on the omics-specific encoders, the embeddings of different

omics layers are separately extracted. However, accurately deciphering sophisticated gene regulation

process requires integrating the status of biological macromolecules in cells from different omics

perspectives[34-37]. Inspired by a recent study that achieved multi-modal entity alignment through

contrastive learning[42], we designed a cross-omics alignment module to model the complex regulatory

interactions between different omics layers.

Given the omics-specific embeddings of spot i obtained by omics-specific encoders, including

$\boldsymbol{h}_i^{ATAC} \in R^{d_{ATAC}}$ for ATAC, $\boldsymbol{h}_i^{RNA} \in R^{d_{RNA}}$ for RNA and $\boldsymbol{h}_i^{ADT} \in R^{d_{ADT}}$ for ADT, we utilized a multi-

layer perception (MLP) module to fuse the embeddings from different omics layers to acquire a cross-

omics representation of spot i, denoted as:

$$\boldsymbol{e}_i = f_{MLP}(\boldsymbol{h}_i^{ATAC} \oplus \boldsymbol{h}_i^{RNA} \oplus \boldsymbol{h}_i^{ADT})$$

where $\boldsymbol{e}_i \in R^d$ represents the final cross-omics representation of spot i, f_{MLP} denotes the transformation

function of the MLP module, d is the latent dimension of PRESENT, and d_{ATAC}, d_{RNA} and d_{ADT} refers

to the output dimensions of omics-specific encoders for ATAC, RNA and ADT data, respectively. The

MLP module is composed of three FCNs, the first two of which are followed by a layer normalization and

an ELU activation function. However, only the MLP fusion module cannot perfectly incorporate

complementary omics-specific knowledge from different omics layers into the final cross-omics

embeddings [42], which is also illustrated in our ablation study shown in Supplementary Fig. 29. Therefore,

we design an inter-omics alignment (IOA) loss to minimize the multi-omics directional KLD over the

output distributions between the joint embeddings and other omics-specific embeddings:

580
$$\mathcal{L}_{IOA} = \frac{1}{M}\left[\frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{ATAC})\right) + \frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{RNA})\right)\right.$$

581
$$\left. + \frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{ADT})\right)\right]$$

582 where M represents the number of omics layers, $\boldsymbol{e}, \boldsymbol{h}^{ATAC}, \boldsymbol{h}^{RNA}$ and $\boldsymbol{h}^{ADT}$ denotes the joint, ATAC,

583 RNA and ADT embeddings respectively, and $q_{ed}^{(i)}(*)$ represents the embedding distribution of spot i

584 over other spots, defined as (taking $\boldsymbol{e}$ as an example):

585
$$q_{ed}^{(i)}(\boldsymbol{e}) = \left(\frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_1})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}, \frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_2})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}, \cdots, \frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_n})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}\right)$$

586
$$\delta_{ed}(\boldsymbol{u}, \boldsymbol{v}) = exp(\frac{\boldsymbol{u}^T\boldsymbol{v}}{\tau})$$

where τ is a temperature parameter and set to 0.1 according to the original study[42].

Omics-specific decoder. The patterns and distributions of sequencing data vary significantly across

different omics layers, revealing complementary biologically regulatory information[34-37]. For example,

ADT sequencing data, which focuses on dozens of surface proteins, provides intuitive and reliable

quantification of cellular activities with low dimensionality, few dropout events and low sparsity[36]. In

contrast, RNA and ATAC data, which measure the expression levels of genes and accessibility scores of

genomic regions respectively, suffer from high dropout rates, dimensionality, and sparsity[91]. Therefore,

we designed omics-specific decoder for each modality to address this challenge.

Inspired by a recently published study which modeled the fragment counts of single-cell ATAC-seq

data with Poisson distribution[44], we fit the fragment counts of spatial ATAC-seq data $\boldsymbol{X}^{ATAC} \in R^{n \times r}$ with

the zero-inflation Poisson (ZIP) distribution to accommodate the lower capture rate of spatial

technologies[10, 14]. Taking the fragment count vector of spot i, i.e., $\boldsymbol{x_i}^{ATAC}$, as an example, the omics-

specific decoder of ATAC data as well as the probability model is described as:

$$\boldsymbol{\pi_i} = \sigma_{sigmoid}\left(f_{MLP}^{ATAC,\pi}(\boldsymbol{e_i})\right) \quad \boldsymbol{\rho_i} = f_{MLP}^{ATAC,\rho}(\boldsymbol{e_i}) \quad \hat{\boldsymbol{x}}_i^c = f_{MLP}^{ATAC,x}(\boldsymbol{e_i})$$

$$\boldsymbol{\omega_i} = softmax(\boldsymbol{\rho_i} + \boldsymbol{r^c})$$

$$\boldsymbol{\lambda_i} = l_i \cdot \boldsymbol{\omega_i}$$

where $\sigma_{sigmoid}$ represents the sigmoid activation function, $\boldsymbol{r^c}$ is designed to capture region specific bias

such as peak length, $l_i = \sum_{j=1}^{r} x_{ij}^c$ refers to the total fragment counts of spot i, $f_{MLP}^{ATAC,\pi}(*)$, $f_{MLP}^{ATAC,\rho}(*)$

and $f_{MLP}^{ATAC,x}(*)$ are three output layers of $f_{MLP}^{ATAC}(*)$, which denotes the MLP module of ATAC omics-

specific decoder. Specifically, $f_{MLP}^{ATAC,\pi}(*)$ and $f_{MLP}^{ATAC,\rho}(*)$ output parameters of the ZIP probability

model, while $f_{MLP}^{ATAC,x}(*)$ outputs reconstructed normalized profiles of ATAC data. Then for each region

$j = 1,2,\dots,r$, the fragment count of i-th spot x_{ij}^{ATAC} satisfies:

$$x_{ij}^{ATAC} \sim ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right)$$

$$ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right) = \pi_{ij}\delta_0\left(x_{ij}^{ATAC}\right) + (1 - \pi_{ij})Poisson(x_{ij}^{ATAC}; \lambda_{ij})$$

$$\delta_0\left(x_{ij}^{ATAC}\right) = \begin{cases} 1 \text{ if } x_{ij}^{ATAC} \text{ equals } 0 \\ 0 \text{ otherwise} \end{cases} \quad Poisson(x_{ij}^{ATAC}; \lambda_{ij}) = \frac{\lambda_{ij}^{x_{ij}^{ATAC}} e^{-\lambda_{ij}}}{x_{ij}^{ATAC}!}$$

Therefore, the negative log likelihood (NLL) loss function of the ZIP probability model is:

$$\mathcal{L}_{NLL}^{ATAC} = \frac{1}{nr}\sum_{i=1}^{n}\sum_{j=1}^{r}\left[-log\,ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right) + \gamma\pi_{ij}^2\right]$$

where the $\gamma\pi_{ij}^2$ refers to a regularization term and γ is set to 0.5 by default. The reconstruction loss of

ATAC omics layer is measured with mean squared error (MSE):

$$\mathcal{L}_{MSE}^{ATAC} = \frac{1}{nr}\sum_{i=1}^{n}\sum_{j=1}^{r}\left(\hat{x}_{ij}^{ATAC} - \tilde{x}_{ij}^{ATAC}\right)^2$$

Similarly, we also model the counts of RNA data $\boldsymbol{X^{RNA}} \in R^{n \times g}$ using the zero-inflation negative

binomial (ZINB) distribution according to a single-cell RNA-seq analytical research[43], denoted as:

$$\boldsymbol{\pi_i} = \sigma_{sigmoid}\left(f_{MLP}^{RNA,\pi}(\boldsymbol{e_i})\right) \quad \boldsymbol{\mu_i} = \sigma_{softplus}\left(f_{MLP}^{RNA,\mu}(\boldsymbol{e_i})\right)$$

$$\boldsymbol{\theta_i} = \sigma_{softplus}\left(f_{MLP}^{RNA,\theta}(\boldsymbol{e_i})\right) \quad \hat{\boldsymbol{x}}_i^r = f_{MLP}^{RNA,x}(\boldsymbol{e_i})$$

where $\sigma_{softplus}$ refers to the softplus activation function, $f_{MLP}^{RNA,\pi}(*)$, $f_{MLP}^{RNA,\mu}(*)$, $f_{MLP}^{RNA,\theta}(*)$ and

$f_{MLP}^{RNA,x}(*)$ are four output layers of f_{MLP}^{RNA}, the MLP module of the RNA omics-specific decoder. Then the

count of spot i on gene j satisfies the following model:

$$x_{ij}^{RNA} \sim ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right)$$

$$ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right) = \pi_{ij}\delta_0\left(x_{ij}^{RNA}\right) + (1 - \pi_{ij})NB(x_{ij}^{RNA}; \mu_{ij}, \theta_{ij})$$

$$NB\left(x_{ij}^{RNA}; \mu_{ij}, \theta_{ij}\right) = \frac{\Gamma\left(x_{ij}^{RNA} + \theta_{ij}\right)}{\Gamma(\theta_{ij})} \left(\frac{\theta_{ij}}{\theta_{ij} + \mu_{ij}}\right)^{\theta_{ij}} \left(\frac{\mu_{ij}}{\theta_{ij} + \mu_{ij}}\right)^{x_{ij}^{RNA}}$$

627   The NLL loss function of the ZINB probability model for RNA data can be denoted as:

$$\mathcal{L}_{NLL}^{RNA} = \frac{1}{ng} \sum_{i=1}^{n} \sum_{j=1}^{g} \left[-\log ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right) + \gamma \pi_{ij}^2 \right]$$

629   where the regularization parameter γ is also set to 0.5 by default. In addition, the reconstruction loss of

630   RNA omics layer is depicted as:

$$\mathcal{L}_{MSE}^{RNA} = \frac{1}{ng} \sum_{i=1}^{n} \sum_{j=1}^{g} \left(\hat{x}_{ij}^{RNA} - \tilde{x}_{ij}^{RNA} \right)^2$$

632   Compared with ATAC and RNA data, lower dropout rates, dimensionality and sparsity make the

633   analysis of ADT data less challenging. Therefore, the ADT omics-specific decoder is built on the MLP

634   module with one output layer to reconstruct the normalized expression level of proteins, denoted as:

$$\hat{x}_i^{ADT} = f_{MLP}^{ADT}(e_i)$$

$$\mathcal{L}_{MSE}^{ADT} = \frac{1}{np} \sum_{i=1}^{n} \sum_{j=1}^{p} \left(\hat{x}_{ij}^{ADT} - \tilde{x}_{ij}^{ADT} \right)^2$$

637   Therefore, the overall NLL loss and reconstruction loss of PRESENT are summarized as:

$$\mathcal{L}_{NLL} = \mathcal{L}_{NLL}^{ATAC} + \mathcal{L}_{NLL}^{RNA}$$

$$\mathcal{L}_{MSE} = \mathcal{L}_{MSE}^{ATAC} + \mathcal{L}_{MSE}^{RNA} + \mathcal{L}_{MSE}^{ADT}$$

641   **Overall loss functions of PRESENT.** Based on the above descriptions, the overall loss function of

642   PRESENT consists of four parts, including the loss of BNN module, the inter-omics alignment loss, the

643   negative log likelihood loss of ZIP and ZINB probability model, and the mean squared error loss for

644   reconstruction of normalized omics profiles, which could be denoted as:

645
$$\mathcal{L}_{repre} = \alpha_1 \mathcal{L}_{BNN} + \alpha_2 \mathcal{L}_{IOA} + \alpha_3 \mathcal{L}_{NLL} + \alpha_4 \mathcal{L}_{MSE}$$

where α_1, α_2, α_3 and α_4 are coefficients of the corresponding loss term, and the default values of them

are all set to 1.

Multi-sample integration with PRESENT-BC

We further extend the PRESENT framework to PRESENT-BC, which not only retains the ability of

PRESENT in incorporating multi-modal information, including various omics layers along with the

necessary spatial coordinate data, but also effectively eliminates batch effects while capturing shared

biological variations across samples. The architecture of PRESENT-BC also includes omics-specific

encoders, a cross-omics alignment module, and omics-specific decoders. However, compared with

PRESENT, the input to the omics-specific decoders in PRESENT-BC is the concatenation of the output

from the cross-omics alignment module and the one-hot encoding of batch labels. Additionally,

PRESENT-BC is enhanced for multi-slice integration based on a two-stage workflow. The first stage

serves as a warmup to pretrain the model, while the second stage leverage a cyclic iterative optimization

strategy to eliminate batch effects while preserving biological variations across samples. Using a scenario

of fusing two spatial multi-omics sequencing samples as an example, where ATAC, RNA, ADT, and

spatial coordinate data are simultaneously sequenced for each spot, we illustrate the workflow of multi-

sample integration with PRESENT-BC in the subsequent sections (Fig. 1b).

Feature sets unification. Before multi-sample integration, the first step is to align the feature sets of

different sequencing samples, which serves as the anchor for multi-sample integration. For ATAC data,

following the merging objects vignette provided by the Signac package[90], a unified set of peak regions is

first obtained by merging all intersecting peaks from all the samples. The fragment counts of each spot

within the unified peak set are then quantified for each sample. For RNA and ADT data, whose feature

sets are predefined genes or proteins, we directly leverage the intersection of gene or protein sets of all the

samples as the unified gene or protein set. After the gene intersection step, highly variable genes will be

selected based on the unified gene set. For convenience, we denoted the ATAC, RNA, ADT and spatial

data of the two samples as $X_1^{ATAC} \in R^{n_1 \times r}$, $X_2^{ATAC} \in R^{n_2 \times r}$, $X_1^{RNA} \in R^{n_1 \times g}$, $X_2^{RNA} \in R^{n_2 \times g}$, $X_1^{ADT} \in$

$R^{n_1 \times p}$, $X_2^{ADT} \in R^{n_2 \times p}$, $M_1 \in R^{n_1 \times 2}$ and $M_2 \in R^{n_2 \times 2}$, respectively. Here, r represents the number of

unified regions, g the number of highly variable genes, p the number of input proteins, and n_1 and n_2

the number of spots of the two samples. The unified ATAC, RNA, ADT data are then concatenated and

preprocessed using the omics-specific methods described in the previous section, obtaining normalized

data, i.e., $\tilde{X}^{ATAC} \in R^{n \times r}$, $\tilde{X}^{RNA} \in R^{n \times g}$, $\tilde{X}^{ADT} \in R^{n \times p}$ and $M \in R^{n \times 2}$, where $n = n_1 + n_2$ represents

the number of spots from the two samples.

Cross-sample graph construction. Due to inconsistent dissection and placement of sequencing tissues,

the spatial coordinates between different samples can hardly be directly compared with each other,

resulting in the failure of representation learning models in capturing the relationship between different

samples[22, 24]. To address this gap, we proposed a strategy to construct a cross-sample graph. Firstly, we

find intra-sample spatial neighbors for each spot based on the spatial coordinate matrix, building

undirected edges between all spots and their k nearest neighbors in Euclidean space from the same

sample. Secondly, we find inter-sample neighbors based on specified feature matrices, such as ATAC,

RNA and ADT profiles, or even latent embeddings, by establishing inter-sample undirected edges

between all spots and their k most similar neighbors in cosine space from other samples. Note that the

parameter k is set to 6 by default. Finally, we acquire a cross-sample graph which consists of both intra-

sample edges and inter-sample edges.

First-stage pretraining. In the first stage, we pretrain the parameter of PRESENT-BC based on a cross-

sample graph and multi-omics sequencing data of different samples, including $\tilde{X}^{ATAC} \in R^{n \times r}$, $\tilde{X}^{RNA} \in$

$R^{n \times g}$ and $\widetilde{X}^{ADT} \in R^{n \times p}$. Specifically, the intra-sample edges in the cross-sample graph are built based on

the Euclidean distance between spots using their spatial coordinates, while the inter-sample edges are

establish based on the cosine similarity between different spots using their ATAC, RNA and ADT profiles.

The objective function and training process of PRESENT-BC in the first pretraining stage are identical to

those of PRESENT, as described in the previous section.

Second-stage integration. The first pretraining stage enables PRESENT-BC to capture biological

variations within each sample, while the technical variations between different samples, i.e., batch effects,

have not been fully addressed. In the second stage, we design effective mechanisms, including an inter-

batch alignment loss, a batch-adversarial learning strategy and an intra-batch preserving loss, to eliminate

the batch effects across samples while simultaneously preserving the shared and sample-specific

biological signals, which are described in detail in the subsequent subsections.

Inter-batch alignment. We designed a inter-batch alignment (IBA) loss to align spots from different

samples based on contrastive learning[45]. Firstly, we find mutual nearest neighbors from other samples as

the positive set of for spot i, denoted as N_i^{pos}, referring to the inter-sample neighbors that are connected

to spot i in the cross-sample graph. Secondly, we randomly select $1/4$ proportion of spots from the

same sample as the negative set of spot i, depicted as N_i^{neg}. Finally, the IBA loss of spot i is:

707
$$\mathcal{L}_{IBA}^{(i)} = -log \frac{\sum_{j \in N_i^{pos}} exp(\delta_{sim}(\boldsymbol{e_i}, \boldsymbol{e_j}))}{\sum_{j \in N_i^{pos}} exp(\delta_{sim}(\boldsymbol{e_i}, \boldsymbol{e_j})) + \sum_{j \in N_i^{neg}} exp(\delta_{sim}(\boldsymbol{e_i}, \boldsymbol{e_j}))}$$

708   where $\boldsymbol{e_i} \in \boldsymbol{R}^d$ denotes the latent representation of spot i, d is the output dimension of PRESENT-BC

709   and δ_{sim} denotes the cosine similarity. Therefore, the inter-batch alignment loss of all the spots is:

710
$$\mathcal{L}_{IBA} = \frac{1}{n} \sum_{i=1}^{n} \mathcal{L}_{IBA}^{(i)}$$

Batch-adversarial learning. Inspired by a domain-adversarial learning research[92] and its's success in

multi-batch integration of single-cell ATAC-seq data[46], we implement a batch-adversarial learning

strategy to remove batch effect contained in joint representations. Specifically, we introduce a batch-

discriminator, aiming to predict the batch index of each spot based on its latent embedding, while the

omics-specific encoders and the cross-omics alignment module of PRESENT-BC are treated as the

generator, aiming to produce latent embeddings without batch effect, thereby confusing the batch-

discriminator in an adversarial manner. The loss function of the batch-adversarial learning is:

718
$$\mathcal{L}_{ADV} = \max_G \min_D CE(P, T)$$

where G refers to the generator, D denotes the batch-discriminator, CE represents the cross-entropy

loss, P denotes the predicted batch indices and T refers to the ground truth batch labels.

Intra-batch preserving. The first pretraining stage empowers PRESENT-BC to capture the biological

variations within each sample, while the inter-batch alignment loss and the batch-adversarial learning

strategy in the second integration stage aims for batch correction across samples. However, the key

challenge of multi-sample integration is how to eliminate batch effects while simultaneously preserving

biological signals. We therefore proposed an intra-batch preserving loss to iteratively maintain the pattern

of latent representations within each batch during batch correction, denoted as:

727
$$\mathcal{L}_{IBP} = \frac{1}{S} \sum_{k=1}^{S} \frac{1}{|s_k|} \sum_{i \in s_k} KLD\left(q_{s_k}^{(i)}(e^{(t)}), q_{s_k}^{(i)}(e^{(t-1)})\right)$$

728   where S represents the number of samples, s_k denotes the spots in sample k, $|s_k|$ denotes the number

729   of spots in s_k, $e^{(t)}$ and $e^{(t-1)}$ refer to the joint embeddings of all spots in current epoch and the last

730   epoch, respectively. $q_{s_k}^{(i)}(*)$ represents the embedding distribution of spot i over other spots in sample k,

731   defined as (taking the embeddings of current iteration $e^{(t)}$ as an example):

732
$$q_{s_k}^{(i)}(e^{(t)}) = \left(\frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(1)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}, \frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(2)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}, \dots, \frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(|s_k|)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}\right)$$

733
$$\delta_{ed}(u, v) = exp(\frac{u^T v}{\tau})$$

where $s_k^{(j)}$ denotes the j-th spot in sample k, τ is a temperature parameter and set to 0.1 by default[42].

Loss functions of PRESENT-BC and the parameter updating process. Batch-adversarial learning

strategy separate the parameter updating process of each epoch into two steps. In the first step, we update

the parameters of the batch-discriminator using the cross-entropy loss $\mathcal{L}_{ADV}^{disc} = \beta_3 CE(P, G)$, with

parameters of the generator and omics-specific decoders fixed. The generator denotes the omics-specific

encoders and cross-omics alignment module of PRESENT-BC. In the second step, we fix the parameters

of batch-discriminator and update the parameters of generator and omics-specific decoders based on an

overall loss function, consisting of the cross-omics representation loss, the inter-batch alignment loss, the

intra-batch preserving loss and a negative cross-entropy loss for the batch-adversarial learning process:

$$\mathcal{L}_{integ} = \mathcal{L}_{repre} + \beta_1 \mathcal{L}_{IBA} + \beta_2 \mathcal{L}_{IBP} + \beta_3 \mathcal{L}_{ADV}^{gen}$$

$$\mathcal{L}_{ADV}^{gen} = -CE(P, G)$$

where β_1, β_2 and β_3 are coefficients of the corresponding loss function, and set to 1 by default.

Cyclic iterative optimization. The training process of the second integration stage is carried out in a

cyclic iterative manner. First, we obtain the joint embeddings of multiple samples using the trained

PRESENT-BC. Next, we construct a cross-sample graph according to the strategy described in the

previous section, where intra-sample connections are established based on the spatial coordinates of spots

and the inter-sample edges are based on the cosine similarity between the newly acquired joint

embeddings of spots from different samples. Finally, we train the PRESENT-BC model for one epoch

using the newly constructed cross-sample graph. The cyclic process, aiming to iteratively refine the cross-

sample graph, is repeated until the model converges. The cross-sample graph is not only significant for

integration of cross-sample relationships based on the GAT module in omics-specific encoders, but also

important for the inter-batch alignment loss, which utilizes the inter-sample edges in cross-sample graph

to construct positive neighbor sets for spots.

Implementation details

With the development of spatial omics sequencing technologies, the number of spots in sequenced tissue has been quickly increasing from thousands to tens and even hundreds of thousands[10, 12, 15, 16, 93]. PRESENT and its extension PRESENT-BC are built upon graph neural networks, which face challenges such as memory explosion and neighborhood expansion when the number of nodes in graph grows large[94, 95]. To address these problems, we adopted necessary sparse adjacency matrix implementation based on the PyG framework when deploying our model. Additionally, we implemented the ClusterGCN algorithm to divide the entire graph into multiple small subgraphs[94], which enables our model to be trained on GPUs with limited memory and extend to larger spatial omics data.

The input dimensions of ATAC and RNA data are generally much higher than the ADT data, resulting in different hidden dimension settings for them. Specifically, the hidden and output dimensions of omics-specific encoders for ATAC and RNA data were (1024, 512) and 50, respectively, while those of omics-specific encoder for ADT data were $\left(2d_{adt}^{in}/3 + d_{adt}^{out}/3, d_{adt}^{in}/3 + 2d_{adt}^{out}/3\right)$ and d_{adt}^{out}. Here d_{adt}^{in} denotes the input dimension of ADT data and $d_{adt}^{out} = \min\left(50, d_{adt}^{in}\right)$ refers to the output dimension of omics-specific encoder for ADT data. The final output dimensions of both PRESENT and PRESENT-BC were set to 50 by default. Additionally, the hidden dimensions of omics-specific decoders were symmetric to the corresponding omics-specific encoders, i.e., $(2d_{adt}^{in}/3 + d_{adt}^{out}/3, d_{adt}^{in}/3 + 2d_{adt}^{out}/3)$ for ADT data and (512, 1024) for ATAC and RNA data. We trained the model for at most 100 epochs using the Adam optimizer, updating the parameters with an initial learning rate of 1e-3 and a weight decay of 1e-4.

Data collection and description

Single-nucleus ATAC-seq and spatial ATAC mouse embryo data. The single-nucleus ATAC-seq mouse embryo data[10] measured the chromatin accessibility of 1,879 cells from E12, E13 and E15

developmental stages. The spatial ATAC mouse embryo data[10] is composed of six samples, including

E12.5-S1 (3,274 spots), E12.5-S2 (3,903 spots), E13.5-S1 (3,160 spots), E13.5-S2 (2,970 spots), E15.5-

S1 (2,246 spots) and E15.5-S2 (2,234 spots). The original study[10] merged the single-cell ATAC-seq

datasets and spatial ATAC datasets and called 248,047 peak regions for the merged dataset based on

MACS2 v2.2.6 package.

Stereo-seq axolotl brain data. We collected the Stereo-seq axolotl brain developmental datasets from a

recent study which explored the regeneration and development of axolotl telencephalon using spatial

transcriptomic atlas[7]. The datasets consist of six sections from developmental stage 44 (1,477 spots),

stage 54 (2,929 spots), stage 57 (4,410 spots), juvenile (11,698 spots), adult (8,243 spots), and

metamorphosis (7,441 spots), measuring the expression level of 12,704 genes.

10x Visium mouse brain data. The spatial transcriptomic postnatal day 56 (P56) mouse brain data,

including a sagittal anterior section and a sagittal posterior section, are collected from the 10x Genomics

Visium platform. The mouse sagittal anterior section contains the expression level of 32,285 genes on

2,823 spots, while the posterior section provides the expression level of 32,285 genes on 3,289 spots.

SPOTS mouse spleen data. The mouse spleen data[13] is generated by a high-throughput spatial

transcriptomics and proteomics co-profiling technology, i.e., Spatial PrOtein and Transcriptome

Sequencing (SPOTS). This dataset provides the spatial coordinates and the expression level of 32,285

genes and 21 proteins on 2,568 spots.

10x Visium RNA-Protein human lymph node data. The human lymph node dataset, generated by 10x

Genomics Visium transcriptomics and proteomics co-profiling technology, is collected from a spatial

multi-omics representation learning study[32]. The in-house dataset contains the spatial coordinates and the

expression level of 18,085 genes and 31 proteins on 3,484 spots.

MISAR-seq mouse brain data. The MISAR-seq mouse brain data[14] provides the paired sequencing data

of spatial gene expression and chromatin accessibility during mouse brain development. We collected

four manually annotated samples from original study[14], including E11.0 (1,258 spots), E13.5 (1,777

spots), E15.5 (1,949 spots) and E18.5 (2,129 spots). We called peak for each sample based on the MACS2

v2.2.9.1 package, obtaining 242,024, 271,126, 265,014 and 294,734 peak regions, respectively.

Experiment setup and baseline methods

Cross-modality representation of spatial ATAC mouse embryo data. The spatial ATAC mouse embryo

data firstly went through a quality control process, where peaks accessible in less than 0.5% spots were

filtered out. Using the filtered ATAC count matrices as input, we then benchmarked the performance of

PRESENT with four methods designed for single-cell ATAC-seq data, including SCALE[49], cisTopic[50] and

SnapATAC2[51], and three methods designed for SRT data, i.e., SpaGCN[29], STAGATE[30] and PAST[17]. All

the baseline methods were implemented according to default settings except for STAGATE and PAST. In

original tutorials, the two SRT methods adopted a feature selection process which is specially designed

for transcriptomic data and only selected 3,000 features, resulting significantly lower performance

(Supplementary Fig. 31). Since other methods utilized all the input peaks for dimension reduction, we

ignored the feature selection step of these two methods for fairness and better performance.

Cross-modality representation of Stereo-seq axolotl brain data. The genes with zero expression in all

spots were filtered out. The filtered count matrices and the spatial locations are fed into PRESENT and

other baseline methods, including SpaGCN[29], STAGATE[30], PAST[17] and SpatialPCA[18]. All baseline

methods are implemented with default settings in their tutorials. Following the workflow of processing

single-cell[96] and spatial[17, 18, 30] RNA-seq data, PRESENT also employed a gene selection procedure,

obtaining 3,000 highly variable genes as input.

Cross-modality representation of 10x Visium RNA-Protein human lymph node data. We filtered out

genes and proteins with zero expression in all spots. Baseline methods includes SpatialGlue[32], totalVI[57],

CiteFuse[58] and Seurat-v4[59]. All baseline methods were conducted according to their tutorials.

Cross-modality representation of MISAR-seq mouse brain data. We filtered out peaks accessible in

less than 3% spots and genes with zero expression in all spots for quality control. The filtered count

matrices of ATAC data and RNA data along with the spatial coordinate matrix were then fed into the

workflow of PRESENT and other baseline methods, involving SpatialGlue[32], MOFA+[68], scAI[69],

scMDC[70], scMVP[19], Seurat-v4[59] and SnapATAC2[51]. All these baseline methods are implemented

according to their default settings. PRESENT again selected 3,000 highly variable genes as the

transcriptomic input for further analysis.

Multi-sample integration of 10x Visium mouse brain data. The intersected gene set of the two mouse

brain samples, including a sagittal anterior section and a sagittal posterior section, were utilized as the

unified gene set. Afterwards, gene count matrices of the two samples were concatenated and genes with

zero expression in all spots were filtered out. Next, 3,000 highly variable genes were selected as the input

for PRESENT-BC. PRESENT-BC was compared with two recently developed SRT methods, i.e.,

STAligner[25] and GraphST[23], which were conducted according to their default settings. Specifically, we

utilized PASTE[22] to accomplish the initial coordinate alignment step for GraphST according to the

recommendation of the original study[23].

Multi-sample integration of spatial ATAC mouse embryo data. The three spatial ATAC mouse embryo

samples from three developmental stages have the same peak sets and were directly concatenated.

Following the experimental settings in cross-modality representation of a single spatial sample, peaks

accessible in less than 0.5% spots were filtered out. Using the filtered concatenated matrix, concatenated

spatial matrix and batch indices as input, we benchmarked PRESENT-BC with six integration methods

developed for single-cell data (MNN[79], Scanorama[39], Harmony[38], SCALEX[80], scBasset-BC[81] and

SnapATAC2[51]) and two recently proposed SRT methods (STAligner[25] and GraphST[23]). MNN, Scanorama

and Harmony performed batch correction based on the latent embeddings obtained by a widely used

epigenomic toolkit, EpiScanpy[97]. All the methods are implemented according to their default settings

except for STAligner. Due to similar reasons in the cross-modality representation for a single sample, we

ignored the feature selection step of STAligner to ensure better performance and fairness.

Multi-sample integration of MISAR-seq mouse brain data. We merged the peak sets of different

mouse brain samples according to the merging objects vignette provided by the Signac package[90], which

first obtained a unified peak set by merging all intersecting peaks from all the samples and then quantified

the fragment count of each spot on the newly obtained peak set. Consistent with the experimental settings

in cross-modality representation, peaks accessible in less than 3% spots in the concatenated matrix were

filtered out and 3,000 highly variable genes of transcriptomic data were selected for PRESENT-BC. Due

to the lack of methods specially designed for the integration of spatial multi-omics data, we compared the

batch effect removal and biological variance conservation performance of PRESENT-BC with six

methods developed for single-cell data (MNN[79], Scanorama[39], Harmony[38], SCALEX[80], scBasset-BC[81]

and SnapATAC2[51]) and three recently proposed SRT multi-sample integration methods (STAligner[25],

Stitch3D[31] and GraphST[23]). Specifically, MNN, Scanorama, Harmony, SCALEX could be applied to both

transcriptomic and epigenomic data, denoted as RNA-* or ATAC-*, respectively, where * denotes the

different methods. MNN, Scanorama and Harmony performed batch correction based on low-dimensional

embeddings extracted from RNA or ATAC data, which were accomplished based on the workflow of

SCANPY[96] and EpiScanpy[97], respectively. All the methods were implemented using default settings.

Evaluation metrics

Spatial clustering metrics. We implemented four commonly used metrics[53, 54] to quantitatively evaluate

the performance of spatial clustering, including adjusted rand index (ARI), adjusted mutual information

(AMI), normalized mutual information (NMI) and homogeneity (Homo).

ARI, ranging from 0 to 1, denotes the consistent adjustment of the expectations of rand index:

877
$$\text{ARI} = \frac{RI(P,T) - E[RI(P,T)]}{\max[RI(P,T)] - E[RI(P,T)]}$$

where $RI(\cdot)$ denotes the function of rand index, T refers to the ground truth labels, and P represents

the clustering labels. Rand index is the probability of the agreement between the clustering result and the

ground truth labels on a randomly selected spot set.

Mutual Information (MI) accesses the similarity between the clustering labels and ground truth

labels, from which both AMI and NMI originate. AMI ranges from -1 to 1, while NMI ranges from 0 to 1.

AMI is calculated as:

884
$$\text{AMI} = \frac{MI(P,T) - E[MI(P,T)]}{avg[H(P), H(T)] - E[MI(P,T)]}$$

885 where $MI(\cdot)$ refers to the mutual information, $E(\cdot)$ represents the expectation function and $H(\cdot)$

886 denotes the function of information entropy. NMI can be calculated as:

887
$$\text{NMI} = \frac{MI(P,T)}{\sqrt{H(P)H(T)}}$$

888 Homogeneity is another metric to measure the clustering performance. Specifically, if all the clusters

889 involve only data points that are members of a single class in the ground truth labels, the clustering results

890 could be interpreted as satisfying homogeneity, calculated as:

891
$$\text{Homo} = 1 - \frac{H(T|P)}{H(T)}$$

Integration metrics. According to a recent benchmarking framework for single-cell omics data

integration, two important perspectives need to be considered when evaluating the performance of multi-

sample integration, namely the batch effect removal and the biological variance conservation[82].

To quantitatively evaluate the performance of batch effect removal, we introduced several metrics

used in the benchmarking framework[82], including graph connectivity(GC), k-nearest-neighbor batch-

effect test (kBET) and batch average silhouette width (bASW). In addition, we also implemented a cross-

validation strategy to evaluate the cross-batch annotation performance based on latent representations

obtained by different integration methods, which is also an important assessment of batch correction[98].

Specifically, using the spots from a specific sample as the test set and those from other samples along

with the domain labels as the training set, we transferred the domain labels of the training set into the test

set based on a k-NN classifier and the joint embeddings of all the samples. k-NN classifier was used as

its performance mainly depended on the quality of joint embeddings obtained by different methods. The

performance of label transfer was evaluated with four commonly used metrics implemented in the scikit-

learn package, i.e., accuracy (Acc), kappa, macro F1 score (mF1) and the weighted F1 score (wF1). After

all the samples were used as the test set in a cross-validation manner, we calculated the average score for

each metric to evaluate the performance of cross-batch annotation.

The quantitative evaluation for biological variance conservation involves the accessment of

representative quality and the clustering performance. The representative quality was accessed via the

mean average precision (MAP)[99] and two metrics included in the benchmarking framework[82], namely the

spot average silhouette width (sASW) and spot local inverse Simpson's index (sLISI) , while the

evaluation of clustering performance was based on the four metrics, i.e., ATI, AMI, NMI and Homo.

We calculated category scores of the batch effect removal and the biological variance conservation

through averaging the metrics mentioned in the two aspects, respectively. We also calculated an overall

score using a 40/60 weighted mean of the two category scores[82, 99].

Downstream analyses

Visualization. Based on the latent representations obtained by various methods, we utilized the uniform

manifold approximation and projection (UMAP)[55] algorithm to further reduce the dimension of latent

representation matrices into two-dimensional matrices for visualization. In addition, the two-dimensional

UMAP and spatial coordinate matrices were visualized using the functions *scanpy.pl.umap()* and

scanpy.pl.spatial() wrapped in the SCANPY package[96].

Spatial clustering. We implemented a binary search strategy to tune the resolution of Leiden algorithm[52],

a graph-based clustering method, to match the number of spatial clusters as closely as possible with the

number of ground truth domains. Specifically, we set a minimum resolution res_{min} and a maximum

resolution res_{max}, and obtain clustering results using a resolution equal to $\frac{res_{min}+res_{max}}{2}$. If the number

of clusters exceeds the number of ground truth domains, res_{max} is updated to $\frac{res_{min}+res_{max}}{2}$. Otherwise,

res_{min} is updated to $\frac{res_{min}+res_{max}}{2}$. The iterative process continues until the number of clusters matches

the number of ground truth domains or the current iteration reaches the maximum number of iterations.

Differentially expressed gene and protein analysis. We implemented differential analysis based on the

Wilcoxon rank-sum test[61] wrapped in *scanpy.tl.rank_genes_groups()* function of the SCANPY package[96].

Gene ontology analysis. Given a specific gene set, we conducted gene ontology (GO) analysis using the

clusterProfiler package[100], which automated the process of biological-term classification and the

enrichment analysis of gene clusters.

Data availability

The single- nucleus ATAC-seq and spatial ATAC mouse embryo datasets[10] are available at Gene

Expression Omnibus (GEO) using the accession code GSE214991. The Stereo-seq axolotl brain datasets[7]

as well as the reference dataset, i.e., axolotl brain data at 20 days post-injury, can be obtained through

their database https://db.cngb.org/stomics/artista/. The 10x Visium mouse brain datasets, including a

sagittal anterior section and a sagittal posterior section, are accessible at the 10x Genomics websites

https://www.10xgenomics.com/datasets/mouse-brain-serial-section-2-sagittal-anterior-1-standard and

https://www.10xgenomics.com/datasets/mouse-brain-serial-section-2-sagittal-posterior-1-standard,

respectively. The SPOTS mouse spleen data[13] is accessible at GEO with accession code GSE198353. The

10x Visium RNA & protein human lymph node dataset[32] can be downloaded from the Zenodo database

https://zenodo.org/records/10362607. The MISAR-seq mouse brain data[14] is available from the National

Genomics Data Center via accession number OEP003285.

Code availability

949   The code for implementing PRESENT and PRESENT-BC can be found at the Github website

950   https://github.com/lizhen18THU/PRESENT.

Acknowledgements

952   This work was supported by the National Key Research and Development Program of China (grant nos.

953   2023YFF1204802 and 2021YFF1200902) and the National Natural Science Foundation of China (grant

954   nos. 62273194).

Author contributions

956   R.J. conceived the study and supervised the project. Z.L. designed, implemented and validated the

957   PRESENT framework. X.J.C., X.Y.C., Z.G., Y. L., Y. P. and S.C. helped with analyzing the results. Z.L.

958   and R.J. wrote the manuscript, with input from all the authors.

Competing interests

960   The authors declare no competing interests.

References

1. Baysoy, A. et al. The technological landscape and applications of single-cell multi-omics. *Nat. Rev. Mol. Cell Biol.* **24**, 695-713 (2023).

2. Kashima, Y. et al. Single-cell sequencing techniques from individual to multiomics analyses. *Exp. Mol. Med.* **52**, 1419-1427 (2020).

3. Adema, K. et al. Lost in space: what single-cell RNA sequencing cannot tell you. *Trends Plant Sci.* (2024).

4. Armingol, E. et al. Deciphering cell-cell interactions and communication from gene expression. *Nat. Rev. Genet.* **22**, 71-88 (2021).

5. Cang, Z.X. et al. Screening cell-cell communication in spatial transcriptomics via collective optimal transport. *Nat. Methods* **20**, 218-228 (2023).

6. Shao, X. et al. Knowledge-graph-based cell-cell communication inference for spatially resolved transcriptomic data with SpaTalk. *Nat. Commun.* **13**, 4429 (2022).

7. Wei, X. et al. Single-cell Stereo-seq reveals induced progenitor cells involved in axolotl brain regeneration. *Science* **377**, eabp9444 (2022).

8. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat. Methods* **19**, 534-546 (2022).

9. Deng, Y.X. et al. Spatial profiling of chromatin accessibility in mouse and human tissues. *Nature* **609**, 375-383 (2022).

10. Llorens-Bobadilla, E. et al. Solid-phase capture and profiling of open chromatin by spatial ATAC. *Nat. Biotechnol.* **41**, 1085-1088 (2023).

11. Vandereyken, K. et al. Methods and applications for single-cell and spatial multi-omics. *Nat. Rev. Genet.* **24**, 494-515 (2023).

12. Liu, Y. et al. High-plex protein and whole transcriptome co-mapping at cellular resolution with spatial CITE-seq. *Nat. Biotechnol.* **41**, 1405-1409 (2023).

13. Ben-Chetrit, N. et al. Integration of whole transcriptome spatial profiling with protein markers. *Nat. Biotechnol.* **41**, 788-193 (2023).

14. Jiang, F.Q. et al. Simultaneous profiling of spatial gene expression and chromatin accessibility during mouse brain development. *Nat. Methods* **20**, 1048-1057 (2023).

15. Yuan, Z.Y. et al. SODB facilitates comprehensive exploration of spatial omics data. *Nat. Methods* **20**, 387-399 (2023).

16. Xu, Z. et al. STOmicsDB: a comprehensive database for spatial transcriptomics data sharing, analysis and visualization. *Nucleic Acids Res.* **52**, D1053-D1061 (2024).

17. Li, Z. et al. Latent feature extraction with a prior-based self-attention framework for spatial transcriptomics. *Genome Res.* **33**, 1757-1773 (2023).

18. Shang, L. & Zhou, X. Spatially aware dimension reduction for spatial transcriptomics. *Nat. Commun.* **13**, 7203 (2022).

19. Li, G.Y. et al. A deep generative model for multi-view profiling of single-cell RNA-seq and ATAC-seq data. *Genome Biol.* **23**, 20 (2022).

20. Swanton, C. Intratumor heterogeneity: evolution through space and time. *Cancer Res.* **72**, 4875-4882 (2012).

21. Janiszewska, M. The microcosmos of intratumor heterogeneity: the space-time of cancer evolution. *Oncogene* **39**, 2031-2039 (2020).

22. Zeira, R. et al. Alignment and integration of spatial transcriptomics data. *Nat. Methods* **19**, 567-575 (2022).

23. Long, Y. et al. Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nat. Commun.* **14**, 1155 (2023).

24. Guo, T.T. et al. SPIRAL: integrating and aligning spatially resolved transcriptomics data across different experiments, conditions, and technologies. *Genome Biol.* **24**, 241 (2023).

1007   25.   Zhou, X., Dong, K.N. & Zhang, S.H. Integrating spatial transcriptomics data across different conditions,
1008   technologies and developmental stages. *Nat. Comput. Sci.* **3**, 894-906 (2023).

1009   26.   Chen, S. et al. Spatially resolved transcriptomics reveals genes associated with the vulnerability of middle
1010   temporal gyrus in Alzheimer's disease. *Acta Neuropathol. Commun.* **10**, 188 (2022).

1011   27.   Cheng, A., Hu, G. & Li, W.V. Benchmarking cell-type clustering methods for spatially resolved
1012   transcriptomics data. *Brief. Bioinform.* **24**, bbac475 (2023).

1013   28.   Yuan, Z. et al. Benchmarking spatial clustering methods with spatially resolved transcriptomics data. *Nat.*
1014   *Methods* **21**, 1-11 (2024).

1015   29.   Hu, J. et al. SpaGCN: Integrating gene expression, spatial location and histology to identify spatial domains
1016   and spatially variable genes by graph convolutional network. *Nat. Methods* **18**, 1342-1351 (2021).

1017   30.   Dong, K.N. & Zhang, S.H. Deciphering spatial domains from spatially resolved transcriptomics with an
1018   adaptive graph attention auto-encoder. *Nat. Commun.* **13**, 1739 (2022).

1019   31.   Wang, G. et al. Construction of a 3D whole organism spatial atlas by joint modelling of multiple slices with
1020   deep neural networks. *Nat. Mach. Intell.* **5**, 1200-1213 (2023).

1021   32.   Long, Y. et al. Integrated analysis of spatial multi-omics with SpatialGlue. Preprint at *bioRxiv*
1022   https://doi.org/10.1101/2023.04.26.538404 (2023).

1023   33.   Tian, T. et al. Dependency-aware deep generative models for multitasking analysis of spatial omics data. *Nat.*
1024   *Methods* (2024).

1025   34.   Berger, S.L. The complex language of chromatin regulation during transcription. *Nature* **447**, 407-412 (2007).

1026   35.   MacNeil, L.T. & Walhout, A.J.M. Gene regulatory networks and the role of robustness and stochasticity in
1027   the control of gene expression. *Genome Res.* **21**, 645-657 (2011).

1028   36.   Stoeckius, M. et al. Simultaneous epitope and transcriptome measurement in single cells. *Nat. Methods* **14**,
1029   865-868 (2017).

1030   37.   Peterson, V.M. et al. Multiplexed quantification of proteins and transcripts in single cells. *Nat. Biotechnol.*
1031   **35**, 936-939 (2017).

1032   38.   Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods*
1033   **16**, 1289-1296 (2019).

1034   39.   Hie, B., Bryson, B. & Berger, B. Efficient integration of heterogeneous single-cell transcriptomes using
1035   Scanorama. *Nat. Biotechnol.* **37**, 685-691 (2019).

1036   40.   Veličković, P. et al. Graph attention networks. Preprint at *arXiv* https://doi.org/10.48550/arXiv.1710.10903
1037   (2017).

1038   41.   Blundell, C. et al. Weight uncertainty in neural network. In *International conference on machine learning*
1039   (2015).

1040   42.   Lin, Z. et al. Multi-modal Contrastive Representation Learning for Entity Alignment. In *Proceedings of the*
1041   *29th International Conference on Computational Linguistics* (2022).

1042   43.   Eraslan, G. et al. Single-cell RNA-seq denoising using a deep count autoencoder. *Nat. Commun.* **10**, 390
1043   (2019).

1044   44.   Martens, L.D. et al. Modeling fragment counts improves single-cell ATAC-seq analysis. *Nat. Methods* **21**,
1045   28-31 (2023).

1046   45.   Chen, T. et al. A simple framework for contrastive learning of visual representations. In *International*
1047   *conference on machine learning* (2020).

1048   46.   Kopp, W., Akalin, A. & Ohler, U. Simultaneous dimensionality reduction and integration for single-cell
1049   ATAC-seq data using deep learning. *Nat. Mach. Intell.* **4**, 162-168 (2022).

1050   47.   Heumos, L. et al. Best practices for single-cell analysis across modalities. *Nat. Rev. Genet.* **24**, 550-572
1051   (2023).

1052   48.   Zhang, H. et al. Extensive evaluation of ATAC-seq protocols for native or formaldehyde-fixed nuclei. *BMC*

1053   *Genomics* **23**, 214 (2022).

49.   Xiong, L. et al. SCALE method for single-cell ATAC-seq analysis via latent feature extraction. *Nat. Commun.* **10**, 4576 (2019).

50.   González-Blas, C.B. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-seq data. *Nat. Methods* **16**, 397-400 (2019).

51.   Zhang, K. et al. A fast, scalable and versatile tool for analysis of single-cell omics data. *Nat. Methods* **21**, 217-227 (2024).

52.   Traag, V.A., Waltman, L. & Van Eck, N.J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci. Rep.* **9**, 5233 (2019).

53.   Chen, S. et al. VPAC: Variational projection for accurate clustering of single-cell transcriptomic data. *BMC Bioinformatics* **20**, 139-151 (2019).

54.   Jiang, R. et al. SINFONIA: scalable identification of spatially variable genes for deciphering spatial domains. *Cells* **12**, 604 (2023).

55.   McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. Preprint at *arXiv* https://doi.org/10.48550/arXiv.1802.03426 (2018).

56.   Pierson, E. & Yau, C. ZIFA: Dimensionality reduction for zero-inflated single-cell gene expression analysis. *Genome Biol.* **16**, 1-10 (2015).

57.   Gayoso, A. et al. Joint probabilistic modeling of single-cell multi-omic data with totalVI. *Nat. Methods* **18**, 272-282 (2021).

58.   Kim, H.J. et al. CiteFuse enables multi-modal analysis of CITE-seq data. *Bioinformatics* **36**, 4137-4143 (2020).

59.   Hao, Y.H. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 (2021).

60.   Lewis, S.M., Williams, A. & Eisenbarth, S.C. Structure and function of the immune system in the spleen. *Sci. Immunol.* **4**, eaau6085 (2019).

61.   Soneson, C. & Robinson, M.D. Bias, robustness and scalability in single-cell differential expression analysis. *Nat. Methods* **15**, 255-261 (2018).

62.   Schmid, M.C. & Varner, J.A. Myeloid cells in tumor inflammation. *Vasc. Cell* **4**, 1-7 (2012).

63.   Borges, M.D. & Sesti-Costa, R. Macrophages: key players in erythrocyte turnover. *Hematol. Transfus. Cell Ther.* **44**, 574-581 (2022).

64.   Seyfried, A.N., Maloney, J.M. & MacNamara, K.C. Macrophages orchestrate hematopoietic programs and regulate HSC function during inflammatory stress. *Front. Immunol.* **11**, 548463 (2020).

65.   Cui, K. et al. Endocytic adaptors in cardiovascular disease. *Front. Cell Dev. Biol.* **8**, 624159 (2020).

66.   Baranov, M.V. et al. Modulation of immune responses by particle size and shape. *Front. Immunol.* **11**, 607945 (2021).

67.   Kurup, S.P., Butler, N.S. & Harty, J.T. T cell-mediated immunity to malaria. *Nat. Rev. Immunol.* **19**, 457-471 (2019).

68.   Argelaguet, R. et al. MOFA+ : a statistical framework for comprehensive integration of multi-modal single-cell data. *Genome Biol.* **21**, 1-17 (2020).

69.   Jin, S.Q., Zhang, L.H. & Nie, Q. scAI: an unsupervised approach for the integrative analysis of parallel single-cell transcriptomic and epigenomic profiles. *Genome Biol.* **21**, 1-19 (2020).

70.   Lin, X. et al. Clustering of single-cell multi-omics data with a multimodal deep learning method. *Nat. Commun.* **13**, 7705 (2022).

71.   Sunkin, S.M. et al. Allen Brain Atlas: an integrated spatio-temporal portal for exploring the central nervous system. *Nucleic Acids Res.* **41**, D996-D1008 (2012).

72.   Teicher, M.H., Anderson, C.M. & Polcari, A. Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proc. Natl. Acad. Sci.* **109**, E563-E572 (2012).

73. Matsuda, K. et al. Transsynaptic modulation of kainate receptor functions by C1q-like proteins. *Neuron* **90**, 752-767 (2016).

74. Park, S.-Y. et al. Hippocalcin promotes neuronal differentiation and inhibits astrocytic differentiation in neural stem cells. *Stem Cell Rep.* **8**, 95-111 (2017).

75. Hachem, S. et al. Expression of S100B during embryonic development of the mouse cerebellum. *BMC Dev. Biol.* **7**, 1-15 (2007).

76. Hirasawa, M. et al. Carbonic anhydrase related protein 8 mutation results in aberrant synaptic morphology and excitatory synaptic function in the cerebellum. *Mol. Cell. Neurosci.* **35**, 161-170 (2007).

77. Lassuthova, P. et al. Mutations in ATP1A1 cause dominant Charcot-Marie-Tooth type 2. *Am. J. Hum. Genet.* **102**, 505-514 (2018).

78. Pacey, L.K. et al. Delayed myelination in a mouse model of fragile X syndrome. *Hum. Mol. Genet.* **22**, 3920-3930 (2013).

79. Haghverdi, L. et al. Batch effects in single-cell RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat. Biotechnol.* **36**, 421-427 (2018).

80. Xiong, L. et al. Online single-cell data integration through projecting heterogeneous datasets into a common cell-embedding space. *Nat. Commun.* **13**, 6118 (2022).

81. Yuan, H. & Kelley, D.R. scBasset: sequence-based modeling of single-cell ATAC-seq using convolutional neural networks. *Nat. Methods* **19**, 1088-1096 (2022).

82. Luecken, M.D. et al. Benchmarking atlas-level data integration in single-cell genomics. *Nat. Methods* **19**, 41-50 (2022).

83. Argelaguet, R. et al. Computational principles and challenges in single-cell data integration. *Nat. Biotechnol.* **39**, 1202-1215 (2021).

84. Chen, S.Q. et al. RA3 is a reference-guided approach for epigenetic characterization of single cells. *Nat. Commun.* **12**, 13 (2021).

85. Chen, X. et al. Cell type annotation of single-cell chromatin accessibility data via supervised Bayesian embedding. *Nat. Mach. Intell.* **4**, 116-126 (2022).

86. Cusanovich, D.A. et al. Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**, 910-914 (2015).

87. Cusanovich, D.A. et al. A single-cell atlas of in vivo mammalian chromatin accessibility. *Cell* **174**, 1309-1324 (2018).

88. Mulè, M.P., Martins, A.J. & Tsang, J.S. Normalizing and denoising protein expression data from droplet-based single cell profiling. *Nat. Commun.* **13**, 2099 (2022).

89. Stoeckius, M. et al. Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. *Genome Biol.* **19**, 1-12 (2018).

90. Stuart, T. et al. Single-cell chromatin state analysis with Signac. *Nat. Methods* **18**, 1333-1341 (2021).

91. Qiu, P. Embracing the dropouts in single-cell RNA-seq analysis. *Nat. Commun.* **11**, 1169 (2020).

92. Ganin, Y. et al. Domain-adversarial training of neural networks. *J. Mach. Learn. Res.* **17**, 1-35 (2016).

93. Zhang, D. et al. Spatial epigenome-transcriptome co-profiling of mammalian tissues. *Nature* **616**, 113-122 (2023).

94. Chiang, W.-L. et al. Cluster-GCN: An efficient algorithm for training deep and large graph convolutional networks. In *Proceedings of the 25th ACM SIGKDD international conference on knowledge discovery & data mining* (2019).

95. Bai, J., Ren, Y. & Zhang, J. Ripple walk training: A subgraph-based training framework for large and deep graph neural network. In *2021 International Joint Conference on Neural Networks (IJCNN)* (2021).

96. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol.* **19**, 1-5 (2018).

1145   97.   Danese, A. et al. EpiScanpy: integrated single-cell epigenomic analysis. *Nat. Commun.* **12**, 5228 (2021).

1146   98.   Song, Y. et al. Benchmarking strategies for cross-species integration of single-cell RNA sequencing data. *Nat.*
1147   *Commun.* **14**, 6495 (2023).

1148   99.   Cao, Z.-J. & Gao, G. Multi-omics single-cell data integration and regulatory inference with graph-linked
1149   embedding. *Nat. Biotechnol.* **40**, 1458-1466 (2022).

1150   100.   Yu, G. et al. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS: J.*
1151   *Integrative Biol.* **16**, 284-287 (2012).

1152
1153

Figures

1155

 Fig. 1 | The framework of PRESENT. a, Cross-modality representation with PRESENT. PRESENT is capable to integrate
1157 spatial coordinate data and an arbitrary combination of different omics data, including proteomic data sequenced by
1158 antibody-derived tags (ADT), transcriptomic data by RNA sequencing (RNA), and epigenomic data generated by the assay
1159 for transposase-accessible chromatin with sequencing (ATAC). Here we illustrate the cross-modality representation of
1160 spatial information and all the three omics layers as an example. Firstly, an omics-specific encoder is used to extract

specific biological variations contained in each omics layer. Each omics-specific encoder is composed of a graph attention neural network (GAT) and Bayesian neural network (BNN) module. All the omics-specific feature matrices obtained by encoders are fused by a cross-omics alignment module consisting of an MLP module, which integrates multi-omics information via a contrastive learning inter-omics alignment loss function. The output of cross-omics alignment module refers to the final representation matrix of PRESENT. The distinct patterns of different omics layers are addressed via omics-specific decoders, which take the output of cross-omics alignment module as input and model the count matrices of RNA and ATAC data using the zero-inflation negative binomial (ZINB) and the zero-inflation Poisson (ZIP) distribution, respectively. **b,** Multi-sample integration with PRESENT-BC. PRESENT-BC adopts a two-stage workflow. In the first pretraining stage, PRESENT-BC has similar architecture as PRESENT and is pretrained using omics data and a cross-sample graph. The intra-sample edges of the cross-sample graph are built using spatial coordinates, while inter-sample connections are established based on the similarity of omics data. In the second integration stage, PRESENT-BC implements a cyclic iterative strategy to refine the cross-sample graph using the joint embeddings obtained by the trained model. PRESENT-BC introduces an inter-batch alignment loss, a batch-adversarial learning strategy and an intra-batch preserving loss in the second stage to eliminate batch effects while preserving biological signals. **c,** Applications of PRESENT in cross-modality representation and PRESENT-BC in multi-sample integration of spatially resolved omics data. The representation obtained by PRESENT can be applied to various downstream analysis, including but not limited to visualization and spatial domain identification. On the other hand, PRESENT-BC enables effective integration of both horizontal and vertical spatial samples generated by various technologies, facilitating integrated analysis of spatial datasets across different dissection regions and developmental stages.

Fig. 2 | PRESENT facilitates accurate spatial domain identification in spatial ATAC-seq and SRT data. a, The ridge plot and violin plot of log10 total fragment counts, namely log10(nFrags), across the six spatial ATAC mouse embryo samples. **b,** The ridge plot of TSS enrichment scores for spots and the line plot of the enrichment profiles of fragments around the TSS across the six mouse embryo samples. **c,** The quantitative evaluation of spatial clustering performance of

different methods using four metrics, including ARI, AMI, NMI and Homo. The center line, box limits and whiskers in the
box and violin plots are the median, upper and lower quartiles, and 1.5× interquartile range, respectively. **d,** The spatial
visualization colored by ground truth domain labels, **e,** the UMAP visualization of the representations obtained by different
methods colored by ground truth domain labels and **f,** the spatial visualization of clusters identified using the latent
representations obtained by different methods on the E15.5-S1 mouse embryo sample. **g,** The detailed comparison of
performance of PRESENT and PRESENT-SN & SP on the six mouse embryo samples based on radar plots. **h,** The UMAP
visualization of the latent representations obtained by different methods colored by ground truth domain labels on E12.5-
S2 mouse embryo sample. **i,** The spatial clustering performance of baseline methods, PRESENT and PRESENT-SN & SP
on the six mouse embryo samples under different level of dropout rate. Note that SCALE encountered error when the
dropout rate was larger than 0.2.
1196

Fig. 3 | PRESENT leverages complementary omics information from paired spatial RNA-ADT data. a, The spatial visualization of the 10x Genomics Visium RNA-Protein human lymph node sample colored by ground truth domain labels. **b,** The quantitative comparison of spatial domain identification performance between PRESENT and other baseline methods based on bar plot and, **c,** the comparison between PRESENT utilizing both RNA and ADT data (RNA & ADT) and PRESENT using only RNA (RNA-only) or ADT data (ADT-only) based on radar plot on the human lymph node dataset. **d,** The UMAP visualization of latent representations obtained by different methods colored by ground truth domain labels and **e,** spatial cluster labels identified based on different methods on the human lymph node dataset. **f,** The spatial visualization of clusters identified based on different methods on the human lymph node dataset. **g,** The histology image and spatial visualization of spatial clusters identified based on PRESENT on the SPOTS mouse spleen dataset. **h,** The dot plot demonstrating the difference of ADT expression levels and **i,** RNA expression levels between each domain and the rest domains. **j,** The volcano plot illustrating the difference of RNA expression levels between Mac1 and Mac2-enriched domains. The vertical dashed line represents the threshold for $\log(\text{FC}) = \pm 0.2$, while the horizontal dashed line denotes the threshold for $-\log(\text{FDR} = 0.05)$. **k,** The chordal plots of the Mac1-enriched domain (left) and the Mac2-enriched domain (right) demonstrating the linkage of differential expressed genes (DEGs) in each domain and the corresponding enriched biological processes. In each chordal plot, the left semicircle represents DEGs while the right semicircle denotes the enriched biological processes.

Fig. 4 | PRESENT extracts cross-modality representations from parallel spatial RNA-ATAC datasets. a, The
quantitative comparison between PRESENT and other baseline methods and **b,** the comparison between PRESENT
utilizing both RNA and ATAC data (RNA & ATAC) and PRESENT using only RNA (RNA-only) or ATAC (ATAC-only)

data based on box plot on the MISAR-seq mouse brain dataset. The center line, box limits and whiskers in the box plots

are the median, upper and lower quartiles, and 1.5× interquartile range, respectively. **c,** The spatial visualization of spots

from the E18.5 mouse brain sample colored by ground truth domain labels. **d,** The heatmap visualization of latent

representations obtained by PRESENT. **e,** The UMAP visualization of latent representations obtained by different methods

colored by ground truth domain labels. **f,** The UMAP visualization of latent representations obtained by different methods

and **g,** the spatial visualization colored by spatial clusters identified based on different methods on the E18.5 mouse brain

sample. The UMAP visualization of latent embeddings obtained by PRESENT using only RNA data, only ATAC data and

both RNA & ATAC data, colored by ground truth domain labels on the **h,** E11.0 and **i,** E15.5 mouse brain samples.

1229

1230

Fig. 5 | PRESENT-BC enables effective integration of horizontal and vertical spatially resolved samples. a, The anatomic annotation of the sagittal region in P56 mouse brain provided by Allen Reference Atlas. The joint spatial clustering results based on the latent embeddings obtained by **b,** STAligner, **c,** GraphST and **d,** PRESENT-BC on the two horizontal mouse brain sagittal samples generated by the 10x Genomics Visium platform. **e,** The quantitative evaluation of different integration methods on the three spatial ATAC mouse embryo samples using 14 metrics divided into two categories, namely batch effect removal and biological variance conservation. The category scores of these two aspects were calculated by averaging the metrics within each category. An overall score for each integration method was computed

1239   using a 40/60 weighted mean of the category scores for batch effect removal and biological variance conservation

1240   (Methods). **f,** The spatial visualization of spots across the three spatial ATAC mouse embryo samples colored by ground

1241   truth spatial domains and **g,** the spatial clusters identified based on different integration methods. The first, second and

1242   third row of **f** and **g** denotes the samples from E12.5, E13.5 and E15.5 stages, respectively.

1243

1244

Fig. 6 | PRESENT-BC captures biological signals while eliminating batch effects in spatial RNA-ATAC data. a, The
quantitative evaluation of different integration methods on the four MISAR-seq mouse brain samples using 14 metrics
divided into two categories, namely batch effect removal and biological variance conservation. The category scores of
these two aspects were calculated by averaging the metrics within each category. An overall score for each integration
method was computed using a 40/60 weighted mean of the category scores for batch effect removal and biological variance
conservation. **b,** The UMAP visualization of spots across the four MISAR-seq mouse brain samples colored by sample
indices, **c,** ground truth domain labels and **d,** spatial clustering labels identified based on different integration methods.