Cross-modality representation and multi-sample integration of spatially resolved omics data

Zhen Li[1], Xuejian Cui[1], Xiaoyang Chen[1], Zijing Gao[1], Yuyao Liu[1], Yan Pan[1], Shengquan Chen[2], Hairong Lv[1], Lei Zhai[3,*], and Rui Jiang[1,*]

[1] Ministry of Education Key Laboratory of Bioinformatics, Bioinformatics Division at the Beijing National Research Center for Information Science and Technology, Center for Synthetic and Systems Biology, Department of Automation, Tsinghua University, Beijing 100084, China

[2] School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, China

[3] Department of Critical Rehabilitation，Beijing Bo'ai Hospital, China Rehabilitation Research Center, Beijing 100068, China

[*] Corresponding author: drzhailei@163.com (Lei Zhai), ruijiang@tsinghua.edu.cn (Rui Jiang)

Abstract

Spatially resolved sequencing technologies have revolutionized our understanding of biological regulatory processes within the microenvironment by accessing the states of genomic regions, genes and proteins as well as spatial coordinates of cells. However, discrepancies between different modalities and samples hinder the analysis of spatial omics data, necessitating the development of advanced computational methods. In this article, we propose PRESENT, an effective and scalable contrastive learning framework, for the cross-modality representation and multi-sample integration of spatial multi-omics, epigenomics and transcriptomics data. Through comprehensive experiments on spatial datasets, PRESENT demonstrates superior performance across various species, tissues, and technologies. Specifically, PRESENT effectively integrates spatial dependency and omics information simultaneously, facilitating the detection of spatially functional domains and the exploration of biological regulatory mechanisms. Furthermore, PRESENT can be extended for the integrative analysis of tissue samples across different dissected regions or developmental stages, promoting the identification of hierarchical structures from systematic and spatiotemporal perspectives.

Introduction

Single-cell sequencing technologies have enabled the characterization of states and activities within individual cells, promoting molecular cell biology and disease research in areas such as cell atlas production, tumor immunology, and cancer genetics[1]. However, an indispensable step of single-cell sequencing techniques involves the dissociation of cells from the tissue of interest, resulting in the loss of spatial information[2]. Complex biological regulatory processes and cellular activities within the microenvironment are driven by intricate cell-cell interactions mediated by ligand-receptor partners, which heavily depend on the spatial distance between cells[3-5]. Recently, rapid advancements in spatially resolved sequencing technologies have enabled the simultaneous profiling of both spatial and omics data, such as the spatially resolved transcriptomics (SRT)[6, 7], spatial epigenomics assay for transposase-accessible chromatin sequencing (ATAC-seq)[8, 9] and spatial multi-omics such as spatial CITE-seq[10], SPOTS[11] and MISAR-seq[12]. These technologies have accumulated substantial spatially resolved omics data in repositories[9, 12] and databases[13, 14], thereby granting us a new perspective to decipher biological regulatory mechanisms in the context of microenvironment.

However, spatial coordinate data and omics data exhibit significant modality discrepancies, posing substantial challenges for the analysis of spatial omics data. Specifically, spatial coordinate data anchors the relative positions of cells within tissues, while omics data depicts the states of chromatin regions, genes and proteins involved in the gene regulatory processes within cells. Additionally, different layers of omics data further present distinct patterns and distributions[15-18]. For example, compared with RNA-seq, ATAC-seq datasets suffer from larger dimensionality, more severe sparsity and higher dropout rates, while proteomics datasets, sequenced by antibody-derived tags (ADT), demonstrate opposite characteristics, requiring targeted modeling for each omics layer. Cross-modality representation, a critical step in spatial omics data analysis, addresses these challenges by integrating multi-modal data from individual spatial

datasets into a low-dimensional embedding matrix. This approach generates efficient and informative features for various downstream applications, including the detection of spatial domains, identification of domain-specific markers, and elucidation of domain-related functions, thereby facilitating the understanding of biological regulatory mechanisms within the microenvironment[19-21].

Due to technical limitations, such as size restrictions of the captured area during data acquisition, individual spatial sequencing samples typically focus on a specific dissected region or developmental stage[22, 23]. Consequently, discrepancies in dissection and placement prevent the direct comparison of spatial coordinates between different samples[23]. Additionally, the intricate coupling between batch effects and biological variations hinders the applications of single-sample representation methods to the joint embedding of multiple samples, impeding the exploration of tissues under different conditions[23-25]. Therefore, the integrative analysis of multiple samples, known as multi-sample integration, is another critical focus in the research of spatially resolved omics data. While facing greater challenges compared to representation of individual samples, multi-sample integration offers a more comprehensive and systematic perspective for deciphering the hierarchical structures of cellular organizations[22, 23, 26, 27]. For example, literatures have demonstrated that integrated analysis of samples under different stages reveals developmental dynamics in the early mouse embryo data[27, 28]. Moreover, recent studies have confirmed that integrating disease and normal samples facilitates the identification of disease-specific domains and promotes the investigation of disease onset and progression mechanisms[27, 29].

Many computational methods have been proposed to address the integrative analysis of SRT data[30], including single-sample representation algorithms like SpaGCN[31], STAGATE[32] and PAST[19], as well as multi-sample integration methods like STAligner[27], Stitch3D[33] and GraphST[22]. Despite the achievements, the development of algorithms for spatial ATAC-seq[8, 9] and spatial multi-omics data[11, 12, 34] remains in preliminary stage. Recently, SpatialGlue[35] was introduced as the first attempt for the single-sample

representation of spatial multi-omics data using an attentional feature fusion mechanism. Tian et al.

proposed a dependency-aware generative variational autoencoder framework for the single-sample

representation of SRT, spatial ATAC-seq and spatial CITE-seq data[36]. While these studies have addressed

the cross-modality representation of individual spatial epigenomics and multi-omics samples, they have

overlooked the integrated analysis of multiple samples across different conditions or time points,

impeding the understanding of tissue structures from a spatiotemporal perspective. To address these

challenges, there is an urgent need for advanced computational algorithms capable of integrating and

analyzing cross-modality and multi-sample spatially resolved omics data.

To this end, we propose PRESENT, an advanced and scalable contrastive learning framework for the

cross-modality representation and multi-sample integration of s**P**atially **RE**solved omics data, including

spatial multi-omic**S, E**pigenomics, tra**N**scrip**T**omics (Fig. 1). Through comprehensive experiments on

datasets generated by various spatial sequencing technologies, including spatial multi-omics[11, 12, 34],

epigenomics[8, 9], and transcriptomics[6, 7], we demonstrated that PRESENT can simultaneously capture

spatially coherent patterns and complementary multi-omics information, obtaining interpretable

representations for various downstream analyses, including spatial visualization and the detection of

spatial domains. Based on the Bayesian neural networks, PRESENT also offers the potential to

incorporate reference data generated by single-cell or spatial technologies, addressing issues including

low sequencing depth and signal-to-noise ratio associated with spatial data. Furthermore, PRESENT can

be extended for the integrative analysis of multiple samples across different dissection regions and

developmental stages, effectively eliminating batch effects while maintaining shared and sample-specific

biological variations, paving the way to explore the hierarchical tissue structures and sophisticated

cellular activities from systematic and spatiotemporal perspectives.

Results

Overview of PRESENT

Using a spatial multi-omics dataset that includes spatial coordinates, epigenomic profiles, transcriptomic expressions, and proteomic profiles, we illustrate the cross-modality representation architecture of PRESENT in Fig. 1a. Key challenges in the cross-modality representation of spatial multi-omics data include recognizing patterns within individual modalities and integrating multi-omics biological signals. Built on a multi-view autoencoder, PRESENT recognizes spatially biological variations contained in each omics layer via an omics-specific encoder consisting of graph attention neural networks[37] (GAT) and a Bayesian neural network[38] (BNN). The GAT module aims to incorporates spatial dependencies and omics information, while the BNN module, known for its effectiveness in non-linear regression problems[38], brings the potential to leverage reference data from existing sequencing data[19]. To further capture the distinct patterns of various omics layers, PRESENT employs omics-specific decoders which model the counts of transcriptomic and epigenomic data via zero-inflation negative binomial (ZINB) distribution[39] and zero-inflation Poisson (ZIP) distribution[40] (Methods), respectively. Moreover, to facilitate the integration of multi-omics biological signals, PRESENT fuses omics-specific embeddings from the encoders using a cross-omics alignment module which implements a multi-modal contrastive learning loss function, namely the inter-omics alignment (IOA) loss[41] (Methods).

Based on the above cross-modality representation framework, PRESENT can be further extended to the multi-sample integration of spatial multi-omics data, as illustrated in Fig. 1b. Critical challenges in multi-sample integration include addressing the spatial coordinate discrepancies between samples to capture cross-sample relationships[23] and distinguishing complexly coupled biological variations from technical batch effects[22, 26, 27, 33]. PRESENT adopts a two-stage training workflow to address these issues. In the first stage, PRESENT is pretrained based on the cross-sample graph and omics data acquired from

all the input samples. Specifically, the intra-sample edges of the cross-sample graph are established using

the spatial coordinates, while the inter-sample connections are built using omics information (Methods).

In the second stage, to align spots from multiple samples and eliminate technical batch effects from the

coupled biological variations, we introduce a contrastive learning strategy[42] and a batch-adversarial

learning strategy[43]. Additionally, we incorporate an intra-batch preserving loss during the training process

to maintain biological signals within each sample while integrating multiple samples. Moreover, the

second stage employs a cyclic iterative optimization process to refine the cross-sample graph using the

joint embeddings generated by the newly trained model at each epoch, enabling a progressively improved

characterization of cross-sample relationships (Methods).

On the other hand, PRESENT can be seamlessly adapted from spatial multi-omics data to spatial

single-omics data, such as SRT and spatial epigenomics data, by simplifying the multi-view autoencoder

framework into a single-view autoencoder (Methods). Through experiments on massive spatial datasets

generated by spatial multi-omics, epigenomics and transcriptomics technologies, we have demonstrated

that the joint embeddings obtained by PRESENT facilitate various downstream applications, including

spatial visualization, domain detection, the integrative analysis of horizontal slices from different

dissection regions, and vertical integration of samples from different developmental stages (Fig. 1c),

paving the way to explore the cellular activities from spatiotemporal perspectives.

PRESENT facilitates accurate spatial domain identification in spatial RNA-ADT data

Spatial domain identification, which aims to decipher tissue regions with distinct anatomical structures,

cell-type compositions, and cell-cell interactions[44, 45], is a fundamental step in revealing gene regulatory

processes and tissue functions in spatial omics data[31, 32]. We first collected a spatial RNA-ADT human

lymph node dataset generated by 10x Genomics Visium technology[35]. This dataset simultaneously

accessed the spatial locations, transcriptomic and proteomic expression of spots and was manually

annotated by the clinician group according to the histological image[35] (Fig. 2a). We benchmarked

PRESENT with SpatialGlue[35] and spaMultiVAE[36], two representation learning methods recently designed

for spatial multi-omics data, and several methods developed for single-cell RNA-ADT paired sequencing

data, including totalVI[46], CiteFuse[47] and Seurat-v4[48]. Based on the latent embeddings generated by these

methods, we implemented the Leiden algorithm[49] to acquire the same number of clusters as the ground

truth domains (Methods). Four commonly used metrics[50, 51], including adjusted rand index (ARI),

adjusted mutual information (AMI), normalized mutual information (NMI) and homogeneity (Homo),

were utilized to quantitatively evaluate the performance of spatial domain detection. As shown in Fig. 2b,

PRESENT achieved the best performance on all the four metrics for spatial domain identification. As a

qualitative evaluation and intuitive illustration of detected domains, we also demonstrated the uniform

manifold approximation and projection (UMAP) manifold structure[52] and spatial visualization results

(Fig. 2d, e, f). Specifically, CiteFuse and Seurat-v4 obtained unsatisfactory visualization results, where

domains such as medulla cords and sinuses scattered in the UMAP plot. spaMultiVAE detected stripe-like

spatial domains, resulting in biased spatial pattern identification in the lymph node tissue. Although

SpatialGlue and totalVI achieved relatively better results, they failed to recognize capsule from

pericapsular adipose tissue. In contrast, PRESENT successfully distinguished the pericapsular adipose

tissue (cluster 8), capsule (cluster 1), follicle (cluster 3) and cortex (cluster2 and cluster 10). Additionally,

we applied PRESENT to only a single omics layer of the human lymph node dataset, thus leveraging

expression information from only RNA data or ADT data, denoted as RNA-only and ADT-only,

respectively. As shown in Fig. 2c, PRESENT utilizing multiple omics layers (RNA & ADT) achieved

improved performance compared to the model that only leveraged RNA or ADT data, highlighting the

capabilities of PRESENT in integrating cross-modality spatial multi-omics information.

As the largest secondary lymphoid organ in the body, the spleen plays a significant role in

hematopoiesis and the clearance of erythrocytes, contributing significantly to immune functions[53]. We

further applied PRESENT to a SPOTS mouse spleen dataset[11], which measured spatial transcriptomic and

proteomic profiles simultaneously, to decipher the functionality of mouse spleen in the context of cellular

organizations. Through cross-modality representation and spatial clustering, PRESENT identified four

main spatial domains (Fig. 2g), denoted as the B cell-enriched domain, the T cell-enriched domain and

two macrophage-enriched subdomains, i.e., Mac1-enriched and Mac2-enriched domains, according to our

differentially expressed analysis (Fig. 2h, i). The differentially expressed analysis compared the protein

expression level of each domain with those of the rest domains based on the Wilcoxon rank-sum test[54]

(Fig. 2h), identifying protein markers distinguishing the B cell-enriched domain (CD3, CD4, CD8), the T

cell-enriched domain (CD19, B220, lgD) and the macrophage-enriched domain (F4-80, CD163, CD68),

consistent with the original study[11]. Specifically, the Mac1-enriched and Mac2-enriched domains had

nearly identical protein markers (Fig. 2h) as the macrophage markers identified in the original study[11],

making it hard to distinguish the two subdomains purely by ADT data. Therefore, we further conducted

differentially expressed analyses on the transcriptomic data between domains identified based on

PRESENT (Fig. 2i). Results demonstrated that Mac1-enriched domain could be marked by genes

including *Hbb-bt, Hbb-a1, Rsad2, Hbb-bs, Car2* and *Slc4a1*, while Mac-2 enriched domain could be

recognized with *Igkc, Jchain, Igha, Ighm, Gm2a* and *Cox8a.* The RNA expression levels between the

Mac1 and Mac2-enriched domains were also directly compared and numerous significantly differentially

expressed genes (DEGs) were found (Fig. 2j). These results showed that, despite the similarity of the two

macrophage-enriched subdomains in ADT expression levels, the distinction of transcriptomic patterns

between them could still be identified by PRESENT, emphasizing the ability of PRESENT in leveraging

complementary proteomic and transcriptomic information for effective domain identification.

To further decipher the functional distinction between the two macrophage-enriched subdomains, we

conducted Gene Ontology (GO) enrichment analysis based on the identified DEGs (Methods).

Specifically, for the Mac1-enriched domain, we enriched biological processes including erythrocyte

development, erythrocyte differentiation, erythrocyte homeostasis, myeloid cell development, myeloid

cell differentiation and myeloid cell homeostasis with false discovery rates (FDRs) of 1.29e-8, 5.48e-6,

6.70e-6, 3.59e-7, 8.71e-5 and 2.06e-5, respectively (Fig. 2k and Supplementary Table 1). Macrophages

are myeloid lineage cells which derive from bone marrow[55], reasonably explaining the enrichment of

development, differentiation and homeostasis of myeloid cell in the Mac1-enriched domain. In addition,

previous studies have stressed the importance of macrophages in the entire life of erythrocytes by

providing necessary factors for the survival, proliferation and development of erythrocytes[56, 57], agreeing

with the development, differentiation and homeostasis of erythrocyte found in the Mac1-enriched domain.

In comparison, the Mac2-enriched domain found biological processes including phagocytosis &

engulfment, plasma membrane invagination, membrane invagination, phagocytosis, humoral immune

response and lymphocyte mediated immunity with FDRs of 7.11e-5, 7.11e-5, 7.11e-5, 4.34e-4, 5.58e-4

and 5.94e-4, respectively (Fig. 2k and Supplementary Table 2). Phagocytosis and other related processes,

called the process of 'cell eating', are commonly considered as classical functions of macrophages[58, 59],

validating the enrichment of macrophages in the Mac2-enriched domain. On the other hand, compared

with the Mac1-enriched domain, the Mac2-enriched domain was spatially close to the B cell-enriched and

T cell-enriched domains (Fig. 2g), thereby explaining the enrichment of lymphocyte mediated immunity

process[60]. In conclusion, PRESENT effectively integrates multi-omics information and spatially coherent

patterns in spatial RNA-ADT data, thereby facilitating accurate spatial domain identification and

systematic exploration of biological implications in the identified domains.

PRESENT integrates spatially aware multi-omics biological signals in spatial RNA-ATAC data

Next, we collected MISAR-seq mouse brain datasets[12], which simultaneously measured the epigenomics and transcriptomics, i.e., paired spatial RNA-ATAC profiles, of four mouse brain samples from four developmental stages, involving E11.0, E13.5, E15.5 and E18.5. Compared with spatial-RNA-ADT co-profiling data, spatial RNA-ATAC datasets suffer from higher dimension and sparsity. We firstly compared the performance of PRESENT in spatial domain identification with the spatial multi-omics method SpatialGlue[35] and six methods for the representation of single-cell RNA-ATAC data, including MOFA+[61], scAI[62], scMDC[63], scMVP[21], Seurat-v4[48] and SnapATAC2[64]. Here spaMultiVAE[36], totalVI[46] and CiteFuse[47] were not used as they are specially designed for single-cell or spatial RNA-ADT data. As shown in Fig. 3a, PRESENT achieved overall the best spatial clustering performance, illustrating the its superiority in extracting spatial multi-omics information into effective cross-modality embeddings for accurate spatial domain identification. We also used PRESENT to obtain latent representations based on only ATAC data or RNA data, denoted as ATAC-only and RNA-only respectively, which led to decreased performance, highlighting the importance of integrating multi-omics information in deciphering biological variations in the context of cellular organizations (Fig. 3b).

Taking the E18.5 mouse brain sample with the most complex structures as an example (Fig. 3c), we intuitively visualized the latent embeddings learned by PRESENT using heatmap (Fig. 3d). The spots from different spatial domains could be successfully distinguished via the combination of several latent features or even a single feature in the latent feature space, including large domains such as cartilage 1-4, mantle zone of dorsal pallium (DPallm), ventricular zone of dorsal pallium (DPallv), diencephalon & hindbrain and subpallium 1, and small domains like midbrain and thalamus. We further demonstrated the UMAP visualization of latent representations obtained by different methods on the E18.5 mouse brain sample (Fig. 3e, f). Most baseline methods, including MOFA+, scAI, scMDC, scMVP and Seurat-v4,

could only roughly separate large domains such as DPallm, DPallv, diencephalon & hindbrain and

subpallium 1. SnapATAC2 could additionally recognize the cartilage 1 but still failed to distinguish

others. In comparison, spatially aware methods, including SpatialGlue and PRESENT, not only identified

large spatial domains such as diencephalon & hindbrain, subpallium 1, DPallm, DPallv, cartilage 1-4, but

also accurately distinguished the two smaller spatial domains, i.e., midbrain and thalamus. Furthermore,

domains identified by these two methods demonstrated enhanced spatial smoothness, highlighting the

importance of integrating spatial information (Fig. 3g). The UMAP and spatial visualization of other three

samples of the MISAR-seq mouse brain data[12] achieved similar results (Supplementary Figs. 1-4).

Additionally, we have conducted comprehensive experiments to investigate the impact of biological

signals from different omics on model performance to provide deeper insights into the model's ability to

extract multi-omics biological signals. Specifically, PRESENT utilizing only the transcriptomic data of

E11.0 brain sample, denoted as RNA-only, successfully captured the difference between the muscle and

the cartilage 2 domain, while the model based solely on epigenomic data, depicted as ATAC-only, failed

to do so (red circles in Fig. 3h). On the other hand, compared with the failure of PRESENT using only

transcriptomic data (RNA-only) in the E15.5 brain sample, the model based on the epigenomic data

(ATAC-only) contributed more to the differentiation between cartilage 2, cartilage 3 and mesenchyme

domains (blue circles in Fig. 3i). Therefore, relying solely on a single omics layer might lead to a biased

interpretation of biological signals in spatial domain detection. When taking both omics layers as input

(RNA & ATAC), PRESENT successfully captured variations among those domains in both E11.0 and

E15.5 mouse brain samples (Fig. 3h, i), highlighting the capabilities of PRESENT in leveraging spatially

aware transcriptomic & epigenomic multi-omics information for cross-modality representation and the

accurate identification of spatial domains from a comprehensive perspective.

PRESENT eliminates batch effects while preserving biological signals in spatial multi-omics data

The MISAR-seq mouse brain dataset simultaneously measured the transcriptomic and epigenomic

profiles of developmental brain samples from four stages, including E11.0, E13.5, E15.5 and E18.5,

providing a multi-omics landscape of the developmental mouse brain in the context of cellular

organizations[12]. However, varying experimental conditions and sequencing batches often introduce

potential technical biases, resulting in significant discrepancies in omics data across samples. These

discrepancies, commonly referred to as batch effects in single-cell studies, pose challenges to the

integrative analysis of multiple samples from different conditions and developmental stages[24, 26]. For

illustration, we applied joint principal component analysis (PCA) and the UMAP algorithm to raw RNA

and ATAC data of the four MISAR-seq mouse brain samples, revealing pronounced batch effects (Fig.

4a). Moreover, during the development of mouse brain, several functional structures might be specific to

certain developmental stages (e.g., primary brain), while others may persist throughout the entire

developmental process (e.g., midbrain and muscle), as shown in Fig. 4b. Therefore, the key challenges in

integrating the MISAR-seq mouse brain samples lied not only in the incorporation of spatial multi-omics

information and the correction of batch effects across samples, but also in the conservation of shared and

sample-specific biological variations.

Due to the lack of multi-sample integration methods specially designed for spatial multi-omics data,

we compared the integration performance of PRESENT with six methods developed for single-cell data

(MNN[65], Scanorama[25], Harmony[24], SCALEX[66], scBasset-BC[67] and SnapATAC2[64]) and three recently

proposed methods designed for SRT data (STAligner[27], Stitch3D[33] and GraphST[22]). Specifically, MNN,

Scanorama, Harmony and SCALEX could be applied to batch integration of either RNA-seq or ATAC-

seq data, denoted as RNA-* and ATAC-*, respectively, where * represents these methods. scBasset-BC

was featured as a sequence-based method specially designed for single-cell ATAC-seq data, while

STAligner, Stitch3D and GraphST were intended for integrative analysis of SRT data. SnapATAC2 has

provided guidance for incorporating multi-omics (i.e., transcriptomics & epigenomics) information for

integrated analysis, serving as a multi-sample integration workflow for single-cell multi-omics.

We first applied the UMAP algorithm to the joint embeddings generated by PRESENT and other

baseline methods for visualization, with the spots colored by batch indices (Fig. 4c and Supplementary

Fig. 5a) and ground truth domain labels (Fig. 4d and Supplementary Fig. 5b). PRESENT successfully

integrated spots from different developmental stages (red boxes in Fig. 4c) while maintaining the

biologically distinct patterns of DPallm, DPallv, subpallium 1 & 2, thalamus and muscle (red boxes in

Fig. 4d). Specifically, spots of muscle domain persisted throughout the entire life of mouse brain

development, while those of subpallium 1&2, DPallm, DPallv and thalamus only existed in specific

stages. The success of PRESENT in distinguishing these domains illustrated its capabilities in eliminating

batch effects while capturing shared and stage-specific biological variance. Additionally, we conducted

joint spatial clustering across the four stages based on the joint embeddings obtained by different

methods, as shown in Supplementary Fig. 6. Except for the scBasset-BC and SnapATAC2, other baseline

methods delivered limited clustering results. In contrast, PRESENT successfully detected not only large

domains like DPallm (cluster 7), but also small domains including thalamus (cluster 16) and muscle

(cluster 11). It is worth noting that when employing individual analysis of the sample from the E18.5

stage (Supplementary Fig. 4), PRESENT was unable to identify these small domains, underscoring the

advantages of integrative analysis across multiple samples.

Furthermore, to intuitively evaluate the integration and biological conservation capabilities of

PRESENT and other baseline methods, we systematically benchmarked them using 14 metrics, evenly

divided into two categories: batch effect removal and biological variance conservation, following a recent

benchmarking framework for single-cell omics data integration[68] (Methods). As shown in Fig. 4e,

scBasset-BC and SnapATAC2 provided satisfactory performance through incorporating sequence and

multi-omics information, respectively. PRESENT, which simultaneously integrated spatial dependency

and multi-omics information of all the samples, achieved the best performance in both batch effect

removal and biological variance conservation among all the 14 methods, illustrating its superiority in

multi-sample integration for spatial multi-omics data.

PRESENT facilitates effective cross-modal representation in spatial single-omics data

PRESENT is highly extensive and can be seamlessly adapted from spatial multi-omics data to spatial

single-omics data, such as spatial epigenomics and transcriptomics data, by simplifying the multi-view

autoencoder into a single-view model (Methods). We first accessed the performance of PRESENT on

spatial epigenomics data by collecting the spatial ATAC mouse embryo datasets which consist of three

developmental stages[9], including E15.5, E13.5 and E12.5. The datasets include six samples, with each

developmental stage containing two replicated samples, denoted as *-S1 and *-S2. Initially, we evaluated

the data quality of the six samples using logarithmic total fragment counts, denoted as log10(nFrags), and

transcription starting site (TSS) enrichment scores. The former metric represents the cellular sequencing

depth of samples[69], while the latter reflects the signal-to-noise ratio of datasets[70]. As shown in Fig. 5a, b,

the samples of E15.5 stage, i.e., E15.5-S1 and E15.5-S2, achieved the highest sequencing depth and

signal-to-noise ratio, while the E13.5-S1 sample and E12.5-S2 sample suffered from the lowest

sequencing depth and signal-to-noise ratio, respectively.

For spatial domain identification, we benchmarked PRESENT and other six baseline methods,

including three methods for single-cell ATAC-seq data (SCALE[71], cisTopic[72] and SnapATAC2[64]) and

three methods for SRT data (SpaGCN[31], STAGATE[32] and PAST[19]). As shown in Fig. 5c, PRESENT

achieved the best performance, with improvements of more than 8.60%, 26.13%, 25.91%, and 22.94%

compared to the second-best baseline method across the clustering metrics, respectively. Using the E15.5-S1 sample as an example, we further demonstrated the UMAP and spatial visualization of spots colored by ground truth domains (Fig. 5d, e) and the spatial clusters identified based on different methods (Fig. 5f). Baseline methods like cisTopic and STAGATE could only identify the forebrain domain. Although SCALE, SnapATAC2, SpaGCN and PAST roughly recognized the subdomains of central nervous system, i.e., forebrain, hindbrain and midbrain, the low-dimensional representations acquired by them are decentralized and scattered, impeding their performance for spatial clustering (Fig. 5c, f). In contrast, PRESENT effectively distinguished domains including the forebrain, hindbrain, midbrain, facial & bone, muscle & heart, and meningeal & peripheral nervous system (PNS) 2. Visualization of the remaining five samples obtained similar results (Supplementary Figs. 7-13), demonstrating the capabilities of PRESENT in cross-modal representation of spatial epigenomics data.

Despite the superiority of PRESENT over existing baseline methods, low sequencing depth and signal-to-noise ratio stills constrained the effectiveness of the model, as illustrated in Fig. 5a, b and Supplementary Fig. 13. A previous study has shown that integrating external reference data through BNN module can effectively address issues related to high noise levels in SRT data[19]. Inspired by this, we collected a single-nucleus ATAC-seq mouse embryo dataset involving cells from three developmental stages, including E12, E13 and E15, which was sequenced in conjunction with the spatial ATAC mouse embryo data[9]. We utilized the single-nucleus ATAC-seq dataset (denoted as PRESENT-SN) and the other five spatial ATAC samples (denoted as PRESENT-SP) as the reference data for PRESENT when analyzing each spatial ATAC mouse embryo sample (Methods). As shown in Fig. 5g and Supplementary Fig. 14, the introduction of prior information from reference data facilitated enhanced performance of PRESENT, especially in samples with lower sequencing depth and signal-to-noise ratio, such as the E13.5-S1 and E12.5-S2 mouse embryo samples. This finding was further validated by the UMAP

visualization of the two low-quality samples (Fig. 5h and Supplementary Figs. 9 and 12). Specifically,

PRESENT-SN and PRESENT-SP successfully identified spatial domains including liver, limb, forebrain,

hindbrain and midbrain, which other baseline methods and PRESENT failed to distinguish. For further

validation, we increased the noise level of datasets by converting lower fragment counts to zero with

growing dropout rates for each sample, according to a study that models the dropout events in single-cell

data[73]. As shown in Fig. 5i, PRESENT-SN and PRESENT-SP gained greater advantages as the noise level

increased, highlighting the improved benefits of integrating prior information from reference data with

the growing dropout rates in spatial ATAC-seq datasets.

We also carried out extensive experiments on SRT axolotl brain datasets generated by Stereo-seq,

which include six samples from different developmental stages, including stage 44, stage 54, stage 57,

juvenile, adult and metamorphosis[6]. The performance of PRESENT was benchmarked with several

recently developed SRT methods, involving SpaGCN[31], STAGATE[32], PAST[19] and SpatialPCA[74]. As

shown in Supplementary Figs. 15-21, experimental results demonstrated the consistent superiority of

PRESENT in cross-modal representation, spatial visualization and domain identification of SRT data. In

addition, utilizing an SRT regenerated axolotl brain data at 20 days post-injury as reference data[6],

PRESENT-SP again gained promoted performance through leveraging prior information contained in

reference data (Supplementary Fig. 15). In summary, PRESENT not only promotes effective cross-modal

representation in both spatial ATAC-seq and SRT data, but also empowers the integration of reference

data to address the low sequencing depth and signal-to-noise ratio of spatial datasets.

PRESENT enables vertical and horizontal integration of spatial single-omics samples

Integrated analysis of multiple samples across different developmental stages (vertically serial tissue

samples) and dissected areas (horizontally adjacent tissue samples), commonly referred to as vertical and

horizontal integration, provides a comprehensive approach for deciphering the hierarchical structures of cellular organizations[22]. We first evaluated the performance of PRESENT in the integrative analysis of vertically serial tissue samples, using the mouse embryo samples from three developmental stages (E15.5, E13.5 and E12.5) generated by spatial ATAC technique[9]. Due to the lack of methods specially designed for the multi-sample integration of spatial ATAC data, we compared PRESENT with six methods developed for single-cell data (MNN[65], Scanorama[25], Harmony[24], SCALEX[66], scBasset-BC[67] and SnapATAC2[64]) and two recently proposed SRT methods (STAligner[27] and GraphST[22]). We employed the benchmarking framework consisting of 14 metrics to simultaneously accessed the batch effect removal and biological variance conservation capabilities of these integration methods (Methods). As shown in Fig. 6a, PRESENT achieved the best overall score for multi-sample vertical integration, ranking first in biological variance conservation and second in batch effects removal among all the integration methods. The UMAP visualization of spot embeddings across different developmental stages also demonstrated that, compared with other methods, PRESENT gained better performance in aggregating spots from the same domain but in different batches (Supplementary Fig. 22). We further performed spatial clustering based on the joint embeddings produced by different methods. As shown in Fig. 6b, c, Scanorama and STAligner could hardly distinguish different spatial domains across different developmental stages. MNN, Harmony, SCALEX and SnapATAC2 provided improved results by roughly recognizing forebrain and liver in E12.5 and E13.5 stages, but they still failed in other domains. In comparison, PRESENT not only identified forebrain (cluster 1) and liver (cluster 5) in all the three stages, but also gained satisfactory results in domains including hindbrain (cluster 2) and midbrain (cluster 8).

Furthermore, we collected two horizontal adjacent SRT mouse brain samples generated by the 10x Genomics Visium platform, which involves a sagittal anterior dissected section and a sagittal posterior dissected section. We applied two SRT integration methods, namely STAligner[27] and GraphST[22], and

PRESENT to perform integrated analysis of these horizontally adjacent brain samples and utilized the

anatomic annotation of the sagittal region in postnatal day 56 (P56) mouse brain provided by Allen

Reference Atlas[75] to intuitively compare the performance of these methods (Fig. 6d). As shown in Fig. 6e-

g, compared with STAligner and GraphST, analytical results of PRESENT achieved improvements in

shared and sample-specific domain identification. For example, PRESENT captured the classical

hierarchical structures, i.e., from outer layer 1 to inner layer 6, of dorsal pallium in both sagittal anterior

and posterior sections, as illustrated in the red rectangle of Fig. 6d. In addition, PRESENT successfully

distinguished the typical cortex Ammonis (CA, cluster 18) and dentate gyrus (DG, cluster 25) structures

simultaneously in the hippocampus (black frames in Fig. 6d), which have been proved to be associated

with mood, anxiety and personality disorders caused by various factors[76]. We further conducted

differentially expressed analysis for domain CA and domain DG, acquiring marker genes including *C1ql2*

and *Hpca*, respectively (Supplementary Figs. 23 and 24). According to previous studies, *C1ql2* serves as a

marker gene for dentate gyrus granule cells (DGCs) enriched in the DG structure[77], while *Hpca* is the

encoding gene for a high-affinity calcium-binding protein restricted to the central nervous system,

abundant in pyramidal cells of the hippocampus CA region[78].

Moreover, PRESENT also exhibited great potential in capturing sample-specific functional domains

and uncovering biological implications in identified spatial domains. PRESENT successfully revealed the

sophisticated substructures of the cerebellar vermis, which only exists in the sagittal posterior section of

mouse brain and plays important roles in coordinating movements of the central body such as trunk, head,

and proximal limbs[79]. Specifically, PRESENT successfully detected the molecular layer (cluster 3),

Purkinje cell layer (cluster 22) and internal granular layer (cluster 17) of cerebellar vermis cortex and the

white matter (cluster 20) of cerebellar vermis. The differentially expressed genes in the corresponding

domains have also been validated in previous studies, such as *S100b*[80], *Car8*[81], *Atp1a1*[82] and *Mbp*[83]

(Supplementary Figs. 25-28). For example, *Car8* and *Atp1a1*, the mutation of which cause the motor

coordination defect and dominant Charcot-Marie-Tooth type 2, were proved to be highly expressed in the

granular layers and Purkinje cells in recent studies[81, 82], respectively. In summary, these experimental

results demonstrate the superior performance of PRESENT in the integrative analysis of both vertically

serial and horizontally adjacent spatial samples generated by various spatial omics technologies, thereby

enabling the discovery of biological insights within identified spatial domains across multiple samples.

Discussion

Recent breakthroughs in spatial omics sequencing technologies have enabled the simultaneous acquisition

of spatial coordinate information and the regulatory status of the biomolecules within cells, such as

chromatin regions, genes, and proteins, accumulating massive spatial omics data in repositories and

databases[13, 14]. In this article, we proposed PRESENT, an effective and scalable contrastive learning

framework, to address the cross-modality representation and multi-sample integration of spatial omics

data, including spatial multi-omics, epigenomics and transcriptomics data. To capture the patterns of

various omics layers, we designed omics-specific encoders and decoders for different omics data, where

the transcriptomic and epigenomic decoders model the fragment counts via the zero-inflation negative

binomial (ZINB) distribution[39] and zero-inflation Poisson (ZIP) distribution[40], respectively (Methods). In

addition, a cross-omics alignment module built on a contrastive learning inter-omics alignment loss

function[41] and graph attention networks[37] (GAT) were also introduced to capture the spatial dependency

and complementary multi-omics information (Methods). We further extended the framework to the multi-

sample integration of spatial omics data, addressing the challenges including the inconsistency of spatial

coordinates between samples as well as the intricate coupling between batch effects and biological

signals. Specifically, we adopted a contrastive learning-based inter-batch alignment strategy[42] and a

batch-adversarial learning strategy[43] to eliminate batch effects as well as an intra-batch preserving loss to

preserve biological signals simultaneously. To validate of the model design, we have conducted a

comprehensive ablation study, as shown in Supplementary Figs. 29 and 30.

Through extensive experiments on different species involving human, mouse and axolotl, we

highlighted the superior performance of PRESENT in cross-modality representation and multi-sample

integration of spatial omics data generated by various spatial sequencing technologies, encompassing

spatial multi-omics techniques (spatial RNA-Protein[11, 12] and spatial RNA-ATAC[12]) and spatial single-

omics technologies (spatial ATAC[9] and spatial transcriptomics[6]). Specifically, PRESENT effectively

integrated the spatial dependency and complex multi-omics information contained in spatial multi-omics

or spatial single-omics techniques (Figs. 2, 3 and 5). Additionally, PRESENT also showed great potential

for leveraging various reference data based on the BNN module to address issues related to low

sequencing depth and signal-to-noise ratio in spatial omics data (Fig. 5). More importantly, PRESENT

can be further extended to the integrative analysis of multiple spatially resolved samples across distinct

dissection areas and developmental stages, thereby promoting the detection of hierarchical structures and

biological functions within cellular organizations from spatiotemporal views (Figs. 4 and 6).

Nevertheless, several avenues remain for further improvement of PRESENT. First, Due to technical

and cost limitations, the number of unpaired spatial samples exceeds that of paired spatial samples[84],

emphasizing the importance of unpaired integration. While PRESENT can anchor shared spots to analyze

paired multi-omics sequencing data or leverage shared genomic features to integrate multiple spatial

samples, it still struggles with the integration of unpaired samples with distinct feature space[84], such as

the integration of two samples generated by spatial ATAC-seq and SRT techniques, respectively. Second,

explorative analyses based on single-cell data and spatial data are limited by lack of spatial information

and low spatial resolution, respectively. As an integrative framework for spatially resolved omics data,

PRESENT could be further extended to integrate single-cell and spatial data, computationally addressing

these technological limitations. Third, beyond integrating samples into the latent space, the alignment of

spatial coordinates in physical space also plays an important role in joint analysis of multiple spatial

samples[23, 26]. Finally, with the development of spatially resolved sequencing technology, imaged-based

spatial techniques, such as 10x Genomics Xenium, produce high-resolution histology image of dissected

samples, providing additional morphological features for enhanced analysis of spatial samples.

Methods

This study aims to investigate spatially functional domains as well as biological regulatory processes within microenvironments through incorporating cross-modality information, including spatial coordinate data and omics data, which involves epigenomic data generated by the assay for transposase-accessible chromatin with sequencing (ATAC), transcriptomic data measured by RNA sequencing (RNA), and proteomic data sequenced by antibody-derived tags (ADT). To achieve this goal, we have established a self-supervised contrastive learning framework, namely PRESENT, for the cross-modality representation and multi-sample integration of spatial multi-omics, epigenomics and transcriptomics data. Furthermore, we have also systematically evaluated the performance of PRESENT on massive real-world spatial datasets across various sequencing technologies, species and tissues.

Cross-modality representation of spatial multi-omics data

First, we detail the modeling approach of PRESENT for cross-modality representation of spatial multi-omics data, using a complex scenario as an example, which involves the comprehensive integration of spatial coordinate data, epigenomic data (ATAC data), transcriptomic data (RNA data), and proteomic data (ADT data). As shown in Fig. 1a, PRESENT is based on a multi-view autoencoder framework and comprises omics-specific encoders, a cross-omics alignment module, and omics-specific decoders, aiming to incorporate the cross-modality information into a low-dimensional feature matrix.

Multi-modality input of PRESENT. The input of PRESENT is a spatial coordinate matrix $M \in R^{n \times 2}$ and three fragment count matrices, namely the ATAC data $X^{ATAC} \in R^{n \times r}$, RNA data $X^{RNA} \in R^{n \times g}$ and ADT data $X^{ADT} \in R^{n \times p}$. Here n denotes the number of spots, r the number of genomic regions, g the number of highly variable genes and p the number of input proteins. Before feeding the spatial and omics data into the omics-specific encoders, we transform the spatial coordinate matrix M into an

undirected neighborhood graph $\boldsymbol{G}$ through adding undirected edges between all spots and their k nearest

neighbors, where k is set to 6 by default. Afterwards, we apply term frequency-inverse document

frequency (TF-IDF) normalization to ATAC data[85-88], library size normalization followed by log-

transformation to RNA data[19, 31, 32] and the centered log ratio (CLR) transformation to ADT data [17, 89, 90],

obtaining normalized omics data $\widetilde{X}^{ATAC} \in R^{n \times r}$, $\widetilde{X}^{RNA} \in R^{n \times g}$ and $\widetilde{X}^{ADT} \in R^{n \times p}$, respectively.

Omics-specific encoder. Graph neural networks, such as graph attention networks[37] (GAT), have

showcased successful application in integrating spatial and transcriptomic information for spatial

transcriptomics[19, 22, 31, 32]. On the other hand, research has proved that introducing weight uncertainty into

models through Bayesian neural networks[38] (BNN) improves generalization in nonlinear regression

problems[38] and offers an extension for leveraging prior information from existing sequencing data[19].

Inspired by these studies, we design an omics-specific encoder, consisting of a GAT module and a BNN

module, to capture spatially organized and biologically regulatory signals embedded in each omics layer.

In the following two subsections, taking the ATAC omics-specific encoder as an example, we delineate

the detailed design of GAT and BNN module.

Graph attention network. Given the input undirected neighborhood graph $\boldsymbol{G}$ and the normalized

ATAC matrix $\widetilde{X}^{ATAC} \in R^{n \times r}$, we denote the normalized accessibility vector of spot i in $\widetilde{X}^{ATAC}$ as

$\widetilde{x}_i^{ATAC} \in R^r$ and the spatial neighborhood of spot i as $\mathcal{N}_i$. Initially, we acquire the embedding from the

first hidden layer of spot i, i.e., $\boldsymbol{h}_i^1 \in R^{d_1}$ using a fully connected network (FCN):

$$\boldsymbol{h}_i^1 = \sigma(\boldsymbol{W}_1 \widetilde{x}_i^{ATAC} + \boldsymbol{b}_1)$$

where d_1 represents dimension of the first hidden layer, σ indicates the combination of layer

normalization and exponential linear unit (ELU) activation function, and $\boldsymbol{W}_1 \in R^{d_1 \times r}$ and $\boldsymbol{b}_1 \in R^{d_1}$

indicate the weight and bias parameters of the FCN, respectively. Afterwards, The second hidden-layer

embedding of spot i, namely $\boldsymbol{h}_i^2 \in R^{d_2}$, is obtained based on a graph attention layer[37], depicted as:

$$h_i^2 = \sigma\left(\sum_{j \in \mathcal{N}_i} \alpha_{ij} W_2 h_j^1\right)$$

$$\alpha_{ij} = \frac{\sigma_{LeakyReLU}\left(a^T\left(W_2 h_i^1 \oplus W_2 h_j^1\right)\right)}{\sum_{k \in \mathcal{N}_i} \sigma_{LeakyReLU}\left(a^T\left(W_2 h_i^1 \oplus W_2 h_k^1\right)\right)}$$

where α_{ij} is the attention score between spot i and j, d_2 denotes the dimension of the second hidden

layer, $W_2 \in R^{d_1 \times d_2}$ and $a \in R^{2d_2}$ are learnable parameters of the graph attention layer, $\sigma_{LeakyReLU}$

denotes the LeakyReLU activation function with a negative slope of 0.2, and $\oplus$ refers to the

concatenation function. Finally, h_i^2 undergoes transformation by another graph attention layer to obtain

$h_i^{gat} \in R^{d_{gat}}$ as the output, where d_{gat} denotes the output dimension of GAT module.

Bayesian neural network. According to a previous work, BNN module provides an opportunity to

leverage biological prior information from existing sequencing data to improve the analytical

performance in spatial transcriptomic data[19]. Inspired by this, PRESENT includes the BNN module,

which assumes that the parameters of weight matrix, denoted as $W \in R^{d_{bnn} \times r}$, follow Gaussian

distributions. Specifically, each element of W is governed by a Gaussian distribution parameterized by

the corresponding elements in the mean matrix μ^b and variance matrix σ^{b^2}, depicted as:

$$W_{ij} \sim \mathcal{N}\left(\mu_{ij}^b, \sigma_{ij}^{b^2}\right)$$

531 In addition, W_{ij} is also governed by a Gaussian prior parameterized by mean μ_{ij}^p and variance $\sigma_{ij}^{p^2}$.

532 Inheriting the original settings[19], we calculate the PCA projection weight matrix of normalized ATAC

533 matrix $\tilde{X}^{ATAC}$ as the mean matrix μ^p and utilize the standard deviation of all the elements in mean

534 matrix μ^p to construct the standard deviation matrix σ^p. The distance between the prior and the

535 distributions of the parameters in BNN module is restricted with Kullback-Leibler divergence (KLD):

$$\mathcal{L}_{BNN} = KLD(\mathcal{N}\left(\mu^b, \sigma^{b^2}\right), \mathcal{N}(\mu^p, \sigma^{p^2}))$$

It is notable that the parameters of prior Gaussian distributions, i.e., the mean matrix μ^p and the variance

matrix $\sigma_{ij}^{p^2}$, can also be constructed with reference ATAC data, enabling PRESENT to incorporate prior

information from existing ATAC-seq data generated by various technologies, including single-cell ATAC-

seq or spatial ATAC-seq. The unification of peak sets between reference ATAC-seq data and target ATAC-

seq data is conducted according to the merging objects vignette provided by the Signac package[91].

The output embeddings of spot i from BNN module, i.e., $\boldsymbol{h}_i^{bnn} \in R^{d_{bnn}}$, is denoted as:

543
$$\boldsymbol{h}_i^{bnn} = \boldsymbol{W}\tilde{\boldsymbol{x}}_i^{ATAC} = (\boldsymbol{\mu}^b + \boldsymbol{\eta}\odot\boldsymbol{\sigma}^b)\tilde{\boldsymbol{x}}_i^{ATAC}$$

544   where d_{bnn} denotes the output dimension of BNN module, $\boldsymbol{\eta}$ is the standard Gaussian noise matrix and

545     $\odot$ denotes element-wise product.

546   Finally, the output embedding of spot i from the ATAC omics-specific encoder, denoted as $\boldsymbol{h}_i^{ATAC} \in$

547     $R^{d_{ATAC}}$, is the concatenation of output embeddings from the GAT and BNN module:

548
$$\boldsymbol{h}_i^{ATAC} = \boldsymbol{h}_i^{gat}\oplus\boldsymbol{h}_i^{bnn}$$

where $d_{ATAC} = d_{gat} + d_{bnn}$ is the output dimension of omics-specific encoder for ATAC data.

Cross-omics alignment module. Based on the omics-specific encoders, the embeddings of different

omics layers are separately extracted. However, accurately deciphering sophisticated gene regulation

process requires integrating the status of biological macromolecules in cells from different omics

perspectives[15-18]. Inspired by a recent study that achieved multi-modal entity alignment through

contrastive learning[41], we designed a cross-omics alignment module to model the complex regulatory

interactions between different omics layers.

Given the omics-specific embeddings of spot i obtained by omics-specific encoders, including

$\boldsymbol{h}_i^{ATAC} \in R^{d_{ATAC}}$ for ATAC, $\boldsymbol{h}_i^{RNA} \in R^{d_{RNA}}$ for RNA and $\boldsymbol{h}_i^{ADT} \in R^{d_{ADT}}$ for ADT, we utilized a multi-

layer perception (MLP) module to fuse the embeddings from different omics layers to acquire a cross-

omics representation of spot i, denoted as:

560
$$\boldsymbol{e}_i = f_{MLP}(\boldsymbol{h}_i^{ATAC}\oplus\boldsymbol{h}_i^{RNA}\oplus\boldsymbol{h}_i^{ADT})$$

where $\boldsymbol{e}_i \in R^d$ represents the final cross-omics representation of spot i, f_{MLP} denotes the transformation

function of the MLP module, d is the latent dimension of PRESENT, and d_{ATAC}, d_{RNA} and d_{ADT} refers

to the output dimensions of omics-specific encoders for ATAC, RNA and ADT data, respectively. The

MLP module is composed of three FCNs, the first two of which are followed by a layer normalization and

an ELU activation function.

However, only the MLP module cannot perfectly incorporate omics-specific knowledge from

different omics layers into the final cross-omics embeddings[41], which is also illustrated in our ablation

study shown in Supplementary Fig. 29. Therefore, we design an inter-omics alignment (IOA) loss to

minimize the multi-omics directional KLD over the output distributions between the joint embeddings

and other omics-specific embeddings:

571
$$\mathcal{L}_{IOA} = \frac{1}{M}\left[\frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{ATAC})\right) + \frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{RNA})\right)\right.$$

572
$$\left. + \frac{1}{n}\sum_{i=1}^{n} KLD\left(q_{ed}^{(i)}(\boldsymbol{e}), q_{ed}^{(i)}(\boldsymbol{h}^{ADT})\right)\right]$$

573 where M represents the number of omics layers, $\boldsymbol{e}, \boldsymbol{h}^{ATAC}, \boldsymbol{h}^{RNA}$ and $\boldsymbol{h}^{ADT}$ denotes the joint, ATAC,

574 RNA and ADT embeddings respectively, and $q_{ed}^{(i)}(*)$ represents the embedding distribution of spot i

575 over other spots, defined as (taking $\boldsymbol{e}$ as an example):

576
$$q_{ed}^{(i)}(\boldsymbol{e}) = \left(\frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_1})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}, \frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_2})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}, \dots, \frac{\delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_n})}{\sum_{j=1}^{n} \delta_{ed}(\boldsymbol{e_i}, \boldsymbol{e_j})}\right)$$

577
$$\delta_{ed}(\boldsymbol{u}, \boldsymbol{v}) = exp(\frac{\boldsymbol{u}^T\boldsymbol{v}}{\tau})$$

where τ is a temperature parameter and set to 0.1 according to the original study[41].

Omics-specific decoder. The patterns and distributions of sequencing data vary significantly across

different omics layers, revealing complementary biologically regulatory information[15-18]. For example,

ADT sequencing data, which focuses on dozens of surface proteins, provides intuitive and reliable

quantification of cellular activities with low dimensionality, few dropout events and low sparsity[17]. In

contrast, RNA and ATAC data, which measure the expression levels of genes and accessibility scores of

genomic regions respectively, suffer from high dropout rates, dimensionality, and sparsity[92]. Therefore,

we designed omics-specific decoder for each modality to address the discrepancies between omics layers.

Inspired by a recently published study which modeled the fragment counts of single-cell ATAC-seq

data with Poisson distribution[40], we fit the fragment counts of spatial ATAC-seq data $\boldsymbol{X^{ATAC}} \in R^{n \times r}$ with

the zero-inflation Poisson (ZIP) distribution to accommodate the lower capture rate of spatial

technologies[9, 12]. Taking the fragment count vector of spot i, i.e., $\boldsymbol{x_i^{ATAC}}$, as an example, the omics-

specific decoder of ATAC data as well as the probability model is described as:

591
$$\boldsymbol{\pi_i} = \sigma_{sigmoid}\left(f_{MLP}^{ATAC,\pi}(\boldsymbol{e_i})\right) \quad \boldsymbol{\rho_i} = f_{MLP}^{ATAC,\rho}(\boldsymbol{e_i}) \quad \boldsymbol{\hat{x}_i^c} = f_{MLP}^{ATAC,x}(\boldsymbol{e_i})$$

592
$$\boldsymbol{\omega_i} = softmax(\boldsymbol{\rho_i} + \boldsymbol{r^c})$$

593
$$\boldsymbol{\lambda_i} = l_i \cdot \boldsymbol{\omega_i}$$

where $\sigma_{sigmoid}$ represents the sigmoid activation function, $\boldsymbol{r^c}$ is designed to capture region specific bias

such as peak length, $l_i = \sum_{j=1}^{r} x_{ij}^c$ refers to the total fragment counts of spot i, $f_{MLP}^{ATAC,\pi}(*)$, $f_{MLP}^{ATAC,\rho}(*)$

and $f_{MLP}^{ATAC,x}(*)$ are three output layers of $f_{MLP}^{ATAC}(*)$, the MLP module of ATAC omics-specific decoder.

Specifically, $f_{MLP}^{ATAC,\pi}(*)$ and $f_{MLP}^{ATAC,\rho}(*)$ output parameters of the ZIP probability model, while

$f_{MLP}^{ATAC,x}(*)$ outputs reconstructed normalized profiles of ATAC data. Then for each region $j = 1,2,...,r$,

the fragment count of i-th spot x_{ij}^{ATAC} satisfies:

600
$$x_{ij}^{ATAC} \sim ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right)$$

601
$$ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right) = \pi_{ij}\delta_0\left(x_{ij}^{ATAC}\right) + (1 - \pi_{ij})Poisson(x_{ij}^{ATAC}; \lambda_{ij})$$

602
$$\delta_0\left(x_{ij}^{ATAC}\right) = \begin{cases} 1 \ if \ x_{ij}^{ATAC} \ equals \ 0 \\ 0 \ otherwise \end{cases} \quad Poisson\left(x_{ij}^{ATAC}; \lambda_{ij}\right) = \frac{\lambda_{ij}^{x_{ij}^{ATAC}} e^{-\lambda_{ij}}}{x_{ij}^{ATAC}!}$$

603   Therefore, the negative log likelihood (NLL) loss function of the ZIP probability model is:

604
$$\mathcal{L}_{NLL}^{ATAC} = \frac{1}{nr}\sum_{i=1}^{n}\sum_{j=1}^{r}\left[-log \ ZIP\left(x_{ij}^{ATAC}; \pi_{ij}, \lambda_{ij}\right) + \gamma\pi_{ij}^2\right]$$

where the $\gamma\pi_{ij}^2$ refers to a regularization term and γ is set to 0.5 by default. The reconstruction loss of

ATAC omics layer is measured with mean squared error (MSE):

$$\mathcal{L}_{MSE}^{ATAC} = \frac{1}{nr}\sum_{i=1}^{n}\sum_{j=1}^{r}\left(\hat{x}_{ij}^{ATAC} - \tilde{x}_{ij}^{ATAC}\right)^{2}$$

Similarly, we also model the counts of RNA data $X^{RNA} \in R^{n \times g}$ using the zero-inflation negative

binomial (ZINB) distribution according to a single-cell RNA-seq analytical research[39], denoted as:

$$\boldsymbol{\pi_i} = \sigma_{sigmoid}\left(f_{MLP}^{RNA,\pi}(\boldsymbol{e_i})\right) \quad \boldsymbol{\mu_i} = \sigma_{softplus}\left(f_{MLP}^{RNA,\mu}(\boldsymbol{e_i})\right)$$

$$\boldsymbol{\theta_i} = \sigma_{softplus}\left(f_{MLP}^{RNA,\theta}(\boldsymbol{e_i})\right) \quad \hat{\boldsymbol{x}}_i^r = f_{MLP}^{RNA,x}(\boldsymbol{e_i})$$

where $\sigma_{softplus}$ refers to the softplus activation function, $f_{MLP}^{RNA,\pi}(*)$, $f_{MLP}^{RNA,\mu}(*)$, $f_{MLP}^{RNA,\theta}(*)$ and

$f_{MLP}^{RNA,x}(*)$ are four output layers of f_{MLP}^{RNA}, the MLP module of the RNA omics-specific decoder. Then the

count of spot i on gene j satisfies the following model:

$$x_{ij}^{RNA} \sim ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right)$$

$$ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right) = \pi_{ij}\delta_0\left(x_{ij}^{RNA}\right) + (1-\pi_{ij})NB(x_{ij}^{RNA}; \mu_{ij}, \theta_{ij})$$

$$NB\left(x_{ij}^{RNA}; \mu_{ij}, \theta_{ij}\right) = \frac{\Gamma\left(x_{ij}^{RNA} + \theta_{ij}\right)}{\Gamma(\theta_{ij})}\left(\frac{\theta_{ij}}{\theta_{ij} + \mu_{ij}}\right)^{\theta_{ij}}\left(\frac{\mu_{ij}}{\theta_{ij} + \mu_{ij}}\right)^{x_{ij}^{RNA}}$$

The NLL loss function of the ZINB probability model for RNA data can be denoted as:

$$\mathcal{L}_{NLL}^{RNA} = \frac{1}{ng}\sum_{i=1}^{n}\sum_{j=1}^{g}\left[-\log ZINB\left(x_{ij}^{RNA}; \pi_{ij}, \mu_{ij}, \theta_{ij}\right) + \gamma\pi_{ij}^2\right]$$

where the regularization parameter γ is also set to 0.5 by default. In addition, the reconstruction loss of

RNA omics layer is depicted as:

$$\mathcal{L}_{MSE}^{RNA} = \frac{1}{ng}\sum_{i=1}^{n}\sum_{j=1}^{g}\left(\hat{x}_{ij}^{RNA} - \tilde{x}_{ij}^{RNA}\right)^2$$

Compared with ATAC and RNA data, lower dropout rates, dimensionality and sparsity make the

analysis of ADT data less challenging. Therefore, the ADT omics-specific decoder is built on the MLP

module with one output layer to reconstruct the normalized expression level of proteins, denoted as:

$$\hat{\boldsymbol{x}}_i^{ADT} = f_{MLP}^{ADT}(\boldsymbol{e_i})$$

$$\mathcal{L}_{MSE}^{ADT} = \frac{1}{np}\sum_{i=1}^{n}\sum_{j=1}^{p}\left(\hat{x}_{ij}^{ADT} - \tilde{x}_{ij}^{ADT}\right)^2$$

Therefore, the overall NLL loss and reconstruction loss of PRESENT are summarized as:

$$629 \quad \mathcal{L}_{NLL} = \mathcal{L}_{NLL}^{ATAC} + \mathcal{L}_{NLL}^{RNA}$$

$$630 \quad \mathcal{L}_{MSE} = \mathcal{L}_{MSE}^{ATAC} + \mathcal{L}_{MSE}^{RNA} + \mathcal{L}_{MSE}^{ADT}$$

631   **Loss function for cross-modality representation of spatial multi-omics data.** Based on the above

632   descriptions, the overall loss function for cross-modality representation of spatial multi-omics data

633   consists of four parts, including the loss of BNN module, the inter-omics alignment loss, the negative log

634   likelihood loss of ZIP and ZINB probability model, and the mean squared error loss for reconstruction of

635   normalized omics profiles, denoted as:

$$636 \quad \mathcal{L}_{repre} = \alpha_1 \mathcal{L}_{BNN} + \alpha_2 \mathcal{L}_{IOA} + \alpha_3 \mathcal{L}_{NLL} + \alpha_4 \mathcal{L}_{MSE}$$

where α_1, α_2, α_3 and α_4 are coefficients of the corresponding loss term. The default values of α_1, α_2,

α_3 and α_4 are all set to 1.

Multi-sample integration of spatial multi-omics data

Multi-sample integration offers a comprehensive perspective for deciphering the structures of cellular

organizations[22, 23, 26, 27]. For example, integrated analysis of samples under different stages can reveal

developmental dynamics[27, 28], while integrating disease and normal samples facilitates the identification

of disease-specific domains and promotes the investigation of disease mechanisms[27, 29]. Therefore, we

have extended the PRESENT framework to multi-sample integration of spatial multi-omics data and

further incorporated mechanisms to address the discrepancies of spatial coordinates as well as batch

effects between samples. Overall, the multi-sample integration workflow based on the PRESENT

framework involves two stages. The first stage serves as a warmup to pretrain the PRESENT model,

while the second stage leverages a cyclic iterative optimization progress to eliminate batch effects while

preserving biological variations across samples. Analogous to cross-modality representation of spatial

multi-omics data, we use a scenario of integrating two spatial multi-omics sequencing samples as an

example to illustrate the workflow of multi-sample integration with PRESENT, where ATAC, RNA, ADT,

and spatial coordinate data are simultaneously provided for each sample (Fig. 1b).

Feature sets unification. The first step in multi-sample integration is to align the feature sets of different

sequencing samples, ensuring that all samples share a common feature space. For ATAC data, following

the merging objects vignette provided by the Signac package[91], a unified set of peak regions is first

obtained by merging all intersecting peaks from all the samples. The fragment counts of each spot within

the unified peak set are then quantified. For RNA and ADT data, whose feature sets are predefined genes

or proteins, we directly leverage the intersection of gene or protein sets of all the samples as the unified

gene or protein set. After the gene intersection step, highly variable genes will be selected based on the

unified gene set. For convenience, we denoted the ATAC, RNA, ADT and spatial data of the two samples

as $X_1^{ATAC} \in R^{n_1 \times r}$, $X_2^{ATAC} \in R^{n_2 \times r}$, $X_1^{RNA} \in R^{n_1 \times g}$, $X_2^{RNA} \in R^{n_2 \times g}$, $X_1^{ADT} \in R^{n_1 \times p}$, $X_2^{ADT} \in R^{n_2 \times p}$,

$M_1 \in R^{n_1 \times 2}$ and $M_2 \in R^{n_2 \times 2}$, respectively. Here, r represents the number of unified peak regions, g

the number of highly variable genes, p the number of unified proteins, and n_1 and n_2 the number of

spots of the two samples. The unified ATAC, RNA, ADT data are then concatenated and preprocessed

using the methods described in the cross-modality representation section, obtaining normalized data, i.e.,

$\widetilde{X}^{ATAC} \in R^{n \times r}$, $\widetilde{X}^{RNA} \in R^{n \times g}$, $\widetilde{X}^{ADT} \in R^{n \times p}$ and $M \in R^{n \times 2}$, where $n = n_1 + n_2$ represents the number

of spots from the two samples.

Cross-sample graph construction. Due to inconsistent dissection and placement of sequencing samples,

the spatial coordinates between these samples cannot be directly compared with each other, resulting in

the failure of cross-modality representation methods in capturing relationships across samples[23, 26]. To

address this gap, we proposed a strategy to construct a cross-sample graph. Firstly, we find intra-sample

spatial neighbors for each spot based on the spatial coordinate matrix, building undirected edges between

all spots and their k nearest neighbors in Euclidean space from the same sample. Secondly, we find inter-

sample neighbors based on omics features, including ATAC, RNA and ADT profiles, to establish inter-

sample undirected edges between all spots and their k most similar neighbors in cosine space from other

samples. Note that the parameter k is set to 6 by default. Finally, we acquire a cross-sample graph which

consists of both intra-sample edges and inter-sample edges.

First-stage pretraining. In the first stage, we pretrain the PRESENT framework based on the cross-

sample graph and the concatenated omics data of different samples, including $\widetilde{X}^{ATAC} \in R^{n \times r}$, $\widetilde{X}^{RNA} \in$

$R^{n \times g}$ and $\widetilde{X}^{ADT} \in R^{n \times p}$. To integrate batch information into the PRESENT model, we concatenate the

one-hot encoded batch labels with the output of the cross-omics alignment module, using the combined

matrix as the input of the omics-specific decoders. The objective function and training process of

PRESENT model during the first pretraining stage are identical to those used for cross-modality

representation of individual samples, as detailed in the previous section.

Second-stage integration. The technical noise between different samples, i.e., batch effects, have not

been fully addressed in the first stage. Therefore, we incorporate a contrastive learning strategy and a

batch-adversarial learning strategy in the second stage to eliminate the batch effects across samples while

preserving the biological signals simultaneously, as detailed in the following subsections.

Inter-batch alignment. We designed an inter-batch alignment (IBA) loss based on contrastive

learning[42], aiming to align spots with similar omics features across different samples. Specifically, we

first find mutual nearest neighbors from other samples as the positive set of for spot i, denoted as N_i^{pos},

referring to the inter-sample neighbors that are connected to spot i in the cross-sample graph. Second, we

randomly select $1/4$ proportion of spots from the same sample as the negative set of spot i, depicted as

N_i^{neg}. Finally, the IBA loss of spot i is:

696
$$\mathcal{L}_{IBA}^{(i)} = -log \frac{\sum_{j \in N_i^{pos}} exp(\delta_{sim}(e_i, e_j))}{\sum_{j \in N_i^{pos}} exp(\delta_{sim}(e_i, e_j)) + \sum_{j \in N_i^{neg}} exp(\delta_{sim}(e_i, e_j))}$$

697 where $e_i \in R^d$ denotes the latent representation of spot i, d is the output dimension of PRESENT and

698 δ_{sim} denotes the cosine similarity. The inter-batch alignment loss of all the spots is:

699
$$\mathcal{L}_{IBA} = \frac{1}{n} \sum_{i=1}^{n} \mathcal{L}_{IBA}^{(i)}$$

Batch-adversarial learning. Inspired by a domain-adversarial learning research[93] and its's success in

multi-batch integration of single-cell ATAC-seq data[43], we implement the batch-adversarial learning

strategy to remove batch effect across spatial samples. Specifically, we introduce a batch-discriminator,

aiming to predict the batch index of each spot based on its latent embedding, while the omics-specific

encoders and the cross-omics alignment module are treated as the generator, aiming to produce latent

embeddings without batch effect, thereby confusing the batch-discriminator in an adversarial manner. The

loss function of the batch-adversarial learning is:

707
$$\mathcal{L}_{ADV} = \max_{G} \min_{D} CE(P, T)$$

where G refers to the generator, D denotes the batch-discriminator, CE represents the cross-entropy

loss, P denotes the predicted batch indices and T refers to the ground truth batch labels.

Intra-batch preserving. The contrastive learning-based inter-batch alignment strategy and the batch-

adversarial learning strategy aim for batch correction across samples. However, a key challenge of multi-

sample integration lies in eliminating batch effects while simultaneously preserving the biological signals

within each individual sample. We therefore proposed an intra-batch preserving (IBP) loss to maintain the

biological pattern of latent representations within each sample during training process, denoted as:

715
$$\mathcal{L}_{IBP} = \frac{1}{S} \sum_{k=1}^{S} \frac{1}{|s_k|} \sum_{i \in s_k} KLD \left(q_{s_k}^{(i)}(e^{(t)}), q_{s_k}^{(i)}(e^{(t-1)}) \right)$$

716 where S represents the number of samples, s_k denotes the spots in sample k, $|s_k|$ denotes the number

717 of spots in sample k, $e^{(t)}$ and $e^{(t-1)}$ refer to the joint embeddings of all spots in current epoch and the

718 last epoch, respectively. $q_{s_k}^{(i)}(*)$ represents the embedding distribution of spot i over other spots within

719   sample k, defined as (taking the embeddings of current iteration $e^{(t)}$ as an example):

720
$$q_{s_k}^{(i)}\left(e^{(t)}\right) = \left(\frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(1)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}, \frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(2)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}, \ldots, \frac{\delta_{ed}\left(e_i^{(t)}, e_{s_k^{(|s_k|)}}^{(t)}\right)}{\sum_{j=1}^{|s_k|} \delta_{ed}\left(e_i^{(t)}, e_{s_k^{(j)}}^{(t)}\right)}\right)$$

721
$$\delta_{ed}(u, v) = exp(\frac{u^T v}{\tau})$$

where $s_k^{(j)}$ denotes the j-th spot in sample k, τ is a temperature parameter and set to 0.1 by default[41]. As

shown in the above formula, the IBP loss is designed to preserve the overall manifold structure of spots

within each sample during training process.

Training process of the second stage. The training process of the second stage is carried out in a

cyclic iterative manner. First, we obtain the joint embeddings of multiple samples using the trained

PRESENT model. Next, we construct a cross-sample graph according to the strategy described above,

where intra-sample connections are established based on the spatial coordinates of spots and the inter-

sample edges are based on the cosine similarity between the newly acquired joint embeddings of spots

from different samples. Finally, we train the PRESENT model for one epoch using the newly constructed

cross-sample graph. This cyclic iterative optimization process, aiming to iteratively refine the cross-

sample graph, is repeated until the model converges. When updating the PRESENT model, we first

update the parameters of the batch-discriminator using the cross-entropy loss $\mathcal{L}_{ADV}^{disc} = \beta_3 CE(P, G)$, with

parameters of the generator (i.e., the omics-specific encoders and the cross-omics alignment module) and

omics-specific decoders fixed. Next, we fix the parameters of batch-discriminator and update the

parameters of generator and omics-specific decoders based on an overall loss function, including the

cross-modality representation loss, the inter-batch alignment loss, the intra-batch preserving loss and the

negative cross-entropy loss for the batch-adversarial learning strategy:

739
$$\mathcal{L}_{integ} = \mathcal{L}_{repre} + \beta_1 \mathcal{L}_{IBA} + \beta_2 \mathcal{L}_{IBP} + \beta_3 \mathcal{L}_{ADV}^{gen}$$

$$\mathcal{L}_{ADV}^{gen} = -CE(P, G)$$

where β_1, β_2 and β_3 are coefficients of the corresponding loss function and set to 1 by default.

Applications of PRESENT to spatial single-omics data

PRESENT is highly versatile and can be seamlessly adapted from spatial multi-omics data to spatial

single-omics data, such as SRT and spatial epigenomics data, by simplifying the multi-view autoencoder

framework into a single-view autoencoder. Specifically, for cross-modality representation or multi-sample

integration of spatial single-omics data, only the respective omics-specific encoder and decoder are

required. For example, only the ATAC omics-specific encoder and decoder are required when modeling

spatial single-omics ATAC-seq data, while only the RNA omics-specific encoder and decoder are needed

when modeling spatial transcriptomic data. Additionally, both the cross-omics alignment module and the

associated inter-omics alignment loss are excluded from the PRESENT model in these cases, as spatial

single-omics data consist of only a single omics layer.

Implementation details

With the development of spatial omics sequencing technologies, the number of spots in sequenced tissue

has been quickly increasing from thousands to tens and even hundreds of thousands[9, 10, 13, 14, 94].

PRESENT is built upon graph neural networks, which face challenges such as memory explosion and

neighborhood expansion when the number of nodes in graph grows large[95, 96]. To address these problems,

we adopted necessary sparse adjacency matrix implementation based on the PyG framework when

deploying our model. Additionally, we implemented the ClusterGCN algorithm to divide the entire graph

into multiple small subgraphs[95], which enables our model to be trained on GPUs with limited memory

and extend to larger spatial omics data.

The input dimensions of ATAC and RNA data are generally much higher than the ADT data, resulting

in different hidden dimension settings for them. Specifically, the hidden and output dimensions of omics-

specific encoders for ATAC and RNA data were (1024, 512) and 50, respectively, while those of omics-

specific encoder for ADT data were $\left(2d_{adt}^{in}/3 + d_{adt}^{out}/3, \ d_{adt}^{in}/3 + 2d_{adt}^{out}/3 \right)$ and d_{adt}^{out}. Here d_{adt}^{in}

denotes the input dimension of ADT data and $d_{adt}^{out} = \min\left(50, d_{adt}^{in}\right)$ refers to the output dimension of

omics-specific encoder for ADT data. The final output dimension of PRESENT was set to 50 by default.

Additionally, the hidden dimensions of omics-specific decoders were symmetric to the corresponding

omics-specific encoders, i.e., $\left(2d_{adt}^{in}/3 + d_{adt}^{out}/3, \ d_{adt}^{in}/3 + 2d_{adt}^{out}/3 \right)$ for ADT data and (512, 1024) for

ATAC and RNA data. We trained the model for at most 100 epochs using the Adam optimizer, updating

the parameters with an initial learning rate of 1e-3 and a weight decay of 1e-4.

Data collection and description

10x Visium RNA-Protein human lymph node data. The human lymph node dataset, generated by 10x

Genomics Visium transcriptomics and proteomics co-profiling technology, was collected from a spatial

multi-omics study[35]. This dataset was manually annotated by the clinician group according to the

histological image[35], containing the spatial coordinates and the expression level of 18,085 genes and 31

proteins on 3,484 spots.

SPOTS mouse spleen data. The mouse spleen data[11] was generated by a high-throughput spatial

transcriptomics and proteomics co-profiling technology, namely Spatial PrOtein and Transcriptome

Sequencing (SPOTS). This dataset provides the spatial coordinates and the expression level of 32,285

genes and 21 proteins on 2,568 spots.

MISAR-seq mouse brain data. The MISAR-seq mouse brain data[12] provides the paired sequencing data

of spatial gene expression and chromatin accessibility during mouse brain development. We collected

four manually annotated samples from original study[12], including E11.0 (1,258 spots), E13.5 (1,777

spots), E15.5 (1,949 spots) and E18.5 (2,129 spots). The manual annotations were produced with

reference to Kaufman's Atlas of Mouse Development and Allen Brain Atlas[12]. We called peak for each

sample based on the MACS2 v2.2.9.1 package, obtaining 242,024, 271,126, 265,014 and 294,734 peak

regions for downstream analyses, respectively.

Single-nucleus ATAC-seq and spatial ATAC mouse embryo data. The single-nucleus ATAC-seq

mouse embryo data[9] measured the chromatin accessibility of 1,879 cells from three developmental stages,

including E12, E13 and E15. The spatial ATAC mouse embryo data[9] is composed of six samples,

including E12.5-S1 (3,274 spots), E12.5-S2 (3,903 spots), E13.5-S1 (3,160 spots), E13.5-S2 (2,970

spots), E15.5-S1 (2,246 spots) and E15.5-S2 (2,234 spots). Associated domain annotations were also

provided within these samples, which have been validated through extensive experiments in the original

study[9]. The single-cell ATAC-seq datasets and spatial ATAC datasets were merged and 248,047 peak

regions were called based on MACS2 v2.2.6 package.

Stereo-seq axolotl brain data. We collected the Stereo-seq axolotl brain developmental datasets from a

recent study which explored the regeneration and development of axolotl telencephalon using spatial

transcriptomic atlas[6]. The datasets consist of six sections from developmental stage 44 (1,477 spots),

stage 54 (2,929 spots), stage 57 (4,410 spots), juvenile (11,698 spots), adult (8,243 spots), and

metamorphosis (7,441 spots), measuring the expression level of 12,704 genes. These samples were

manually annotated with reference to eHistology Kaufman Annotations and Allen Brain Atlas[12].

10x Visium mouse brain data. The spatial transcriptomic postnatal day 56 (P56) mouse brain data,

including a sagittal anterior section and a sagittal posterior section, were collected from the 10x Genomics

Visium platform. The mouse sagittal anterior section contains the expression level of 32,285 genes on

2,823 spots, while the posterior section provides the expression level of 32,285 genes on 3,289 spots.

Experiment setup and baseline methods

Cross-modality representation of 10x Visium RNA-Protein human lymph node data. We excluded

genes or proteins with zero expression across all spots. Baseline methods includes totalVI[46], CiteFuse[47],

Seurat-v4[48], SpatialGlue[35] and spaMultiVAE[36], all implemented following their respective tutorials.

Cross-modality representation of MISAR-seq mouse brain data. We filtered out peaks accessible in

less than 3% spots and genes with zero expression in all spots for quality control. The filtered count

matrices of ATAC data and RNA data along with the spatial coordinate matrix were then fed into the

workflow of PRESENT and other baseline methods, involving SpatialGlue[35], MOFA+[61], scAI[62],

scMDC[63], scMVP[21], Seurat-v4[48] and SnapATAC2[64]. Note that totalVI[46], CiteFuse[47] and spaMultiVAE[36]

were not used as they are specially designed for single-cell or spatial RNA-ADT data. All the baseline

methods were implemented following the respective tutorials and default settings. Additionally, following

the workflow of processing single-cell[97] and spatial[19, 32, 74] RNA-seq data, PRESENT selected 3,000

highly variable genes as the transcriptomic input for further analysis.

Multi-sample integration of MISAR-seq mouse brain data. We merged the peak sets of different

mouse brain samples according to the merging objects vignette provided by the Signac package[91], which

first obtained a unified peak set by merging all intersecting peaks from all the samples and then quantified

the fragment count of each spot on the newly obtained peak set. Consistent with the experimental settings

in cross-modality representation, peaks accessible in less than 3% spots in the concatenated matrix were

filtered out and 3,000 highly variable genes of transcriptomic data were selected for PRESENT. Due to

the lack of methods specially designed for the integration of spatial multi-omics data, we compared the

batch effect removal and biological variance conservation performance of PRESENT with six methods

developed for single-cell data (MNN[65], Scanorama[25], Harmony[24], SCALEX[66], scBasset-BC[67] and

SnapATAC2[64]) and three recently proposed SRT multi-sample integration methods (STAligner[27],

Stitch3D[33] and GraphST[22]). Specifically, MNN, Scanorama, Harmony, SCALEX could be applied to both

transcriptomic and epigenomic data, denoted as RNA-* or ATAC-*, respectively, where * denotes

respective methods. MNN, Scanorama and Harmony performed batch correction based on embeddings

extracted from RNA or ATAC data, which were accomplished based on the workflow of SCANPY[97] and

EpiScanpy[98], respectively. All the methods were implemented using default settings.

Cross-modality representation of spatial ATAC mouse embryo data. The spatial ATAC mouse embryo

data firstly went through a quality control process, where peaks accessible in less than 0.5% spots were

filtered out. Using the filtered ATAC count matrices as input, we then benchmarked the performance of

PRESENT with four methods designed for single-cell ATAC-seq data, including SCALE[71], cisTopic[72] and

SnapATAC2[64], and three methods designed for SRT data, i.e., SpaGCN[31], STAGATE[32] and PAST[19]. All

the baseline methods were implemented using their default settings, except for STAGATE and PAST. In

original tutorials, the two SRT methods adopted a feature selection process which is specially designed

for transcriptomic data and only selected 3,000 features, resulting significantly lower performance

(Supplementary Fig. 31). Since other methods utilized all the input peaks for dimension reduction, we

ignored the feature selection step of these two methods for fairness and better performance.

Cross-modality representation of Stereo-seq axolotl brain data. The genes with zero expression in all

spots were filtered out. The filtered count matrices and the spatial locations were fed into PRESENT and

other baseline methods, including SpaGCN[31], STAGATE[32], PAST[19] and SpatialPCA[74]. All baseline

methods were implemented with default settings in their tutorials. PRESENT again employed a gene

selection procedure, obtaining 3,000 highly variable genes as input.

Multi-sample integration of spatial ATAC mouse embryo data. The three spatial ATAC mouse embryo

samples from three developmental stages have the unified peak sets and were directly concatenated.

Following the experimental settings in cross-modality representation of a single spatial sample, peaks

accessible in less than 0.5% spots were filtered out. We benchmarked PRESENT with six integration

methods developed for single-cell data (MNN[65], Scanorama[25], Harmony[24], SCALEX[66], scBasset-BC[67]

and SnapATAC2[64]) and two recently proposed SRT methods (STAligner[27] and GraphST[22]). Here

Stitch3D[33] was not used as it encountered error. MNN, Scanorama and Harmony performed batch

correction based on the latent embeddings obtained by EpiScanpy[98]. All the methods were implemented

according to their default settings except for STAligner. Since other methods utilized all the input peaks,

we ignored the feature selection step of STAligner to ensure better performance and fairness.

Multi-sample integration of 10x Visium mouse brain data. The intersected gene set of the two mouse

brain samples, including a sagittal anterior section and a sagittal posterior section, were utilized as the

unified gene set. Next, gene count matrices of the two samples were concatenated and genes with zero

expression in all spots were filtered out. Similarly, 3,000 highly variable genes were selected as the input

for PRESENT. PRESENT was compared with two recently developed SRT methods, i.e., STAligner[27] and

GraphST[22], which were conducted according to their default settings. Specifically, we utilized PASTE[23] to

accomplish the initial coordinate alignment step for GraphST according to the original study[22].

Evaluation metrics

Spatial clustering metrics. We implemented four commonly used metrics[50, 51] to quantitatively evaluate

the performance of spatial clustering, including adjusted rand index (ARI), adjusted mutual information

(AMI), normalized mutual information (NMI) and homogeneity (Homo).

ARI, ranging from 0 to 1, denotes the consistent adjustment of the expectations of rand index:

876
$$\text{ARI} = \frac{RI(P,T) - E[RI(P,T)]}{\max[RI(P,T)] - E[RI(P,T)]}$$

where $RI(\cdot)$ denotes the function of rand index, T refers to the ground truth labels, and P represents

the clustering labels. Rand index is the probability of the agreement between the clustering result and the

ground truth labels on a randomly selected spot set.

Mutual Information (MI) accesses the similarity between the clustering labels and ground truth

labels, from which both AMI and NMI originate. AMI ranges from -1 to 1, while NMI ranges from 0 to 1.

AMI is calculated as:

883
$$\text{AMI} = \frac{MI(P,T) - E[MI(P,T)]}{avg[H(P), H(T)] - E[MI(P,T)]}$$

884   where $MI(\cdot)$ refers to the mutual information, $E(\cdot)$ represents the expectation function and $H(\cdot)$

885   denotes the function of information entropy. NMI can be calculated as:

886
$$\text{NMI} = \frac{MI(P,T)}{\sqrt{H(P)H(T)}}$$

887   Homogeneity is another metric to measure the clustering performance. Specifically, if all the clusters

888   involve only data points that are members of a single class in the ground truth labels, the clustering results

889   could be interpreted as satisfying homogeneity, calculated as:

890
$$\text{Homo} = 1 - \frac{H(T|P)}{H(T)}$$

Integration metrics. According to a recent benchmarking framework for single-cell omics data

integration, two important perspectives need to be considered when evaluating the performance of multi-

sample integration, namely the batch effect removal and the biological variance conservation[68].

To quantitatively evaluate the performance of batch effect removal, we introduced several metrics

used in the benchmarking framework[68], including graph connectivity(GC), k-nearest-neighbor batch-

effect test (kBET) and batch average silhouette width (bASW). In addition, we also implemented a cross-

validation strategy to evaluate the cross-batch annotation performance based on latent representations

obtained by different integration methods, which is also an important assessment of batch correction[99].

Specifically, using the spots from a specific sample as the test set and those from other samples along

with the domain labels as the training set, we transferred the domain labels of the training set into the test

set based on a k-NN classifier and the joint embeddings of all the samples. k-NN classifier was used as

its performance mainly depended on the quality of joint embeddings obtained by different methods. The

performance of label transfer was evaluated with four commonly used metrics implemented in the scikit-

learn package, i.e., accuracy (Acc), kappa, macro F1 score (mF1) and the weighted F1 score (wF1). After

all the samples were used as the test set in a cross-validation manner, we calculated the average score for

each metric to evaluate the performance of cross-batch annotation.

The quantitative evaluation for biological variance conservation involves the assessment of

representative quality and the clustering performance. The representative quality was accessed via the

mean average precision (MAP)[100] and two metrics included in the benchmarking framework[68], namely

the spot average silhouette width (sASW) and spot local inverse Simpson's index (sLISI) , while the

evaluation of clustering performance was based on the four metrics, i.e., ATI, AMI, NMI and Homo.

We calculated category scores of the batch effect removal and the biological variance conservation

through averaging the metrics mentioned in the two aspects, respectively. We also calculated an overall

score using a 40/60 weighted mean of the two category scores following previous studies[68, 100].

Downstream analyses

Visualization. Based on the latent representations obtained by various methods, we utilized the uniform

manifold approximation and projection (UMAP)[52] algorithm to further reduce the dimension of latent

representation matrices into two-dimensional matrices for visualization. In addition, the two-dimensional

UMAP and spatial coordinate matrices were visualized using the functions *scanpy.pl.umap()* and

scanpy.pl.spatial() wrapped in the SCANPY package[97].

Spatial clustering. We implemented a binary search strategy to tune the resolution of Leiden algorithm[49],

a graph-based clustering method, to match the number of spatial clusters as closely as possible with the

number of ground truth domains. Specifically, we set a minimum resolution res_{min} and a maximum

resolution res_{max}, and obtain clustering results using a resolution equal to $\frac{res_{min}+res_{max}}{2}$. If the number

of clusters exceeds the number of ground truth domains, res_{max} is updated to $\frac{res_{min}+res_{max}}{2}$. Otherwise,

res_{min} is updated to $\frac{res_{min}+res_{max}}{2}$. The iterative process continues until the number of clusters matches

the number of ground truth domains or the current iteration reaches the maximum number of iterations.

Differentially expressed gene and protein analysis. We implemented differential analysis based on the

Wilcoxon rank-sum test[54] wrapped in *scanpy.tl.rank_genes_groups()* function of the SCANPY package[97].

Gene ontology analysis. Given a specific gene set, we conducted gene ontology (GO) analysis using the

clusterProfiler package[101], which automated the process of biological-term classification and the

enrichment analysis of gene clusters.

Data availability

The single-nucleus ATAC-seq and spatial ATAC mouse embryo datasets[9] are available at Gene

Expression Omnibus (GEO) using the accession code GSE214991. The Stereo-seq axolotl brain datasets[6]

as well as the reference dataset, i.e., axolotl brain data at 20 days post-injury, can be obtained through

their database https://db.cngb.org/stomics/artista/. The 10x Visium mouse brain datasets, including a

sagittal anterior section and a sagittal posterior section, are accessible at the 10x Genomics websites

https://www.10xgenomics.com/datasets/mouse-brain-serial-section-2-sagittal-anterior-1-standard and

https://www.10xgenomics.com/datasets/mouse-brain-serial-section-2-sagittal-posterior-1-standard,

respectively. The SPOTS mouse spleen data[11] is accessible at GEO with accession code GSE198353. The

10x Visium RNA & protein human lymph node dataset[35] can be downloaded from the Zenodo database

https://zenodo.org/records/10362607. The MISAR-seq mouse brain data[12] is available from the National

Genomics Data Center via accession number OEP003285.

Code availability

The code as well as the detailed tutorials for implementing PRESENT can be found at the Github website https://github.com/lizhen18THU/PRESENT.

Acknowledgements

This work was partly supported by the National Key Research and Development Program of China (grant nos. 2022YFF1202400 and 2023YFF1204802), the National Natural Science Foundation of China (grant nos. 62273194), and Beijing Natural Science Foundation (L242026).

Author contributions

R.J. and L.Z. conceived the study and supervised the project. Z.L. designed, implemented and validated the PRESENT framework. X.J.C., X.Y.C., Z.G., Y.L., Y.P., S.C. and H.L. helped with analyzing the results. Z.L. and R.J. wrote the manuscript, with input from all the authors.

Competing interests

The authors declare no competing interests.

References

1. Baysoy, A. et al. The technological landscape and applications of single-cell multi-omics. *Nat. Rev. Mol. Cell Biol.* **24**, 695-713 (2023).

2. Adema, K. et al. Lost in space: what single-cell RNA sequencing cannot tell you. *Trends Plant Sci.* (2024).

3. Armingol, E. et al. Deciphering cell-cell interactions and communication from gene expression. *Nat. Rev. Genet.* **22**, 71-88 (2021).

4. Cang, Z.X. et al. Screening cell-cell communication in spatial transcriptomics via collective optimal transport. *Nat. Methods* **20**, 218-228 (2023).

5. Shao, X. et al. Knowledge-graph-based cell-cell communication inference for spatially resolved transcriptomic data with SpaTalk. *Nat. Commun.* **13**, 4429 (2022).

6. Wei, X. et al. Single-cell Stereo-seq reveals induced progenitor cells involved in axolotl brain regeneration. *Science* **377**, eabp9444 (2022).

7. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat. Methods* **19**, 534-546 (2022).

8. Deng, Y.X. et al. Spatial profiling of chromatin accessibility in mouse and human tissues. *Nature* **609**, 375-383 (2022).

9. Llorens-Bobadilla, E. et al. Solid-phase capture and profiling of open chromatin by spatial ATAC. *Nat. Biotechnol.* **41**, 1085-1088 (2023).

10. Liu, Y. et al. High-plex protein and whole transcriptome co-mapping at cellular resolution with spatial CITE-seq. *Nat. Biotechnol.* **41**, 1405-1409 (2023).

11. Ben-Chetrit, N. et al. Integration of whole transcriptome spatial profiling with protein markers. *Nat. Biotechnol.* **41**, 788-193 (2023).

12. Jiang, F.Q. et al. Simultaneous profiling of spatial gene expression and chromatin accessibility during mouse brain development. *Nat. Methods* **20**, 1048-1057 (2023).

13. Yuan, Z.Y. et al. SODB facilitates comprehensive exploration of spatial omics data. *Nat. Methods* **20**, 387-399 (2023).

14. Xu, Z. et al. STOmicsDB: a comprehensive database for spatial transcriptomics data sharing, analysis and visualization. *Nucleic Acids Res.* **52**, D1053-D1061 (2024).

15. Berger, S.L. The complex language of chromatin regulation during transcription. *Nature* **447**, 407-412 (2007).

16. MacNeil, L.T. & Walhout, A.J.M. Gene regulatory networks and the role of robustness and stochasticity in the control of gene expression. *Genome Res.* **21**, 645-657 (2011).

17. Stoeckius, M. et al. Simultaneous epitope and transcriptome measurement in single cells. *Nat. Methods* **14**, 865-868 (2017).

18. Peterson, V.M. et al. Multiplexed quantification of proteins and transcripts in single cells. *Nat. Biotechnol.* **35**, 936-939 (2017).

19. Li, Z. et al. Latent feature extraction with a prior-based self-attention framework for spatial transcriptomics. *Genome Res.* **33**, 1757-1773 (2023).

20. Shang, L. & Xiang, Z. Spatially aware dimension reduction for spatial transcriptomics. *Nat. Commun.* **13**, 7203 (2022).

21. Li, G.Y. et al. A deep generative model for multi-view profiling of single-cell RNA-seq and ATAC-seq data. *Genome Biol.* **23**, 20 (2022).

22. Long, Y. et al. Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST. *Nat. Commun.* **14**, 1155 (2023).

23. Zeira, R. et al. Alignment and integration of spatial transcriptomics data. *Nat. Methods* **19**, 567-575 (2022).

24. Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289-1296 (2019).

25. Hie, B., Bryson, B. & Berger, B. Efficient integration of heterogeneous single-cell transcriptomes using
Scanorama. *Nat. Biotechnol.* **37**, 685-691 (2019).

26. Guo, T.T. et al. SPIRAL: integrating and aligning spatially resolved transcriptomics data across different
experiments, conditions, and technologies. *Genome Biol.* **24**, 241 (2023).

27. Zhou, X., Dong, K.N. & Zhang, S.H. Integrating spatial transcriptomics data across different conditions,
technologies and developmental stages. *Nat. Comput. Sci.* **3**, 894-906 (2023).

28. Chen, A. et al. Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned
arrays. *Cell* **185**, 1777-+ (2022).

29. Chen, S. et al. Spatially resolved transcriptomics reveals genes associated with the vulnerability of middle
temporal gyrus in Alzheimer's disease. *Acta Neuropathol. Commun.* **10**, 188 (2022).

30. Yuan, Z. et al. Benchmarking spatial clustering methods with spatially resolved transcriptomics data. *Nat.*
Methods **21**, 1-11 (2024).

31. Hu, J. et al. SpaGCN: Integrating gene expression, spatial location and histology to identify spatial domains
and spatially variable genes by graph convolutional network. *Nat. Methods* **18**, 1342-1351 (2021).

32. Dong, K.N. & Zhang, S.H. Deciphering spatial domains from spatially resolved transcriptomics with an
adaptive graph attention auto-encoder. *Nat. Commun.* **13**, 1739 (2022).

33. Wang, G. et al. Construction of a 3D whole organism spatial atlas by joint modelling of multiple slices with
deep neural networks. *Nat. Mach. Intell.* **5**, 1200-1213 (2023).

34. Vandereyken, K. et al. Methods and applications for single-cell and spatial multi-omics. *Nat. Rev. Genet.* **24**,
494-515 (2023).

35. Long, Y. et al. Deciphering spatial domains from spatial multi-omics with SpatialGlue. *Nat. Methods* (2024).

36. Tian, T. et al. Dependency-aware deep generative models for multitasking analysis of spatial omics data. *Nat.*
Methods (2024).

37. Veličković, P. et al. Graph attention networks. Preprint at *arXiv* https://arxiv.org/abs/1710.10903 (2017).

38. Blundell, C. et al. Weight uncertainty in neural network. In *32nd International Conference on Machine*
Learning (2015).

39. Eraslan, G. et al. Single-cell RNA-seq denoising using a deep count autoencoder. *Nat. Commun.* **10**, 390
(2019).

40. Martens, L.D. et al. Modeling fragment counts improves single-cell ATAC-seq analysis. *Nat. Methods* **21**,
28-31 (2023).

41. Lin, Z. et al. Multi-modal Contrastive Representation Learning for Entity Alignment. In *29th International*
Conference on Computational Linguistics (2022).

42. Chen, T. et al. A simple framework for contrastive learning of visual representations. In *37th International*
Conference on Machine Learning (2020).

43. Kopp, W., Akalin, A. & Ohler, U. Simultaneous dimensionality reduction and integration for single-cell
ATAC-seq data using deep learning. *Nat. Mach. Intell.* **4**, 162-168 (2022).

44. Swanton, C. Intratumor heterogeneity: evolution through space and time. *Cancer Res.* **72**, 4875-4882 (2012).

45. Janiszewska, M. The microcosmos of intratumor heterogeneity: the space-time of cancer evolution.
Oncogene **39**, 2031-2039 (2020).

46. Gayoso, A. et al. Joint probabilistic modeling of single-cell multi-omic data with totalVI. *Nat. Methods* **18**,
272-282 (2021).

47. Kim, H.J. et al. CiteFuse enables multi-modal analysis of CITE-seq data. *Bioinformatics* **36**, 4137-4143
(2020).

48. Hao, Y.H. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 (2021).

49. Traag, V.A., Waltman, L. & Van Eck, N.J. From Louvain to Leiden: guaranteeing well-connected
communities. *Sci. Rep.* **9**, 5233 (2019).

50. Chen, S. et al. VPAC: Variational projection for accurate clustering of single-cell transcriptomic data. *BMC*
Bioinformatics **20**, 139-151 (2019).

51. Jiang, R. et al. SINFONIA: scalable identification of spatially variable genes for deciphering spatial domains.
Cells **12**, 604 (2023).

52. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension
reduction. Preprint at *arXiv* https://arxiv.org/abs/1802.03426 (2018).

53. Lewis, S.M., Williams, A. & Eisenbarth, S.C. Structure and function of the immune system in the spleen. *Sci.*
Immunol. **4**, eaau6085 (2019).

54. Soneson, C. & Robinson, M.D. Bias, robustness and scalability in single-cell differential expression analysis.
Nat. Methods **15**, 255-261 (2018).

55. Schmid, M.C. & Varner, J.A. Myeloid cells in tumor inflammation. *Vasc. Cell* **4**, 1-7 (2012).

56. Borges, M.D. & Sesti-Costa, R. Macrophages: key players in erythrocyte turnover. *Hematol. Transfus. Cell*
Ther. **44**, 574-581 (2022).

57. Seyfried, A.N., Maloney, J.M. & MacNamara, K.C. Macrophages orchestrate hematopoietic programs and
regulate HSC function during inflammatory stress. *Front. Immunol.* **11**, 548463 (2020).

58. Cui, K. et al. Endocytic adaptors in cardiovascular disease. *Front. Cell Dev. Biol.* **8**, 624159 (2020).

59. Baranov, M.V. et al. Modulation of immune responses by particle size and shape. *Front. Immunol.* **11**, 607945
(2021).

60. Kurup, S.P., Butler, N.S. & Harty, J.T. T cell-mediated immunity to malaria. *Nat. Rev. Immunol.* **19**, 457-471
(2019).

61. Argelaguet, R. et al. MOFA+ : a statistical framework for comprehensive integration of multi-modal single-
cell data. *Genome Biol.* **21**, 1-17 (2020).

62. Jin, S.Q., Zhang, L.H. & Nie, Q. scAI: an unsupervised approach for the integrative analysis of parallel
single-cell transcriptomic and epigenomic profiles. *Genome Biol.* **21**, 1-19 (2020).

63. Lin, X. et al. Clustering of single-cell multi-omics data with a multimodal deep learning method. *Nat.*
Commun. **13**, 7705 (2022).

64. Zhang, K. et al. A fast, scalable and versatile tool for analysis of single-cell omics data. *Nat. Methods* **21**,
217-227 (2024).

65. Haghverdi, L. et al. Batch effects in single-cell RNA-sequencing data are corrected by matching mutual
nearest neighbors. *Nat. Biotechnol.* **36**, 421-427 (2018).

66. Xiong, L. et al. Online single-cell data integration through projecting heterogeneous datasets into a common
cell-embedding space. *Nat. Commun.* **13**, 6118 (2022).

67. Yuan, H. & Kelley, D.R. scBasset: sequence-based modeling of single-cell ATAC-seq using convolutional
neural networks. *Nat. Methods* **19**, 1088-1096 (2022).

68. Luecken, M.D. et al. Benchmarking atlas-level data integration in single-cell genomics. *Nat. Methods* **19**,
41-50 (2022).

69. Heumos, L. et al. Best practices for single-cell analysis across modalities. *Nat. Rev. Genet.* **24**, 550-572
(2023).

70. Zhang, H. et al. Extensive evaluation of ATAC-seq protocols for native or formaldehyde-fixed nuclei. *BMC*
Genomics **23**, 214 (2022).

71. Xiong, L. et al. SCALE method for single-cell ATAC-seq analysis via latent feature extraction. *Nat. Commun.*
10, 4576 (2019).

72. González-Blas, C.B. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-seq data. *Nat.*
Methods **16**, 397-400 (2019).

73. Pierson, E. & Yau, C. ZIFA: Dimensionality reduction for zero-inflated single-cell gene expression analysis.
Genome Biol. **16**, 1-10 (2015).

74.   Shang, L. & Zhou, X. Spatially aware dimension reduction for spatial transcriptomics. *Nat. Commun.* **13**, 7203 (2022).

75.   Sunkin, S.M. et al. Allen Brain Atlas: an integrated spatio-temporal portal for exploring the central nervous system. *Nucleic Acids Res.* **41**, D996-D1008 (2012).

76.   Teicher, M.H., Anderson, C.M. & Polcari, A. Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proc. Natl. Acad. Sci.* **109**, E563-E572 (2012).

77.   Matsuda, K. et al. Transsynaptic modulation of kainate receptor functions by C1q-like proteins. *Neuron* **90**, 752-767 (2016).

78.   Park, S.-Y. et al. Hippocalcin promotes neuronal differentiation and inhibits astrocytic differentiation in neural stem cells. *Stem Cell Rep.* **8**, 95-111 (2017).

79.   Coffman, K.A., Dum, R.P. & Strick, P.L. Cerebellar vermis is a target of projections from the motor areas in the cerebral cortex. *Proc. Natl. Acad. Sci.* **108**, 16068-16073 (2011).

80.   Hachem, S. et al. Expression of S100B during embryonic development of the mouse cerebellum. *BMC Dev. Biol.* **7**, 1-15 (2007).

81.   Hirasawa, M. et al. Carbonic anhydrase related protein 8 mutation results in aberrant synaptic morphology and excitatory synaptic function in the cerebellum. *Mol. Cell. Neurosci.* **35**, 161-170 (2007).

82.   Lassuthova, P. et al. Mutations in ATP1A1 cause dominant Charcot-Marie-Tooth type 2. *Am. J. Hum. Genet.* **102**, 505-514 (2018).

83.   Pacey, L.K. et al. Delayed myelination in a mouse model of fragile X syndrome. *Hum. Mol. Genet.* **22**, 3920-3930 (2013).

84.   Argelaguet, R. et al. Computational principles and challenges in single-cell data integration. *Nat. Biotechnol.* **39**, 1202-1215 (2021).

85.   Chen, S.Q. et al. RA3 is a reference-guided approach for epigenetic characterization of single cells. *Nat. Commun.* **12**, 13 (2021).

86.   Chen, X. et al. Cell type annotation of single-cell chromatin accessibility data via supervised Bayesian embedding. *Nat. Mach. Intell.* **4**, 116-126 (2022).

87.   Cusanovich, D.A. et al. Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**, 910-914 (2015).

88.   Cusanovich, D.A. et al. A single-cell atlas of in vivo mammalian chromatin accessibility. *Cell* **174**, 1309-1324 (2018).

89.   Mulè, M.P., Martins, A.J. & Tsang, J.S. Normalizing and denoising protein expression data from droplet-based single cell profiling. *Nat. Commun.* **13**, 2099 (2022).

90.   Stoeckius, M. et al. Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. *Genome Biol.* **19**, 1-12 (2018).

91.   Stuart, T. et al. Single-cell chromatin state analysis with Signac. *Nat. Methods* **18**, 1333-1341 (2021).

92.   Qiu, P. Embracing the dropouts in single-cell RNA-seq analysis. *Nat. Commun.* **11**, 1169 (2020).

93.   Ganin, Y. et al. Domain-adversarial training of neural networks. *J. Mach. Learn. Res.* **17**, 1-35 (2016).

94.   Zhang, D. et al. Spatial epigenome-transcriptome co-profiling of mammalian tissues. *Nature* **616**, 113-122 (2023).

95.   Chiang, W.-L. et al. Cluster-GCN: An efficient algorithm for training deep and large graph convolutional networks. In *25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining* (2019).

96.   Bai, J., Ren, Y. & Zhang, J. Ripple walk training: A subgraph-based training framework for large and deep graph neural network. In *2021 International Joint Conference on Neural Networks (IJCNN)* (2021).

97.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol.* **19**, 1-5 (2018).

98.   Danese, A. et al. EpiScanpy: integrated single-cell epigenomic analysis. *Nat. Commun.* **12**, 5228 (2021).

99.   Song, Y. et al. Benchmarking strategies for cross-species integration of single-cell RNA sequencing data. *Nat.*
1145   *Commun.* **14**, 6495 (2023).
1146   100.   Cao, Z.-J. & Gao, G. Multi-omics single-cell data integration and regulatory inference with graph-linked
1147   embedding. *Nat. Biotechnol.* **40**, 1458-1466 (2022).
1148   101.   Yu, G. et al. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS: J.*
1149   *Integrative Biol.* **16**, 284-287 (2012).
1150
1151

Figures

Fig. 1 | The framework of PRESENT. a, Cross-modality representation of spatial multi-omics data. PRESENT is capable to integrate spatial coordinate data and multi-omics data, including proteomic data sequenced by antibody-derived tags (ADT), transcriptomic data by RNA sequencing (RNA), and epigenomic data generated by the assay for transposase-accessible chromatin with sequencing (ATAC). Firstly, an omics-specific encoder is used to extract specific biological variations contained in each omics layer. Each omics-specific encoder is composed of a graph attention neural network

(GAT) and a Bayesian neural network (BNN) module. All the omics-specific feature matrices extracted by encoders are
fused by a cross-omics alignment module, which integrates multi-omics information via the contrastive learning inter-
omics alignment loss function. The output of cross-omics alignment module refers to the final representation matrix of
PRESENT. The distinct patterns of different omics layers are addressed via omics-specific decoders, which take the output
of cross-omics alignment module as input and model the count matrices of RNA and ATAC data using the zero-inflation
negative binomial (ZINB) and the zero-inflation Poisson (ZIP) distribution, respectively. **b,** Multi-sample integration of
spatial omics data. PRESENT adopts a two-stage workflow for the integration of multiple spatial samples. In the first
pretraining stage, PRESENT is pretrained using omics data and the cross-sample graph obtained from all the samples. The
intra-sample edges of the cross-sample graph are built using spatial coordinates, while inter-sample connections are
established based on the similarity of omics features across samples. In the second integration stage, PRESENT implements
a cyclic iterative strategy to refine the cross-sample graph using the joint embeddings obtained by the trained model.
PRESENT further introduces a contrastive learning-based inter-batch alignment loss, a batch-adversarial learning strategy
and an intra-batch preserving loss in the second stage to eliminate batch effects while preserving biological signals. **c,**
Applications of PRESENT in spatial omics data. PRESENT can be easily adapted from spatial multi-omics to spatial
single-omics data by simplifying the multi-view autoencoder into a single-view autoencoder. The joint embeddings
obtained by PRESENT can be applied to various downstream analyses, including spatial visualization, domain
identification and the integrative analysis of samples across conditions and developmental stages.
1176

Fig. 2 | PRESENT facilitates accurate spatial domain identification in spatial RNA-ADT data. a, The spatial visualization of the 10x Genomics Visium RNA-Protein human lymph node sample colored by ground truth domain labels. **b,** The quantitative comparison of spatial domain identification performance between PRESENT and other baseline methods based on bar plot and, **c,** the comparison between PRESENT utilizing both RNA and ADT data (RNA & ADT) and PRESENT using only RNA (RNA-only) or ADT data (ADT-only) based on radar plot on the human lymph node dataset. **d,** The UMAP visualization of latent embeddings obtained by different methods colored by ground truth domain labels and **e,** cluster labels identified based on different methods on the human lymph node dataset. **f,** The spatial visualization of clusters identified based on different methods on the human lymph node dataset. **g,** The histology image and spatial visualization of spatial clusters identified based on PRESENT on the SPOTS mouse spleen dataset. **h,** The dot plot demonstrating the difference of ADT expression levels and **i,** RNA expression levels between each domain and the rest domains. **j,** The volcano plot illustrating the difference of RNA expression levels between Mac1 and Mac2-enriched domains. The vertical dashed line represents the threshold for $\log(\text{FC}) = \pm 0.2$, while the horizontal dashed line denotes the threshold for $-\log(\text{FDR} = 0.05)$. **k,** The chordal plots of the Mac1-enriched domain (left) and the Mac2-enriched domain (right) demonstrating the linkage of differential expressed genes (DEGs) in each domain and the corresponding enriched biological processes. In each chordal plot, the left semicircle represents DEGs while the right semicircle denotes the enriched biological processes.

Fig. 3 | PRESENT integrates spatially aware multi-omics biological signals in spatial RNA-ATAC data. a, The quantitative comparison between PRESENT and other baseline methods and **b,** the comparison between PRESENT

utilizing both RNA and ATAC data (RNA & ATAC) and PRESENT using only RNA (RNA-only) or ATAC (ATAC-only) data based on box plot on the MISAR-seq mouse brain dataset. The center line, box limits and whiskers in the box plots are the median, upper and lower quartiles, and 1.5× interquartile range, respectively. **c,** The spatial visualization of spots from the E18.5 mouse brain sample colored by ground truth domain labels. **d,** The heatmap visualization of latent representations obtained by PRESENT. **e,** The UMAP visualization of latent representations obtained by different methods colored by ground truth domain labels. **f,** The UMAP visualization of latent representations obtained by different methods and **g,** the spatial visualization colored by spatial clusters identified based on different methods on the E18.5 mouse brain sample. The UMAP visualization of latent embeddings obtained by PRESENT using only RNA data, only ATAC data and both RNA & ATAC data, colored by ground truth domain labels on the **h,** E11.0 and **i,** E15.5 mouse brain samples.

Fig. 4 | PRESENT eliminates batch effects while preserving biological signals in spatial multi-omics data. The UMAP visualization of spots across the four MISAR-seq mouse brain samples based on raw RNA or ATAC data colored by **a,** sample indices and **b,** ground truth domain labels. Note that prior to applying the UMAP algorithm to raw RNA or ATAC data, we first reduce the dimensionality to 50 using PCA transformation. **c,** The UMAP visualization of spots across the four MISAR-seq mouse brain samples based on the joint embeddings produced by different integration methods colored

by **c,** sample indices and **d,** ground truth domain labels. **e,** The quantitative evaluation of different integration methods on the four MISAR-seq mouse brain samples using 14 metrics divided into two categories, namely batch effect removal and biological variance conservation. The category scores of these two aspects were calculated by averaging the metrics within each category. An overall score for each integration method was computed using a 40/60 weighted mean of the category scores for batch effect removal and biological variance conservation.

Fig. 5 | PRESENT facilitates effective cross-modal representation in spatial single-omics data. a, The ridge plot and violin plot of log10 total fragment counts, namely log10(nFrags), across the six spatial ATAC mouse embryo samples. **b,** The ridge plot of TSS enrichment scores for spots and the line plot of the enrichment profiles of fragments around the TSS across the six mouse embryo samples. **c,** The quantitative evaluation of spatial clustering performance of different methods

using four metrics, including ARI, AMI, NMI and Homo. The center line, box limits and whiskers in the box and violin plots are the median, upper and lower quartiles, and 1.5× interquartile range, respectively. **d,** The spatial visualization colored by ground truth domain labels, **e,** the UMAP visualization of the embeddings obtained by different methods colored by ground truth domain labels and **f,** the spatial visualization of clusters identified using the embeddings obtained by different methods on the E15.5-S1 mouse embryo sample. **g,** The detailed comparison of performance of PRESENT and PRESENT-SN & SP on the six mouse embryo samples based on radar plots. **h,** The UMAP visualization of the latent representations obtained by different methods colored by ground truth domain labels on E12.5-S2 mouse embryo sample. **i,** The spatial clustering performance of baseline methods, PRESENT and PRESENT-SN & SP on the six mouse embryo samples under different levels of dropout rate. Here SCALE encountered error when the dropout rate was larger than 0.2.

Fig. 6 | PRESENT enables vertical and horizontal integration of spatial single-omics samples. a, The quantitative evaluation of different integration methods on the three spatial ATAC mouse embryo samples using 14 metrics divided into two categories, namely batch effect removal and biological variance conservation. The category scores of these two aspects were calculated by averaging the metrics within each category. An overall score for each integration method was computed using a 40/60 weighted mean of the category scores for batch effect removal and biological variance conservation (Methods). **b,** The spatial visualization of spots across the three spatial ATAC mouse embryo samples colored by ground truth spatial domains and **c,** the spatial clusters identified based on different integration methods. The first, second and third row of **b** and **c** denotes the samples from E12.5, E13.5 and E15.5 stages, respectively. **d,** The anatomic

1249   annotation of the sagittal region in P56 mouse brain provided by Allen Reference Atlas. The joint spatial clustering results

1250   based on the latent embeddings obtained by **e,** STAligner, **f,** GraphST and **g,** PRESENT on the two horizontal mouse brain

1251   sagittal samples generated by the 10x Genomics Visium platform.

1252