Chromosome-level reference genome assembly for the mountain hare (*Lepus timidus*)

Zsófia Fekete[1,2], Dominic E. Absolon[3], Craig Michell[4], Jonathan M. D. Wood[3], Steffi Goffart[1] and Jaakko L. O. Pohjoismäki*[1]

[1]University of Eastern Finland, Department of Environmental and Biological Sciences, Joensuu, Finland
[2]Hungarian University of Agriculture and Life Sciences, Institute of Genetics and Biotechnology, Gödöllő, Hungary
[3]Tree of Life, Wellcome Sanger Institute, Hinxton, Cambridge, UK.
[4]King Abdullah University of Science and Technology, Red Sea Research Center, Biological and Environmental Science and Engineering Division, Thuwal, Kingdom of Saudi Arabia

*Corresponding author
Correspondence: jaakko.pohjoismaki@uef.fi

Abstract

We present here a high-quality genome assembly of a male mountain hare (*Lepus timidus* Linnaeus), from Ilomantsi, Eastern Finland, utilizing an isolated fibroblast cell line as the source for fresh DNA and RNA. Following the previously published brown hare reference genome assembly, the mountain hare is the second Finnish pilot species for the European Reference Genome Atlas (ERGA) initiative, a collaborative effort to generate reference genomes for European biodiversity.

The genome was assembled using 21× PacBio HiFi sequencing data and scaffolded using the Hi-C chromosome structure capture approach. After manual curation, the primary assembly length was 2,695,305,354 bp with N50 125,755,317 bp. The largest scaffold was 181 Mbp and the scaffold N50 127 Mbp, contributing to a primary assembly consisting of 85 scaffolds and an alternate assembly with 109 scaffolds. The scaffolds include 23 autosomes, numbered according to their size, as well as X and Y chromosome, matching the known karyotype. Telomeric regions were present on at least one end of 19 of the chromosomes. The genome has a high degree of completeness based on the BUSCO score (mammalia_odb10 database), Complete: 95.1 % [Single copy: 92.3 %, Duplicated: 2.7 %], Fragmented 0.8 %, and Missing 4.1 %. The mitochondrial genome of the cell line was sequenced and assembled separately.

Compared to the previous pseudo-reference genome assembly of *L. timidus* ssp. *hibernicus* Bell, assembled using the rabbit genome, this new reference genome represents the nominal subspecies and the species-specific chromosomal conformation. The published genome assembly can help to understand the genomic basis of adaptations to snowy and cold environments and opens possibilities for experimental analysis of mountain hare genes, as well as provides a solid basis for future genomic work on Lagomorpha.

Keywords: Lagomorpha, reference genome, chromosome structure, phylogeny, fibroblast cell line

Introduction

The mountain hare (*Lepus timidus* L.), including several subspecies, is the most widespread hare
species in the world, with its distribution ranging from Ireland to Japan (Angerbjörn & Schai-Braun,
2022). The species is adapted to the cold and snowy conditions of northern Eurasia by having wide
snowshoe feet and white winter pelage. The ongoing climate change-induced shortening of the snow-
covered season is detrimental for the mountain hare, especially due to camouflage mismatch (Zimova
et al., 2016; Zimova et al., 2020), resulting in range contraction to the North or higher altitudes in
mountains, while benefitting the expansion of its more southern relative, the brown hare (*L. europaeus*)
(C.-G. Thulin, 2003; Jansson & Pehrson, 2007; Reid, 2011; Levänen, Kunnasranta, et al., 2018; Rehnus et
al., 2018).

The mountain hare has shown resilience during the past warm periods of the Holocene (Smith et al.,
2017), diverging also to more temperate climate-adapted subspecies (Giska et al., 2022). Notable
examples include the Irish hare (*L. t. hibernicus* Bell) and the heath hare (*L. t. sylvaticus* Nilsson), which
are likely relatively old evolutionary lineages that are genetically distinct from the widespread nominal
subspecies (Giska et al., 2022; Michell et al., 2022). In addition, post-glacial contact with other congeneric
hare species has resulted in recurrent hybridization and genetic introgression (Melo-Ferreira et al., 2009;
Levänen, Thulin, et al., 2018), which appears to have adaptive significance (Giska et al., 2019;
Pohjoismäki et al., 2021; Giska et al., 2022).

Besides the fact that geographically restricted, relic-like subspecies are threatened (Caravaggi et al.,
2017; C. G. Thulin et al., 2021) and thus interesting in the context of conservation genetics, hares provide
ample opportunities to study local adaptations (Giska et al., 2022; Michell et al., 2022), genomic
introgression (Levänen, Thulin, et al., 2018; Giska et al., 2019; Pohjoismäki et al., 2021) and genetic
constituents of species boundaries (Gaertner et al., 2022). Such studies are greatly facilitated by the
availability of reference-level genome assemblies (Michell et al., 2024).

Relatively recent advances in sequencing technologies, so-called third generation sequencing or
high-throughput sequencing of long molecules, has provided affordable means to produce highly
continuous genome assemblies. Combined with techniques for sequence contact maps for
chromosomal scaffolding, these methods enable reference-level genome assemblies for almost any
species (Lawniczak et al., 2022), a level previously achieved only by a few model organisms. This in turn
has resulted in emergence of several sequencing initiatives to produce reference genome assemblies for
various taxa, including the European Reference Genome Atlas (ERGA) and the Darwin Tree of Life (DToL)
(Blaxter et al., 2022). In the presented work, we use these methods to provide a reference-level sequence
assembly for the mountain hare. Together with our previously released brown hare genome
assembly(Michell et al., 2024), the mountain hare genome represents another Finnish contribution to
the ERGA pilot project, aimed to demonstrate the feasibility of a continent-wide collaboration to
sequence and assemble reference genomes for local species. Through resource and knowledge sharing,
this collaborative approach not only accelerates the creation of reference-grade genomes but also
enhances biological discovery, benefiting the broader scientific community (Mc Cartney et al., 2024).

Since the success of long-molecule sequencing depends on intact, high molecular weight DNA,
requiring fresh or flash-frozen tissue samples, we used a living fibroblast cell line of a male mountain
hare specimen was collected from Ilomantsi, Eastern Finland (Gaertner et al., 2022) as the source. The
sex of the specimen was important to cover the Y-chromosome. The sampling locality was chosen
outside of the brown hare range to minimize the probability for hybridization (Levänen, Kunnasranta, et
al., 2018; Levänen, Thulin, et al., 2018). Furthermore, the sampled population represents a continuum
across the vast eastern Taiga zone, making our sample very representative geographically for the
nominal mountain hare subspecies. It is worth noting that Linné did not formally designate a type
specimen for the mountain hare, and the type locality has been later decided to be Uppsala in Sweden
(Angerbjörn & Schai-Braun, 2022). The Swedish hares belong to the same larger, geographically

connected Fennoscandian population as our specimen (Levänen, Thulin, et al., 2018; Michell et al., 2022).
This is important because sampling from or near the type locality is considered desirable for the
representativeness of the reference genome assembly (Lawniczak et al., 2022).

Following the same approach as with our brown hare reference genome assembly (Michell et al.,
2024), we assembled a highly complete genome of the nominal mountain hare subspecies (*Lepus timidus*
timidus) using 21× genome coverage of PacBio HiFi read data. The 2.69 Gbp assembly was further
scaffolded with Hi-C sequencing data to chromosome-level, including all of the expected 23 autosomes,
and X and Y sex chromosomes (2n = 48) (Gustavsson, 1972). Telomeric regions were present on at least
one end in 19 of the assembled chromosomes, further demonstrating the high continuity of the
assembly.

While there is an existing genome assembly for the mountain hare (GCA_009760805), it represents
the Irish subspecies *L. t. hibernicus* (from here on the "Irish hare") and is based on a female specimen,
thus lacking the Y-chromosome. Besides having much lower continuity than the assembly presented
here, the Irish hare genome assembly is a "pseudoreference", scaffolded using the rabbit (*Oryctolagus*
cuniculus L.) reference genome assembly (Marques et al., 2020), and thus retaining the rabbit 2n = 44
karyotype (Korstanje et al., 1999). Besides representing a significant improvement over the Irish hare
genome assembly, providing a robust reference for future studies on adaptation, hybridization, and
conservation of the mountain hare, this assembly can offer insights into the evolutionary history of
hares, genetic basis of physiological adaptations as well as expand the molecular toolkit for
experimental investigation of genes and their functions.

Methods

Sampling and confirming of the species identity

A young male mountain hare was hunted in October 2018 in Kelovaara, Ilomantsi, Finland (63.0609
N, 30.6248 E). The sampling had minimal impact on the local mountain hare population and no impact
on the habitat. As mountain hares are legal game animals in Finland and the hunting followed the
regional hunting seasons and legislation (Metsästyslaki [Hunting law] 1993/615/5§), the sampling
adheres to the ARRIVE guidelines and no ethical assessment for the sampling was required. Sampling
also did not involve International Trade in Endangered Species of Wild Fauna and Flora (CITES) or other
export of specimens, as defined by the Convention on Biological Diversity (CBD).

The species identity was confirmed at site based on the morphological features distinguishing the
mountain hare from the brown hare, the only other hare species in northern Europe. The collection
location is remote wilderness, dominated by boreal Taiga Forest (Figure 1A) and devoid of brown hares,
which in Finland are dependent on man-made environments (Levänen, Kunnasranta, et al., 2018).
Further analysis of the coding part of the genome and mitochondrial DNA haplotyping showed no
ancestral admixture with brown hares (Gaertner et al., 2022).

Generation and vouchering of the cell line

The fibroblast cell line (LT1) was isolated from the specimen as described earlier (Gaertner et al.,
2022) and is deposited under voucher number ZFMK-TIS-69753 into the biobank of the Stiftung Leibniz-
Institut zur Analyse des Biodiversitätswandels, Zoological Research Museum Alexander König (ZFMK),
Bonn, Germany. The cell line has been immortalized using SV40 large T-antigen transformation. The cells
used for the DNA extraction represent passage number 10, as counted from the isolation of the primary
cells from the tissue biopsy. Cell lines with passage number under 20 are considered to have a low
passage number (ATCC, 2024).

High molecular weight DNA extraction and PacBio HiFi sequencing

High molecular weight DNA was purified using the classical phenol:chloroform:IAA extraction method (Sambrook & Russell, 2006). Briefly, the cells were grown to confluency on a 10 cm cell culture dish and detached using trypsin, followed by centrifugation and two washes with PBS. The cell pellet was resuspended in 2 ml of DNA extraction buffer (25 mM EDTA pH 8.0, 75 mM NaCl, 1 SDS and RNAseA), followed by Proteinase K digestion and phenol extraction, as detailed in the protocol (Sambrook & Russell, 2006). As previously (Michell et al., 2024), the DNA was sequenced on the PacBio Sequel II by DNA Sequencing and Genomics Laboratory, Institute of Biotechnology, University of Helsinki. The DNA was quantified using Qubit and the fragment length was assessed with the Fragment Analyzer (HS large fragment kit) and then sheared to a length of 24 kb using the Megaruptor 3 (Diagenode) (45/200/30). Following this, the buffer was replaced with PacBio's Elution buffer using AMPure beads. The DNA underwent repair and A-tailing, adapter ligation, nuclease treatment, and cleanup with SMRTbell cleanup beads, utilizing the SMRTbell prep kit 3.0. Fragments larger than 10,000 bp were purified with the BluePippin (Sage Science) (0.75% DF Marker S1 high pass 6-10kb v3), and the DNA was further purified using PacBio AMPure beads. It was then treated with DNA damage repair mix (PacBio) at 37°C for 60 minutes, followed by another purification with AMPure cleanup beads and elution into 11 µl of PacBio's Elution buffer. Libraries were sequenced on two SMRT flow cells of the PacBio Sequel II at a concentration of 90 pM according to PacBio's instructions, as provided by the SMRTlink software.

Mitochondrial DNA (mtDNA) sequencing

The mitochondrial genomes of this hare cell line has been sequenced for the purposes of another study by PCR amplification of approximately 2 kb overlapping mtDNA fragments (Tapanainen et al., 2024) (GenBank access# OR915850).

157   **Figure 1.** A) The geographic location and a view of the sampling biotope in Ilomantsi. Photo by Jaakko
158 Pohjoismäki. B) Although the cell line used in this study originates from a killed specimen, fibroblasts can
159 be isolated from e.g. ear clippings taken during tagging of animals for behavioral studies. Photo by Mervi
160 Kunnasranta. C) The genome presented here was generated using the LT1 cell line, exhibiting a typical
161 fibroblast morphology.

Hi-C library preparation

As with the brown hare genome assembly (Michell et al., 2024), Hi-C sequencing libraries were
prepared following the protocol of (Belaghzal et al., 2017) with the following changes: 1.) The Hi-C
protocol was performed in triplicates to produce a diverse sequencing library; 2.) NEBNext Ultra II FS
DNA module was used for size fractionation; 3.) the NEBNext Ultra II library preparation kit for Illumina
was utilized to obtain the sequencing libraries and 4.) triplicate PCR reactions with six cycles of PCR were
used for library enrichment. The PCR reactions were purified using Ampure XP beads at a ratio of 0.9X.
The final clean libraries were quantified using Qubit, followed by agarose gel electrophoresis to confirm
the fragment size. The sequencing was performed on a single lane of the Illumina NovaSeq 6000 using
the SP flowcell with paired-end chemistry 2 x 150bp.

Genome assembly

Quality of the sequencing data was assessed by FastQC v0.11.9, and cutadapt v4.6 was used to
remove any adapters or low-quality sequences. Prior to assembly, k-mer profiling was done using Meryl

v1.3 (https://github.com/marbl/meryl) and GenomeScope v2.0 (Vurture et al., 2017; Ranallo-Benavidez
et al., 2020), and assembly parameters adjusted based on the expected genome size and coverage.
HiFiasm version 0.16.1 (Cheng et al., 2021) was used to assemble the PacBio HiFi reads using the
arguments -m 10000000 -p 100000 –hom-cov 20, and –hic1 –hic2 to integrate the Hi-C read data and
produce two raw assemblies. Statistics and k-mer profiles of both assemblies were checked using
gfastats v1.3.6 (Formenti et al., 2022) and Merqury v1.3 (Rhie et al., 2020), and the most complete
assembly was chosen as primary, with the other kept as alternate. Duplicate contigs were removed using
purge_dups v1.2.6 (Guan et al., 2020) from both assemblies. We then continued with the scaffolding of
the assemblies. To process the HiC data, we followed the base pipeline by ArimaGenomics
(https://github.com/ArimaGenomics/mapping_pipeline). Briefly, we mapped the HiC data to the
genome assemblies using bwa-mem2 v2.2.1 (Vasimuddin et al., 2019). The mapped reads were then
parsed and filtered using samtools v1.18 (Danecek et al., 2021) and the scripts provided on the
ArimaGenomics github page above, and duplicate reads were removed using the MarkDuplicates tool of
the GATK toolkit v4.3.0.0 (O'Connor & van der Auwera, 2020).
For scaffolding, YaHS v1.2 (Zhou et al., 2023) was run using parameters -e GATC -l 5000 –no-contig-
ec for the assembly and the filtered Hi-C bam file. Contiguity and general genome statistics were
calculated using QUAST v5.2.0 (Mikheenko et al., 2018). We assessed the completeness of the genome
by calculating the number of complete single copy orthologs with BUSCO v5.5.0 (Manni et al., 2021),
using using both the glires_odb10 and mammalia_odb10 databases.

Telomere and repeat annotation

Repeat annotation of the genome was performed with RepeatModeler v2.0.5 (Flynn et al., 2020).
Using the repeat library produced by RepeatModeler, we masked the scaffolded genome using
RepeatMasker v4.1.6 (Smit et al., 2013-2015).
Telomeric sequences were identified using the Telomere Identification ToolKit (tidk v0.2.0) by first
running tidk explore to identify potential telomeric repeat sequences, and then tidk search with the
sequence of "AACCCT".

Manual curation

202   The assembled and annotated genome was manually curated to further improve its quality as
203 described in (Howe et al., 2021) using PretextView https://github.com/sanger-tol/PretextView/ and the
204 rapid curation workflow https://gitlab.com/wtsi-grit/rapid-curation. The manual curation allows the
205 identification and fixing of erroneous scaffold assemblies and contig duplications.

Comparison with previous assembly

207   We performed a comparison of our scaffolded assembly to the current *L. timidus* genome assembly
208 (NCBI Accession number: GCA_009760805) as well as the *L. europaeus* reference genome assembly (NCBI
209 accession number: GCF_033115175.1). Mapping to the genome was performed using minimap2 version
210 2.21 (Li, 2018) with the arguments *-asm5*. A dot plot of the alignment was created using the R script
211 pafCoordsDotPlotly.R (https://github.com/tpoorten/dotPlotly).

Results

213   The genome assemblies can be accessed via BioSample accession SAMN41430840 as well as
214 BioProject accession PRJNA1112569 for the primary assembly and raw data of the PacBio and Hi-C
215 sequencing, and PRJNA1112568 for the alternative assembly.

Genome assembly

The expected haploid genome size of *L. timidus* is 3.25 pg (3,178,500,000 bp) (Vinogradov, 1998),
containing 23 autosomal chromosomes as well as X and Y sex chromosomes (Gustavsson, 1972). PacBio
HiFi sequencing produced data with sequence N50 of 19.97 kb with 21× coverage. Based on the PacBio
HiFi data, the expected genome size using k-mer length k =28 is 2.65 Gbp (Figure 2A). The data enabled
haplotype-phasing of a primary and an alternative assembly (Figure 2B). The illumina sequencing of the
Hi-C data produced 995,011,507 paired reads representing about 56 × coverage of the genome. The
duplication rate of the Hi-C data was 18 %. Assembly with HiFiasm yielded a primary contig assembly of
2,838,807,612 bp made up of 2332 contigs and an alternate assembly of 2,654,293,704 bp of 1989 contigs
(The longest contigs for the assemblies were 33 Mbp and 26.6 Mbp, and the contig N50s 4.7 Mbp and 4.8
Mbp, respectively. Using the uniquely mapped Hi-C data, we were able to scaffold the contigs and fix
misassembled contigs. The primary Hi-C scaffolded assembly was 2,654,826,856 bp in size, the largest
scaffold was 181 Mbp and the scaffold N50 127 Mbp.
To further improve the assembly, manual curation was performed at the Sanger Institute using Hi-C
maps (Figure 2C). The manual curation changed the scaffold N50 of 127.6 Mb to 125.8 Mb and caused a
reduction in scaffold count of 103 to 85 in the primary and 128 to 109 in the alternate assembly. Of the
finalized assembly, 93.16 % could be assigned to 25 identified chromosomes (23 autosomes plus X and
Y) (Table 1). Autosomes were named according to their size. After decontamination and manual curation,
the scaffold counts are 85 for the primary, and 109 for the alternate assembly. Detailed final assembly
statistics are shown in Table 2 The curated genome length was 2,695,305,354 bp. The mitochondrial
genome assembly and the annotation of the functional loci have been presented elsewhere (Tapanainen
et al., 2024).
The BUSCO scores of the *L. timidus* assembly suggest that it is near-complete, with the following
results: Complete: 95.1 % [Single copy: 92.3 %, Duplicated: 2.7 %] Fragmented 0.8 %, and Missing 4.1 %
based on the mammalia_odb10 database and Complete: 93.2% [Single copy: 90.6 %, Duplicated: 2.7 %],
Fragmented 1.1 % and Missing 5.6 % based on the glires_odb10 database. The number of total groups
searched were 9226 and 13798, respectively (Table 3, Figure 3). As of note, T-antigen vector insertion was
detected on the chromosomes 17 and 19. The insertions were in intergenic regions and have been
replaced with equal length of Ns in the final assembly.

Figure 2. An overview of the mountain hare genome assembly. A) K-mer frequency and coverage
distribution obtained from the GenomeScope analysis. B) k-mer distribution profile after the first
assembly step with Hifiasm showing also the amount of unique k-mers for the primary (01) and
alternative (02) assemblies. C) Contact map of Hi-C scaffolding of the primary assembly after manual
curation, generated with PretextView.

259 **Table 1.** Sequence assignment of the primary mountain hare genome assembly.

	Length	% of total length	Count
Autosomes	2,519,453,057	93.48	23
Unlocalized on autosomes	1,279,896	0.05	1
Other scaffolds	11,202,567	0.42	58
X	139,721,860	5.18	1
Y	24,927,870	0.92	1
MT	17,482		1
TOTAL	2,695,305,354		85

260
261 **Table 2.** Assembly statistics for the Hi-C scaffolded, post-curation sequencing data for
262 the mountain hare reference genome.

Assembly statistics	Primary	Alternate
# scaffolds	85	109
Total scaffold length	2,695,305,354	2,556,802,277
Average scaffold length	31,709,474.75	23,456,901.62
Scaffold N50	125,755,317	126,936,470
Scaffold auN	125,197,910.08	126,023,208.11
Scaffold L50	9	9
Largest scaffold	182,608,553	185,135,719
Smallest scaffold	10,771	21,369
# contigs	1,414	1,454
Total contig length	2,695,169,802	2,556,665,250
Average contig length	1,906,060.68	1,758,366.75
Contig N50	4,885,811	5,114,768
Contig auN	6,632,874.02	6,902,786.36
Contig L50	157	136
Largest contig	32,947,725	26,587,001
Smallest contig	967	21,369
# gaps in scaffolds	1,329	1,345
Total gap length in scaffolds	135,552	137,027
Average gap length in scaffolds	102.00	101.88
Gap N50 in scaffolds	100	100
Gap auN in scaffolds	104.10	103.78
Gap L50 in scaffolds	650	660
Largest gap in scaffolds	200	200
Smallest gap in scaffolds	26	27
Base composition (A:C:G:T)		
A	758,096,831	716,813,053
C	589,442,155	561,406,584
G	589,603,553	561,320,925
T	758,027,263	717,124,688

GC content %	43.75	43.91
# soft-masked bases	0	0
# segments	1,414	1,454
Total segment length	2,695,169,802	2,556,665,250
Average segment length	1,906,060.68	1,758,366.75
# gaps	1,329	1,345
# paths	85	109

264 **Table 3.** BUSCO completeness statistics for the mountain hare reference genome
265 assembly.

BUSCO	Primary		Alternate	
Mammalia	**Count**	**%**	**Count**	**%**
Complete BUSCOs (C)	8770	95.06	8601	93.23
Complete and single-copy BUSCOs (S)	8517	92.32	8378	90.81
Complete and duplicated BUSCOs (D)	253	2.74	223	2.42
Fragmented BUSCOs (F)	76	0.82	84	0.91
Missing BUSCOs (M)	380	4.12	541	5.86
Total BUSCO groups searched	9226		9226	
Glires				
Complete BUSCOs (C)	12863	93.22	12553	90.98
Complete and single-copy BUSCOs (S)	12495	90.56	12226	88.61
Complete and duplicated BUSCOs (D)	368	2.67	327	2.37
Fragmented BUSCOs (F)	157	1.14	149	1.08
Missing BUSCOs (M)	778	5.64	1096	7.94
Total BUSCO groups searched	13798		13798	

Record statistics
- Log10 record count (total 85)
- Record length (total 2.70G)
- Longest record (183M)
- N50 length (126M)
- N90 length (69M)

BUSCO glires_odb10 (13798)
- Comp. (93.2%)
- Frag. (1.1%)
- Dupl. (2.7%)
- Missing (5.6%)

Scale
- 2.70G
- 183M

Dataset:

Composition
- GC (43.8%)
- AT (56.2%)
- N (0.0%)

Figure 3. A snail plot summary of assembly statistics for the final, manually curated 2,695,305,354 bp primary assembly. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1 % of the final assembly. The distribution of sequence lengths is shown in dark grey with the plot radius scaled to the longest sequence present in the assembly (182,608,553 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 sequence lengths (125,755,317 and 68,953,458 bp), respectively. The pale grey spiral shows the cumulative sequence count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the glires_odb10 set is shown in the top right.

Repeat detection with RepeatModeler produced a curated custom repeat library of 28,386 unique repetitive elements classified into 809 repeat families. Interestingly, the proportion of the genome masked as repetitive elements was very high, at 42.35 % of the genome, similar to the situation observed with our brown hare genome assembly (46.8 %) (Michell et al., 2024). In contrast, in the Irish hare genome assembly (GCA_009760805), only 28 % of the genome was masked (Marques et al., 2020). The *k*-mer based estimate of the repetitive elements in our mountain hare was 23.0 %, which is slightly less than the 26.7 % observed for the brown hare (Michell et al., 2024).

Telomeric sequences were found at both ends of 9 chromosomes, and only on one end of 10 more chromosomes (Figure 4). More than half of the chromosomes (13) also contain telomeric elements not only at their ends but also at various internal locations.

Figure 4. Telomeric repeat arrays on the mountain hare chromosome assemblies, calculated in windows
of 200 kb. The location and relative length of the repeat is presented by the red bars. Short telomeric
repeats are not visible in the graph.

Comparison to previous assemblies

Minimap2 was able to align 99.95 % (97.57 % primary mapped) of the contig sequences from the Irish
hare genome assembly with the presented mountain hare genome assembly (Figure 5A). The two
assemblies have a high level of sequence similarity as well as a high degree of synteny. As pointed out
earlier, the Irish hare genome assembly retains the chromosomal arrangement of the rabbit genome
(Beklemisheva et al., 2011) used for its scaffolding (Marques et al., 2020). As a result, a few notable
differences are present in the chromosome assignments, which were also noted when comparing the
Irish hare assembly with the brown hare assembly (Michell et al., 2024). Specifically, hare chromosomes
chr7 and chr12 correspond to Chr1 of the rabbit genome, and chromosomes chr13 and chr16 correspond
to rabbit Chr2 (Figure 5A), explaining the karyotype difference (2n = 44 vs. 2n = 48).

Consequently, the brown hare genome (Michell et al. 2023, NCBI RefSeq
GCF_033115175.1) shows an even higher degree of synteny with our mountain hare genome assembly
(Figure 5B), with Minimap2 aligning 99.87 % (78.77 % primary mapped) of the sequences. However, while
the visual representation of the two genomes doesn't show major rearrangements, the relatively lower
percentage of primary mapped contigs with minimap2 also means a higher amounts of chimeric
alignments, pointing to minor differences in chromosome sequence organization. Despite the high levels
of synteny, the sequence similarity between the brown hare and mountain hare assemblies is lower than
that between the mountain hare subspecies. This is illustrated by the sequence identity colors in the
alignments (Figure 5A vs. B). The mean sequence similarity, as calculated from the number of matches
between each of the aligned scaffolds, is 96.77 % for the mountain hare vs. Irish hare and 93.09 % for the
mountain hare vs. brown hare.

Figure 5. A comparison of the LT1 mountain hare (*Lepus timidus timidus*) genome assembly with A) the
previous Irish hare (*Lepus timidus hibernicus*) pseudoreference assembly (GCA_009760805, examples of
karyotype differences are highlighted in red) and B) our previously published European brown hare
(*Lepus europaeus*) genome assembly (GCF_033115175). Note the higher synteny but lower sequence
similarity compared to (A).

Discussion

To complement our previously published brown hare reference genome assembly (Michell et al.,
2024), we have sequenced and produced a reference-quality genome assembly for the mountain hare.
Although large and complex mammalian genomes can pose problems for assembly, our final genome
assembly is highly continuous (85 scaffolds in the primary assembly, with an N50 of 126 Mbp, and contig
N50 4.9 Mbp) and complete (BUSCO complete 95.1 % mammalia_odb10) (Tables 1–3, Figures 1–3). The
genome assembly is at chromosome scale, with all expected 23 autosomes as well as X and Y scaffolded.
Most of the chromosome assemblies represent telomere-to-telomere continuity (Figure 4). High copy
number telomeric sequences were also present within interstitial positions of some chromosomes,
confirming previous observations using FISH staining of mountain hare cells (Forsyth et al., 2005). These
regions might be remnants of ancestral chromosome translocation events that gave rise to the current
2n = 46 karyotype in hares.
As in the case of our brown hare genome assembly (Michell et al., 2024), we utilized living cultured
fibroblasts to overcome the technical challenges in obtaining large quantities of intact, high molecular
weight DNA for HiFi sequencing, essential for high quality and highly continuous genome assemblies.
Importantly, our mountain hare genome assembly did not show any genomic rearrangements or
aneuploidies that would have been obtained during cell culture. Although genomic changes can occur
during cell culture (Didion et al., 2014), these are more likely in cancer cell lines with compromised DNA
repair and cell proliferation pathways (He et al., 2023). Genetic changes need time as they appear by
chance and increase their prevalence through drift. Therefore, long-term culturing of the cells should be
avoided, and low-passage number cells should be preferred for analyses. However, despite being devoid
of karyotypic changes, we noted the insertion of the T-antigen vector used for immortalization of the
cells (Gaertner et al., 2022) on chromosome 17 and 19. Interestingly, such insertions are not present in
our brown hare genome assembly (Michell et al., 2024), indicating that the immortalization of this brown
hare cell line has not worked. Although the insertions were in the intergenic region and we have removed

the vector sequences from the final genome assembly, we recognize that this is not an ideal situation as we cannot control for possible insertional effects on the surrounding chromosomal region. Therefore, we recommend that primary cell lines should be used for reference genome work. For other types of studies it needs to be taken into consideration that non-immortalized cell lines have limited lifetime and can show signs of senescence, which can influence analyses.

The presented mountain hare genome assembly shares a high degree of similarity with the existing Irish hare pseudo-reference genome assembly. The observed differences in synteny (Figure 5A) can be attributed to use of the karyotypically different European rabbit (*Oryctolagus cuniculus)* for the chromosomal scaffolding of the Irish hare genome assembly (Marques et al., 2020). Compared to the recently published brown hare genome assembly (Michell et al., 2024), the presented mountain hare genome assembly shows almost identical synteny (Figure 5B), reflecting their close evolutionary relationship. However, when compared to the Irish hare, mountain hare – brown hare comparison yielded higher amounts of chimeric alignments, likely due to minor differences in chromosome structure. These structural differences are unlikely to have a major biological significance, given that the two species can hybridize and produce fertile offspring.

The 93.09 % sequence similarity of the mountain hare and brown hare genomes is interesting to contrast with the 96.77 % similarity between the mountain hare subspecies. It should be noted that these similarities are based on matches between scaffold alignments, and comparing specific elements, such as coding sequences alone, would likely yield higher similarity percentages. Nonetheless, these figures highlight the relative differences between species and subspecies genomes, particularly as the Irish hare is known to be genetically distinct from Fennoscandian mountain hares (Giska et al., 2022). More detailed population-level comparisons might allow to pinpoint functional loci contributing to local adaptations (Giska et al., 2019) as well as shed light onto speciation processes (Gaertner et al., 2022).

The mountain hare is an iconic boreoalpine mammalian species with unique physiological and morphological adaptations to cold, snowy winter conditions, it is also an important part of the local food webs. Consequently, the population genetics, evolution and genomics of mountain hares has been intensively studied [e.g. (Melo-Ferreira et al., 2009; Smith et al., 2017; Levänen, Thulin, et al., 2018; Marques et al., 2020; Ferreira et al., 2021; Pohjoismäki et al., 2021; Gaertner et al., 2022; Giska et al., 2022; Michell et al., 2022)]. Together, this genome and the previously published brown hare genome assembly will provide a solid base for future studies on hares. Annotated genome assemblies also enable effective use of reverse genetics, allowing experimental manipulation of any gene or functional feature to study its phenotypic effects. This capability opens opportunities to experimentally demonstrate the functional significance of genetic variants controlling interesting traits, which otherwise could only be indirectly inferred from population genetic analyses. For population genetics, chromosomally scaffolded genome assemblies enable fast and reliable identification of informative SNPs and their linkage. Specifically, the mountain hare genome assembly may offer crucial insights for the conservation of threatened subspecies such as the heath hare (*Lepus timidus sylvaticus*) (C. G. Thulin et al., 2021; Michell et al., 2022) and Irish hare (*Lepus timidus hibernicus*) (Reid, 2011; Caravaggi et al., 2017).

In conclusion, the successful sequencing and assembly of the mountain hare genome, alongside the previously published brown hare genome, mark significant advancements in our understanding of these species' genomic landscapes. These high-quality genome assemblies not only provide invaluable resources for exploring the genetic basis of adaptation and resilience in hares, but also pave the way for future research in evolutionary biology, conservation genetics, and ecological genomics.

Acknowledgements

We acknowledge the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology, University of Helsinki for the PacBio and Hi-C sequencing. We are grateful to the European Reference

Genome Atlas (ERGA) for the opportunity to include our mountain hare genome into their pilot project. Mr Jussi Mähönen, MSc Manu Soininmäki and Finnish hound Hupi are thanked for their contribution in obtaining the mountain hare specimen. We are indebted also to Ms Anita Kervinen for her valuable laboratory assistance and Dr. Mervi Kunnasranta for kindly allowing to use her mountain hare photo in the publication.

Funding

This study belongs to the xHARES consortium funded by the R'Life initiative of the Academy of Finland, grant no. 329264.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

The genome assembly and sequencing data is available from the NCBI database under BioProject ID PRJNA1112569 and PRJNA1112568 and accession numbers JBEDNU000000000 and JBEDNT000000000. The cell line is available through the corresponding author as well as from the ZFMK biobank.

References

Angerbjörn A, Schai-Braun SC (2022) Mountain Hare Lepus timidus Linnaeus, 1758. . In: *Handbook of the Mammals of Europe. Handbook of the Mammals of Europe* eds Hackländer K & Zachos FE), pp. 1-29. Springer, Cham.

ATCC. (2024). Passage number effects in cell lines. . *ATCC Techinical Documents.* Retrieved from https://www.atcc.org/-/media/resources/technical-documents/passage-number-effects-in-cell-lines.pdf

Beklemisheva VR, Romanenko SA, Biltueva LS, Trifonov VA, Vorobieva NV, Serdukova NA, Rubtsova NV, Brandler OV, O'Brien PC, Yang F, Stanyon R, Ferguson-Smith MA, Graphodatsky AS (2011) Reconstruction of karyotype evolution in core Glires. I. The genome homology revealed by comparative chromosome painting. *Chromosome Res,* **19**, 549-565. https://doi.org/10.1007/s10577-011-9210-y

Belaghzal H, Dekker J, Gibcus JH (2017) Hi-C 2.0: An optimized Hi-C procedure for high-resolution genome-wide mapping of chromosome conformation. *Methods,* **123**, 56-65. https://doi.org/10.1016/j.ymeth.2017.04.004

Blaxter M, Archibald JM, Childers AK, Coddington JA, Crandall KA, Di Palma F, Durbin R, Edwards SV, Graves JAM, Hackett KJ, Hall N, Jarvis ED, Johnson RN, Karlsson EK, Kress WJ, Kuraku S, Lawniczak MKN, Lindblad-Toh K, Lopez JV, Moran NA, Robinson GE, Ryder OA, Shapiro B, Soltis PS, Warnow T, Zhang GJ, Lewin HA (2022) Why sequence all eukaryotes? *Proceedings of the National Academy of Sciences of the United States of America,* **119**. https://doi.org/ARTN e2115636118

10.1073/pnas.2115636118

Caravaggi A, Leach K, Santilli F, Rintala J, Helle P, Tiainen J, Bisi F, Martinoli A, Montgomery WI, Reid N (2017) Niche overlap of mountain hare subspecies and the vulnerability of their ranges to invasion by the European hare; the (bad) luck of the Irish. *Biological Invasions,* **19**, 655-674. https://doi.org/10.1007/s10530-016-1330-z

Cheng H, Concepcion GT, Feng X, Zhang H, Li H (2021) Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nat Methods,* **18**, 170-175. https://doi.org/10.1038/s41592-020-01056-5

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM, Li H (2021) Twelve years of SAMtools and BCFtools. *GigaScience,* **10**. https://doi.org/10.1093/gigascience/giab008

Didion JP, Buus RJ, Naghashfar Z, Threadgill DW, Morse HC, de Villen FPM (2014) SNP array profiling of mouse cell lines identifies their strains of origin and reveals cross-contamination and widespread aneuploidy. *Bmc Genomics,* **15**. https://doi.org/Artn 847

10.1186/1471-2164-15-847

Ferreira MS, Jones MR, Callahan CM, Farelo L, Tolesa Z, Suchentrunk F, Boursot P, Mills LS, Alves PC, Good JM, Melo-Ferreira J (2021) The Legacy of Recurrent Introgression during the Radiation of Hares. *Syst Biol,* **70**, 593-607. https://doi.org/10.1093/sysbio/syaa088

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF (2020) RepeatModeler2 for automated genomic discovery of transposable element families. *Proc Natl Acad Sci U S A,* **117**, 9451-9457. https://doi.org/10.1073/pnas.1921046117

Formenti G, Abueg L, Brajuka A, Brajuka N, Gallardo-Alba C, Giani A, Fedrigo O, Jarvis ED (2022) Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs. *Bioinformatics,* **38**, 4214-4216. https://doi.org/10.1093/bioinformatics/btac460

Forsyth NR, Elder FF, Shay JW, Wright WE (2005) Lagomorphs (rabbits, pikas and hares) do not use telomere-directed replicative aging in vitro. *Mech Ageing Dev,* **126**, 685-691. https://doi.org/10.1016/j.mad.2005.01.003

Gaertner K, Michell C, Tapanainen R, Goffart S, Saari S, Soininmaki M, Dufour E, Pohjoismaki JLO (2022) Molecular phenotyping uncovers differences in basic housekeeping functions among closely related species of hares (Lepus spp., Lagomorpha: Leporidae). *Mol Ecol*. https://doi.org/10.1111/mec.16755

Giska I, Farelo L, Pimenta J, Seixas FA, Ferreira MS, Marques JP, Miranda I, Letty J, Jenny H, Hacklander K, Magnussen E, Melo-Ferreira J (2019) Introgression drives repeated evolution of winter coat color polymorphism in hares. *Proc Natl Acad Sci U S A,* **116**, 24150-24156. https://doi.org/10.1073/pnas.1910471116

Giska I, Pimenta J, Farelo L, Boursot P, Hacklander K, Jenny H, Reid N, Montgomery WI, Prodohl PA, Alves PC, Melo-Ferreira J (2022) The evolutionary pathways for local adaptation in mountain hares. *Mol Ecol,* **31**, 1487-1503. https://doi.org/10.1111/mec.16338

Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R (2020) Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics,* **36**, 2896-2898. https://doi.org/10.1093/bioinformatics/btaa025

Gustavsson I (1972) Mitotic and meiotic chromosomes of the variable hare (Lepus timidus L.), the common hare (Lepus europaeus Pall.) and their hybrids. *Hereditas,* **67**, 27-34. https://doi.org/10.1111/j.1601-5223.1971.tb02355.x

He ZC, Wilson A, Rich F, Kenwright D, Stevens A, Low YS, Thunders M (2023) Chromosomal instability and its effect on cell lines. *Cancer Reports,* **6**. https://doi.org/10.1002/cnr2.1822

Howe K, Chow W, Collins J, Pelan S, Pointon D-L, Sims Y, Torrance J, Tracey A, Wood J (2021) Significantly improving the quality of genome assemblies through curation. *GigaScience,* **10**. https://doi.org/10.1093/gigascience/giaa153

Jansson G, Pehrson A (2007) The recent expansion of the brown hare (Lepus europaeus) in Sweden with possible implications to the mountain hare (L. timidus). *European Journal of Wildlife Research,* **53**, 125-130. https://doi.org/10.1007/s10344-007-0086-2

Korstanje R, O'Brien PC, Yang F, Rens W, Bosma AA, van Lith HA, van Zutphen LF, Ferguson-Smith MA (1999) Complete homology maps of the rabbit (Oryctolagus cuniculus) and human by reciprocal chromosome painting. *Cytogenet Cell Genet,* **86**, 317-322. https://doi.org/10.1159/000015325

Lawniczak MKN, Durbin R, Flicek P, Lindblad-Toh K, Wei X, Archibald JM, Baker WJ, Belov K, Blaxter ML, Marques Bonet T, Childers AK, Coddington JA, Crandall KA, Crawford AJ, Davey RP, Di Palma F, Fang Q, Haerty W, Hall N, Hoff KJ, Howe K, Jarvis ED, Johnson WE, Johnson RN, Kersey PJ, Liu X, Lopez JV, Myers EW, Pettersson OV, Phillippy AM, Poelchau MF, Pruitt KD, Rhie A, Castilla-Rubio JC, Sahu SK, Salmon NA, Soltis PS, Swarbreck D, Thibaud-Nissen F, Wang S, Wegrzyn JL, Zhang G, Zhang H, Lewin HA, Richards S (2022) Standards recommendations for the Earth BioGenome Project. *Proc Natl Acad Sci U S A,* **119**. https://doi.org/10.1073/pnas.2115639118

Levänen R, Kunnasranta M, Pohjoismäki J (2018) Mitochondrial DNA introgression at the northern edge of the brown hare (Lepus europaeus) range. *Annales Zoologici Fennici,* **55**, 15-24

Levänen R, Thulin CG, Spong G, Pohjoismäki JLO (2018) Widespread introgression of mountain hare genes into
Fennoscandian brown hare populations. *Plos One,* **13**. https://doi.org/ARTN e0191790
10.1371/journal.pone.0191790
Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics,* **34**, 3094-3100.
https://doi.org/10.1093/bioinformatics/bty191
Manni M, Berkeley MR, Seppey M, Zdobnov EM (2021) BUSCO: Assessing Genomic Data Quality and Beyond.
Curr Protoc, **1**, e323. https://doi.org/10.1002/cpz1.323
Marques JP, Seixas FA, Farelo L, Callahan CM, Good JM, Montgomery WI, Reid N, Alves PC, Boursot P, Melo-
Ferreira J (2020) An Annotated Draft Genome of the Mountain Hare (Lepus timidus). *Genome Biol Evol,*
12, 3656-3662. https://doi.org/10.1093/gbe/evz273
Mc Cartney AM, Formenti G, Mouton A, De Panis D, Marins LS, Leitao HG, Diedericks G, Kirangwa J, Morselli M,
Salces J, Escudero N, Iannucci A, Natali C, Svardal H, Fernandez R, De Pooter T, Joris G, Strazisar M,
Wood J, Herron KE, Seehausen O, Watts PC, Shaw F, Davey RP, Minotto A, Fernandez Gonzalez JM,
Bohne A, Alegria C, Alioto T, Alves PC, Amorim IR, Aury J-M, Backstrom N, Baldrian P, Ballarin L,
Baltrunaite L, Barta E, BedHom B, Belser C, Bergsten J, Bertrand L, Bilandija H, Binzer-Panchal M,
Bista I, Blaxter M, Borges PA, Borges Dias G, Bosse M, Brown T, Bruggmann R, Buena-Atienza E, Burgin
J, Buzan E, Cariani A, Casadei N, Chiara M, Chozas S, Ciampor F, Crottini A, Cruaud C, Cruz F, Dalen L,
De Biase A, del Campo J, Delic T, Dennis AB, Derks MF, Diroma MA, Djan M, Duprat S, Eleftheriadi K,
Feulner PG, Flot J-F, Forni G, Fosso B, Fournier P, Fournier-Chambrillon C, Gabaldon T, Garg S, Gissi C,
Giupponi L, Gomez-Garrido J, Gonzalez J, Grilo ML, Gruening B, Guerin T, Guiglielmoni N, Gut M,
Haesler MP, Hahn C, Halpern B, Harrison P, Heintz J, Hindrikson M, Hoglund J, Howe K, Hughes G,
Istace B, Cock MJ, Jancekovic F, Jonsson ZO, Joye-Dind S, Koskimaki JJ, Krystufek B, Kubacka J, Kuhl
H, Kusza S, Labadie K, Lahteenaro M, Lantz H, Lavrinienko A, Leclere L, Lopes RJ, Madsen O,
Magdelenat G, Magoga G, Manousaki T, Mappes T, Marques JP, Martinez Redondo GI, Maumus F,
McCarthy SA, Megens H-J, Melo-Ferreira J, Mendes SL, Montagna M, Moreno J, Mosbech M-B, Moura
M, Musilova Z, Myers E, Nash WJ, Nater A, Nicholson P, Niell M, Nijland R, Noel B, Noren K, Oliveira PH,
Olsen R-A, Ometto L, Oomen RA, Ossowski S, Palinauskas V, Palsson S, Panibe JP, Pauperio J, Pavlek
M, Payen E, Pawlowska J, Pellicer J, Pesole G, Pimenta J, Pippel M, Pirttila AM, Poulakakis N, Rajan J,
Rego RM, Resendes R, Resl P, Riesgo A, Rodin-Morch P, Soares AE, Rodriguez Fernandes C, Romeiras
MM, Roxo G, Ruber L, Ruiz-Lopez MJ, Saarma U, Silva LP, Sim-Sim M, Soler L, Sousa VC, Sousa Santos
C, Spada A, Stefanovic M, Steger V, Stiller J, Stock M, Struck THH, Sudasinghe H, Tapanainen R,
Tellgren-Roth C, Trindade H, Tukalenko Y, Urso I, Vacherie B, Van Belleghem SM, van Oers K, Vargas-
Chavez C, Velickovic N, Vella N, Vella A, Vernesi C, Vicente S, Villa S, Vinnere Pettersson O, Volckaert
FA, Voros J, Wincker P, Winkler S, Ciofi C, Waterhouse RM, Mazzoni CJ (2024) The European Reference
Genome Atlas: piloting a decentralised approach to equitable biodiversity genomics. *bioRxiv*,
2023.2009.2025.559365. https://doi.org/10.1101/2023.09.25.559365
Melo-Ferreira J, Alves PC, Freitas H, Ferrand N, Boursot P (2009) The genomic legacy from the extinct Lepus
timidus to the three hare species of Iberia: contrast between mtDNA, sex chromosomes and
autosomes. *Mol Ecol,* **18**, 2643-2658. https://doi.org/10.1111/j.1365-294X.2009.04221.x
Michell CT, Collins J, Laine PK, Fekete Z, Tapanainen R, Wood JMD, Goffart S, Pohjoismäki JLO (2024) High
quality genome assembly of the brown hare (Lepus europaeus) with chromosome-level scaffolding.
Peer Community Journal, **4**, e26. https://doi.org/10.24072/pcjournal.393
Michell CT, Pohjoismäki JLO, Spong G, Thulin CG (2022) Mountain- and brown hare genetic polymorphisms to
survey local adaptations and conservation status of the heath hare (Lepus timidus sylvaticus, Nilsson
1831). *Sci Data,* **9**, 667. https://doi.org/10.1038/s41597-022-01794-5
Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A (2018) Versatile genome assembly evaluation with
QUAST-LG. *Bioinformatics,* **34**, i142-i150. https://doi.org/10.1093/bioinformatics/bty266
O'Connor BD, van der Auwera G (2020) *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*. O'Reilly
Media, Incorporated.
Pohjoismäki JLO, Michell C, Levänen R, Smith S (2021) Hybridization with mountain hares increases the
functional allelic repertoire in brown hares. *Sci Rep,* **11**, 15771. https://doi.org/10.1038/s41598-021-
95357-0
Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020) GenomeScope 2.0 and Smudgeplot for reference-free
profiling of polyploid genomes. *Nat Commun,* **11**, 1432. https://doi.org/10.1038/s41467-020-14998-3

Rehnus M, Bollmann K, Schmatz DR, Hacklander K, Braunisch V (2018) Alpine glacial relict species losing out to climate change: The case of the fragmented mountain hare population (Lepus timidus) in the Alps. *Glob Chang Biol,* **24**, 3236-3253. https://doi.org/10.1111/gcb.14087

Reid N (2011) European hare (Lepus europaeus) invasion ecology: implication for the conservation of the endemic Irish hare (Lepus timidus hibernicus). *Biological Invasions,* **13**, 559-569. https://doi.org/10.1007/s10530-010-9849-x

Rhie A, Walenz BP, Koren S, Phillippy AM (2020) Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biol,* **21**, 245. https://doi.org/10.1186/s13059-020-02134-9

Sambrook J, Russell DW (2006) Purification of Nucleic Acids by Extraction with Phenol:Chloroform. *Cold Spring Harbor Protocols,* **1**, pdb.prot4455. https://doi.org/doi:10.1101/pdb.prot4455

Smit AFA, Hubley R, Green P. (2013-2015). RepeatMasker Open-4.0. http://www.repeatmasker.org.

Smith S, Sandoval-Castellanos E, Lagerholm VK, Napierala H, Sablin M, Von Seth J, Fladerer FA, Germonpre M, Wojtal P, Miller R, Stewart JR, Dalen L (2017) Nonreceding hare lines: genetic continuity since the Late Pleistocene in European mountain hares (Lepus timidus). *Biological Journal of the Linnean Society,* **120**, 891-908

Tapanainen R, Aasumets K, Fekete Z, Goffart S, Dufour E, Pohjoismäki JLO (2024) Species-specific variation in mitochondrial genome tandem repeat polymorphisms in hares (Lepus spp., Lagomorpha, Leporidae) provides insight into their evolution. *bioRxiv*, 2024.2003.2004.583303. https://doi.org/10.1101/2024.03.04.583303

Thulin C-G (2003) The distribution of mountain hares (Lepus timidus, L. 1758) in Europe: A challenge from brown hares (L. europaeus, Pall. 1778)? *Mammal Review,* **33**, 29-42. https://doi.org/DOI 10.1046/j.1365-2907.2003.00008.x

Thulin CG, Winiger A, Tallian AG, Kindberg J (2021) Hunting harvest data in Sweden indicate precipitous decline in the native mountain hare subspecies Lepus timidus sylvaticus (heath hare). *Journal for Nature Conservation,* **64**. https://doi.org/ARTN 126069

10.1016/j.jnc.2021.126069

Vasimuddin M, Misra S, Li H, Aluru S. (2019). *Efficient architecture-aware acceleration of BWA-MEM for multicore systems*. Paper presented at the IEEE International Parallel and Distributed Processing Symposium (IPDPS).

Vinogradov AE (1998) Genome size and GC-percent in vertebrates as determined by flow cytometry: The triangular relationship. *Cytometry,* **31**, 100-109. https://doi.org/Doi 10.1002/(Sici)1097-0320(19980201)31:2<100::Aid-Cyto5>3.0.Co;2-Q

Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, Schatz MC (2017) GenomeScope: fast reference-free genome profiling from short reads. *Bioinformatics,* **33**, 2202-2204. https://doi.org/10.1093/bioinformatics/btx153

Zhou C, McCarthy SA, Durbin R (2023) YaHS: yet another Hi-C scaffolding tool. *Bioinformatics,* **39**. https://doi.org/10.1093/bioinformatics/btac808

Zimova M, Giery ST, Newey S, Nowak JJ, Spencer M, Mills LS (2020) Lack of phenological shift leads to increased camouflage mismatch in mountain hares. *Proc Biol Sci,* **287**, 20201786. https://doi.org/10.1098/rspb.2020.1786

Zimova M, Mills LS, Nowak JJ (2016) High fitness costs of climate change-induced camouflage mismatch. *Ecology Letters,* **19**, 299-307. https://doi.org/10.1111/ele.12568