Molecular tension authenticates apoptotic cells being phagocytosed

Chanhyuk Min[1,2,3]†, Hyeokjin Cho[1,2]†, Jeongmin Yu[1,2]†, Byeongjin Moon[1], Jaeseon Jeon[1],

Nafeesa Shahdab[1]‡, Hyunji Moon[1] , Susumin Yang[1], Juyeon Lee[1], Jaeseong Jin[4], Mingi Hong[1,2],

Won-jin Chung[4], Jihwan Park[1], Gwangrog Lee[2,3]*, Daeho Park[1]*

[1]School of Life Sciences, Gwangju Institute of Science and Technology; Gwangju 61005, Korea

[2]Protein Engineering & Mechano-ImmunoTheraphy Laboratory, Korea Advanced Institute of Science and

Technology; Daejeon 34141, Korea

[3]Department of Biological Sciences, Korea Advanced Institute of Science and Technology; Daejeon

34141, Korea

[4]Department of Chemistry, Gwangju Institute of Science and Technology; Gwangju 61005, Korea

†These authors contributed equally: Chanhyuk Min, Hyeokjin Cho, Jeongmin Yu

‡Current Affiliation: School of Health and Life Sciences, Teesside University; Middlesbrough, United

Kingdom

19 *Correspondence and requests for materials should be addressed to:

20 G.L. (email: ifglee@kaist.ac.kr) or D.P. (email: daehopark@gist.ac.kr)

Abstract

Profound cytoskeletal reorganization and plasma membrane deformation in phagocytes is indispensable for phagocytosis of massive apoptotic cells, but whether these dynamics accompany a mechanical signal modulating signaling during efferocytosis remains largely unexplored. Here, we report that tension between phosphatidylserine (PS) and PS receptors (PSR) generated at the phagocytic synapse serves as a signal to determine whether PS-exposing cells should be phagocytosed. Mechanistically, increased membrane tension of phagocytes via Rac1-dependent actin polymerization and membrane stiffness of apoptotic cells caused tension between PS and PSRs, leading to phosphoinositide 3-kinase recruitment to PSRs, which resulted in Rac1 inactivation and myosin II phosphorylation required for phagocytic cup closure. Our observations imply that tension between PS and PSRs acts as a mechanical fail-safe to prevent removal of all PS-exposing cells for the integrity of efferocytosis.

One-Sentence Summary

Tension between phosphatidylserine and its receptors acts as a final decision-maker determining phagocytosis of apoptotic cells.

Main Text

Phagocytosis of apoptotic cells, called efferocytosis, differs from other types of phagocytosis in

terms of the endogeny and size of the targets. Specifically, this process removes endogenously

generated apoptotic cells that are as large as phagocytes (*1, 2*). Elaborately coordinated

signaling between two types of biochemical signals, called 'eat-me' and 'don't eat-me' signals,

enables phagocytes to remove such endogenous and massive targets. Phagocytes sense cells

to be phagocytosed through eat-me signals, of which phosphatidylserine (PS) is the most well-

known, and cells not to be phagocytosed through don't eat-me signals (*3-8*). Accordingly, when

eat-me signals are present and don't eat-me signals are nullified on the cell surface, phagocytes

are convinced that the cells are apoptotic cells that should be phagocytosed, and ultimately

efferocytosis may occur (*9, 10*). Furthermore, concerted signaling between these two signals

induces cytoskeletal rearrangement, which results in plasma membrane deformation, allowing

phagocytes to engulf the massive targets (*11-13*). Nevertheless, it has not been sufficiently

assessed whether the two signals are decisive for efferocytosis to occur. In particular, the

drastic plasma membrane deformation and cytoskeletal rearrangement in phagocytes may

induce a tensile force (tension) between molecules interacting intra- and intercellularly on the

plasma membrane (e.g., PS on apoptotic cells and its receptors (PSRs) on phagocytes). This

force may interconnect with the conventional signaling of efferocytosis. However, it remains

unclear whether such a force exists and, if so, what role it plays in efferocytosis. In this study,

we investigated the existence, generation, and role of tension between PS and PSRs (referred

to as tension of PS-PSRs) in efferocytosis using various approaches including surrogates of

apoptotic cells that can be used to adjust tension of PS-PSRs and time-lapse confocal

microscopy.

To investigate whether tension of PS-PSRs exists during and is required for

efferocytosis, we devised beads with PS connected via a tension gauge tether (*14*) (PS-TGT) as

surrogates of apoptotic cells. In PS-TGT beads, PS is covalently conjugated to the Cy3- or Cy5-

labeled upper strand of a double-stranded DNA tether that is immobilized on neutravidin beads

and the TGT ruptures at a critical force, termed tension tolerance (Fig. 1A). When J774A.1 cells,

a murine macrophage cell line, were incubated with PS-TGT beads, the cells engulfed beads

with a 56 pN tension tolerance TGT (PS-TGT^{56pN}) more efficiently than beads with a 12 pN

tension tolerance TGT (PS-TGT^{12pN}), as measured by the number of engulfed beads per

phagocyte and the percentage of phagocytes engulfing the beads (Fig. 1, B and C, and movie

S1). Other macrophages, including peritoneal macrophages, bone marrow-derived

macrophages, and RAW264.7 cells, also engulfed PS-TGT^{56pN} beads more efficiently than PS-

TGT^{12pN} beads (fig. S1, A to C). However, the superior engulfment of TGT^{56pN} beads relative to

TGT^{12pN} beads by macrophages was not observed when phosphatidylcholine (PC) was

conjugated to the TGT instead of PS (Fig. 1D). Notably, these results were not dependent on

the amount of TGT immobilized on the beads because PS-TGT^{56pN} beads were consistently

engulfed more efficiently than PS-TGT^{12pN} beads and engulfment of PC-TGT^{56pN} beads was still

comparable with that of PC-TGT^{12pN} beads when various amounts of the TGT were immobilized

on the beads (fig. S1D).

We then investigated whether the TGT is indeed ruptured during engulfment of PS-TGT

beads. Several observations indicated that PS-TGT^{12pN}, but not PS-TGT^{56pN}, was ruptured

during engulfment of the beads. First, the Cy5 mean fluorescence intensity (MFI) of non-

engulfing phagocytes was higher upon incubation with PS-TGT^{12pN} beads than upon incubation

with PS-TGT^{56pN} or control TGT beads (Fig. 1E), implying that the ruptured Cy5-labeled upper

strand of PS-TGT^{12pN} is present in non-engulfing phagocytes. Second, after macrophages were

incubated on a polyethylene glycol (PEG)-passivated glass plate on which PS-TGT was

immobilized via a biotin-neutravidin interaction (*15*), cell detachment from the plate was

evaluated. The cell number was comparable on the PS-TGT^{12pN} and PS-TGT^{56pN} plates after

incubation for 5 min. However, after incubation for 60 min, few cells were present on the PS-

TGT^{12pN} plate, whereas most cells remained attached to the PS-TGT^{56pN} plate (Fig. 1F). This

also implies that PS-TGT^{12pN}, but not PS-TGT^{56pN}, is ruptured because the cells were attached

to the plate only through PS-TGT (fig. S1E). Third, the loss of Cy5 fluorescence, indicative of

rupture of the Cy5-labeled upper strand of the TGT, was greater with PS-TGT^{12pN} than with PS-

TGT^{56pN} when macrophages were incubated on glass plates (Fig. 1G and fig. S1F). Collectively,

these data imply that the degree of rupture of the PS-TGT differs depending on tension

tolerance during engulfment of PS-TGT beads, indicating that tension is loaded between PS

and PSRs during and is essential for efferocytosis.

The rupture of PS-TGT^{12pN} may cause the ligand to separate from the target, preventing

further signaling and potentially leading to inefficient engulfment of the PS-TGT^{12pN} beads. To test

this, we designed hairpin PS-TGTs with variable loop sizes, created using DNA or PEG, that

prevent the ligand from separating from the phagocytic target even when TGT rupture occurs. In

this measurement, the upper and lower strands of 12 pN TGT were connected by loops made of

PEG or ssDNA, creating TGTs with various tensions depending on the length of the loop (Fig. 1H).

For the hairpin TGTs with ssDNA after rupture (Fig. 1I), they are subjected to minimal entropic

tension of ~2.3 and ~3.6 pN for the 16-bp- and 50-bp-ssDNA loops, respectively, based on the

length-dependent elasticity of ssDNA. The results showed that macrophages still engulfed the

hairpin PS-TGT^{12pN} beads with significantly lower efficiency than hairpin or normal PS-TGT^{56pN}

beads, and there were no differences between engulfment of normal PS-TGT^{12pN} and hairpin PS-

TGT^{12pN} beads. Nevertheless, in the case of the PEG loop (Fig. 1J), as the loop size increased,

engulfment decreased at both 12 pN and 56 pN, which is believed to be due to increased steric

hindrance as loop size increases. Regardless of the loop size and type, the engulfment of hairpin

PS-TGT^{12pN} beads remained less efficient compared to the engulfment of hairpin PS-TGT^{56pN}

beads (Fig. 1, I and J), implying that inefficient engulfment of PS-TGT^{12pN} beads is not due to the

termination of signaling caused by the disassembly of the ligand from the phagocytic target.

We next investigated how tension of PS-PSRs affects engulfment of PS-TGT beads. To

this end, we scrutinized the binding, internalization, and degradation steps of efferocytosis. The

numbers of PS-TGT^{12pN} and PS-TGT^{56pN} beads bound to macrophages were comparable when

cells were incubated with the beads at 4°C (fig. S2A). F-actin formation, which is required for

internalization of apoptotic cells during efferocytosis and was monitored by phalloidin staining,

was also comparable around PS-TGT^{12pN} and PS-TGT^{56pN} beads (fig. S2B). In addition, neither

the acidification of phagolysosomes nor the degradation of phagolysosomal cargos, as measured

by the change rates of the fluorescence intensities of pHrodo and Cy3, respectively, was altered

according to the tension tolerance of PS-TGT beads (fig. S2, C and D, and movies S2 and S3).

However, unexpectedly, time-lapse observation of F-actin using SiR-actin, a live F-actin labeling

probe, revealed that F-actin persisted longer around PS-TGT^{12pN} beads than around PS-TGT^{56pN}

beads (Fig. 2, A and D, and movie S4). Disassembly of F-actin is synchronized with phagocytic

cup closure during efferocytosis (*16, 17*); therefore, we next monitored phagocytic cup closure

depending on tension tolerance during engulfment of PS-TGT beads. Macrophages often failed

to close the phagocytic cup to completely engulf PS-TGT^{12pN} beads and consequently it took

longer to close the phagocytic cup for engulfment of PS-TGT^{12pN} beads than for engulfment of

PS-TGT^{56pN} beads (Fig. 2, C and D, and movie S5). These data imply that tension of PS-PSRs

affects efferocytosis by modulating phagocytic cup closure but not by modulating the binding

capacity, F-actin formation, or phagolysosomal function of phagocytes during efferocytosis.

We then investigated how tension of PS-PSRs modulates phagocytic cup closure

during engulfment of PS-TGT beads. Rac1 is involved in an early signaling pathway leading to

efferocytosis and its activity regulates F-actin dynamics (*11, 18*). In particular, spatiotemporal

regulation of Rac1 activity at the phagocytic cup is crucial for phagocytic cup formation and

closure (*16*). Therefore, we investigated whether the tension affects Rac1 activity. To this end,

we used Raichu-Rac1, a biosensor that monitors Rac1 activity via fluorescence resonance

energy transfer (FRET) (*19*). Macrophages expressing Raichu-Rac1 were incubated with PS-

TGT beads, and FRET was measured at the interface between macrophages and the beads

from 0° to 180° in a circular fashion as a function of time. Rac1 activation was initiated at the

junction of macrophages and the beads and proceeded along the tip of the phagocytic cup. At

0–10 min, Rac1 was activated with similar FRET levels in phagocytes incubated with PS-

TGT^{12pN} and PS-TGT^{56pN} beads. By contrast, at 10–20 min, Rac1 activation was instantly

followed by its inactivation at the phagocytic cup of phagocytes incubated with PS-TGT^{56pN}

beads, but not of phagocytes incubated with PS-TGT^{12pN} beads (Fig. 2, E and F, and movies S6

and S7). In addition, given that constitutively active Rac1 (Rac1(CA)) has been reported to

interfere with phagocytic cup closure and inhibit efferocytosis, Rac1(CA) can mitigate the effect

of tension of PS-PSRs on engulfment of PS-TGT^{56pN} beads. To test this, we overexpressed

Rac1(CA) in phagocytes and evaluated its effect. As expected, engulfment of PS-TGT^{56pN} by

phagocytes overexpressing Rac1(CA) was impeded in a manner dependent on the level of

Rac1(CA) expression (fig. S2, E and F). These data suggest that tension of PS-PSRs

orchestrates phagocytic cup closure via spatiotemporal inactivation of Rac1 during

efferocytosis.

Along with Rac1 inactivation, myosin II-mediated contraction is necessary for

phagocytic cup closure (*20*). To investigate whether the tension also affects myosin II activity,

we performed a frustrated phagocytosis assay that can quantitatively evaluate phagocytic cup

extension and contraction (*21*). Macrophages were incubated on PS-TGT-coated plates and

their extension and contraction were monitored. During the extension phase, cells spread

comparably on PS-TGT^{12pN} and PS-TGT^{56pN} plates. However, during the contraction phase,

phagocytes contracted more slowly on the PS-TGT^{12pN} plate than on the PS-TGT^{56pN} plate. In

addition, treatment of cells with blebbistatin, a myosin II inhibitor, drastically delayed the

contraction of phagocytes on the PS-TGT^{56pN} plate such that it was similar to that on the PS-

TGT^{12pN} plate (Fig. 2G and movies S8 to S10). Higher levels of myosin light chain (MLC)

phosphorylation were observed around PS-TGT^{56pN} beads and at the leading edge of cells on

the PS-TGT^{56pN} plate when macrophages were incubated with PS-TGT beads and on PS-TGT

plates, respectively (Fig. 2H, fig. S3, A to C, and movie S11). Overall, these data imply that

tension of PS-PSRs causes Rac1 inactivation at the phagocytic cup, leading to both myosin II-

mediated contraction and F-actin disassembly required for phagocytic cup closure.

Next, we investigated how tension of PS-PSRs modulates Rac1 activity.

Phosphoinositide 3-kinase (PI3K) recruits GTPase-activating proteins (GAPs) for Rac1 by

generating PIP$_3$ at the phagocytic cup during efferocytosis, leading to Rac1 inactivation and thus

phagocytic cup closure (22). Therefore, we reasoned that the tension can alter PI3K activity. To

test whether PI3K affects tension tolerance-dependent engulfment, macrophages were treated

with LY294002, a PI3K inhibitor, and engulfment of the beads was evaluated. The inhibitor

completely abrogated the superior engulfment of PS-TGT^{56pN} beads relative to PS-TGT^{12pN}

beads (Fig. 3A), suggesting that PI3K is involved in tension tolerance-dependent engulfment of

PS-TGT beads. To further understand the relationship between the tension and PI3K activity, we

used the GFP-tagged PH domain of Akt (PH-Akt-GFP) that binds to PIP$_3$, which represents

PI3K activity. When PH-Akt-GFP-expressing macrophages were incubated with PS-TGT beads,

the GFP signal was higher in phagocytic cups around PS-TGT^{56pN} beads than in phagocytic

cups around PS-TGT^{12pN} beads, and this difference was nullified in the presence of the PI3K

inhibitor (Fig. 3, B and C, and movies S12 to S14). This implies that tension of PS-PSRs

increases the activity of PI3K, leading to Rac1 inactivation in the phagocytic cup.

We then explored how the tension modulates PI3K activity. Given that PI3K associates

with some engulfment receptors (e.g., Tim-1 and Mertk) and all experiments in this study were

performed under serum-free conditions in which only direct PSRs could participate in

engulfment of the beads (*23-27*), we investigated whether PI3K associates with PSRs and, if so,

whether the association is affected by tension of PS-PSRs. PI3K robustly associated with Tim-4

among members of the Tim family and BAI1 (Fig. 3D and fig. 4, A and B). In addition, to confirm

the subcellular location of the association between Tim-4 and PI3K at the endogenous level, a

proximity ligation assay (PLA) was performed. As a result, in macrophages incubated with PS-

TGT^{12pN} beads, PLA signals were not confined to the phagocytic cup and were observed

ubiquitously. However, in macrophages incubated with PS-TGT^{56pN} beads, most signals were

observed around the phagocytic cup (Fig. 3, E and F). Collectively, these data imply that tension

of PS-PSRs increases the association of PI3K with PSRs, e.g., Tim-4, sequentially resulting in

Rac1 inactivation, F-actin disassembly, and myosin II-mediated contraction, which ultimately

causes phagocytic cup closure.

To investigate how tension is loaded on PS-PSRs, actin and myosin were examined

because they generate force in cells and are essential for phagocytic cup formation and closure

during efferocytosis (*20, 28*). In addition, F-actin disassembly and myosin II-mediated

contraction were modulated in a tension tolerance-dependent manner during engulfment of PS-

TGT beads (Fig. 2, A, B, G, and H). This suggests that tension of PS-PSRs at the phagocytic

cup during efferocytosis can be loaded by actin dynamics and/or myosin II-mediated

contraction. To test this possibility, cell detachment from PS-TGT plates was evaluated in the

presence or absence of cytochalasin D and blebbistatin, which are inhibitors of actin

polymerization and myosin II-mediated contraction, respectively. Cytochalasin D blocked

detachment of macrophages from the glass plate to which they were connected via PS-TGT^{12pN}

(Fig. 4, A and B) and loss of Cy5 signals in Cy5-labeled PS-TGT^{12pN}, but blebbistatin did not

(Fig. 4C and fig. S5A). These data imply that actin polymerization, rather than myosin II-

mediated contraction, is crucial for tension loading between PS and PSRs at the phagocytic

cup. In addition, the tension is not due to an allosteric effect of PSRs induced by PS binding

because cytochalasin D still allows PS to bind to its receptors (Fig. 4A).

Given that membrane tension increases during phagocytic cup formation driven by actin

polymerization (*29*), this membrane tension may play an important role in loading tension on

PS-PSRs. To validate this, we decreased the membrane tension of macrophages using

hypertonic solution (*30, 31*). Indeed, the more efficient engulfment of PS-TGT^{56pN} beads relative

to PS-TGT^{12pN} beads was abrogated by treating macrophages with hypertonic solution (Fig. 4D).

The treatment also reduced the Cy5 MFI of non-engulfing phagocytes upon incubation with PS-

TGT^{12pN} beads, but that of non-engulfing phagocytes upon incubation with PS-TGT^{56pN} beads

was unaltered (fig. S5B). Furthermore, treatment with hypertonic solution diminished the GFP

signal, representing PI3K activity, around PS-TGT^{56pN} beads in macrophages expressing PH-

Akt-GFP (Fig. 4, E and F, and movie S15). Consistently, when apoptotic cells were used instead

of PS-TGT beads to increase the physiological relevance, phagocytosis of apoptotic cells was

drastically impaired when macrophages were treated with hypertonic solution (Fig. 4G). Overall,

these data imply that an increase in the membrane tension of phagocytes is coupled with an

increase in tension or loading rate exerted on PS-PSRs during efferocytosis.

The membrane stiffness of apoptotic cells is increased (*32, 33*), which is expected to be

necessary for loading tension on PS-PSRs during efferocytosis. To apply tension of PS-PSRs,

PS on the surface of apoptotic cells must be pulled by PSRs on phagocytes. In this situation,

the membrane stiffness of apoptotic cells may be crucial to reach a certain level of tension

generated by the pulling of PSRs within the limited phagocytic synaptic space. To validate this,

we altered the membrane stiffness of apoptotic cells using cytochalasin D, which decreases

membrane stiffness (*33*), and performed an efferocytosis assay. Treatment of apoptotic cells

with cytochalasin D substantially impeded efferocytosis *in vitro* and *in vivo* (Fig. 4, H and I).

Notably, the conditioned medium of apoptotic cells treated with cytochalasin D did not affect

efferocytosis, demonstrating that the direct effect of cytochalasin D released from apoptotic cells

on phagocytes was minimal (fig. S5C). In addition, cytochalasin D did not alter the status of

apoptotic cells (fig. S5D). These data imply that increased membrane stiffness in apoptotic cells

also contributes to tension generation between PS and PSRs during efferocytosis.

PS is exposed not only on apoptotic cells but also on some live cells, but only apoptotic

cells are phagocytosed (*9, 34-36*). In general, this is thought to be due to don't eat-me signals

such as CD47 on live cells (*10*). In addition to these signals, our observations strongly suggest

that tension of PS-PSRs is another decisive signal for efferocytosis. To investigate this, we

generated PS-exposing Jurkat cells using A23187, a calcium ionophore that stimulates PS

exposure on the cell surface without inducing apoptosis (fig. S5, E and F), and artificially

increased the membrane stiffness of Jurkat cells by depleting cholesterol using methyl-β-

cyclodextrin (MβCD) (*37, 38*). Macrophages efficiently engulfed Jurkat cells treated with both

A23187 and MβCD, but engulfment of Jurkat cells treated with A23187 or MβCD alone was

minimally increased compared with that of control Jurkat cells (Fig. 4J). To further validate the

physiological relevance of these findings, we used platelets, which expose PS on their surface

when they are activated for blood coagulation (fig. S5G) (*39, 40*). Engulfment of activated

platelets by macrophages was indistinguishable from that of non-activated platelets, but

activated platelets treated with MβCD were efficiently engulfed (Fig. 4K). Taken together, these

data suggest that tension of PS-PSRs, mediated by both increased membrane tension at the

phagocytic cup in phagocytes and membrane stiffness in apoptotic cells, acts as a checkpoint to

determine whether a PS-exposing cell should be phagocytosed and this tension is thus

essential for completion of efferocytosis.

In sum, we report that efferocytosis, a type of phagocytosis, is a novel example of a

process mediated by chemo-mechanical pathways. Upon binding between PS and PSRs, Rac1

is activated, resulting in actin polymerization to form the phagocytic cup around apoptotic cells

and thus increasing membrane tension concomitant with tension of PS-PSRs at the phagocytic

cup. When the tension reaches a substantial level (e.g., 56 pN), PI3K is recruited to PSRs at the

phagocytic cup, which results in Rac1 inactivation, leading to F-actin disassembly and myosin II-

mediated contraction required for phagocytic cup closure. Ultimately, tension of PS-PSRs must

be generated to successfully complete phagocytosis of apoptotic cells (fig. S6A).

Our findings provide insights into how chemical and mechanical signals generated

during phagocytosis of PS-exposing cells are delicately coordinated to enhance the integrity of

efferocytosis without errors. More specifically, biochemical signaling of efferocytosis initiated by

binding of PS to PSRs is converted into a mechanical signal (i.e., tension of PS-PSRs). The

tension is eventually converted back into a biochemical signal that determines whether PS-

exposing cells being engulfed are completely phagocytosed (fig. S6B). By generating a

mechanical signal during phagocytosis of PS-exposing cells, phagocytes can avoid removing all

PS-exposing cells and precisely remove cells to be cleared, e.g., apoptotic cells. Indeed, many

live cells expose PS on their surfaces including some cancer cells (*34-36, 41-44*). Cancer

immunotherapy based on innate immunity has recently attracted much attention (*45*). Cancer

cells may evade phagocytosis by immune cells via nullifying this type of mechanical signal as a

checkpoint during phagocytosis. It would be interesting to investigate the correlation of cancer

cell phagocytosis with tension between a ligand on cancer cells and its corresponding receptor

on immune cells, which may provide insight to develop cancer immunotherapy.

281
282

References and Notes

1. S. Arandjelovic, K. S. Ravichandran, Phagocytosis of apoptotic cells in homeostasis. *Nat Immunol* **16**, 907-917 (2015).

2. A. C. Doran, A. Yurdagul, Jr., I. Tabas, Efferocytosis in health and disease. *Nat Rev Immunol* **20**, 254-267 (2020).

3. V. A. Fadok, D. L. Bratton, S. C. Frasch, M. L. Warner, P. M. Henson, The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell Death Differ* **5**, 551-562 (1998).

4. V. A. Fadok *et al.*, Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J Immunol* **148**, 2207-2216 (1992).

5. P. A. Oldenborg *et al.*, Role of CD47 as a marker of self on red blood cells. *Science* **288**, 2051-2054 (2000).

6. A. A. Barkal *et al.*, CD24 signalling through macrophage Siglec-10 is a target for cancer immunotherapy. *Nature* **572**, 392-396 (2019).

7. S. Brown *et al.*, Apoptosis disables CD31-mediated cell detachment from phagocytes promoting binding and engulfment. *Nature* **418**, 200-203 (2002).

8. A. A. Barkal *et al.*, Engagement of MHC class I by the inhibitory receptor LILRB1 suppresses macrophages and is a target of cancer immunotherapy. *Nat Immunol* **19**, 76-84 (2018).

9. K. Segawa, J. Suzuki, S. Nagata, Constitutive exposure of phosphatidylserine on viable cells. *Proc Natl Acad Sci U S A* **108**, 19246-19251 (2011).

10. S. Jaiswal *et al.*, CD47 is upregulated on circulating hematopoietic stem cells and leukemia cells to avoid phagocytosis. *Cell* **138**, 271-285 (2009).

11. J. M. Kinchen, K. S. Ravichandran, Journey to the grave: signaling events regulating

308   removal of apoptotic cells. *J Cell Sci* **120**, 2143-2149 (2007).

309 12. R. K. Tsai, D. E. Discher, Inhibition of "self" engulfment through deactivation of myosin-II

310   at the phagocytic synapse between human cells. *J Cell Biol* **180**, 989-1003 (2008).

311 13. S. Mylvaganam, S. A. Freeman, S. Grinstein, The cytoskeleton in phagocytosis and

312   macropinocytosis. *Curr Biol* **31**, R619-R632 (2021).

313 14. X. Wang, T. Ha, Defining single molecular forces required to activate integrin and notch

314   signaling. *Science* **340**, 991-994 (2013).

315 15. R. Roy, S. Hohng, T. Ha, A practical guide to single-molecule FRET. *Nat Methods* **5**, 507-

316   516 (2008).

317 16. M. Nakaya, M. Kitano, M. Matsuda, S. Nagata, Spatiotemporal activation of Rac1 for

318   engulfment of apoptotic cells. *Proc Natl Acad Sci U S A* **105**, 9198-9203 (2008).

319 17. H. Moon *et al.*, Crbn modulates calcium influx by regulating Orai1 during efferocytosis.

320   *Nat Commun* **11**, 5489 (2020).

321 18. M. R. Elliott, K. M. Koster, P. S. Murphy, Efferocytosis Signaling in the Regulation of

322   Macrophage Inflammatory Responses. *J Immunol* **198**, 1387-1394 (2017).

323 19. R. E. Itoh *et al.*, Activation of rac and cdc42 video imaged by fluorescent resonance

324   energy transfer-based single-molecule probes in the membrane of living cells. *Mol Cell*

325   *Biol* **22**, 6582-6591 (2002).

326 20. Y. Ikeda *et al.*, Rac1 switching at the right time and location is essential for Fcgamma

327   receptor-mediated phagosome formation. *J Cell Sci* **130**, 2530-2540 (2017).

328 21. D. T. Kovari *et al.*, Frustrated Phagocytic Spreading of J774A-1 Macrophages Ends in

329   Myosin II-Dependent Contraction. *Biophys J* **111**, 2698-2710 (2016).

330 22. D. Schlam *et al.*, Phosphoinositide 3-kinase enables phagocytosis of large particles by

331   terminating actin assembly through Rac/Cdc42 GTPase-activating proteins. *Nat*

332   *Commun* **6**, 8623 (2015).

23.   P. Sen *et al.*, Apoptotic cells induce Mer tyrosine kinase-dependent blockade of NF-

kappaB activation in dendritic cells. *Blood* **109**, 653-660 (2007).

24.   A. J. de Souza *et al.*, T cell Ig and mucin domain-1-mediated T cell activation requires

recruitment and activation of phosphoinositide 3-kinase. *J Immunol* **180**, 6518-6526

(2008).

25.   G. X. Ruan, A. Kazlauskas, Axl is essential for VEGF-A-dependent activation of

PI3K/Akt. *EMBO J* **31**, 1692-1703 (2012).

26.   M. Miyanishi *et al.*, Identification of Tim4 as a phosphatidylserine receptor. *Nature* **450**,

435-439 (2007).

27.   D. Park *et al.*, BAI1 is an engulfment receptor for apoptotic cells upstream of the

ELMO/Dock180/Rac module. *Nature* **450**, 430-434 (2007).

28.   K. Carvalho *et al.*, Actin polymerization or myosin contraction: two ways to build up

cortical tension for symmetry breaking. *Philos Trans R Soc Lond B Biol Sci* **368**,

20130005 (2013).

29.   V. Jaumouille, C. M. Waterman, Physical Constraints and Forces Involved in

Phagocytosis. *Front Immunol* **11**, 1097 (2020).

30.   A. R. Houk *et al.*, Membrane tension maintains cell polarity by confining signals to the

leading edge during neutrophil migration. *Cell* **148**, 175-188 (2012).

31.   T. A. Masters, B. Pontes, V. Viasnoff, Y. Li, N. C. Gauthier, Plasma membrane tension

orchestrates membrane trafficking, cytoskeletal remodeling, and biochemical signaling

during phagocytosis. *Proc Natl Acad Sci U S A* **110**, 11875-11880 (2013).

32.   N. I. Nikolaev, T. Muller, D. J. Williams, Y. Liu, Changes in the stiffness of human

mesenchymal stem cells with the progress of cell death as measured by atomic force

microscopy. *J Biomech* **47**, 625-630 (2014).

33.   W. A. Lam, M. J. Rosenbluth, D. A. Fletcher, Chemotherapy exposure increases

leukemia cell stiffness. *Blood* **109**, 3505-3508 (2007).

34.   J. I. Elliott *et al.*, Membrane phosphatidylserine distribution as a non-apoptotic signalling

mechanism in lymphocytes. *Nat Cell Biol* **7**, 808-816 (2005).

35.   S. Riedl *et al.*, In search of a novel target - phosphatidylserine exposed by non-apoptotic

tumor cells and metastases of malignancies with poor treatment efficacy. *Biochim*

Biophys Acta **1808**, 2638-2645 (2011).

36.   S. M. Qadri, R. Bissinger, Z. Solh, P. A. Oldenborg, Eryptosis in health and disease: A

paradigm shift towards understanding the (patho)physiological implications of

programmed cell death of erythrocytes. *Blood Rev* **31**, 349-361 (2017).

37.   J. Suzuki, M. Umeda, P. J. Sims, S. Nagata, Calcium-dependent phospholipid

scrambling by TMEM16F. *Nature* **468**, 834-838 (2010).

38.   A. Biswas, P. Kashyap, S. Datta, T. Sengupta, B. Sinha, Cholesterol Depletion by

MbetaCD Enhances Cell Membrane Tension and Its Variations-Reducing Integrity.

Biophys J **116**, 1456-1468 (2019).

39.   S. M. Schoenwaelder *et al.*, Two distinct pathways regulate platelet phosphatidylserine

exposure and procoagulant function. *Blood* **114**, 663-666 (2009).

40.   B. R. Lentz, Exposure of platelet membrane phosphatidylserine regulates blood

coagulation. *Prog Lipid Res* **42**, 423-438 (2003).

41.   K. Fischer *et al.*, Antigen recognition induces phosphatidylserine exposure on the cell

surface of human CD8+ T cells. *Blood* **108**, 4094-4101 (2006).

42.   J. I. Elliott *et al.*, Phosphatidylserine exposure in B lymphocytes: a role for lipid packing.

Blood **108**, 1611-1617 (2006).

43.   K. J. de Vries, T. Wiedmer, P. J. Sims, B. M. Gadella, Caspase-independent exposure of

aminophospholipids and tyrosine phosphorylation in bicarbonate responsive human

sperm cells. *Biol Reprod* **68**, 2122-2134 (2003).

44.   J. H. Stafford, P. E. Thorpe, Increased exposure of phosphatidylethanolamine on the

384   surface of tumor vascular endothelium. *Neoplasia* **13**, 299-308 (2011).

385   45.   M. Feng *et al.*, Phagocytosis checkpoints as new targets for cancer immunotherapy. *Nat*

386   *Rev Cancer* **19**, 568-586 (2019).

Acknowledgments

We thank Michiyuki Matsuda (Kyoto University) and Jihye Seong (Korea Institute of Science

and Technology) for kindly providing the plasmids of Raichu-Rac1 and PH-Akt-GFP,

respectively.

Funding

This work was supported by the National Research Foundation of Korea funded by the Korea

government (MSIP) (2022R1A2C1008334 and 2022R1A4A2000790) and by GIST research

Institute (GRI) IIBR.

Author contributions

Conceptualization, G.L. and D.P.; Methodology, C.M., H.C, J.Y., B.M., J.J., N.S., H.M., S.Y., J.L.,

and M.H.; Formal analysis, C.M., H.C, J.Y., P.J., G.L., and D.P.; Resources, J.J. and W.-J.C.;

Writing, C.M., H.C, J.Y., G.L., and D.P.; Funding acquisition, G.L. and D.P.

Competing interests

The authors declare no competing interests.

Data and materials availability

Data from this study are available within the main text and the supplementary materials or from

the corresponding authors (G.L. and D.P.) upon reasonable request.

Supplementary Materials

Materials and Methods

Supplementary Text (Uncropped blots of western blot panels, Gating strategies used for flow

cytometry analysis, Movie legends, Synthesis of 18:1-12:0(maleimide) phosphatidylcholine)

Figs. S1 to S6

Movies S1 to S15

Figure 1

Fig. 1. Tension-dependent engulfment of PS-TGT beads.

(**A**) Schematic diagram of PS-TGT beads. PS and biotin were conjugated to the Cy3- or Cy5-labeled upper strand and the lower strand of 21 bp DNA, respectively, which was immobilized on neutravidin-coated polystyrene beads via a neutravidin-biotin linker. Yellow arrows indicate the direction of force (F). (**B**) J774A.1 cells incubated with Cy3-labeled PS-TGT^{12pN} or PS-TGT^{56pN} beads for 30 min were observed by time-lapse confocal microscopy (left) and engulfed beads were counted (right). Asterisks indicate ingested beads. Scale bar, 10 μm. Data are mean ± s.e.m. from n=165 cells for PS-TGT^{12pN} and n=124 cells for PS-TGT^{56pN}. ***$P < 0.001$; two-tailed unpaired Student's t test. (**C** and **D**) J774A.1 cells were incubated with the indicated beads for 10 min and analyzed by flow cytometry. Ctrl, control; NA, neutravidin bead. Data are mean ± s.e.m. from n=5 (C) and n=3 (D) independent experiments. *$P < 0.05$; ***$P < 0.001$; NS, not significant; one-way ANOVA. (**E**) J774A.1 cells incubated with the indicated beads for 10 min were analyzed by flow cytometry. Representative histograms of phagocytes (left) and MFIs of non-engulfing phagocytes (right) are shown. The box and dotted lines in the histograms indicate non-engulfing phagocytes and the peak of the fluorescence intensity of non-engulfing phagocytes, respectively. Data are mean ± s.e.m. from n=5 independent experiments. ***$P < 0.001$; NS, not significant; one-way ANOVA. (**F**) J774A.1 cells were incubated on PS-TGT plates, on which the indicated PS-TGTs were immobilized, for 5 min, washed, further incubated for 60 min, and washed to remove detached cells. Scale bar, 200 μm. Data are mean ± s.e.m. from n=5 independent experiments. ***$P < 0.001$; NS, not significant; two-way ANOVA. (**G**) J774A.1 cells stably expressing membrane-targeted GFP (GFP-CAAX) were incubated on PS-TGT plates, on which a mixture of the indicated Cy5-labeled PS-TGTs and Cy3-labeled PS-TGT^{56pN} was immobilized, for 60 min and observed by confocal microscopy. Relative Cy5 intensities were calculated. The intensity of Cy5 in the area where cells were absent was set to 0. Scale bar, 10 μm. Data are represented in the box-and-whisker plot. n=26 cells for Ctrl

TGT^{12pN}, n=30 cells for Ctrl TGT^{56pN}, n=29 cells for PS-TGT^{12pN}, and n=49 cells for PS-TGT^{56pN}.

***$P < 0.001$; NS, not significant; one-way ANOVA. (**H**) Schematic diagram of normal PS-

TGT^{12pN} (top) and hairpin PS-TGT^{12pN} (bottom). The upper and lower strands of hairpin PS-

TGT^{12pN} were connected by PEG or DNA loops. Yellow arrows indicate the direction of force (F).

(**I** and **J**) J774A.1 cells were incubated with the indicated beads for 15 min. Percentage of

engulfing phagocytes were measured by flow cytometry. Data are mean ± s.e.m. from n=3

independent experiments for (I), n=4 independent experiments for (J). *$P < 0.05$; NS, not

significant; two-way ANOVA.

Figure 2

Fig. 2. The tension affects engulfment of PS-TGT beads through Rac1.

(**A** and **B**) J774A.1 cells stained with SiR-actin were incubated with Cy3-labeled PS-TGT beads for 30 min and observed by time-lapse confocal microscopy (A). The duration of F-actin labeling was measured (B). Asterisks indicate PS-TGT beads with SiR-actin signals. Scale bar, 10 μm. Data are mean ± s.e.m. from n=29 beads for PS-TGT^{12pN} and n=62 beads for PS-TGT^{56pN}. ***P < 0.001; two-tailed unpaired Student's *t* test. (**C** and **D**) J774A.1 cells stably expressing membrane-targeted GFP were incubated with Cy3-labeled PS-TGT beads for 30 min and observed by time-lapse confocal microscopy (C). Asterisks indicate beads with extended phagocytic cups. The amount of time required for phagocytic cup closure was measured (D). Scale bar, 10 μm. Data are mean ± s.e.m. from n=36 beads for PS-TGT^{12pN} and n=54 beads for PS-TGT^{56pN}. ***P < 0.001; two-tailed unpaired Student's *t* test. (**E** and **F**) J774A.1 cells stably expressing Raichu-Rac1 were incubated with PS-TGT beads for 30 min and observed by time-lapse confocal microscopy (E). Asterisks indicate beads with extended phagocytic cups. Rac1 activity in phagocytic cups around the beads was calculated by YFP/CFP ratio FRET. Scale bar, 5 μm. Schematic diagram depicting the analysis of Rac1 FRET according to the phagocytic cup position (F, left). Rac1 activity was measured at the phagocytic cup around the beads (F, right). n=33 beads for PS-TGT^{12pN} and n=23 beads for PS-TGT^{56pN}. (**G**) J774A.1 cells stably expressing membrane-targeted GFP were incubated on the indicated PS-TGT plates with or without blebbistatin (10 μM) and observed by time-lapse confocal microscopy. When the cells were maximally expanded, the time and relative area were set to 0 min and 1, respectively. The areas of cells on the plates were normalized to the maximum area. Scale bar, 10 μm. Blebb, blebbistatin. Data are mean ± s.e.m. from n=8 cells for PS-TGT^{12pN} (DMSO), n=11 cells for PS-TGT^{56pN} (DMSO), n=16 cells for PS-TGT^{12pN} (Blebb.), and n=8 cells for PS-TGT^{56pN} (Blebb.). Statistical analysis was performed between the end points of PS-TGT^{12pN} (DMSO) and PS-TGT^{56pN} (DMSO). *P < 0.05; one-way ANOVA. (**H**) J774A.1 cells were incubated with both Cy5-

labeled PS-TGT^{12pN} and unlabeled PS-TGT^{56pN} beads for 10 min, stained with an anti-phospho-

MLC antibody and Alexa Fluor 594-conjugated phalloidin, and observed by confocal

microscopy. Magenta and white arrows indicate PS-TGT^{12pN} and PS-TGT^{56pN} beads,

respectively. The intensities of phosphorylated MLC (pMLC) staining were quantified across the

phagocytic cups. Scale bar, 10 µm. Data are mean ± s.e.m. from n=26 beads for PS-TGT^{12pN}

and PS-TGT^{56pN}.

Figure 3

Fig. 3. The tension modulates PI3K activity.

(**A**) J774A.1 cells pre-incubated with LY294 (LY294002, 50 µM) for 10 min were incubated with

the indicated Cy3-labeled beads and analyzed by flow cytometry. Data are mean ± s.e.m. from

n=4 independent experiments. ***P < 0.001; NS, not significant; two-way ANOVA. (**B** and **C**)

J774A.1 cells stably expressing PH-Akt-GFP were incubated with Cy3-labeled PS-TGT beads

for 30 min and observed by time-lapse confocal microscopy. Asterisks indicate beads with

extended phagocytic cups (B). The intensities of GFP at phagocytic cups were measured (C).

Scale bar, 10 µm. n=67 beads for PS-TGT^{12pN} (DMSO), n=41 beads for PS-TGT^{56pN} (DMSO),

n=70 beads for PS-TGT^{12pN} (LY294002), and n=54 beads for PS-TGT^{56pN} (LY294002). (**D**)

Lysates of 293T cells transfected with the indicated plasmids were incubated with anti-HA

antibody-conjugated agarose beads. Bead-bound proteins were detected by the indicated

antibodies. IP, immunoprecipitation; TCL, total cell lysate. (**E** and **F**) Peritoneal macrophages

were incubated with Cy3-labeled PS-TGT beads for 10 min, fixed, blocked, and incubated with

anti-Tim4 and anti-PI3K antibodies. Then, PLA probes were stained and incubated in

amplification solution. PLA signals were observed using confocal microscopy (E), and PLA

signals for each bead were quantified (F). Asterisks indicate beads with extended phagocytic

cups. Scale bar, 10 µm. Data are mean ± s.e.m. from n=126 beads for PS-TGT^{12pN} and n=153

beads for PS-TGT^{56pN}. ***P < 0.001; two-tailed unpaired Student's *t* test.

Figure 4

Fig. 4. Membrane tension increased by actin polymerization loads tension on PS-PSRs.

(**A**) J774A.1 cells incubated on the indicated PS-TGT plates for 5 min were washed and then further incubated in the absence or presence of cytochalasin D (1 µM) or blebbistatin (10 µM) for 60 min. The cell number at 60 min relative to that at 5 min was calculated. Representative images after incubation for 60 min are shown. Scale bar, 200 µm. Cyto.D, cytochalasin D; Blebb, blebbistatin. Data are mean ± s.e.m. from n=3 independent experiments. ***P < 0.001; NS, not significant; two-way ANOVA. (**B** and **C**) J774A.1 cells stably expressing membrane-targeted GFP were incubated on the indicated PS-TGT plates for 60 min and Cy5 fluorescence was observed by confocal microscopy (B). Relative Cy5 intensities were calculated. The intensity of Cy5 in the area where cells were absent was set to 0 (C). Data are presented in the box-and-whisker plot. n=55 cells for PS-TGT^{12pN} (DMSO), n=172 cells for PS-TGT^{12pN} (Cyto.D), n=115 cells for PS-TGT^{12pN} (Blebb.), n=92 cells for PS-TGT^{56pN} (DMSO), n=189 cells for PS-TGT^{56pN} (Cyto.D), and n=161 cells for PS-TGT^{56pN} (Blebb.). ***P < 0.001; NS, not significant; two-way ANOVA. (**D**) J774A.1 cells pre-incubated in medium containing 150 mM sucrose (hypertonic medium) were incubated with the indicated beads for 10 min and analyzed by flow cytometry. Data are mean ± s.e.m. from n=3 independent experiments. ***P < 0.001; NS, not significant; two-way ANOVA. (**E** and **F**) J774A.1 cells stably expressing PH-Akt-GFP were pre-incubated in medium containing 150 mM sucrose and then incubated with Cy3-labeled PS-TGT^{56pN} beads and observed by time-lapse confocal microscopy. Asterisks indicate beads with extended phagocytic cups (E). The intensities of GFP fluorescence at phagocytic cups around the beads were measured (F). Scale bar, 10 µm. n=40 beads for PS-TGT^{56pN} (control medium) and n=51 beads for PS-TGT^{56pN} (hypertonic medium). (**G**) J774A.1 cells pre-incubated in medium containing 150 mM sucrose were incubated with apoptotic Jurkat cells for 30 min and analyzed by flow cytometry. Data are mean ± s.e.m. from n=3 independent experiments. ***P < 0.001; two-tailed unpaired Student's *t* test. (**H**) J774A.1 cells were incubated with live or

apoptotic Jurkat cells treated with or without cytochalasin D (2 μM) for 30 min and analyzed by

flow cytometry. Data are mean ± s.e.m. from n=4 independent experiments. ***$P < 0.001$; NS,

not significant; one-way ANOVA. (**I**) TAMRA-stained apoptotic Jurkat cells treated with or without

cytochalasin D (2 μM) were intraperitoneally injected into 8-week-old mice. Peritoneal exudates

from the mice were stained with an anti-F4/80 antibody and analyzed by flow cytometry. Data

are mean ± s.e.m. from n=7 mice for Apop. and n=8 mice for Apop. (+Cyto.D). ***$P < 0.001$; two-

tailed unpaired Student's *t* test. (**J**) J774A.1 cells were incubated with Jurkat cells treated with

MβCD (1 mM), A23187 (10 μM), or MβCD and A23187 for 30 min and analyzed by flow

cytometry. Data are mean ± s.e.m. from n=3 independent experiments. ***$P < 0.001$; one-way

ANOVA. (**K**) J774A.1 cells were incubated with platelets treated with MβCD (10 mM) or

activated platelets treated with MβCD for 30 min and analyzed by flow cytometry. Data are

mean ± s.e.m. from n=6 independent experiments. *$P < 0.05$, **$P < 0.01$; one-way ANOVA.