NMDA receptor antagonist memantine selectively affects recurrent processing during perceptual inference

Samuel Noorman[1,2*], Timo Stein[1,2], Jasper Zantvoord[3,4],

Johannes J. Fahrenfort[1,2,5**], Simon van Gaal[1,2**]

[1]Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands

[2]Amsterdam Brain and Cognition, University of Amsterdam, Amsterdam, The Netherlands

[3]Amsterdam UMC location University of Amsterdam, Department of Psychiatry, Amsterdam, The Netherlands

[4]Amsterdam Neuroscience, Amsterdam, The Netherlands

[5]Department of Applied and Experimental Psychology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[*]Correspondence: sgnoorman@gmail.com

[**]Shared senior authorship

This research was supported by a grant from the H2020 European Research Council (ERC STG 715605, SVG).

Abstract

Perceptual inference requires the integration of visual features through recurrent processing, the dynamic exchange of information between higher and lower level cortical regions. While animal research has demonstrated a crucial role of NMDA receptors in recurrent processing, establishing a causal link between NMDA-mediated recurrent processing and human perception has remained challenging. Here, we report two pharmacological studies with randomized, double-blind, crossover designs in which we administered the NMDA antagonist memantine, while collecting human electroencephalography (EEG). We trained and tested EEG classifiers to reflect the processing of specific stimulus features with increasing levels of complexity, namely differences in stimulus contrast, collinearity between local line elements, and illusory surfaces of a Kanizsa triangle. In two experiments involving different participants and visual tasks, we found that memantine selectively affected decoding of the Kanizsa illusion, known to depend on recurrent processing, while leaving decoding of contrast and collinearity largely unaffected. Interestingly, the results from an attentional blink (experiment 1) and task-relevance manipulation (experiment 2) showed that memantine was only effective when the stimulus was attended and consciously accessed. These findings demonstrate that NMDA inhibition selectively affects recurrent processing, especially for attended objects, and thereby provide a crucial step toward bridging animal and human research, shedding light on the neural mechanisms underpinning perceptual inference and conscious perception.

Introduction

Different theories of conscious perception converge on a central principle: conscious perception entails integration of information through recurrent neural processes that exceeds the capabilities of mere feedforward processing (Mashour et al., 2020; Seth & Bayne, 2022). Recurrent neural mechanisms encompass feedback connections between higher- and lower-order cortical areas as well as lateral connections within sensory areas. Studies in mice and monkeys have revealed a causal role of feedback from higher cortical areas to sensory areas for tactile perception (Manita et al., 2015; Sachidhanandam et al., 2013; Takahashi et al., 2016), detection of odors (Allen et al., 2017), and visual stimulus attributes (Ciceri et al., 2023; Lamme et al., 1998; van Vugt et al., 2018; Zatka-Haas et al., 2021). In macaques, general anesthetics targeted at reducing the overall level of consciousness of the animal reduce global cortical recurrent interactions (Mashour & Hudetz, 2018) and block feedback connections to V1, while sparing feedforward processing (Lamme et al., 1998). Although in humans indirect evidence for a role of recurrent processing in conscious perception and perceptual integration has been provided, for example by focusing on neural markers of figure-ground segregation or backward masking (Fahrenfort et al., 2007, 2008, 2012, 2017; Kok et al., 2016; Kok & de Lange, 2014; Noorman et al., 2023; Scholte et al., 2008; Wyatte et al., 2014), the underlying biochemistry has remained so far elusive. In this study, we aimed to provide direct evidence for a causal role of NMDA receptors in recurrent processing (through lateral and feedback connections) and their role in perceptual inference and conscious access, factoring in potential modulating influences of attention (Flohr et al., 1998; Lamme, 2010; Sachidhanandam et al., 2013; Self et al., 2012; Thiele, 2012; Thiele & Bellgrove, 2018; Van Kerkoerle et al., 2014; Wang, 2008) .

NMDA has a slower time constant than most other receptors (decay time constant ~100 ms, voltage-gated) and is known to be densely populated in recurrent connections in the cortex (Wang, 2008). In macaque monkeys, blocking NMDA receptors selectively disrupted late onset modulations of neural activity in V1 (i.e., recurrent processing) during figure-ground detection, whereas blocking AMPA receptors did not, but instead reduced the strength of the first feedforward sweep (Self et al., 2012; Van Kerkoerle et al., 2014). This suggests that feedback connections target neuronal synapses that are rich in NMDA receptors (among other receptors), whereas feedforward connections target synapses especially rich in AMPA receptors (Thiele, 2012; Thiele & Bellgrove, 2018). In mice, blocking NMDA receptors strongly reduced hit rates for the detection of whisker movements (Sachidhanandam et al., 2013). In humans, few studies have investigated the role of NMDA receptors in recurrent

processing and (conscious) perception. Blocking NMDA receptors via administration of ketamine resulted in impaired feature integration in a texture discrimination task (Meuwese et al., 2013) and a distortion of the neural representations of objects in early visual cortex (van Loon et al., 2016). However, as these human studies did not clearly distinguish between markers of feedforward and recurrent processing, they could not determine whether these effects were specific to recurrent processing or reflected other non-specific effects. Ketamine has rather broad adverse side effects, such as sedation and dissociation, which could potentially explain the observed behavioral deficits and neural distortions. Also, ketamine is known for its complex pharmacological dynamics, having effects on several neurotransmitter systems, including catecholaminergic, cholinergic, nicotinic, muscarinic and GABAergic receptors (Kocsis et al., 2013; Mathew et al., 2012; Müller et al., 2011).

In this study, we used a novel Kanizsa stimulus whose different visual features were designed to target distinct processing stages with increasing levels of complexity, namely local contrast differences, differences in collinearity of local line elements and the presence/absence of an illusory triangle (**Fig. 1A**) (see also Noorman et al., 2023). The processing of collinearity of line elements predominantly relies on lateral connections (Bosking et al., 1997; Gilbert & Wiesel, 1979; Li, 1998; Liang et al., 2017; Schmidt et al., 1997; Stettler et al., 2002), while processing of the illusory surface of the Kanizsa illusion relies on feedback connections (Halgren et al., 2003; Kok et al., 2016; Kok & de Lange, 2014; Lee & Nguyen, 2001; Pak et al., 2020; Wokke et al., 2013). Therefore, multivariate EEG decoding of the presence/absence of the illusionary Kanizsa triangle served as a marker of feedback processing (Fahrenfort et al., 2017; Noorman et al., 2023). The ancillary features of the stimulus, the two-legged white circles, could be aligned to form a non-illusionary triangle, enabling us to use collinearity decoding as a marker of lateral processing. By rotating the Kanizsa by 180 degrees, thus either having one black Pacman on top of the screen or two, we created differences in contrast between the upper and lower visual field. Because local contrast differences are processed early in the visual system, this allowed us to employ local contrast decoding as a marker of feedforward processing. In recent work we have validated that the decoding profiles of these features of different complexities can indeed serve as EEG markers of feedforward, lateral and recurrent processes (Fahrenfort et al., 2017; Noorman et al., 2023).

We tested how these neural markers were affected by memantine, a low-affinity, voltage-dependent, uncompetitive NMDA antagonist. Memantine is well tolerated and has minimal side effects compared to ketamine. We performed two separate experiments. In experiment 1, we perturbed conscious access through masking and inattention using a modified attentional-blink (AB) paradigm (**Fig. 1B**). In experiment 2, we manipulated whether the illusion, collinearity, or local contrast were task-relevant or not. We hypothesized that blocking NMDA receptors by memantine would selectively affect illusion and possibly collinearity decoding, as putative markers of recurrent and lateral processing, but would spare the decoding of local contrast differences, our marker of feedforward processing.

Figure 1. Experiment 1: design, behavior, and training data. (A) Target stimulus set and schematic of the markers for the different types of processing (showing example stimuli). (B) Trial design. (C) Effects of memantine on subjective state assessed with the visual analogue scale. The categories alertness, contentedness, and calmness are based on Bond and Lader (1974), sedation on Danion et al. (1989). For details see Methods. (D) Participants' ability to detect the Kanizsa illusion in the second target stimulus (T2). (E) Tenfold cross-validation decoding scheme applied to an independent training dataset. Mean decoding performance, area under the receiver operating characteristic curve (AUC), over time ± standard error of the mean (SEM) is shown. Thick lines differ from chance: P<0.05, cluster-based permutation test. Local contrast, collinearity, and illusion decoding accuracy peaked at 86, 242, and 273 ms after stimulus onset and these peaks were encompassed by a 78-94, 192-292, and 223-323 ms time window, respectively. Error bars are mean ± SEM. Individual data points are plotted using low contrast. Ns is not significant (*P*>0.05). **P=0.005.

Results

Memantine caused minimal side effects

The study had a randomized, double-blind, crossover design in which a placebo was administered during one session and memantine (20 mg) during another. Drug order was counterbalanced across the 29 healthy participants. Potential side effects of memantine on different aspects of the participant's subjective state were assessed with the visual analog scale (VAS) (Bond & Lader, 1974; Danion et al., 1989). The only significant effect was a moderate decrease in the category "calmness" for memantine compared to placebo (t_{28}=-23.07, *P*=0.005, BF_{10}=8.71; **Fig. 1C**, for details see Methods). Note that in experiment 2 (**Fig. 4**) none of these subjective measures differed between memantine and placebo, indicating that overall subjective side effects of memantine were minimal.

Memantine selectively affects decoding of the Kanizsa illusion

We recorded EEG signals from participants who detected illusory surfaces in two black targets (T1 and T2) amongst red distractors in a rapid serial visual presentation (RSVP) task (**Fig. 1B**). Visibility of T2 was manipulated in two ways, resulting in a 2×2 factorial task design. First, T2 was either masked or unmasked. Second, shortening the interval (lag) between the two targets from 900 to 200/300 ms induced an attentional blink (AB), which reduced T2 visibility as well. At the end of a trial, participants indicated for each target whether they had perceived an illusory surface or not. For all main analyses of behavior and EEG decoding we conducted repeated-measures ANOVAs with the factors drug (memantine vs. placebo), T1-T2 lag (short, inducing an AB, vs. long, no AB), and masks (masked vs. unmasked). The complete results of these ANOVAs are presented in the corresponding results figures. In the text, we only highlight the key results and report selectively additional statistics such as post-hoc tests.

As expected, both masking and the AB impaired participant's ability to detect the Kanizsa illusion in the second target (T2). Our masking procedure was highly effective, leading to poor T2 performance on masked trials (red lines in **Fig. 1D**). The AB was also effective, leading to a sharp drop in T2 performance on short lag versus long lag trials (green lines in **Fig. 1D**). Memantine had however no effect on perceptual sensitivity (**Fig. 1D**).

To establish our markers for the different neural processes of interest, the processing of different visual features (local contrast, collinearity, and the Kanizsa illusion) were decoded (**Fig. 1A**). For these decoding analyses, classifiers were trained using an independent training set obtained from a separate experimental task where the same Kanizsa target stimulus was presented, but outside the context of the RSVP task (see Methods for details). In brief, participants had to match the target stimulus to different example stimuli ("templates"). For successful performance in this template-matching task, all three features of the stimulus had to be attended, and all features were thus task-relevant. We also used the independent training set from this template-matching task to determine the time windows for local contrast, collinearity, and illusion decoding. First, we decoded each visual feature from the training set using a tenfold cross-validation scheme by training linear discriminant classifiers to categorize the feature as either present or absent based on the raw EEG activity across all electrodes. After testing the classifiers, each decoded visual feature showed a clear peak in classifier performance (area under the receiver operating characteristic curve [AUC]) (Hand & Till, 2001) over time: local contrast at 86 ms after stimulus onset, collinearity at 242 ms, and the illusion at 273 ms. In the following, we refer to "time window-based analyses" when we focus on the averages of the time windows encompassing these peaks (for the exact time windows used see **Fig. 1E**). The timing and mostly occipital nature (see the covariance/class separability maps of **Fig. 2A**) (Haufe et al., 2014) of these windows were consistent with the neural processes they are thought to reflect and in line with our previous EEG decoding studies (Fahrenfort et al., 2017; Noorman et al., 2023). Because of this topology, subsequent decoding analyses were restricted to occipitoparietal electrodes. To increase power, we averaged each time window in the localizer task's raw EEG data. Classifiers were trained on these averages and tested on all time samples of each condition of the main RSVP task's data (see Methods) (King & Dehaene, 2014). Note that this approach therefore reflects "off-diagonal decoding" in the generalization across time approach. We show all "on-diagonal" analyses in the **supplementary figures S2-S4**, which revealed very similar results.

We first examined the effect of memantine on the decoding of the different features of the first target (T1) from the RSVP task. For each time sample, we tested whether the memantine conditions differed in classifier performance from the placebo conditions. Cluster-based permutation testing was used to correct for multiple comparisons (Maris & Oostenveld, 2007). Crucially, only illusion decoding showed a significant cluster ranging from 211 to 352 ms after stimulus onset (**Fig. 2A**, blue lines). A similar result was obtained from our time window-based analysis (**Fig. 2B**, ANOVA results in the figure). Memantine significantly improved overall T1 decoding ($F_{1,28}$=8.33, P=0.007) and drug interacted with visual feature ($F_{1,28}$=5.16, P=0.011). Post hoc t-tests confirmed that memantine

selectively affected illusion decoding accuracy (t_{28}=-4.14, P<0.001, BF_{10}=101.92), thus leaving local
contrast (t_{28}=-0.60, P=0.560, BF_{10}=0.23) and collinearity decoding unaffected (t_{28}=-1.04, P=0.306,
BF_{10}=0.32). As memantine is an NMDA blocker, we had initially hypothesized that it would impair
recurrent processing, and therefore decrease illusion decoding accuracy. Surprisingly however,
although memantine specifically affected illusion decoding, it *increased* instead of decreased illusion
decoding. Recent studies using memantine have however mostly observed increases in task
performance and/or neural markers of sensory processing (Korostenskaja et al., 2007; Light et al.,
2017; Swerdlow et al., 2016, 2020), and are therefore mostly in line with our observations here. To
foreshadow our further results, this effect was robust as we obtained the same *increase* in Kanizsa
decoding for T2's in this study (**Fig. 3**), and the effect was replicated in study 2 (**Fig. 5**). We address this
issue further in the discussion.

Figure 2. Decoding applied to the T1 data. (A) Local contrast, collinearity, and Kanizsa illusion
decoding. These three visual features' time windows: 78-94, 192-292, and 223-323 ms, respectively.
Covariance/class separability maps with arbitrary units (AU) reflecting underlying neural sources are
shown. Below these maps: mean decoding performance, area under the receiver operating
characteristic curve (AUC), over time ± standard error of the mean (SEM). Thick lines differ from
chance: P<0.05, cluster-based permutation test. (B) Mean AUC for every time window. Error bars are
mean ± SEM. Individual data points are plotted using low contrast. Ns is not significant (*P*>0.05).
***P*<0.001.

Memantine's effect on illusion decoding requires attention

To determine how memantine interacted with our manipulations of consciousness and attention, we
examined the decoding of the different features of the second target (T2) from the RSVP task (**Fig. 3A**,
C, and **E**). First, we tested whether masking and the attentional blink (AB) affected local contrast,
collinearity and illusion decoding in the way we expected. Indeed, local contrast decoding was
unaffected by the AB (**Fig. 3B**), and although it was impaired by masking, the drop in decoding accuracy
was numerically small (0.012 AUC on average) and considerably smaller than the effect of masking on
collinearity (0.026 AUC) and illusion (0.123 AUC) decoding. These results are in line with theoretical
proposals and empirical findings that suggest limited effects of masking and attentional manipulations
on feedforward processes (Dehaene et al., 2006; Fahrenfort et al., 2007, 2017; Lamme, 2010; Lamme
et al., 2002). As would be expected for markers of lateral and feedback processing, both collinearity
and illusion decoding were abolished by masking (**Fig. 3C-F**). The AB exclusively impaired illusion
decoding, but decoding accuracy for the illusion remained well above chance during the AB (**Fig. 3E-**
F).

Similar to T1, also for T2 decoding, memantine increased illusion decoding, but only when T2
was presented outside the AB and when it was unmasked (**Fig. 3E**, top left, note that masking fully

abolished illusion decoding). Also our time window-based analyses showed a significant effect of
memantine only when the illusion was both unmasked and fully attended (t_{28}=-2.76, P=0.010,
BF_{10}=4.53; **Fig. 3F**). Although these memantine effects were slightly less robust than for T1, probably
due to reduced trial counts, these results suggest a selective effect of memantine on illusion-related
feedback processing that depends on the availability of attention. One potential concern with this
interpretation is that in the RSVP task, the illusion was always task-relevant and hence attended,
whereas local contrast and collinearity were never task-relevant. It is therefore an open question
whether local contrast and collinearity decoding were unaffected by memantine, because these visual
features were unattended (not task-relevant) and/or because memantine exclusively targets feedback
processes. To alleviate this potential concern, and to replicate the surprising result of increased illusion
decoding under memantine, we carried out a second experiment. A new group of participants was
recruited and we systematically varied task-relevance of local contrast, collinearity, and the illusion in
this task. This allowed us to determine whether memantine indeed specifically affected feedback
processing.

Figure 3. Decoding of the T2 data. (A) Local contrast decoding: mean decoding performance, area under the receiver operating characteristic curve (AUC), over time ± standard error of the mean (SEM). Thick lines differ from chance: P<0.05, cluster-based permutation test. (B) Mean AUC for local contrast (rotation) decoding's time window: 78-94 ms. Error bars are mean ± SEM. Individual data points are plotted using low contrast. Ns is not significant (*P*>0.05). (C) Collinearity decoding: mean AUC over time. (D) Mean AUC for collinearity decoding's time window: 192-292 ms. (E) Kanizsa illusion decoding: mean AUC over time. (F) Mean AUC for Kanizsa illusion's time window: 223-323 ms. **P*=0.010.

Experiment 2 replicates memantine's selective effect on illusion decoding

In experiment 2, the pharmacological design was the same as before and VAS measurements now revealed no side effects from memantine (**Fig. 4B**). Twenty-seven new participants performed a discrimination task in which each visual feature, in separate blocks, was task-relevant (**Fig. 4A**). Approximately every 900-1100 ms an individual target stimulus (no RSVP) was presented for 83 ms when it was unmasked, or for 33 ms when it was masked. For each target, participants indicated whether the current task-relevant feature (e.g., the Kanizsa illusion) was present or absent. In experiment 2, each visual feature was task-relevant and thus attended in some blocks, while it was task-irrelevant and thus less attended in other blocks. This allowed us to directly examine whether memantine's effects were specific to feedback processes or dependent on task-relevance of the stimulus features. For all main analyses of behavior and EEG decoding we conducted repeated-measures ANOVAs with the factors drug (memantine vs. placebo), masking (masked vs. unmasked), and task-relevant feature (local contrast, collinearity, illusion). The complete results of these ANOVAs are presented in the corresponding results figures. We found that masking impaired perceptual sensitivity, most prominently for the detection of the Kanizsa illusion, and less so for collinearity and rotation. Note that we employed less strong masking in this experiment than in experiment 1, leading to overall above chance perceptual sensitivity for the masked trials. For unmasked trials performance was at ceiling for all visual features (no differences in performance). Perceptual sensitivity was unaffected by memantine (**Fig. 4C**).

Figure 4. Experiment 2: design, behavior, and training data. (A) Trial design. (B) Effects of memantine
on subjective state assessed with the visual analogue scale. The categories alertness, contentedness,
and calmness are based on Bond and Lader (1974), sedation on Danion et al. (1989). For details see
Methods. (C) Participants' ability to detect the task-relevant feature. (D) Tenfold cross-validation
decoding scheme applied to an independent training dataset. Mean decoding performance, area
under the receiver operating characteristic curve (AUC), over time ± standard error of the mean (SEM)
is shown. Thick lines differ from chance: P<0.05, cluster-based permutation test. Local contrast,
collinearity, and illusion decoding accuracy peaked at 86, 297, and 266 ms after stimulus onset and
these peaks were encompassed by a 78-94, 247-347, and 217-317 ms time window, respectively. Error
bars are mean ± SEM. Individual data points are plotted using low contrast. Ns is not significant
(*P*>0.05).

The EEG decoding schemes were the same as for experiment 1 and similar time windows were
obtained from the independent training dataset (**Fig. 4D**). Overall, the timing of the decoding of the
three different features in the main experiment replicated our findings from experiment 1 (**Fig. 5**). For
local contrast, task-relevance had no effect and the significant drop in decoding accuracy caused by
masking was again numerically small (0.010 AUC on average); smaller than for collinearity (0.015 AUC)
and illusion decoding (0.069 AUC). Collinearity (**Fig. 5C-D**) and illusion (**Fig. 5E-F**) decoding were
impaired by masking and boosted when the corresponding feature was task-relevant. Crucially, our
cluster-based analyses again revealed that memantine significantly improved decoding only for the
illusion and only when the illusion was unmasked and task-relevant (**Fig. 5E**, top left), thus replicating

the results from experiment 1. Also the time window-based analysis revealed an overall main effect
of drug for illusion decoding, and post-hoc test revealed that the drug effect was only significant when
the illusion was unmasked and task-relevant (post hoc *t*-tests comparing memantine vs. placebo, t_{28}=-
3.36, *P*=0.002, BF_{10}=15.66; **Fig. 5F**). Our time window-based ANOVAs now also revealed a main effect
of memantine on collinearity decoding (**Fig. 5D**), in line with the hypothesized role of NMDA receptors
in recurrent (feedback and lateral) processing.

Figure 5. Decoding results of experiment 2. (A) Local contrast decoding: mean decoding performance, area under the receiver operating characteristic curve (AUC), over time ± standard error of the mean (SEM). Thick lines differ from chance: P<0.05, cluster-based permutation test. (B) Mean AUC for local contrast decoding's time window: 78-94 ms. Error bars are mean ± SEM. Individual data points are plotted using low contrast. Ns is not significant (*P*>0.05). (C) Collinearity decoding: mean AUC over time. (D) Mean AUC for collinearity decoding's time window: 217-317 ms. (E) Kanizsa illusion decoding: mean AUC over time. (F) Mean AUC for Kanizsa illusion's time window: 223-323 ms. *P=0.002.

Discussion

In this study, we investigated the impact of NMDA receptor blockade on multivariate EEG decoding of different features of a visual stimulus (Kanizsa illusion, collinearity, and local contrast). We used the NMDA receptor antagonist memantine to selectively target recurrent processing while minimizing non-specific side effects, a notable improvement over previous studies blocking NMDA receptors using ketamine. Two separate experiments demonstrated selective effects of memantine on decoding accuracy of the Kanizsa illusion, our marker of recurrent processing, while leaving collinearity and contrast decoding relatively unaffected, our markers for lateral and feedforward processing. Interestingly, in both experiments the effect of memantine on illusion decoding was specific to attended, task-relevant stimuli presented without a backward mask. In experiment 1, the illusion was task-relevant and the effect was specific to stimuli presented unmasked and outside the attentional blink (AB), when full attention was available. In experiment 2, we obtained the effect only when the stimulus was unmasked and the illusion was task-relevant. Notably, rather than impairing illusion decoding, in both experiments memantine improved illusion decoding. These findings provide new insight into the role of NMDA-receptor mediated recurrent processing in human visual perception and will be further discussed below.

The effects of memantine are broadly consistent with previous studies indicating that ketamine, another NMDA receptor antagonist, affects perception and neural coding in visual tasks thought to rely on recurrent mechanisms (Meuwese et al., 2013; van Loon et al., 2016). The effect of ketamine in these studies likely affected intra-area lateral connections in early visual cortex (Meuwese et al., 2013) and inter-area feedback mechanisms (van Loon et al., 2016), respectively. The effect of memantine in our study was specific to illusion decoding but the direction of this effect was in the opposite direction: whereas ketamine impaired perceptual organization and visual cortex representations (see also Self et al., 2012; Van Kerkoerle et al., 2014), memantine improved illusion decoding. Although both drugs are known to inhibit NMDA receptors by occupying the receptor's ion channel and are thereby blocking current flow (Glasgow et al., 2017; Molina et al., 2020), previous studies have also found opposite effects of the two drugs. For example, while ketamine diminishes the amplitude of the so-called mismatch negativity (MMN) EEG response to (auditory) oddball stimuli (Rosburg & Kreitschmann-Andermahr, 2016), a single dose of memantine enhances MMN amplitude (Korostenskaja et al., 2007; Light et al., 2017; Swerdlow et al., 2016). Although memantine was found to impair learning as well as markers of perceptual and neural processing in some tasks (e.g., Rammsayer, 2001, 2006; Schugens et al., 1997), most studies found memantine to improve learning and to boost perceptual and neural processing, including EEG markers of the excitation-inhibition (E/I) balance (1/f slope) (Golisch et al., 2017; Korostenskaja et al., 2007; Light et al., 2017; Swerdlow et al., 2016, 2020). As these previous studies typically used auditory stimuli, our results with visual stimuli add to a growing body of literature indicating that memantine can enhance sensory function, instead of decreasing it, and is the first study explicitly linking memantine to recurrent processing.

At this point we can only speculate why NMDA receptor blockade via memantine increased (rather than decreased) decoding accuracy in the condition measuring illusion-specific feedback. Our behavioral results suggest that improved decoding accuracy under memantine was not associated with boosted conscious perception. If anything, to the contrary. In experiment 1 we found that memantine numerically reduced perceptual sensitivity in the same experimental condition in which

we observed improved decoding accuracy under memantine (unmasked, outside the AB, t_{28}=1.58,
P=0.126, BF$_{10}$=0.60). One speculative possibility is that NMDA receptor blockade may have reduced
signal variability or entropy, resulting in more stereotypical neural processing, which may have
improved classifier performance. This may potentially be related to the possibility that NMDA
blockade is known to modulate processing at other pre- or post-synaptic receptors present at NMDA-
rich synapses. Memantine and ketamine both have different actions at receptors other than NMDA,
for instance ketamine also acts on dopamine D2 and serotonin 5-H2 receptors, whereas memantine
is known to inhibit several subtypes of the acetylcholine (ACh) receptor as well as serotonin 5-HT3
receptors. Memantine's therapeutic effects likely involve non-glutamatergic targets (Glasgow et al.,
2017; Johnson et al., 2015). Local increases in ACh levels have been shown to increase firing rates and
reduce firing rate variability in macaques (Herrero et al., 2013, 2008). Further, memantine has been
shown to affect the excitation/inhibition (E/I) balance in human participants with schizophrenia, as
measured noninvasively by computing the aperiodic, 1/f profile of the EEG power spectrum (Molina
et al., 2020). A flatter 1/f power spectrum is thought to reflect noisier signal processing. Changes in
E/I balance have been linked to variations in states of consciousness (He et al., 2010; Nir et al., 2008),
and may also affect decoding performance due to changes in neural noise. Clearly, further research is
needed to elucidate the precise mechanisms by which memantine modulates feedback processing,
how EEG decoding may relate to signal variability, and to explore its implications for perception.

Interestingly, the effect of memantine on illusion decoding was specific to temporally
attended (experiment 1, AB task) and task-relevant Kanizsa stimuli (both experiments) that were
presented without a backward mask. This suggests that object-based attention is crucial for
memantine to exert its effects. Previous work has revealed that object-based attention primarily
enhances neuronal firing rates compared to unattended objects, especially during the recurrent
processing phase (Roelfsema, 2023; Thiele & Bellgrove, 2018). Attention related neural enhancement
can be observed in many regions throughout cortex, e.g., in early and intermediate visual regions, as
well as inferotemporal, parietal, and even frontal cortices (reviewed in Roelfsema, 2023). It is
hypothesized that when an object is attended, this facilitates the integration of its local elements and
that it enables figure-ground segmentation to occur, crucial for the perception of coherent shapes and
for the phenomenological properties of the Kanizsa illusion (Narayanan et al., 2013). The observation
that the effect of inhibiting NMDA receptors by memantine is most pronounced for attended objects,
suggests that memantine is most effective in conditions where recurrent processing is strongest.
Previous work has shown that memantine also affects cholinergic receptors (Glasgow et al., 2017;
Johnson et al., 2015), known to be involved in attentional processes (Herrero et al., 2013, 2008; Thiele
& Bellgrove, 2018), which may partly explain why the effect of memantine is dependent on attentional
allocation. This novel finding deserves future investigation to further understand its mechanistic
origin.

In conclusion, our study demonstrates that memantine exerts selective effects on feedback
processing during human visual perception, and these effects are contingent upon the allocation of
attention. These findings contribute to our understanding of the neural mechanisms underpinning
perception and highlight the importance of considering attentional factors in the study of recurrent
processing. Further research is needed to elucidate the precise mechanisms by which memantine
modulates feedback processing and to explore its implications for our understanding of perception
and consciousness.

Methods

Experiment 1

Participants

Thirty-three right-handed men with normal, or corrected-to-normal, vision were recruited after they
had passed extensive physical and mental screening. Three of them failed at the screening session's

training task and did not proceed to the two experimental sessions (see Procedure). One dropped out
during his first experimental session because of a headache, which was unrelated to memantine as he
had taken the placebo. The remaining 29 participants (22±3 years old) were analyzed. The study was
approved by the Medical Ethical Committee of the Amsterdam Medical Centre (AMC) and the local
ethics committee of the University of Amsterdam. Participants gave informed consent and received
monetary compensation.

Stimuli

The target stimulus set had a 2 Kanizsa illusion levels (absent/present) × 2 collinearity levels
(absent/present) × 2 rotation levels (absent/present) design, resulting in eight stimuli (**Fig. 1A**). Three
aligned Pac-Men induced the Kanizsa illusion. Collinearity was present when the stimuli's three other
elements (the "two-legged white circles") were aligned. The controls for both the illusion and
collinearity were made by rotating their elements 90 degrees. Differences in local contrast were
created by rotating the entire stimulus 180 degrees. The targets spanned 7.5 degrees by 8.3 degrees
of visual angle. The distance between the three Pac-Man stimuli as well as between the three aligned
two-legged white circles was 2.8 degrees of visual angle. Although neuronal responses to collinearity
in primary visual cortex are most robust when this distance is smaller (Kapadia et al., 1995, 2000),
longer-range horizontal connections between neurons with similar orientation selectivity can span
distances corresponding to visual angles considerably greater than 2.8 degrees (Bosking et al., 1997;
Stettler et al., 2002).

The distractor stimulus set was the same as the target stimulus set, with two exceptions. First,
the distractors were red instead of black. Second, the distractors' six elements were rotated 180
degrees relative to the targets', so neither the illusion nor collinearity were present in the distractors.
Masks consisted of six differently shaped elements, all capable of covering the targets' elements. Six
masks were created by rotating the original mask five times by 60 degrees. They spanned 8.5 degrees
by 9.1 degrees of visual angle. Finally, the design of the fixation cross was adapted from Thaler et al.
(2013).

Task designs

The training dataset for our decoding analyses was obtained from a separate "template matching"
task. Here, every 900-1100 ms, a target stimulus was presented for 50 ms. Participants had to indicate
whether the target matched the "template", one of the eight target stimuli that was presented before
the start of a block of trials. Each stimulus served as the template once. In short, the task design was
8 templates × (7 mismatches × 16 presentations + 1 match × 48 presentations), totaling 1280 trials,
3/10 of which were matches.

The main task was a rapid serial visual presentation (RSVP) task. Here, on every trial,
participants were shown two targets (T1 and T2) within a RSVP of distractors (**Fig. 1B**). The targets and
distractors had a stimulus onset asynchrony of 100 ms. T2 and each distractor were presented for 17
ms. With the aim of inducing an attentional blink, attending to T1 was made easier by presenting it
for 67 ms. T1 was preceded by five distractors and T2 was followed by six distractors. For the first six
participants, T2 and each distractor were presented for 10 ms and T1 for 40 ms (using the monitor's
100 Hz refresh rate). These durations were changed to make the task easier, so fewer volunteers
would need to be excluded (see Procedure).

T2 visibility was manipulated in two ways. First, half the T2s were unmasked. The other half
was immediately followed by three masks, each presented for 17 ms with an interstimulus interval of
0 ms. The three masks were selected randomly, but all differed from each other. Masking drastically
reduced T2 visibility. Second, half the lags between the two targets were long: eight distractors (i.e.,
900 ms). The other half was short: one or two distractors (i.e., 200 or 300 ms). The participant's short
lag duration was determined beforehand, during the screening session (see Procedure). Short lags
often induced an attentional blink (i.e., missing T2s). Following the RSVP sequence, participants

indicated for each target whether it contained the Kanizsa illusion or not. Response screens lasted
until the response. Response button mapping was counterbalanced within the task. In short, each
experimental session had an 8 T1 stimuli × 8 T2 stimuli × 2 masking levels (absent/present) × 2 lags
(long/short) task design, resulting in 256 conditions. Each condition was presented four times, totaling
1024 trials.

For both the RSVP and the template matching task, participants were instructed to fixate on
the fixation cross.

Procedure

The experiment consisted of three separate sessions conducted on different days: a four-hour
screening session and two seven-hour experimental sessions. At the beginning of the screening
session, the volunteers were screened for contraindications to using memantine. If there were no
contraindications, participants proceeded to performing the template matching task and training for
the RSVP task. To proceed to the experimental sessions, participants had to score well enough on this
training task. Three participants were excluded for failing to achieve this. The training task was also
used to discover the duration of the short lag that led to the lowest score to reliably induce an
attentional blink. For all participants T2 performance at short lags was more than 5% lower than at
long lags.

Memantine (20 mg) and the placebo were administered in different experimental sessions,
separated by at least a week. These sessions were conducted using a randomized, double-blind
crossover design. The order of drug administration (placebo or memantine first) was counterbalanced
across participants. Participants were required to refrain from recreational drugs four weeks before
the experiment and to refrain from alcohol two days before each experimental session. Experimental
sessions started at 9:00 with the pill intake and briefly practicing the RSVP task. The actual RSVP task
was performed when memantine reached peak plasma level, four hours after the pharmaceutical was
administered. After this task, five minutes of resting state data were collected and another task, the
local-global task (Bekinschtein et al., 2009), was performed. These latter two paradigms are beyond
the scope of this paper.

Before pill intake (~9:00) and before (~13:00) and after (~15:30) the tasks, participants filled
in the visual analogue scale (VAS) to measure the effects of memantine on subjective state (Bond &
Lader, 1974; Danion et al., 1989). Tasks were programmed in Presentation software (Neurobehavioral
Systems) and displayed on a 23 inch, 60 Hz, 1920×1080 pixels monitor, viewed at an 80 cm distance
with the use of a chin rest.

Behavioral analysis

To investigate behavioral performance on the RSVP task, we used perceptual sensitivity (d') and bias
(criterion, see **Fig. S1**) derived from signal detection theory. The attentional blink requires T1 to be
attended, so trials with incorrect T1s were excluded when T2 performance was investigated. To
investigate the VAS measurements, the mean scores of subsets of the scales were calculated and
taken as measures of alertness, contentedness, and calmness following Bond and Lader (1974) and
of sedation following Danion et al. (1989). VAS scores of the two measurements after pill intake
were averaged and corrected by calculating percentage change from the baseline measurement. For
all behavioral analyses, repeated measures ANOVAs and double-sided (Bayesian) t-tests were used
to test the differences between our conditions.

EEG recording and preprocessing

EEG was recorded at 1024 Hz using a 64 channel ActiveTwo system (BioSemi). Four
electrooculographic (EOG) electrodes measured horizontal and vertical eye movements. The data
were analyzed with MATLAB (MathWorks). For most of the preprocessing steps, EEGLAB was used
(Delorme & Makeig, 2004). The data were re-referenced to the earlobes. Poor channels were
interpolated. High-pass filtering can cause artifacts in decoding analyses, so for slow-drift removal we

used trial-masked robust detrending (van Driel et al., 2021). Each trial was epoched from -250 to 1000
ms relative to target onset. To improve our independent component analysis (ICA), baseline correction
was applied using the whole epoch as baseline (Groppe et al., 2009). ICA was used to remove blinks.
Blink components were removed manually. Baseline correction was applied, now using a -250 to 0 ms
window relative to target onset. Trials with values outside of a -300 to 300 microvolts range were
removed. We used an adapted version of FieldTrip's ft_artifact_zvalue funct ion to detect and remove
trials with muscle artifacts (Oostenveld et al., 2011). As in the behavioral analyses, T2s preceded by
incorrect T1s were excluded. Finally, the data were downsampled to 128 Hz.

Multivariate pattern analysis

We used the Amsterdam Decoding and Modeling (ADAM) toolbox to perform multivariate pattern
analyses (MVPAs) on the EEG data (Fahrenfort et al., 2018). Using a tenfold cross-validation scheme,
we first analyzed the template matching task's data. We decoded each visual feature (e.g., the Kanizsa
illusion) by training linear discriminant classifiers to discriminate between the feature's absence and
presence based on the raw EEG activity [across all electrodes]. Individual participants' data were split
into ten equal-sized folds after randomizing the task's trial order. A classifier was then trained on nine
folds and tested on the tenth one, ensuring independence of the training and testing sets. This
procedure was repeated until each fold served as the test set once. Classifier performance, here area
under the receiver operating characteristic (AUC) (Hand & Till, 2001), was averaged across all ten
iterations. We applied "diagonal decoding": classifier performance was calculated for every time
sample and the classifiers were tested on the same time sample they had been trained on.

Each decoded visual feature showed a clear peak in classifier performance over time (**Fig. 1E**).
Many of our analyses focus on the time windows encompassing these peaks: from 78 to 94 ms for
local contrast, 192-292 ms for collinearity, and 223-323 ms for the illusion. To obtain topographic maps
showing the neural sources of the classifier performance, we multiplied the classifier weights with the
data covariance matrix, yielding covariance/class separability maps (Haufe et al., 2014). The
covariance/class separability maps of the three time windows showed that the classifiers depended
on occipitoparietal electrodes during peak decoding accuracy (**Fig. 2A**). We restricted the decoding
analyses to using only these electrodes to ensure that any effects we observed were not due to poor
signal-to-noise ratio.

Next, classifiers were trained on the template m atchings task's data and tested on each
condition of the RSVP task's data. We averaged each visual feature's time window in the training set's
raw EEG data to increase power. Classifiers were trained on these averages and tested on all time
samples of the test set, resulting in "off-diagonal decoding" (King & Dehaene, 2014). To establish the
putative markers for the different forms of neural processing, each visual feature was decoded in this
manner. When decoding a visual feature, the average of the same feature's time window was used as
the training set. Off-diagonal decoding allowed us to focus on the neural processes of interest. For
completeness, we also performed a diagonal decoding version of this train-test scheme. Here,
classifiers trained on a time sample of the template matching task's data were tested on the same
time sample of the RSVP task's data. The diagonal decoding results were qualitatively similar to the
off-diagonal ones (**Fig. S2-4**).

For the time samples from -100 to 700 ms relative to target onset, we used a double-sided *t*-
test to evaluate whether classifier performance differed from chance and whether the memantine
conditions' classifier performance differed from the placebo conditions'. We used cluster-based
permutation testing (10,000 iterations at a threshold of 0.05) to correct for multiple comparisons
(Maris & Oostenveld, 2007). As in the behavioral analyses, repeated measures ANOVAs and double-
sided (Bayesian) *t*-tests—here applied to the time windows' average classifier performance—were
used to test the differences between our conditions.

Experiment 2

Participants

Thirty right-handed men with normal, or corrected-to-normal, vision were recruited after they had passed extensive physical and mental screening. Two of them dropped out after the first experimental session because of personal reasons unrelated to memantine. EEG recording failed during an experimental session of one participant. The remaining 27 participants (22±2 years old) were analyzed. The study was approved by the Medical Ethical Committee of the Amsterdam Medical Centre (AMC) and the local ethics committee of the University of Amsterdam. Participants gave informed consent and received monetary compensation.

Stimuli

Stimuli were the same as in experiment 1, except for a small change to the visual feature collinearity, for which we removed the "feet" from the two-legged white circles (**Fig. S5**).

Task designs

The training dataset for our decoding analyses was obtained from a separate "localizer" task. Here, in separate blocks of trials each visual feature was task-relevant. On every trial, a target stimulus was presented for 83 ms. Participants had to fixate on the fixation cross and indicate whether the current task-relevant feature (e.g., the Kanizsa illusion) was present or absent. The (first) response window started at target onset and ended after 900 ms. If no response was made, "Respond faster!" was presented until response. Trials ended with 0-200 ms jitter and at least 300 ms between response and jitter. The order of the task-relevant features was counterbalanced over participants and, because each participant performed the task multiple times, kept the same within participants. Response button mapping was counterbalanced within the task. In short, the task design was 3 task-relevant features × 8 target stimuli × 72 presentations, totaling 1728 trials.

The main task the participants performed was the "task-relevance" task (**Fig. 4A**), which was the same as the localizer task, except for the following changes. Half the trials were masked and on these trials, the target stimulus was presented for 33 ms and immediately followed by three masks, the same way as in the RSVP task. As in the localizer task each visual feature was at one point task-relevant and thus attended, while at another point task-irrelevant and thus less attended. The task also contained a base-rate manipulation, whose effects are beyond the scope of this paper. In one block of trials, the task-relevant feature was present on 75% of the trials, while on the next block, it was present on 25% of the trials, or vice versa. Overall, the task-relevant feature's was equally often present and absent, which made it possible to ignore the base-rate manipulation for the purpose of the present analyses. In short, the task design was 3 task-relevant features × 8 target stimuli × 2 masking levels (absent/present) × 80 presentations, totaling 3840 trials.

To achieve masked performance of ~75% correct (making it more comparable to a typical attentional blink performance), we staircased mask contrast using the weighted up-down method (Kaernbach, 1991). Staircasing was done when the Kanizsa illusion was task-relevant, because masking affected this feature more than the other two. Contrast levels ranged from 0 (black) to 255 (white). Mask contrast started at level 230. Each correct response made the task more difficult: masks got darker by downward step size S_{down}. Each incorrect response made the task easier: masks got lighter by upward step size S_{up}. Step sizes were determined by $S_{up} \times p = S_{down} \times (1 - p)$, where p was the desired accuracy of 0.75. The smallest step size was always 13 contrast levels. The staircase ended after 25 reversals (a mistake after a correct response, or a correct response after a mistake). The mask contrast that was used during the experimental sessions was the average contrast level of the last 20 reversals.

Procedure

The procedure was the same as for experiment 1, except for the following changes. During the screening session, if the volunteers had no contraindications to using memantine, they now proceeded to performing the localizer task and practicing the task-relevance task. This practice

included our staircase procedure. The mask strength obtained here was used during the experimental sessions. To increase the power of the decoding analyses, the pill intake at the start of the experimental sessions was now immediately followed by another session of the localizer task. The localizer task was followed by briefly practicing the task-relevance task. The actual task-relevance task was performed four hours after the pharmaceutical was administered. No additional tasks were performed. The experiment was conducted in two different types of labs. For 18 participants, tasks were displayed on a 23 inch, 1920 × 1080 pixels monitor and for the other participants, on a 27 inch, 2560 × 1440 pixels monitor.

Behavioral analysis
Behavioral analysis was the same as for experiment 1. One of the baseline VAS forms of one participant was not filled in, so he was excluded from the VAS analysis.

EEG recording and preprocessing
EEG recording and preprocessing were the same as for experiment 1.

Multivariate pattern analysis
Multivariate pattern analysis was the same as for experiment 1, except for the three instances of the localizer task now together serving the role of the template matching task and the task-relevance task serving the role of the RSVP task. When decoding a visual feature (e.g., the Kanizsa illusion), not all of the localizer task's trials were used, only those during which the same visual feature was task-relevant. The tenfold cross-validation scheme was applied to the localizer task and each decoded visual feature again showed a clear peak in classifier performance over time (**Fig. 4D**). Here, the time windows encompassing these peaks were from 78 to 94 ms for local contrast, 247-347 ms for collinearity, and 217-317 ms for the illusion. Covariance/class separability maps again showed that the classifiers depended on occipitoparietal electrodes during peak decoding accuracy, so our decoding analyses were restricted to using only these electrodes.

References

Allen, W. E., Kauvar, I. V., Chen, M. Z., Richman, E. B., Yang, S. J., Chan, K., Gradinaru, V., Deverman, B. E., Luo, L., & Deisseroth, K. (2017). Global Representations of Goal-Directed Behavior in Distinct Cell Types of Mouse Neocortex. *Neuron*, *94*(4), 891-907.e6.

Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., & Naccache, L. (2009). Neural signature of the conscious processing of auditory regularities. *Proceedings of the National Academy of Sciences*, *106*(5), 1672–1677.

Bond, A., & Lader, M. (1974). The use of analogue scales in rating subjective feelings. *The British Journal of Medical Psychology*, *47*(3), 211–218.

Bosking, W. H., Zhang, Y., Schofield, B., & Fitzpatrick, D. (1997). Orientation selectivity and the arrangement of horizontal connections in tree shrew striate cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *17*(6), 2112–2127.

Ciceri, S., Oude Lohuis, M. N., Rottschäfer, V., Pennartz, C. M. A., Avitabile, D., van Gaal, S., & Olcese, U. (2023). The neural and computational architecture of feedback dynamics in mouse cortex during stimulus report. In *bioRxiv* (p. 2023.07.19.549692). https://doi.org/10.1101/2023.07.19.549692

Danion, J. M., Zimmermann, M. A., Willard-Schroeder, D., Grangé, D., & Singer, L. (1989). Diazepam induces a dissociation between explicit and implicit memory. *Psychopharmacology*, *99*(2), 238–243.

Dehaene, S., Changeux, J.-P., Naccache, L., Sackur, J., & Sergent, C. (2006). Conscious, preconscious, and subliminal processing: a testable taxonomy. *Trends in Cognitive Sciences*, *10*(5), 204–211.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21.

Fahrenfort, J. J., Scholte, H. S., & Lamme, V. A. F. (2007). Masking disrupts reentrant processing in
human visual cortex. *Journal of Cognitive Neuroscience*, *19*(9), 1488–1497.
Fahrenfort, J. J., Scholte, H. S., & Lamme, V. A. F. (2008). The spat iotemporal profile of cortical
processing leading up to visual perception. *Journal of Vision*, *8*(1), 12.1-12.
Fahrenfort, J. J., Snijders, T. M., Heinen, K., van Gaal, S., Scholte, H. S., & Lamme, V. A. F. (2012).
Neuronal integration in visual cortex eleva tes face category tuning to conscious face
perception. *Proceedings of the National Academy of Sciences of the United States of America*,
109(52), 21504–21509.
Fahrenfort, J. J., van Driel, J., van Gaal, S., & Olivers, C. N. L. (2018). From ERPs to MVPA Usin g the
Amsterdam Decoding and Modeling Toolbox (ADAM). *Frontiers in Neuroscience*, *12*, 368.
Fahrenfort, J. J., van Leeuwen, J., Olivers, C. N. L., & Hogendoorn, H. (2017). Perceptual integration
without conscious access. *Proceedings of the National Academy of Sciences of the United*
States of America, *114*(14), 3744–3749.
Flohr, H., Glade, U., & Motzko, D. (1998). The role of the NMDA synapse in general anesthesia.
Toxicology Letters, *100–101*, 23–29.
Gilbert, C. D., & Wiesel, T. N. (1979). Morphology and intracortical projections of functionally
characterised neurones in the cat visual cortex. *Nature*, *280*(5718), 120–125.
Glasgow, N. G., Povysheva, N. V., Azofeifa, A. M., & Johnson, J. W. (2017). Memant ine and Ketamine
Differentially Alter NMDA Receptor Desensitization. *The Journal of Neuroscience: The Official*
Journal of the Society for Neuroscience, *37*(40), 9686–9704.
Golisch, A., Heba, S., Glaubitz, B., Tegenthoff, M., & Lissek, S. (2017). Enhancing E ffects of NMDA-
Receptor Blockade on Extinction Learning and Related Brain Activation Are Modulated by BMI.
Frontiers in Behavioral Neuroscience, *11*, 34.
Groppe, D. M., Makeig, S., & Kutas, M. (2009). Identifying reliable independent components via split-
half comparisons. *NeuroImage*, *45*(4), 1199–1211.
Halgren, E., Mendola, J., Chong, C. D. R., & Dale, A. M. (2003). Cortical activation to illusory shapes as
measured with magnetoencephalography. *NeuroImage*, *18*(4), 1001–1009.
Hand, D. J., & Till, R. J. (2001). A Simple Generalisation of the Area Under the ROC Curve for Multiple
Class Classification Problems. *Machine Learning*, *45*(2), 171–186.
Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J. -D., Blankertz, B., & Bießmann, F. (2014). On
the interpretation of weight vectors of linear models in multivariate neuroimaging.
NeuroImage, *87*, 96–110.
He, B. J., Zempel, J. M., Snyder, A. Z., & Raichle, M. E. (2010). The temporal structures and functional
significance of scale-free brain activity. *Neuron*, *66*(3), 353–369.
Herrero, J. L., Gieselmann, M. A., Sanayei, M., & Thiele, A. (2013). Attention -induced variance and
noise correlation reduction in macaque V1 is mediated by NMDA receptors. *Neuron*, *78*(4),
729–739.
Herrero, J. L., Roberts, M. J., Delicato, L. S., Gieselmann, M. A., Dayan, P., & Thiele, A. (2008).
Acetylcholine contributes through muscarinic receptors to attentional modulation in V1.
Nature, *454*(7208), 1110–1114.
Johnson, J. W., Glasgow, N. G., & Povy sheva, N. V. (2015). Recent insights into the mode of action of
memantine and ketamine. *Current Opinion in Pharmacology*, *20*, 54–63.
Kaernbach, C. (1991). Simple adaptive testing with the weighted up-down method. *Perception &*
Psychophysics, *49*(3), 227–229.
Kapadia, M. K., Ito, M., Gilbert, C. D., & Westheimer, G. (1995). Improvement in visual sensitivity by
changes in local context: parallel studies in human observers and in V1 of alert monkeys.
Neuron, *15*(4), 843–856.
Kapadia, M. K., Westheimer, G., & Gilbe rt, C. D. (2000). Spatial distribution of contextual interactions
in primary visual cortex and in visual perception. *Journal of Neurophysiology*, *84*(4), 2048–
2062.

King, J.-R., & Dehaene, S. (2014). Characterizing the dynamics of mental representations: the temporal generalization method. *Trends in Cognitive Sciences*, *18*(4), 203–210.

Kocsis, B., Brown, R. E., McCarley, R. W., & Hajos, M. (2013). Impact of ketamine on neuronal network dynamics: translational modeling of schizophrenia-relevant deficits. *CNS Neuroscience & Therapeutics*, *19*(6), 437–447.

Kok, P., Bains, L. J., van Mourik, T., Norris, D. G., & de Lange, F. P. (2016). Selective Activation of the Deep Layers of the Human Primary Visual Cortex by Top-Down Feedback. *Current Biology: CB*, *26*(3), 371–376.

Kok, P., & de Lange, F. P. (2014). Shape perception simultaneously up- and downregulates neural activity in the primary visual cortex. *Current Biology: CB*, *24*(13), 1531–1535.

Korostenskaja, M., Nikulin, V. V., Kicić, D., Nikulina, A. V., & Kähkönen, S. (2007). Effects of NMDA receptor antagonist memantine on mismatch negativity. *Brain Research Bulletin*, *72*(4–6), 275–283.

Lamme, V. A. F. (2010). How neuroscience will change our view on consciousness. *Cognitive Neuroscience*, *1*(3), 204–220.

Lamme, V. A. F., Zipser, K., & Spekreijse, H. (1998). Figure-ground activity in primary visual cortex is suppressed by anesthesia. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(6), 3263–3268.

Lamme, V. A. F., Zipser, K., & Spekreijse, H. (2002). Masking interrupts figure-ground signals in V1. *Journal of Cognitive Neuroscience*, *14*(7), 1044–1053.

Lee, T. S., & Nguyen, M. (2001). Dynamics of subjective contour formation in the early visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(4), 1907–1911.

Li, Z. (1998). A neural model of contour integration in the primary visual cortex. *Neural Computation*, *10*(4), 903–940.

Liang, H., Gong, X., Chen, M., Yan, Y., Li, W., & Gilbert, C. D. (2017). Interactions between feedback and lateral connections in the primary visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(32), 8637–8642.

Light, G. A., Zhang, W., Joshi, Y. B., Bhakta, S., Talledo, J. A., & Swerdlow, N. R. (2017). Single-Dose Memantine Improves Cortical Oscillatory Response Dynamics in Patients with Schizophrenia. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *42*(13), 2633–2639.

Manita, S., Suzuki, T., Homma, C., Matsumoto, T., Odagawa, M., Yamada, K., Ota, K., Matsubara, C., Inutsuka, A., Sato, M., Ohkura, M., Yamanaka, A., Yanagawa, Y., Nakai, J., Hayashi, Y., Larkum, M. E., & Murayama, M. (2015). A Top-Down Cortical Circuit for Accurate Sensory Perception. *Neuron*, *86*(5), 1304–1316.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

Mashour, G. A., & Hudetz, A. G. (2018). Neural Correlates of Unconsciousness in Large-Scale Brain Networks. *Trends in Neurosciences*. https://doi.org/10.1016/j.tins.2018.01.003

Mashour, G. A., Roelfsema, P., Changeux, J.-P., & Dehaene, S. (2020). Conscious Processing and the Global Neuronal Workspace Hypothesis. *Neuron*, *105*(5), 776–798.

Mathew, S. J., Shah, A., Lapidus, K., Clark, C., Jarun, N., Ostermeyer, B., & Murrough, J. W. (2012). Ketamine for treatment-resistant unipolar depression: current evidence. *CNS Drugs*, *26*(3), 189–204.

Meuwese, J. D. I., van Loon, A. M., Scholte, H. S., Lirk, P. B., Vulink, N. C. C., Hollmann, M. W., & Lamme, V. A. F. (2013). NMDA receptor antagonist ketamine impairs feature integration in visual perception. *PloS One*, *8*(11), e79326.

Molina, J. L., Voytek, B., Thomas, M. L., Joshi, Y. B., Bhakta, S. G., Talledo, J. A., Swerdlow, N. R., & Light, G. A. (2020). Memantine Effects on Electroencephalographic Measures of Putative

Excitatory/Inhibitory Balance in Schizophrenia. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, *5*(6), 562–568.

Müller, C. P., Pum, M. E., Amato, D., Schüttler, J., Huston, J. P., & Silva, M. A. D. S. (2011). The in vivo neurochemistry of the brain during general anesthesia. *Journal of Neurochemistry*, *119*(3), 419–446.

Narayanan, B., Stevens, M. C., Jiantonio, R. E., Krystal, J. H., & Pearlson, G. D. (2013). Effects of memantine on event-related potential, oscillations, and complexity in individuals with and without family histories of alcoholism. *Journal of Studies on Alcohol and Drugs*, *74*(2), 245–257.

Nir, Y., Mukamel, R., Dinstein, I., Privman, E., Harel, M., Fisch, L., Gelbard-Sagiv, H., Kipervasser, S., Andelman, F., Neufeld, M. Y., Kramer, U., Arieli, A., Fried, I., & Malach, R. (2008). Interhemispheric correlations of slow spontaneous neuronal fluctuations revealed in human sensory cortex. *Nature Neuroscience*, *11*(9), 1100–1108.

Noorman, S., Stein, T., Fahrenfort, J. J., & van Gaal, S. (2023). Isolating feedforward, lateral, and feedback mechanisms underlying perceptual and attentional impairments of conscious access. In *bioRxiv* (p. 2023.10.16.562523). https://doi.org/10.1101/2023.10.16.562523

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869.

Pak, A., Ryu, E., Li, C., & Chubykin, A. A. (2020). Top-Down Feedback Controls the Cortical Representation of Illusory Contours in Mouse Primary Visual Cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *40*(3), 648–660.

Rammsayer, T. H. (2001). Effects of pharmacologically induced changes in NMDA-receptor activity on long-term memory in humans. *Learning & Memory*, *8*(1), 20–25.

Rammsayer, T. H. (2006). Effects of pharmacologically induced changes in NMDA receptor activity on human timing and sensorimotor performance. *Brain Research*, *1073–1074*, 407–416.

Roelfsema, P. R. (2023). Solving the binding problem: Assemblies form when neurons enhance their firing rate-they don't need to oscillate or synchronize. *Neuron*, *111*(7), 1003–1019.

Rosburg, T., & Kreitschmann-Andermahr, I. (2016). The effects of ketamine on the mismatch negativity (MMN) in humans--a meta-analysis. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *127*(2), 1387–1394.

Sachidhanandam, S., Sreenivasan, V., Kyriakatos, A., Kremer, Y., & Petersen, C. C. H. (2013). Membrane potential correlates of sensory perception in mouse barrel cortex. *Nature Neuroscience*, *16*(11), 1671–1677.

Schmidt, K. E., Goebel, R., Löwel, S., & Singer, W. (1997). The perceptual grouping criterion of colinearity is reflected by anisotropies of connections in the primary visual cortex. *The European Journal of Neuroscience*, *9*(5), 1083–1089.

Scholte, H. S., Jolij, J., Fahrenfort, J. J., & Lamme, V. A. F. (2008). Feedforward and recurrent processing in scene segmentation: electroencephalography and functional magnetic resonance imaging. *Journal of Cognitive Neuroscience*, *20*(11), 2097–2109.

Schugens, M. M., Egerter, R., Daum, I., Schepelmann, K., Klockgether, T., & Löschmann, P. A. (1997). The NMDA antagonist memantine impairs classical eyeblink conditioning in humans. *Neuroscience Letters*, *224*(1), 57–60.

Self, M. W., Kooijmans, R. N., Supèr, H., Lamme, V. A. F., & Roelfsema, P. R. (2012). Different glutamate receptors convey feedforward and recurrent processing in macaque V1. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(27), 11031–11036.

Seth, A. K., & Bayne, T. (2022). Theories of consciousness. *Nature Reviews. Neuroscience*. https://doi.org/10.1038/s41583-022-00587-4

Stettler, D. D., Das, A., Bennett, J., & Gilbert, C. D. (2002). Lateral connectivity and contextual interactions in macaque primary visual cortex. *Neuron*, *36*(4), 739–750.

Swerdlow, N. R., Bhakta, S., Chou, H.-H., Talledo, J. A., Balvaneda, B., & Light, G. A. (2016). Memantine Effects On Sensorimotor Gating and Mismatch Negativity in Patients with Chronic Psychosis. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *41*(2), 419–430.

Swerdlow, N. R., Bhakta, S. G., Talledo, J., Kotz, J., Roberts, B. Z., Clifford, R. E., Thomas, M. L., Joshi, Y. B., Molina, J. L., & Light, G. A. (2020). Memantine effects on auditory discrimination and training in schizophrenia patients. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *45*(13), 2180–2188.

Takahashi, N., Oertner, T. G., Hegemann, P., & Larkum, M. E. (2016). Active cortical dendrites modulate perception. *Science*, *354*(6319), 1587–1590.

Thaler, L., Schütz, A. C., Goodale, M. A., & Gegenfurtner, K. R. (2013). What is the best fixation target? The effect of target shape on stability of fixational eye movements. *Vision Research*, *76*, 31–42.

Thiele, A. (2012). NMDA receptors figure it out [Review of *NMDA receptors figure it out*]. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(27), 10749–10750.

Thiele, A., & Bellgrove, M. A. (2018). Neuromodulation of Attention. *Neuron*, *97*(4), 769–785.

van Driel, J., Olivers, C. N. L., & Fahrenfort, J. J. (2021). High-pass filtering artifacts in multivariate classification of neural time series data. *Journal of Neuroscience Methods*, *352*, 109080.

Van Kerkoerle, T., Self, M. W., Dagnino, B., Gariel-Mathis, M.-A., Poort, J., Van Der Togt, C., & Roelfsema, P. R. (2014). Alpha and gamma oscillations characterize feedback and feedforward processing in monkey visual cortex. *Proceedings of the National Academy of Sciences*, *111*(40), 14332–14341.

van Loon, A. M., Fahrenfort, J. J., van der Velde, B., Lirk, P. B., Vulink, N. C. C., Hollmann, M. W., Scholte, H. S., & Lamme, V. A. F. (2016). NMDA Receptor Antagonist Ketamine Distorts Object Recognition by Reducing Feedback to Early Visual Cortex. *Cerebral Cortex*, *26*(5), 1986–1996.

van Vugt, B., Dagnino, B., Vartak, D., Safaai, H., Panzeri, S., Dehaene, S., & Roelfsema, P. R. (2018). The threshold for conscious report: Signal loss and response bias in visual and frontal cortex. *Science*, *360*(6388), 537–542.

Wang, X.-J. (2008). Decision making in recurrent neuronal circuits. *Neuron*, *60*(2), 215–234.

Wokke, M. E., Vandenbroucke, A. R. E., Scholte, H. S., & Lamme, V. A. F. (2013). Confuse your illusion: feedback to early visual cortex contributes to perceptual completion. *Psychological Science*, *24*(1), 63–71.

Wyatte, D., Jilk, D. J., & O'Reilly, R. C. (2014). Early recurrent feedback facilitates visual object recognition under challenging conditions. *Frontiers in Psychology*, *5*, 674.

Zatka-Haas, P., Steinmetz, N. A., Carandini, M., & Harris, K. D. (2021). Sensory coding and the causal impact of mouse cortex in a visual decision. *ELife*, *10*. https://doi.org/10.7554/eLife.63163