A high-light tolerant alga from the desert is protected from oxidative stress

by NPQ-independent responses

Guy Levin[1]*,**, Michael Yasmin[1]*, Oded Liran[2], Rawad Hanna[1], Oded Kleifeld[1], Guy

Horev[3], Francis-André Wollman[4], Gadi Schuster[1,5] and Wojciech J. Nawrocki[4] **

[1] Faculty of Biology, Technion, Haifa, 32000 Israel

[2] Kineret Limnological Laboratory, Israel Oceanographic & Limnological Research, Migdal,

14950 Israel

[3] Migal Galilee Research Institute Ltd, Qiryat Shemona, 11016 Israel

[4] Unité Mixte de Recherche 7141, Centre National de la Recherche Scientifique, Sorbonne

Université, Institut de Biologie Physico-Chimique, Paris, France

[5] Grand Technion Energy Program, Technion, Haifa, 32000 Israel

*These authors have equally contributed to this work

**Corresponding authors: guylevin@campus.technion.ac.il or guylevin5@gmail.com (GL),

wojciech.nawrocki@cnrs.fr (WN).

Running title: NPQ-independent photoprotection in a desert alga

Abstract

Non-photochemical quenching (NPQ) mechanisms are crucial for protecting photosynthesis from photoinhibition in plants, algae, and cyanobacteria, and their modulation is a long-standing goal for improving photosynthesis and crop yields. The current work demonstrates that *Chlorella ohadii*, a green micro-alga that thrives in the desert under high light intensities which are fatal to many photosynthetic organisms, does not perform nor require NPQ to protect photosynthesis under constant high light. Instead of dissipating excess energy, it minimizes its uptake by eliminating the photosynthetic antenna of photosystem II. In addition it accumulates antioxidants that neutralize harmful reactive oxygen species (ROS) and ramps up cyclic electron flow around PSI. These NPQ-independent responses proved efficient in preventing ROS accumulation and reducing oxidative damage to proteins in high-light-grown cells.

Introduction

Light drives photosynthesis, but it also damages the photosynthetic apparatus in a number of ways. Particularly, illumination facilitates the generation of reactive oxygen species (ROS) which damage the molecular players in photosynthesis, thereby inhibiting the process – an effect known as photoinhibition (Ögren, 1988; Ögren and Sjöström, 1990). Under high light (HL) conditions, non-photochemical quenching (NPQ) mechanisms through which excess absorbed light energy is harmlessly dissipated as heat, become critical for photosynthesis protection (photoprotection)(Delosme, 1967; Wolff and Witt, 1969; Demmig-Adams et al., 1989; Demmig-Adams et al., 1990). The major NPQ component, q_E, is located within the membrane-intrinsic major light-harvesting complex (LHCII) for photosystem II (PSII) and is activated by the acidification of the lumen compartment which results from photosynthetic electron transport. Typically, q_E is the strongest and the most rapidly triggered component of NPQ in the green lineage, undergoing activation in seconds, as recently reviewed by (Bassi and Dall'Osto, 2021). During periods of high photosynthetic activity, the thylakoid lumen is acidified due to proton translocation across the thylakoid membrane by the combined activity of linear and cyclic photosynthetic electron flows (LEF and CEF). PsbS and LhcSR in plants and algae, respectively act as proton sensors which, upon acidification of the lumen, induce q_E to dissipate excess energy as heat (Li et al., 2000; Li et al., 2004; Peers et al., 2009; Liguori et al., 2013). Simultaneously, the xanthophyll cycle is activated, and newly formed zeaxanthin interacting in the PSII-LHCII chlorophyll pool provides pathways for energy dissipation. (Demmig et al., 1987; Niyogi et al., 1998; Xu et al., 2015; Sacharz et al., 2017).

Chlorella ohadii is a green micro-alga that thrives in the desert, where it is exposed to extremely high light intensities (Treves et al., 2013; Levin et al., 2021). Accordingly, *C. ohadii* is highly tolerant to photoinhibition (Treves et al., 2016; Levin et al., 2023; Levin et al., 2024). The adaptation of *C. ohadii* to HL involves marked changes in thylakoid content of which the most pronounced is a decrease in LHCII, as well as the massive accumulation of carotenoids and several photoprotective proteins (Levin et al 2021, 2023). HL-grown *C. ohadii* does not undergo state transitions, a mechanism of antenna exchange between PSII and PSI (Bonaventura and Myers, 1969; Chow et al., 1981; Delosme et al., 1996), which also has been implicated in photoprotection (Allorent et al., 2013; Nawrocki et al., 2016). Moreover, *C. ohadii* also lack genes encoding LhcSR proteins nor does it accumulate the PsbS protein (Treves et al., 2016; Levin et al., 2023; Murik et al., 2024), suggesting the absence of q_E. Interestingly, a carotenoid biosynthesis-related protein (CBR), which belongs to the LHC-like protein family with a structure similar to PsbS, accumulates to a great extent in HL-grown *C. ohadii* (Levy et al., 1993; Levin et al., 2023; Levin and Schuster, 2023). Structural modeling suggested that one of the proton-sensing glutamates of PsbS was conserved in CBR, raising the possibility of its role in lumen acidification sensing. Moreover, fractionation of the photosynthetic protein complexes by sucrose density gradient centrifugation showed that CBR from HL-grown cells co-localized with a detached fraction of LHCII and carotenoids, both of which are associated with NPQ and ROS scavenging (Levin et al., 2023). Given that LHCII is the major site of NPQ (Ruban et al., 1999; Elrad et al., 2002; Sacharz et al., 2017; Nicol and Croce, 2021), its elimination and the lack of LhcSR raise doubts as to the contribution of a q_E regulation which require to investigate further the mode of photoprotection in *C. ohadii*.

In this work, the contribution of the decrease in PSII antenna to HL acclimation, as well as the interplay between NPQ and CEF on ROS accumulation and oxidative damage in *C. ohadii* are investigated. We demonstrate that q_E is actually absent and replaced with an NPQ-independent photoprotection in HL-grown *C. ohadii* cells. This mechanism, rather than allowing energy dissipation, mainly relies limiting the absorption of light. It is concomitantly coupled to a dramatic enhancement of CEF around PSI and a vast accumulation of CBR, carotenoids, and other antioxidants, functionally reducing ROS accumulation. Thus, *C. ohadii,* adapted to grow at extremely high light intensity, employs both active and passive mechanisms of resistance to photoinhibition. The present work demonstrates that NPQ is not required to withstand excess illumination in HL-grown *C. ohadii* and that under conditions of prolonged

exposure to HL intensities, minimizing the PSII-LHCII absorption cross-section and ROS pro-

duction while maximizing antioxidant activity has been selected as the most successful strategy

to reduce photoinhibition.

Results

HL-grown cells upregulate CBR and enzymatic antioxidant enzymes.

To understand the remarkable resistance of *C. ohadii* to HL intensities, we grew it under

low and high light conditions (LL: 50 µmol photons m^{-2} s^{-1}; HL: 2000 µmol photons m^{-2} s^{-1}),

and then subjected the cells to label-free quantification mass spectrometry (LFQ-MS) to detect

possible enzymatic antioxidants that participate in the detoxification of ROS and determine

their effect on chloroplast protein oxidation. As was observed for isolated thylakoid membranes

(Levin et al., 2023), the most prominent protein in HL-grown cells was CBR, with a 995-fold

change compared to LL-grown cells. This was accompanied by a drastic loss in LHCII (Fig.

1A). Changes in LHCI abundance were much smaller than in LHCII, which suggests that PSII

light-harvesting was more affected than PSI. Additionally, we identified an increase in 15 pro-

teins associated with ROS detoxification or with energy dissipation via alternative electron

transfer pathways, (Figs. 1A and 1B). Specifically, we detected two isoforms of superoxide

dismutase (SOD), which catalyzes the detoxification of superoxide (O_2^-) by converting it to

H_2O_2, , one of which showing a 1.8 higher accumulation in HL-grown cells than in LL-grown

cells. Of glutathione peroxidase (GPX) and ascorbate peroxidase (APX), which can further

detoxify H_2O_2 by catalyzing its conversion to water (H_2O), only GPX was significantly accu-

mulated in the HL-grown cells (3.3-fold change). Dehydroascorbate reductase (DHAR) and

two isoforms of glutathione-s-transferase (GST), which participate in detoxifying the ascor-

bate-glutathione cycle, were also significantly accumulated in HL-grown cells (3.8-, 1.9- and

1.2-fold change, respectively). There were no statistically significant changes for catalase,

which catalyzes the conversion of H_2O_2 to H_2O in HL-grown cells. ω-3 fatty acid desaturase

(ω3FAD), which catalyzes the generation of polyunsaturated fatty acids (PUFAs), underwent

a 2.1-fold increased accumulation in HL- compared to LL-grown cells. PUFAs have been sug-

gested to contribute to ROS detoxification by interacting with ROS during lipid peroxidation

(Schmid-Siegert et al., 2016). Flavodiiron proteins (Flv1 and Flv2), which act as electron sinks

possibly involved in PSII and/or PSI photoprotection (Chaux et al., 2015; Burlacot et al., 2022;

Bag et al., 2023), were significantly accumulated in HL-grown cells (2.9- and 8.4-fold change,

respectively), providing an additional route for energy dissipation. In contrast, PTOX, another

putative electron sink (Nawrocki et al., 2015), was not differentially accumulated in HL-grown
cells (0.9-fold change) (Figs. 1A and 1B).

To additionally understand whether the strong change in the detoxification enzyme
abundance was not simply a reason of their absence in LL, we relied on absolute quantification
of the enzyme concentration. To this end, intensity-based absolute quantification (iBAQ)
(Schwanhäusser et al., 2011) was used (Fig. 1B). Out of the abovementioned enzymes, SOD
was the most abundant (top 0.7% percentile compared to all detected proteins), followed by
GPX (6.7%) and CBR (7.9%). (Fig. 1B). This further highlights the contribution of CBR and
enzymatic antioxidants for photoprotection in HL-grown *C. ohadii* cells.

HL-grown cells accumulate less oxidatively damaged proteins.

To functionally assess the effects of ROS detoxification, the extent of oxidative damage
was assessed by quantifying the irreversible oxidation of tryptophan – itself leading to protein
degradation (Kato et al., 2023). Remarkably, there were more oxidized peptides from PSII,
LHCII, PSI, and LHCI subunits, accumulating in LL-grown cells when compared to HL-grown
cells (Fig. 1C), despite the 20-fold lower illumination intensity. This observation underlines
the high efficiency of ROS scavenging in HL-grown cells. Taken together, our proteomics
analysis suggests that, in addition to the robust downregulation of LHCII and marked accumu-
lation of carotenoid in HL-grown cells (Levin et al., 2021; Levin et al., 2023), SOD and GPX
– as well as potentially CBR - play a major role in reducing ROS accumulation and oxidative
damage in HL-grown cells.

High light acclimation results in LHCII loss and reduced PSII functional antenna.

To understand the extent of losses in LHCII complexes, thylakoid membranes isolated
from *C. ohadii* grown under LL and HL conditions were solubilized and separated by centrif-
ugation on sucrose density gradients. As shown in Fig. 2A, LL-grown cells displayed, in addi-
tion to a PSI-LHCI supercomplex, a PSII-LHCII super-complex, and some free LHCII. In con-
trast, while we observed no changes in PSI-LHCI between cells grown at either light regime,
we found no PSII-LHCII super-complex in HL-grown *C. ohadii*. In contrast, only the PSII core
complex was visible on the gradient.

To examine how the loss in LHCII and the accumulation of CBR could affect light
harvesting in PSII, we performed measurements of its functional antenna size. To this end, we
studied the fluorescence rise in the presence of the PSII inhibitor DCMU, which measures the

Q_A reduction rate upon continuous illumination. The functional antenna size is approximated

by calculating the reciprocal of the area above the fluorescence rise curve. A larger area relates

to a smaller antenna size, given the slower reduction of Q_A. As shown in Fig. 2B, the area

above the fluorescence rise curves was larger in HL-cells at all actinic light intensities used.

The comparison of the slopes of linear regression fits across the measured rate points indicates

a 40% difference in PSII absorption capacity between those two conditions (Fig. 2C). This

demonstrates that the loss in LHCII induces a strong decrease in the amount of light absorbed

by HL-adapted *C. ohadii*. Nonetheless, this is a more limited effect than expected from the

massive loss of LHCII observed by the sucrose density gradient assay. These seemingly con-

flicting observations may be resolved if the CBR protein associated with a large number of

carotenoids would act as a PSII antenna that substitutes in part for the loss in LHCII., Alterna-

tively, the LL-grown cells may exhibit already small supercomplex size or accommodate a

significant fraction of quenched, free LHCII which do not transfer their excitation energy to

the PSII reaction centers.

To understand the effect of a lower PSII antenna size on the PSII electron transfer rate

(ETR II) under continuous illumination, the PSII yield (Φ_{PSII}) was measured after a 30-second

illumination at various light intensities. The yield was then multiplied by the maximal PSII rate

at these intensities to provide the ETR II. The results are shown in Fig. 2D and demonstrate

that in LL-cells the saturation of PSII ETR is reached at about 400 µmol photons m^{-2} s^{-1}, similar

to other algae and plants. However, at HL-cells that lack the PSII-antenna, the saturation is

reached at much higher light intensities, well above 1500 µmol photons m^{-2} s^{-1}, and the ETR

amplitude is higher than in LL cells.

Absence of the q_E component of NPQ in Chlorella ohadii

Since the PSII peripheral antenna in green algae and plants is a major site of reversible

NPQ, we investigated this photoprotective mechanism in *C. ohadii*. LL- and HL-grown cells

were dark-adapted and then subjected to high actinic light intensity (2000 µmol photons m^{-2} s^{-}

^{1}), after which, their fluorescence behavior were recorded. Strikingly, both LL- and HL-grown

cells showed a negligible NPQ response, as indicated by the limited decrease at F_M', with little

if any NPQ calculated following the equation: NPQ = $(F_M/F_M')-1$, where F_M and F_M' are the

maximal fluorescence of dark-adapted cells or cells exposed to light, respectively, during a

saturating light pulse (Fig. 3A). The decrease in F_M' was very limited in *C. ohadii* compared

to (HL-grown) *C. reinhardtii* tested under similar light conditions (Fig. 3A). At 2000 µmol

photons m^{-2} s^{-1}, NPQ reached a value of ~0.2 (Fig. 3B), in *C. ohadii* compared to ~2.3 in *Chlamydomonas*, owing to its well established q_E NPQ response of high amplitude, facilitated by the proton-sensing capabilities of LhcSR (Fig. 3A). When *C. ohadii* was exposed to even higher light intensities (3000 µmol photons m^{-2} s^{-1}), or when grown under strict photoautotrophic conditions, LL-grown cells developed only a limited NPQ response (~0.7), with similar levels of NPQ in HL-grown cells (~0.2) (Supplementary Figs. 1B, 1C, and 2)., Thus under all tested conditions, virtually no light-induced q_E quenching was detected in *C. ohadii*.

q_E is the major and most rapid component of NPQ which responds to light in seconds and is activated by a pH decrease of the thylakoid lumen. Lumen acidification occurs due to the translocation of protons resulting from high photosynthetic electron flow under HL conditions. These protons bind conserved luminal glutamate residues in LhcSR (algae) and PsbS (plants) and activate q_E. Even though *C. ohadii* lacks the gene for LhcSR and shows no evidence for PsbS expression (Treves et al., 2016; Levin et al., 2021; Levin et al., 2023), CBRs - which largely accumulate in HL-cells - harbour a conserved glutamate that may function as the proton sensor activating the q_E (Levin et al., 2023; Levin and Schuster, 2023). To exclude that there was a limitation to the lumen acidification due to the strong decrease of PSII antenna size, and to verify if CBR can trigger qE at a low lumen pH, we used externally-added acid method, as described in (Tian et al., 2019). The fluorescence traces of LL and HL-grown cells were followed after artificial lumen acidification to pH 5.5. In agreement with the above in vivo analysis of light-induced NPQ, lumen acidification had little effect on the fluorescence signal, indicating the absence of q_E activation in both LL- and HL-grown *C. ohadii* cells cultured in either mixotrophic (Fig. 3C) or strictly photoautotrophic (Supplementary Fig. 3) conditions. When the same assay was applied to *C. reinhardtii* cells as a positive control, fluorescence changes, indicative of a strong q_E were observed (Fig. 3C). The lack of q_E agrees with the absence of the lhcSR gene and the lack of PsbS accumulation in *C. ohadii*, and mirrors the behavior of a *C. reinhardtii* mutant lacking the major NPQ effector, LhcSR3 (Tian et al., 2019), and the psbS mutant of Arabidopsis (Wilson et al., 2024). Importantly, we conclude that despite a conserved glutamate (E131 in *C. ohadii*, E122 in *A. thaliana*) residue on the lumenal side of the predicted CBR structure (Levin et al., 2023; Levin and Schuster, 2023), this protein does not act as a pH-induced q_E trigger.

Together, these results indicate that *C. ohadii* does not rely on NPQ mechanisms to survive when grown under intense HL. While LL-grown cells still showed a very limited,

219 slowly developing NPQ effect, possibly due to q_I quenching, HL-grown cells were fully lack-
220 ing in the two types of NPQ.

HL-grown cells accumulate less damaging ROS than LL-grown cells

To quantify the photoprotective capacity of *C. ohadii* to limit ROS accumulation in the
absence of NPQ, we used confocal microscopy coupled with the detection of Singlet Oxygen
Sensor Green (SOSG). This dye emits green fluorescence upon interaction with singlet oxygen
($^1O_2$) (Supplementary Fig. 4). LL- and HL-grown cells were incubated with SOSG and exposed
to 10 min of high red actinic light, followed by quantification of the accumulated $^1O_2$. HL-
grown, as compared to LL-grown cells, accumulated significantly lower levels of $^1O_2$ (Figs.
4A-C). To evaluate the contribution of $^1O_2$ that was generated via photosynthesis-related pro-
cesses to this difference, we analyzed the SOSG fluorescence that was emitted specifically
from within the chloroplast (Supplementary Fig. 4). The analysis of 24 and 54 individual chlo-
roplasts from LL and HL-grown cells, respectively, showed that individual chloroplasts of LL-
grown cells accumulated much higher levels of $^1O_2$ (Fig. 4C), with a 324% increase following
10 min of red-light illumination, while in HL-grown cells, the increase was limited to an aver-
age of 175% (Fig. 4C). These results are in line with the increased accumulation of O_2^- in LL-
grown as compared with HL-grown algae, while the two cell types accumulated similar
amounts of H_2O_2 (Fig. 4D) (Levin et al., 2023). Kinetic analyses further confirmed that LL-
grown cells accumulated $^1O_2$ at a significantly higher rate (Fig. 5). Moreover, a significant
proportion of chloroplasts of LL-grown cells accumulated 10 times more $^1O_2$ than those in HL-
grown cells (Fig. 4B).

To determine how changes in peripheral antenna between the two light growth condi-
tions were involved in these differences in photosensitivity and ROS production, $^1O_2$ produc-
tion was compared between detached LHCII fractions (Fig. 2A) of LL- vs. HL-grown cells at
the same chlorophyll concentration. Importantly, detached LHCII fractions from HL-grown
cells also contain large amounts of carotenoids and CBR, which may contribute to reduced $^1O_2$
accumulation via direct $^1O_2$ scavenging (Levin et al., 2023). The results shown in Fig. 4E
demonstrate that the CBR-containing detached LHCII fractions from HL-grown cells indeed
accumulated less $^1O_2$ compared to the same fraction from LL-grown cells. However, the dif-
ference was smaller than in whole-cell measurements, suggesting that both the antenna size
difference and the differences in the detached antenna properties or the accumulation of CBR
and carotenoids had a paramount influence on the rate of ROS synthesis upon photosynthesis.

These observations argue for photoprotective effects mainly originating from the limitation of both light absorption and $^1O_2$ accumulation by the peripheral antenna.

Cyclic electron flow is highly activated in HL-grown cells

A decrease in PSII antenna relative to that of PSI creates a favorable environment for CEF around PSI (Fig. 6A) (Tagawa et al. 1963; Joliot and Joliot 2005; Alric 2015; Nawrocki et al. 2019b). This led us assess the kinetics of light-induced P_{700} oxidation to determine whether CEF exhibited a different behavior in LL- vs. HL-grown *C. ohadii*. P_{700} cation absorption at 830 nm serves as a proxy for the activity of CEF when PSII is inactivated by the addition of the herbicide DCMU, or when both CEF and LEF are blocked by DBMIB – a cytochrome b_6f inhibitor (Nawrocki et al., 2019a). Drastically different behaviors of P_{700} oxidation were observed between the LL- and HL-grown algae (Fig. 6B). When exposed to the highest tested actinic light intensities in the presence of DCMU, a long lag in P_{700} oxidation was noted when cells were grown in HL, suggesting a high CEF activity (Fig. 6B upper panels). Remarkably, this lag was still observed in presence of methyl viologen, an electron acceptor from PSI, which also decreases the frequency of back reactions while preserving CEF to some extent (Nawrocki et al., 2019a; Joliot et al., 2022). The half-time of PSI oxidation by CEF under these conditions of extremely strong illumination, as determined by the signal obtained following the addition of DCMU, reached 4 seconds, at least 4 times longer than in LL-grown *C. ohadii*, and more than one order of magnitude longer than in *Chlamydomonas* (Nawrocki et al., 2019a). Indeed, the steady state of P_{700} oxidation under continuous illumination at 2000 µmol photons m^{-2} s^{-1}– itself a commonly-used proxy for CEF activity (Takahashi et al., 2013; Alric, 2014) - was only around 80%, indicating that even in the absence of PSII activity, a very high rate of CEF can be maintained in *C. ohadii* adapted to HL conditions (Fig. 6B). As expected, this effect was abolished when DBMIB rather than DCMU was added, demonstrating the requirement of electron flow through the cytochrome b_6f to maintain CEF. Finally, at lower actinic light intensities in the absence of inhibitors, there was no accumulation of P_{700}^+ due to its efficient reduction by PSII (Fig. 6B). The effect of DCMU was visible in cells grown in LL, while no effect was detected in HL-grown *C. ohadii*, further indicating that a highly efficient CEF can maintain P_{700} mostly reduced. The inability to completely oxidize P_{700} at low light in the presence of DBMIB hints to additional pathways of P_{700}^+ reduction, as substantiated by the relatively rapid reduction of the cation at the end of the illumination period.

The stimulation of CEF in HL-grown cells was consistent with mass spectrometry data.
According to our proteomics analysis, proton gradient regulation 5 (PGR5) and PRG5-like
photosynthetic phenotype 1 (PGRL1), which regulate CEF (Nawrocki et al., 2019b), showed
enhanced accumulation in HL-grown cells (1.5-fold for both enzymes) (Fig. 6C). Also, the
enhanced accumulation of ferredoxin (Fd) (3.7-fold), together with the decrease in Fd-NADP-
reductase (FNR) levels (0.4-fold change), suggested a shift towards CEF rather than to LEF
(Fig. 6C).

Together, these results imply that CEF is induced and is a major electron pathway dur-
ing low light and high light exposure in HL-grown cells, Therefore, aside from the natural
requirement to balance the photosynthetic electron flow according to the imbalanced absorp-
tion cross-sections of PSII and PSI, the highly active CEF in HL-grown cells likely contributes
to extra ATP generation that would be required in these environmental conditions.

Discussion

An efficient NPQ-independent photoprotective mechanism underlines *C. ohadii* remark-
able resistance to high-light stress

Extremophile species are fascinating forms of living matter, even more so that - their
growth environments being exacerbated versions of stressful situations - they may serve as
models for the discovery of engineering strategies for the generation of more resistant crops.
Photoprotection is thought to be a primary photosynthesis defense mechanism. The various
forms of NPQ-controlled photoprotection processes are widely studied, from their molecular
mechanism, by numerous molecular and chemical effectors, to their impact on gene expression
(for recent reviews, see (Niyogi and Truong, 2013; Murchie and Ruban, 2020; Bassi and
Dall'Osto, 2021)). While similar NPQ pathways are widespread in most plants and algae, the
current work shows that an extremely light-tolerant microalga, *C. ohadii*, does not require any
NPQ to thrive under extreme HL intensities (Fig. 7). When these cells are grown at LL inten-
sities, PSII harvests light both through its core and peripheral antennae and generates ROS, as
the membranes contain limited amount of carotenoids (Fig. 7A). However, when these LL-
grown cells are exposed to HL intensities before being acclimated to HL conditions, even more
ROS are produced, leading to protein oxidation and a high-light acclimation response (Fig.
7B). Indeed, this acclimation is efficient: we observed large differences in the levels of oxidized
peptides between the LL and HL cells, suggestive of higher levels of protein damage in LL-
grown cells (Fig. 1C). Acclimation to HL conditions leads to a major loss in PSII peripheral

antennae while several antioxidant enzymes, CBR, and a large number of carotenoids accumu-

late. The combined effect of these processes is a marked decrease in the generation and accu-

mulation of ROS, which results in less protein oxidation and turnover (Fig. 7C). Thus, although

the elimination of the LHCII in HL-grown *C. ohadii* leads to a potential loss of NPQ sites, it

significantly reduces the absorption cross-section of PSII which, in turn, keeps ROS production

to a low level under high light. Together with ROS detoxification carried out by the increased

accumulation of antioxidants, it drastically reduces the rate of damage that could be caused by

the concurrent energy conversion process (Fig. 7).

322 Minimizing light absorption may be a more efficient approach to photoprotection than
323 NPQ induction under steady illumination.

A drop in LHCII is potentially a risky acclimation strategy for photoprotection upon

exposure to high light conditions for at least two reasons. It leads to a loss in most NPQ regu-

latory processes which are critical to mitigate the effects abrupt changes in light intensities

under a fluctuating light regime. In addition, it decreases the light-harvesting capabilities dur-

ing LL periods. However, *C. ohadii* thrives in the soil crusts of the desert under direct sunlight,

where few clouds or vegetation can provide shading. Therefore it is placed in a very stable light

environment with extended time periods under constant HL. In these conditions the small core

antenna of PSII provides enough light energy to fuel PSII-mediated electron transfer and elim-

ination of LHCII allows a drastic reduction in light-induced ROS production (Fig. 7). It is of

note that alongside LHCII elimination, the loss of NPQ in *C. ohadii* also is driven by the ab-

sence of the LhcSR gene and PsbS protein (Treves et al., 2016; Levin et al., 2023). Thus, *C.*

ohadii provides a remarkable demonstration of what NPQ is not suited for: it does not deliver

an appropriate photoprotection against long-lived illumination at very high light intensities, as

encountered among extremophiles. NPQ should be regarded as a series of photoprotection re-

sponses that evolved with oxygenic photosynthesis exposed to fluctuating light ranging from

frequent non-saturating illumination to transient exposures to saturating light as is the case for

most crop species and microalgae studied so far. This interpretation is in line with the recent

data of photosynthetic response in the frequency domain, with strongest NPQ effect taking

place on ~minutes timescale (Niu et al., 2023).

The extent of the antenna loss, estimated from mass spectrometry data upon solubiliza-

tion of photosynthetic membranes, looked more extensive that the significant but more limited

~40% functional decrease in PSII light harvesting capacity during a switch between LL and

HL growth (Figs. 1 and 2). This apparent discrepancy may stem from a functional antenna size

in LL conditions being smaller than suggested by the amount of LHCII in the sample, in which
case a fraction of this peripheral antenna would be disconnected from the reaction centers but
aggregated in a quenched state. These quenched LHCII would not contribute to PSII cross-
section, and they would barely affect the F_V/F_M in cells, and their loss upon HL acclimation
would be consistent with the decrease of chlorophyll content. As an alternative hypothesis, a
substitution mechanism for LHCII may take place under HL condition: another light-harvest-
ing complex would contribute to the PSII absorption cross-section, CBRs being the primary
candidates given their impressive increased abundance. In the future, time- and spectrally-re-
solved fluorescence methods could reveal the lifetimes of chlorophyll excited states in each
fraction of the antenna in vivo and help decipher which of the two hypotheses is correct.

Minimizing light absorbance while accumulating antioxidants leads to reduced oxidative
damage

In addition to lowering ROS production, the decrease in LHCII content in *C. ohadii* is
accompanied by the accumulation of antioxidants such as carotenoids (Levin et al., 2021; Levin
et al., 2023) and specialized enzymes, specifically GPX, SOD, and CBR (Fig. 1A). The role of
the latter protein remains elusive, but the marked increase in accumulation under HL suggests
a key role in *C. ohadii* photoacclimation (Fig. 1A). Despite its structural similarities with PsbS
and LhcSR, CBR does not induce q_E or other NPQ processes via lumen acidity sensing (Fig.
3B and Supplementary Fig. 3). It was suggested that CBR binds carotenoids (Levy et al., 1993)
and, indeed, it was found to co-localize with carotenoids and the detached LHCII fraction of
HL-grown *C. ohadii* (Levin et al., 2023). Carotenoids are efficient ROS scavengers (Tri-
antaphylidès and Havaux, 2009), and CBR may contribute to these activities while it is bound
to carotenoids. Moreover, the significant accumulation of carotenoids in the thylakoid mem-
brane of HL-grown *C. ohadii*, predominantly zeaxanthin, and lutein, may contribute to this
effect (Levin et al., 2023). In addition to being a ROS target, CBR may be an internal UV light
screen, expressed by the cells to prevent donor-side damage to PSII.

The overall level of detectable ROS was significantly decreased upon illumination of
HL-grown alga. Given the elimination of the PSII antenna and antioxidant accumulation, it is
likely that the decline in ROS levels is a product of the smaller PSII absorption cross-section
and of the detoxifying effect of the upregulated ROS-scavenging enzymes (Fig. 7). More spe-
cifically, the smaller cross-section results in a mild decrease in the amount of absorbed light,
which limits ROS synthesis (Santabarbara et al., 1999; Santabarbara et al., 2001). Then, the

decreased concentration of nascent ROS due to scavenging activity or using CBR-bound carotenoids as an oxidation target, minimizes the overall accumulation of these deleterious species.

What is the role of highly active cyclic electron flow in *C. ohadii*?

Finally, CEF around PSI is often thought of as a transient process that occurs before the Calvin-Benson cycle becomes active at the onset of illumination of vascular plants. In addition, because CEF drives acidification of the thylakoid lumen, it contributes to photoprotection of PSII through NPQ activation but also to protecting PSI from photoinhibition since it slows down P_{700}^+ re-reduction by the cytochrome b_6f complex (Harris and Boekema, 2018). Last, it has also been implicated at a means to maintain ATP production in anaerobic conditions in microalgae, when mitochondrial respiration is inhibited (for recent reviews on CEF see (Suorsa, 2015; Labs et al., 2016; Nawrocki et al., 2019b)) The finding that CEF is highly active particularly in HL-grown *C. ohadii* calls for a specific role in this alga. Since, little or no lumen acidification-induced NPQ was detected in the alga (Fig. 3B), it shows that this highly active CEF is not used to boost PSII photoprotection by the classic NPQ mechanism. Additionally, P_{700} is primarily maintained in its neutral form even under high light (Fig. 6B)(Caspy et al., 2021). These findings suggest that CEF is not employed to boost lumen acidification – neither for PSII, nor for PSI photoprotection.

It remains that CEF in *C. ohadii* should enhance ATP synthesis in HL conditions. This may be needed to cope with high rates of photoinhibition under desert growth conditions and the heavy energetic cost of PSII repair. Generally speaking, CEF-generated ATP would provide the required energy for keeping with intracellular metabolism in non-growing or slowly-growing conditions, whether for protein and DNA repair, protein translation and energy storage for dark periods.

It is a long-standing technical difficulty in the field of functional photosynthesis to ascertain that transient CEF in the absence of PSII activity – as is the case in our measurements – reflects the situation when LEF is functioning (Fan et al., 2016). The long-lived CEF at the onset of illumination in *C. ohadii* could nonetheless serve as a model for the development of novel methodologies for its quantification, a critical feat for photosynthesis physiology in the coming years.

Materials and Methods

Cell culture and acclimation to low or high light conditions

Exponential-phase *Chlorella ohadii* cells were diluted to optical density (OD)$_{750nm}$ 0.2 in a final

volume of 400 ml tris-acetate phosphate (TAP) medium and then split: 200 ml were placed

under LL (50 µmol photons m^{-2} s^{-1}) conditions and 200 ml were placed under HL (2,000 µmol

photons m^{-2} s^{-1}) conditions, in 1 L Erlenmeyer under shaking (100 rpm), for 24 h, at 28 °C. To

prevent self-shading, the cells were maintained below an OD$_{750nm}$ value of 0.8 by diluting them

to OD$_{750nm}$ 0.2 every 8 h during the growth period. After 24 h, the LL- and HL-grown cells

were harvested for further experimentation. For experiments under strict photoautotrophic con-

ditions, the cells were grown in a tris-phosphate (TP) medium, with no acetate as a carbon

source.

Chlamydomonas stt7-9 strain, lacking the kinase activity required for State transitions, was

grown in continuous 400 µmol photons m^{-2} s^{-1} condition in MIN media.

Photosynthetic complexes separation by sucrose density gradient centrifugation

LL- and HL-grown cells were pelleted and washed twice with sucrose (300 mM)-tris-tricine

(15 mM each pH 8)-NaCl$_2$ (15 mM) (STN) buffer. The cells were broken in a MicroFluidizer

(50 psi); unbroken cells were pelleted (15,000 g, 10 min, 4 °C). The supernatant was collected

and thylakoid membranes were pelleted in an ultracentrifuge (250,000g, 2 h, 4 °C). Thylakoid

membranes were resuspended in 25 mM MES (pH 6.5) to a chlorophyll concentration of 0.2

or 0.4 mg/ml for LL- or HL-grown samples, respectively, and solubilized with 1% *n*-dodecyl-

α-D-maltopyranoside (α-DDM) for 15 min, on ice, under dark conditions. Insolubilized mate-

rial was pelleted (9,500 g, 10 min, 4 °C) and the supernatant was loaded on a sucrose density

gradient (20 mM MES pH 6.5, 0.02% α-DDM, 4%-45% sucrose). Individual complexes were

separated by ultracentrifugation (100,000 g, 20 h, 4 °C)

Functional analyses

434   LL- and HL-grown cells were pelleted and resuspended in fresh medium to the equivalent of

435   10 µg/ml chlorophyll for PSII chlorophyll fluorescence analysis, or 50 µg/ml for P$_{700}$ (PSI)

436   absorption analysis. The cells were dark-adapted for at least 30 min with air bubbling, then

437   loaded to a quartz cuvette and analyzed with a Dual-PAM-100 fluorometer (Walz, Germany).

438   Red actinic light was used in all measurements.

PSII electron transport rate saturation curve

440   After exposure of LL- and HL-grown cells to different light intensities for 30 sec, the PSII

441   operating efficiency (Φ_{PSII}) was measured using the following equation: $\Phi_{PSII} = (F_M' - F')/F_M'$.

PSII ETR was calculated following the equation: $\text{ETR(II)} = \sigma_{II} \cdot \Phi_{PSII}$, where σ_{II} is the maximal, light-limited rate of PSII at each light intensity ("PSII antenna size"). The latter is the reciprocal of the area above fluorescence-DCMU curve, delimited by F_M value normalized to 1 and by the t_0.

NPQ analysis

The fluorescence traces of LL- and HL-grown cells were monitored upon exposure to various light intensities (50, 400, 2000, and 3000 μmol photons m^{-2} s^{-1}) for 10 min, followed by 3 min of darkness for recovery. When indicated, acetic acid (1 mM final) was added directly to the cuvette to induce q_E (Tian et al., 2019). NPQ was calculated following the equation: NPQ = (F_M/F_M')-1 in both *C. ohadii* and *C. reinhardtii*.

P_{700} redox measurements

P_{700} oxidation kinetics were monitored in LL- and HL-grown cells upon illumination with various light intensities: 50, 400, or 2000 μmol photons m^{-2} s^{-1}, in the presence or absence of the photosynthetic electron transfer inhibitors 3-(3,4-dichlorophenyl)-1,1-dimethylurea (100 μM, DCMU) or 2,5-dibromo-3-methyl-6-isopropylbenzoquinone (100 μM, DBMIB). Methyl viologen (6 mM) was added to all samples to prevent charge recombination in PSI.

ROS accumulation assay

Live-cell $^1O_2$ quantification with singlet oxygen sensor green

LL- and HL-grown *C. ohadii* cells were pelleted and resuspended to OD$_{750nm}$ 0.2 in TAP medium containing 50μM singlet oxygen sensor green (SOSG). Subsequently, the cells were incubated for 30 min under LL conditions, at 28°C, and pelleted and resuspended in fresh TAP medium. To visualize the formation of $^1O_2$, the cells were loaded to a spinning disk confocal microscope (Nikon Eclipse Ti2, Confocal scanner unit Yokogawa CSU-W1, Camera Photometrics PrimS-BSI). Light stimulation was achieved using a red LED integrated into the microscope, providing an intensity of 3.9 mW (660 nm) for a duration of 1-10 min. The SOSG was excited at 488 nm, and the fluorescence emission was recorded at 530 nm. Chlorophyll fluorescence was recorded with excitation at 430 nm and emission at 680 nm to identify the chloroplasts. IMARIS software (Oxford Instruments) was employed to reconstruct the chloroplast structure based on the chlorophyll fluorescence. Subsequently, only SOSG fluorescence that corresponded to the reconstructed chloroplast was measured, ensuring a specific and accurate assessment of the effects of light stress on the observed photosynthetically derived $^1O_2$

formation. The graphs were prepared with the R environment for statistical computing and graphics (https://www.R-project.org/) using the ggplot2 package (Wickham, 2016). The linear mixed model was calculated with the lme4 package (Bates et al., 2015). SEs and post-hoc analysis were calculated using the emmeans package (https://CRAN.R-project.org/package=emmeans). Mixed-model ANOVA was used to determine statistical significance (see Fig. 5B)

$^1O_2$ accumulation in isolated LHCII complexes

Isolated LHCII from LL-grown cells and isolated LHCII together with CBR from HL-grown cells were resuspended in STNM buffer (50 mM Tricine-KOH pH 7.6, 15mM $NaCl_2$, 300 mM sucrose, and 5mM $MgCl_2$) with SOSG (10 mM) to a final chlorophyll concentration of 0.028 µg/µl. The samples were loaded onto a black 96-well plate and SOSG fluorescence was measured before and after 10 min of light illumination in a fluorescence plate reader (ClarioStar 500, BMG Labtech, Ortenberg, Germany) with excitation at 488 nm and an emission range of 530-600nm. The samples were placed on ice during the illumination period to avoid protein denaturation. STNM buffer with SOSG was used as a blank to measure and exclude the background SOSG fluorescence. Three biological repeats were tested for each treatment condition.

O_2^- and H_2O_2 accumulation in low-light- and high-light-grown C. ohadii cells

Nitro-blue tetrazolium (NBT), which upon reduction creates blueish–greyish insoluble precipitates, and 3,3′-diaminobenzidine (DAB), which upon oxidation by H_2O_2 becomes dark brown, were used to measure O_2^- and H_2O_2 accumulation in LL- and HL-grown *C. ohadii* cells. The cell OD_{750nm} was adjusted to 0.7. One millilitre of each sample was subjected to a 10 min incubation under HL conditions, with the addition of either 5 mM NBT or 5 mM DAB. Thereafter, chlorophyll was extracted with 100% methanol, at 40°C. Cells were then pelleted (3000 g, 5 min), and resuspended in 200 µl methanol (100%). The resulting cell suspension was applied as spots onto the Whatman 3 paper filter. ROS were quantified with ImageJ, based on the colour intensity.

Mass spectrometry whole-cell sample preparation and proteolysis

Three biological repeats of LL- and HL-grown *C. ohadii* cells were harvested, washed twice with STN, and broken under 50 psi with a MicroFluidizer. Samples were suspended in 8.5 M urea, 100 mM ammonium bicarbonate, and 10 mM DTT. Protein content was estimated using

Bradford readings. The samples were reduced (60 ºC, 30 min), modified with 35.2 mM iodoa-
cetamide in 100 mM ammonium bicarbonate (room temperature for 30 min in the dark), and
digested in 1.5 M urea and 17.6 mM ammonium bicarbonate with modified trypsin (Promega),
overnight, at 37 °C, at a 1:50 (M/M) enzyme-to-substrate ratio. A second digestion with trypsin
was performed for 4 h, at 37 °C, at a 1:100 (M/M) enzyme-to-substrate ratio. The tryptic pep-
tides were desalted using Oasis HLB 96-well µElution Plate (Waters), dried, and re-suspended
in 0.1% formic acid.

Mass spectrometry analysis

The resulting peptides were analysed by LC-MS/MS using a Q Exactive HFX mass spectrom-
eter (Thermo) fitted with an HPLC capillary (Ultimate 3000, Thermo Scientific). The peptides
were loaded in solvent A (0.1% formic acid in water) on a homemade capillary column (30
cm, 75-micron ID) packed with Reprosil C18-Aqua (Dr. Maisch GmbH, Germany). The pep-
tide mixture was resolved with a 5-28% linear gradient of solvent B (99.99% acetonitrile with
0.1% formic acid) for 180 min, followed by a gradient of 28-95% for 15 min, and then 15
minutes at 95% acetonitrile with 0.1% formic acid in water at flow rates of 0.15 µl/min. Mass
spectrometry was performed in a positive mode (m/z 350-1200, resolution 120,000 for MS1
and 15,000 for MS2), using repetitively full MS scan followed by high collision dissociation
(HCD, at 27 normalized collision energy) of the 30 most-dominant ions (>1 charges) selected
from the first MS scan. The AGC settings were $3x10^6$ for the full MS and $1x10^5$ for the MS/MS
scans. A dynamic exclusion list was enabled with an exclusion duration of 20 s.

Mass spectrometry data analysis

Raw data was processed with MaxQuant version 2.5.2 (Tyanova et al., 2016) for protein
accumulation analysis and MSfragger version 4.0 (Kong et al., 2017) via FragPipe version 21.1
(https://fragpipe.nesvilab.org/) for detection of tryptophan oxidized peptides. In the absence of
a fully annotated proteome for *C. ohadii*, we utilized a combined dataset of protein sequences
in our identification process. This dataset includes all available Chlorella protein sequences
(Tax ID = 3701) downloaded from UniProt as of July 2019, comprising a total of 20973 se-
quences. Additionally, we incorporated all available *C. ohadii* protein sequences (11470) from
NCBI, as downloaded on October 2023. All searches included enzymatic cleavage with trypsin
(cleavage after K/R but not P) and two missed cleavages and a clip of N-term methionine unless
specified otherwise. Cysteine carbamidomethyl (+57.02146) was set as a fixed modification,

and methionine oxidation (+15.994915) and protein N-terminal acetylation were set as variable modifications.

MaxQuant searches were performed using tryptic digestion mode with a minimal peptide length of 7 amino acids. Search criteria included up to 2 missed cleavages, precursor and fragments tolerance of 20 ppm, oxidation of methionine, and protein N-terminal acetylation set as variable modifications. Default settings were used for all other parameters. Candidates were filtered to obtain an FDR of 1% at the peptide and protein levels. For quantification, the match between runs (MBR) module of MaxQuant was used with the Label-Free Quantification (LFQ) normalization method enabled.

Identification and quantification of tryptophan oxidation was performed using Frag-Pipe's 'LFQ-MBR' workflow. The workflow default settings were used with the additions of tryptophan oxidation (+15.994915) and deoxidation (+31.989829) as variable modifications for identifying tryptophan oxidation.

Further analysis was performed using the Perseus software v.2.0.11 (Tyanova et al., 2016). Label-free quantification (LFQ) intensity values were used to calculate the relative abundance of total proteins and tryptophan oxidized peptides in LL and HL whole-cell samples. The results were filtered to exclude contaminants, reverse, and only identified by site proteins. Label-free quantification values were transformed (Log2) and filtered to include only proteins that were detected in three repeats of at least one group: LL or HL samples. Missing values were then added from a normal distribution (https://cox-labs.github.io/coxdocs/replace-missingfromgaussian.html). A two-sample t-test was used to determine the statistical significance of differences between groups, applying a permutation-based false discovery rate (FDR) of 0.05 and an S0 parameter of 0.1. Intensity-based absolute quantification (iBAQ) (Schwanhäusser et al., 2011) was used to determine the abundance of a given protein in HL-grown cells. A protein with a higher iBAQ value is more abundant compared to a protein with a low iBAQ value. The proteins were listed according to their average iBAQ values (high to low) in HL-grown cells. The percentile of the protein was determined by dividing its number (position) on the list, by the total number of proteins, and multiplying by 100. For example, if protein X is 40[th] on the list, and a total of 4000 proteins were detected in the analysis, then protein X is in the 1[st] percentile, i.e. in the top 1% most abundant proteins in the cell.

Data availability

The mass spectrometry proteomics RAW data have been deposited to the ProteomeXchange

Consortium via the PRIDE (Perez-Riverol et al., 2022) partner repository with the dataset iden-

tifier PXD053918

Funding:

Funding was provided by an Israel Science Foundation (ISF) grant (2199/22).

Acknowledgments:

We thank Dr. Nitzan Dahan from the Technion LS&E microscopy unit for his assistance and

guidance. We thank the Smoler Proteomics Center at the Technion for performing the mass

spectrometry experiments. We thank Nir Keren and Iftach Yacobi for their helpful advice and

discussions.

Author contributions: GL, GS, FAW, and WN planned the research and wrote this manu-

script. GL and MS performed the experiments and data analysis unless stated otherwise. OK

and RH assisted with proteomics data analysis. GH helped analyze the singlet oxygen accumu-

lation data and performed statistical analyses. OL assisted with NPQ measurements.

Declaration of interests:

The authors declare no competing interests.

Dedication

This paper is dedicated to the late Professor Itzhak Ohad, an outstanding and beloved scientist

and mentor, who made a substantial contribution to our understanding of photoinhibition and

paved the way to the isolation of *C. ohadii*.

References

Allorent G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D, Petroutsos
D, Kuntz M, Breyton C, et al (2013) A Dual Strategy to Cope with High Light in *Chlamydomo-*
nas reinhardtii. The Plant Cell **25**: 545–557

Alric J (2014) Redox and ATP control of photosynthetic cyclic electron flow in Chlamydomonas rein-
hardtii. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1837**: 825–834

Bag P, Shutova T, Shevela D, Lihavainen J, Nanda S, Ivanov AG, Messinger J, Jansson S (2023) Fla-
vodiiron-mediated O2 photoreduction at photosystem I acceptor-side provides photoprotec-
tion to conifer thylakoids in early spring. Nat Commun **14**: 3210

Bassi R, Dall'Osto L (2021) Dissipation of Light Energy Absorbed in Excess: The Molecular Mecha-
nisms. Annu Rev Plant Biol **72**: 47–76

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using **lme4**. J
Stat Soft. doi: 10.18637/jss.v067.i01

Bonaventura C, Myers J (1969) Fluorescence and oxygen evolution from Chlorella pyrenoidosa. Bio-
chimica et Biophysica Acta (BBA) - Bioenergetics **189**: 366–383

Burlacot A, Dao O, Auroy P, Cuiné S, Li-Beisson Y, Peltier G (2022) Alternative photosynthesis path-
ways drive the algal CO2-concentrating mechanism. Nature **605**: 366–371

**Caspy I, Neumann E, Fadeeva M, Liveanu V, Savitsky A, Frank A, Kalisman YL, Shkolnisky Y, Murik
O, Treves H, et al** (2021) Cryo-EM photosystem I structure reveals adaptation mechanisms
to extreme high light in Chlorella ohadii. Nat Plants **7**: 1314–1322

Chaux F, Peltier G, Johnson X (2015) A security network in PSI photoprotection: regulation of photo-
synthetic control, NPQ and O2 photoreduction by cyclic electron flow. Front Plant Sci. doi:
10.3389/fpls.2015.00875

Chow W, Telfer A, Chapman DJ, Barber J (1981) State 1-state 2 transition in leaves and its associa-
tion with ATP-induced chlorophyll fluorescence quenching. Biochimica et Biophysica Acta
(BBA) - Bioenergetics **638**: 60–68

Delosme R (1967) Étude de l'induction de fluorescence des algues vertes et des chloroplastes au dé-
but d'une illumination intense. Biochimica et Biophysica Acta (BBA) - Bioenergetics **143**:
108–128

Delosme R, Olive J, Wollman F-A (1996) Changes in light energy distribution upon state transitions:
an in vivo photoacoustic study of the wild type and photosynthesis mutants from Chlamydo-
monas reinhardtii. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1273**: 150–158

Demmig B, Winter K, Krüger A, Czygan F-C (1987) Photoinhibition and Zeaxanthin Formation in In-
tact Leaves: A Possible Role of the Xanthophyll Cycle in the Dissipation of Excess Light En-
ergy. Plant Physiol **84**: 218–224

**Demmig-Adams B, Adams WW, Heber U, Neimanis S, Winter K, Krüger A, Czygan F-C, Bilger W,
Björkman O** (1990) Inhibition of Zeaxanthin Formation and of Rapid Changes in Radiationless
Energy Dissipation by Dithiothreitol in Spinach Leaves and Chloroplasts. Plant Physiol **92**:
293–301

Demmig-Adams B, Winter K, Krüger A, Czygan F-C (1989) Zeaxanthin Synthesis, Energy Dissipation,
and Photoprotection of Photosystem II at Chilling Temperatures. Plant Physiol **90**: 894–898

Elrad D, Niyogi KK, Grossman AR (2002) A Major Light-Harvesting Polypeptide of Photosystem II
Functions in Thermal Dissipation[W]. The Plant Cell **14**: 1801–1816

Fan D-Y, Fitzpatrick D, Oguchi R, Ma W, Kou J, Chow WS (2016) Obstacles in the quantification of
the cyclic electron flux around Photosystem I in leaves of C3 plants. Photosynth Res **129**:
239–251

Harris JR, Boekema EJ, eds (2018) Membrane Protein Complexes: Structure and Function. doi:
10.1007/978-981-10-7757-9

Joliot P, Sellés J, Wollman F-A, Verméglio A (2022) High efficient cyclic electron flow and functional
supercomplexes in Chlamydomonas cells. Biochimica et Biophysica Acta (BBA) - Bioenerget-
ics **1863**: 148909

Kato Y, Kuroda H, Ozawa S-I, Saito K, Dogra V, Scholz M, Zhang G, de Vitry C, Ishikita H, Kim C, et al
(2023) Characterization of Tryptophan Oxidation Affecting D1 Degradation by FtsH in the
Photosystem II Quality Control of Chloroplasts. doi: 10.7554/eLife.88822.2

Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI (2017) MSFragger: ultrafast
and comprehensive peptide identification in mass spectrometry–based proteomics. Nat
Methods **14**: 513–520

Labs M, Rühle T, Leister D (2016) The antimycin A-sensitive pathway of cyclic electron flow: from
1963 to 2015. Photosynth Res **129**: 231–238

Levin G, Kulikovsky S, Liveanu V, Eichenbaum B, Meir A, Isaacson T, Tadmor Y, Adir N, Schuster G
(2021) The desert green algae *Chlorella ohadii* thrives at excessively high light intensities by
exceptionally enhancing the mechanisms that protect photosynthesis from photoinhibition.
The Plant Journal **106**: 1260–1277

Levin G, Schuster G (2023) LHC-like Proteins: The Guardians of Photosynthesis. IJMS **24**: 2503

Levin G, Yasmin M, Liveanu V, Burstein C, Hanna R, Kleifeld O, Simanowitz MC, Meir A, Tadmor Y,
Hirschberg J, et al (2023) A desert *Chlorella* sp. that thrives at extreme high-light intensities
using a unique photoinhibition protection mechanism. The Plant Journal **115**: 510–528

Levin G, Yasmin M, Pieńko T, Yehishalom N, Hanna R, Kleifeld O, Glaser F, Schuster G (2024) The
protein phosphorylation landscape in photosystem I of the desert algae *Chlorella* sp. New
Phytologist nph.19603

Levy H, Tal T, Shaish A, Zamir A (1993) Cbr, an algal homolog of plant early light-induced proteins, is
a putative zeaxanthin binding protein. Journal of Biological Chemistry **268**: 20892–20896

Li X-P, Björkman O, Shih C, Grossman AR, Rosenquist M, Jansson S, Niyogi KK (2000) A pigment-
binding protein essential for regulation of photosynthetic light harvesting. Nature **403**: 391–
395

Li X-P, Gilmore AM, Caffarri S, Bassi R, Golan T, Kramer D, Niyogi KK (2004) Regulation of Photosyn-
thetic Light Harvesting Involves Intrathylakoid Lumen pH Sensing by the PsbS Protein. Jour-
nal of Biological Chemistry **279**: 22866–22874

Liguori N, Roy LM, Opacic M, Durand G, Croce R (2013) Regulation of Light Harvesting in the Green
Alga *Chlamydomonas reinhardtii* : The C-Terminus of LHCSR Is the Knob of a Dimmer Switch.
J Am Chem Soc **135**: 18339–18342

Murchie EH, Ruban AV (2020) Dynamic non-photochemical quenching in plants: from molecular
mechanism to productivity. The Plant Journal **101**: 885–896

Murik O, Geffen O, Shotland Y, Fernandez-Pozo N, Ullrich KK, Walther D, Rensing SA, Treves H
(2024) Genomic imprints of unparalleled growth. New Phytologist **241**: 1144–1160

Nawrocki WJ, Bailleul B, Cardol P, Rappaport F, Wollman F-A, Joliot P (2019a) Maximal cyclic electron flow rate is independent of PGRL1 in Chlamydomonas. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1860**: 425–432

Nawrocki WJ, Bailleul B, Picot D, Cardol P, Rappaport F, Wollman F-A, Joliot P (2019b) The mechanism of cyclic electron flow. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1860**: 433–438

Nawrocki WJ, Santabarbara S, Mosebach L, Wollman F-A, Rappaport F (2016) State transitions redistribute rather than dissipate energy between the two photosystems in Chlamydomonas. Nat Plants **2**: 16031

Nawrocki WJ, Tourasse NJ, Taly A, Rappaport F, Wollman F-A (2015) The Plastid Terminal Oxidase: Its Elusive Function Points to Multiple Contributions to Plastid Physiology. Annu Rev Plant Biol **66**: 49–74

Nicol L, Croce R (2021) The PsbS protein and low pH are necessary and sufficient to induce quenching in the light-harvesting complex of plants LHCII. Sci Rep **11**: 7415

Niu Y, Lazár D, Holzwarth AR, Kramer DM, Matsubara S, Fiorani F, Poorter H, Schrey SD, Nedbal L (2023) Plants cope with fluctuating light by frequency-dependent nonphotochemical quenching and cyclic electron transport. New Phytologist **239**: 1869–1886

Niyogi KK, Grossman AR, Björkman O (1998) Arabidopsis Mutants Define a Central Role for the Xanthophyll Cycle in the Regulation of Photosynthetic Energy Conversion. The Plant Cell **10**: 1121–1134

Niyogi KK, Truong TB (2013) Evolution of flexible non-photochemical quenching mechanisms that regulate light harvesting in oxygenic photosynthesis. Current Opinion in Plant Biology **16**: 307–314

Ögren E (1988) Photoinhibition of photosynthesis in willow leaves under field conditions. Planta **175**: 229–236

Ögren E, Sjöström M (1990) Estimation of the effect of photoinhibition on the carbon gain in leaves of a willow canopy. Planta. doi: 10.1007/BF00193011

Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M, Niyogi KK (2009) An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. Nature **462**: 518–521

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, et al (2022) The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic Acids Research **50**: D543–D552

Ruban AV, Lee PJ, Wentworth M, Young AJ, Horton P (1999) Determination of the Stoichiometry and Strength of Binding of Xanthophylls to the Photosystem II Light Harvesting Complexes. Journal of Biological Chemistry **274**: 10458–10465

Sacharz J, Giovagnetti V, Ungerer P, Mastroianni G, Ruban AV (2017) The xanthophyll cycle affects reversible interactions between PsbS and light-harvesting complex II to control non-photochemical quenching. Nature Plants **3**: 16225

Santabarbara S, Garlaschi FM, Zucchelli G, Jennings RC (1999) The effect of excited state population in Photosystem II on the photoinhibition-induced changes in chlorophyll fluorescence parameters. Biochimica et Biophysica Acta **1409**: 165–170

Santabarbara S, Neverov KV, Garlaschi FM, Zucchelli G, Jennings RC (2001) Involvement of uncoupled antenna chlorophylls in photoinhibition in thylakoids. FEBS Letters **491**: 109–113

Schmid-Siegert E, Stepushenko O, Glauser G, Farmer EE (2016) Membranes as Structural Antioxidants. Journal of Biological Chemistry **291**: 13005–13013

Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M (2011) Global quantification of mammalian gene expression control. Nature **473**: 337–342

Suorsa M (2015) Cyclic electron flow provides acclimatory plasticity for the photosynthetic machinery under various environmental conditions and developmental stages. Front Plant Sci. doi: 10.3389/fpls.2015.00800

Takahashi H, Clowez S, Wollman F-A, Vallon O, Rappaport F (2013) Cyclic electron flow is redox-controlled but independent of state transition. Nat Commun **4**: 1954

Tian L, Nawrocki WJ, Liu X, Polukhina I, van Stokkum IHM, Croce R (2019) pH dependence, kinetics and light-harvesting regulation of nonphotochemical quenching in *Chlamydomonas*. Proc Natl Acad Sci USA **116**: 8320–8325

Treves H, Raanan H, Finkel OM, Berkowicz SM, Keren N, Shotland Y, Kaplan A (2013) A newly isolated *Chlorella* sp. from desert sand crusts exhibits a unique resistance to excess light intensity. FEMS Microbiol Ecol **86**: 373–380

Treves H, Raanan H, Kedem I, Murik O, Keren N, Zer H, Berkowicz SM, Giordano M, Norici A, Shotland Y, et al (2016) The mechanisms whereby the green alga *Chlorella ohadii* , isolated from desert soil crust, exhibits unparalleled photodamage resistance. New Phytol **210**: 1229–1243

Triantaphylidès C, Havaux M (2009) Singlet oxygen in plants: production, detoxification and signaling. Trends in Plant Science **14**: 219–228

Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J (2016) The Perseus computational platform for comprehensive analysis of (prote)omics data. Nat Methods **13**: 731–740

Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis, 2nd ed. doi: 10.1007/978-3-319-24277-4

Wilson S, Clarke CD, Carbajal MA, Buccafusca R, Fleck RA, Daskalakis V, Ruban AV (2024) Hydrophobic Mismatch in the Thylakoid Membrane Regulates Photosynthetic Light Harvesting. Journal of the American Chemical Society. doi: https://doi.org/10.1021/jacs.4c05220

Wolff Ch, Witt HT (1969) On Metastable States of Carotenoids in Primary Events of Photosynthesis: — Registration by repetitive ultra-short-flash photometry —. Zeitschrift für Naturforschung B **24**: 1031–1037

Xu P, Tian L, Kloz M, Croce R (2015) Molecular insights into Zeaxanthin-dependent quenching in higher plants. Sci Rep **5**: 13679

Figures and figure legends

Figure 1. Adaptation to high light conditions in *Chlorella ohadii* as revealed by mass spectrometry. **A)** Label-free quantification (LFQ) Liquid chromatography-tandem mass spectrometry (LC-MS/MS) in low and high light (LL and HL)-grown cells reveals the accumulation of known enzymatic antioxidants and CBR in HL-grown cells. Note the substantial reduction in LHCII subunits in HL-grown *C. ohadii*. **B)** List of the detected antioxidants and other related enzymes. Values in the 'LFQ HL/LL' column represent the fold-change value of a given protein in HL over LL-grown cells. Values in the 'Percentile in HL (iBAQ)' column represent the abundance of the protein within HL-grown cells. A value of 10 means this protein is amongst the 10% most abundant proteins in these cells. **C)** Detected peptides with oxidized tryptophan as a mark for oxidative damage. Note the accumulation of almost all detected oxidized peptides in LL-grown cells.

SOD – Superoxide dismutase. GPX – Glutathione peroxidase. CBR – Carotenoid biosynthesis related. Ω3FAD – omega 3 fatty acid desaturase. GR – Glutathione reductase. DHAR - Dehydroascorbate reductase. FLV – Flavodiiron protein. APX – Ascorbate peroxidase. GST – Glutathione S transferase. PTOX – Plastid/Plastoquinol terminal oxidase.

Figure 2. High light-grown *C. ohadii* drastically decreases PSII antenna size. **A)** Thylakoid membranes from low and high-light (LL and HL) grown cells were solubilized with detergents and the photosynthetic complexes were separated by a sucrose density gradient. Samples containing the same amount of chlorophyll (200 μg) were loaded on the gradients. Note the accumulation of a PSII core, stripped from LHCII, in HL-grown cells. **B)** The kinetics of Q_A reduction upon exposure to actinic light in the presence of DCMU was measured with a fluorometer and is directly correlated to the PSII antenna size. Y(II) values (Φ_{PSII}) were determined under the same experimental conditions, without adding DCMU. Fluorescence data was normalized to maximum ($F_M = 1$) and minimum ($F_0 = 0$) values for each sample. The maximal photochemical rate of PSII at each light intensity was calculated by taking the reciprocal of the area delimited by t0, the fluorescence rise curve, and the value of 1 (i.e. F_M) on the ordinate axis. μE refers to light intensity m^{-2} s^{-1}. **C)** Comparison of the slopes of linear regression fits across the measured rate points indicates an approximately 40% decrease in PSII absorption capacity in high-light-grown *C. ohadii*. **D)** Electron transport rates of photosystem II (ETRII) were normalized according to the calculated antenna size and tested under various light intensities. PAR – Photosynthetic active radiation, in μmol photons m^{-2} s^{-1}. HL-grown *C. ohadii* ETRII reaches peak values at higher light intensities due to its truncated photosystem II antenna.

Figure 3. *Chlorella ohadii* exhibits virtually no protective NPQ **A) Left:** Fluorescence traces of dark-adapted LL (top) and HL (bottom) *C. ohadii* grown cells, measured under 2000 µmol photons m^{-2}s^{-1}. Black and yellow colors at the line below the figure indicate dark and actinic light conditions. The red line indicates the time points when saturating pulses were employed to induce maximum fluorescence (F_M and F_M'). Low NPQ was detected as indicated by the little change in F_M' in response to exposure to high light. **Right:** Fluorescence traces of dark-adapted high light grown *stt7-9 C. reinhardtii* cells, exposed to 2000 µmol photons m^{-2}s^{-1}. Note the high NPQ levels. **B)** NPQ values were calculated from the fluorescence traces of dark-adapted LL and HL cells as shown in (A). Down-pointing arrows indicate time of turning off of the actinic light. **C)** NPQ is triggered by high light and acidification of the lumen in *C. reinhardtii* but not in *C. ohadii.* **Left:** Fluorescence traces of dark-adapted LL (top) and HL (bottom) *C. ohadii* grown cells, measured under 2000 µmol photons m^{-2}s^{-1} or after the acidification of the media to pH 5.5. Black arrows indicate the point of acid addition. **Right:** Fluorescence traces of dark-adapted *stt7-9 C. reinhardtii* cells, exposed to acidic conditions. KOH was added to the cell following 200 sec with no actinic light. The acidification of the medium induced high NPQ levels, mimicking the high light activation of NPQ. This is reversed upon adding KOH, mimicking the light-off action (compare the left panels of **A** and **C**). Fluorescence was normalized between to F_M in the dark-adapted cells for all samples.

Figure 4. Decreased $^1O_2$ accumulation in high-light adapted *Chlorella ohadii*. **A)** Confocal micros-copy wide-field vision of low and high light (LL and HL)-grown cells. Their intrinsic chlorophyll fluorescence, showed on the left panels, was used to localize the chloroplasts. Note the decreased fluorescence of the HL-grown cells. In the middle and right panels, the fluorescence of the $^1O_2$ marker SOSG is shown, originating from the same chloroplasts that are shown in the left panels. The SOSG fluorescence has been detected before (T0) and after 10 min (T10) of high red-light treatment. **B)** Quantification of $^1O_2$ SOSG signal of the chloroplasts shown in A. Only the SOSG fluorescence that overlaps the intrinsic chlorophyll fluorescent marking the chloroplast has been counted for the quantification (see Supplemented Figure 4). **C)** Violin plots of $^1O_2$ formation fol-lowing 10 min of illumination with high-intensity red light in 54 and 24 individual chloroplasts of LL and HL-grown cells, respectively. The internal point indicates the mean value and the error bars the standard error (SE). The mean numbers are: LL T0: 223, LL T10: 723, HL T0: 204, HL T10: 358. Note the log scale of the y-axis. The statistical analysis was done as described in the methods section. **D)** Accumulation of O_2^- and H_2O_2 in cells after 10 min of strong light illumination. The formation in HL cells was calculated as a percentage of the formation in LL cells (See methods). **E)** $^1O_2$ accumulation in isolated LHCII complexes after 10 min of strong light treatment was meas-ured by quantifying SOSG fluorescence.

Figure 5. Kinetics of $^1O_2$ production in *Chlorella ohadii*. **A)** Low and high-light (LL and HL)-grown cells were placed in a medium containing the $^1O_2$ marker SOSG in the confocal microscope and chlorophyll fluorescence was detected to locate the chloroplasts in the cells (chl). The cells were then illuminated with strong red light for the times indicated at the bottom and the accumulated $^1O_2$ was detected by the fluorescence intensity of SOSG. **B)** Quantification of the SOSG accumulation in 54 and 24 individual chloroplasts from LL and HL-grown cells, respectively, as shown in panel A. Statistical analysis was performed with a linear mixed model of the log10 transformed response with light as fixed effect and time as both fixed and random effect. Shadowed area represents +/- 2*SE. Mixed model ANOVA showed a significant effect of time (p= 0.0027, table x1) and a significant interaction between light and time (p = 3.215E-06, table x1). Post hoc comparisons between treatments in each time point resulted in significant differences in time points 4,6,8,10 (table x2).

Figure 6. P_{700} redox measurements suggest a strong increase of cyclic electron flow efficiency upon high light acclimation of *C. ohadii*. **A)** In high light (HL)-grown *C. ohadii* the light-harvesting complex II (LHCII) is removed from photosystem II (PSII) while photosystem I (PSI) maintains all or most of its antenna subunits. As a result, PSI is expected to absorb higher quantities of light energy. **B)** P700 oxidation kinetics analysis in dark-adapted cells shows that in the presence of DCMU which blocks linear electron flow (LEF), PSI of HL-grown cells is oxidized at a slower rate compared to that of LL-grown cells in all tested light intensities. When cyclic electron flow (CEF) is blocked by DBMIB, similar PSI oxidation rates are observed in both growth conditions. Note that in LL-grown cells, CEF becomes apparent only by high-light treatment, while in HL-grown cells, CEF is highly activated at all light intensities, and LEF is not activated at very low light intensities. 6 mM methyl viologen was added to all samples. **C)** PGR5, PGRL1, and Fd accumulate in HL-grown cells while FNR accumulates in LL-grown cells, as measured with label-free quantification (LFQ) Liquid chromatography-tandem mass spectrometry (LC-MS/MS).

Figure 7. *Chlorella ohadii* does not require NPQ for photoprotection. **A)** Under low light (LL) conditions, linear electron flow is favored and small amounts of reactive oxygen species (ROS) are generated. **B)** Upon a shift to high light (HL) conditions, large amounts of ROS are produced in the photosystem II (PSII) and by its antenna the light-harvesting complex II (LHCII). Cyclic electron flow (CEF) around PSI is relatively low. **C)** To avoid light-induced oxidative damage, HL-grown *C. ohadii* eliminates the LHCII, massively reducing the absorption of excess light energy. Additionally, carotenoids, superoxide dismutase (SOD), glutathione peroxidase (GPX), and possibly the carotenoid biosynthesis-related (CBR) protein and other enzymatic antioxidants act as ROS scavengers. CBR may quench light energy that is absorbed by the detached LHCII, before their degradation. CEF is highly active – at least in the absence of LEF - and may provide extra ATP for metabolic processes.

802   **Supplemental information**

Supplementary Figure 1. *Chlorella ohadii* lacks NPQ under strict photoautotrophic conditions. **A)** Fluorescence traces of dark-adapted LL (top) and HL (bottom) cells grown in strict photoautotrophic conditions, measured under 2000 μmol photons $m^{-2} s^{-1}$. Negligible NPQ was detected as indicated by the little change in maximal fluorescence (F_M) in LL and no change in HL in response to exposure to the pulses of high light. **B)** NPQ and Y(II) values were calculated from fluorescence signals as shown in panel *A*.

LL-grown cells   HL-grown cells

3000 µmol photons m⁻²s⁻¹

LL   2000 µmol photons m⁻²s⁻¹   HL

LL   400 µmol photons m⁻²s⁻¹   HL

LL   50 µmol photons m⁻²s⁻¹   HL

Supplementary Figure 2. *Chlorella ohadii* lacks NPQ under low or high light intensities. Fluorescence traces of dark-adapted low and high light (LL, left, and HL, right)-grown cells. The fluorescence measurement was carried out under different light intensities, as indicated in the Methods section. Low NPQ was detected as indicated by the little change in F_M' in response to exposure to low or high light.

Supplementary Figure 3. NPQ is not induced by high light or acidifying conditions in *Chlorella ohadii* grown under strict photoautotrophic conditions. Fluorescence traces of dark-adapted low and high light (LL, left, and HL, right)-grown cells. First under high light exposure (2000 μmol photons $m^{-2} s^{-1}$), and then quenching was detected in response to the acidification of the media to pH 5.5 with acetic acid. Blue arrows indicate turning the light on/off. Black arrows indicate the point of acid addition. The numbers above each FM signal indicate the NPQ values which are very small compared to *C. reinhardtii* where they routinely attain 2.2-2.5 (see Figure 3).

Supplementary Figure 4. $^1O_2$ detection with singlet oxygen sensor green (SOSG) probe workflow. **A)** SOSG was introduced to LL and HL cells by incubation for 30 min at the light intensity of 100 μmol photons m^{-2} sec^{-1}, with gentle shaking. **B)** The cells were pelleted and resuspended in a fresh TAP medium and loaded onto a microscope slide and applied to a spinning disk confocal microscope with a Z stacking of 8 μm around the cell central plain. **C)** Induction of high light stress was initiated within the microscope using red LED illumination. This wavelength selection (~600 nm) was chosen to specifically activate photosynthesis. **D)** SOSG fluorescence (excitation at 488 nm/emission at 530 nm) was recorded. To locate the chloroplast, chlorophyll fluorescence (excitation at 430 nm/emission at 680 nm) was concurrently recorded. **E)** Chlorophyll fluorescence was detected in different focused plains before and after the red high light treatment and a 3D model of the fluorescence was created by Z-stacking in the IMARIS program. Due to the limited permeability of SOSG, green fluorescence appeared on the cell membrane throughout the experiments. To subtract the access fluorescence that was not contributed by the photosynthetic-driven $^1O_2$ formation, a chloroplast volume was constructed from the chlorophyll fluorescence. **F)** SOSG fluorescence was only calculated from the chloroplast volume. This approach ensures the isolation of photosynthetic singlet oxygen formation, offering a comprehensive insight into the dynamics of singlet oxygen accumulation within the cellular context.