Senescent fibroblasts support tumor growth by inducing immune cell death in humanized models

Monica Cruz[1,2], Joshua Dulong[1,2], Anthony Sonn[1,2], Gaël Moquin-Beaudry[3], Basma Benabdallah[1], Oanh Le[1], Christian Beauséjour[1,2, *]

[1]Centre de recherche du CHU Sainte-Justine, Montréal, Québec, Canada

[2]Département de pharmacologie et physiologie Université de Montréal, Montréal, Québec, Canada

[3]Institut Gustave Roussy Paris, France

Financial Support: This work was supported by a project grant from the Terry Fox Foundation and by the Cancer Research Society (CRP-159385). M.C. is supported by a fellowship from le Fonds de recherche Santé Québec and J.D. is supported partially by a post-doctoral fellowship from the Cole Foundation

Running title: Senescent fibroblasts can kill immune cells

* **Correspondence**:
Christian Beauséjour PhD
CHU Ste-Justine, 3175 Côte Sainte-Catherine
Montréal, Québec, Canada, H3T 1C5
(Phone) 514-345-4931 x4385,
e-mail: c.beausejour@umontreal.ca

ABSTRACT

Syngeneic mouse models revealed that senescence influences the tumor immune response in many ways by, for example, inducing an immunosuppressive microenvironment or by favoring immune cell recruitment. Yet, the impact of senescence on the tumor immune response in a humanized setting is mostly unknown. Here, using tumor spheroids and humanized mice bearing tumors immunogenic to autologous human immune cells, we show that senescent fibroblasts exert a dual impact on the tumor immune response by attracting more immune cells but also by inducing the death of T and NK cells. Mechanistically, we demonstrate that this was mostly the result of increased Fas ligand (FasL) expression at the surface of senescent fibroblasts and that the deletion of FasL was sufficient to prevent immune cell death and increase tumor cell killing. Our results highlight the importance of evaluating the impact of therapy-induced senescence in humanized models to better understand and predict the outcome of cancer treatments.

INTRODUCTION

Cancer treatments can induce senescence of cancer cells and other cell types within the tumor microenvironment (TME) [1]. This phenotype is believed to have mostly detrimental consequences, as senescent cells were shown to play a significant role in the development and progression of cancer [2,3]. Indeed, Krtolica and colleagues were the first to demonstrate that senescent human fibroblasts favor tumor growth in immune-deficient mice [2]. The accumulation of senescent fibroblasts was also shown to contribute to local and systemic inflammation, promoting adverse effects and cancer relapse [4]. These effects are believed to be mediated mainly by the senescence-associated secretion phenotype (SASP), which includes several factors linked to inflammation and malignancy [5,6].

Several studies using syngeneic tumor models in immune-competent mice showed that the SASP can prevent tumor rejection by inducing an immunosuppressive TME [7-10]. For example, prolonged exposure to SASP factors such as type 1 interferon and IL-6 were shown to downregulate immune cell functions [11-13] and to interfere with the tumor immune response mediated by T and NK cells [7-9]. On the other hand, other studies showed that the SASP can stimulate and recruit immune cells [14-17]. Hence, while the SASP can undoubtedly promote tumor growth, its impact on immune cells is variable and was shown to depend on the tumor model.

Several immune cell populations impact tumor growth, including cytotoxic T cells, whose intra-tumoral presence correlates with a survival benefit [18]. However, the tumor immune response is known to be significantly diminished by myeloid-derived suppressor cells (MDSC) and other immune-inhibitory cells or molecules within the TME [19,20]. One of

these inhibitory molecule is FasL, which can trigger apoptosis of Fas-expressing effector cells such as T cells and natural killer (NK) cells [21]. FasL is expressed by several cell types, including tumor cells, MDSC [20,21], tumor-associated macrophages, regulatory T cells [21], and cancer-associated fibroblasts [22]. Elevated FasL levels in solid tumors correlate with disease progression, increased metastasis, poor overall survival [23], and decreased numbers of CD8[+] infiltrating T cells [24].

Nevertheless, the impact of senescence on the tumor immune response in a humanized setting remains undetermined. This is because, until recently, there was no adequate humanized tumor model where human cancer cells could be rejected or their growth delayed in the presence of immune cells. Such models are essential to evaluate whether senescent cells interfere with the tumor immune response and if so, through which mechanism. To this end, we recently developed iPSC-derived humanized mouse models where tumors can be rejected following the injection of autologous human immune cells [25]. Using these models, we show here that senescent human dermal fibroblasts (HDF) have a dual impact on the tumor immune response. On the one hand, senescent HDF attract immune cells and on the other hand, they induce their death. Mechanistically, we demonstrate that this is mostly the result of increased FasL expression at the surface of senescent HDF. By providing a better understanding of the complex interactions between senescent stromal cells and immune cells, our results should help develop better-targeted therapeutic interventions.

RESULTS

Senescent fibroblasts promote tumor growth and stimulate the infiltration of immune cells in humanized mice.

To determine the role of a senescent stroma on the tumor-immune response, we first induced senescence of HDF by exposure to ionizing radiation (IR) or the expression of K-RAS^{V12}, two proven methods that we and others routinely use (**Supplementary Fig. 1A and B**) [26,27]. We then injected pre-senescent or senescent HDF together with tumor cells into the flanks of NSG-SGM3 mice (ratio of 1:4). Here, we used two tumor cell lines (HEPA-4T and LEC-4T derived from hepatic and lung iPSC-derived progenitor cells respectively) which we previously showed are immunogenic to autologous immune cells [25]. The day after tumor inoculation, mice were injected intraperitoneally with human immune cells (PBMCs and granulocytes, 5×10^6 of each). Note that tumor cells, HDF, and immune cells were all derived from the same donor to evaluate the impact of senescence in a humanized autologous setting (**Fig. 1A**). HEPA-4T and LEC-4T tumor cells were transduced to express mPlum and HDF were stained with the NIR790 cytoplasmic membrane dye, allowing the visualization of all cell types, and tracking of tumor growth by quantitative *in vivo* imaging (**Fig. 1B**).

As expected, the co-injection of tumor cells with pre-senescent or senescent HDF resulted in accelerated tumor growth compared to mice injected with tumor cells alone (**Fig. 1C and 1D**). The injection of immune cells was able to delay tumor growth in all groups. Surprisingly, the presence of HDF did not seem to interfere with the tumor rejection, except for RAS-induced HDF, which slightly hampered the rejection of LEC-4T tumors. Indeed, for this group, tumor size was reduced only by about 45% in the presence

of immune cells compared to 60-80% in the other groups (**Fig. 1C**). These results were confirmed at the time of sacrifice when residual tumors were excised and weighted (**Fig. 1E and 1F**). *In vivo*, imaging of NIR790-labeled HDF over four weeks following the injection of immune cells revealed that most HDF persists over time, suggesting they are not the target of immune cells (**Supplementary Fig. 1C and D**). Overall, these results demonstrate that senescent HDF have either no effect or only a moderate effect on the tumor immune rejection in these humanized models.

We were expecting that senescent HDF, through a combination of their mitogenic effect and immunosuppressive features, would interfere with tumor rejection. Hence, to better understand the impact of senescence, we characterized the infiltration of human immune cells (hCD45$^+$) in dissociated tumors at the time of sacrifice. Cell counts obtained by flow cytometry and normalized to the tumor weight showed that tumors injected with HDF, especially senescent HDF, tended to have an increased infiltration of hCD45$^+$ immune cells (**Fig. 2A and 2C**). However, while the ratio of tumor-infiltrating immune cell types was slightly different between the two tumor cell lines, it was not significantly changed between groups and was composed mostly of CD4$^+$ and CD8$^+$ T cells (**Fig. 2B and 2D**). Because residual tumors were of different sizes at the time of sacrifice, which likely impacted infiltration, we then measured immune cell infiltration using a more defined spheroid model (**Fig. 2E**). Hence, LEC-4T and HEPA-4T cell lines expressing mPlum were cultured in 3D spheres with or without pre-senescent or senescent HDF expressing GFP (**Fig. 2F**). Spheroids were co-cultured for 24 hours with PBMCs, after which the localization of infiltrated immune cells into the spheroids was assessed by immunofluorescence. Images showed an increase in the number of infiltrating immune

cells at 24 hours in the presence of senescent HDF, particularly towards the core of the spheroids (**Fig. 2G**). To better quantify the infiltrated cells, we repeated the experiment, but this time, we dissociated the spheres with trypsin and confirmed the increase infiltration of immune cells by flow cytometry (**Fig. 2H**). These observations led us to hypothesize that senescent HDF may have opposing effects as they can induce the recruitment of immune cells yet without enhancing tumor rejection (**Fig. 1C and 1D**).

Senescent fibroblasts trigger apoptosis of human immune cells through upregulation of FasL.

To better evaluate the effects of senescent HDF on immune cells, we used a model where PBMCs collected from three distinct donors were co-cultured for up to 72 hours with pre-senescent or senescent HDF. Immune cells were then counted either by flow cytometry or alternatively, immune cell death was assessed with propidium iodide (PI) staining and quantified using live-cell imaging (**Fig. 3A**). For all donors, we found a substantial decrease in the absolute number of live CD45$^+$ in contact with senescent HDF, mostly RAS-induced senescent HDF (**Fig. 3B**). Such a decrease in cell viability was primarily due to dying CD3$^+$ T cells and to a lesser extent CD3$^-$CD56$^+$ NK cells although the effect was donor dependant (**Fig. 3B**). This ability of senescent HDF to kill immune cells was also confirmed by counting dying (PI positive) immune cells using live-cell imaging (**Fig. 3C and 3D**).

We next wanted to investigate the mechanism leading to immune cell death. Among the pathways we investigated, we found that senescent HDF consistently increased FasL expression independently of the inducer (**Fig. 3E**). Such an increase in

the proportion of cells expressing FasL was also accompanied by a significant two and five-fold increase in the mean fluorescence intensity in IR/doxorubicin and RAS-induced senescent HDF respectively (**Fig. 3E**). Similar results were observed using other human fibroblast cell lines (**Supplementary Fig. 2**). To confirm the involvement of FasL in immune cell killing, we generated FasL knockout (KO) HDF using CRISPR-Cas9 and confirmed loss of FasL expression by flow cytometry in the absence or presence of IFN-γ which increases FasL expression in wild-type cells (**Fig. 3F and 3G**). We then performed co-culture studies using HDF and human PBMCs, and we observed that immune cells survival was not compromised in contact with FasL KO HDF as determined by counting the number of PI+ cells per image using the IncuCyte software (**Fig. 3H**). Increased immune cell survival in contact with FasL KO senescent HDF was also confirmed using flow cytometry (**Fig. 3I**). Staining for Annexin V showed that immune cell death was induced mainly by apoptosis, especially by RAS-induced HDF (**Fig. 3J**). Overall, these results demonstrate that FasL expression is induced in senescent HDF and that knocking out FasL expression greatly diminishes the ability of senescent HDF to kill immune cells.

Therapy-induced senescence increases FasL expression in tumor spheroids and decreases immune cell survival.

We next wanted to evaluate if increased FasL expression would occur in the context of therapy-induced senescence and if this would negatively impact immune cell survival. Therefore, we generated tumor spheroids by mixing an equal number of HDF and A549 tumor cells (**Fig. 4A**). We chose to work with the A549 lung epithelial carcinoma cell line for this experiment because A549 cells adopt a senescence phenotype when exposed to

IR or doxorubicin. Indeed, we observed that A549 spheroids increased SA-β-gal activity and significantly reduced their growth upon treatments (**Fig. 4B and C**). Immunofluorescence imaging performed on cryosections showed an increase in FasL expression in response to treatments in tumor spheroids (**Fig. 4D**). Given the relatively low number of HDF in our spheres at day 11 (day 6 post treatment), such an increase in FasL expression was surprising. It is possible FasL expression is increased in a 3D environment or that it arises from the infiltrated immune cells. Such increased expression is unlikely to arise from tumor cells as IR or doxorubicin-induced senescent A549 show only a modest increase in FasL expression in culture as detected by flow cytometry or by immunofluorescence in spheroids grown in absence of HDF and immune cells (**Fig. 4E and Supplementary Fig. 3**). Nonetheless, when we analyzed the proportion of live immune cells (CD45[+], PI[-]) in dissociated spheroids we observed significantly less viable immune cells in the context of therapy-induced senescence (**Fig. 4F**).

The SASP is sufficient to induce immune cell death and to promote tumor cells survival *in vitro*.

The SASP allows for a crosstalk between senescent cells and neighboring cells. Therefore, we next wanted to determine if the SASP alone could induce immune cell death, knowing that FasL can be found soluble [28,29]. We observed that soluble FasL was indeed present in the conditioned media (CM) of RAS-induced senescent HDF but not in any other conditions, including in CM from IR-induced senescent HDF (**Fig. 5A**). Therefore, to determine the impact of the SASP we incubated immune cells in the presence of CM for 48 hours and counted the number of viable immune cells (CD45[+] and

CD3$^+$) by flow cytometry. In accordance with the results of the ELISA, we found a significant decrease in the number of viable immune cells in the presence of CM collected from RAS-induced-senescence HDF (**Fig. 5B**). However, we also observed a slight decrease in cell survival in the presence of CM collected from IR-induced senescent HDF, suggesting that the SASP is toxic to immune cells independently of soluble FasL, although at a lower extent (**Fig. 5B**).

We also observed that the SASP had immunosuppressive properties as it prevented the proliferation of CFSE-labelled immune cells in a mixed lymphocyte reaction. For this experiment, PBMCs from two unrelated donors were mixed in the presence or absence of CM collected from pre-senescent or senescent HDF, and the proliferation of T cells was analyzed by flow cytometry. The results showed that the SASP can interfere with the proliferation of T cells (**Supplementary Fig. 4**). Considering these observations, we reasoned that increased immune cell killing combined with their reduced proliferation should lead to reduced killing of tumor cells. To test this hypothesis, we incubated PBMCs with LEC-4T tumor cells (expressing mPlum) in the absence or presence of CM for 72 hours. Using live-cell imaging, we tracked tumor cell growth and apoptosis (Caspase 3/7 specific dye in green). As expected, images showed less apoptosis of tumor cells and a consequent increase in their number in the presence of CM collected from senescent HDF (**Fig. 5C-E**). In agreement with soluble FasL being present only in CM collected from RAS-induced senescent HDF, tumor cell counts were higher in this group compared to IR-induced senescent HDF. Counting live tumor cells by flow cytometry confirmed the results (**Fig. 5F**). Together, this demonstrates that the SASP

can protect tumor cells either by killing immune cells or by limiting their proliferation using both FasL-dependent and independent mechanisms.

FasL KO senescent fibroblasts do not interfere with the tumor immune response in humanized mice.

Finally, we evaluated if tumors would be better rejected by immune cells in the absence of FasL expression from senescent HDF. To do so, we used LEC-4T tumor cells co-injected with RAS-induced senescent HDF, the only combination that impaired tumor-immune rejection (**Fig. 1C**). Hence, we co-injected subcutaneously LEC-4T cells with either WT or FasL KO RAS-induced senescent HDF in NSG-SGM3 mice (**Fig. 6A**). First, we confirmed that in the absence of immune cells, that wildtype or FasL KO senescent HDF had a similar mitogenic effect on tumor growth when compared to tumor cells injected alone (**Fig. 6B and 6C**). However, in the presence of immune cells (PBMCs and granulocytes), we observed that tumors containing FasL KO senescent HDF were rejected like tumors without HDF (**Fig. 6B and 6C**). Tumor weight at the time of sacrifice confirmed these results (**Fig. 6D**). However, the number and proportion of tumor-infiltrating immune cells were surprisingly not significantly changed in the absence of FasL expression (**Fig. 6E and 6F**). We speculate that this may be the consequence of the relatively small size of residual tumors in the FasL KO group at the time of sacrifice, which may not accurately represent the peak of immune cell infiltration during the tumor immune response.

DISCUSSION

Using immunogenic autologous tumors, we were able to uncover a dual role of senescence on the tumor immune response in humanized mice. Indeed, we observed that senescent HDF, aside from their mitogenic effect on tumor cells, have a mixed effect on the tumor immune response by first recruiting immune cells and then inducing their death through a mechanism that involves FasL. Importantly, we observed increased FasL expression on various HDF lines in response to different senescence inducers. Highlighting the importance of the tumor model, we found that subcutaneous A549 tumors were not rejected despite being infiltrated by immune cells at similar levels to those observed in HEPA-4T and LEC-4T tumors (**Supplementary Fig. 5**). The reasons for this are unknown, but it is likely that tumor cell lines derived from patients have evolved to evade the immune system and thus are not suitable to measure the possible detrimental impact of senescence.

Interestingly, soluble FasL was only found in the CM of RAS-induced senescent HDF. The reason for this may be that the release of soluble FasL requires the action of metalloproteinases (MMP) [29] and that RAS-induced senescent HDF expresses much higher MMP levels than IR-induced senescent HDF [3]. As MMP can also be highly expressed by cancer cells [30-32], it is possible that soluble FasL is released from all types of senescent cells in the TME. Soluble FasL has been reported to inhibit Fas-mediated apoptosis induced by cytotoxic T cells, suggesting that senescent HDF can protect tumor cells by preventing their Fas-mediated killing by T cells or by inducing apoptosis of immune cells directly [33]. Indeed, FasL has emerged as an important factor in the TME that can limit the anti-tumor immune response and the efficacy of immunotherapy by

inducing the apoptosis of T cells [34,35]. Consistent with these results, we found that ablating FasL expression on senescent HDF abrogated their ability to induce T and NK cells apoptosis and consequently enhanced tumor cell survival in vitro and tumor rejection in mice. Interestingly, Motz et al. reported that FasL expression on the tumor endothelium could protect tumor cells [36]. Overall, our data support a mechanism of tumor resistance where FasL expressed by senescent cells induces apoptosis of T and NK cells. Given the relatively limited number of immune cells injected in mice and the fact that most NK cells do not survive more than a few days in NSG mice in absence of appropriate cytokines (for example human IL-15), it is likely that tumor cells were mostly killed by T cells in mice. However, we cannot rule out the implication of NK cells, given we previously showed that the adoptive transfer of a limited number of NK cells in NSG mice is sufficient to reject iPSC-derived teratomas [37]. Moreover, we also observed that the killing of A549 tumor cells by NK cells *in vitro* is inhibited by the co-culture of wild type but not FasL KO HDF (**Supplementary Fig. 6**).

Other immune resistance mechanisms have been reported to be induced by senescence. For example, in mouse syngeneic models, senescent stromal cells can induce an immunosuppressive environment by favoring the formation of myeloid-derived suppressive cells [8]. In our study, it was not possible to verify if this mechanism of resistance is also induced in our tumors, given the short half-life of human granulocytes in humanized mice. Another limitation of our study is that we measured the impact of therapy-induced senescence in tumor spheroids and not in humanized mice. The reason for this is that it is difficult to control tumor growth *in vivo* in response to therapy and that our tumors would be contaminated with host fibroblasts and endothelial cells. More

experiments will be necessary to understand the extent and in which context the increased expression of FasL occurs. The relationship between the molecular pathways involved in senescence and the expression of FasL also remains to be elucidated. We speculate that cytotoxic stress and DNA-damaging agents that can trigger the activation of the c-Jun N-terminal kinase (JNK) pathway may be involved since this pathway was shown to regulate FasL expression [38]. Moreover, emerging evidence indicates that JNK signaling is involved in regulating cellular senescence [39]. Another possibility is that the nuclear factor-kappa B (NF-κB) transcription factor, which plays an important role in regulating the SASP, is involved as it was also shown to control FasL gene expression [40,41].

In summary, our results uncover a role for FasL during senescence and suggest it plays an important role in the tumor immune response. It will be interesting to determine the extent to which FasL expression is increased in the context of therapy-induced senescence in different types of human tumors and if targeting FasL can improve the efficiency of immunotherapy.

METHODS

Cell culture

Human dermal fibroblasts (HDF) from an adult donor were isolated as previously described [25,42]. Briefly, an adult skin biopsy was collected in accordance with the ethics committee from the Centre Hospitalier Universitaire (CHU) Sainte-Justine. The biopsy was cleaned out, cut into 1–5-mm^2 pieces, and digested with collagenase D (Roche) for 1 h at 37°C with agitation, centrifugated at 400 × g for 5 min, and washed with DMEM (Wisent Bio Products). HDF were maintained in DMEM with 10% FBS and 0.2% primocin. HDF senescence was induced by exposing cells to a 12 Gy dose of ionizing radiation (1 Gy/min using a Faxitron CP-160), or after their stable transduction with lentiviral particles carrying an inducible version of K-RAS, or after treatment with 0.1μM of Doxorubicin. Conditioned media (CM) was prepared from pre-senescent or senescent HDF cultured in monolayer with RPMI 1640 (Wisent Bio Products) without serum for 24 hours after which it was then collected, filtered, and centrifuged. HDF were counted at the time of collection, and the CM normalized to the lowest number of cells in a particular condition by dilution with fresh RPMI.

iPSC-derived tumor models

iPSCs reprogramming and cellular transformation were performed as previously described [25]. In brief, iPSC-derived hepatocyte progenitors (HEPA-4T) and iPSC-derived lung epithelial progenitors (LEC-4T) were generated and transduced using lentiviral particles carrying the SV40 large T antigen and the neomycin resistance gene. Three days later, 300 µg/ml G418 (Thermo Fisher Scientific) selection was applied. Surviving

cells were subsequently transduced with lentiviral particles carrying the H-RAS^{V12} and puromycin-resistance genes. Cells were selected for three days post-transduction with 2 µg/mL of puromycin (Thermo Fisher Scientific). Surviving cells were then transduced with lentiviral particles carrying the human telomerase gene. Finally, transformed cells were transduced to express the mPlum gene. All transductions were carried out using fresh culture media containing 8 µg/ml polybrene (Sigma-Aldrich).

Animals and solid tumor models

Animal experiments were performed under a protocol approved by the institutional committee for good laboratory practices for animal research (protocol #2022-3508). Nine-week-old female and male NSG-SGM3 (expressing human IL3, GM-CSF, and SCF) mice originally obtained from The Jackson Laboratory (Bar Harbor, ME) and bred at the animal care facility at the CHU Sainte-Justine Research Center were used. Mice were housed under strict specific pathogen-free conditions and handled with aseptic techniques under anesthesia (2% isoflurane) to inject tumor cells.

For the subcutaneous injections, 5×10^4 transformed cells (HEPA-4T or LEC-4T) alone or mixed with 2×10^5 pre-senescent or senescent HDF were diluted in 100 µl of RPMI 1640 (Wisent Bio Products) and implanted in the right and left flank of mice previously anesthetized and shaved. Tumor growth was monitored for four to five weeks using the Q-Lumi In Vivo imaging system (MediLumine, Montreal, QC, Canada) by fluorescent tracking of mPlum-expressing tumor cells (Ex. 562-40 nm and Em. 641-75 nm). Pre-senescent or senescent HDF were stained with a fluorescent dye before their injection in mice. In brief, cells were diluted at 1×10^6 cells/ml, and CellBrite NIR790 Cytoplasmic Membrane Dye (catalog 30079; Biotium) was added at the concentration of

1 µM and incubated at 37°C for 20 min. Cells were then washed twice in serum-free RPMI 1640 (Wisent Bio Products) before being resuspended in 100µL of cold serum-free RPMI 1640 (Wisent Bio Products) for injection in mice. HDF were tracked using near-infrared filters (excitation 769-41 nm and emission 832-37 nm). The tumor fluorescent signal was analyzed and normalized using Fiji macros for picture processing and expressed in fluorescence-integrated density or radiance (photons $\cdot$ s^{-1} $\cdot$ sr^{-1} $\cdot$ cm^{-2}) integrated density. For tumor growth analysis, mice were sacrificed when one limit point was reached according to our animal comity guidelines. Our comity established limit points as no more than 10% weight loss, no distress signs such as alopecia or decreasing activity and that tumor size does not reach more than 1500 mm3 or become ulcerated. Tumors were resected and weighed after sacrifice. Tumors were considered eliminated when they were unpalpable or too small to be harvested at sacrifice.

Mouse immune reconstitution and characterization of the tumor-immune infiltrate.

For the adoptive transfer of human immune cells, peripheral blood mononuclear cells (PBMCs) were isolated using the Ficoll-Paque gradient (GE Healthcare) from the blood of healthy donors after informed consent. Granulocytes were collected by lysing the red blood cells (RBC) by resuspending the Ficoll-Paque cell pellet in 19mL of sterile deionized water for 20 seconds before adding 1 ml of sterile 20× PBS solution. PBMCs and granulocytes were counted and mixed at a 1:1 ratio before being injected into mice; a total of 1×10^7 human immune cells (5×10^6 each) were injected i.p. with 200 µl of RPMI 1640 (Wisent Bio Products). Tumors were digested using the human Tumor Dissociation Kit and gentleMACS Octo Dissociation with Heaters (Miltenyi Biotec), according to the manufacturer's instructions. The cell suspension was filtered on 40 µm MACS

SmartStrainers (Miltenyi Biotec) and washed using RPMI 1640 (Wisent Bio Products) containing 10% FBS. Finally, resuspended cells were stained with the following antibodies to analyze the tumor-infiltrating immune cell: mouse CD45/PE/Cy7, human CD45/BUV395, human CD3/AF700, human CD19/PE-CF594, human CD4/BB515, human CD8/BV421, human CD14/APC/H7, human CD56/BV786, and human CD127/BB700, all from BD Biosciences except human CD33/BV510 and human CD25/BV711 who were purchased from Biolegend. All data were acquired on LSR Fortessa (BD Biosciences) and data analysis was done on FlowJo V10 (v10; Tree Star).

CRISPR-Cas9 knockout cell lines

FasL Knock-out (KO) HDF and A549 cells were generated by the Gene Editing Platform of the CHU Sainte-Justine Research Center using CRISPR-Cas-mediated genome editing. Briefly, a guide RNA (gRNA) targeting a DNA sequence within the first exon of the FasL gene (5'-CTGGGCACAGAGGTTGGACA-3') was cloned into the BsmBI restriction site of the 3^{rd} generation LentiCRISPR.V2 (Addgene, #52961) vector. Lentiviral particles were produced by transfecting LentiCRISPR.V2 (pLC-EF1a-SpCas9-U6-gRNA1-FasL 450 ng) together with the pMDL (750 ng), pREV (300 ng), and pVSV-G (390 ng) packaging plasmids in 5 x 10^5 HEK293T cells with Lipofectamine 2000 (Invitrogen). The supernatant was collected 40 hours later, centrifuged 5min at 3000 rpm on a Sorvall LegendMicro17 (ThermoFisher) to remove debris and then frozen at -80°C. FasL KO HDF were produced by transducing 5 x 10^5 cells with 100 µL of viral particles. After 24 hours, Puromycin (2 mg/mL) (Wisent) was added to select positive clones expressing Cas9 and the gRNA. 72 hours later, 10 cells were transferred into separate wells of a 96-

well plate and left to establish a positive clone mixture. PCR and sequencing were performed to characterize knock-out populations. Finally, ICE Analysis Software (Synthego) showed that 1 cell population no longer expressed FasL. The latter was composed of 37% and 63% of cells with indels of +1 bp and -11 bp, respectively. The loss of expression of FasL was confirmed by FACS analysis (LSR Fortessa) (BD Biosciences) using a FasL/PE conjugated antibody from BioLegend.

3D spheroids formation, invasion assay, and image analysis

Spheroids were formed by centrifugating 5000 tumor cells/well (LEC-4T, HEPA-4T, A549) alone or together with 5000 HDF/well (pre-senescent and senescent) for 5 min at 150g in ultra-low attachment 96-well plates and incubated at 37°C under 5% CO_2 for 6 days. Where indicated, 5×10^5 peripheral blood mononuclear cells (PBMCs) were added per well and 24 or 48 hours later spheroids were dissociated into single cells for subsequent flow cytometric analysis. Alternatively, spheroids were embedded in OCT, frozen and 12 µm sections were generated using a Leica cryostat. Sections were transferred to gelatinized slides and immunofluorescence staining was performed using a human CD45 antibody (Rat. YAML501.4; Thermo Fisher Scientific, USA), GFP Polyclonal Antibody (A-11122; Thermo Fisher Scientific, USA), FAS-L Antibody (NOK-1. sc-19681; Santa Cruz Biotechnology), and DAPI to counterstain DNA.

3D tumor spheroids growth after therapy-induced senescence

Spheroids (A549 mPlum cells alone or together with GFP-expressing HDF) were allowed to form for five days after which they were treated with 0.1µM of doxorubicin or a single dose of 15 Gy IR. Spheroids size was monitored using the IncuCyte S3 (Essen

Bioscience) through 4X images taken every 3 hours for a total of 96 hours. Likewise, some spheroids were fixed with 4% formaldehyde and stained for SA-β-gal activity.

Immunophenotyping of HDF by Flow Cytometry

HDF were harvested, re-suspended in phosphate-buffered saline (PBS, Wisent Bio Products), and stained with the LIVE/DEAD™ Fixable Far-Red Dead Cell Stain Kit (Thermo Fisher Scientific) for 15 min on ice. Cells were then washed twice and re-suspended in flow cytometry staining buffer (BD Biosciences) and incubated for 30 min at 4°C with 1 µL of PE anti-human FasL antibody (clone NOK-1) from BioLegend. Nonspecific background signals were measured by incubating cells with the appropriate isotype-matched antibody. Unstained, viability dye only, and single-stained compensation beads (Invitrogen) served as controls. Doublets were gated out using forward-scatter width/height and sideward-scatter width/height event characteristics. All stained cells were analyzed using an LSR Fortessa cytometer (BD Bioscience), and the obtained results were analyzed using FlowJo v10 (v10; Tree Star).

Detection of apoptosis

To detect apoptotic cells, we used the IncuCyte S3 (Essen Bioscience) live cell imaging system in combination with propidium iodide (PI) or the caspase 3/7 specific activity dye. In brief, PBMCs (3×10^5) were co-culture with pre-senescent or senescent HDF (3×10^4 cells) in a 48-well plate. Alternatively, PBMCs were cultured in the presence of CM collected from HDF in a 48-well plate. PI (25µg) and IL-2 (150 IU/mL) were added to the media and cell viability was monitored for 72 hours. Cells were treated with 7 µM of camptothecin as a positive control for cell death. Dead cells were counted via red object

counts at indicated time points, and data was analyzed using the IncuCyte software. To measure apoptosis of tumor cells we used the caspase 3/7 activity dye. Tumor cells (1×10^4) were seeded in a 24-well plate and treated with 200 ng/mL IFNγ overnight. The next day, PBMCs were added at a ratio of 1:20 (tumor cells: PBMCs) in the presence or absence of CM collected from pre-senescent or senescent HDF and the Incucyte® green Caspase-3/7 activity reagent (Sartorius, Ann Arbor, MI, USA). Plates were analyzed at a two-hour interval for a total of 72 hours.

Apoptosis of PBMCs was also measured by flow cytometry using the Apotracker(TM) Green (Biolegend) apoptosis probe in combination with PI (Sigma Aldrich). Both floating and adherent cells were collected after 3 days of co-culture, washed twice with PBS, and stained with Apotracker(TM) Green following the manufacturer's protocol. Cells were stained with the following antibodies: human CD45/BUV395, human CD3/AF700, human CD8/BV421, and human CD56/BV786 from BioLegend and then immediately analyzed using the LSR Fortessa (BD Biosciences). Cells were considered alive when not stained with PI or Apotracker, while apoptotic cells were defined as those stained with Apotracker. Absolute counting beads (Invitrogen) were used to normalize FACS event acquisition and to calculate absolute cell numbers.

NK cell culture conditions and cytotoxicity assay

NK cells were isolated from PBMCs of healthy donors by depleting CD3 cells using an EasySep human CD3 positive selection kit II (StemCell Technologies). The negative cell fraction was expanded using irradiated (100 Gy) K562 mbIL21 4-1BBL for a minimum of 2 weeks in RPMI1640 supplemented with 10% FBS, 1% penicillin/streptomycin (Wisent Bio Products), and 100 UI/mL IL-2 (SteriMax) until the proportion of NK cells (CD3⁻,

CD56$^+$) reached close to 100% as determined by flow cytometry. Cells were rested for one week after their expansion prior to be used in a cytotoxicity assay. In brief, A549 mPlum cells were co-cultured with GFP-expressing HDF (pre-senescent or senescent) at a 1:2 ratio in a 96 well plate for 72 hours. NK cells were then added at different effector: target ratios in presence of 20 UI/mL IL-2. Cell death was determined by counting cells by flow cytometry. Before the acquisition, the supernatant was gently removed, and cells were stained with Zombie NIR™ Fixable Viability Kit (BioLegend) according to the manufacturer's instructions and then detached using trypsin. Cytotoxicity was assessed by flow cytometry and was calculated as Lysis (%) = [1-live targets (sample)/live targets (control)] × 100%. Cytotoxicity against A549 was performed using NK cells isolated from the same donor in presence of pre-senescent or senescent HDF. A total of three different donors of NK cells were used.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8.0 (GraphPad Software, Inc., San Diego, CA). Data are represented as mean ± SEM. For comparisons of three or more groups, data were subjected to t-tests or one-way ANOVA analysis, followed by Dunnett's multiple comparisons test when comparing every mean to a control mean or Tukey's multiple comparisons test when comparing every mean to every other mean. The tumor growth kinetics was analyzed using mixed-effects, followed by Tukey's multiple comparison test. *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$, and ****$p < 0.0001$.

COMPETING INTERESTS:

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS:

M.C., J. D., A.S., performed the experiments; G.M.B., and B.B., provided study material;

O.L., provided technical guidance; C.B., designed and supervised the study; M.C., and

C.B., wrote the manuscript with the contributions from all the authors.

REFERENCES

D'Ambrosio, M. & Gil, J. Reshaping of the tumor microenvironment by cellular senescence: An opportunity for senotherapies. *Dev Cell* **58**, 1007-1021 (2023). https://doi.org:10.1016/j.devcel.2023.05.010

Krtolica, A., Parrinello, S., Lockett, S., Desprez, P. Y. & Campisi, J. Senescent fibroblasts promote epithelial cell growth and tumorigenesis: a link between cancer and aging. *Proc Natl Acad Sci U S A* **98**, 12072-12077 (2001). https://doi.org:10.1073/pnas.211053698

Muñoz, D. P. *et al.* Targetable mechanisms driving immunoevasion of persistent senescent cells link chemotherapy-resistant cancer to aging. *JCI Insight* **5** (2019). https://doi.org:10.1172/jci.insight.124716

Demaria, M. *et al.* Cellular Senescence Promotes Adverse Effects of Chemotherapy and Cancer Relapse. *Cancer Discov* **7**, 165-176 (2017). https://doi.org:10.1158/2159-8290.CD-16-0241

Coppe, J. P. *et al.* Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol* **6**, 2853-2868 (2008). https://doi.org:08-PLBI-RA-2566 [pii]10.1371/journal.pbio.0060301

Omer, A. *et al.* G3BP1 controls the senescence-associated secretome and its impact on cancer progression. *Nat Commun* **11**, 4979 (2020). https://doi.org:10.1038/s41467-020-18734-9

Eggert, T. *et al.* Distinct Functions of Senescence-Associated Immune Responses in Liver Tumor Surveillance and Tumor Progression. *Cancer Cell* **30**, 533-547 (2016). https://doi.org:10.1016/j.ccell.2016.09.003

Ruhland, M. K. *et al.* Stromal senescence establishes an immunosuppressive microenvironment that drives tumorigenesis. *Nat Commun* **7**, 11762 (2016). https://doi.org:10.1038/ncomms11762

Toso, A. *et al.* Enhancing chemotherapy efficacy in Pten-deficient prostate tumors by activating the senescence-associated antitumor immunity. *Cell reports* **9**, 75-89 (2014). https://doi.org:10.1016/j.celrep.2014.08.044

Ye, J. *et al.* Senescent CAFs Mediate Immunosuppression and Drive Breast Cancer Progression. *Cancer Discov*, Of1-of22 (2024). https://doi.org:10.1158/2159-8290.Cd-23-0426

Harker, J. A., Dolgoter, A. & Zuniga, E. I. Cell-intrinsic IL-27 and gp130 cytokine receptor signaling regulates virus-specific CD4(+) T cell responses and viral control during chronic infection. *Immunity* **39**, 548-559 (2013). https://doi.org:10.1016/j.immuni.2013.08.010

Cunningham, C. R. *et al.* Type I and Type II Interferon Coordinately Regulate Suppressive Dendritic Cell Fate and Function during Viral Persistence. *PLoS Pathog* **12**, e1005356 (2016). https://doi.org:10.1371/journal.ppat.1005356

Wilson, E. B. *et al.* Blockade of chronic type I interferon signaling to control persistent LCMV infection. *Science* **340**, 202-207 (2013). https://doi.org:10.1126/science.1235208

Kansara, M. *et al.* Immune response to RB1-regulated senescence limits radiation-induced osteosarcoma formation. *J Clin Invest* **123**, 5351-5360 (2013). https://doi.org:10.1172/JCI70559

Ruscetti, M. *et al.* NK cell-mediated cytotoxicity contributes to tumor control by a cytostatic drug combination. *Science* **362**, 1416-1422 (2018). https://doi.org:10.1126/science.aas9090

Kang, T. W. *et al.* Senescence surveillance of pre-malignant hepatocytes limits liver cancer development. *Nature* **479**, 547-551 (2011). https://doi.org:10.1038/nature10599

Xue, W. *et al.* Senescence and tumour clearance is triggered by p53 restoration in murine liver carcinomas. *Nature* **445**, 656-660 (2007).

Gooden, M. J., de Bock, G. H., Leffers, N., Daemen, T. & Nijman, H. W. The prognostic influence of tumour-infiltrating lymphocytes in cancer: a systematic review with meta-analysis. *Br J Cancer* **105**, 93-103 (2011). https://doi.org:10.1038/bjc.2011.189

Hou, A. J., Chen, L. C. & Chen, Y. Y. Navigating CAR-T cells through the solid-tumour microenvironment. *Nat Rev Drug Discov* **20**, 531-550 (2021). https://doi.org:10.1038/s41573-021-00189-2

Zhu, J. *et al.* Resistance to cancer immunotherapy mediated by apoptosis of tumor-infiltrating lymphocytes. *Nature Communications* **8**, 1404 (2017). https://doi.org:10.1038/s41467-017-00784-1

Bajgain, P. *et al.* Secreted Fas Decoys Enhance the Antitumor Activity of Engineered and Bystander T Cells in Fas Ligand-Expressing Solid Tumors. *Cancer Immunol Res* **10**, 1370-1385 (2022). https://doi.org:10.1158/2326-6066.Cir-22-0115

Lakins, M. A., Ghorani, E., Munir, H., Martins, C. P. & Shields, J. D. Cancer-associated fibroblasts induce antigen-specific deletion of CD8+T Cells to protect tumour cells. *Nature Communications* **9**, 948 (2018). https://doi.org:10.1038/s41467-018-03347-0

Peter, M. E. *et al.* The role of CD95 and CD95 ligand in cancer. *Cell Death Differ* **22**, 549-559 (2015). https://doi.org:10.1038/cdd.2015.3

Zietz, C., Rumpler, U., Stürzl, M. & Löhrs, U. Inverse relation of Fas-ligand and tumor-infiltrating lymphocytes in angiosarcoma: indications of apoptotic tumor counterattack. *Am J Pathol* **159**, 963-970 (2001). https://doi.org:10.1016/s0002-9440(10)61772-5

Moquin-Beaudry, G. *et al.* Autologous humanized mouse models of iPSC-derived tumors enable characterization and modulation of cancer-immune cell interactions. *Cell Rep Methods* **2**, 100153 (2022). https://doi.org:10.1016/j.crmeth.2021.100153

Goyer, M.-L. *et al.* Induced Pluripotent Stem Cell-Derived Fibroblasts Efficiently Engage Senescence Pathways but Show Increased Sensitivity to Stress Inducers. *Cells* **13**, 849 (2024).

Wang, B., Kohli, J. & Demaria, M. Senescent Cells in Cancer Therapy: Friends or Foes? *Trends Cancer* **6**, 838-857 (2020). https://doi.org:10.1016/j.trecan.2020.05.004

Tanaka, M., Suda, T., Takahashi, T. & Nagata, S. Expression of the functional soluble form of human fas ligand in activated lymphocytes. *Embo j* **14**, 1129-1135 (1995). https://doi.org:10.1002/j.1460-2075.1995.tb07096.x

Kayagaki, N. *et al.* Metalloproteinase-mediated release of human Fas ligand. *J Exp Med* **182**, 1777-1783 (1995). https://doi.org:10.1084/jem.182.6.1777

Köhrmann, A., Kammerer, U., Kapp, M., Dietl, J. & Anacker, J. Expression of matrix metalloproteinases (MMPs) in primary human breast cancer and breast cancer cell lines: New findings and review of the literature. *BMC Cancer* **9**, 188 (2009). https://doi.org:10.1186/1471-2407-9-188

Forget, M. A., Desrosiers, R. R. & Béliveau, R. Physiological roles of matrix metalloproteinases: implications for tumor growth and metastasis. *Can J Physiol Pharmacol* **77**, 465-480 (1999).

Curran, S. *et al.* Matrix metalloproteinase/tissue inhibitors of matrix metalloproteinase phenotype identifies poor prognosis colorectal cancers. *Clin Cancer Res* **10**, 8229-8234 (2004). https://doi.org:10.1158/1078-0432.Ccr-04-0424

Yamada, A., Arakaki, R., Saito, M., Kudo, Y. & Ishimaru, N. Dual Role of Fas/FasL-Mediated Signal in Peripheral Immune Tolerance. *Front Immunol* **8**, 403 (2017). https://doi.org:10.3389/fimmu.2017.00403

Pitt, J. M. *et al.* Targeting the tumor microenvironment: removing obstruction to anticancer immune responses and immunotherapy. *Ann Oncol* **27**, 1482-1492 (2016). https://doi.org:10.1093/annonc/mdw168

Zhu, J., Petit, P. F. & Van den Eynde, B. J. Apoptosis of tumor-infiltrating T lymphocytes: a new immune checkpoint mechanism. *Cancer Immunol Immunother* **68**, 835-847 (2019). https://doi.org:10.1007/s00262-018-2269-y

Motz, G. T. *et al.* Tumor endothelium FasL establishes a selective immune barrier promoting tolerance in tumors. *Nat Med* **20**, 607-615 (2014). https://doi.org:10.1038/nm.3541

Benabdallah, B. *et al.* Natural Killer Cells Prevent the Formation of Teratomas Derived From Human Induced Pluripotent Stem Cells. *Front Immunol* **10**, 2580 (2019). https://doi.org:10.3389/fimmu.2019.02580

Dhanasekaran, D. N. & Reddy, E. P. JNK signaling in apoptosis. *Oncogene* **27**, 6245-6251 (2008). https://doi.org:10.1038/onc.2008.301

Deng, Y. *et al.* c-Jun N-terminal kinase signaling in cellular senescence. *Arch Toxicol* **97**, 2089-2109 (2023). https://doi.org:10.1007/s00204-023-03540-1

Kavurma, M. M. & Khachigian, L. M. Signaling and transcriptional control of Fas ligand gene expression. *Cell Death & Differentiation* **10**, 36-44 (2003). https://doi.org:10.1038/sj.cdd.4401179

Chien, Y. *et al.* Control of the senescence-associated secretory phenotype by NF-κB promotes senescence and enhances chemosensitivity. *Genes Dev* **25**, 2125-2136 (2011). https://doi.org:10.1101/gad.17276711

Moquin-Beaudry, G. *et al.* The Tumor-Immune Response Is Not Compromised by Mesenchymal Stromal Cells in Humanized Mice. *J Immunol* **203**, 2735-2745 (2019). https://doi.org:10.4049/jimmunol.1900807

Figure 1. **Senescent fibroblasts promote tumor growth without compromising the tumor immune response in humanized mice**.

A) Schematic of the *in vivo* experimental design. NSG-SGM3 mice were subcutaneously injected with LEC-4T or HEPA-4T tumor cells (5×10^4) and pre-senescent or senescent HDF (2×10^5 cells) on day 0. The next day, mice were injected intraperitoneally with PBMCs (5×10^6) and granulocytes (5×10^6). Mice were sacrificed on day 32 and tumors were collected for analysis.

B) Representative images of LEC-4T tumors on day 22 expressing mPlum and HDF stained with the NIR790 dye after their injection.

C) Growth curves for LEC-4T tumors injected alone (in black n=12) or co-injected with pre-senescent HDF (in red n=12-14), senescent HDF induced by irradiation (in blue n=12-14) or induced by RAS (in green n=14-18) in mice without (solid line) and with (dashed line) autologous immune cells. Each line represents the mean tumor growth (± SEM) over 32 days. Statistical analyses were performed using a mixed-effects model, followed by Tukey's multiple comparison test.

D) Growth curves for HEPA-4T tumors injected alone (in black n=18-20) or co-injected with pre-senescent HDF (in red n=18-20), senescent HDF induced by irradiation (in blue n=18-20) or induced by RAS (in green n=18-20) in mice without (solid line) and with (dashed line) autologous immune cells. Each line represents the mean tumor growth ± SEM over 28 days. Statistical analyses were performed using a mixed-effects model, followed by Tukey's multiple comparison test.

E and F) Graphs representing the weights of LEC-4T or HEPA-4T tumors collected at sacrifice. Each dot represents an individual tumor. Values represent the mean ± SEM. A one-way ANOVA with Dunnett's multiple comparisons tests was used to determine statistical significance.

Figure 2. Cruz et al.

Figure 2. **Senescent fibroblasts increase immune cell infiltration in spheroids and in tumors.**

A) Graph showing the absolute counts of tumor-infiltrating human CD45[+] cells per gram of dissociated LEC-4T tumors as determined by flow cytometry. Each dot represents the count from a single tumor. Shown is the mean ± SEM. Statistical analysis between groups was performed with one-way ANOVA followed by Dunnett's multiple comparisons tests.

B) Stacked bar graph showing the proportion of immune cell subset populations in LEC-4T tumors (the same as in panel A) from each group. Shown is the mean ± SEM.

C) Graph showing the absolute counts of tumor-infiltrating human CD45[+] cells per gram of dissociated HEPA-4T tumors as determined by flow cytometry. Each dot represents the count from a single tumor. Shown is the mean ± SEM. Statistical analysis between groups was performed with one-way ANOVA Dunnett's multiple comparisons tests.

D) Stacked bar graph showing the proportion of immune subset cell populations in HEPA-4T tumors (the same as in panel C) from each group. Shown is the mean ± SEM.

E) Schematic representation of the 3D tumor spheroid invasion model. In brief, HDF were exposed or not to IR or RAS and 10 days later, an equal number of tumor cells (LEC-4T or HEPA-4T) were mixed with pre-senescent or senescent HDF (5000 cells each) to form mixed-cell spheroids. Once the spheroid assembled and matured for 6 days, 5×10^5 human PBMCs were added and the infiltration of immune cells into the spheroid was quantified 24 hours later by flow cytometry.

F) Representative images (at 4X) of spheroids at day 16 before adding immune cells. Tumor cells are shown in red (mPlum), HDF in green (EGFP) and in yellow the merge of the two signals.

G) Graphs showing the number of human immune cells infiltrated in spheroids as determined by flow cytometry (CD45[+]) from each indicated group. Each dot represents the count from a single dissociated sphere obtained in the course of three independent experiments. Statistical analysis between groups was performed with one-way ANOVA with Dunnett's multiple comparisons tests.

H) Representative images of immunostained sections from the indicated spheroids showing infiltrated human immune cells (CD45[+] in red) and nuclei (DAPI in blue). The scale bar represents 100 µm.

Figure 3. Cruz et al.

Figure 3. **Senescent fibroblasts induce immune cell death through up-regulation of FasL**.

A) Schematic representation of the experimental design. In brief, HDF were exposed to IR or RAS and 3 days later cells (3×10^4) were seeded in a 48-well plate. An equal number of non-senescent HDF were seeded only on day 7 and the day after PBMCs (3×10^5) were added. Cell death as determined by propidium iodide (PI$^-$) incorporation tracked over 72 hours at 2-hour intervals using an IncuCyte live-cell analysis system or alternatively, absolute cell counts were determined by flow cytometry at 72 hours.

B) Graphs showing the proportion of live CD45$^+$, CD3$^+$, and NK cells (CD3$^-$CD56$^+$) after 72 hours of coculture with senescent HDF normalized to counts obtained in presence of pre-senescent HDF. Data are presented as mean ± SEM from three independent experiments using PBMCs from three different donors. Statistical analysis between groups was performed, followed by one-way ANOVA with Dunnett's multiple comparisons tests.

C) Representative images captured by the IncuCyte using a 10X objective showing PI$^+$ (in red) dying cells at 72 hours. The scale bar represents 400 µm.

D) Quantification of PI$^+$ dying immune cells from each group. CPT was added as a positive control. Data are presented as mean ± SEM from three independent experiments. Statistical analysis between groups was performed with one-way ANOVA followed by Dunnett's multiple comparisons tests.

E) Quantification by flow cytometry of the proportion and the mean fluorescence intensity (MFI) of pre-senescent and senescent HDF expressing FasL at their surface. The MFI was calculated by subtracting the fluorescence of the isotype control. Data are presented as mean ± SEM from three independent experiments. Statistical analysis between groups was performed with one-way ANOVA with Dunnett's multiple comparisons tests.

F) FasL expression as detected by flow cytometry on wild type (WT) and FasL KO HDF either pre-senescent (PRE in red) or senescent (IR in blue or RAS in green). Also shown are non-stained cells (in gray) and cells stained with the isotype controls (in black). Cells were stimulated or not with IFN-γ (200ng/mL). Representative histograms of three independent experiments are shown.

G) Bar graphs showing the proportion of cells expressing FasL in the different HDF populations presented in panel F. Data are presented as mean ± SEM from three independent experiments. Statistical analysis between groups was performed with one-way ANOVA with Dunnett's multiple comparisons tests.

H) Quantification of PI$^+$ dying immune cells after 72 hours of co-culture with the indicated HDF populations. Data was acquired with the IncuCyte® live-cell analysis system. CPT was added as a positive control. Data are presented as mean ± SEM from three independent experiments. The p-value was calculated by multiple t-tests.

I) Quantification of live (Annexin V$^-$/PI$^-$) CD45$^+$ cells after 72 hours of co-culture with the indicated HDF populations as determined by flow cytometry. CPT was used as a positive control. Data are presented as mean ± SEM from three independent experiments. Statistical analysis was calculated by multiple t-tests.

J) Quantification of apoptotic (Annexin V$^+$/PI$^+$) CD45$^+$ cells after 72 hours of co-culture with the indicated HDF populations as determined by flow cytometry. CPT was used as a positive control. Data are presented as mean ± SEM from three independent experiments. Statistical significance was calculated by multiple t-tests.

Figure 4. Cruz et al.

Figure 4. Therapy-induced senescence in tumor spheroids impairs the viability of infiltrated immune cells.

A) Schematic representation of the 3D tumor spheroid model. In brief, A549 tumor cells alone or with an equal number of HDF (5000 cells for each) were mixed to form monospheroids or mixed-cell spheroids, respectively. Five days after, spheroids were treated with doxorubicin (0.1µM) or irradiated (15 Gy) to induce senescence. Then, four days later PBMCc (5×10^5) were added and their infiltration into the spheroid was quantified 48 hours later by flow cytometry.

B) Representative images of spheroids stained for β-galactosidase activity six days after being treated with doxorubicin or irradiation. Unstained spheroids or spheroids not exposed to therapy were used as controls. The scale bar represents 100 µm.

C) Graph showing the average size of spheroids represented by the largest brightfield object area metric (μm^2) over time following treatments as detected by IncuCyte imaging. Shown is the mean ± SEM of three independent experiments. Statistical differences were identified by mixed-effects modeling with Tukey's multiple comparisons.

D) Representative images of sections of immunostained spheroids showing the infiltration of CD45$^+$ (in white), the presence of HDF (GFP in green), expression of FasL (in red), and cell nuclei (DAPI in blue). The scale bar represents 100 µm.

E) Quantification by flow cytometry of the proportion of A549 tumor cells expressing FasL when grown for 48 hours in co-culture with pre-senescent or senescent HDF (left panel). Also shown is the mean fluorescence intensity (MFI – right panel). The MFI was calculated by subtracting the fluorescence of the isotype control. Statistical analysis between groups was performed with one-way ANOVA with Dunnett's multiple comparisons tests.

F) Graph showing the number of live (Annexin V$^-$/PI$^-$) CD45$^+$ immune cells infiltrated in spheroids from the indicated groups. Each dot represents the average of infiltrated cells in n=6 spheroids collected from independent experiments. Shown is the mean ± SEM. Statistical analysis between groups was performed by a one-way ANOVA with Tukey's multiple comparisons.

Figure 5. The SASP triggers apoptosis of immune cells and promote tumor cells survival in vitro

A) Concentration of soluble FasL (sFasL) as detected by ELISA in the conditioned media (CM) of pre-senescent and senescent HDF populations. Each dot represents an independent experiment (n=3), and t-tests determined statistical significance between groups.

B) Graphs showing the absolute counts of CD45$^+$ (left panel) and CD3$^+$ (right panel) live cells as determined by flow cytometry after 48 hours of culture in CM collected from the indicated HDF populations. Shown is the mean ± SEM from three independent experiments. Multiple t-tests determined statistical significance between groups.

C) Representative images of LEC-4T tumor cells (expressing mPlum in red) after 72 hours co-cultured with PBMCs in CM collected from the indicated HDF populations. The Caspase-3/7 Green reagent was added to detect cells undergoing apoptosis. Scale bars represent 400 µm**.**

D) Scatter dot plot showing the number of caspase 3/7 positive tumor cells (red and green overlap) after 72 hours of co-culture as determined using the IncuCyte® live-cell analysis software. Shown is the mean ± SEM from four independent experiments. Multiple t-tests determined statistical significance between groups.

E) Scatter dot plot showing the number of LEC-4T tumor cells (in red) after 48 hours of co-culture as determined using the IncuCyte® live-cell analysis software. Shown is the mean ± SEM from four independent experiments. Multiple t-tests determined statistical significance between groups.

F) Absolute counts of live LEC-4T cells as measured by flow cytometry at the end of the 72-hours co-culture period. Each dot represents the average of five technical replicates. Shown is the mean ± SEM from five independent experiments. Multiple t-tests determined statistical significance between groups.

Figure 6. Cruz et al.

Figure 6. **FasL KO senescent fibroblasts do not interfere with the tumor immune response in humanized mice.**

A) Schematic of the in vivo experimental design. NSG-SGM3 mice were subcutaneously injected with LEC-4T tumor cells (5 x 10^4) and pre-senescent or senescent HDF (2 x 10^5 cells) on day 0. The next day, mice were injected intraperitoneally with PBMCs (5 x 10^6) and granulocytes (5 x 10^6). Mice were sacrificed on day 32 and tumors were collected for analysis.

B) Representative images of mice bearing subcutaneous LEC-4T tumors (expressing mPlum) alone or co-injected with the indicated HDF populations stained with the NIR790 dye after their injection. Images from day 1 show HDF-stained with the NIR790 dye while days 15 and 32 show mPlum tumor growth.

C) Growth curves for LEC-4T tumors in humanized mice injected with LEC-4T cells alone (n=12) or co-injected with RAS-induced senescent HDF (n=18) or RAS-induced senescent HDF KO for Fas-L (n=16). Also shown is the growth curve of LEC-4T tumors in NSG-SGM3 mice co-injected with RAS-induced senescent HDF (n=14) or RAS-induced senescent HDF KO for Fas-L (n=6). Each line represents the mean tumor growth (±SEM) over 32 days. Statistical analyses were performed using a mixed-effects model, followed by Tukey's multiple comparison test.

D) Shown is the weight of LEC-4T tumors collected at sacrifice on day 32. Each dot represents the weight of an individual tumor. Values represent the mean ± SEM Statistical analysis between groups was performed by t-test or one-way ANOVA with Tukey's multiple comparisons.

E) Bar graph showing the number of tumor-infiltrating hCD45[+] immune cells per gram of tumor collected from both groups at the time of sacrifice. Each dot represents the infiltration in an individual residual tumor large enough to be excised and dissociated (n=6). Values represent the mean ± SEM, and a t-test was performed for statistical analysis.

F) Stacked bar graph showing the proportion of immune cell subset populations in LEC-4T tumors collected from the indicated groups. Shown is the mean ± SEM