Genetic and Environmental Contributions to Epigenetic Aging Across Adolescence and

Young Adulthood

Dmitry V. Kuznetsov[1*], Yixuan Liu[1], Alicia M. Schowe[2,3], Darina Czamara[2], Jana Instinske[4],

Charlotte K. L. Pahnke[5], Markus M. Nöthen[5], Frank M. Spinath[6], Elisabeth B. Binder[2], Martin

Diewald[1], Andreas J. Forstner[5,7], Christian Kandler[4], and Bastian Mönkediek[1]

[1]Bielefeld University, Germany

[2]Max Planck Institute of Psychiatry, Department Genes and Environment, Munich, Germany

[3]Graduate School of Systemic Neuroscience, Ludwig Maximilian University, Munich, Germany

[4]University of Bremen, Germany

[5]Institute of Human Genetics, University of Bonn, School of Medicine & University Hospital

Bonn, Bonn, Germany

[6]Department of Psychology, Saarland University, Saarbrücken, Germany

[7]Institute of Neuroscience and Medicine (INM-1), Research Center Jülich, Germany

*Corresponding author, Dmitry V. Kuznetsov, dmitry.kuznetsov@uni-bielefeld.de

Abstract

Background

Epigenetic aging estimators commonly track chronological and biological aging,

quantifying its accumulation (i.e., epigenetic age acceleration) or speed (i.e., epigenetic aging

pace). Their scores reflect a combination of inherent biological programming and the impact of

environmental factors, which are suggested to vary at different life stages. The transition from

adolescence to adulthood is an important period in this regard, marked by an increasing and,

then, stabilizing epigenetic aging variance. Whether this pattern arises from environmental

influences or genetic factors is still uncertain.

This study delves into understanding the genetic and environmental contributions to

variance in epigenetic aging across these developmental stages. Using twin modeling, we

analyzed four estimators of epigenetic aging, namely Horvath Acceleration, PedBE Acceleration,

GrimAge Acceleration, and DunedinPACE, based on saliva samples collected at two timepoints

approximately 2.5 years apart from 976 twins of four birth cohorts (aged about 9.5, 15.5, 21.5,

and 27.5 years at first and 12, 18, 24, and 30 years at second measurement occasion).

Results

Half to two-thirds (50–68%) of the differences in epigenetic aging were due to unique

environmental factors, indicating the role of life experiences and epigenetic drift, besides

measurement error. The remaining variance was explained by genetic (Horvath Acceleration:

24%; GrimAge Acceleration: 32%; DunedinPACE: 47%) and shared environmental factors

(Horvath Acceleration: 26%; PedBE Acceleration: 47%). The genetic and shared environmental

factors represented the primary sources of stable differences in corresponding epigenetic aging

estimators over 2.5 years.

Age moderation analyses revealed that the variance due to individually-unique

environmental sources was smaller in younger than in older cohorts in epigenetic aging

estimators trained on chronological age (Horvath Acceleration: 47% to 49%; PedBE

Acceleration: 33% to 68%). The variance due to genetic contributions, in turn, potentially

increased across age groups for epigenetic aging estimators trained in adult samples (Horvath

Acceleration: 18% to 39%; GrimAge Acceleration: 24% to 43%; DunedinPACE: 42% to 57%).

Conclusions

Transition to adulthood is a period of the increasing variance in epigenetic aging. Both

environmental and genetic factors contribute to this trend. The degree of environmental and

genetic contributions can be partially explained by the design of epigenetic aging estimators.

Keywords Epigenetic aging, Age-moderation, Adolescence, Gene☐environment

interaction, Horvath clock, PedBE clock, GrimAge, DunedinPACE.

Background

Estimators of epigenetic aging offer insights into how individuals age over time and

across age groups [28]. Their increased estimates are associated with age-related diseases and

mortality in adults [43, 57], and with worse cognitive functioning [58] and more socioeconomic

disadvantage in adolescents and children [59]. By measuring DNA methylation (DNAm) levels

at specific DNA loci, epigenetic aging estimators address, along universal, specific aspects of

aging processes.

Understanding epigenetic aging measures and the aging processes underlying them

requires considering their training models with a variable number of CpGs, training markers, and

age ranges. The Horvath clock, for example, was trained to predict chronological age in samples

with a large age range. It considers the discrepancy between chronological and epigenetic age to

indicate accelerated or decelerated aging across the whole life span [27]. Similarly, in buccal

samples of children and adolescents, the pediatric-buccal-epigenetic (PedBE) clock illuminates

the specific aging processes in younger age groups [47]. Alternatively, epigenetic aging

estimators have been developed on the basis of links between aging and biological health

indicators. Two examples of such measures are the biomarker of human mortality risk GrimAge

[40, 41] and the biomarkers of the pace of aging, DunedinPoAm and DunedinPACE [4, 5].

These measures account for DNAm at locus-specific sites associated, for example, with plasma

proteins and smoking (GrimAge) or biomarkers of organ systems integrity, such as blood

pressure or serum leptin levels (Dunedin). In addition, Dunedin measures are trained on

longitudinal data and represent, in contrast to the degree of epigenetic aging, the pace of

epigenetic aging [4, 5].

The strength of the prediction by epigenetic aging measures stems from their ability to

represent both genetically driven aging processes and the cumulative effects of the environment

on DNAm [35]. The contributions of genetic factors to the variance in epigenetic aging can be

reflected in the narrow- and broad-sense *heritability,* the proportion of the phenotypic variance

attributable to genetic differences at the time and age of measurement in a certain population.

Narrow-sense heritability quantifies only the contribution of additive genetic factors, whereas

broad-sense heritability also includes the interactions between genetic factors such as allelic

dominance (i.e., interactions within specific gene loci of chromosomes) and epistasis (i.e.,

interactions across genetic variants between gene loci).

Figure 1 summarizes previously reported estimates of SNP-[1], pedigree-[2], and twin-based[3]

heritability for the aforementioned epigenetic aging measures (see also Additional file 1: Table

S1). As shown in Figure 1, the narrow-sense heritability of the Horvath clock measured in blood

is not constant across particular ages. According to previous studies, heritability estimates vary

from 39% to 61% in older adults, which is on average lower than the estimates obtained for

young adults, ranging from 60% to 77% [26, 27, 28, 33, 48, 66]. Further, prior research revealed

a lower heritability of the Horvath clock in adolescent than in young adult samples, with a range

from 37% to 43% [43, 67]. However, the estimates in the adolescent samples were derived from

pedigree- and SNP-based studies, whereas the estimates in the young adult samples were derived

from twin studies, which typically result in higher heritability estimates. Given that the twin-

---

[1] The proportion of the variance explained by all single nucleotide polymorphisms (SNPs) used in a genome-wide association study in conventionally unrelated individuals [78].

[2] The proportion of the variance explained by the patterns of genetic resemblance among not-too-distantly related family members [24].

[3] The proportion of the variance explained by the patterns of genetic resemblance between monozygotic and dizygotic twins.

based heritability of the Horvath clock appears to peak at birth [27], the lower heritability

estimates for adolescence may potentially represent missing heritability besides the age-

associated pattern [78].

The possibility of extrapolating the potential age-associated trend in the heritability of the

Horvath clock to other epigenetic aging measures has just started to be explored. For example,

heritability comparisons between epigenetic aging estimators are confined mainly to emerging

adults [34] or older individuals [26]. In the sample of emerging adults, the Horvath clock,

GrimAge, and pace of aging demonstrated similar heritability estimates of 62–73% [34]. Later in

life, the Horvath clock, in comparison to GrimAge and DunedinPACE, demonstrated higher

estimates until age 66 [26]. Knowledge on the possible comparability of heritability estimates

between tissues and resources of DNAm is mainly limited to the comparison of blood tissue and

postmortem brain, with lower estimates for the latter (Additional file 1: Table S1). Studying the

age-associated patterns of heritability across a broader age range and spectrum of epigenetic

aging measures that vary in training ages and phenotypes, therefore, would allow to elucidate

trends that are common and specific to certain epigenetic aging measures. Furthermore, DNAm

based on saliva can be assumed to correlate with DNAm based on both blood and buccal cells

[9]. The examination of the age-associated patterns of heritability based on DNAm derived from

noninvasive saliva samples, therefore, may also be indicative of cross-tissue accuracy and tissue-

specific differences in heritability.

The trend of epigenetic aging heritability across particular ages may arise from

environmental sources and their interaction with genetic factors. A design of twins reared

together allows to disentangle environmental factors shared by twins, which increase twin

siblings' similarity, from non-shared or individually-unique environmental factors, which

increase twin siblings' dissimilarity in DNAm. Particularly in younger age, when twins live

together in one household, shared environments may be more important. These environments

can encompass, for instance, prenatal factors, social backgrounds, common parental and peer

influences, and shared neighborhoods. Research on complex traits suggests that shared

environmental and genetic factors may interact, particularly in younger individuals [6, 32]. With

respect to the variance in epigenetic aging, the role of shared environmental factors was found to

increase before adulthood when twins lived together [37]. Individually-unique environmental

factors are assumed to be important across the whole life span and can obtain greater importance

when twins live in different households and have more life experiences of their own. These

environments can include non-shared perinatal factors, stochastic changes in life circumstances,

and individual-specific experiences, such as different peer groups or differences in experiences

of objectively shared environments. Individually-unique environmental factors have been shown

to moderate the influences of genetic factors on DNAm [22] and possibly have the same effect

on DNAm-based epigenetic aging during adulthood [30]. These shifts in genetic and

environmental contributions to the variance in epigenetic aging over the life course suggest a

possible link between life stage processes and gene□environment interplay in epigenetic aging.

Adolescence and the transition to adulthood, so-called emerging adulthood, are potential

periods of dynamic genetic and environmental contributions to the variance of epigenetic aging.

The time before transition is a period of growth and development with a faster change rate in

DNAm and epigenetic age [1, 28]. Studies have shown, for instance, that this change rate peaks

in infancy, decreases nonlinearly until approximately age 20, and then stabilizes [27, 28].

Variance in epigenetic aging can also undergo changes. Starting small at birth, the variance in

the epigenetic age of the Horvath clock increases with chronological age until adulthood onset,

and then stabilizes [37]. It remains unclear whether these trends of later stabilization in change

rates correspond to trends in the relative contributions of genetic and non-genetic factors and

which sources, genetic or environmental, are responsible for the growing and stabilizing variance

of epigenetic aging during adolescence and the transition to adulthood.

To address these gaps in the epigenetic literature, this study aimed to investigate the

trends in genetic and environmental contributions to the variance in epigenetic aging across

adolescence, emerging, and young adulthood. With our design, which includes four twin birth

cohorts and saliva-based DNAm from two measurement occasions, it is possible to examine how

genetic and environmental sources contribute to the variance in epigenetic aging and how these

sources shift or remain stable across age groups and over time (approximately two and a half

years in this study). DNAm was obtained from DNA extracted from saliva samples of early

adolescent (M(age) = 10.7 years), late adolescent (M(age) = 16.6 years), emerging adult (M(age)

= 22.7 years), and young adult (M(age) = 28.8 years) twins. The epigenetic aging estimators

included two acceleration measures trained on chronological age, the Horvath clock [27] and the

PedBE clock [47], and two epigenetic aging measures trained on biological indicators of aging,

one of which, GrimAge [40], represents biological age acceleration and the other,

DunedinPACE, represents the pace of biological aging [5].

With the use of these epigenetic aging estimators, we were able to answer the following

research questions: (a) Do genetic and environmental contributions to the variance differ

between differently developed epigenetic aging estimators? (b) To what extent do genetic and

environmental contributions account for stable differences in epigenetic aging estimators across

two measurement occasions? (c) Are there age-associated trends in the genetic and

environmental contributions to the variance in epigenetic aging from early adolescence to young

adulthood and do they differ for different epigenetic aging estimators? In addition, the findings

allow comparisons of heritability estimates of epigenetic aging in young adults assessed using

saliva samples with those that have previously been reported in other studies based on blood

samples (see Figure 1 and Additional file 1: Table S1).

Figure 1

The SNP-, pedigree-, and twin-based heritability of epigenetic aging estimated in previous

studies

Note. The estimates of SNP-, pedigree-, and twin-based heritability measured in blood DNAm

are presented for Horvath Acceleration (purple), GrimAge Acceleration (green), and epigenetic

aging pace measures, DunedinPoAm and DunedinPACE (yellow).

Methods

Sample

The research questions were examined in a sample of 976 twins from the TwinLife

Epigenetic Change Satellite (TECS) project, which is a subsample of the German twin family

panel TwinLife. TwinLife is a longitudinal study of four cohorts of same-sex twin pairs and their

families recruited as a representative sample from the general population of Germany [15]. The

study focuses on longitudinal measures of social inequality and the potential underlying

biopsychosocial mechanisms [25]. The current analysis sample consisted of 263 monozygotic

(MZ) and 225 dizygotic (DZ) complete twin pairs aged 8 to 29 years at the time of the first

epigenetic measurement ($M = 16.0$) and aged 10 to 31 years at the second measurement ($M =$

18.3). Zygosity was confirmed by genotyping. The age difference between cohorts was

approximately 6 years. At the time of assessment, members of the four cohorts were in the ages

of early adolescence, late adolescence, emerging adulthood, and young adulthood, respectively

(for more details see Table 1).

DNA methylation

Saliva sampling and DNA extraction

Saliva sampling for DNA extraction was conducted as a part of two TwinLife satellite

projects, namely TwinSNPs and TECS. Twins and their family members were invited to

participate in the molecular genetic analysis during TwinLife face-to-face interviews carried out

in 2018–2020. During face-to-face interviews, interviewers offered to participate in the

molecular genetic studies, which involved genotyping and epigenetic profiling. A short video

explaining the study goals and procedures of the saliva sampling accompanied the recruitment

procedure. The second saliva collection took place in 2021. This time, participants were

contacted by mail. In addition to the invitation to participate, respondents received consent

forms, saliva collection toolkits, and corresponding instructions. For all participants who

provided valid consent, saliva samples were sent to the researchers at the University Hospital

Bonn, Germany. The samples were collected using Oragene® saliva self-collection kits (OG-500

and OG-600; DNA Genotek, Canada) and DNA was extracted following the manufacturer's

protocol.

DNA methylation assessment

The assessment of DNA methylation profiles was conducted in a set of 2 128 samples, of

which the present sample constituted a subset. The samples were randomized by age, sex,

zygosity, and complete versus single twin pair across 23 plates (266 arrays) using the R package

"Omixer" (version 1.4.0; [68]). The family identifier was used as a blocking variable to ensure

that all samples and timepoints of a twin pair ran on the same batch/array. Bisulfite-conversion

of 500 ng of DNA was performed using the D5033 EZ-96 DNA Methylation-Lightning Kit

(Deep-Well; Zymo Research Corp., Irvine, USA), and methylation profiles were assessed using

the Infinium MethylationEPIC BeadChip v1.0 (Illumina, San Diego, CA, USA).

214   Preprocessing of the methylation samples ($n = 2\ 128$) was performed following a

standard pipeline by Maksimovic et al. [42] using the R package "minfi" (version 1.40.0; [2]).

The following sample exclusion criteria were applied: a mean detection p value $> .05$ ($n = 3$) and

sex mismatches between estimated sex from methylation data and confirmed phenotypic sex ($n = $

14). To normalize the beta values, we applied stratified quantile normalization[4] [72], followed by

BMIQ [70]. In further analysis, we removed probes containing SNPs ($n = 24\ 038$), X or Y

chromosome probes ($n = 16\ 263$), cross-hybridizing probes ($n = 46\ 867$) and polymorphic probes

($n = 297$) according to Chen et al. [11] and McCartney et al. [45], and probes with a detection p

value $> .01$ in at least one sample ($n = 79\ 922$). Afterwards, beta-values were transformed into

M-values, and batch-effects were removed using Combat [36]. For this purpose, we examined

the strength of associations between three possible batches (plate, slide, and array) and the first

five principal components via principal component analysis. The strongest batch effects were

iteratively removed. Sample mix-ups were checked with MixupMapper [74]. The comparison of

beta values with the existing genotype data (assessed using Global Screening Arrays [GSA+MD-

24v3.0-Psych-24v1.1, Illumina, San Diego, CA, USA]; for further details see [13]) revealed four

mix-ups. For these four cases, both samples (first and second measurements) were removed

(total of $n = 8$ individual samples). The resulting DNA methylation dataset comprises 2 102

---

[4] The chemical chip used to measure DNA methylation signal from extracted DNA consists of two types of probes (I and II). These two types of probes produce different distributions of signals that can introduce a technical noise when analyzed together. Stratified quantile normalization fixes this by stratifying the probes into type I and II, then using type I signals as 'anchors' to normalize Infinium II signals at the level of probe coverage categories [72].

samples of $n = 1\,055$ participants (among which $n = 488$ complete twin pairs or 976 individuals

with two timepoints of DNAm data) and 698 472 probes.

The cell type composition was estimated using the R package "EpiDish" (version 2.10;

[71]) and the epidish() function. Because estimated cell type proportions of saliva consisted of

three (highly correlated) cell types (leukocytes, epithelial cells, and fibroblasts), we obtained the

first principal component (for each epigenetic measurement, explaining 99.6% and 99.7 % of

cell type variation in the three estimated cell types at the first and second epigenetic

measurements, respectively) to control for cell type composition in the following analyses.

Calculating epigenetic age, epigenetic age acceleration, and pace of epigenetic aging

There are several types of epigenetic aging estimators, each based on a unique training

model with a variable number of CpGs, source tissues, and age ranges [3, 7]. To capture

potential differences in genetic and environmental contributions between different epigenetic

aging measures, we selected four commonly used ones (two trained on chronological age, and

two trained on biological health indicators) for our analysis: the Horvath clock [27], the PedBE

clock [47], GrimAge2 (an improved version of the original GrimAge; [40,41]), and

DunedinPACE ([5]; Additional file 2: Table S2 and Figure S1). All four epigenetic aging

measures have previously been shown to be associated with aging-related outcomes and to be

sensitive to external drivers of aging (e.g., stress; [12, 19, 41, 46, 50, 63]). In particular,

GrimAge and DunedinPACE have previously been associated with external drivers of aging in

saliva of children and adolescents [17, 58, 60]. The specific background of the training sample of

each epigenetic aging estimator is provided in Additional file 2: Table S2.

By the inclusion of the Horvath clock and the PedBE clock, we could address how

differences in the training age range and tissue can potentially correspond to the estimates of the

genetic and environmental contributions to the variance in epigenetic aging. The Horvath clock

was trained as a pan-tissue epigenetic aging estimator in a large age range (from 0 to 101 years;

[27]), while the PedBE clock was developed in buccal cells in younger age groups (from 0 to 20

years; [47]). By the inclusion of GrimAge2 and DunedinPACE, we addressed differences in the

effects of the training phenotypes: both of these estimators, in contrast to Horvath and PedBE

clocks, were trained on biological health indicators. In particular, GrimAge is trained on plasma

proteins and smoking [41] and DunedinPACE is trained on biomarkers of organ systems

integrity, such as blood pressure or serum leptin levels [5]. Finally, the inclusion of

DunedinPACE allowed, in addition, to address the differences in the study design. Namely, the

Horvath and PedBE clock, as well as GrimAge are trained on the cross-sectional data, while

DunedinPACE accounts for longitudinal patterns of aging.

Epigenetic aging measures were calculated based on the normalized batch-corrected beta

values. The Horvath and the PedBE epigenetic age were calculated using the R package

"methylclock" (version 1.01; [54]). A total of 334 (94.6%) of the original 353 Horvath clock

CpG sites and all 94 CpG sites of the PedBE clock were present in the dataset. Version 2 of

GrimAge was computed based on 1 029 CpG sites present in our dataset (out of 1 030 mentioned

in the publication) using analysis code from Lu et al. [40] provided by A. Lu and S. Horvath via

personal correspondence. The DunedinPACE values were calculated with the DunedinPACE R

package (https://github.com/danbelsky/DunedinPACE/) using all of 173 CpG sites.

The four epigenetic age estimators were significantly positively correlated with

chronological age (r(Horvath epigenetic age) = .80, $p < .001$; r(PedBE epigenetic age) = .77, $p <$

$.001$; r(GrimAge) = .80, $p < .001$; and r(DunedinPACE) = .08, $p < .001$; Additional file 2: Figure

S2). For the Horvath epigenetic age, the Median Absolute Error (MAE) estimates ranged from

2.75 to 3.70 (3.26 to 3.65 for early adolescents, from 2.75 to 2.93 for late adolescents, from 3.33

to 3.34 for emerging adults, and from 2.89 to 3.71 for young adults). For the PedBE epigenetic

age, the MAE estimates ranged from 1.07 to 12.72 (2.22 to 2.91 for early adolescents, from 1.08

to 2.01 for late adolescents, from 6.49 to 7.51 for emerging adults, and from 11.61 to 12.72 for

young adults). For GrimAge, the MAEs ranged from 19.38 to 25.60 (22.96 to 24.38 for early

adolescents, from 25.42 to 25.60 for late adolescents, from 21.84 to 22.57 for emerging adults,

and from 19.39 to 20.25 for young adults; Additional file 2: Figure S3).

284   Acceleration estimates for the Horvath, PedBE, and GrimAge epigenetic ages were

obtained as follows: epigenetic age was regressed on chronological age (in years and decimal

months), biological sex, and cell type composition within each measurement occasion. In

addition, the Horvath and PedBE epigenetic ages were regressed on smoking exposure (i.e., beta

values of the cg05575921 probe which has previously been shown to predict cigarette

consumption in both blood and saliva [55]). This method has been chosen to overcome the

missing data on smoking exposure in our sample (especially in the age group of the onset of

adolescence), the limited reliability of the self-reports on smoking status and to capture, beyond

active smoking, the possible effects of passive smoking exposure. The resulting residuals (i.e.,

Horvath Acceleration, PedBE Acceleration, and GrimAge Acceleration), defined as *age*

acceleration (positive score) or *age deceleration* (negative score) according to the direction of

the deviation, were used in our analyses. The DunedinPACE values were controlled for smoking

probe, biological sex, and cell type composition. For DunedinPACE, a value of one reflects a

pace of aging in line with one year of chronological aging. Correspondingly, values greater than

one reflect an increased and values lower than one a decreased pace of aging.

Analytical strategy

Main analyses

302   To examine the contributions of genetic and environmental factors to the variance in

epigenetic aging, we applied three biometrical variance decomposition models, including (1)

univariate twin models, (2) bivariate twin models across measurement occasions, and (3)

univariate twin models with age as a linear continuous moderator (Additional file 3: Figure S4;

[51]). MZ twins share close to 100% of their genetic makeup, while DZ twins share, on average,

50% of genetic variants that differ among humans. As a consequence, the correlation of any

genetic factors is approximately 1 for MZ twins, and is assumed to approach .50 for additive

genetic factors and .25 for non-additive genetic factors due to allelic dominance deviation for DZ

twins. Therefore, a greater correlation within MZ twin pairs than within DZ twin pairs signals

the presence of genetic influences on the variance in a specific phenotype, such as epigenetic

aging in the current study.

Furthermore, both MZ and DZ twins are assumed to share some prenatal, early life

familial, and extra-familial factors [8] that act to increase their resemblance beyond genetic

factors. These shared environmental factors are assumed to contribute to the similarity within

MZ twins to the same degree as within DZ twins. Only individually-unique environmental

factors are assumed to increase the dissimilarity between MZ twin siblings. Under the fulfillment

of these assumptions, the classical biometrical model allows for a decomposition of the observed

variance into variance attributable to genetic factors (A – additive genetic and D – non-additive

genetic), environmental influences shared by twins (C), and environmental influences unique to

each twin (E, including random error of measurement; [8]). For identification purposes, only

three of the variance components can be estimated simultaneously in the classical twin model.

Depending on the pattern of the correlations within the MZ and DZ twins, either D (if $r\text{MZ} < 2 \times$

$r\text{DZ}$) or C (if $r\text{MZ} > 2 \times r\text{DZ}$) components have to be fixed to zero, resulting in the specification

of either ACE or ADE models [16].

As outlined, we started our analysis with univariate twin models to decompose the

variance in each epigenetic aging estimator into genetic and environmental components

(Additional file 3: Figure S4a). Here, we pooled the data for the twin pairs across the two

measurement occasions, resulting in a sample of 976 twin pairs for each univariate twin model.

We set the variance of all variance components (A, C/D, and E) to 1 for identification purposes

and ensured that the statistical conditions specified for twin modeling, namely nonsignificant

differences in means and variances across co-twins and zygosity groups, were met in our sample.

Then, we decided for each of the four epigenetic aging measures whether to use ACE or ADE

models.

After that, among the different possible model variants (ACE, AE, CE, and E; or ADE,

AE, and E), we identified the respective model that provided the best fit to the data considering

model parsimony based on the chi-square difference test. In cases where alternative models (e.g.,

AE and CE) were nested in one more complex model (ACE) and did not differ significantly from

the latter model based on the chi-square test, the model with a lower Akaike information

criterion (AIC) was considered to fit the data better. Among the two nested models (e.g., AE vs.

ADE) that differed significantly based on the chi-square test, the model with the higher number

of parameters (e.g., ADE) was considered as the model that fits the data best [21]. For the best-

fitting models, we calculated the standardized coefficients with 90% confidence intervals.

In the second step, we specified bivariate twin models across the two measurement

occasions (488 twin pairs) to examine the genetic and environmental sources of variance in

epigenetic aging considering the repeated measurement (Additional file 3: Figure S4b). In this

bivariate factor model, we analyzed the first measurement of epigenetic aging as the first

variable and the second measurement of epigenetic aging as the second variable. Genetic and

environmental factors were specified for each measurement occasion and were allowed to

correlate over time in the sense of a *correlated factors model* [39]. For identification purposes,

the path coefficients of the latent factors were set to 1. This model specification allowed us to

estimate, along with genetic and environmental contributions to the variance in epigenetic aging

at the first and second measurement occasions, the genetic and environmental covariances

between the two timepoints. First, we ensured that the statistical conditions specified for twin

modeling were met in our sample and specified the full model. The full model included three

estimates per factor and two estimates of expected means. As was done for the univariate twin

models, we have chosen among the ACE, AE, CE, and E models or, alternatively, among the

ADE, AE, and E models which of these models provided the best balance between parsimony

and fit. The most parsimonious model showed the lowest AIC and did not significantly fit the

data worse than a more complex model based on the chi-square difference test. Using the most

parsimonious model, unstandardized and standardized variance components and genetic and

environmental correlations were estimated. To examine whether or not the genetic and

environmental contributions to the variance could be considered constant over time, we then

restricted the genetic and environmental variance components to be equal across the first and

second measurements of epigenetic aging. The model with equal variances was compared to the

less constrained model with unequal variance components over time using the chi-square

difference test.

In the third step, we specified twin models with age as a linear continuous moderator of

the variance components. Four age cohorts with two measures each were pooled together,

resulting in eight unique age groups from 9.5 to 30 (see Table 1) with age gaps of about two and

a half years between measurement occasions and age gaps of about three and a half years

between the second measure of the younger cohort and the first measure of the subsequent

cohort. Here, we tested whether the age of participants moderated genetic and environmental

contributions to the variance in epigenetic aging to unveil possible hints on gene☐environment

interactions (Additional file 3: Figure S4c; [56]). Since gene☐environment interaction effects

cannot be directly assessed in classical twin models, the estimates of genetic and environmental

contributions could be confounded with interaction effects [56]. In the case of interactions

between additive genetic factors (*A*) and environmental factors that are shared by siblings reared

together (*C*), unconsidered interaction effects would be confounded with estimates of additive

genetic effects. Potential but unconsidered interaction effects between genetic factors (*A*) and

environmental factors not shared by siblings reared together (*E*) would be confounded with

estimates of nonshared environmental effects. While the first interaction pattern is more

plausible in younger twins, who are still living in a common household, the latter is more

plausible in adult years, when twins live in different households and gain more separate life

experiences. As a consequence, increasing estimates of genetic contributions to the variance in

epigenetic aging are more plausible in adolescence and emerging adulthood than in adulthood

when accumulating effects of individually-unique experiences of twins may or may not interact

with genetic factors [10, 32].

To investigate the age-related moderation of the genetic and environmental variance

components of epigenetic aging, age was added as a continuous linear moderator to the

univariate models. This partitioned the ACE/ADE variance components into non-moderated

baseline ACE/ADE components, age-moderated ACE/ADE components, a baseline intercept

level, and independent linear and quadratic effects of age, producing a model with 9 parameters

in total. The model was still identified as the saturated model had in total 10 measured

parameters. We started with the full models (with 9 estimated parameters) and then dropped

nonsignificant paths to obtain more parsimonious models with more degrees of freedom. We

excluded parameters until we reached the model that fitted the data best, taking model parsimony

into account (i.e., a reduced model with a lower AIC that did not fit the data significantly worse

based on the chi-square test). The variance components, confidence intervals, and standardized

estimates were then derived.

Sensitivity analyses

In the final step, two sensitivity analyses were performed. As a first sensitivity analysis,

we scrutinized the models of the main analysis further not controlling the epigenetic aging

measures either for sex, smoking probe, or cell type composition. We repeated the univariate,

bivariate, and univariate age moderation analyses, first, for all chosen epigenetic aging measures

unadjusted for sex, second, for three epigenetic aging measures unadjusted for smoking, and,

finally, for four epigenetic aging measures unadjusted for cell type composition. For the second

sensitivity analysis, we rerun the age-moderated univariate models excluding the second

timepoint to support the patterns observed in age-moderated univariate twin models using the

pooled sample. By doing this, we could check if the trends in the genetic and environmental

contributions to the variance in epigenetic aging across age cohorts were similar to what we

observed in the main analysis with the pooled data across age cohorts and measurement

occasions, ensuring that drawing the same participants twice in the analysis sample, using the

first and second timepoints, did not affect the main analysis results.

The analyses were performed in R studio 2023.06.2 with R version 4.2.3. For data

preparation, we used the R packages "tidyverse" [75], "ltm" [62], "dlyr" [23], and "psych" [61].

For the variance decomposition, we used "OpenMX" [52].

Power analyses

For the estimation of power to detect the model with the best balance between parsimony

and fit to the data, we applied the functions proposed by Verhulst [73]. In detail, we applied the

function powerValue() along with acePow() function for univariate and bivPow() for bivariate

twin models. To address moderation, we applied a function developed for binary moderators,

sexLimPower(), and fitted paired age groups. We formulated two possible expectations

regarding genetic factors (A) based on estimates obtained in larger sample studies. Previous

research suggests that the contribution of genetic factors (A) to the variance of the Horvath clock

is 13% across the lifespan [37] or equals 37% at age 15 (combination with middle-aged mothers;

[67]). Our expectation regarding shared environmental factors (C) was based on plots for MZ

and DZ twins in Li et al. ([37]; page 8, Figure 4, cohabitation-dependent CE model). Here, we

identified approximated estimates of C for our sample's mean age (17.1 years) of about 35%.

The expectations for A and C were set to be the same across time. A power calculation for the

univariate models without moderation demonstrated that 976 pairs (526 MZ and 450 DZ twin

pairs) were sufficient for detecting the *C* component (power .99), while power for detecting the *A*

component varied from .40 to .99 depending on the estimates of *A*. In the bivariate case without

moderation, 488 pairs (263 MZ and 225 DZ twin pairs) were sufficient (.96 to .99) to detect the

C component. For the detecting of the *A* component, the power ranged from .27 to .99 depending

on the estimates of *A*. Finally, power calculation for models with moderation based on the power

calculation for the binary moderation demonstrated that the power to detect the *A* component

across the age groups varied in the univariate analysis from .12 to .61. The power to detect *C*

varied from .52 to .94.

Results

Descriptive analysis

The means and standard deviations for the epigenetic aging measures across age groups

are presented in Table 1. The twin correlations of epigenetic aging measures for MZ and DZ

twins for each cohort and timepoint separately are given in Table 2 (see also Additional file 3:

Table S3 for 95% confidence intervals). Rank-order stability estimates within and across twins'

measurements for MZ and DZ twins are shown in Additional file 3: Table S4.

(please add Table 1 here)

(please add Table 2 here)

Table 1

Descriptive statistics for epigenetic aging measures

Variable	Early adolescence		Late adolescence		Emerging adulthood		Young adulthood	
	Time 1: age 9.5	Time 2: age 12	Time 1: age 15.5	Time 2: age 18	Time 1: age 21.5	Time 2: age 24	Time 1: age 27.5	Time 2: age 30
N (MZ pairs/DZ pairs)	326 (82/81)		372 (87/99)		152 (42/34)		126 (52/11)	
Chronological Age *M* (*SD*)	9.51 (0.33)	11.83 (0.58)	15.48 (0.31)	17.75 (0.58)	21.58 (0.37)	23.89 (0.62)	27.60 (0.70)	29.92 (1.04)
Horvath Accel. *M* (*SD*)	-0.02 (2.64)	-0.25 (3.35)	-0.07 (3.56)	0.28 (3.99)	0.47 (4.30)	0.29 (4.38)	-0.30 (4.17)	-0.52 (4.20)
PedBE Accel. *M* (*SD*)	-0.10 (0.91)	-0.21 (0.98)	0.10 (0.95)	0.24 (1.14)	0.16 (1.21)	0.17 (1.36)	-0.22 (1.24)	-0.39 (1.30)
GrimAge Accel. *M* (*SD*)	0.31 (2.03)	-0.08 (2.41)	-0.44 (2.32)	0.11 (2.40)	0.09 (2.23)	0.01 (2.35)	0.37 (2.59)	-0.14 (2.61)
DunedinPACE *M* (*SD*)	0.00 (0.07)	-0.01 (0.08)	-0.01 (0.08)	0.00 (0.08)	0.00 (0.08)	0.00 (0.08)	0.02 (0.08)	0.02 (0.09)

Note. Four cohorts (early adolescents, late adolescents, emerging adults, and young adults) were measured twice, approximately 2.5 years apart. MZ, monozygotic; DZ, dizygotic; *M*, mean; *SD*, standard deviation; Accel., Acceleration.

455 **Table 2**

456 *Spearman correlations for monozygotic and dizygotic twins for epigenetic aging measures*

| Variable | Zygosity | Cross-cohort correlation | Early adolescence | | Late adolescence | | Emerging adulthood | | Young adulthood | |
			Time 1: age 9.5	Time 2: age 12	Time 1: age 15.5	Time 2: age 18	Time 1: age 21.5	Time 2: age 24	Time 1: age 27.5	Time 2: age 30
Horvath Accel.	MZ	.51***	.44***	.53***	.46***	.49***	.56***	.69***	.59***	.43**
	DZ	.37***	.37***	.59***	.34***	.40***	.25	.08	.52	.36
PedBE Accel.	MZ	.45***	.51***	.61***	.61***	.48***	.76***	.59***	.11	.06
	DZ	.49***	.60***	.45***	.55***	.61***	.01	.26	.23	.33
GrimAge Accel.	MZ	.32***	.27*	.30**	.35***	.16	.43**	.34*	.45***	.39**
	DZ	.17***	.17	.10	.11	.31**	.23	.04	.12	.36
DunedinPACE	MZ	.48***	.44***	.50***	.40***	.36***	.65***	.62***	.53***	.48***
	DZ	.19***	.32**	.23*	.16	.13	.22	.23	.26	-.13

Note. MZ, monozygotic; DZ, dizygotic; Accel., Acceleration; *, $p < .05$; **, $p < .01$; ***, $p < .001$.

Horvath and PedBE Acceleration, clocks trained on the chronological age, demonstrated

age acceleration in the groups of late adolescents and emerging adults and age deceleration in the

two other groups of early adolescents and young adults (see Table 1). For GrimAge

Acceleration, estimates were higher across several periods, including the onset of adolescence,

late adolescence, and young adulthood. The estimates of DunedinPACE showed a faster pace

during young adulthood compared to other age periods.

The standard deviations of the four epigenetic aging estimators across age groups

indicated patterns of increased variance across time and age stages (Table 1 and Additional file

3: Figure S5). The two measures trained on chronological age, Horvath and PedBE Acceleration,

showed a slightly decreasing trend during young adulthood, though. In the eldest age group ($M =$

29.92) compared to the youngest age group ($M = 9.51$), the standard deviation was 60% greater

for Horvath Acceleration, 43% greater for PedBE Acceleration, and 29% greater for both

GrimAge Acceleration and DunedinPACE.

The twin correlations for Horvath and PedBE Acceleration demonstrated substantial

similarity for adolescent twins, with only marginal differences between MZ and DZ twins (Table

2), indicating contributions of shared environmental influences to the variance in epigenetic

aging during this period of life. In groups of emerging adults and later, the correlations for MZ

twins were greater than the correlations for DZ twins (except the nonsignificant MZ and DZ twin

correlations for PedBE Acceleration in young adulthood), indicating genetic influences on the

variance in epigenetic aging in adulthood. In contrast, Spearman correlations for GrimAge

Acceleration and DunedinPACE were greater for MZ twins than for DZ twins across almost all

age groups, indicating genetic contributions to the variance in these epigenetic aging measures.

The rank-order stability across two measurement points was approximately $r = .50$ with

no differences between ages and between MZ and DZ twin groups for all measures except for

GrimAge Acceleration, where rank-order stability was lower in younger age groups, and PedBE

Acceleration, where rank-order stability was greater for DZ twins as compared to MZ twins in

young adulthood (Additional file 3: Table S4). Rank-order stability across matched MZ twin and

co-twin measurements tended to be greater than those of DZ twins in adulthood, indicating

declining contributions of shared environmental factors and increasing genetic contributions to

rank-order stability with age.

Univariate twin model analyses of epigenetic aging

To perform univariate twin modeling, we first examined whether the statistical conditions

specified for twin modeling were met (Additional file 3: Table S5). We observed small

differences in means across co-twins for Horvath Acceleration and DunedinPACE, but the

models with the expected means and variances constrained to be equal across twin siblings'

order and zygosity did not fit the data significantly worse.

To make a choice regarding the ACE or ADE model, we examined if the correlations for

MZ twins were more than two times higher than the DZ twin correlations. For Horvath, PedBE,

and GrimAge Acceleration, the cross-cohort MZ twin correlations were not more than twice as

strong as the DZ twin correlations, indicating the absence of non-additive genetic factors (*D*) and

the presence of additive genetic factors (*A*) and shared environmental factors (*C*) instead: rMZ =

.51 and rDZ = .37 regarding Horvath Acceleration, rMZ= .45 and rDZ = .49 regarding PedBE

Acceleration, rMZ = .32 and rDZ = .17 regarding GrimAge Acceleration (see Table 2). The

cross-cohort correlations for MZ twins for DunedinPACE were more than two times higher than

correlations for DZ twins indicating, besides additive genetic factors (*A*), the presence of non-

additive genetic factors (*D*): rMZ = .48 and rDZ = .19.

The results of the univariate twin model analyses of the epigenetic aging measures are

presented in the Additional file 3: Tables S6-S7. Cross-cohort standardized genetic and

environmental variance components for the full models are presented in Additional file 3: Figure

S6 and the best-fitting models in Figure 2. For Horvath Acceleration, we observed a better fit of

the full model (ACE), indicating that the variance in Horvath Acceleration was best explained by

additive genetic as well as environmental factors shared and not shared by twins. PedBE

Acceleration, GrimAge Acceleration, and DunedinPACE estimates in the univariate analysis

were better represented by restricted models: a CE model for PedBE Acceleration and an AE

model for both GrimAge Acceleration and DunedinPACE. The variance in PedBE Acceleration

was thus best explained by environmental factors shared and not shared by twins, whereas the

variance in GrimAge Acceleration and DunedinPACE was best explained by additive genetic

and individually-unique environmental factors.

Figure 2

Cross-cohort standardized genetic and environmental variance components of epigenetic aging

measures in the best-fitting model

Note. The estimates are derived from the best-fitting models in the univariate twin models

analyses: ACE for Horvath Acceleration, CE for PedBE Acceleration, and AE for GrimAge

Acceleration and DunedinPACE (see Additional file 3: Tables S6-S7).

Bivariate twin model analyses of epigenetic aging

In a further step, we decomposed the variance in the two repeated measures of epigenetic

aging with the use of bivariate twin models. The models with the expected means and variances

to be equal across co-twins and zygosity within each measurement occasion did not fit the data

significantly worse than the saturated models (Additional file 3: Table S5). Thus, the results

confirmed that the statistical conditions specified for bivariate twin modeling were met.

Following the previously reported MZ and DZ twin correlations (Table 2), we similarly

specified ACE models for Horvath, PedBE, and GrimAge Acceleration, and an ADE model for

DunedinPACE. As for the univariate analyses, a bivariate ACE model for Horvath, a CE model

for PedBE, and an AE model for GrimAge and DunedinPACE provided the best balance

between parsimony and fit (see Additional file 3: Tables S8-S9). Figure 3 presents the results

from the variance decompositions of both timepoints based on the best-fitting reduced models.

The results from the full models can be found in Additional file 3: Figure S7. The best-fitting

reduced models were used as a basis to test whether genetic and environmental variance

components could be constrained to be equal over time.

For each of the four epigenetic aging measures, restricting the variance components of all

three or two genetic and environmental factors (*A*, *C*, and *E*) to be equal across the two

timepoints, respectively, resulted in a significantly worse fit to the data (see Additional file 3:

Table S8). Accordingly, our results suggested that the relative contributions of the genetic and/or

environmental factors changed over time. However, setting only the variance component due to

additive genetic factors to be equal across the two timepoints did not result in a significantly

worse fit to the data for the epigenetic aging measures trained in adult samples (i.e., Horvath,

GrimAge, DunedinPACE). These results suggested almost equal absolute contributions of

additive genetic factors over time. In contrast, variance components due to environmental factors

that are not shared by twins increased from the first to the second measurement timepoint for all

epigenetic aging measures, resulting in increasing variance over time (see Figure 3). In addition,

for epigenetic aging measures trained on the chronological age, the variance components due to

shared environmental factors seemed to increase over time. This pattern was not observed for the

epigenetic aging measures trained on health biomarkers, for which shared environmental factors

were negligible.

Then, we estimated the degree to which genetic and environmental factors correlated over

time. The three epigenetic aging measures trained in adult samples demonstrated significant

positive genetic correlations from the first to the second measurement timepoint. The highest

positive correlation of additive genetic factors was observed for Horvath Acceleration ($r_A = .90$,

$p < .001$), followed by GrimAge Acceleration ($r_A = .82$, $p < .001$) and DunedinPACE ($r_A = .74$, p

$< .001$). Epigenetic aging estimators trained to predict chronological age, in turn, demonstrated

statistically significant positive correlations of the shared environmental factors over time ($r_C =$

$.92$, $p < .001$ for Horvath Acceleration and $r_C = .87$, $p < .001$ for PedBE Acceleration). Unique

environmental factors correlated significantly for PedBE Acceleration ($r_E = .22$, $p < .001$),

DunedinPACE ($r_E = .18$, $p < .001$), and Horvath Acceleration ($r_E = .15$, $p < .001$), but not

significantly for GrimAge Acceleration ($r_E = .06$, $p = .19$). Generally, the correlations of

individually-unique environmental factors were weak ($\leq .22$), while the correlations of additive

genetic factors and shared environmental factors were substantial ($\geq .74$). In line with the a priori

power analysis, the power to distinguish between the ACE/ADE, CE, and AE models varied

from .41 to 1 for Horvath Acceleration, from .40 to 1 for PedBE Acceleration, from .39 to .93 for

GrimAge Acceleration, and from .41 to .76 for DunedinPACE. The power to distinguish the

573   best-fitting model from the E model was 1 for all epigenetic aging measures (Additional file 3:

574   Table S8).

576   **Figure 3**

577   *Unstandardized genetic and environmental variance components of epigenetic aging measures*

578   *over 2.5 years*

579
Note. The estimates are derived from the best-fitting bivariate twin models: ACE for Horvath

Acceleration, CE for PedBE Acceleration, and AE for GrimAge Acceleration and DunedinPACE

(see also Additional file 3: Tables S8-S9). tp1, timepoint 1; tp2, timepoint 2.

Age moderation of genetic and environmental differences in epigenetic aging

In the final step, we explored whether the contributions of the *A*, *C* (or *D*), and *E*

components to the variance of epigenetic aging differed across age groups (Table 3, Figure 4,

Additional file 3: Tables S10-S11, Figure S8). Here, we only present the parameter estimates of

the models that fitted the data best. The model fit statistics can be found in Additional file 3:

Table S10.

When moderation by age in the univariate models was allowed, statistically significant

moderation was observed for all epigenetic aging measures resulting in generally increasing

variance across age groups for all epigenetic aging measures. However, different factors turned

out to be relevant for this increase in different epigenetic aging measures. For acceleration

measures trained on chronological age, including Horvath and PedBE Acceleration, we observed

the moderation of the individually-unique environmental component (E). That is, the unique

environmental variance increased across age groups regarding both estimators. The contribution

of the unique environmental factors to the variance in the Horvath and PedBE Acceleration was

larger for the oldest age group (young adults) compared to the youngest ones (4.18 at the onset

of adolescence versus 11.73 in young adults for Horvath Acceleration; 0.29 at the onset of

adolescence versus 1.23 in young adults for PedBE Acceleration, see Table 3). Moderation of the

variance attributable to genetic factors, with higher contribution in young adulthood compared to

early adolescence, was found for Horvath and GrimAge Acceleration (1.59 versus 9.37 for

Horvath Acceleration and 1.17 versus 2.81 for GrimAge Acceleration) as well as for

DunedinPACE (non-additive genetic component was 0.00056 in younger adolescents and

0.00251 in young adults). Thus, the narrow-sense heritability potentially increased across age

groups from 18% to 39% for Horvath Acceleration and from 24% to 43% for GrimAge

Acceleration. The broad-sense heritability ($A + D$) of DunedinPACE increased from 42% to

57%.

In contrast to the bivariate model and the univariate twin model without age moderation,

the CE model for PedBE (with moderated e effect) showed a higher AIC value (5 550.13) in

comparison to the best-fitting model in the age-moderated univariate analysis (moderated e

effect in ACE model; AIC = 5 548.89). However, the power to distinguish between these two

models was limited (.30) and both models did not fit significantly worse to the data in

comparison to the corresponding more complex models. Likewise, the AE model for

DunedinPACE yielded a significantly worse fit to the data compared to the full model (-2LL = -4

562.38, df = 1 947, AIC = -4 552.38, p = .048) in the univariate analysis with age moderation.

(please add Table 3 here)

618 **Table 3**

619 *Variance components derived from age-moderated univariate models*

Epigenetic aging measure	Age group	*A* (%)	*C/D* (%)	*E* (%)	*V*
	9.5	**1.59 (18)**	3.04 (35)	**4.18 (47)**	8.82
	12	**2.20 (22)**	3.04 (30)	**4.90 (48)**	10.14
Horvath Accel.	15.5	**3.20 (26)**	3.04 (25)	**6.00 (49)**	12.24
(ACE model with	18	**4.03 (29)**	3.04 (22)	**6.85 (49)**	13.92
moderated *a* and *e*	21.5	**5.36 (32)**	3.04 (19)	**8.14 (49)**	16.54
effects)	24	**6.42 (35)**	3.04 (16)	**9.13 (49)**	18.59
	27.5	**8.08 (37)**	3.04 (14)	**10.60 (49)**	21.72
	30	**9.37 (39)**	3.04 (12)	**11.73 (49)**	24.14
	9.5	0.16 (18)	0.43 (49)	**0.29 (33)**	0.88
	12	0.16 (16)	0.43 (45)	**0.37 (39)**	0.96
	15.5	0.16 (14)	0.43 (40)	**0.50 (46)**	1.09
PedBE Accel.	18	0.16 (13)	0.43 (36)	**0.60 (51)**	1.19
(ACE model with	21.5	0.16 (11)	0.43 (32)	**0.76 (57)**	1.35
moderated *e* effect)	24	0.16 (11)	0.43 (29)	**0.89 (60)**	1.48
	27.5	0.16 (9)	0.43 (26)	**1.08 (65)**	1.67
	30	0.16 (8)	0.43 (24)	**1.23 (68)**	1.82
	9.5	**1.17 (24)**	-	3.78 (76)	4.95
GrimAge Accel.	12	**1.33 (26)**	-	3.78 (74)	5.11
(AE model with	15.5	**1.57 (29)**	-	3.78 (71)	5.35
moderated *a* effect)	18	**1.76 (32)**	-	3.78 (68)	5.54

Epigenetic aging measure	Age group	A (%)	C/D (%)	E (%)	V
	21.5	**2.04 (35)**	-	3.78 (65)	5.82
	24	**2.25 (37)**	-	3.78 (63)	6.03
	27.5	**2.57 (40)**	-	3.78 (60)	6.35
	30	**2.81 (43)**	-	3.78 (57)	6.59
	9.5	0.002 (32)	**0.001 (10)**	0.003 (58)	0.005
	12	0.002 (30)	**0.001 (13)**	0.003 (57)	0.006
	15.5	0.002 (29)	**0.001 (17)**	0.003 (54)	0.006
DunedinPACE	18	0.002 (28)	**0.001 (20)**	0.003 (52)	0.006
(ADE model with	21.5	0.002 (27)	**0.001 (24)**	0.003 (49)	0.006
moderated d effect)	24	0.002 (26)	**0.002 (27)**	0.003 (47)	0.007
	27.5	0.002 (24)	**0.002 (31)**	0.003 (45)	0.007
	30	0.002 (23)	**0.002 (34)**	0.003 (43)	0.007

Note. The estimates are derived from the best-fitting univariate models with age moderation of

genetic and environmental components of the four epigenetic aging measures, which are

depicted in parentheses. *A* (%), component due to additive genetic factors; *D* (%), component

due to non-additive genetic factors; *C* (%), component due to shared environmental factors; *E*

(%), component due to individually-unique environmental factors including measurement error;

V, total variance; Accel., Acceleration. Components that are significantly moderated by age are

shown in bold.

Figure 4

Unstandardized and standardized variance components derived from age-moderated models of

epigenetic aging measures

Note. The estimates are derived from the best-fitting univariate models of age moderation of

genetic and environmental components in the four epigenetic aging measures: ACE model with

moderated *a* and *e* effects for Horvath Acceleration, ACE model with moderated *e* effect for

PedBE Acceleration, AE model with moderated *a* effect for GrimAge Acceleration, and ADE

model with moderated *d* effect for DunedinPACE (see also Table 3 and Additional file 3: Tables

S10-S11).

Sensitivity analysis

In the first part of the sensitivity analysis, we performed three separate analyses for

univariate twin modeling, bivariate twin modeling, and univariate twin modeling with age as a

moderator in epigenetic aging measures unadjusted either for biological sex, the smoking probe,

or cell type composition (Additional file 4: Tables S12-S29). Not controlling for sex or smoking

did not lead to different conclusions compared to those from the main analysis. Not controlling

for cell type composition obscured some of the findings from the main analysis (Additional file

4: Tables S24-S29). For Horvath Acceleration, the AE model demonstrated a better fit in the

univariate analysis without moderation, while the ACE model still yielded a better fit in the

bivariate analysis and the analysis with age moderation. For PedBE Acceleration, we observed,

along with the moderation of the *E* component, the moderation of the *A* component in the better

fitting ACE age-moderated model. GrimAge Acceleration demonstrated a better fit of the CE

models across the univariate and bivariate model in the sensitivity as opposed to the main

analysis with the latter indicating a better fit of the AE models. Not controlling for cell type

composition in DunedinPACE demonstrated, finally, a better fit of the ADE model in the

univariate and bivariate twin models, as well as the moderation of the *E* instead of the *D*

component in the age-moderated univariate twin model. The analysis of epigenetic aging

measures unadjusted for cell type composition did not reach sufficient power to distinguish

between models. The observed pattern is potentially explained by the fact that different tissues

have heterogeneous cell type compositions [69], which, in turn, can have strong independent

associations with age [76]. Therefore, the measures of epigenetic aging unadjusted for cell type

composition might reflect, in addition to the genuine age-associated signal of DNAm,

heterogeneity in different cell type compositions regarding DNAm as well as a link between cell

type composition and age.

As the second part of the sensitivity analysis, we specified and fitted the univariate

models of the variance decomposition with age moderation for the first epigenetic measurement

to explore if drawing the data from two measurements to increase statistical power could affect

the results. The models' goodness-of-fit statistics are presented in the Additional file 4: Table

S30. The same models were observed as the best-fitting to the data for Horvath Acceleration,

GrimAge Acceleration, and DunedinPACE. Similar to the main analysis, we observed that the

models with the moderated *a* and *e* paths for Horvath Acceleration, with the moderated *a* path

for GrimAge Acceleration, and with the moderated *d* path for DunedinPACE showed a better fit

to the data. For PedBE Acceleration, we did observe the moderation of the *e* path as also

reported in the main analysis. In contrast to the main analysis, however, the model without *A*

factors demonstrated a better fit to the data. The observed difference in results probably could

originate from the halved sample size and decreased power to detect lower contributions of

additive genetic factors to the variance.

Discussion

In this study, we investigated the genetic and environmental contributions to the variance

in epigenetic aging from early adolescence to young adulthood. Three main results can be

summarized. First, genetic and environmental contributions to the variance in epigenetic aging

varied between differently developed epigenetic aging estimators. Second, stable differences in

epigenetic aging across two measurement occasions, 2.5 years apart, were mainly attributable to

genetic and shared environmental factors, while individually-unique environmental factors

represented the primary source of instability (i.e., rank-order change) over time. Third, the

variance in epigenetic aging increased from early adolescence to young adulthood (i.e., from age

9.5 to 30 years), which was due to increasing genetic or environmental contributions to the

variance, depending on the specific estimator for epigenetic aging.

Genetic and environmental contributions to the variances of differently developed

epigenetic aging estimators

691   The performed univariate and bivariate twin model analyses demonstrated both

similarities and differences in the genetic and environmental contributions across the four

differently developed epigenetic aging measures. One of the similarities was that at least half of

the differences in every epigenetic aging estimator was attributable to individually-unique

environmental factors not shared by twins reared together. The contributions of these factors to

the variance could emphasize the role of individually-unique life experiences contributing to

individual differences in epigenetic aging. Childhood and adolescence are a sequence of

sensitive and critical developmental periods, during which the effects of positive and negative

experiences can be particularly influential. Experiences of violence in children or institutional

caring [29], for example, have already been mentioned to predict greater epigenetic age,

affecting ongoing brain and emotional regulation development [65]. Individually-unique

environmental factors can also reflect the *epigenetic drift*, stochastic changes in DNAm. When

epigenetic drift occurs in epigenetic aging estimators' locus-specific DNAm, the individually-

unique environmental variance estimated in biometrical variance decomposition models would

partly reflect epigenetic drift [20]. Epigenetic drift with age could also explain the greater power

of epigenetic age to predict chronological age at the onset of life compared to childhood and

adolescence [47, 67].

Genetic and shared environmental factors, in turn, contributed differently to the four

epigenetic aging measures. In particular, the variance in the best-fitting univariate models was

explained either by genetic (GrimAge Acceleration and DunedinPACE), by shared

environmental (PedBE Acceleration), or by both genetic and shared environmental factors

(Horvath Acceleration). The developmental design of the investigated epigenetic aging

estimators suggests that these differences are not random. The larger contribution of the shared

environmental factors to the variance in Horvath Acceleration and PedBE Acceleration

corresponds to the training outcome of these estimators. Both are trained to predict chronological

age, which is by definition shared by twins of one pair. The transition to adulthood, in turn, could

explain why genetic contributions were observed for all epigenetic aging measures except for

PedBE Acceleration. In contrast to the PedBE clock, which was initially developed in a sample

of individuals up to 20 years of age, the Horvath clock, GrimAge, and DunedinPACE were

trained in adults (see Additional file 2: Table S2). The genetic contributions we observed for the

latter three epigenetic aging estimators, but not for PedBE Acceleration, might potentially

characterize intrinsic epigenetic processes that are activated later in life and shared to a greater

extent by genetically identical MZ twins compared to DZ twins. These processes can include the

processes of physical development, for example the transition from puberty to the maintenance

of biological systems and the high organism functionality. As these processes are less

pronounced during adolescence, they may be not fully reflected in the genetic contributions to

the variance in PedBE Acceleration.

The patterns observed for the genetic and environmental contributions to the variance in

epigenetic aging were also reflected in stable differences across measurements, as shown in the

bivariate twin analyses. Here, substantial positive genetic correlations over time were found in

three estimators trained in adults. In contrast to environmental differences, the absolute size of

genetic variance in epigenetic aging estimators remained constant over time, suggesting that

genetic factors represent a primary source of stable differences in the epigenetic aging rate, at

least across 2.5 years in our study. Estimators trained to predict chronological age (Horvath

Acceleration and PedBE Acceleration) also demonstrated substantial positive correlations of

shared environmental factors. However, their contributions to the variance increased from the

first to the second measurement occasion, respectively. This indicates that the similarity of same-

aged siblings and differences between families in epigenetic aging relative to chronological age

tend to increase over time. While genetic and shared environmental factors represent the primary

source of stability, individually-unique environmental factors were found to be only marginally

stable over the period of 2.5 years. Moreover, individual differences in all epigenetic aging

estimators increased over time due to an increase in those environmental influences not shared

by twins. This indicates that individually-unique environmental sources represent the primary

source of inter-individual differences in intra-individual change in epigenetic aging over time.

Finally, the training design could potentially explain the specific variance patterns for

DunedinPACE, for which we observed a smaller variance in general and its modest increase.

This is in line with previous findings suggesting a remarkable stability of DunedinPoAm, the

predecessor of DunedinPACE, for the age period from 7 to 20 [53]. Beyond that, DunedinPACE

was characterized by the greatest contribution of genetic factors to the variance and was the only

epigenetic aging estimator for which the presence of non-additive genetic factors was indicated

to some extent. These findings could also signal the differences between cross-sectional and

longitudinal training designs. While the GrimAge, Horvath, and PedBE clocks were developed

by analyzing cross-sectional samples, DunedinPACE was trained on longitudinal data,

accounting for the within-individual trends of aging. Representing the speed of epigenetic aging

based on 12 months, DunedinPACE does not directly relate to chronological age and the

accumulation of aging processes, which are commonly represented by measures of epigenetic

age acceleration. The smaller changes in the variance in DunedinPACE, along with the largest

contribution of the genetic factors, suggest that their role in the pace of epigenetic aging across

adolescence and young adulthood is potentially more stable than in epigenetic acceleration

measures.

Genetic and environmental contributions to age-associated differences in epigenetic aging

estimators

The analysis of the age-associated differences demonstrated that the genetic and

environmental contributions to the variance were not constant across age groups. These findings

can be discussed in the context of potential gene☐environment correlations and

gene☐environment interactions relevant for variance in epigenetic aging from early adolescence

to young adulthood. That is, in addition to the patterns observed in the univariate and bivariate

models, the three estimators Horvath Acceleration, GrimAge Acceleration, and DunedinPACE,

trained in adults, demonstrated an increased variance across age groups due to genetic (additive

or non-additive) factors. First, this potential increase is consistent with longitudinal studies on

the SNP-based heritability of epigenetic aging and the heritability of other complex phenotypes.

Specifically, in the sample of children and their parents, the SNP-based heritability was

demonstrated to be 0% during the onset of life but higher later in life [67]. Similarly, genetic

contributions to the variance in complex phenotypes, such as intelligence, depressive symptoms,

and personality traits, demonstrate increasing heritability from childhood to adulthood [6, 32,

49].

The observed phenomenon of the increased role of genetic factors during this age is

commonly explained by active and evocative gene☐environment correlation in the presence of

gene☐environment interaction, "[…] when people become more and more autonomous and self-

directed during adolescence. They have increasing opportunities to pave their own way, evoke,

select and create environments that match their heritable tendencies. These environments in turn

can provide experiences that have the potential to reinforce the pre-existing tendencies" ([32]; p.

238). One illustrative example of an active gene☐environment correlation is the self-selection of

individuals in environments that fit their genetic predispositions [6, 31]. For instance, when

individuals with genetic predispositions for a slower epigenetic aging rather develop healthy

habits and search for health-promoting environments that fit this genetic tendency, this would

reflect processes of active gene☐environment transactions resulting in non-random gene–

environment correlation. As the increasing importance of active gene☐environment correlation

would come along with an increase of genetic differences, while a declining role of passive

gene☐environment correlation (i.e., genetic and environmental factors on offspring's phenotypes

are correlated due to parental phenotypes and behavior) would be reflected by decreasing effects

of shared environmental influences [10, 56], our findings can be considered in line with this

interpretation.

Since we did not detect increasing genetic variance over an interval of 2.5 years in our

sample, which is not entirely compatible with the interpretation of gene–environment correlation,

alternative explanations must also be considered. Specifically, the gene☐environment interplay

underlying estimates of genetic variance can also reflect interactions of genetic and

environmental factors shared by twins, which would also account for the finding of non-perfect

correlations of genetic factors over the 2.5-year period in our sample [32, 56]. This form of

gene–environment interplay might be more relevant for twins reared together and, thus, more

likely to occur during adolescence, when twins still share a common family household. Previous

studies illuminating the role of shared environmental factors emphasized the driving role of the

prenatal environment [38] and cohabitation for differences in epigenetic aging, implying that the

longer relatives live together, the more similar they are epigenetically [37]. From adolescence to

adulthood, the role of these factors seems to be steadily replaced by genetic and individually-

unique environmental factors as well as by interactions between them. As cohabiting is more

common before adulthood, our findings extend previous conclusions, implying an increasing role

of active gene☐environment correlation in the presence of interactions between genetic and

environmental factors, when relatives, on average, transit to adulthood and tend to start living

separately. These processes are reflected in the trajectories of gene☐environment interactions

and the relationships between individuals and their immediate environments. This would include,

for instance, the transition from the sorting into fitting environments to the individual adjustment

to changing environments, or finding a balance between person☐environment fitting and

adjustment [31].

Furthermore, the contributions of individually-unique environmental influences to the

variance notably increased during adolescence and emerging adulthood in Horvath and PedBE

Acceleration, epigenetic aging estimators trained on chronological age. In addition to the

discussed accumulated effects of unique life experiences and epigenetic drift, the observed

pattern can signal increasing measurement error in epigenetic aging measures, which was

particularly evident for PedBE Acceleration. Being trained in a younger population [47], its

application in adults was less precise. In our analysis, PedBE Acceleration had a larger error in

predicting chronological age in older cohorts, while its MZ and DZ twin correlation estimates

varied largely. Measurement error, however, is likely not the sole driver for the increasing

variance in epigenetic aging due to individually-unique environmental factors, as the same trend

of increasing environmental variance was observed for Horvath Acceleration. This estimator was

also trained to predict chronological age and its mean prediction error was consistent across

analyzed age stages and groups.

Although we observed that the individually-unique environmental factors increase from

the first to the second timepoint in GrimAge Acceleration and DunedinPACE, the contributions

of individually-unique environmental factors to variance in both estimators were not found to be

moderated by age in the pooled univariate analysis. In contrast to epigenetic aging estimators

trained on chronological age, the two latter measures were developed in (exclusively adult)

blood samples and rely on intrinsic biological functioning. In particular, GrimAge is linked to

the level of plasma proteins [41] and DunedinPACE reflects, among other factors, the level of

glycated hemoglobin, leptin, and serum C-reactive proteins [5]. Because these two estimators are

also intrinsically driven, these processes may be more elusive and less sensitive to trends in

individually-unique environmental contributions when they are measured in saliva.

The heritability of epigenetic aging based on saliva samples

The age-specific patterns of genetic contributions to the variance in epigenetic aging in

saliva obtained in our analysis were rather consistent with the assessments reported in blood.

However, our heritability estimates were, on average, lower than those reported in studies based

on blood tissue samples. For example, previous research reported that genetic differences

accounted for 69%, 73%, and 68% of the variance in Horvath Acceleration, GrimAge

Acceleration, and DunedinPACE in 22-year-olds, respectively [34]. The broad-sense heritability

for the 21.5-year-old twins in our analysis was substantially lower (32%, 35%, and 51%

respectively). The comparison of the heritability of PedBE Acceleration in saliva with its

heritability in other sources of DNAm (e.g., buccal cells) was not possible, because no study so

far, to our knowledge, provided heritability estimates for PedBE Acceleration and our study

indicated decreasing or zero heritability. The observed differences in heritability estimates across

sources of DNAm for epigenetic aging measures could be related to variations in the age

compositions of the studies' samples, the presence of age moderation, or the genuine differences

in epigenetic aging across tissues [77]. Therefore, further studies are needed to investigate

whether the observed pattern of the lower heritability of epigenetic aging based on saliva can be

replicated.

Limitations and future research

Among the strengths of our study are the relatively large sample size of twin pairs in the

field of epigenetics, the focus on both epigenetic age acceleration and the pace of epigenetic

aging measures, the examination of genetic and environmental contributions to the covariance in

epigenetic aging measures over two timepoints, and, finally, the consideration of age differences

during the second and third decades of life. Nevertheless, our study is not without limitations.

First, differences in the performance of epigenetic aging measures across different ages

and tissues may have introduced additional sources of measurement error. The Horvath clock,

for example, is less precise in younger samples, while the PedBE clock could be more biased in

adults [47]. Similarly, our study exclusively utilized saliva samples, but not all applied

epigenetic aging estimators were trained on saliva-based DNA methylation profiles. While the

Horvath clock is a pantissue epigenetic clock and the PedBE clock was developed in buccal

cells, DunedinPACE and GrimAge were trained in blood. Although the pace of aging can be

assessed based on saliva [58, 59] and the second version of GrimAge demonstrated good

performance in saliva-based DNA methylation data [40], applying these epigenetic aging

estimators in a tissue different from the initial design may affect the estimates. The Horvath

clock, at the same time, being a pantissue clock, could lose some precision when applied to any

tissue.

Second, despite the advantages of a comparatively large sample size with epigenetic data

and the use of two measurement points in the biometrical variance decomposition, the post hoc

power analysis demonstrated that the differentiation between several nested models was limited,

as confidence intervals overlapped. Previous literature has suggested a minimum sample size of

600 twin pairs for univariate analysis and larger samples for detecting possible non-additive

genetic factors, which, unfortunately, could not be reached [44, 64]. Nevertheless, we had

enough power to detect the presence of genetic or environmental age moderation in epigenetic

aging measures.

Third, the utilization of measurements from two timepoints in four birth cohorts could

introduce additional sources of variance. Different age groups, for instance, might introduce

cohort effects that cannot be entirely disentangled from age effects. In addition, the second

measurement was realized during the COVID-19 pandemic. While prior studies did not confirm

specific effects of the COVID-19 pandemic on epigenetic aging in adolescents [14], we observed

an increased contribution of the individually-unique environmental factors to variance in

epigenetic aging at the second measurement. It thus might be that pandemic-related factors, such

as changes in health status, lockdowns, and restrictions, formed potential new influential

environmental sources that could act very individually. Hence, the role of these factors in

epigenetic aging dynamics should be clarified further.

Fourth, classical twin analysis is an efficient way to estimate genetic and environmental

sources of variance but at the cost of several assumptions. One assumption is that environmental

experiences shared by twins contribute to the similarity within MZ twin pairs to the same degree

as within DZ twin pairs. One study proposed that this assumption might not hold true for

epigenetic aging, as MZ and DZ pairs can possibly share their preadult environmental effects to

different extents [37]. Such phenomenon can have both an environmental or genetic origin [18].

In addition, the classical twin design does not allow, so far, the estimation of shared

environmental (C) and non-additive genetic (D) contributions in one model, while both sources

of variance can be relevant for variance in epigenetic aging at the same or during different

developmental stages. These aspects should be addressed in further investigations.

Conclusion

Altogether, the present study demonstrated that the genetic and environmental sources of

variance in epigenetic aging differ across different epigenetic aging estimators. For estimators

trained in adult samples, larger contributions of genetic factors to the variance in epigenetic

aging were observed. Shared environmental factors accounted for variance in epigenetic aging

measures trained on chronological age only. Genetic factors and environmental factors shared by

twins reared together represented the primary sources contributing to the stability of epigenetic

aging differences in corresponding estimators over time, while individually-unique

environmental factors represented the primary source of rank-order change.

The increase of the variance in epigenetic aging measures from early adolescence to

young adulthood was associated with differences in the contributions of genetic and

environmental factors. The variance of epigenetic aging estimators trained in adult samples was

characterized by increasing contributions of genetic factors across age. Increasing differences in

epigenetic aging measures trained on chronological age, in addition, were driven by increasing

contributions of individually-unique environmental factors. These findings can be interpreted in

terms of an interplay between genetic and environmental factors and contribute to a better

understanding of the epigenetic aging processes during the transition to adulthood.

Declarations

Ethics approval and consent to participate

The TwinLife study was reviewed and approved by the German Psychological Society (Deutsche Gesellschaft für Psychologie; protocol number: RR 11.2009). The TwinSNPs project received ethical approval from the Ethical Committee of Bielefeld University (No. 2020-180) and the TwinSNPs and TECS projects received ethical approval from the Ethical Committee of the Medical Faculty of the University of Bonn (Lfd. Nr. 113/18). All participants provided written informed consent. Participation was voluntary and participants could withhold participation at any timepoint. The personal data collection and analysis complied with the corresponding data protection regulations. The statistical analysis was performed on pseudonymized data from the servers of Bielefeld University.

Availability of data and materials

The TwinLife data are archived in the GESIS data catalog: https://search.gesis.org/research_data/ZA6701. The data are released for academic research and teaching after the data depositor's written authorization. The epigenetic data are available for collaborators via TwinLife data management.

Funding

This work was supported by the German Research Foundation (DFG) as a part of the long-term project TwinLife under Grant [number 220286500], the project "Genetic and social causes of social inequalities: Integration of molecular genetic data into the longitudinal twin family study TwinLife" (TwinSNPs) under Grant [number 428902522], and the TwinLife Epigenetic Change Satellite (TECS) project under Grant [number 458609264].

Competing interests

The authors report that there are no competing interests to declare.

Authors' contributions

DVK, CK, and BM were involved in the study conception and design. DVK analyzed the data and interpreted the results. DVK, YL, AMS, and CK drafted the manuscript. DVK, YL, AMS, DC, JI, CKLP, MMN, EBB, MD, AJF, CK, and BM provided valuable feedback for the manuscript revision. MD & MMN were responsible for obtaining ethical approval. MD, BM, MMN, AJF, CKLP, FMS, EBB, DC, and CK collected the data. AMS, DC, DVK, and CKLP assessed DNA methylation and generated the epigenetic aging scores. BM supervised the study. All the authors have read and approved the final manuscript.

Acknowledgments

We thank M. Ruks, A. Kottwitz, and K. Krell from TwinLife data management team for their support; S. Zare, A. Vogelsang, and N. Fricker for their contribution to the DNA extraction and epigenetic profiling; M. Ruks and L. Weigel for valuable advice on the manuscript.

References

1. Alisch RS, Barwick BG, Chopra P, Myrick LK, Satten GA, Conneely KN, et al. Age-associated DNA methylation in pediatric populations. Genome Research. 2012;22(4):623–32. https://doi.org/10.1101/gr.125187.111

2. Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al. Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. Bioinformatics. 2014;30(10):1363–9. https://doi.org/10.1093/bioinformatics/btu049

3. Bell CG, Lowe R, Adams PD, Baccarelli AA, Beck S, Bell JT, et al. DNA methylation aging clocks: challenges and recommendations. Genome Biology. 2019;20(1):249. https://doi.org/10.1186/s13059-019-1824-y

4. Belsky DW, Caspi A, Arseneault L, Baccarelli A, Corcoran DL, Gao X, et al.

Quantification of the pace of biological aging in humans through a blood test, the

DunedinPoAm DNA methylation algorithm. eLife. 2020;9,e54870.

https://doi.org/10.7554/eLife.54870

5. Belsky DW, Caspi A, Corcoran DL, Sugden K, Poulton R, Arseneault L, et al.

DunedinPACE, a DNA methylation biomarker of the pace of aging. Elife.

2022;11:e73420. https://doi.org/10.7554/eLife.73420

6. Bergen SE, Gardner CO, Kendler KS. Age-Related Changes in Heritability of Behavioral

Phenotypes Over Adolescence and Young Adulthood: A Meta-Analysis. Twin Research

and Human Genetics. 2007;10(3):423–33. https://doi.org/10.1375/twin.10.3.423

7. Bergsma T, Rogaeva E. DNA Methylation Clocks and Their Predictive Capacity for

Aging Phenotypes and Healthspan. Neuroscience Insights. 2020;15:2633105520942221.

https://doi.org/10.1177/2633105520942221

8. Boomsma D, Busjahn A, Peltonen L. Classical twin studies and beyond. Nature Reviews

Genetics. 2002;3(11):872–82. https://doi.org/10.1038/nrg932

9. Braun PR, Han S, Hing B, Nagahama Y, Gaul LN, Heinzman JT, et al. Genome-wide

DNA methylation comparison between live human brain and peripheral tissues within

individuals. Translational Psychiatry. 2019;9(1):47. https://doi.org/10.1038/s41398-019-

0376-y

10. Briley DA, Livengood J, Derringer J. Behavior genetic frameworks of causal reasoning

for personality psychology. European Journal of Personality. 2018;32:202–20.

https://doi.org/10.1002/per.2153

11. Chen Y, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, et al.

Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium

HumanMethylation450 microarray. Epigenetics. 2013;8(2):203–9.

https://doi.org/10.4161/epi.23470

12. Dammering F, Martins J, Dittrich K, Czamara D, Rex-Haffner M, Overfeld J, et al. The

pediatric buccal epigenetic clock identifies significant ageing acceleration in children

with internalizing disorder and maltreatment exposure. Neurobiology of Stress.

2021;15:100394. https://doi.org/10.1016/j.ynstr.2021.100394

13. Deppe M, Pahnke CKL, Maj C, Aldisi R, Nöthen M, Diewald M, et al. Is happiness

always a personal(ity) thing? A quasi-replication and extension of previous well-being

studies. [Internet]. 2024 [cited 2024 Jun 3]. Available from: https://osf.io/6ctm4

14. deSteiguer AJ, Raffington L, Sabhlok A, Tanksley P, Tucker-Drob EM, Harden K.P.

Stability of DNA-Methylation Profiles of Biological Aging in Children and Adolescents

[Preprint]. Developmental Biology; 2023. Available from:

http://biorxiv.org/lookup/doi/10.1101/2023.10.30.564766

15. Diewald M, Kandler C, Riemann R, Spinath FM, Andreas A, Baier T, et al.

TwinLifeTwinLife (6.1.0) [dataset]. GESIS; 2022. Available from:

https://search.gesis.org/research_data/ZA6701?doi=10.4232/1.13987

16. Dolan CV, Boomsma DI, Nivard MG, Neal MC. A Note on Jöreskog's ACDE Twin

Model: A Solution, Not the Solution. Structural Equation Modeling: A Multidisciplinary

Journal. 2022;29(6):933–4. https://doi.org/10.1080/10705511.2022.2095640

17. Dos Santos Oliveira NC, Katrinli S, De Assis SG, Smith AK, Serpeloni F. Community

and domestic violence are associated with DNA methylation GrimAge acceleration and

heart rate variability in adolescents. European Journal of Psychotraumatology.

2023;14(2):2202054. https://doi.org/10.1080/20008066.2023.2202054

18. Evans DM, Martin NG. The validity of twin studies. GeneScreen. 2000;1(2):77–9.

https://doi.org/10.1046/j.1466-9218.2000.00027.x

19. Faul JD, Kim JK, Levine ME, Thyagarajan B, Weir DR, Crimmins EM. Epigenetic-

based age acceleration in a representative sample of older Americans: Associations with

aging-related morbidity and mortality. Proc Natl Acad Sci USA.

2023;120(9):e2215840120. https://doi.org/10.1073/pnas.2215840120

20. Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, et al. Epigenetic

differences arise during the lifetime of monozygotic twins. Proceedings of the National

Academy of Sciences. 2005;102(30):10604–9. https://doi.org/10.1073/pnas.0500398102

21. Gana K, Broc G. Structural equation modeling with lavaan. London□: Hoboken, NJ:

ISTE Ltd□; John Wilery & Sons, Inc; 2019.

22. Gaunt TR, Shihab HA, Hemani G, Min JL, Woodward G, Lyttleton O, et al. Systematic

identification of genetic influences on methylation across the human life course. Genome

Biology. 2016;17(1):61. https://doi.org/10.1186/s13059-016-0926-z

23. Hadley W, Romain F, Lionel H, Kirill M, Davis V. dplyr: A Grammar of Data

Manipulation [Software]. 2023. Available from: https://dplyr.tidyverse.org,

https://github.com/tidyverse/dplyr

24. Hagenbeek FA, Hirzinger JS, Breunig S, Bruins S, Kuznetsov DV, Schut K, et al.

Maximizing the value of twin studies in health and behaviour. Nature Human Behaviour.

2023;7(6):849–60. https://doi.org/10.1038/s41562-023-01609-6

25. Hahn E, Gottschling J, Bleidorn W, Kandler C, Spengler M, Kornadt AE, et al. What

Drives the Development of Social Inequality Over the Life Course? The German

TwinLife Study. Twin Research and Human Genetics. 2016;19(6):659–72.

https://doi.org/10.1017/thg.2016.76

26. Hong X, Cao H, Cao W, Lv J, Yu C, Huang T, et al. Trends of genetic contributions on

epigenetic clocks and related methylation sites with aging: A population□based adult

twin study. Aging Cell. 2024;e14403. https://doi.org/10.1111/acel.14403

27. Horvath S. DNA methylation age of human tissues and cell types. Genome Biology.

2013;14(10):R115. https://doi.org/10.1186/gb-2013-14-10-r115

28. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of

ageing. Nature Reviews Genetics. 2018;19(6):371–84. https://doi.org/10.1038/s41576-

018-0004-3

29. Jovanovic T, Vance LA, Cross D, Knight AK, Kilaru V, Michopoulos V, et al. Exposure

to Violence Accelerates Epigenetic Aging in Children. Scientific Reports.

2017;7(1):8962. https://doi.org/10.1038/s41598-017-09235-9

30. Jylhävä J, Hjelmborg J, Soerensen M, Munoz E, Tan Q, Kuja-Halkola R, et al.

Longitudinal changes in the genetic and environmental influences on the epigenetic

clocks across old age: Evidence from two twin cohorts. EBioMedicine. 2019;40:710–6.

https://doi.org/10.1016/j.ebiom.2019.01.040

31. Kandler C, Kühn S, Mönkediek B, Forstner AJ, Bleidorn W. A Multidisciplinary

Perspective on Person-Environment Fit: Relevance, Measurement, and Future Directions.

Current Directions in Psychological Science. 2024;0(0).

https://doi.org/10.1177/09637214241242451

32. Kandler C, Waaktaar T, Mõttus R, Riemann R, Torgersen S. Unravelling the interplay

between genetic and environmental contributions in the unfolding of personality

differences from early adolescence to young adulthood. European Journal of Personality.

2019;33:221–44. http://dx.doi.org/10.1002/per.2189

33. Kankaanpää A, Tolvanen A, Bollepalli S, Leskinen T, Kujala UM, Kaprio J, et al.

Leisure-Time and Occupational Physical Activity Associates Differently with Epigenetic

Aging. Medicine & Science in Sports & Exercise. 2021;53(3):487–95.

https://doi.org/10.1249/MSS.0000000000002498

34. Kankaanpää A, Tolvanen A, Heikkinen A, Kaprio J, Ollikainen M, Sillanpää E. The role

of adolescent lifestyle habits in biological aging: A prospective twin study. eLife.

2022;11:e80729. https://doi.org/10.7554/eLife.80729

35. Kilpinen H, Dermitzakis ET. Genetic and epigenetic contribution to complex traits.

Human Molecular Genetics. 2012;21(R1):R24–8. https://doi.org/10.1093/hmg/dds383

36. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing

batch effects and other unwanted variation in high-throughput experiments.

Bioinformatics. 2012;28(6):882–3. https://doi.org/10.1093/bioinformatics/bts034

37. Li S, Nguyen TL, Wong EM, Dugué PA, Dite GS, Armstrong NJ, et al. Genetic and

environmental causes of variation in epigenetic aging across the lifespan. Clinical

Epigenetics. 2020;12(1):158. https://doi.org/10.1186/s13148-020-00950-1

38. Li S, Wong EM, Dugué PA, McRae AF, Kim E, Joo JHE, et al. Genome-wide average

DNA methylation is determined in utero. International Journal of Epidemiology.

2018;47(3):908–16. https://doi.org/10.1093/ije/dyy028

39. Loehlin JC. The Cholesky approach: A cautionary note. Behavior Genetics.

1996;26(1):65–9. https://doi.org/10.1007/BF02361160

40. Lu AT, Binder AM, Zhang J, Yan Q, Reiner AP, Cox SR, et al. DNA methylation

GrimAge version 2. Aging. 2022. Available from: https://www.aging-

us.com/lookup/doi/10.18632/aging.204434

41. Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, et al. DNA methylation

GrimAge strongly predicts lifespan and healthspan. Aging. 2019;11(2):303–27.

https://doi.org/10.18632/aging.101684

42. Maksimovic J, Phipson B, Oshlack A. A cross-package Bioconductor workflow for

analysing methylation array data. F1000Research. 2017;5:1281.

https://doi.org/10.12688/f1000research.8839.3

43. Marioni RE, Shah S, McRae AF, Chen BH, Colicino E, Harris SE, et al. DNA

methylation age of blood predicts all-cause mortality in later life. Genome Biology.

2015;16(1):25. https://doi.org/10.1186/s13059-015-0584-6

44. Martin NG, Eaves LJ, Kearsey MJ, Davies P. The power of the classical twin study.

Heredity. 1978;40(1):97–116. https://doi.org/10.1038/hdy.1978.10

45. McCartney DL, Walker RM., Morris SW, McIntosh AM, Porteous DJ, Evans KL.

Identification of polymorphic and off-target probe binding sites on the Illumina Infinium

MethylationEPIC BeadChip. Genomics Data. 2016;9:22–4.

https://doi.org/10.1016/j.gdata.2016.05.012

46. McCrory C, Fiorito G, Hernandez B, Polidoro S, O'Halloran AM, Hever A, et al.

GrimAge Outperforms Other Epigenetic Clocks in the Prediction of Age-Related Clinical

Phenotypes and All-Cause Mortality. Le Couteur D, editor. The Journals of Gerontology:

Series A. 2021;76(5):741–9. https://doi.org/10.1093/gerona/glaa286

47. McEwen LM, O'Donnell KJ, McGill MG, Edgar RD, Jones MJ, MacIsaac JL, et al.

(2020). The PedBE clock accurately estimates DNA methylation age in pediatric buccal

cells. Proceedings of the National Academy of Sciences. 2020;117(38):23329–35.

https://doi.org/10.1073/pnas.1820843116

48. Miao K, Liu S, Cao W, Lv J, Yu C, Huang T, et al. Five years of change in adult twins:

longitudinal changes of genetic and environmental influence on epigenetic clocks. BMC

Med. 2024;22(1):289. https://doi.org/10.1186/s12916-024-03511-y

49. Mõttus R, Briley DA, Zheng A, Mann FD, Engelhardt LE, Tackett, JL, et al. Kids

becoming less alike: A behavioral genetic analysis of developmental increases in

personality variance from childhood to adolescence. Journal of Personality and Social

Psychology. 2019;117:635–58. http://dx.doi.org/10.1037/pspp0000194

50. Musci RJ, Raghunathan RS, Johnson SB, Klein L, Ladd-Acosta C, Ansah R, et al. Using

Epigenetic Clocks to Characterize Biological Aging in Studies of Children and

Childhood Exposures: a Systematic Review. Prevention Science. 2023;24(7):1398–423.

https://doi.org/10.1007/s11121-023-01576-4

51. Neale MC. Biometrical Models in Behavioral Genetics. In: Kim YK, editor. Handbook of

Behavior Genetics. New York, NY: Springer New York; 2009. p. 15–33.

https://doi.org/10.1007/978-0-387-76727-7_2

52. Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM. OpenMx 2.0:

Extended Structural Equation and Statistical Modeling. Psychometrika. 2016;81(2):535–

49. https://doi.org/10.1007/s11336-014-9435-8

53. Niccodemi G, Menta G, Turner J, D'Ambrosio C. Pace of aging, family environment and

cognitive skills in children and adolescents. SSM - Population Health. 2022;20:101280.

https://doi.org/10.1016/j.ssmph.2022.101280

54. Pelegí-Sisó D, de Prado P, Ronkainen J, Bustamante M, González JR. methylclock: A

Bioconductor package to estimate DNA methylation age. Bioinformatics.

2021;37(12):1759–60. https://doi.org/10.1093/bioinformatics/btaa825

55. Philibert R, Dogan M, Beach SRH, Mills JA, Long JD. AHRR methylation predicts

smoking status and smoking intensity in both saliva and blood DNA. American Journal

of Medical Genetics Part B: Neuropsychiatric Genetics. 2020;183(1):51–60.

https://doi.org/10.1002/ajmg.b.32760

56. Purcell S. Variance Components Models for Gene–Environment Interaction in Twin

Analysis. Twin Research. 2002;5(6):554–71.

https://doi.org/10.1375/136905202762342026

57. Raffington L, Belsky DW. Integrating DNA Methylation Measures of Biological Aging

into Social Determinants of Health Research. Current Environmental Health Reports.

2022;9(2):196–210. https://doi.org/10.1007/s40572-022-00338-8

58. Raffington L, Tanksley PT, Sabhlok A, Vinnik L, Mallard T, King LS, et al. Socially

Stratified Epigenetic Profiles Are Associated With Cognitive Functioning in Children

and Adolescents. Psychological Science. 2023;34(2):170–85.

https://doi.org/10.1177/09567976221122760

59. Raffington L, Tanksley PT, Vinnik L, Sabhlok A, Patterson MW, Mallard T, et al.

Associations of DNA-Methylation Measures of Biological Aging With Social Disparities

in Child and Adolescent Mental Health. Clinical Psychological Science. 2023;0(0).

https://doi.org/10.1177/21677026231186802

60. Raffington L, Schwaba T, Aikins M, Richter D, Wagner GG, Harden KP, et al.

Associations of socioeconomic disparities with buccal DNA-methylation measures of

biological aging. Clinical Epigenetics. 2023;15(1):70. https://doi.org/10.1186/s13148-

023-01489-7

61. Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research

(R package version 2.3.6) [Software]. Evanston, Illinois: Northwestern University; 2023.

Available from: https://CRAN.R-project.org/package=psych

62. Rizopoulos D. ltm: An R Package for Latent Variable Modeling and Item Response

Theory Analyses. Journal of Statistical Software. 2006;17(5). Available from:

http://www.jstatsoft.org/v17/i05/. https://doi.org/10.18637/jss.v017.i05

63. Ryan J, Wrigglesworth J, Loong J, Fransquet PD, Woods RL. A Systematic Review and

Meta-analysis of Environmental, Lifestyle, and Health Factors Associated With DNA

Methylation Age. Anderson R, editor. The Journals of Gerontology: Series A.

2020;75(3):481–94. https://doi.org/10.1093/gerona/glz099

64. Sham PC, Purcell SM, Cherny SS, Neale MC, Neale BM. Statistical Power and the

Classical Twin Design. Twin Research and Human Genetics. 2020;23(2):87–9.

https://doi.org/10.1017/thg.2020.46

65. Shonkoff JP, Garner AS. The Lifelong Effects of Early Childhood Adversity and Toxic

Stress. Pediatrics. 2012;129(1):e232–e246. https://doi.org/10.1542/peds.2011-2663

66. Sillanpää E, Ollikainen M, Kaprio J, Wang X, Leskinen T, Kujala UM, et al. Leisure-

time physical activity and DNA methylation age—A twin study. Clinical Epigenetics.

2019;11(1):12. https://doi.org/10.1186/s13148-019-0613-5

67. Simpkin AJ, Hemani G, Suderman M, Gaunt TR, Lyttleton O, Mcardle WL, et al.

Prenatal and early life influences on epigenetic age in children: A study of mother–

offspring pairs from two cohort studies. Human Molecular Genetics. 2016;25(1):191–

201. https://doi.org/10.1093/hmg/ddv456

68. Sinke L, Cats D, Heijmans BT. Omixer: Multivariate and reproducible sample

randomization to proactively counter batch effects in omics studies. Bioinformatics.

2021;37(18):3051–2. https://doi.org/10.1093/bioinformatics/btab159

69. Teschendorff AE. Cell-Type Heterogeneity in DNA Methylation Studies: Statistical

Methods and Guidelines. In: Michels KB, editor. Epigenetic Epidemiology. Springer

International Publishing; 2022. p. 67–96. https://doi.org/10.1007/978-3-030-94475-9_4

70. Teschendorff AE, Marabita F, Lechner M, Bartlett T, Tegner J, Gomez-Cabrero D, et al.

A beta-mixture quantile normalization method for correcting probe design bias in

Illumina Infinium 450 k DNA methylation data. Bioinformatics. 2013;29(2):189–96.

https://doi.org/10.1093/bioinformatics/bts680

71. Teschendorff AE, Zheng SC. EpiDISH [Software]. Bioconductor; 2017. Available from:

https://doi.org/10.18129/B9.BIOC.EPIDISH

72. Touleimat N, Tost J. Complete pipeline for Infinium [®] Human Methylation 450K

BeadChip data processing using subset quantile normalization for accurate DNA

methylation estimation. Epigenomics. 2012;4(3):325–41.

https://doi.org/10.2217/epi.12.21

73. Verhulst B. A Power Calculator for the Classical Twin Design. Behavior Genetics. 2017;47(2):255–61. https://doi.org/10.1007/s10519-016-9828-9

74. Westra HJ, Jansen RC, Fehrmann RSN, te Meerman GJ, van Heel D, Wijmenga C, et al. MixupMapper: Correcting sample mix-ups in genome-wide datasets increases power to detect small genetic effects. Bioinformatics. 2011;27(15):2104–11. https://doi.org/10.1093/bioinformatics/btr323

75. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the Tidyverse. Journal of Open Source Software. 2019;4(43):1686. https://doi.org/10.21105/joss.01686

76. Wong YT, Tayeb MA, Stone TC, Lovat LB, Teschendorff AE, Iwasiow R, et al. A comparison of epithelial cell content of oral samples estimated using cytology and DNA methylation. Epigenetics. 2022;17(3):327–34. https://doi.org/10.1080/15592294.2021.1950977

77. Wu HC, Wang Q, Chung WK, Andrulis IL, Daly MB, John EM, et al. Correlation of DNA methylation levels in blood and saliva DNA in young girls of the LEGACY Girls study. Epigenetics. 2014;9(7):929–33. https://doi.org/10.4161/epi.28902

78. Yang J, Zeng J, Goddard ME, Wray NR, Visscher PM. Concepts, estimation and interpretation of SNP-based heritability. Nature Genetics. 2017;49(9):1304–10. https://doi.org/10.1038/ng.3941

Additional files

Additional file 1. The SNP-, pedigree-, and twin-based heritability of epigenetic aging estimated in previous studies.

Additional file 2. Performance and training characteristics of epigenetic aging measures. The

figures and tables represent CpG sites overlap between the epigenetic aging measures, Median

Absolute Errors and mean deviations for three epigenetic age estimators.

Additional file 3. Supplementary Tables and Figures.

Additional file 4. Sensitivity analysis results. Tables S12-29 report goodness-of-fit statistics and

standardized variance components for univariate, bivariate, and univariate age-moderated twin

models of epigenetic aging measures unadjusted for biological sex, smoking probe, or cell type

composition. Table S30 reports the univariate age-moderated twin models of epigenetic aging

measures for the first epigenetic measurement.