Age-Associated Genetic and Environmental Contributions to Epigenetic Aging Across

Adolescence and Emerging Adulthood

Dmitry V. Kuznetsov[1*], Yixuan Liu[1], Alicia M. Schowe[2,3], Darina Czamara[2], Jana Instinske[4],

Charlotte K. L. Pahnke[5], Markus M. Nöthen[5], Frank M. Spinath[6], Elisabeth B. Binder[2], Martin

Diewald[1], Andreas J. Forstner[5,7], Christian Kandler[4], and Bastian Mönkediek[1]

[1]Bielefeld University, Germany

[2]Max Planck Institute of Psychiatry, Department Genes and Environment, Munich, Germany

[3]Graduate School of Systemic Neuroscience, Ludwig Maximilian University, Munich, Germany

[4]University of Bremen, Germany

[5]Institute of Human Genetics, University of Bonn, School of Medicine & University Hospital

Bonn, Bonn, Germany

[6]Department of Psychology, Saarland University, Saarbrücken, Germany

[7]Institute of Neuroscience and Medicine (INM-1), Research Center Jülich, Germany

*corresponding author

Abstract

Background

Epigenetic aging estimators commonly track chronological and biological aging,

quantifying its accumulation (i.e., epigenetic age acceleration) or its speed (i.e., epigenetic aging

pace). These estimates reflect a combination of inherent biological programming and the impact

of environmental factors, which are suggested to vary at different stages of life. The transition

from adolescence to adulthood is an important period in this regard, marked by an increasing

and, then, stabilizing epigenetic aging variance. Whether this pattern arises from environmental

influences or genetic factors remains uncertain.

This study delves into understanding the age-associated genetic and environmental

contributions to differences both between and within individuals across these developmental

stages. For this purpose, we analyzed four differently developed epigenetic aging estimators,

namely, Horvath Acceleration, GrimAge Acceleration, PedBE Acceleration, and DunedinPACE,

which were collected from 976 twins aged 8 to 31 years (M=16.0, SD=6.05) measured twice,

two years apart.

Results

Approximately half (33-76%) of the between-individual differences in epigenetic aging

and most (70-99%) of the variance in within-individual changes across the two years were

attributable to unique environmental factors. The contribution of these factors to the variance of

epigenetic estimators trained on chronological age notably increased with time (Horvath

Acceleration: from 80% to 90%; PedBE Acceleration: from 70% to 99%).

Genetic contributions were more pronounced in the between-individual differences than

in the variance of within-individual changes. For epigenetic aging estimators trained on

chronological age, both additive genetic factors (8%-39%) and shared environmental influences

(13%-49%) contributed to the variance. For epigenetic estimators trained on biological

indicators, either additive (GrimAge Acceleration) or additive and non-additive genetic

(DunedinPACE) factors were relevant. The variance of three estimators, initially developed in

adult samples, demonstrated increasing contributions of genetic factors to the variance across

adolescence and emerging adulthood (Horvath Acceleration: from 18% to 39%; GrimAge

Acceleration: from 24% to 43%; DunedinPACE: from 42% to 57%).

Conclusions

Our findings suggest that increasing between-individual differences in epigenetic aging

during adolescence and emerging adulthood are the result of both unique life experiences and

genetic underpinnings, whereas variance in within-individual changes is primarily attributable to

unique life experiences.

Keywords Epigenetic aging, Age-moderation, Adolescence, Gene-environment

interaction, the Horvath clock, the PedBE clock, GrimAge, DunedinPACE.

Background

Estimators of epigenetic aging offer insights into how individuals age over time and

across age groups [20]. They are predictive of age-related diseases and mortality in adults [34,

45], worse cognitive functioning [46] and socioeconomic disadvantage [47] in adolescents and

children. By measuring DNA methylation (DNAm) levels at specific DNA loci, epigenetic

estimators address universal and specific aspects of aging processes. For example, the Horvath

clock, which was trained to predict chronological age in samples with a large age range,

considers differences between chronological and epigenetic age to indicate accelerated or

decelerated aging across the whole life span [19]. Similarly, in buccal samples of children and

adolescents, the pediatric-buccal-epigenetic (PedBE) clock illuminates the specific aging

processes in younger age groups [37]. Alternatively, epigenetic aging estimators have been

developed on the basis of links between aging and biological health indicators. Examples of such

measures include the biomarker of human mortality risk GrimAge [31, 32] and the biomarkers of

the pace of aging, DunedinPoAm and DunedinPACE [3, 4]. These measures account for DNAm

at locus-specific sites associated, for example, with plasma proteins and smoking (GrimAge) or

biomarkers of organ systems integrity, such as blood pressure or serum leptin levels (Dunedin).

In addition, Dunedin measures are trained on longitudinal data and represent, in contrast to the

degree of epigenetic aging, the pace of epigenetic aging [3, 4].

The strength of the prediction by epigenetic aging measures stems from their ability to

represent both genetically driven aging processes and the cumulative effects of the environment

on DNAm [27]. The contributions of genetic factors to the variance in epigenetic aging can be

reflected in the narrow- and broad-sense *heritability,* the proportion of the phenotypic variance

attributable to genetic differences at the time and age of measurement in a certain population.

Narrow-sense heritability quantifies only the contribution of additive genetic factors, while

broad-sense heritability also includes the interactions between genetic factors such as allelic

dominance (i.e., interactions within specific gene loci of chromosomes) and epistasis (i.e.,

interactions across genetic variants between gene loci).

Figure 1 and Additional file 1: Table S1, summarizes previously reported estimates of

SNP- [1], pedigree-[2], and twin-based[3] heritability for the aforementioned epigenetic measures. As

shown in Figure 1, the narrow-sense heritability of the Horvath clock is not constant across

particular ages. According to previous studies, heritability estimates vary from 39% to 61% in

older adults, which is on average lower than the estimates obtained for young adults, ranging

from 62% to 74% [19, 22, 25, 52]. In adolescents, pedigree- and SNP-based research revealed a

lower heritability of the Horvath clock than in young adult samples, with a range from 37% to

43% [34, 53]. The possibility of extrapolating this reverse U-shaped trend in the heritability of

the Horvath clock to other epigenetic aging measures and various DNAm resources remains less

explored. For example, comparisons of heritability between epigenetic aging estimators are

mainly confined to emerging adults where the Horvath clock, GrimAge, and pace of aging

demonstrated similar heritabilities of 62-73% [26]. Knowledge on the comparability of

heritability between tissues is rather limited, as most studies have focused on blood tissue.

---

[1] Variance explained by all single nucleotide polymorphisms used in a genome-wide association study in conventionally unrelated individuals [64].

[2] Variance explained by the patterns of genetic resemblance among not-too-distantly related family members [17].

[3] Variance explained by the patterns of genetic resemblance between monozygotic and dizygotic twins.

Studying the age-dependent patterns of heritability across a broader spectrum of epigenetic aging

measures based on noninvasive saliva samples could elucidate universal trends or tissue- and

measure-specific differences in epigenetic aging.

The trend of epigenetic aging heritability across particular ages may arise from non-

genetic environmental sources and their interaction with genetic factors. A design of twins reared

together, for example, allows to disentangle environmental factors shared by twins, which

increase twin siblings' similarity, from non-shared or individually unique environmental factors,

which increase twin siblings' dissimilarity. Particularly in younger age, when twins live together

in one household, shared environments may be important. These environments can encompass

prenatal factors, social backgrounds, common parental and peer influences, and shared

neighborhoods. Research on complex traits suggests that shared environmental and genetic

factors may interact, particularly in younger people. With respect to the variance in epigenetic

aging, the role of shared environmental factors was found to increase before adulthood when

twins lived together [29]. Individually unique environmental factors are presented across the

whole life span and can obtain greater importance when twins live in different households and

have more separate life experiences. These environments can include stochastic changes and

individual-specific experiences, such as different peer groups or differences in experiences of

objectively shared environments. Unique environmental factors have been shown to moderate

the influence of genetic factors on DNAm [15] and possibly have the same effect on epigenetic

aging during adulthood [22]. These shifts in genetic and environmental contributions to the

variance of epigenetic aging over the life course suggest the possible link between life stage

processes and gene☐environment interplay in epigenetic aging.

Adolescence and the transition to adulthood, so-called emerging adulthood, is potentially

one of the periods of dynamic genetic and environmental contributions to the variance of

epigenetic aging. The time before transition is a period of growth and development with a faster

change rate in DNAm and epigenetic age [1, 20]. Studies have shown, for instance, that this

change rate peaks in infancy, decreases nonlinearly until approximately age 20, and then

stabilizes [19, 20]. Variance in epigenetic aging can also undergo changes. Starting small at

birth, the variance in the epigenetic age of the Horvath clock increases with chronological age

until adulthood onset, and then stabilizes [29]. It remains unclear whether these trends of later

stabilization in change rates correspond to trends in the relative contributions of genetic and non-

genetic factors and which sources, genetic or environmental, are responsible for the growing and

stabilizing variance of epigenetic aging during adolescence and the transition to adulthood.

To address this gap in the epigenetic literature, this study aimed to investigate the trends

in genetic and environmental contributions to the variance in epigenetic aging across

adolescence and emerging adulthood. We focused on between- and within-individual

differences. With between-individual differences, we can examine how genetic and

environmental sources contribute to the variance in epigenetic aging and how these sources shift

across age groups. Analyzing within-individual differences would reveal, in addition, the genetic

and environmental contributions to the variance in individual changes in epigenetic aging

(approximately two years in this study), which allows us to investigate individual deviations

from the common trends of epigenetic aging for different developmental stages. We examined

four epigenetic aging estimates measured twice, two years apart. DNAm was obtained from

DNA extracted from saliva samples of early (*M(age)* = 10.7 years) and late adolescents (*M(age)*

= 16.6 years), emerging adults (*M(age)* = 22.7 years), and young adult twins (*M(age)* = 28.8

years). The epigenetic aging estimates included two acceleration measures trained on

chronological age, the Horvath epigenetic clock [19] and the PedBE epigenetic clock [37], and

two epigenetic aging measures trained on biological indicators of aging, one of which, GrimAge

[31], represents acceleration and the other, DunedinPACE, represents the pace of aging [4].

Comparing these epigenetic aging estimates allowed us to (a) assess the age-associated trends in

the genetic and environmental contributions to between-individual differences and within-

individual changes across adolescence and emerging adulthood, (b) compare the genetic and

environmental contributions to the variance in differently developed epigenetic aging estimators,

and (c) examine whether heritability estimates of epigenetic aging assessed based on saliva are

similar to those that have previously been observed in other studies of blood samples (see Figure

1 and Additional file 1: Table S1).

Figure 1

The SNP-, pedigree-, and twin-based heritability of epigenetic aging estimated in previous

studies

Note. The estimates of SNP-, pedigree-, and twin-based heritability are presented for Horvath

Acceleration (purple), GrimAge Acceleration (green), and epigenetic aging pace measures,

DunedinPoAm and DunedinPACE (yellow).

Methods

Sample

The research questions were examined in a sample of 976 twins from the TwinLife

Epigenetics Change Satellite (TECS) project, which is a subsample of the German Twin Family

Panel (TwinLife). TwinLife is a longitudinal study of four cohorts of same-sex twin pairs and

their families recruited from the general population of Germany [11]. The study focuses on

longitudinal measures of social inequality and the potential underlying biopsychosocial

mechanisms [18]. The current analysis sample consisted of 263 monozygotic (MZ) and 225

dizygotic (DZ) complete twin pairs aged 8 to 29 years at the time of the first epigenetic

measurement ($M = 16.0$) and aged 10 to 31 years at the second measurement ($M = 18.3$). At the

time of assessment, members of the four cohorts were in the ages of early adolescence, late

adolescence, emerging adulthood, and young adulthood, respectively (for more details see Table

1).

DNA methylation

Saliva sampling and DNA extraction

Saliva sampling for DNA extraction was conducted as a part of two TwinLife satellite

projects, namely TwinSNPs and TECS. Twins and their family members were invited to

participate in the molecular genetic analysis during TwinLife face-to-face interviews carried out

in 2018–2020. During face-to-face interviews, interviewers offered to participate in the

molecular genetic studies, which involved genotyping and epigenetic profiling. A short video

explaining the study goals and procedures of the saliva sampling accompanied the recruitment

procedure. The second saliva collection took place in 2021. This time, participants were

contacted by mail. In addition to the invitation to participate, respondents received consent

forms, saliva collection toolkits, and corresponding instructions. For all participants who

provided valid consent, saliva samples were sent to the researchers at the University Hospital

Bonn, Germany. The samples were collected using Oragene® saliva self-collection kits (OG-500

and OG-600; DNA Genotek, Canada) and DNA was extracted following the manufacturer's

protocol.

DNA methylation assessment

The assessment of DNA methylation profiles was conducted in a set of 2 128 samples, of

which the present sample constituted a subset. The samples were randomized by age, sex,

zygosity, and complete versus single twin pair across 23 plates (266 arrays) using the R package

"Omixer" (version 1.4.0; [54]). The family identifier was used as a blocking variable to ensure

that all samples and timepoints of a twin pair ran on the same batch/array. Bisulfite-conversion

of 500 ng of DNA was performed using the D5033 EZ-96 DNA Methylation-Lightning Kit

(Deep-Well; Zymo Research Corp., Irvine, USA), and methylation profiles were assessed using

the Infinium MethylationEPIC BeadChip v1.0 (Illumina, San Diego, CA, USA).

Preprocessing of the methylation samples ($n = 2\ 128$) was performed following a

standard pipeline by Maksimovic et al. [33] using the R package "minfi" (version 1.40.0; [2]).

The following sample exclusion criteria were applied: a mean detection p value > .05 ($n = 3$) and

sex mismatches between estimated sex from methylation data and confirmed phenotypic sex ($n =$

14). To normalize the beta values, we applied stratified quantile normalization [58], followed by

BMIQ [56]. In further analysis, we removed probes containing SNPs ($n = 24\ 038$), X or Y

chromosome probes ($n = 16\ 263$), cross-hybridizing probes ($n = 46\ 867$) and polymorphic probes

($n = 297$) according to Chen et al. [8] and McCartney et al. [36], and probes with a detection p

value > .01 in at least one sample ($n = 79\ 922$). The batch effects were corrected for transformed

M-values using Combat [28]. For this purpose, we examined the strength of associations

between three possible batches (plate, slide, array) and the first five principal components via

principal component analysis. The strongest batch effects were iteratively removed. Sample mix-

ups were checked with MixupMapper [60]. The comparison of beta values with the existing

genotype data (assessed using Global Screening Arrays [GSA+MD-24v3.0-Psych-24v1.1,

Illumina, San Diego, CA, USA]; for further details see [9]) revealed four mix-ups. For these four

cases, both samples (first and second measurements) were removed (total of $n = 8$ individual

samples). The resulting DNA methylation dataset comprises 2 102 samples of $n = 1\ 055$

participants (among which $n = 488$ complete twin pairs or 976 individuals with two timepoints of

DNAm data) and 698 472 probes.

The cell type composition was estimated using the R package "EpiDish" (version 2.10;

[57]) and the epidish() function. Because saliva consists of three (highly correlated) cell type

estimates (leukocytes, epithelial cells, fibroblasts), we obtained the first principal component at

each occasion separately to control for cell type composition in the following analyses.

Calculating epigenetic age, epigenetic age acceleration, and pace of epigenetic aging

Four measures of epigenetic aging were calculated based on the normalized batch-

corrected beta values: the Horvath clock [19], the PedBE clock [37], GrimAge2 (an improved

version of the original GrimAge; [31,32]), and DunedinPACE estimate ([4]; Additional file 2:

Figure S1). The Horvath and PedBE epigenetic age were calculated using the R package

"methylclock" (version 1.01; [42]). A total of 334 (94.6%) of the original 353 Horvath clock

CpG sites and all 94 CpG sites of the PedBE clock were present in the dataset. Version 2 of

GrimAge was computed based on 1 029 CpG sites present in our dataset (out of 1 030 mentioned

in the publication) using analysis code from Lu et al. [31] provided by A. Lu and S. Horvath via

personal correspondence. The DunedinPACE values were calculated with the DunedinPACE R

package (https://github.com/danbelsky/DunedinPACE/) using all of 173 CpG sites.

The four epigenetic age estimates were significantly positively correlated with

chronological age (r(Horvath epigenetic age) = .80 ($p < .001$), r(PedBE epigenetic age) = .77 (p

$< .001$), r(GrimAge) = .80 ($p < .001$), and r(DunedinPACE) = .08 ($p < .001$); Additional file 2:

Figure S2). For the Horvath epigenetic age, the Median Absolute Error (MAE) estimates ranged

from 2.75 to 3.70 (3.26 to 3.65 for early adolescents, from 2.75 to 2.93 for late adolescents, from

3.33 to 3.34 for emerging adults, and from 2.89 to 3.71 for young adults). For the PedBE

epigenetic age, the MAE estimates ranged from 1.07 to 12.72 (2.22 to 2.91 for early adolescents,

from 1.08 to 2.01 for late adolescents, from 6.49 to 7.51 for emerging adults, and from 11.61 to

12.72 for young adults). For GrimAge, the MAEs ranged from 19.38 to 25.60 (22.96 to 24.38 for

early adolescents, from 25.42 to 25.60 for late adolescents, from 21.84 to 22.57 for emerging

adults, and from 19.39 to 20.25 for young adults; Additional file 2: Figure S3).

Acceleration estimates for the Horvath, PedBE, and GrimAge epigenetic ages were

obtained as follows: epigenetic age was regressed on chronological age (in years and decimal

months), biological sex, and cell-type composition within each measurement occasion. In

addition, the Horvath and PedBE epigenetic ages were regressed on smoking exposure, which

was estimated based on hypomethylation of the AHRR probe [43]. The resulting residuals (i.e.,

Horvath Acceleration, PedBE Acceleration, and GrimAge Acceleration), defined as "age

acceleration" (positive score) or "age deceleration" (negative score) according to the direction of

the deviation, were used in our analyses. The DunedinPACE values were controlled for smoking

probe, biological sex, and cell type composition. For DunedinPACE, a value of one reflects a

pace of aging in line with 1 year of chronological aging, with a value > 1 reflecting an increased

pace and values < 1 reflecting a decreased pace of aging.

Analytical strategy

To examine the role of genetic and environmental factors in epigenetic aging, biometrical

twin variance decomposition in structural equation modeling was applied [39]. MZ twins share

close to 100% of their genetic makeup, while DZ twins share, on average, 50% of genetic

variants that differ among humans. As a consequence, the correlation of genetic factors is

approximately 1 for MZ twins, whereas it is assumed to approach .5 for additive genetic factors

and .25 for non-additive genetic factors due to allelic dominance deviation for DZ twins.

Therefore, a greater correlation within MZ twin pairs than in DZ twin pairs signals the presence

of genetic influences on the variance in a specific phenotype, such as epigenetic aging.

Furthermore, both MZ and DZ twins are assumed to share some prenatal and familial factors [6]

that act to increase their resemblance beyond genetic factors to the same extent. Therefore, these

shared environmental factors contribute to the similarity within MZ twin pairs to the same

degree as within DZ twins. Only individual-specific unique environmental factors are assumed to

increase the dissimilarity between monozygotic twin siblings. These assumptions lie in the

essence of the classical biometrical model, which allows for a decomposition of the observable

variance into variance attributable to genetic factors (*A* – additive genetic and *D* – non-additive

genetic effects) and environmental influences (*C* – shared by twins and *E* – unique environment;

[6]). For identification purposes, only three sources of variance can be estimated simultaneously

in the classical twin model. Depending on the pattern of the correlations within the MZ and DZ

twins, either *D* (if $r\text{MZ} < 2 \times r\text{DZ}$) or *C* (if $r\text{MZ} > 2 \times r\text{DZ}$) components have to be fixed to zero,

resulting in the specification of either ACE or ADE models [12].

First, we started with the analysis of the between-individual differences and specified

univariate variance decomposition models for each epigenetic aging estimator. Four age groups

with two measures each produced eight unique timepoints analyzed in one model. We set the

variance of all latent variables (A, C/D, E) to 1 for identification purposes and ensured that the

statistical conditions specified for twin modeling, namely nonsignificant differences in means

and variances across twin order and zygosity, were met in our sample. Then, we decided whether

to use ACE or ADE models. After that, the model that fit to the data at best for each of the four

epigenetic aging measures was identified, and the standardized coefficients with 90% confidence

intervals were calculated. We have chosen among the ACE, AE, CE, and E models or,

alternatively, among the ADE, AE, and E models. Among the two models that did not differ

significantly based on the chi-square test, the model with a lower Akaike information criterion

(AIC) was considered to fit the data better. Among the two models that differed significantly

based on the chi-square test, the model with the higher number of parameters was considered as

the model that best fit the data [14].

In a second step, we tested whether the age of participants moderated genetic and

environmental contributions to the variance in epigenetic aging to unveil possible

gene☐environment interaction [44]. Since gene☐environment interaction effects cannot be

directly assessed in classical twin models, the estimates of genetic and environmental

contributions could be confounded with interaction effects of these two [44]. In the case of

interactions between additive genetic factors (*A*) and environmental factors that are shared by

siblings reared together (*C*), interaction effects would be confounded with estimates of additive

genetic effects. Potential interaction effects between genetic factors (*A*) and environmental

factors not shared by siblings reared together (*E*) would be confounded with estimates of

nonshared environmental effects. While the first interaction pattern is more plausible in younger

people, when twins are still living in a common household, the latter is more plausible in adult

years, when twins are living in different households and having more separate life experiences.

As a consequence, higher estimates of genetic contributions to the variance in epigenetic aging

are plausible in adolescence and emerging adulthood with declining trends across adulthood due

to accumulating effects of individually unique experiences that may or may not interact with

genetic factors.

Adding age as a moderator within the estimated structural equation models partitioned

the variance of ACE/ADE components into 3 paths of non-moderated ACE/ADE components, 3

paths of the moderated ACE/ADE components, age moderating effect on means, and

independent linear and quadratic effects of age, producing in total a model with 9 parameters.

The model was still identified as the saturated model had in total 10 measured parameters. We

started with the full models (with 9 estimated parameters) and then dropped nonsignificant paths

to obtain more parsimonious models with more degrees of freedom. We excluded parameters

until we reached the model that fit the data best, taking model parsimony into account (i.e., a

model with a lower AIC that did not fit the data significantly worse based on the chi-square test).

The variance components, confidence intervals, and standardized estimates were then derived.

To investigate the role of genetic and environmental sources of variance in within-

individual two-year changes, we first calculated the individual change in epigenetic aging

measures, obtained as the difference between the second and first measurements of epigenetic

aging (referred to as Δ in the following). The means and standard deviations for two-year

changes in four epigenetic aging measures within each age group were calculated. Then, we

repeated the statistical analysis described above. Similar to the analysis of between-individual

differences, we started with univariate variance decomposition models for each epigenetic aging

estimator and then examined the moderation of age. This resulted in four estimates of the genetic

and environmental factors across ages from 8 to 31 for each epigenetic aging estimator.

For sensitivity analysis, we scrutinized the models with age moderation further not

controlling the epigenetic aging measures either for sex, smoking probe, or cell type

composition. We repeated the age moderation analysis, first, for all chosen epigenetic aging

measures unadjusted for sex, second, for three epigenetic aging measures unadjusted for

smoking, and, finally, for four epigenetic aging measures unadjusted for cell type composition.

The analyses were performed in R studio 2023.06.2 with R version 4.2.3. For data

preparation, we used the R packages "tidyverse" [61], "ltm" [49], "dlyr" [16], and "psych" [48].

For the variance decomposition, we used "OpenMX" [40].

Prior power analysis

For the estimation of power to detect model misspecification, we applied the functions

acePow() and powerValue() proposed by Verhulst [59] to univariate ACE models. For that

purpose, we formulated two possible expectations regarding genetic factors (*A*) based on

estimates obtained in larger sample studies. Previous research suggested a contribution of *A* 13%

over the life course [29] and 37% at age 15 (combination with middle-aged mothers; [53]). Our

expectation regarding shared environmental factors (*C*) was based on plots for MZ and DZ twins

in Li et al. ([29]; page 8, Figure 4, cohabitation-dependent CE model). Here, we identified

approximated estimates of *C* for our sample's mean age of about 35% (for age 17.1). A power

calculation demonstrated that 976 pairs (526 MZ and 450 DZ twin pairs) in the between-

individual difference analysis and 488 pairs (263 MZ and 225 DZ twin pairs) in the within-

individual change analysis were sufficient to detect the genetic contributions mentioned in the

literature (13% or 37%), given that 35% of the variance was attributable to shared environmental

factors. The power for the analysis of between-individual differences was equal to .99 for both

expected genetic variance components. For the analysis of within-individual changes, the power

was good for higher estimates of genetic contributions to the variance (.96 given 37% of additive

genetic factors) and still acceptable for lower estimates of genetic contributions (.90 given 13%

of additive genetic factors).

Results

Descriptive analysis

The means and standard deviations for the epigenetic aging measures across age groups

are presented in Table 1, the twin correlations of epigenetic aging measures for MZ and DZ

twins for each timepoint are given in Table 2, and within-individual and within-twin pair rank-

order stability estimates for MZ and DZ twins are shown in Additional file 3: Table S2.

(please add Table 1 here)

(please add Table 2 here)

Trained on the chronological age, Horvath and PedBE Acceleration demonstrated

acceleration in the groups of late adolescents ($M = 16.6$) and emerging adults ($M = 22.7$) and

deceleration in two other groups of early adolescents ($M = 10.7$) and young adults ($M = 28.8$; see

Table 1). GrimAge Acceleration was higher in late adolescents with a mean age of 17.75,

emerging adults (aged from 20 to 29), and at the onset of adolescence. The estimates of

DunedinPACE showed a faster pace during age from 26 to 31 compared to other age periods.

For the two-year changes in epigenetic aging measures, we observed a greater positive change

for the period of late adolescence (change from 14 to 18 years of age) and a higher negative

change in the groups of early adolescents (change from 8 to 12 years of age) and young adults

(change from 26 to 31 years of age), which potentially reflects the growing within-individual

differences during these age periods.

Plotting the spread and locality of four epigenetic aging estimates across age groups

demonstrated patterns of increased variance (Table 1 and Additional file 3: Figure S4). Two

measures trained on chronological age, Horvath and PedBE Acceleration also showed a slightly

decreasing trend during young adulthood. In the eldest age group ($M = 29.92$) compared to the

youngest age group ($M = 9.51$), the standard deviation was 60% greater for Horvath

Acceleration, 43% greater for PedBE Acceleration, 29% greater for both GrimAge Acceleration

and DunedinPACE. For the individual two-year change in epigenetic aging, increasing standard

deviation was also observed for Δ Horvath (30%) and Δ PedBE (50%) Acceleration. The

standard deviation in Δ GrimAge Acceleration decreased by 9% and remained relatively stable

for Δ DunedinPACE.

Twin correlations for Horvath and PedBE Acceleration demonstrated substantial

similarity for adolescent twins, with only marginal differences between MZ and DZ twins (Table

2), indicating contributions of shared environmental influences to variance during this period of

life. In groups of emerging adults and later, the correlations for MZ twins were greater than the

correlations for DZ twins, indicating genetic influences on variance in epigenetic aging in

adulthood. In contrast, Spearman correlations for GrimAge Acceleration and DunedinPACE

were greater for MZ twins than for DZ twins across almost all age groups.

The within-individual rank-order stability across two measurement points was

approximately .50 with no differences between the MZ and DZ twin groups for all measures

except GrimAge Acceleration, where within-individual rank-order stability was lower in younger

age groups. MZ within-twin pair rank-order stability estimates tended to be greater than those of

DZ twins in adulthood (Additional file 3: Table S2), indicating declining contributions of shared

environmental factors and increasing genetic contributions to rank-order stability with age.

Finally, the twin correlations for two-year individual change in epigenetic aging were much

lower than the correlations regarding individual differences within measurement occasions for

different ages, suggesting either proneness to measurement error in the prediction of

chronological and biological age or the greater role of unique environmental factors contributing

to variance in within-individual changes than to between-individual differences.

Univariate twin model analysis of between-individual differences in epigenetic aging

403   To perform univariate variance decomposition analysis of the between-individual

differences in epigenetic aging measures, we first examined whether the statistical conditions

specified for twin modeling were met (Additional file 3: Table S3). We observed small

differences in means across twin siblings' order for Horvath Acceleration and DunedinPACE,

but the models with the expected means and variances to be equal across twin siblings' order and

zygosity did not fit the data significantly worse. For Horvath, PedBE and GrimAge Acceleration,

cross-cohort MZ twin correlations were not more than twice as strong as DZ twin correlations,

indicating that shared environmental factors (C) were more likely to contribute to the variance

than non-additive genetic factors: $r\text{MZ} = .51 < 2 \times r\text{DZ} = .74$ in the case of Horvath

Acceleration, $r\text{MZ}= .45 < 2 \times r\text{DZ} = .96$ in the case of PedBE Acceleration and $r\text{MZ} = .32 < 2 \times$

$r\text{DZ} = .34$ in the case of GrimAge Acceleration (see Table 2). The correlational patterns for MZ

twins and DZ twins for DunedinPACE indicated, besides additive genetic factors, the presence

of non-additive genetic factors (D): $r\text{MZ} = .48 > 2 \times r\text{DZ} = .38$. Therefore, we focused on ACE

variance decomposition in the following analysis for Horvath, PedBE, and GrimAge

Acceleration and on ADE variance decomposition for DunedinPACE.

The results of the univariate twin model analyses of the epigenetic aging measures are

presented in Figure 2 and Additional file 3: Tables S4-5. For Horvath Acceleration, we observed

a better fit of the full model (ACE), indicating that the variance in Horvath Acceleration was best

explained by additive genetic as well as unique and shared environmental factors. PedBE

Acceleration, GrimAge Acceleration, and DunedinPACE estimates in the univariate analysis

were better represented by restricted models (the CE model for PedBE Acceleration and the AE

models for GrimAge Acceleration and DunedinPACE). The variation in PedBE acceleration was

thus best explained by shared and unique environmental factors, whereas the variance in

GrimAge Acceleration and DunedinPACE was best explained by additive genetic and unique

environmental factors.

Figure 2

Cross-cohort standardized genetic and environmental variance components of epigenetic aging

measures

Note. The estimates are derived from the best-fitting models in the univariate twin model

analysis for between-individual differences (A) and variance in two-year changes (B) in

epigenetic aging measures.

Age moderation of genetic and environmental between-individual differences in epigenetic

aging

Then, we explored whether the relative contributions of the *A*, *C* (or *D*), and *E*

components to the between-individual differences of epigenetic aging differed across age groups

(Table 3, Figure 3, Additional file 3: Tables S6-7, Figure S5). Here, we only present the models

that fit the data best (the model fit statistics can be found in Additional file 3: Table S6).

442    When moderation by age in the between-individual differences models was allowed,

443 moderation was observed for all epigenetic aging measures. For acceleration measures trained on

444 chronological age, Horvath and PedBE Acceleration, we observed the moderation of the unique

445 environmental component (E), which increased for both estimators. The proportion of the unique

446 environmental factors in the between-individual variance of the Horvath and PedBE

447 Acceleration increased from 47% to 49% and from 33% to 68% respectively. Moderation of the

448 variance attributable to genetic components, which increased from early adolescence to

449 adulthood, was found for Horvath and GrimAge Acceleration (increasing additive genetic

450 component) and for DunedinPACE (increasing non-additive genetic component). The narrow-

451 sense heritability increased from 18% to 39% for Horvath Acceleration and from 24% to 43%

452 for GrimAge Acceleration. The broad-sense heritability ($A + D$) of DunedinPACE increased

453 from 42% to 57%.

454    *(please add Table 3 here)*

455 **Figure 3**

456 *Unstandardized and standardized variance components derived from age-moderated models of*

457 *between-individual differences*

458 *Note.* The estimates are derived from the best-fitting univariate models of age moderation of

459 genetic and environmental components of between-individual differences in the four epigenetic

460 aging measures. Unstandardized variance components moderated by age are depicted as curves

461 with filled areas.

463 **Univariate twin model analysis of variance in two-year changes of epigenetic aging**

464    In a further step, we decomposed the variance in two-year within-individual changes,

465 quantified as the difference in epigenetic aging measures across two timepoints (Figure 2,

466 Additional file 3: Tables S8-9). The models with the expected means and variances to be equal

across twin order and zygosity did not fit the data significantly worse than did the saturated

models (Additional file 3: Table S3) confirming that the statistical conditions specified for twin

modeling were met. MZ correlations were not more than twice as large as DZ correlations

indicating the possible contributions of shared environmental factors (*C*) rather than non-additive

genetic factors (*D*) to the variance in the four epigenetic aging measures.

The results showed that the variance of two-year within-individual changes in the

epigenetic acceleration measures, Δ Horvath, Δ GrimAge, and Δ PedBE Acceleration, was best

explained by shared and unique environmental factors (CE model), whereas the variation in two-

year changes of DunedinPACE was best explained by additive genetic and unique environmental

factors (AE model). For all epigenetic aging measures, however, unique environmental factors

not shared by twins were found to be the primary source of differences in changes in epigenetic

aging (Additional file 3: Table S9). It is important to note that, in contrast to the a priori power

analysis, we did not have enough power to distinguish between the CE and AE models and

between the CE and E models for Δ Horvath and Δ GrimAge Acceleration.

Age moderation of genetic and environmental variance in two-year changes in epigenetic

aging

In the final step, we investigated whether the genetic and environmental contributions to

the variance in the two-year changes in epigenetic aging estimates varied with chronological age

(Table 4, Figure 4, Additional file 3: Tables S10-11, Figure S6). For three epigenetic

acceleration measures, Δ Horvath, Δ PedBE, and Δ GrimAge, the models with age moderation of

the non-genetic component demonstrated a better fit to the data. It was the AE model with the

moderation of the unique environment component (*E*; increasing trend) for Δ Horvath

Acceleration, the CE model including moderation of the shared environmental component (*C*;

decreasing trend) and the unique environment component (*E*; increasing trend) for Δ PedBE

Acceleration, and the CE model including moderation of the unique environment component (E;

decreasing trend) for Δ GrimAge Acceleration. In two models (Δ Horvath and Δ PedBE

Acceleration), we had sufficient power to identify the presence of the moderation of unique

environmental contributions. For Δ DunedinPACE, the CE model without moderation of the

genetic and environmental factors showed no worse fit to the data, possibly reflecting the

difference in the estimations of epigenetic age acceleration and pace of epigenetic aging.

(please add Table 4 here)

Figure 4

Unstandardized and standardized variance components derived from age-moderated models of

the two-year changes

Note. The estimates are derived from the best-fitting univariate models of age moderation of

genetic and environmental components of two-year changes in the four epigenetic aging

measures. Unstandardized variance components moderated by age are depicted as curves with

filled areas.

Sensitivity analysis

For sensitivity analysis, we performed three separate analyses of age moderation of between-

individual differences and variance in within-individual changes in epigenetic aging measures

unadjusted either for biological sex, a smoking probe, or cell-type composition. The model's

goodness-of-fit statistics and the estimates of the squared standardized coefficients are presented

in the Additional file 4. Noncontrolling for sex or smoking did not affect the conclusions of the

observed moderation in the main analysis, with the exclusion of two models for within-

individual changes in epigenetic aging where the power to distinguish between models was

limited (Δ PedBE Acceleration and Δ DunedinPACE, Additional file 4: Tables S12-19). Not

controlling for cell type composition obscured some of the main analysis findings (Additional

file 4: Tables S20-23). While the results were robust for Horvath Acceleration, not accounting

for cell type composition in between-individual differences introduced moderation of the unique

environment component for other epigenetic aging measures, eliminating the moderation of

genetic components for GrimAge Acceleration and DunedinPACE. The analysis of the within-

individual differences not controlled for cell type composition confirmed the moderation of

unique environmental contributions to variance in Δ PedBE Acceleration, introduced it to

DunedinPACE variance, and eliminated it in Δ GrimAge Acceleration. The analysis of

epigenetic aging measures unadjusted for cell type composition did not reach sufficient power to

distinguish between models with or without moderation. The observed pattern is potentially

explained by the fact that different tissues have heterogeneous cell type compositions [55],

which, in turn, can have a strong independent association with age [62]. Therefore, the

unadjusted for cell-type composition measure of epigenetic aging reflects, in addition to the

genuine age-associated signal of DNAm, heterogeneity in different cell types regarding DNAm,

and the link between age and cell-type composition.

Discussion

In this study, we investigated age-associated differences in genetic and environmental

contributions to the variance of epigenetic aging across adolescence and emerging adulthood.

Two main results can be reported. First, common trends of increasing between-individual

differences in epigenetic aging from adolescence to young adulthood were driven by both

environmental and genetic influences, while individual deviations from this general trend of

variance expansion were the result of unique environmental factors. Second, the proportions of

genetic and environmental contributions to the variance varied for different epigenetic aging

estimators.

Unique environmental contributions to age-associated differences in epigenetic estimators

tracking chronological age and biological aging.

The results of the analysis suggest that during adolescence and emerging adulthood

between-individual differences and variance in within-individual changes in epigenetic aging

estimators trained on chronological age are driven by unique experiences. One explanation for

these findings is the accumulated effect of unique life experiences. Childhood and adolescence

are a sequence of sensitive and critical periods in development, during which the effects of

positive and negative factors can be particularly influential. Experiences of violence in children

or institutional caring [21], for example, have already been mentioned to predict greater

epigenetic age, affecting ongoing brain and emotional regulation development [51]. An

alternative explanation for our findings might be the *epigenetic drift*, stochastic changes in

DNAm. These random changes in DNAm have been shown to contribute to progressively

increasing epigenetic differences between genetically identical MZ twins over time [13]. When

epigenetic drift happens in epigenetic estimators' locus-specific DNAm, the increase in unique

environmental factors estimated in twin modeling would partly reflect epigenetic drift. This

would also explain the greater power of epigenetic age to predict chronological age at the onset

of life compared to childhood and adolescence [37, 53].

While we observed an increasing unique environmental contribution to the variance of

the epigenetic aging measures tracking chronological age, unique environmental contributions

did not increase in the epigenetic aging estimators trained on biological health indicators. For

GrimAge Acceleration, we did not observe moderation of the unique environmental component

in between-individual differences, while the environmentally driven variance in individual two-

year changes demonstrated a decreasing trend. Consistently, no age-moderation of unique

environmental factors was found for the pace of epigenetic aging DunedinPACE in either of the

two analyses. In contrast to epigenetic aging estimators trained on chronological age, the two

later measures were developed in (exclusively adult) blood samples and rely on intrinsic

biological functioning. In particular, GrimAge is linked to the level of plasma proteins [31] and

DunedinPACE reflects among other factors, the level of glycated hemoglobin, leptin, and serum

C-reactive proteins [4]. Because they are also intrinsically driven, these processes may be more

elusive and less sensitive to trends in unique environmental contributions when they are

measured in saliva.

The absence of age-moderation of unique environmental factors in DunedinPACE could

also signal the difference between training designs across epigenetic aging estimators. While

GrimAge, Horvath, and PedBE clocks were developed by analyzing cross-sectional data,

DunedinPACE estimate was trained on longitudinal data, accounting for the within-individual

trends of aging. Representing the speed of aging based on 12 months, DunedinPACE does not

directly relate to chronological age and the accumulation of aging processes, which are

commonly represented by acceleration measures. This could also explain the observed trends in

the variances in epigenetic aging across age groups. In our analysis, late adolescents and

emerging adults both demonstrated increasing variance in epigenetic age acceleration measures.

In particular, the between-individual differences increased up to the ages 20-24 for Horvath and

PedBE Acceleration and continued to do so in the older age group for GrimAge Acceleration. A

similar pattern was found for Δ Horvath and Δ PedBE Acceleration. In contrast, for

DunedinPACE, we observed, if at all, only a slight increase in variance. This was in line with the

previous findings of the remarkable stability of DunedinPoAm, the predecessor of

DunedinPACE, for the age period from 7 to 20 [41]. The smaller changes in variance and the

absence of moderation in DunedinPACE suggest that the roles of unique environmental factors

in the pace of epigenetic aging are potentially more stable than the environmental contributions

to epigenetic acceleration measures.

The genetic contribution to age-associated between-individual differences in epigenetic

acceleration and the pace of aging

Three out of four estimators, Horvath Acceleration, GrimAge Acceleration, and

DunedinPACE, trained in adults, demonstrated an increase in the variance of between-individual

differences across age due to genetic (additive or non-additive) factors. First, this pattern is

consistent with longitudinal studies of the SNP-heritability of epigenetic aging and studies of

complex phenotypes. SNP-heritability in children and their parents demonstrated that, starting at

0% during the onset of life, it increases up to 43% later in life [34, 53]. Similarly, genetic

contributions to the variance in complex phenotypes, such as intelligence, depressive symptoms,

and personality traits, demonstrate increasing heritability from the transition to adolescence and

adulthood [5, 24, 38].

The observed phenomenon of the increased role of genetic factors during this age is

commonly explained by active and evocative gene☐environment correlation in the presence of

gene☐environment interaction, "…when people become more and more autonomous and self-

directed during adolescence. They have increasing opportunities to pave their own way, evoke,

select and create environments that match their heritable tendencies. These environments in turn

can provide experiences that have the potential to reinforce the pre-existing tendencies" ([24]; p.

238). One illustrative example of an active gene☐environment correlation is the selection of the

suitable environments by people that fit their genetic predispositions that prevail over the passive

gene☐environment correlation effects created by the genetic background of parents [5, 23]. This

would happen when, for instance, individuals with genetic predispositions for a slower

epigenetic aging rather develop healthy habits and search for communities of similar individuals,

eventually supporting and exacerbating this genetic tendency. As the increasing importance of

active gene☐environment correlation would come along with an increase of genetic differences,

while a declining role of passive gene☐environment correlation would be reflected by

decreasing effects of shared environmental influences [7, 44], our findings can be considered in

line with this interpretation.

Alternatively, the gene☐environment interplay underlying estimates of genetic factors

can reflect the interaction of genetic and shared environmental factors [24, 44], which is more

plausible for twins reared together and, thus, more probable for them during adolescence, when

they still share a common family household. Previous studies illuminating the role of shared

environmental factors [30] prioritized the role of the prenatal environment and the driving role of

cohabitation in epigenetic aging, implying that the longer relatives live together, the more similar

they were epigenetically [29]. From adolescence to adulthood, the role of these factors seems to

be steadily replaced by genetic and unique environmental forces as well as by interactions

between them. As cohabiting is more common before adulthood, our findings extend previous

conclusions, implying an increasing role of active gene☐environment correlation in the presence

of interactions between genetic and environmental factors, when relatives, on average, transit to

adulthood and tend to start living separately.

The transition to adulthood could explain why an increasing trend of genetic moderation

was observed for all epigenetic aging measures but not for PedBE Acceleration. The age

composition of the initial training samples in the chosen epigenetic estimators differed. In

contrast to the PedBE clock, which was initially developed in a sample of individuals up to age

20, the Horvath clock, GrimAge, and DunedinPACE were trained in adults. Therefore, the

increasing genetic contributions we observed for these three epigenetic measures might

potentially characterize the intrinsic epigenetic processes that are activated later in life. These

processes can include the trajectories of the physical development, for example, the transition

from puberty to the maintenance of the biological systems and the high organism functionality.

Alternatively, these processes can reflect the trajectories in gene☐environment interaction and

the relationships between individuals and their immediate environments. This would include, for

instance, the transition from the sorting into fitting environments to the active creation and

modification of environments or finding a balance between person☐environment fitting and

adjustment [23]. Because these processes are less pronounced during adolescence, they may be

not fully reflected in the genetic contributions to the variance in PedBE Acceleration.

The heritability of epigenetic aging based on saliva samples

The age-specific pattern of genetic contributions to the variance in epigenetic aging in

saliva obtained in our analysis was rather consistent with the assessments reported in blood.

However, our heritability estimates were, on average, lower than those reported in studies based

on blood tissue samples. For example, previous research reported that genetic differences

accounted for 69%, 73%, and 68% of the variance in Horvath Acceleration, GrimAge

Acceleration, and DunedinPACE in 22-year-olds, respectively [26]. The broad-sense heritability

for 21.5-year old twins in our analysis was substantially lower (32%, 35%, and 51%

respectively) (see Additional file 3: Figure S7). This difference in results could be related to

variations in the age compositions of the studies' samples, the presence of age moderation, or the

genuine differences in epigenetic aging across tissues [63]. Therefore, further studies are

required to investigate whether the observed pattern of the lower heritability of epigenetic aging

based on saliva can be replicated.

Limitations and future research

Among the strengths of our study are the relatively large sample size, the focus on both

epigenetic age acceleration and the pace of epigenetic aging measures, the examination of

additive and non-additive genetic sources, and, finally, the consideration of age differences

during the second and third decades of life. Nevertheless, our study is not without limitations.

First, despite the advantage of a comparatively large sample size with epigenetic data and

usage of both measurement points in the biometrical variance decomposition, the post hoc power

analysis demonstrated that the differentiation between several nested models with age as a

moderator was limited, as confidence intervals overlapped. Earlier, power analyses in twin

studies have suggested a minimum sample size of 600 twin pairs for univariate analysis and

larger samples for detecting possible non-additive genetic factors, which, unfortunately, could

not be reached [35, 50]. Nevertheless, we had enough power to detect the presence of genetic

and environmental age moderation in the between-individual differences in epigenetic measures

and the moderation of unique environmental factors contributing to the variance in within-

individual two-year changes for epigenetic age acceleration.

Second, differences in the performance of epigenetic aging measures across different

ages and tissues could have introduced additional sources of measurement error. The Horvath

clock, for example, is less precise in younger samples, while the PedBE clock could be more

biased in adults [37]. Similarly, our study exclusively utilized saliva samples, but not all applied

epigenetic aging estimators were trained on saliva-based methylation profiles. While the Horvath

clock is a pantissue epigenetic clock and the PedBE clock was developed in buccal cells,

DunedinPACE and GrimAge were trained in blood. Although the pace of aging can be assessed

based on saliva [46, 47] and the second version of GrimAge demonstrated good performance in

saliva methylation data [31], applying these epigenetic estimators in a tissue different from the

initial design may affect the estimates. The Horvath clock, at the same time, being a pantissue

clock, could lose some precision when applied to any tissue.

Third, the utilization of measurements from two timepoints in four age groups could

cause additional sources of variance. Different age groups, for instance, might introduce cohort

effects that cannot be entirely disentangled from age effects. In addition, the second

measurement occurred during the COVID-19 pandemic. While other research groups have not

confirmed the effect of the COVID-19 pandemic on epigenetic aging in adolescents [10], we

assume that changes in health status, lockdowns, and restrictions are potential new influential

environmental sources. Hence, the role of these factors in epigenetic aging dynamics should be

clarified further.

Fourth, classical twin analysis is an efficient way to estimate genetic and environmental

sources of variance but at the cost of several assumptions. One assumption is that environmental

experiences shared by twins contribute to the similarity within MZ twin pairs to the same degree

as within DZ twin pairs. One study showed that this assumption might not hold true for

epigenetic aging, as MZ and DZ pairs can possibly share their preadult environmental effects to

different extents [29]. In addition, the classical twin design does not allow, so far, the estimation

of shared environmental (C) and non-additive genetic (D) contributions in one model, while both

sources of variance can be relevant for variance in epigenetic aging at the same or during

different developmental stages. These aspects should be addressed in further investigations.

Conclusion

Altogether, the present study demonstrated that the role of genetic and environmental

sources of variance in epigenetic aging is not constant across adolescence and emerging

adulthood. Increasing between-individual differences in epigenetic aging during the transition to

adulthood were rather driven by increasing contributions of genetic and individually unique

environmental factors. The individual trajectories of epigenetic aging were mostly independent

of genetic sources and primarily driven by unique individual experiences. These findings

contribute to a better understanding of the epigenetic aging process during the transition to

adulthood.

Declarations

Ethics approval and consent to participate

The TwinLife study was reviewed and approved by the German Psychological Society (Deutsche Gesellschaft für Psychologie; protocol number: RR 11.2009). The TwinSNPs project received ethical approval from the Ethical Committee of Bielefeld University (No. 2020-180) and the TwinSNPs and TECS projects received ethical approval from the Ethical Committee of the Medical Faculty of the University of Bonn (Lfd. Nr. 113/18). All participants provided written informed consent. Participation was voluntary and participants could withhold participation at any timepoint. The personal data collection and analysis complied with the corresponding data protection regulations. The statistical analysis was performed on pseudonymized data from the servers of Bielefeld University.

Availability of data and materials

The TwinLife data are archived in the GESIS data catalog: https://search.gesis.org/research_data/ZA6701. The data are released for academic research and teaching after the data depositor's written authorization. The epigenetic data are available for collaborators via TwinLife data management.

Funding

This work was supported by the German Research Foundation (DFG) as a part of the long-term project TwinLife under Grant [number 220286500], the project "Genetic and social causes of social inequalities: Integration of molecular genetic data into the longitudinal twin family study TwinLife" (TwinSNPs) under Grant [number 428902522], and the TwinLife Epigenetic Change Satellite (TECS) project under Grant [number 458609264].

Competing interests

The authors report that there are no competing interests to declare.

Authors' contributions

DVK & BM were involved in the study conception and design. DVK analyzed the data and

interpreted the results. DVK, YL, AMS, and CK drafted the manuscript. DVK, YL, AMS, DC,

JI, CKLP, MMN, EBB, MD, AJF, CK, and BM provided valuable feedback for the manuscript

revision. MD & MMN were responsible for obtaining ethical approval. MD, BM, MMN, AJF,

CKLP, FMS, EBB, DC, and CK collected the data. AMS, DC, DVK, and CKLP assessed DNA

methylation and generated the epigenetic aging scores. BM supervised the study. All the authors

have read and approved the final manuscript.

Acknowledgments

We thank M. Ruks, A. Kottwitz, and K. Krell from TwinLife data management team for their

support; S. Zare, A. Vogelsang, and N. Fricker for their contribution to the DNA extraction and

epigenetic profiling; M. Ruks and L. Weigel for valuable advice on the manuscript.

References

1.   Alisch RS, Barwick BG, Chopra P, Myrick LK, Satten GA, Conneely KN, et al. Age-

associated DNA methylation in pediatric populations. Genome Research.

2012;22(4):623–32. https://doi.org/10.1101/gr.125187.111

2.   Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al.

Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium

DNA methylation microarrays. Bioinformatics. 2014;30(10):1363–9.

https://doi.org/10.1093/bioinformatics/btu049

3.   Belsky DW, Caspi A, Arseneault L, Baccarelli A, Corcoran DL, Gao X, et al.

Quantification of the pace of biological aging in humans through a blood test, the

DunedinPoAm DNA methylation algorithm. eLife. 2020;9,e54870.

https://doi.org/10.7554/eLife.54870

4. Belsky DW, Caspi A, Corcoran DL, Sugden K, Poulton R, Arseneault L, et al.

DunedinPACE, a DNA methylation biomarker of the pace of aging. Elife.

2022;11:e73420. https://doi.org/10.7554/eLife.73420

5. Bergen SE, Gardner CO, Kendler KS. Age-Related Changes in Heritability of Behavioral

Phenotypes Over Adolescence and Young Adulthood: A Meta-Analysis. Twin Research

and Human Genetics. 2007;10(3):423–33. https://doi.org/10.1375/twin.10.3.423

6. Boomsma D, Busjahn A, Peltonen L. Classical twin studies and beyond. Nature Reviews

Genetics. 2002;3(11):872–82. https://doi.org/10.1038/nrg932

7. Briley DA, Livengood J, Derringer J. Behavior genetic frameworks of causal reasoning

for personality psychology. European Journal of Personality. 2018;32:202–20.

https://doi.org/10.1002/per.2153

8. Chen Y, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, et al.

Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium

HumanMethylation450 microarray. Epigenetics. 2013;8(2):203–9.

https://doi.org/10.4161/epi.23470

9. Deppe M, Pahnke CKL, Maj C, Aldisi R, Nöthen M, Diewald M, et al. Is happiness

always a personal(ity) thing? A quasi-replication and extension of previous well-being

studies. [Internet]. 2024 [cited 2024 Jun 3]. Available from: https://osf.io/6ctm4

10. deSteiguer AJ, Raffington L, Sabhlok A, Tanksley P, Tucker-Drob EM, Harden K.P.

Stability of DNA-Methylation Profiles of Biological Aging in Children and Adolescents

[Preprint]. Developmental Biology; 2023. Available from:

http://biorxiv.org/lookup/doi/10.1101/2023.10.30.564766

11. Diewald M, Kandler C, Riemann R, Spinath FM, Andreas A, Baier T, et al.

TwinLifeTwinLife (6.1.0) [dataset]. GESIS; 2022. Available from:

https://search.gesis.org/research_data/ZA6701?doi=10.4232/1.13987

12. Dolan CV, Boomsma DI, Nivard MG, Neal MC. A Note on Jöreskog's ACDE Twin

Model: A Solution, Not the Solution. Structural Equation Modeling: A Multidisciplinary

Journal. 2022;29(6):933–4. https://doi.org/10.1080/10705511.2022.2095640

13. Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, et al. Epigenetic

differences arise during the lifetime of monozygotic twins. Proceedings of the National

Academy of Sciences. 2005;102(30):10604–9. https://doi.org/10.1073/pnas.0500398102

14. Gana K, Broc G. Structural equation modeling with lavaan. London□: Hoboken, NJ:

ISTE Ltd□; John Wilery & Sons, Inc; 2019.

15. Gaunt TR, Shihab HA, Hemani G, Min JL, Woodward G, Lyttleton O, et al. Systematic

identification of genetic influences on methylation across the human life course. Genome

Biology. 2016;17(1):61. https://doi.org/10.1186/s13059-016-0926-z

16. Hadley W, Romain F, Lionel H, Kirill M, Davis V. dplyr: A Grammar of Data

Manipulation [Software]. 2023. Available from: https://dplyr.tidyverse.org,

https://github.com/tidyverse/dplyr

17. Hagenbeek FA, Hirzinger JS, Breunig S, Bruins S, Kuznetsov DV, Schut K, et al.

Maximizing the value of twin studies in health and behaviour. Nature Human Behaviour.

2023;7(6):849–60. https://doi.org/10.1038/s41562-023-01609-6

18. Hahn E, Gottschling J, Bleidorn W, Kandler C, Spengler M, Kornadt AE, et al. What

Drives the Development of Social Inequality Over the Life Course? The German

TwinLife Study. Twin Research and Human Genetics. 2016;19(6):659–72.

https://doi.org/10.1017/thg.2016.76

19. Horvath S. DNA methylation age of human tissues and cell types. Genome Biology.

2013;14(10):R115. https://doi.org/10.1186/gb-2013-14-10-r115

20. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of

ageing. Nature Reviews Genetics. 2018;19(6):371–84. https://doi.org/10.1038/s41576-

018-0004-3

21. Jovanovic T, Vance LA, Cross D, Knight AK, Kilaru V, Michopoulos V, et al. Exposure

to Violence Accelerates Epigenetic Aging in Children. Scientific Reports.

2017;7(1):8962. https://doi.org/10.1038/s41598-017-09235-9

22. Jylhävä J, Hjelmborg J, Soerensen M, Munoz E, Tan Q, Kuja-Halkola R, et al.

Longitudinal changes in the genetic and environmental influences on the epigenetic

clocks across old age: Evidence from two twin cohorts. EBioMedicine. 2019;40:710–6.

https://doi.org/10.1016/j.ebiom.2019.01.040

23. Kandler C, Kühn S, Mönkediek B, Forstner AJ, Bleidorn W. A Multidisciplinary

Perspective on Person-Environment Fit: Relevance, Measurement, and Future Directions.

Current Directions in Psychological Science. 2024;0(0).

https://doi.org/10.1177/09637214241242451

24. Kandler C, Waaktaar T, Mõttus R, Riemann R, Torgersen S. Unravelling the interplay

between genetic and environmental contributions in the unfolding of personality

differences from early adolescence to young adulthood. European Journal of Personality.

2019;33:221–44. http://dx.doi.org/10.1002/per.2189

25. Kankaanpää A, Tolvanen A, Bollepalli S, Leskinen T, Kujala UM, Kaprio J, et al.

Leisure-Time and Occupational Physical Activity Associates Differently with Epigenetic

Aging. Medicine & Science in Sports & Exercise. 2021;53(3):487–95.

https://doi.org/10.1249/MSS.0000000000002498

26. Kankaanpää A, Tolvanen A, Heikkinen A, Kaprio J, Ollikainen M, Sillanpää E. The role

of adolescent lifestyle habits in biological aging: A prospective twin study. eLife.

2022;11:e80729. https://doi.org/10.7554/eLife.80729

27. Kilpinen H, Dermitzakis ET. Genetic and epigenetic contribution to complex traits.

Human Molecular Genetics. 2012;21(R1):R24–8. https://doi.org/10.1093/hmg/dds383

28. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing

batch effects and other unwanted variation in high-throughput experiments.

Bioinformatics. 2012;28(6):882–3. https://doi.org/10.1093/bioinformatics/bts034

29. Li S, Nguyen TL, Wong EM, Dugué PA, Dite GS, Armstrong NJ, et al. Genetic and

environmental causes of variation in epigenetic aging across the lifespan. Clinical

Epigenetics. 2020;12(1):158. https://doi.org/10.1186/s13148-020-00950-1

30. Li S, Wong EM, Dugué PA, McRae AF, Kim E, Joo JHE, et al. Genome-wide average

DNA methylation is determined in utero. International Journal of Epidemiology.

2018;47(3):908–16. https://doi.org/10.1093/ije/dyy028

31. Lu AT, Binder AM, Zhang J, Yan Q, Reiner AP, Cox SR, et al. DNA methylation

GrimAge version 2. Aging. 2022. Available from: https://www.aging-

us.com/lookup/doi/10.18632/aging.204434

32. Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, et al. DNA methylation

GrimAge strongly predicts lifespan and healthspan. Aging. 2019;11(2):303–27.

https://doi.org/10.18632/aging.101684

33. Maksimovic J, Phipson B, Oshlack A. A cross-package Bioconductor workflow for

analysing methylation array data. F1000Research. 2017;5:1281.

https://doi.org/10.12688/f1000research.8839.3

34. Marioni RE, Shah S, McRae AF, Chen BH, Colicino E, Harris SE, et al. DNA

methylation age of blood predicts all-cause mortality in later life. Genome Biology.

2015;16(1):25. https://doi.org/10.1186/s13059-015-0584-6

35. Martin NG, Eaves LJ, Kearsey MJ, Davies P. The power of the classical twin study.

Heredity. 1978;40(1):97–116. https://doi.org/10.1038/hdy.1978.10

36. McCartney DL, Walker RM., Morris SW, McIntosh AM, Porteous DJ, Evans KL.

Identification of polymorphic and off-target probe binding sites on the Illumina Infinium

MethylationEPIC BeadChip. Genomics Data. 2016;9:22–4.

https://doi.org/10.1016/j.gdata.2016.05.012

37. McEwen LM, O'Donnell KJ, McGill MG, Edgar RD, Jones MJ, MacIsaac JL, et al.

(2020). The PedBE clock accurately estimates DNA methylation age in pediatric buccal

cells. Proceedings of the National Academy of Sciences. 2020;117(38):23329–35.

https://doi.org/10.1073/pnas.1820843116

38. Mõttus R, Briley DA, Zheng A, Mann FD, Engelhardt LE, Tackett, JL, et al. Kids

becoming less alike: A behavioral genetic analysis of developmental increases in

personality variance from childhood to adolescence. Journal of Personality and Social

Psychology. 2019;117:635–58. http://dx.doi.org/10.1037/pspp0000194

39. Neale MC. Biometrical Models in Behavioral Genetics. In: Kim YK, editor. Handbook of

Behavior Genetics. New York, NY: Springer New York; 2009. p. 15–33.

https://doi.org/10.1007/978-0-387-76727-7_2

40. Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM. OpenMx 2.0:

Extended Structural Equation and Statistical Modeling. Psychometrika. 2016;81(2):535–

49. https://doi.org/10.1007/s11336-014-9435-8

41. Niccodemi G, Menta G, Turner J, D'Ambrosio C. Pace of aging, family environment and

cognitive skills in children and adolescents. SSM - Population Health. 2022;20:101280.

https://doi.org/10.1016/j.ssmph.2022.101280

42. Pelegí-Sisó D, de Prado P, Ronkainen J, Bustamante M, González JR. methylclock: A

Bioconductor package to estimate DNA methylation age. Bioinformatics.

2021;37(12):1759–60. https://doi.org/10.1093/bioinformatics/btaa825

43. Philibert R, Dogan M, Beach SRH, Mills JA, Long JD. AHRR methylation predicts

smoking status and smoking intensity in both saliva and blood DNA. American Journal

of Medical Genetics Part B: Neuropsychiatric Genetics. 2020;183(1):51–60.

https://doi.org/10.1002/ajmg.b.32760

44. Purcell S. Variance Components Models for Gene–Environment Interaction in Twin

Analysis. Twin Research. 2002;5(6):554–71.

https://doi.org/10.1375/136905202762342026

45. Raffington L, Belsky DW. Integrating DNA Methylation Measures of Biological Aging

into Social Determinants of Health Research. Current Environmental Health Reports.

2022;9(2):196–210. https://doi.org/10.1007/s40572-022-00338-8

46. Raffington L, Tanksley PT, Sabhlok A, Vinnik L, Mallard T, King LS, et al. Socially

Stratified Epigenetic Profiles Are Associated With Cognitive Functioning in Children

and Adolescents. Psychological Science. 2023;34(2):170–85.

https://doi.org/10.1177/09567976221122760

47. Raffington L, Tanksley PT, Vinnik L, Sabhlok A, Patterson MW, Mallard T, et al.

Associations of DNA-Methylation Measures of Biological Aging With Social Disparities

in Child and Adolescent Mental Health. Clinical Psychological Science. 2023;0(0).

https://doi.org/10.1177/21677026231186802

48. Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research

(R package version 2.3.6) [Software]. Evanston, Illinois: Northwestern University; 2023.

Available from: https://CRAN.R-project.org/package=psych

49. Rizopoulos D. ltm: An R Package for Latent Variable Modeling and Item Response

Theory Analyses. Journal of Statistical Software. 2006;17(5). Available from:

http://www.jstatsoft.org/v17/i05/. https://doi.org/10.18637/jss.v017.i05

50. Sham PC, Purcell SM, Cherny SS, Neale MC, Neale BM. Statistical Power and the

Classical Twin Design. Twin Research and Human Genetics. 2020;23(2):87–9.

https://doi.org/10.1017/thg.2020.46

51. Shonkoff JP, Garner AS. The Lifelong Effects of Early Childhood Adversity and Toxic

Stress. Pediatrics. 2012;129(1):e232–e246. https://doi.org/10.1542/peds.2011-2663

52. Sillanpää E, Ollikainen M, Kaprio J, Wang X, Leskinen T, Kujala UM, et al. Leisure-

time physical activity and DNA methylation age—A twin study. Clinical Epigenetics.

2019;11(1):12. https://doi.org/10.1186/s13148-019-0613-5

53. Simpkin AJ, Hemani G, Suderman M, Gaunt TR, Lyttleton O, Mcardle WL, et al.

Prenatal and early life influences on epigenetic age in children: A study of mother–

offspring pairs from two cohort studies. Human Molecular Genetics. 2016;25(1):191–

201. https://doi.org/10.1093/hmg/ddv456

54. Sinke L, Cats D, Heijmans BT. Omixer: Multivariate and reproducible sample

randomization to proactively counter batch effects in omics studies. Bioinformatics.

2021;37(18):3051–2. https://doi.org/10.1093/bioinformatics/btab159

55. Teschendorff AE. Cell-Type Heterogeneity in DNA Methylation Studies: Statistical

Methods and Guidelines. In: Michels KB, editor. Epigenetic Epidemiology. Springer

International Publishing; 2022. p. 67–96. https://doi.org/10.1007/978-3-030-94475-9_4

56. Teschendorff AE, Marabita F, Lechner M, Bartlett T, Tegner J, Gomez-Cabrero D, et al.

A beta-mixture quantile normalization method for correcting probe design bias in

Illumina Infinium 450 k DNA methylation data. Bioinformatics. 2013;29(2):189–96.

https://doi.org/10.1093/bioinformatics/bts680

57. Teschendorff AE, Zheng SC. EpiDISH [Software]. Bioconductor; 2017. Available from:

https://doi.org/10.18129/B9.BIOC.EPIDISH

58. Touleimat N, Tost J. Complete pipeline for Infinium ® Human Methylation 450K

BeadChip data processing using subset quantile normalization for accurate DNA

methylation estimation. Epigenomics. 2012;4(3):325–41.

https://doi.org/10.2217/epi.12.21

59. Verhulst B. A Power Calculator for the Classical Twin Design. Behavior Genetics.

2017;47(2):255–61. https://doi.org/10.1007/s10519-016-9828-9

60. Westra HJ, Jansen RC, Fehrmann RSN, te Meerman GJ, van Heel D, Wijmenga C, et al.

MixupMapper: Correcting sample mix-ups in genome-wide datasets increases power to

detect small genetic effects. Bioinformatics. 2011;27(15):2104–11.

https://doi.org/10.1093/bioinformatics/btr323

61. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to

the Tidyverse. Journal of Open Source Software. 2019;4(43):1686.

https://doi.org/10.21105/joss.01686

62. Wong YT, Tayeb MA, Stone TC, Lovat LB, Teschendorff AE, Iwasiow R, et al. A

comparison of epithelial cell content of oral samples estimated using cytology and DNA

methylation. Epigenetics. 2022;17(3):327–34.

https://doi.org/10.1080/15592294.2021.1950977

63. Wu HC, Wang Q, Chung WK, Andrulis IL, Daly MB, John EM, et al. Correlation of

DNA methylation levels in blood and saliva DNA in young girls of the LEGACY Girls

study. Epigenetics. 2014;9(7):929–33. https://doi.org/10.4161/epi.28902

64. Yang J, Zeng J, Goddard ME, Wray NR, Visscher PM. Concepts, estimation and

interpretation of SNP-based heritability. Nature Genetics. 2017;49(9):1304–10.

https://doi.org/10.1038/ng.3941

Additional files

Additional file 1. Previously published estimates of SNP-, pedigree-, and twin-based heritability

of the analyzed epigenetic aging measures.

Additional file 2. Performance characteristics of epigenetic aging measures. The figures and

tables represent CpG sites overlap between the epigenetic aging measures, Median Absolute

Errors and mean deviations for three epigenetic age estimators.

Additional file 3. Supplementary Tables and Figures.

Additional file 4. Sensitivity analysis results. Tables S12-23 report goodness-of-fit statistics and

standardized variance components for univariate age-moderated twin models of epigenetic aging

measures unadjusted for biological sex, smoking probe, or cell type composition.

Between-individual differences

Within-individual differences

A

B

Proportion of the variance (A, B axes)

Epigenetic aging measure

A-axis categories: Horvath Accel., PedBE Accel., GrimAge Accel., DunedinPACE

B-axis categories: Δ Horvath Accel., Δ PedBE Accel., Δ GrimAge Accel., Δ DunedinPACE

A values:
- Horvath Accel.: 24%, 26%, 50%
- PedBE Accel.: 47%, 53%
- GrimAge Accel.: 32%, 68%
- DunedinPACE: 47%, 53%

B values:
- Δ Horvath Accel.: 10%, 90%
- Δ PedBE Accel.: 14%, 86%
- Δ GrimAge Accel.: 7%, 93%
- Δ DunedinPACE: 22%, 78%

Relative contribution ■ Additive genetic ■ Shared environmental ■ Unique environmental

| Unstandardized variance components | Standardized variance components |

Horvath Acceleration

PedBE Acceleration

GrimAge Acceleration

DunedinPACE

Relative contribution to variance

— Additive genetic   — Shared environmental   — Variance

— Non-additive genetic   — Unique environmental

Unstandardized variance components

Δ Horvath Acceleration

Δ PedBE Acceleration

Δ GrimAge Acceleration

Δ DunedinPACE

Standardized variance components

Relative contribution to variance

— Additive genetic — Shared environmental — Unique environmental — Variance