UNRAVELING CRP/cAMP-MEDIATED METABOLIC REGULATION IN *ESCHERICHIA COLI* PERSISTER CELLS

Han G. Ngo[1], Sayed Golam Mohiuddin[1], Aina Ananda[2], Mehmet A. Orman[1,*]

[1]Department of Chemical and Biomolecular Engineering, University of Houston, TX, 77204

[2]Department of Biology, Monmouth University, NJ, 07764

[*]Corresponding author: morman@central.uh.edu

ABSTRACT

A substantial gap persists in our comprehension of how bacterial metabolism undergoes rewiring during the transition to a persistent state. Also, it remains unclear which metabolic mechanisms become indispensable for persister cell survival. To address these questions, we directed our efforts towards persister cells in *Escherichia coli* that emerge during the late stationary phase. These cells have been recognized for their exceptional resilience and are commonly believed to be in a dormant state. Our results indicate that the global metabolic regulator Crp/cAMP redirects the metabolism of these antibiotic-tolerant cells from anabolism to oxidative phosphorylation. Although our data demonstrates that persisters exhibit a reduced metabolic rate compared to rapidly growing exponential-phase cells, their survival still relies on energy metabolism. Extensive genomic-level analyses of metabolomics, proteomics, and single-gene deletions consistently highlight the critical role of energy metabolism, specifically the tricarboxylic acid (TCA) cycle, electron transport chain (ETC), and ATP synthase, in sustaining persister levels within cell populations. Altogether, this study provides much-needed clarification regarding the role of energy metabolism in antibiotic tolerance and highlights the importance of using a multipronged approach at the genomic level to obtain a broader picture of the metabolic state of persister cells.

INTRODUCTION

Bacterial persisters within cell cultures constitute a small subpopulation of cells exhibiting a transient antibiotic-tolerant state (Balaban et al., 2004). While persisters have traditionally been characterized as non-growing and dormant phenotypes (Lewis, 2010), recent studies challenge these conventional hallmarks, revealing the heterogeneity of persister cells in terms of growth, metabolism, and other cellular activities (Adams et al., 2011; Arnoldini et al., 2014; Hossain et al., 2023; Ma et al., 2010; Mok, Park, et al., 2015; Orman & Brynildsen, 2013a, 2015; Shan et al., 2015; Wakamoto et al., 2013). Despite potential discrepancies in research outcomes in the field, we think that these variations arise from the intricate and diverse survival mechanisms employed by bacterial cells in response to adverse conditions, such as antibiotic treatments. Furthermore, the interplay of stochastic and deterministic factors associated with these mechanisms adds another layer of complexity, with outcomes highly contingent on factors such as cell types, antibiotics, and experimental and growth conditions (Michiels et al., 2016). The persistence phenomenon presents a significant health concern (Huemer et al., 2020; Murray et al., 2022), as the transient antibiotic-tolerant state of persister cells promotes recurrent infections (Fisher et al., 2017) and establishes them as a reservoir for the emergence

of antibiotic-resistant mutants (Barrett et al., 2019; Levin-Reisman et al., 2017; Windels et al.,
2019).

Drug tolerance is a widespread phenomenon observed in both prokaryotic and eukaryotic cell
types. Otto Warburg's research in the early twentieth century unveiled an intriguing aspect of
mammalian cell metabolism known as "aerobic glycolysis," wherein proliferating cells (e.g.,
tumor cells) derive energy predominantly through glycolysis, even in the presence of oxygen
(Warburg, 1925). This metabolic reprogramming involves restricting entry into the TCA cycle
through precise enzymatic control, diverting glycolytic intermediates towards anabolic
pathways. This adaptation supports the extensive biosynthesis required for active cell
proliferation in tumors (Hanahan & Weinberg, 2011). Remarkably, tumorigenic persisters that
exist in a non-proliferating state may not primarily depend on aerobic glycolysis. Instead, there
is substantial evidence suggesting that these cells rely on energy metabolism (Porporato et al.,
2018; Vasan et al., 2020); however, the presence of this metabolic state in antibiotic-tolerant
bacteria is still a question mark (Orman & Brynildsen, 2015). Reprogramming energy
metabolism seems to be an evolutionarily conserved strategy for cells facing stress or adverse
conditions, as these cells might benefit from the significantly higher ATP production efficiency
provided by oxidative phosphorylation (Hanahan & Weinberg, 2011). The identification of a
mechanism shared by diverse cell types could open avenues for the development of global
strategies to target drug-tolerant cells effectively.

A recent study suggests that bacterial persisters constitute a stochastically formed
subpopulation of low-energy cells, despite some observed overlap in ATP levels between
antibiotic-sensitive and persister cells (Manuse et al., 2021). The persister cells examined in that
study were derived from an aged stationary phase culture (48 hours post-inoculation) (Manuse
et al., 2021), a condition known to elevate the number of non-growing cells, which do not
promptly resume growth upon transfer to a fresh medium (Mohiuddin, Kavousi, et al., 2020).
While the non-growing cells formed during the stationary phase have reduced metabolic
activity compared to growing cells, as demonstrated in our earlier study (Orman & Brynildsen,
2013a), they still exhibit a certain degree of respiratory activity (Orman & Brynildsen, 2015). In
a recent study measuring ATP levels in viable but non-culturable (VBNC; a phenotype that is
antibiotic tolerant but unable to resume growth after antibiotic removal), persister and
antibiotic-sensitive cells from 24-hour stationary phase cultures, a significant overlap in
intracellular ATP concentrations was observed between antibiotic-sensitive and persisters cells
(Li et al., 2024). On the other hand, VBNC cells exhibited drastically lower ATP levels compared
to both persister and antibiotic-sensitive cells (Li et al., 2024). Another independent study,
which utilized single-cell analysis, reported that ofloxacin persisters were metabolically active
cells in exponentially growing cultures before treatment and these cultures were obtained from
16-hour overnight precultures (not aged) (Goormaghtigh & Van Melderen, 2019). These
findings suggest that while persisters formed during exponential growth may initially retain
metabolic activity, they may gradually transition to reduced energy states as they age within
the culture. The nature of persister-cell metabolism in bacteria is a topic that has long been a
point of contention in scientific circles. The controversy surrounding this topic primarily arises
from studies that rely on bacteriostatic chemicals or a limited number of gene deletions, or
direct comparisons to exponentially growing cells, all of which have inherent drawbacks

(Conlon et al., 2016; Manuse et al., 2021; Orman & Brynildsen, 2013a, 2015). The metabolism
of persister cells, a very complex phenomenon, cannot be easily characterized by a simplistic
term such as "metabolic dormancy". Even if persister cells may exhibit a lower metabolism
compared to the vast majority of rapidly growing exponential-phase cells, they may still rely on
energy metabolism for their functioning (Amato et al., 2014; Prax & Bertram, 2014). In fact,
analyzing published studies collectively suggests that bacterial cell metabolism undergoes
intricate alterations or rewiring as they transition into a tolerant state, and these alterations
seem to be highly dependent on the specific conditions tested (Adams et al., 2011; Arnoldini et
al., 2014; Ma et al., 2010; Mok, Park, et al., 2015; Orman & Brynildsen, 2013a, 2015; Shan et al.,
2015; Wakamoto et al., 2013).

To gain a better understanding of the critical role of energy metabolism in persister cell
survival, we have focused our research on antibiotic-tolerant cells formed during the stationary
phase, given that these cells are known to be highly resilient and capable of surviving a variety
of stressors, including antibiotics, and are assumed to be dormant (Balaban et al., 2004; Orman
& Brynildsen, 2015). These cells are also referred to as type I persisters, which cannot readily
resume growth when diluted in fresh media during the lag phase (Balaban et al., 2004).
Previous studies showed that these persister cells can metabolize specific carbon sources that
make them susceptible to aminoglycosides (AG) (Allison et al., 2011; Mok, Orman, et al., 2015;
Orman & Brynildsen, 2013b). Their AG susceptibility is due to increased AG uptake, which is
facilitated by increased ETC activity and membrane potential (Allison et al., 2011). The presence
of active energy metabolism in antibiotic-tolerant, non-growing cells may indeed explain their
rapid killing by AG in the presence of carbon sources (Orman & Brynildsen, 2013b, 2015). When
the knockout strains of global transcriptional regulators (i.e., ArcA, Cra, Crp, DksA, Fnr, Lrp, and
RpoS) were screened using the AG potentiation assay in our previous study, the results showed
that the panel of carbon sources tested potentiated the AG killing of tolerant cells derived from
most knockout strains, except for Δcrp and ΔcyaA (Mok, Orman, et al., 2015). This can be
attributed to the lack of active energy metabolism in these mutant strains, as the Crp/cAMP
potentially shapes persister cell metabolism during the stationary phase. Depletion of primary
carbon sources activates adenylate cyclase (CyaA) (Pastan & Perlman, 1970), increasing cyclic-
AMP (cAMP) levels in cells (Bettenbrock et al., 2007; Park et al., 2006). The cAMP molecules,
along with its receptor protein (Crp), activate genes related to the catabolism of secondary
carbon sources, potentially supporting cellular functions and energy levels (Deutscher, 2008; Fic
et al., 2009; Görke & Stülke, 2008; Kolb et al., 1993). Here, using metabolomics, proteomics,
and high-throughput screening of single-gene deletion strains, we have provided evidence that
the Crp/cAMP regulatory complex maintains an active state of energy metabolism while
downregulating anabolic pathways in the antibiotic-tolerant persister cells.

RESULTS

Disruption of the Crp/cAMP complex affects the formation of persister cells at the late
stationary phase.

Since Crp/cAMP-mediated metabolic changes can be induced by nutrient depletion during the
stationary phase, we wanted to assess the effects of deleting the *crp* and *cyaA* genes (Δ*crp* and

ΔcyaA) on both persister cell formation and metabolism during this phase. As anticipated, the
deletion of the cyaA gene resulted in a notable reduction in intracellular cAMP concentration
(**Figure 1—figure supplement 1**). However, the Δcrp strain exhibited an increase in cAMP
concentration, potentially due to the negative feedback regulatory mechanism of the
Crp/cAMP complex for the cyaA gene promoter (Keseler, 2004; Majerfeld et al., 1981) (**Figure
1—figure supplement 2**). When comparing the growth curves of E. coli wild-type (WT), Δcrp,
and ΔcyaA main cultures under identical conditions studied here (see **Materials and Methods**),
all three strains started to enter the stationary phase around 5 hours, with the mutant strains
exhibiting slightly lower optical density levels than the WT at this time point (**Figure 1—figure
supplement 3**). Also, our data provide evidence of an increase in cAMP levels in WT cells during
their transition into the stationary phase (**Figure 1—figure supplement 4**), aligning with existing
literature (Bettenbrock et al., 2007; Keseler, 2004; Park et al., 2006). For type I persister
quantification, we diluted cells in the fresh medium, consistent with previous studies (Balaban
et al., 2004; Orman & Brynildsen, 2015), at early (t=5 h) and late (t=24 h) stationary phases and
subsequently exposed them to an extended period (20 h) of ampicillin or ofloxacin treatment
(200 µg/mL ampicillin and 5 µg/mL ofloxacin). These treatments were carried out at
concentrations surpassing the minimum inhibitory concentrations (MIC), necessary for the
selection of antibiotic-tolerant persister cells (**Supplementary File 1**) (Keren, Kaldalu, et al.,
2004). Also, type I persisters, formed during the stationary phase, exhibit a slow transition from
a non-growing state to an active state when transferred to a fresh medium in the lag phase
(Balaban et al., 2004; Jõers & Tenson, 2016; Orman & Brynildsen, 2015; Vulin et al., 2018);
therefore, this transition requires longer antibiotic treatment durations. Moreover, we
transferred an equal number of cells from each strain to the fresh medium to ensure
consistency in cell numbers. To assess persistence, we collected samples during antibiotic
treatment, washed them to remove antibiotics, and plated them on agar media to quantify
colony-forming units (CFU) of surviving cells (see **Materials and Methods**). The resulting
biphasic kill curves—plots of CFU levels over treatment time—are characteristic of persistence
phenotypes (**Figure 1**). Notably, the WT strain showed a marked increase in both ampicillin- and
ofloxacin-persister cells during the late stationary phase, in contrast to the mutant strains,
where no such increase was observed (**Figure 1A, B**). However, this trend was not observed in
the early stationary phase (**Figure 1A, B**). To confirm that the observed decrease in persister
levels in the mutant strains in the late stationary phase is solely attributed to the perturbation
of the Crp/cAMP regulatory network, we reintroduced crp expression to the Δcrp strain using a
low-copy plasmid carrying the crp gene and its promoter. As a control, we utilized an empty
vector of the same plasmid. The results demonstrated that the expression of crp restored the
persister level in the mutant strain, while the plasmid itself had no impact on persister levels
(**Figure 1—figure supplement 5A, B**). Altogether, these findings highlight the significant role of
Crp/cAMP in ampicillin and ofloxacin persister formation in the late stationary phase.

Ampicillin and ofloxacin, both broad-spectrum antibiotics with a strong dependence on cell
metabolism (E. J. Zheng et al., 2020), target cell wall synthesis and DNA gyrase activity,
respectively. In addition to these two antibiotics, we also quantified aminoglycoside-persister
levels in both WT and mutant strains. This was achieved by exposing diluted cells from both
early and late stationary phases to 50 µg/mL gentamicin, a concentration exceeding the MIC

levels (**Supplementary File 1**). After prolonged gentamicin exposure (20 h), the tolerant cell
colonies were found to be below the limit of detection for all strains and conditions (**Figure 1C,**
and Figure 1—figure supplement 5C). Although bacterial tolerance can vary significantly
depending on the specific antibiotics and growth phase used (Hofsteenge et al., 2013), this
outcome contrasts starkly with the observed levels of ampicillin and ofloxacin persisters in WT
(**Figure 1**). The mechanism by which aminoglycosides eliminate persister cells in the WT strain
may be linked to their metabolism, given that aminoglycoside uptake is an energy-requiring
process (Allison et al., 2011; Taber et al., 1987), and this aspect will be explored further in the
subsequent section.

We would like to highlight that antibiotic concentrations, washing procedures (to remove
antibiotics), and agar plate incubation times for CFU enumeration may affect experimental
outcomes. However, variations in these parameters, including lower antibiotic concentrations
normalized to the MIC of each strain (5× or 10× MIC; see **Figure 1—figure supplement 6**),
additional washing steps to ensure complete removal of antibiotics (see **Figure 1—figure**
supplement 6), and extended agar plate incubation times of up to 48 hours (see **Figure 1—**
figure supplement 7), did not affect the persistence phenotype. The Δ*crp* and Δ*cyaA* strains
consistently showed reduced ampicillin and ofloxacin persistence compared to WT, whereas all
three strains showed no detectable persisters following gentamicin treatment (**Figure 1—figure**
supplement 6, 7). Moreover, we examined the impact of Crp/cAMP disruption in the HipA7
strain to assess whether the role of Crp/cAMP in persistence extends to a genetically sensitized,
persister-enriched background. Deletion of either *crp* or *cyaA* in the HipA7 background
significantly reduced persistence to both ampicillin and ofloxacin during the late stationary
phase (**Figure 1—figure supplement 8**), mirroring the effects observed in the WT strain.
Altogether, these results show that disruption of the Crp/cAMP regulatory network impairs
persister formation during the late stationary phase, underscoring its critical role in ampicillin
and ofloxacin tolerance.

Crp/cAMP complex governs *E. coli* stationary phase metabolism.

To determine whether the reduced ampicillin and ofloxacin persister levels at the late
stationary phase in the mutant strains (**Figure 1A, B**) are linked to stationary phase metabolism,
we utilized untargeted mass spectrometry (MS). This approach facilitated the quantification of
metabolites in Δ*crp* cells, allowing for a comparison with WT controls in both the early and late
stationary phases (**Figure 2A**). Since both Δ*crp* and Δ*cyaA* strains exhibit the same persistence
phenotype, and *crp* deletion effectively abolishes Crp/cAMP complex function, we used the
Δ*crp* mutant for metabolomic profiling to capture key regulatory changes while maintaining
experimental feasibility. The metabolomics data were subjected to unsupervised hierarchical
clustering, and metabolites identified in independent biological replicates of each strain and
condition were found to cluster together (**Figure 2A**), thus confirming the reproducibility of our
data.

To elucidate the upregulated and downregulated metabolic pathways in the mutant strain as
compared to the WT strain, we performed enrichment analyses utilizing MetaboAnalyst (Lu et
al., 2023). For downregulated pathways, we considered a threshold ratio of 0.5 or lower, where
the ratio indicates metabolite levels in the mutant strain relative to the WT (**Supplementary**

File 2A). Conversely, for upregulated pathways, the threshold ratio was set at 2 or higher (**Supplementary File 2B**). The enrichment ratio for each pathway was calculated based on the number of metabolite hits compared to the expected hits derived from the chemical structure library (Lu et al., 2023). Our extensive comparison of the mutant cells to the WT cells through pathway enrichment analysis (refer to **Figure 2B** and **Figure 2—figure supplement 1-3** for pairwise comparison of different conditions) revealed several important findings:

(i) During the early stationary phase, we observe a slight downregulation in the abundance of TCA cycle metabolites, including citrate and fumarate, in the Δ*crp* strain compared to WT (**Figure 2C** and **Figure 2—figure supplement 1**). However, as the late stationary phase progresses, the downregulation in both TCA cycle and pentose phosphate metabolism becomes more pronounced in the Δ*crp* strain (**Figure 2B, D**).

(ii) The Δ*crp* strain exhibits upregulation of several metabolites compared to WT during both early and late stationary phases, primarily associated with anabolic pathways. Particularly, the upregulation of some of these pathways becomes more pronounced during the late stationary phase. These pathways include crucial metabolites like deoxyribonucleosides and ribonucleosides (which play essential roles in DNA and RNA synthesis), fatty acids and carboxylic acids (the main components of bacterial cell membranes), and peptides (which are linked to protein synthesis) (**Figure 2B**).

(iii) During the early stationary phase, we noticed a significant upregulation in the abundance of intermediate metabolites related to glycolysis, gluconeogenesis, and pyruvate metabolism in mutant cells compared to WT cells (**Figure 2C**). This observation is not surprising, as the inhibition of the TCA cycle in the mutant strain could potentially redirect metabolic fluxes toward glycolysis and lactate metabolism.

Altogether, our metabolic data indicate that, in the stationary phase, WT cells maintain their energy metabolism to some extent while downregulating their anabolic pathways (**Figure 2A**). This metabolic state appears to be regulated by the Crp/cAMP complex, as perturbing its function leads to a significant downregulation of energy metabolism and an upregulation in the abundance of anabolic metabolites (**Figure 2B**).

Proteomics analysis revealed upregulated pathways in the Δ*crp* strain associated with anabolic metabolism, alongside the downregulation of key proteins in energy metabolism.

Since the Crp/cAMP complex acts as a transcriptional regulator affecting the expression of metabolic proteins whose abundance directly affects cellular metabolites, we performed untargeted proteomics, our second genomic-level study, to further validate our results. Considering the noticeable metabolic alterations observed during the late stationary phase, we utilized MS to quantify proteins in Δ*crp* cells and compared them to WT controls at this stage. The resulting proteomics data were subjected to unsupervised hierarchical clustering, and the proteins identified in independent biological replicates of each strain were found to cluster together, confirming the consistency of our findings (**Figure 3—figure supplement 1**). By analyzing protein-protein association networks and employing functional enrichment through STRING (Szklarczyk et al., 2021, 2023), which integrates various functional pathway classification frameworks such as Gene Ontology annotations, KEGG pathways, and UniProt keywords, we pinpointed various upregulated and downregulated pathways in the mutant

strain when compared to the WT (**Supplementary File 3**). The upregulated pathways are associated with anabolic metabolism and encompass peptidoglycan metabolic processes, cell wall organization or biogenesis, cellular component organization or biogenesis, regulation of cell shape, cell cycle, and cell division (**Figure 3A**). Also, the mutant strain displayed downregulated pathways, encompassing glycerol metabolism, TCA cycle, pyruvate metabolism, glycolysis, and various pathways associated with ribosome and transcriptional factor activity (**Figure 3B**). Our analysis specifically pinpointed a cluster of proteins involved in energy metabolism and respiratory processes. Notably, this cluster includes GltA, SdhB, SucC, SucD, FrdB, FrdA, AcnA, AceA, and Mdh proteins, which play crucial roles in either the TCA cycle or as membrane-bound components of ETC (**Figure 3B**). Altogether, the alignment between metabolomics and proteomics analyses provides additional validation for the Crp/cAMP-mediated metabolic state.

Crp/cAMP complex shapes *E. coli* cell proliferation dynamics.

The omics data suggest that the upregulation in the abundance of anabolic metabolites and proteins, particularly those related to cell wall organization or biogenesis, cell cycle, and cell division, in the stationary-phase mutant cells, would likely enhance their ability to resume growth upon transitioning to fresh medium. To investigate this, we utilized a cell proliferation assay that employed an inducible fluorescent protein (mCherry) expression cassette. This assay provided us with the ability to monitor non-growing cells at a single-cell resolution, as described previously (Orman & Brynildsen, 2013a; Roostalu et al., 2008). The mCherry expression cassette is controlled by an isopropyl ß-D-1-thiogalactopyranoside (IPTG) inducible synthetic T5 promoter that was previously inserted into the chromosome of an *E. coli* strain carrying a *lacI^q* promoter mutation (Orman & Brynildsen, 2013a). This configuration allowed for precise regulation of mCherry expression using IPTG. Here, we introduced *crp* and *cyaA* deletions into this strain. These deletions reduced the persistence of the mCherry-expressing *E. coli* strain in the late stationary phase (**Figure 4—figure supplement 1**), consistent with the findings presented in **Figure 1**. To perform the growth assay, we induced mCherry expression in the main cultures and then washed the cells to remove IPTG. The cells were then inoculated into a fresh medium without IPTG and their growth was analyzed with a flow cytometer. This allowed us to track the dilution of mCherry protein within the cells, which served as an indicator of cell proliferation. As shown in **Figure 4A**, initially, all cells exhibited high red fluorescence. However, as cells underwent division, the red fluorescence of the overall population decreased in the absence of the inducer (**Figure 4A**). Notably, within the WT strain, a subpopulation from the late stationary phase cultures displayed constant fluorescence levels, indicating their inability to divide (**Figure 4B**). In contrast to the WT strain, we did not detect similar subpopulations in the mutant strains (**Figure 4B**). Additionally, this subpopulation of non-growing cells does not emerge during the early stationary phase cultures (**Figure 4A**). This observation provides an explanation for the observed reduction in persister levels in these mutant strains in the late stationary phase, as the enrichment of persister cells within these non-growing cell subpopulations was reported in previous studies (Joers et al., 2010; Orman & Brynildsen, 2013b; Roostalu et al., 2008).

To confirm the significance of Crp/cAMP in the formation of non-growing cells, we introduced the expression plasmid carrying the *crp* gene into the Δ*crp* strain. As anticipated, the

introduction of the *crp* expression plasmid resulted in the emergence of a non-growing population within the culture, contrasting with the mutant strain containing the empty plasmid vector used as a control (**Figure 4—figure supplement 2A, B**). The reduced capacity of stationary-phase WT cells to initiate proliferation upon transfer to a fresh medium suggests the possible presence of an extended lag phase in these cells. To investigate this, we employed flow cytometry to precisely quantify cell numbers and generate growth curves for both WT and mutant strains. As anticipated, the growth curve of the WT strain displayed a slower initial growth rate and a prolonged lag phase duration compared to the mutant strains (**Figure 4C**). Conversely, the mutant strains displayed a shorter lag phase, yet they demonstrated an increased doubling time in the exponential phase compared to WT (**Figure 4C**), which was also anticipated, considering their decreased reliance on oxidative phosphorylation due to TCA cycle inhibition. Altogether, these results provide additional support and validation for the findings from our metabolomics and proteomics data, as our data reveals a correlation between the abundance of molecules associated with cell division and the ability of the stationary phase cells to resume growth.

Persister cells rely on energy metabolism.

The Crp/cAMP-mediated metabolic state, characterized by increased respiration in WT compared to mutant strains, was further validated using redox sensor green (RSG) dye and a reporter plasmid measuring the promoter activity of succinate:quinone oxidoreductase (SQR) genes. The SQR reporter system (Zaslaver et al., 2006) employs green fluorescent protein (GFP) expression, regulated by the promoter of the SQR operon, which includes the *sdhA, sdhB, sdhC,* and *sdhD* subunit genes. The SQR complex plays a vital role in cellular metabolism by catalyzing the oxidation of succinate to fumarate concurrently with the reduction of ubiquinone to ubiquinol, thus directly linking the TCA cycle with the respiratory ETC (Keseler, 2004). Our results indicate an upregulation of the SQR promoter activity in WT cells compared to the mutant strains in the stationary phase, validating the findings from our metabolomics and proteomics data (**Figure 5A**). The RSG dye, on the other hand, serves as a well-established metabolic indicator, measuring bacterial oxidation-reduction activity, a crucial function involving the ETC driven by the TCA cycle. Once reduced by bacterial reductases, the RSG dye emits a stable green fluorescent signal (**Figure 5—figure supplement 1**). Our data demonstrate that the redox activities of WT cells are much higher and more heterogeneous compared to those of the mutant strains in the late stationary phase, further corroborating the results from our preceding analyses (**Figure 5B**). Furthermore, we utilized a methodology that integrates the mCherry expression system, flow cytometry, and ampicillin-mediated cell lysis, to determine whether persister cells in WT still maintain their respiration. In this assay, both the WT and mutant strains carrying the mCherry expression system were exposed to ampicillin after transferring them to a fresh medium. The inducer was added to both growth and treatment cultures to sustain the cells' red signals. Unlike other antibiotics, ampicillin disrupts cell wall synthesis, leading to the lysis of cells upon their resumption of growth. As seen in **Figure 5— figure supplement 2**, the cells that were lysed lost their mCherry signals. On the other hand, the resilient, tolerant cells that evaded ampicillin-induced lysis maintained their mCherry levels throughout the treatment. In the mutant strains, ampicillin was effective in lysing almost all cells as anticipated, leaving no or small number of intact cells (**Figure 5—figure supplement 2**).

However, in the WT strain, we detected a subpopulation of intact cells throughout the entire treatment period (**Figure 5C,** and **Figure 5—figure supplement 2**). While the population-level redox activities of these tolerant intact cells in WT are lower than those of exponential phase cells, they still displayed a significant increase in RSG levels compared to cell populations before antibiotic treatments or untreated control cells subjected to identical conditions (**Figure 5D),** suggesting that they maintain steady-state energy metabolism. We want to highlight that not all intact cells in the WT strain reported here are persisters. A significant portion comprises 'viable but non-culturable' (VBNC) cells, and WT cells exhibit markedly higher VBNC levels than Δ*crp* and Δ*cyaA* strains (**Figure 5—figure supplement 3**). VBNC cells can be quantified from intact cells following beta-lactam treatments (Orman & Brynildsen, 2013b; Roostalu et al., 2008). These cells may exhibit metabolic activities but are unable to readily colonize upon transfer to fresh medium (Ayrapetyan et al., 2018; Orman & Brynildsen, 2013b). Collectively, our metabolic measurement data (**Figure 5A-C**) aligns with the findings from our omics analyses, and the number of intact cells observed in WT after beta-lactam treatment is consistent with the count of non-growing cells in WT (**Figure 4B** vs **Figure 5—figure supplement 2**). These non-growing cells are anticipated to be less susceptible to lysis by beta-lactams (Orman & Brynildsen, 2013b; Roostalu et al., 2008).

The genomic-level screening of *E. coli* knockout underscores the significance of energy metabolism in sustaining the viability of persister cells.

Although some metabolic genes, including those encoding the TCA cycle (e.g., *sdhA, sucB, mdh, icd*), have been studied in *E. coli* (Luidalepp et al., 2011; Ma et al., 2010; Manuse et al., 2021; Orman & Brynildsen, 2015; Yu et al., 2019), a comprehensive genomic-level screening strategy is necessary to validate which metabolic pathways are truly associated with antibiotic tolerance. To further underscore the importance of energy metabolism, we conducted a high-throughput screening of 149 different *E. coli* K-12 BW25113 mutant strains from the Keio knockout library (Baba et al., 2006). The selected strains are related to central carbon metabolism, encompassing glycolysis, pentose phosphate pathways, TCA cycle, ETC, ATP synthase, and fermentation pathways (**Figure 5E**). While the deletion of genes related to glycolysis and pentose phosphate pathways did not affect antibiotic tolerance in the cells, some mutant strains associated with cytochrome bo and quinone oxidoreductase complexes (e.g., *cyo* genes, and *nuoL*) exhibited enhanced tolerance (**Supplementary File 4**). However, the mutant strains exhibited reduced tolerance to both antibiotics compared to the control *E. coli* K-12 BW25113 WT strain were found to be largely associated with the TCA pathway (*sucA, sucB, lpd, sucC, sucD, sdhA, sdhB, sdhC, sdhD, gltA, acnB, aceE, fumA, mdh* and *fumC*), ETC (*nuoB, nuoC, nuoI, nuoK, nuoM,* and *narV*), ATP synthesis (*atpA, atpB, atpC, atpD, atpE,* and *atpH*), and mixed acid fermentation pathways (*ldhA, fdhF, pta, adhE,* and *frdC*) (**Figure 5E,** and **Supplementary File 4**). We acknowledge that the Keio strains were generated in a high-throughput manner, and there might be unknown errors in their genomic DNA. To ensure the reproducibility of our findings, we generated knockout strains for key genes associated with the TCA, ETC, ATP synthase, glycolysis, and pentose phosphate pathway (**Figure 5F**, see **Supplementary File 5** for detailed description of genes). We then tested their antibiotic tolerance, and our results were consistent with the omics data and screening outcomes (**Figure 5F**), confirming the critical role of energy metabolism, specifically the TCA cycle, ETC, and ATP

synthase, in bacterial persistence. We note that equal numbers of cells from each strain were
transferred into antibiotic treatment media to ensure consistency in cell numbers, and no
significant growth deficiencies or differences in cell density were observed in the late stationary
phase of the knockout strains (**Figure 5—figure supplement 4**).

DISCUSSION

Our study highlights the crucial role of the Crp/cAMP complex in maintaining the metabolic
state of stationary-phase persister cells, enabling their survival under adverse conditions.
Through metabolomics, proteomics, and high-throughput screening of single-gene deletion
strains, we substantiated that the Crp/cAMP regulatory complex sustains an active respiratory
state while downregulating anabolic pathways in persister cells. This respiratory state is vital for
the survival of persister cells, as perturbing the Crp/cAMP complex or respiration significantly
reduced persister phenotypes, which may explain the previously reported decrease in antibiotic
tolerance in *E. coli* cells cultured under anaerobic conditions (Orman & Brynildsen, 2015).
Notably, we observed an upregulation of anabolic metabolites and proteins when the
Crp/cAMP regulatory complex was perturbed, particularly those associated with cell wall
organization, cell cycle, and cell division, enhancing the ability of stationary-phase mutant cells
to resume growth. Although the literature has shown associations between antibiotic tolerance
with proteins involved in cell division and the TCA cycle (e.g., SdhA, SucB, Mdh) (Luidalepp et
al., 2011; Ma et al., 2010; Orman & Brynildsen, 2015; Yu et al., 2019), our study establishes a
strong link between these critical cellular processes and the Crp/cAMP complex, providing
much-needed clarity in the field. In fact, upon investigating Crp/cAMP regulons via the Ecocyc
database (Keseler, 2004), we identified that certain metabolic genes, deleted in our *E. coli* K-12
MG1655 background, are potentially regulated by Crp/cAMP (**Supplementary File 5**), providing
additional support for the validity of our omics results.

We acknowledge that our metabolomic and proteomic data were obtained at the whole-
population level, rather than from isolated persister cells. Fluorescent reporters combined with
fluorescence-activated cell sorting (FACS) have been utilized to study persister cells, including in
our previous studies (Amato et al., 2013; Orman & Brynildsen, 2013a, 2015). However, this
approach only enriches for persisters rather than isolating a pure population, as persisters still
constitute a small fraction of the sorted cells. Despite these limitations, our population-level
analyses provide valuable insights into the role of the Crp/cAMP complex in regulating non-
growing cell formation and persistence (**Figure 4**). We further validated these findings through
single-gene deletions and flow cytometry-based assays (**Figure 5**), confirming a consistent
reduction in both non-growing and persister cells in Crp/cAMP-deficient strains.

The deletion of *cyaA* resulted in a reduction of cAMP levels, as expected given the role of the
CyaA enzyme in cAMP synthesis (**Figure 1–figure supplement 1**). Conversely, the removal of *crp*
led to an increase in cAMP levels compared to those in wild-type cells. Notably, this increase is
statistically significant (**Figure 1–figure supplement 1**). Although this is an interesting
observation, it is likely due to the feedback regulation of the Crp/cAMP complex on *cyaA*
expression (Aiba, 1985; Keseler et al., 2011; Majerfeld et al., 1981). Specifically, disruption of
Crp function is expected to derepress the *cyaA* promoter (P$_{cyaA}$), leading to increased CyaA
expression and elevated cAMP levels. To test this, we employed a P$_{cyaA}$-*gfp* reporter plasmid

(pMSs201) in *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* strains to monitor promoter activity. As
anticipated, deletion of *crp* led to enhanced *gfp* expression, confirming that Crp/cAMP
negatively regulates *cyaA* expression via feedback. These results clarify the observed cAMP
differences and support the regulatory role of the Crp/cAMP complex (**Figure 1—figure**
supplement 2).

Our findings reveal substantial heterogeneity in metabolism (measured by RSG) among WT
stationary phase cells, contrasting with the more uniform behavior observed in Δ*crp* and Δ*cyaA*
strains (**Figure 5B**). We also demonstrated the presence of two distinct populations in WT cells
during the late stationary phase: one that resumes rapid growth and another subpopulation
that does not resume growth when transferred to a fresh medium (**Figure 4B**). The absence of
this non-growing cell subpopulation in the mutant strains could account for their sensitivity to
aminoglycosides. However, the mechanism by which aminoglycosides kill these non-growing
cells in the WT strain remains perplexing. The underlying reasons might be linked to their
metabolism, as aminoglycoside uptake is an energy-requiring process, relying on the electron
flow through membrane-bound respiratory chains (Allison et al., 2011). Moreover, persister
cells obtained from various antibiotics, such as ampicillin and ofloxacin, in WT *E. coli* were
previously found to exhibit sensitivity to aminoglycosides when sugar molecules were
introduced into the cultures (Allison et al., 2011; Orman & Brynildsen, 2013b). However, the
enhanced sensitivity mediated by sugar molecules was reversed to its original state in a
subsequent study when the Crp/cAMP complex was genetically perturbed (Mok, Orman, et al.,
2015). This can be attributed to the lack of active energy metabolism in these genetically
altered strains, as suggested by our comprehensive genomic-level analyses here. While the
absence of cell division in non-growing cell subpopulations in WT may suggest a
downregulation in anabolic metabolism, their energy metabolism may remain partially active,
which could potentially explain the phenomenon of aminoglycoside potentiation. Indeed, our
results presented in **Figure 5D** support this interpretation.

Antibiotics are generally effective against proliferating bacteria, leading to the notions that
tolerance is linked to temporary growth suppression and that persister cells are dormant
phenotypes with repressed metabolism. Although persister cells are generally considered to
have reduced metabolic activity compared to exponentially growing cells, their survival may still
depend on energy metabolism. Also, the direct comparison of persister cell metabolism to that
of exponentially growing cells may not be the best approach. Growing cells have a very high
energy output and consume metabolites at a fast pace. Therefore, any comparison between
tolerant and non-tolerant cell populations requires proper normalization techniques such as
adjusting cellular metabolic activities to the amount of substrate utilized by cells. An example of
this normalization was conducted by Heinemann's group (Radzikowski et al., 2016),
demonstrating that ATP production rates per substrate in tolerant cells exceed those of
exponentially growing cells.

We diluted cells in fresh medium at early and late stationary phases before antibiotic
treatments. This step is essential for quantifying type I persisters, as these cells do not readily
resume growth upon dilution in the fresh medium during the lag phase (Balaban et al., 2004;
Orman & Brynildsen, 2015). We acknowledge that antibiotic tolerance is influenced by various
factors, including culture dilutions, media, specific strains, antibiotics tested, treatment

durations, and the growth phase during treatment administration. These factors may contribute to variations in reported persister levels observed in the ΔcyaA strain during the exponential phase (Chu et al., 2012; Molina-Quiroz et al., 2018; Parsons et al., 2024; Sulaiman & Lam, 2020; Yamasaki et al., 2020; A. Zeng et al., 2022). To assess how *cyaA* and *crp* deletions affect antibiotic responses under conditions similar to those used by Zeng et al. (J. Zeng et al., 2022) —specifically, exponential-phase *E. coli* BW25113 strains (Keio collection), lower antibiotic concentrations, and short treatments (e.g., 1 hour)—we first tested *E. coli* MG1655 WT, Δ*crp*, and Δ*cyaA* strains in late stationary phase using reduced antibiotic concentrations and shorter exposures. Both knockouts showed decreased survival following ampicillin and ofloxacin treatment compared to WT (**Supplementary File 6**), consistent with our findings in **Figure 1**. In exponential-phase, the knockout strains exhibited reduced survival after ampicillin treatment but increased survival after ofloxacin treatment relative to WT (**Supplementary File 7A**), again mirroring the trends in **Figure 1**. Gentamicin treatment, however, produced variable results in MG1655 knockouts, likely due to the brief 1-hour exposure being insufficient for robust conclusions (**Supplementary File 7A**). Notably, when we tested the corresponding Keio knockout strains in the BW25113 background, we observed increased tolerance in exponential-phase cells, reproducing Zeng et al.'s findings under their specific conditions (**Supplementary File 7B**), although BW25113 and MG1655 exhibited distinct persister phenotypes in exponential phase (**Supplementary File 7A, B**). These results, altogether, highlight the sensitivity of antibiotic tolerance and persistence phenotypes to factors such as strain background, antibiotic concentration, and treatment duration.

While we did not focus on the exponential growth phase in our study, it is noteworthy that the reduced growth rate during this phase in the mutant strains (**Figure 4C**) may explain the antibiotic tolerance observed in previous studies involving the *cyaA* deletion (Chu et al., 2012; Molina-Quiroz et al., 2018; Sulaiman & Lam, 2020; Yamasaki et al., 2020). The growth disparity noted between mutant and WT strains, particularly evident around 5 hours in our results (**Figure 1–figure supplement 3**), may also be linked to the ofloxacin persisters observed in the mutant strains at this specific time point (**Figure 1**). While slow cell growth may indeed correlate with bacterial persistence (Kaldalu & Tenson, 2019), it is important to note that the persistence associated with perturbations of metabolic genes cannot be solely attributed to the slow growth. In fact, the persistence of these mutant strains should depend on many factors (see **Supplementary File 5**) as reported by diverse research groups (Kim et al., 2016; Luidalepp et al., 2011; Ma et al., 2010; Orman & Brynildsen, 2015; Pandey et al., 2021; Shan et al., 2015; Spoering et al., 2006; Wang et al., 2018; Yu et al., 2019; Zalis et al., 2019). For instance, in *E. coli*, TCA inactivation was shown to decrease ampicillin and ofloxacin persistence during the lag phase (Orman & Brynildsen, 2015), yet it enhances gentamicin tolerance in the exponential phase, which remains unexplained by factors such as cell growth, redox activities, proton motive force (PMF), or ATP levels (Shiraliyev & Orman, 2023). Furthermore, gene deletions often trigger pleiotropic effects, leading to unique tolerance mechanisms not evident in wild-type strains. In our Keio screening data analysis, we observed that the deletion of *icd* appeared to enhance persistence (**Supplementary File 4**), in line with a previous study (Manuse et al., 2021). The *icd* gene encodes a TCA cycle enzyme, isocitrate dehydrogenase. While it remains unclear whether this observed outcome is attributable to other unseen pleiotropic effects

stemming from the *icd* deletion, our data consistently indicates that the most significant
reduction in persistence levels occurs with disruptions in energy metabolism. A comprehensive
approach, encompassing omics and knockout screening as presented in this study, offers a
more complete understanding, revealing the consensus behavior within the entire metabolic
network.

Reactive oxygen species (ROS) have been proposed to contribute to antibiotic killing; however,
our prior work using identical experimental conditions demonstrated that ROS are unlikely to
be a major factor in persister formation during the late stationary phase (Orman & Brynildsen,
2015). In that study, we overexpressed catalytically active antioxidant enzymes, including
catalases (KatE, KatG) and superoxide dismutases (SodA, SodB, SodC), yet observed no
significant change in persister levels. To further decouple ROS from respiratory activity in that
study, we performed anaerobic experiments using nitrate as an alternative terminal electron
acceptor. Interestingly, anaerobic respiration enhanced persister formation, and inhibition of
nitrate reductases with KCN reduced it—further supporting a ROS-independent mechanism.
Together, these findings suggest that under our experimental conditions, it is respiratory
activity rather than ROS production that plays a more central role in antibiotic tolerance.

In conclusion, a significant gap in the current literature is the lack of a comprehensive
understanding of how bacterial cell metabolism undergoes changes during the transition to a
tolerant state. Identifying the specific metabolic pathways that gain significance for cell survival
in this context is crucial. This knowledge can pave the way for the development of more
informed and targeted treatment strategies, ultimately enhancing our ability to combat
tolerant cells and improve overall treatment outcomes.

MATERIALS AND METHODS
Bacterial Strains and Plasmids
All experiments were conducted using *E. coli* K-12 MG1655 wild-type (WT) and its derivative
strains. *E. coli* K-12 MG1655, MO strains (carrying the mCherry expression system), *hipA7* (high
persister strain) and pUA66 plasmids were obtained from Mark P. Brynildsen at Princeton
University. *E. coli* MO strain was used to monitor cell proliferation at single cell level due to its
chromosomally integrated isopropyl β-D-1-thiogalactopyranoside (IPTG)-inducible mCherry
expression cassette (Orman & Brynildsen, 2013b, 2013a, 2015). *E. coli* K-12 BW25113 WT and
single deletions were obtained from Dharmacon Keio Collection (Dharmacon, Catalog#
OEC4988, Lafayette, CO, USA). The mutant strains in this study were generated using the
Datsenko-Wanner method (Datsenko & Wanner, 2000). The pUA66-EV was generated by the
removal of *gfp* gene from the plasmid. The *crp* gene with its promoter was cloned into the
modified pUA66 plasmid to obtain the pUA66-*crp* expression system. The *sdhABCD* reporter
(pMSs201-P$_{sdhABCD}$-*gfp*) and the *cyaA* reporter (pMSs201-P$_{cyaA}$-*gfp*) were obtained from a
previous study (Zaslaver et al., 2006). The cloning method was followed according to a standard
method from NEB (Tirabassi & Bio, 2014). Genetic modifications were verified by PCR and gene
sequencing (Genewiz, South Plainfield, NJ, USA). A complete list of strains, plasmids, and
oligonucleotides used in this study is presented in the **Key Resources Table**.
Media, Chemicals, and Culture Conditions

All chemicals used in this study were purchased from Fisher Scientific (Atlanta, GA, USA), VWR International (Pittsburgh, PA, USA), or Sigma Aldrich (St. Louis, MO, USA). Luria-Bertani (LB) medium was prepared by combining 5 g of yeast extract, 10 g of tryptone, and 10 g of sodium chloride in 1 L of autoclaved deionized (DI) water. LB agar media was prepared by mixing 40 g of pre-mixed LB agar with 1 L of autoclaved DI water; LB agar media were used to enumerate colony-forming units (CFUs) (Amato et al., 2013; Keren, Kaldalu, et al., 2004; Orman & Brynildsen, 2015). For washing cells and removing chemicals and antibiotics before plating on agar media, 1X Phosphate-Buffered Saline (PBS) was employed. In the persister assay, concentrations of 5 µg/mL of ofloxacin (OFX), 200 µg/mL of ampicillin (AMP), and 50 µg/mL of gentamicin (GEN) were used (Allison et al., 2011; De Groote et al., 2009; Keren, Kaldalu, et al., 2004; Keren, Shah, et al., 2004). The retention of plasmids necessitated 50 µg/mL of kanamycin (KAN) in the culture media (Orman & Brynildsen, 2015). Fluorescent protein expression was induced using 1 mM IPTG (Orman & Brynildsen, 2015). Overnight pre-cultures were prepared in 14 mL Falcon test tubes containing 2 mL of LB medium, inoculated from a 25% glycerol cell stock stored at -80°C, and incubated for 24 hours at 37°C with shaking at 250 revolutions per minute (rpm). Main cultures were established by diluting the overnight pre-cultures at a ratio of 1:1000 into 2 mL fresh LB medium in 14 mL Falcon test tubes. Experimental cell cultures were prepared by further dilution of the main cultures into either 25 mL fresh LB medium in 250 mL baffled flasks or 2 mL fresh LB medium in 14 mL Falcon test tubes. Cultures at t=5 h and t=24 h were defined as early and late stationary phase cultures, respectively. Detailed experimental procedures are outlined below.

Cell growth and persister assays

Main cultures were prepared by diluting the overnight pre-cultures at a ratio of 1:1000 into 2 mL of fresh LB medium in 14 mL Falcon test tubes. These cultures were then incubated at 37°C with shaking at 250 rpm. Cell growth was monitored by measuring the optical density at 600 nm wavelength (OD_{600}) using a Varioskan LUX Multimode Microplate Reader (Thermo Fisher, Waltham, MA, USA). The plate reader data was collected using SkanIt Software V 5.0. Cell cultures at both early and late stationary phases were collected from the test tubes and transferred to the baffled flasks to achieve ~$5x10^7$ cells/mL. This concentration represents an approximately 100-fold dilution of the WT main culture into a fresh medium within the flask. It is important to highlight that we consistently employed flow cytometry to quantify the initial cell count (refer to the section "Monitoring cell division" for comprehensive details). As needed, adjustments in cell number-to-volume were executed to ensure the same cell number among both WT and mutant strains. These cultures were then treated with antibiotics at the indicated concentrations and were cultured with shaking at 37°C for 20 hours. After treatment, a 1 mL sample from each flask was transferred to a microcentrifuge tube for a rigorous washing protocol to minimize antibiotic carryover. The samples were centrifuged at 13,300 RPM (17,000 × g) for 3 minutes, and >950 µL of supernatant was carefully removed without disturbing the pellet. The pellet was then resuspended in 950 µL of PBS, achieving a >20-fold dilution of residual antibiotics. This wash step was repeated once more, resulting in a cumulative >400-fold dilution. After the final wash, cells were resuspended in 100 µL of PBS. A 10 µL aliquot of this

593 suspension was serially diluted in a 96-well round-bottom plate, and 10 µL from each dilution
594 was spotted onto an agar plate. The remaining 90 µL of the sample was also plated to quantify
595 CFU levels around the limit of detection, (~1 CFU/mL). Plates were incubated at 37 °C for 16
596 hours (or up to 48 hours) to allow CFUs to develop. When necessary, this washing procedure
597 was repeated up to six times to assess its impact on CFU recovery.

Mid-exponential phase persister assays

Overnight cultures of *E. coli* K-12 MG1655 or BW25113 WT, and mutant strains were diluted
1:100 in 2 mL of LB medium in 14 mL Falcon tubes and incubated at 37°C with shaking at 250
rpm. At mid-exponential phase (OD_{600}~0.25), cells were challenged with the indicated
antibiotics and concentrations. At designated time points, cells (1 mL) were collected, washed
with 1X PBS, and plated on agar to determine colony-forming units (CFU). To quantify the initial
cell count, 10 µL of the culture before treatments was serially diluted and plated on LB agar. CFU
levels were assessed after incubating the plates for at least 16 hours at 37°C.

cAMP profile assay

An overnight pre-culture of *E. coli* K-12 MG1655 WT was prepared in test tubes with 2 mL of LB
medium. The culture was incubated at 37°C with shaking at 250 rpm for 24 hours. For the
experimental cultures, a 1:1000 dilution was made in fresh medium. At time-points 0, 2, 4, 6, 8
and 24 h, 100 µL of the experimental culture was washed with cold 1X PBS. After centrifuging at
13,300 rpm (17,000 x gravitational force or g), the supernatant was removed. The cells were
then resuspended in 100 µL of cell lysis buffer from the Cyclic AMP XP® Assay Kit (Catalog#
4339S, Cell Signaling Technology, Danvers, MA, USA) on ice for 10 minutes. Next, 50 µL of lysed
cells was mixed with the kit's horseradish peroxidase (HRP)-linked cAMP solution in the cAMP
assay plate. This mixture was incubated at room temperature for 3 hours on a plate shaker at
250 rpm. Following the 3-hour incubation, the plate content was discarded, and the plate was
washed four times with the kit's Wash Buffer. Then, 100 µL of tetramethylbenzidine (TMB)
substrate was added to allow color development, and after 30 minutes, 100 µL of the stop
solution provided by the kit was added. The absorbance was measured at 450 nm using a plate
reader. The standard curve was prepared using the same conditions and the standard cAMP
solutions provided by the kit to determine cAMP concentrations.

Metabolomics

Metabolites from both Δ*crp* and WT cells were analyzed at the Metabolon, Inc., facility
(Morrisville, NC, USA). Cells were cultured until the early stationary phase and late stationary
phase at 37°C with shaking at 250 rpm. Afterward, cells were collected through centrifugation at
4700 rpm at 37°C for 15 minutes, yielding a pellet of approximately 100 µL containing around
10^{10} cells. Subsequently, the cells were washed once with 1X PBS and then centrifuged (13,000
rpm, 3 minutes at 4°C). Following this, they were frozen in an ethanol/dry ice bath for 10
minutes. The extracts from both mutant and WT cells were subjected to analysis using ultra-
high-performance liquid chromatography-tandem accurate mass spectrometry (MS), a process
aimed at identifying a wide range of metabolites. Sample extraction, preparation, instrument
settings, and conditions for the MS platform adhered to Metabolon's protocols (as detailed in
our previous study) (Mohiuddin et al., 2022). To identify the sample's metabolites among

potential false positives from instrument noise, process artifacts, and redundant ion features,
the results were cross-referenced with Metabolon's extensive metabolite library (standards).
Data normalization was carried out based on protein concentration, determined using the
Bradford assay. The significant difference between mutant and WT was identified using Welch's
two-sample t-test. Further analysis involved pathway enrichment assessment using
MetaboAnalyst (Lu et al., 2023). This involved inputting the upregulated and downregulated
metabolites based on chosen thresholds. A comprehensive overview of metabolite
measurements, pathway enrichment, statistical analyses, and data representations can be
found in our previously published study (Mohiuddin et al., 2022).

Proteomics

Overnight cultures for both *E. coli* K-12 MG1655 WT and Δ*crp* strains were prepared using 2 mL
of LB medium. Incubation was carried out at 37°C and 250 rpm for 24 hours. The following day,
the main cultures were established under the same conditions, using a 1000-fold dilution of the
overnight culture in 2 mL fresh LB medium. After 24 hours, the OD_{600} of both WT and mutant
strains was measured and adjusted to an OD_{600} of 2.5. For further processing, 2 mL of the main
culture was washed twice with cold 1X PBS, maintaining the cold environment throughout.
Centrifugation conditions were set at 4°C, 13,000 rpm for 3 minutes. Before the final
centrifugation, a cell count was conducted using flow cytometry. This involved using 10 μL of
washed culture and 990 μL of 1X PBS. Subsequent to centrifugation, the pellets were collected.
Cell lysis was carried out using 300 μL of NEBExpress® *E. coli* Lysis Reagent (Catalog# P8116S,
Ipswich, MA, USA) at room temperature for 30 minutes. Following this, the lysed samples were
centrifuged at 16,600 x g for 10 minutes, and 250 μL of supernatants were collected from each
sample for the assay. The total protein concentration of the supernatants was determined using
the bicinchoninic acid (BCA) assay (Catalog# 23225, Thermo Fisher Scientific, Waltham, MA,
USA). In a 96-well plate, 25 μL of each cell lysate sample, diluted 5 and 10 times with ultra-pure
DI water, were loaded into each well. Subsequently, 200 μL of the BCA working reagent (50:1,
Reagent A:B) was added to each well and mixed on a plate shaker for 30 seconds. The plate was
then incubated at 37°C for 30 minutes, followed by cooling for 5 minutes at room temperature,
shielded from light. The absorbance was finally measured at 562 nm using a plate reader, and
the total protein concentration in each sample was calculated using a standard curve prepared
from standard protein solutions. Protein analysis for both WT and mutant was conducted by the
proteomics service at UT Health's Clinical and Translational Proteomics Service Center (Houston,
TX). The samples underwent acetone precipitation, during which proteins were precipitated by
exposing them to -20°C for 3 hours. Following this, a centrifugation step at 12,000 x g for 5
minutes separated the precipitated pellets. These pellets were subsequently subjected to
denaturation and reduction using a mixture containing 30 μL of 6 M urea, 20 mM DTT in 150
mM Tris HCl (pH 8.0) at 37°C for 40 minutes. Afterward, alkylation was carried out with 40 mM
iodoacetamide in the absence of light for 30 minutes. To prepare for digestion, the reaction
mixture was diluted 10-fold using 50 mM Tris-HCl (pH 8.0) and then incubated overnight at 37°C
with trypsin at a 1:30 enzyme-to-substrate ratio. The digestion process was terminated by
adding an equal volume of 2% formic acid, followed by desalting using Waters Oasis HLB 1 mL

reverse phase cartridges, following the vendor's recommended procedure. Finally, the elutes were dried using vacuum centrifugation. Approximately 1 µg of the tryptic digest, prepared in a solution containing 2% acetonitrile and 0.1% formic acid in water, underwent analysis using LC/MS/MS. The instrument used was the Orbitrap Fusion™ Tribrid™ mass spectrometer by Thermo Scientific™, connected to a Dionex UltiMate 3000 Binary RSLCnano System. The separation of peptides occurred on an analytical C18 column with dimensions of 100 µm ID x 25 cm, featuring 5 µm particles and an 18 Å pore size. Peptides were eluted at a flow rate of 350 nL/min. The gradient conditions applied were as follows: a gradient starting from 3% B and increasing to 22% B over a duration of 90 minutes, followed by a step to 22%-35% B for 10 minutes, then another step to 35%-90% B for 10 minutes, and finally, maintaining 90% B for an additional 10 minutes (Solvent A was composed of 0.1% formic acid in water, while solvent B contained 0.1% formic acid in acetonitrile). The peptides were analyzed using a data-dependent acquisition method. The Orbitrap Fusion MS operated by measuring FTMS1 spectra with a resolution of 120,000 FWHM, scanning in the m/z range of 350-1500, using an AGC target set to 2E5, and with a maximum injection time of 50 ms. Within a maximum cycle time of 3 sec, ITMS2 spectra were collected in rapid scan mode. High Collision Dissociation (HCD) was employed with a normalized collision energy (NCE) of 34, an isolation window of 1.6 m/z, an AGC target set to 1E4, and a maximum injection time of 35 ms. Dynamic exclusion was implemented for a duration of 35 sec to prevent repeated analysis of the same ions. For the experimental analysis, the Thermo Scientific™ Proteome Discoverer™ software version 1.4 was utilized to process the raw data files. The spectra were subjected to analysis against the *E. coli* proteome database (Swiss-Prot 29,161) through the Sequest HT search engine. Additionally, the spectra were compared against a decoy database, employing a target false discovery rate (FDR) of 1% for stringent criteria and 5% for more relaxed criteria. The enzymatic cleavage allowance for trypsin included up to two potential missed cleavages. The MS tolerance was defined as 10 ppm, while the MS/MS tolerance was set at 0.6 Da. Fixed modification involved carbamidomethylation on cysteine residues, and variable modifications encompassed methionine oxidation and asparagine deamidation. For proteomics data processing and fold change calculations, the approaches were essentially followed by the method paper from Aguilan et al. (Aguilan et al., 2020). Then, the STRING tool V12.0 (Szklarczyk et al., 2021, 2023) was employed to find the significant networks among input proteins. To generate the protein network and pathway enrichment analysis, we input the protein identifiers (Accession numbers) for upregulated or downregulated proteins with at least a 2-fold increase or reduction, respectively. *E. coli* K-12 was selected as the organism of interest. We opted for evidence as the criterion for network edges, prioritizing the type of interaction evidence, helping us conduct an automated pathway-enrichment analysis, centering on the entered proteins and identifying pathways that occurred more frequently than expected. This analysis was grounded in the statistical background of the entire genome and encompasses various functional pathway classification frameworks, such as Gene Ontology annotations, KEGG pathways, and Uniprot keywords, as detailed elsewhere (Szklarczyk et al., 2021, 2023). In pathway enrichment analysis, the "strength score," calculated as Log_{10}(observed/expected), serves to assess the degree or significance of enrichment within a

specific biological pathway. This metric reflects the magnitude of the enrichment effect, with a
higher score indicating stronger enrichment. The score is derived from the ratio of i) annotated
proteins in the network for a given term to ii) the expected number of proteins annotated with
the same term in a random network of equivalent size (Szklarczyk et al., 2021, 2023). To gauge
the significance of enrichment, False Discovery Rate (FDR) is employed. FDR scores represent p-
values corrected for multiple testing within each category using the Benjamini–Hochberg
procedure (Szklarczyk et al., 2021, 2023).

Monitoring cell division

To monitor cell division and quantify non-growing cells, we utilized inducible fluorescent protein
(mCherry) expression. Overnight pre-cultures of *E. coli* MO were prepared with 2 mL of LB
medium containing 1 mM of IPTG. These cultures were grown in test tubes at 37°C with shaking
at 250 rpm for 24 hours. Main cultures were established by diluting the overnight pre-cultures
(at a ratio of 1:1000) into 2 mL of fresh LB medium in 14 mL Falcon test tubes. These cultures
were incubated at 37°C with shaking at 250 rpm. Cells were allowed to grow until they reached
the early stationary phase and the late stationary phase. The mCherry positive cells were then
collected, washed twice with 1X PBS to remove the IPTG from the culture, and subsequently re-
suspended in fresh 2 ml LB media in test tubes to achieve ~$5x10^7$ cells/mL. This concentration
represents an approximately 100-fold dilution. When needed, adjustments in cell number-to-
volume were made to ensure the same cell number among both WT and mutant strains. In the
experimental culture test tubes, 2 mL of LB medium was added, and the volume of washed cells
was inoculated to achieve an OD_{600} of 0.0286. The culture was then incubated at 37°C with
shaking at 250 rpm. At specific time points (0, 1, 2, and 2.5 h), cells were collected and re-
suspended in 1X PBS to measure their fluorescent protein content using flow cytometry. For
flow cytometry analysis, cells were collected and diluted to a desired cell density (~10^6-10^7
cells/mL) in 1 mL of 1X PBS in flow cytometry tubes (5 mL round-bottom Falcon tubes, size: 12 x
75 mm). The flow cytometry analysis was conducted using a NovoCyte 3000RYB instrument
(ACEA Bioscience Inc., San Diego, CA, USA). During flow cytometry analysis, a slow sample flow
rate of 14 µL/min was chosen, along with a sample stream diameter (core diameter) of 7.7 µm.
The instrument maintained a constant sheath flow rate of 6.5 mL/min. The core diameter was
calculated using the ratio of the sample flow rate to the sheath flow rate. These specific
conditions were selected to achieve improved data resolution for the size of *E. coli* cells. Flow
diagrams utilized forward and side scatter signals from viable cells, alongside a control of
solvent devoid of cells, to ascertain the presence of cells. For the flow cytometry analysis, cells
were excited at a 561 nm wavelength, and the red fluorescence was detected using a 615/20
nm bandpass filter.

Flow Cytometry Analysis of Cell Growth

Overnight cultures of *E. coli* K-12 MG1655 MO WT and mutant strains were diluted at a ratio of
1:1000 into 2 mL of fresh LB medium, placed in 14 mL Falcon test tubes, and incubated at 37°C
with shaking at 250 rpm for 24 hours. For the main cultures, a similar strategy was employed.
The cultures were diluted at a ratio of 1:100 into 2 mL of fresh LB medium in 14 mL Falcon test
tubes. These cultures were then incubated at 37°C with shaking at 250 rpm. At specific time

points, including t=0, 20 min, 40 min, and 1 to 5 h, the cell growth was halted. This was
achieved by diluting the cells in 1X PBS containing 25 µg/mL of chloramphenicol (CAM). The
CAM treatment allowed for subsequent analysis without further division. Flow cytometry was
then utilized to measure the number of cells at each of these time points. This approach
provided insight into cell division dynamics and allowed for the quantification of cell
populations under specific conditions.

Fluorescent protein expression assay for reporter genes
Mutant and control strains were derived from *E. coli* K-12 MG1655 and carried pMSs201-*gfp*
plasmids incorporating P$_{sdhABCD}$ or P$_{cyaA}$ gene promoters. Overnight pre-cultures were prepared
using 2 mL of LB medium supplemented with 50 µg/mL KAN. These cultures were incubated in
test tubes within a shaker at 37°C for 24 hours. Main cultures were established by diluting the
overnight pre-cultures at a ratio of 1:1000 into 2 mL of fresh LB medium within 14 mL Falcon
test tubes. These main cultures were maintained at 37°C with shaking at 250 rpm. Cell cultures
at the desired growth phase were collected and then diluted to attain a desired cell density of
around 10^6-10^7 cells/mL in 1 mL of 1X PBS within flow cytometry tubes. This allowed for
subsequent flow cytometry analysis, using the same conditions as described earlier for
monitoring cell division (refer to "Monitoring cell division"). During analysis, a laser emitting
light at 488 nm was used to excite the cells, and the resulting green fluorescence was detected
using a 530/30 nm bandpass filter. This setup enabled the examination of the fluorescence
patterns of the cells, offering insights into their dynamics under different conditions.

Redox Sensor Green assay
To gauge bacterial metabolic activity, we employed the Redox Sensor Green (RSG) dye from
Thermo Fisher (Catalog# B34954, Thermo Fisher Scientific, Waltham, MA). *E. coli* K-12 MG1655
WT and mutant cells from the desired growth phase were diluted at a ratio of 1:100 in 1 mL of
1X PBS. To this solution, 1 µL of the RSG dye was added to flow cytometry tubes. After a brief
vortexing to ensure uniform mixing, the samples were incubated at 37°C in darkness for 10
minutes. Subsequently, these samples were subjected to flow cytometry analysis. For the flow
cytometry analysis, the same methodology as employed in "Monitoring cell division" was
followed, with one variation. Cells were excited at 488 nm during analysis, and the resulting
green fluorescence was detected using a 530/30 nm bandpass filter. This setup allowed us to
assess the fluorescence patterns, reflecting the metabolic activity of the bacterial cells under
different conditions. As a control measure, cells were treated with 20 µM of carbonyl cyanide
m-chlorophenyl hydrazone (CCCP) for 5 minutes before the addition of the RSG dye. This served
to validate the assay's sensitivity to changes in metabolic activity (**Figure 5–figure supplement
1**).

Metabolic activity of non-lysing cells and VBNC cell quantification
Overnight cultures of *E. coli* K-12 MG1655 MO WT and mutant strains were diluted at a ratio of
1:1000 into 2 mL of fresh LB medium supplemented with 1 mM IPTG. These cultures were
established in 14 mL Falcon test tubes and incubated at 37°C with shaking at 250 rpm for 24
hours. Treatment cultures were prepared by diluting the main cultures at a ratio of 1:100 into
25 mL of fresh LB medium supplemented with 1 mM IPTG. These cultures were set up in 250 mL

baffled flasks and contained 200 μg/mL AMP. They were then cultured at 37°C with shaking at
250 rpm for 20 hours. Both before and after the treatment, 1 mL samples were collected from
the cultures. These samples were subjected to a washing procedure with 1X PBS to eliminate
the antibiotic present in the samples. The washed cells were then resuspended in 1 mL of 1X
PBS within flow cytometry tubes. To measure the metabolic activity of the non-lysing cells, the
RSG dye was employed as described above. Intact cells (following antibiotic treatment), stained
as live with RSG, comprised both persister and VBNC cells. Persister levels were quantified by
plating the cells on an agar medium, as described previously (Orman & Brynildsen, 2013b). As
VBNC cells cannot grow on agar medium, their enumeration involved subtracting the number of
persister cells from the total number of intact cells.

Screening *E. coli* (K-12 BW25113) Keio Knockout Collection
Overnight cultures of individual mutant strains, along with their parental strain K-12 BW25113
WT harboring a kanamycin-resistant marker, were diluted at a ratio of 1:1000 in fresh LB
medium containing 50 μg/mL of KAN. This was done in 14 mL Falcon test tubes and the cultures
were then incubated at 37°C with shaking at 250 rpm. Upon reaching the late stationary phase,
cells were further diluted at a ratio of 1:100 in fresh medium supplemented with antibiotics at
specified concentrations. These cultures were once again incubated at 37°C with shaking for 20
hours. Following the 20-hour treatment period, the same methodology described in the section
"Cell growth and persister assays" was employed to quantify the number of persisters. This
approach allowed for an assessment of the impact of antibiotics on the formation of persister
cells for both mutant strains and the parental K-12 BW25113 WT strain.

Persister quantitation in *E. coli* K-12 MG1655 single gene deletions
Overnight cultures of mutant strains were diluted at a ratio of 1:1000 in 14 mL Falcon test tubes
containing 2 mL of LB medium. These cultures were then incubated at 37°C with shaking at 250
rpm. Upon reaching the late stationary phase, cells were diluted at a ratio of 1:100 in fresh
medium supplemented with antibiotics at specified concentrations. The cultures were once
again subjected to shaking at 37°C for 20 hours. The same method described earlier, referred to
as "Cell growth and persister assays," was employed to quantify the number of persister cells
resulting from this treatment. This approach allowed for the assessment of the impact of
antibiotic exposure on persister cell formation within the mutant strains.

Statistics and reproducibility
A nonlinear logarithmic model was employed to create biphasic kill curves (Mohiuddin, Hoang,
et al., 2020; Windels et al., 2019). The significance of these kill curves was determined through
the utilization of F-statistics (Mohiuddin, Hoang, et al., 2020; Windels et al., 2019).
Metabolomics data were subjected to analysis using Welch's two-sample t-test in order to
identify metabolites that significantly differed between the control and mutant groups (YUEN,
1974). For all experiments, a minimum of three independent biological replicates were
conducted, unless explicitly stated otherwise. In each figure (excluding flow diagrams), the data
for each time point are represented as the mean value accompanied by the standard deviation.
In terms of statistical significance analysis, the designated threshold values for P were set as
follows: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. All figures were generated

using GraphPad Prism 10.3.0. The statistical analyses were carried out using the statistical
functions of GraphPad Prism 10.3.0. For the clustering of metabolomics and proteomics data,
the "Clustergram" function of MATLAB (V R2020b) was employed. FlowJo (V 10.8.1) was the tool
used to analyze the data acquired from flow cytometry.
843
844 **Data Availability**
845 All data generated or analyzed during this study are included in the manuscript and supporting
846 files.
847

Article and author information
Author details
Han G Ngo
Cullen College of Engineering, William A. Brookshire Department of Chemical and Biomolecular
Engineering, University of Houston, Houston, United States
Contribution: Conception and design, Acquisition of data, Analysis and interpretation of data,
Drafting or revising the article
Competing interests: No competing interests declared
Sayed G Mohiuddin
Cullen College of Engineering, William A. Brookshire Department of Chemical and Biomolecular
Engineering, University of Houston, Houston, United States
Contribution: Acquisition of data, Analysis and interpretation of data, Drafting or revising the
article
Competing interests: No competing interests declared
Aina Ananda
School of Science, Department of Biology, Monmouth University, West Long Branch, United
States
Contribution: Analysis of proteomics data, Drafting or revising the article
Competing interests: No competing interests declared
Mehmet A Orman
Cullen College of Engineering, William A. Brookshire Department of Chemical and Biomolecular
Engineering, University of Houston, Houston, United States
Contribution: Conception and design, Acquisition of data, Analysis and interpretation of data,
Drafting or revising the article
For correspondence: morman@central.uh.edu
Competing interests: No competing interests declared

Funding
National Science Foundation (2044375)
• Mehmet A Orman

National Institutes of Health (R01AI143643)
• Mehmet A Orman

The funders had no role in study design, data collection and interpretation, or the decision to
submit the work for publication.

Acknowledgements

We would like to express our gratitude to the members of Dr. Orman's laboratory for their support and assistance throughout this study. We also thank Metabolon, Inc. for their expertise in metabolomics experiments and data analysis, as well as UT Health's Clinical and Translational Proteomics Service Center for their contribution to the proteomics experiments.

References

Adams, K. N., Takaki, K., Connolly, L. E., Wiedenhoft, H., Winglee, K., Humbert, O., Edelstein, P. H., Cosma, C. L., & Ramakrishnan, L. (2011). Drug tolerance in replicating mycobacteria mediated by a macrophage-induced efflux mechanism. *Cell*, *145*(1), 39–53. https://doi.org/10.1016/j.cell.2011.02.022

Aguilan, J. T., Kulej, K., & Sidoli, S. (2020). Guide for protein fold change and *p*-value calculation for non-experts in proteomics. *Molecular Omics*, *16*(6), 573–582. https://doi.org/10.1039/D0MO00087F

Aiba, H. (1985). Transcription of the Escherichia coli adenylate cyclase gene is negatively regulated by cAMP-cAMP receptor protein. *The Journal of Biological Chemistry*, *260*(5), 3063–3070.

Allison, K. R., Brynildsen, M. P., & Collins, J. J. (2011). Metabolite-enabled eradication of bacterial persisters by aminoglycosides. *Nature*, *473*(7346), 216–220. https://doi.org/10.1038/nature10069

Amato, S. M., Fazen, C. H., Henry, T. C., Mok, W. W. K., Orman, M. A., Sandvik, E. L., Volzing, K. G., & Brynildsen, M. P. (2014). The role of metabolism in bacterial persistence. *Frontiers in Microbiology*, *5*, 70.

Amato, S. M., Orman, M. A., & Brynildsen, M. P. (2013). Metabolic Control of Persister Formation in Escherichia coli. *Molecular Cell*, *50*(4), 475–487. https://doi.org/10.1016/J.MOLCEL.2013.04.002

Arnoldini, M., Vizcarra, I. A., Peña-Miller, R., Stocker, N., Diard, M., Vogel, V., Beardmore, R. E., Hardt, W.-D., & Ackermann, M. (2014). Bistable Expression of Virulence Genes in Salmonella Leads to the Formation of an Antibiotic-Tolerant Subpopulation. *PLoS Biology*, *12*(8), e1001928. https://doi.org/10.1371/journal.pbio.1001928

Ayrapetyan, M., Williams, T., & Oliver, J. D. (2018). Relationship between the Viable but Nonculturable State and Antibiotic Persister Cells. *Journal of Bacteriology*, *200*(20). https://doi.org/10.1128/JB.00249-18

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L., & Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular Systems Biology*, *2*(1). https://doi.org/10.1038/msb4100050

Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L., & Leibler, S. (2004). Bacterial persistence as a phenotypic switch. *Science*, *305*(5690), 1622–1625. https://doi.org/10.1126/SCIENCE.1099390

Barrett, T. C., Mok, W. W. K., Murawski, A. M., & Brynildsen, M. P. (2019). Enhanced antibiotic
resistance development from fluoroquinolone persisters after a single exposure to
antibiotic. *Nature Communications*, *10*(1), 1177.
Bettenbrock, K., Sauter, T., Jahreis, K., Kremling, A., Lengeler, J. W., & Gilles, E.-D. (2007).
Correlation between growth rates, EIIACrr phosphorylation, and intracellular cyclic AMP
levels in Escherichia coli K-12. *Journal of Bacteriology*, *189*(19), 6891–6900.
Chu, W., Zere, T. R., Weber, M. M., Wood, T. K., Whiteley, M., Hidalgo-Romano, B., Valenzuela
Jr, E., & McLean, R. J. C. (2012). Indole production promotes Escherichia coli mixed-culture
growth with Pseudomonas aeruginosa by inhibiting quorum signaling. *Applied and*
Environmental Microbiology, *78*(2), 411–419.
Conlon, B. P., Rowe, S. E., Gandt, A. B., Nuxoll, A. S., Donegan, N. P., Zalis, E. A., Clair, G., Adkins,
J. N., Cheung, A. L., & Lewis, K. (2016). Persister formation in Staphylococcus aureus is
associated with ATP depletion. *Nature Microbiology*, *1*(5), 1–7.
Cunningham, L., & Guest, J. R. (1998). Transcription and transcript processing in the sdhCDAB-
sucABCD operon of Escherichia coli. *Microbiology (Reading, England)*, *144 (Pt 8)*, 2113–
2123. https://doi.org/10.1099/00221287-144-8-2113
Datsenko, K. A., & Wanner, B. L. (2000). One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of Sciences*,
97(12), 6640–6645. https://doi.org/10.1073/pnas.120163297
De Groote, V. N., Verstraeten, N., Fauvart, M., Kint, C. I., Verbeeck, A. M., Beullens, S., Cornelis,
P., & Michiels, J. (2009). Novel persistence genes in *Pseudomonas aeruginosa* identified by
high-throughput screening. *FEMS Microbiology Letters*, *297*(1), 73–79.
https://doi.org/10.1111/j.1574-6968.2009.01657.x
Deutscher, J. (2008). The mechanisms of carbon catabolite repression in bacteria. *Current*
Opinion in Microbiology, *11*(2), 87–93.
Fic, E., Bonarek, P., Górecki, A., Kedracka-Krok, S., Mikolajczak, J., Polit, A., Tworzydlo, M.,
Dziedzicka-Wasylewska, M., & Wasylewski, Z. (2009). cAMP receptor protein from
Escherichia coli as a model of signal transduction in proteins–a review. *Microbial*
Physiology, *17*(1), 1–11.
Fisher, R. A., Gollan, B., & Helaine, S. (2017). Persistent bacterial infections and persister cells.
Nature Reviews Microbiology, *15*(8), 453–464. https://doi.org/10.1038/nrmicro.2017.42
Gonzalez-Gil, G. (1998). Regulation of crp transcription by oscillation between distinct
nucleoprotein complexes. *The EMBO Journal*, *17*(10), 2877–2885.
https://doi.org/10.1093/emboj/17.10.2877
Goormaghtigh, F., & Van Melderen, L. (2019). Single-cell imaging and characterization of
Escherichia coli persister cells to ofloxacin in exponential cultures. *Science Advances*, *5*(6),
eaav9462.
Görke, B., & Stülke, J. (2008). Carbon catabolite repression in bacteria: many ways to make the
most out of nutrients. *Nature Reviews Microbiology*, *6*(8), 613–624.
Gosset, G., Zhang, Z., Nayyar, S., Cuevas, W. A., & Saier, M. H. (2004). Transcriptome Analysis of
Crp-Dependent Catabolite Control of Gene Expression in *Escherichia coli*. *Journal of*
Bacteriology, *186*(11), 3516–3524. https://doi.org/10.1128/JB.186.11.3516-3524.2004
Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: the next generation. *Cell*, *144*(5),
646–674.

Hofsteenge, N., van Nimwegen, E., & Silander, O. K. (2013). Quantitative analysis of persister
fractions suggests different mechanisms of formation among environmental isolates of E.
coli. *BMC Microbiology*, *13*(1), 25. https://doi.org/10.1186/1471-2180-13-25
Hossain, T., Singh, A., & Butzin, N. C. (2023). *Escherichia coli* cells are primed for survival before
lethal antibiotic stress. *Microbiology Spectrum*, *11*(5).
https://doi.org/10.1128/spectrum.01219-23
Huemer, M., Mairpady Shambat, S., Brugger, S. D., & Zinkernagel, A. S. (2020). Antibiotic
resistance and persistence—Implications for human health and treatment perspectives.
EMBO Reports, *21*(12). https://doi.org/10.15252/embr.202051034
Joers, A., Kaldalu, N., & Tenson, T. (2010). The frequency of persisters in Escherichia coli reflects
the kinetics of awakening from dormancy. *Journal of Bacteriology*, *192*(13), 3379–3384.
Jõers, A., & Tenson, T. (2016). Growth resumption from stationary phase reveals memory in
Escherichia coli cultures. *Scientific Reports*, *6*(1), 24055.
https://doi.org/10.1038/srep24055
Kaldalu, N., & Tenson, T. (2019). Slow growth causes bacterial persistence. *Science Signaling*,
12(592). https://doi.org/10.1126/scisignal.aay1167
Keren, I., Kaldalu, N., Spoering, A., Wang, Y., & Lewis, K. (2004). Persister cells and tolerance to
antimicrobials. *FEMS Microbiology Letters*, *230*(1), 13–18.
Keren, I., Shah, D., Spoering, A., Kaldalu, N., & Lewis, K. (2004). Specialized Persister Cells and
the Mechanism of Multidrug Tolerance in *Escherichia coli*. *Journal of Bacteriology*, *186*(24),
8172–8180. https://doi.org/10.1128/JB.186.24.8172-8180.2004
Keseler, I. M. (2004). EcoCyc: a comprehensive database resource for Escherichia coli. *Nucleic
Acids Research*, *33*(Database issue), D334–D337. https://doi.org/10.1093/nar/gki108
Keseler, I. M., Collado-Vides, J., Santos-Zavaleta, A., Peralta-Gil, M., Gama-Castro, S., Muniz-
Rascado, L., Bonavides-Martinez, C., Paley, S., Krummenacker, M., Altman, T., Kaipa, P.,
Spaulding, A., Pacheco, J., Latendresse, M., Fulcher, C., Sarker, M., Shearer, A. G., Mackie,
A., Paulsen, I., … Karp, P. D. (2011). EcoCyc: a comprehensive database of Escherichia coli
biology. *Nucleic Acids Research*, *39*(Database), D583–D590.
https://doi.org/10.1093/nar/gkq1143
Khankal, R., Chin, J. W., Ghosh, D., & Cirino, P. C. (2009). Transcriptional effects of CRP*
expression in Escherichia coli. *Journal of Biological Engineering*, *3*(1), 13.
https://doi.org/10.1186/1754-1611-3-13
Kim, J.-S., Cho, D.-H., Heo, P., Jung, S.-C., Park, M., Oh, E.-J., Sung, J., Kim, P.-J., Lee, S.-C., Lee,
D.-H., Lee, S., Lee, C. H., Shin, D., Jin, Y.-S., & Kweon, D.-H. (2016). Fumarate-Mediated
Persistence of Escherichia coli against Antibiotics. *Antimicrobial Agents and Chemotherapy*,
60(4), 2232–2240. https://doi.org/10.1128/AAC.01794-15
Kohanski, M. A., Dwyer, D. J., Hayete, B., Lawrence, C. A., & Collins, J. J. (2007). A Common
Mechanism of Cellular Death Induced by Bactericidal Antibiotics. *Cell*, *130*(5), 797–810.
https://doi.org/10.1016/j.cell.2007.06.049
Kolb, A., Busby, S., Buc, H., Garges, S., & Adhya, S. (1993). Transcriptional regulation by cAMP
and its receptor protein. *Annual Review of Biochemistry*, *62*(1), 749–797.
Leatham-Jensen, M. P., Mokszycki, M. E., Rowley, D. C., Deering, R., Camberg, J. L., Sokurenko,
E. V., Tchesnokova, V. L., Frimodt-Møller, J., Krogfelt, K. A., Leth Nielsen, K., Frimodt-
Møller, N., Sun, G., & Cohen, P. S. (2016). Uropathogenic Escherichia coli Metabolite-

Dependent Quiescence and Persistence May Explain Antibiotic Tolerance during Urinary Tract Infection. *MSphere*, *1*(1). https://doi.org/10.1128/mSphere.00055-15

Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., & Balaban, N. Q. (2017). Antibiotic tolerance facilitates the evolution of resistance. *Science*, *355*(6327), 826–830.

Lewis, K. (2010). Persister cells. In *Annual Review of Microbiology* (Vol. 64). https://doi.org/10.1146/annurev.micro.112408.134306

Li, B., Chen, X., Yang, J.-Y., Gao, S., & Bai, F. (2024). Intracellular ATP concentration is a key regulator of bacterial cell fate. *Journal of Bacteriology*, *206*(12). https://doi.org/10.1128/jb.00208-24

Liu, S., Wu, N., Zhang, S., Yuan, Y., Zhang, W., & Zhang, Y. (2017). Variable Persister Gene Interactions with (p)ppGpp for Persister Formation in Escherichia coli. *Frontiers in Microbiology*, *8*. https://doi.org/10.3389/fmicb.2017.01795

Lobritz, M. A., Belenky, P., Porter, C. B. M., Gutierrez, A., Yang, J. H., Schwarz, E. G., Dwyer, D. J., Khalil, A. S., & Collins, J. J. (2015). Antibiotic efficacy is linked to bacterial cellular respiration. *Proceedings of the National Academy of Sciences*, *112*(27), 8173–8180. https://doi.org/10.1073/pnas.1509743112

Lu, Y., Pang, Z., & Xia, J. (2023). Comprehensive investigation of pathway enrichment methods for functional interpretation of LC–MS global metabolomics data. *Briefings in Bioinformatics*, *24*(1), bbac553.

Luidalepp, H., Joers, A., Kaldalu, N., & Tenson, T. (2011). Age of inoculum strongly influences persister frequency and can mask effects of mutations implicated in altered persistence. *Journal of Bacteriology*, *193*(14), 3598–3605.

Lynch, A. S., & Lin, E. C. (1996). Transcriptional control mediated by the ArcA two-component response regulator protein of Escherichia coli: characterization of DNA binding at target promoters. *Journal of Bacteriology*, *178*(21), 6238–6249. https://doi.org/10.1128/jb.178.21.6238-6249.1996

Ma, C., Sim, S., Shi, W., Du, L., Xing, D., & Zhang, Y. (2010). Energy production genes sucB and ubiF are involved in persister survival and tolerance to multiple antibiotics and stresses in Escherichia coli. *FEMS Microbiology Letters*, *303*(1), 33–40.

Majerfeld, I. H., Miller, D., Spitz, E., & Rickenberg, H. V. (1981). Regulation of the synthesis of adenylate cyclase in Escherichia coli by the cAMP — cAMP receptor protein complex. *Molecular and General Genetics MGG*, *181*(4), 470–475. https://doi.org/10.1007/BF00428738

Manuse, S., Shan, Y., Canas-Duarte, S. J., Bakshi, S., Sun, W.-S., Mori, H., Paulsson, J., & Lewis, K. (2021). Bacterial persisters are a stochastically formed subpopulation of low-energy cells. *PLoS Biology*, *19*(4), e3001194.

Michiels, J. E., Van den Bergh, B., Verstraeten, N., & Michiels, J. (2016). Molecular mechanisms and clinical implications of bacterial persistence. *Drug Resistance Updates*, *29*, 76–89. https://doi.org/10.1016/j.drup.2016.10.002

Mohiuddin, S. G., Hoang, T., Saba, A., Karki, P., & Orman, M. A. (2020). Identifying Metabolic Inhibitors to Reduce Bacterial Persistence. *Frontiers in Microbiology*, *11*. https://doi.org/10.3389/fmicb.2020.00472

Mohiuddin, S. G., Kavousi, P., & Orman, M. A. (2020). Flow-cytometry analysis reveals persister resuscitation characteristics. *BMC Microbiology*, *20*(1), 202. https://doi.org/10.1186/s12866-020-01888-3

Mohiuddin, S. G., Nguyen, T. V., & Orman, M. A. (2022). Pleiotropic actions of phenothiazine drugs are detrimental to Gram-negative bacterial persister cells. *Communications Biology*, *5*(1), 217.

Mok, W. W. K., Orman, M. A., & Brynildsen, M. P. (2015). Impacts of global transcriptional regulators on persister metabolism. *Antimicrobial Agents and Chemotherapy*, *59*(5), 2713–2719.

Mok, W. W. K., Park, J. O., Rabinowitz, J. D., & Brynildsen, M. P. (2015). RNA Futile Cycling in Model Persisters Derived from MazF Accumulation. *MBio*, *6*(6). https://doi.org/10.1128/mBio.01588-15

Molina-Quiroz, R. C., Silva-Valenzuela, C., Brewster, J., Castro-Nallar, E., Levy, S. B., & Camilli, A. (2018). Cyclic AMP regulates bacterial persistence through repression of the oxidative stress response and SOS-dependent DNA repair in uropathogenic Escherichia coli. *MBio*, *9*(1), 10–1128.

Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Robles Aguilar, G., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-Woodhouse, G., Kashef Hamadani, B. H., Kumaran, E. A. P., McManigal, B., … Naghavi, M. (2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*, *399*(10325), 629–655. https://doi.org/10.1016/S0140-6736(21)02724-0

Opalek, M., Smug, B. J., & Wloch-Salamon, D. (2022). How to determine microbial lag phase duration? *BioRxiv*, 2011–2022.

Orman, M. A., & Brynildsen, M. P. (2013a). Dormancy is not necessary or sufficient for bacterial persistence. *Antimicrobial Agents and Chemotherapy*, *57*(7), 3230–3239.

Orman, M. A., & Brynildsen, M. P. (2013b). Establishment of a method to rapidly assay bacterial persister metabolism. *Antimicrobial Agents and Chemotherapy*, *57*(9), 4398–4409.

Orman, M. A., & Brynildsen, M. P. (2015). Inhibition of stationary phase respiration impairs persister formation in E. coli. *Nature Communications*, *6*(1), 7983.

Pandey, S., Sahukhal, G. S., & Elasri, M. O. (2021). The msaABCR Operon Regulates Persister Formation by Modulating Energy Metabolism in Staphylococcus aureus. *Frontiers in Microbiology*, *12*. https://doi.org/10.3389/fmicb.2021.657753

Park, Y.-H., Lee, B. R., Seok, Y.-J., & Peterkofsky, A. (2006). In vitro reconstitution of catabolite repression in Escherichia coli. *Journal of Biological Chemistry*, *281*(10), 6448–6454.

Parsons, J. B., Sidders, A. E., Velez, A. Z., Hanson, B. M., Angeles-Solano, M., Ruffin, F., Rowe, S. E., Arias, C. A., Fowler, V. G., Thaden, J. T., & Conlon, B. P. (2024). In-patient evolution of a high-persister Escherichia coli strain with reduced in vivo antibiotic susceptibility. *Proceedings of the National Academy of Sciences of the United States of America*, *121*(3). https://doi.org/10.1073/pnas.2314514121

Pastan, I., & Perlman, R. (1970). Cyclic Adenosine Monophosphate in Bacteria: In many bacteria the synthesis of inducible enzymes requires this cyclic nucleotide. *Science*, *169*(3943), 339–344.

Perrenoud, A., & Sauer, U. (2005). Impact of Global Transcriptional Regulation by ArcA, ArcB, Cra, Crp, Cya, Fnr, and Mlc on Glucose Catabolism in *Escherichia coli*. *Journal of Bacteriology*, *187*(9), 3171–3179. https://doi.org/10.1128/JB.187.9.3171-3179.2005

Porporato, P. E., Filigheddu, N., Pedro, J. M. B.-S., Kroemer, G., & Galluzzi, L. (2018). Mitochondrial metabolism and cancer. *Cell Research*, *28*(3), 265–280.

Prax, M., & Bertram, R. (2014). Metabolic aspects of bacterial persisters. *Frontiers in Cellular and Infection Microbiology*, *4*, 148.

Quail, M. A., Haydon, D. J., & Guest, J. R. (1994). The pdhR-aceEF-lpd operon of Escherichia coli expresses the pyruvate dehydrogenase complex. *Molecular Microbiology*, *12*(1), 95–104. https://doi.org/10.1111/j.1365-2958.1994.tb00998.x

Radzikowski, J. L., Vedelaar, S., Siegel, D., Ortega, Á. D., Schmidt, A., & Heinemann, M. (2016). Bacterial persistence is an active σS stress response to metabolic flux limitation. *Molecular Systems Biology*, *12*(9), 882.

Roostalu, J., J\~oers, A., Luidalepp, H., Kaldalu, N., & Tenson, T. (2008). Cell division in Escherichia coli cultures monitored at single cell resolution. *BMC Microbiology*, *8*, 1–14.

Salgado, H., Peralta-Gil, M., Gama-Castro, S., Santos-Zavaleta, A., Muñiz-Rascado, L., García-Sotelo, J. S., Weiss, V., Solano-Lira, H., Martínez-Flores, I., Medina-Rivera, A., Salgado-Osorio, G., Alquicira-Hernández, S., Alquicira-Hernández, K., López-Fuentes, A., Porrón-Sotelo, L., Huerta, A. M., Bonavides-Martínez, C., Balderas-Martínez, Y. I., Pannier, L., … Collado-Vides, J. (2013). RegulonDB v8.0: omics data sets, evolutionary conservation, regulatory phrases, cross-validated gold standards and more. *Nucleic Acids Research*, *41*(D1), D203–D213. https://doi.org/10.1093/nar/gks1201

Shan, Y., Brown Gandt, A., Rowe, S. E., Deisinger, J. P., Conlon, B. P., & Lewis, K. (2017). ATP-dependent persister formation in Escherichia coli. *MBio*, *8*(1), 10–1128.

Shan, Y., Lazinski, D., Rowe, S., Camilli, A., & Lewis, K. (2015). Genetic Basis of Persister Tolerance to Aminoglycosides in Escherichia coli. *MBio*, *6*(2). https://doi.org/10.1128/mBio.00078-15

Shiraliyev, R. C., & Orman, M. (2023). Metabolic disruption impairs ribosomal protein levels, resulting in enhanced aminoglycoside tolerance. *BioRxiv*, 2012–2023.

Spoering, A. L., Vulić, M., & Lewis, K. (2006). GlpD and PlsB Participate in Persister Cell Formation in *Escherichia coli*. *Journal of Bacteriology*, *188*(14), 5136–5144. https://doi.org/10.1128/JB.00369-06

Sulaiman, J. E., & Lam, H. (2020). Proteomic investigation of tolerant Escherichia coli populations from cyclic antibiotic treatment. *Journal of Proteome Research*, *19*(2), 900–913.

Surmann, K., Stopp, M., Wörner, S., Dhople, V. M., Völker, U., Unden, G., & Hammer, E. (2020). Fumarate dependent protein composition under aerobic and anaerobic growth conditions in Escherichia coli. *Journal of Proteomics*, *212*, 103583. https://doi.org/10.1016/j.jprot.2019.103583

Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T., Legeay, M., Fang, T., Bork, P., Jensen, L. J., & von Mering, C. (2021). The STRING database in 2021: customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Research*, *49*(D1), D605–D612. https://doi.org/10.1093/nar/gkaa1074

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L., Fang,
T., Doncheva, N. T., Pyysalo, S., & others. (2023). The STRING database in 2023: protein–
protein association networks and functional enrichment analyses for any sequenced
genome of interest. *Nucleic Acids Research*, *51*(D1), D638–D646.
Taber, H. W., Mueller, J. P., Miller, P. F., & Arrow, A. S. (1987). Bacterial uptake of
aminoglycoside antibiotics. *Microbiological Reviews*, *51*(4), 439–457.
https://doi.org/10.1128/mr.51.4.439-457.1987
Tirabassi, R., & Bio, B. (2014). Foundations of Molecular Cloning-Past, Present and Future. *NEB,*
New England Bio Labs Inc, Www. Neb. Com/Tools-and-
Resources/Featurearticles/Foundations-of-Molecular-Cloning-Past-Present-and-Future.
Tsai, M.-J., Wang, J.-R., Yang, C.-D., Kao, K.-C., Huang, W.-L., Huang, H.-Y., Tseng, C.-P., Huang,
H.-D., & Ho, S.-Y. (2018). PredCRP: predicting and analysing the regulatory roles of CRP
from its binding sites in Escherichia coli. *Scientific Reports*, *8*(1), 951.
https://doi.org/10.1038/s41598-017-18648-5
Vasan, K., Werner, M., & Chandel, N. S. (2020). Mitochondrial metabolism as a target for cancer
therapy. *Cell Metabolism*, *32*(3), 341–352.
Vogel, R. F., Entian, K. D., & Mecke, D. (1987). Cloning and sequence of the mdh structural gene
of Escherichia coli coding for malate dehydrogenase. *Archives of Microbiology*, *149*(1), 36–
42. https://doi.org/10.1007/BF00423133
Vulin, C., Leimer, N., Huemer, M., Ackermann, M., & Zinkernagel, A. S. (2018). Prolonged
bacterial lag time results in small colony variants that represent a sub-population of
persisters. *Nature Communications*, *9*(1), 4074. https://doi.org/10.1038/s41467-018-
06527-0
Wakamoto, Y., Dhar, N., Chait, R., Schneider, K., Signorino-Gelo, F., Leibler, S., & McKinney, J. D.
(2013). Dynamic Persistence of Antibiotic-Stressed Mycobacteria. *Science*, *339*(6115), 91–
95. https://doi.org/10.1126/science.1229858
Wang, Y., Bojer, M. S., George, S. E., Wang, Z., Jensen, P. R., Wolz, C., & Ingmer, H. (2018).
Inactivation of TCA cycle enhances Staphylococcus aureus persister cell formation in
stationary phase. *Scientific Reports*, *8*(1), 10849. https://doi.org/10.1038/s41598-018-
29123-0
Warburg, O. (1925). The metabolism of carcinoma cells. *The Journal of Cancer Research*, *9*(1),
148–163.
Windels, E. M., Michiels, J. E., Fauvart, M., Wenseleers, T., den Bergh, B., & Michiels, J. (2019).
Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival
and mutation rates. *The ISME Journal*, *13*(5), 1239–1251.
Yamasaki, R., Song, S., Benedik, M. J., & Wood, T. K. (2020). Persister cells resuscitate using
membrane sensors that activate chemotaxis, lower cAMP levels, and revive ribosomes.
Iscience, *23*(1).
Yu, J., Liu, Y., Yin, H., & Chang, Z. (2019). Regrowth-delay body as a bacterial subcellular
structure marking multidrug-tolerant persisters. *Cell Discovery*, *5*(1), 8.
YUEN, K. K. (1974). The two-sample trimmed *t* for unequal population variances. *Biometrika*,
61(1), 165–170. https://doi.org/10.1093/biomet/61.1.165

Zalis, E. A., Nuxoll, A. S., Manuse, S., Clair, G., Radlinski, L. C., Conlon, B. P., Adkins, J., & Lewis, K.
(2019). Stochastic Variation in Expression of the Tricarboxylic Acid Cycle Produces Persister
Cells. *MBio*, *10*(5). https://doi.org/10.1128/mBio.01930-19
Zaslaver, A., Bren, A., Ronen, M., Itzkovitz, S., Kikoin, I., Shavit, S., Liebermeister, W., Surette, M.
G., & Alon, U. (2006). A comprehensive library of fluorescent transcriptional reporters for
Escherichia coli. *Nature Methods*, *3*(8), 623–628.
Zeng, A., Fan, Y., Di, Z., Wang, Y., & Havlin, S. (2022). *Impactful scientists have higher tendency*
to involve collaborators in new topics. https://doi.org/10.1073/pnas
Zeng, J., Hong, Y., Zhao, N., Liu, Q., Zhu, W., Xiao, L., Wang, W., Chen, M., Hong, S., Wu, L., Xue,
Y., Wang, D., Niu, J., Drlica, K., & Zhao, X. (2022). A broadly applicable, stress-mediated
bacterial death pathway regulated by the phosphotransferase system (PTS) and the cAMP-
Crp cascade. *Proceedings of the National Academy of Sciences*, *119*(23).
https://doi.org/10.1073/pnas.2118566119
Zhang, Z., Gosset, G., Barabote, R., Gonzalez, C. S., Cuevas, W. A., & Saier, M. H. (2005).
Functional interactions between the carbon and iron utilization regulators, Crp and Fur, in
Escherichia coli. *Journal of Bacteriology*, *187*(3), 980–990.
https://doi.org/10.1128/JB.187.3.980-990.2005
Zheng, D. (2004). Identification of the CRP regulon using in vitro and in vivo transcriptional
profiling. *Nucleic Acids Research*, *32*(19), 5874–5893. https://doi.org/10.1093/nar/gkh908
Zheng, E. J., Stokes, J. M., & Collins, J. J. (2020). Eradicating Bacterial Persisters with
Combinations of Strongly and Weakly Metabolism-Dependent Antibiotics. *Cell Chemical*
Biology. https://doi.org/10.1016/j.chembiol.2020.08.015

1203

Figure Legends

Figure 1: Crp/cAMP Regulation of Persister Cell Formation in the Stationary Phase. *E. coli* K-12
MG1655 WT and mutant cells at early (t=5 h) and late (t=24 h) stationary phases were
transferred to fresh medium with antibiotics for persister cell quantification. At time points 0, 2,
6, 8, and 20 h, 1 mL of the treated culture was washed with 1X phosphate-buffered saline (PBS)
to remove antibiotics. It was then serially diluted and plated on an agar plate to count the
colony-forming units (CFUs). **(A)** Persister levels of ampicillin-treated culture with an antibiotic
concentration of 200 µg/mL. **(B)** Persister levels of ofloxacin-treated culture with an antibiotic
concentration of 5 µg/mL. **(C)** Persister levels of gentamicin-treated culture with an antibiotic
concentration of 50 µg/mL. The number of biological replicates is n=4 for all panels. Biphasic kill
curves were generated using a non-linear model (see **Materials and Methods**). Statistical
significance tests were conducted using F-statistics (*P<0.05, **P<0.01, ****P<0.0001).
The data for each time point represent the mean value $\pm$ standard deviation.

Figure 2: The effect of Crp/cAMP on Persister Cell Metabolism during Stationary Phase. (A)
MS analysis of *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* at early (t=5 h) and late (t=24 h)
stationary phases. Unsupervised hierarchical clustering was applied to standardized metabolic
data. Each column represents a biological replicate. n=4. **(B)** Pathway enrichment analysis was
conducted using MetaboAnalyst (Lu et al., 2023). Upregulated and downregulated pathways of
the Δ*crp* strain compared to WT in the late stationary growth phase were provided in this
figure. **(C, D)** Pathway enrichment maps comparing metabolites of the TCA cycle, pentose

phosphate metabolism, glycolysis, gluconeogenesis, and pyruvate metabolism in Δ*crp* versus
WT for early and late stationary phase conditions, respectively. Circle size corresponds to the
ratio of normalized metabolite intensities between mutant and control cells. Blue (P ≤ 0.05 for
dark blue; 0.05 < P < 0.10 for light blue) and red (P ≤ 0.05 for dark red; 0.05 < P < 0.10 for
light red) indicate significantly downregulated or upregulated metabolites in the mutant
compared to the control. White signifies no significant difference. n=4 for all panels. ESP: Early
stationary phase, LSP: Late stationary phase.

Figure 3: Validation of Crp/cAMP-Mediated Metabolic State in Persister Cells Through
Proteomics Analysis. Pathway enrichment analysis was conducted in STRING (Szklarczyk et al.,
2021, 2023) for upregulated **(panel A)** and downregulated **(panel B)** proteins. Genes highlighted
in red are linked with the upregulated protein networks, while genes in blue, gray, and purple
correspond to those in the downregulated protein network. The visual network in STRING
illustrates protein interactions. In evidence mode, color in the network represents the
interaction evidence of data support, derived from curated databases, experimental data, gene
neighborhood, gene fusions, co-occurrence, co-expression, protein homology, and text mining
(Szklarczyk et al., 2021, 2023).

Figure 4: The Role of Crp/cAMP in Non-Growing Cell Formation. (A, B) Flow cytometry
histograms depict mCherry expression in *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* at early (t=5
h) and late (t=24 h) stationary phases, respectively. Cells containing an IPTG-inducible mCherry
expression system were cultivated with IPTG. After washing and dilution of early and late
stationary phase cells in IPTG-free fresh media, fluorescence was tracked in non-growing and
growing cells for 2.5 hours. The panel is a representative biological replicate. Consistent results
were seen across all 3 biological replicates. **(C)** Growth curves of WT, Δ*crp*, and Δ*cyaA* cultures
were determined using flow cytometry to calculate lag and doubling times. Lag times were
calculated using the "Microbial lag phase duration calculator" (Opalek et al., 2022). Doubling
times were computed using the formula $t_d = \Delta t/(3.3 \times \mathrm{Log}_{10}(N/N_o))$. n=3. *Statistical significance
observed between control and mutant strains (P < 0.05, 2-tailed t-test). The data for each time
point represent the mean value ± standard deviation.

Figure 5: Crp/cAMP-Mediated Metabolic State of Persister Cells. (A) GFP reporter plasmid
introduced into *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* cells to monitor SQR gene activity.
Flow cytometry was used to detect activity at early (t=5 h) and late (t=24 h) stationary phases.
The panel on the left represents a biological replicate, and the results are consistent across all 3
replicates, as demonstrated in the panel on the right. Statistical significance observed between
control and mutant groups (*P < 0.05, **P < 0.01, ***P < 0.001, 2-tailed t-test). **(B)** Redox
activities of *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* cells were measured at early (t=5 h) and
late (t=24 h) stationary phases by flow cytometry using a RSG dye. This dye fluoresces green
after reduction by bacterial reductases. A representative biological replicate is shown (left),
with consistent results across all 5 replicates (right). Statistical significance observed between
control and mutant groups (*P < 0.05, **P < 0.01, 2-tailed t-test). **(C)** *E. coli* cells with integrated
mCherry expression system used to validate cellular respiration. Cells were diluted into fresh
media and treated with ampicillin (200 µg/mL) for 20 hours. Flow cytometry measured the red
fluorescence of intact surviving cells. A representative biological replicate is shown, with
consistent results across all 3 replicates. **(D)** RSG levels of cells (carrying the mCherry expression

system) at exponential phase (t=3 h); cells before ampicillin treatment; non-lysed (intact) cells
after 20 hours of ampicillin treatment; and untreated cells after 20 hours of culturing. A
representative biological replicate is shown (left), with consistent results across all 4 replicates
(right). Statistical significance observed between intact antibiotic-treated cells and others (*P <
0.05, **P < 0.01, 2-tailed t-test). **(E)** High-throughput screening of mutants from the Keio
collection. The mutant strains selected are associated with central metabolism. Stationary
phase cells were diluted 100-fold in fresh medium and treated with ampicillin (200 µg/mL) or
ofloxacin (5 µg/mL) for 20 hours. Treated cultures were washed, serially diluted, and plated on
agar plates to quantify CFUs. **(F)** Genes related to the TCA cycle, ETC, ATP synthase, glycolysis,
and pentose phosphate pathway (PPP) were knocked out and then treated with ampicillin (200
µg/mL) or ofloxacin (5 µg/mL) to enumerate CFUs. n=4. Biphasic kill curves were generated
using a non-linear model. Statistical significance tests were conducted using F-statistics
(*P < 0.05, and **P < 0.01). Each data point represents the mean value ± standard
deviation.

1281

Supplementary Figure Legends

Figure 1–figure supplement 1. cAMP concentrations normalized to cell numbers for _E. coli_ K-
12 MG1655 WT, Δ_crp_, and Δ_cyaA_. The cAMP levels were measured in late stationary phase
cultures at 450 nm using the Cyclic AMP XP® Assay Kit (Cell Signaling Technology). n=4.
Statistical significance was observed between control and mutant strains (*P < 0.05, ***P <
0.001, One-way ANOVA with Dunnett's multiple comparisons test). The data for each time
point represent the mean value ± standard deviation.

Figure 1–figure supplement 2. Genetic perturbation of Crp/cAMP enhanced P$_{cyaA}$ promoter
activity, resulting in increased _gfp_ expression. Overnight cultures of _E. coli_ K-12 MG1655 WT,
Δ_crp_, and Δ_cyaA_ strains harboring the pMSs201 plasmid, which encodes green fluorescent
protein (GFP) under the control of the P$_{cyaA}$ promoter, were diluted 1:1000 into fresh LB
medium and incubated at 37°C with shaking at 250 rpm for 24 hours. Cells at the late stationary
phase were then collected, diluted in 1X PBS, and analyzed by flow cytometry. n=4. Statistical
significance was observed between control and mutant strains (****P < 0.0001, One-way
ANOVA with Dunnett's multiple comparisons test). The data for each time point represent the
mean value ± standard deviation.

Figure 1–figure supplement 3. Growth curves of _E. coli_ K-12 MG1655 WT, Δ_crp_, and Δ_cyaA_.
Optical densities of cell cultures at 600 nm (OD$_{600}$) were measured every hour using a plate
reader. n=3. The data for each time point represent the mean value ± standard deviation.

Figure 1–figure supplement 4. Normalized cAMP concentrations of _E. coli_ K-12 MG1655 WT.
The cAMP concentrations were measured in growth cultures at the indicated time points. First,
the cAMP concentrations were normalized to the number of cells. Subsequently, the data were
further normalized based on the time point 0 to mitigate errors associated with batch-to-batch
assay kit variations and to capture the trend in cAMP levels across the time points. n=8.
Statistical significance was observed between time points (**P < 0.01, One-way ANOVA with
Dunnett's multiple comparisons test). The data for each time point represent the mean

value ± standard error.

Figure 1–figure supplement 5. Persister levels of cells carrying the Crp expression system.
Cells at early (t=5 h) and late (t=24 h) stationary phases were transferred to fresh media with
antibiotics for persister cell quantification. At time points 0, 2, 6, 8, and 20 h, 1 mL of the
treated culture underwent two washes with 1X PBS to remove antibiotics. It was then serially
diluted and plated on an agar plate to count the CFUs. **(A)** Persister levels of ampicillin-treated
culture (200 μg/mL). **(B)** Persister levels of ofloxacin-treated cultures (5 μg/mL). **(C)** Persister
levels of gentamicin-treated culture (50 μg/mL). n=4. Biphasic kill curves were generated using
a non-linear model (see **Materials and Methods**). Statistical significance tests were conducted
using F-statistics (**P < 0.01 and ****P < 0.0001). The data for each time point represent the
mean value ± standard deviation.

Figure 1–figure supplement 6. Persister levels of *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA*
strains at normalized antibiotic concentrations. Panels show survival following treatment with
(A) ampicillin, **(B)** ofloxacin, and **(C)** gentamicin. After treatment, samples were washed six
times with 1X PBS to minimize antibiotic carryover. Antibiotic concentrations were normalized
to MICs to ensure valid comparisons across strains (The concentrations of 33× MIC for
ampicillin and 100× MIC for ofloxacin and gentamicin match those used in Figure 1). n=4.
Statistical significance was observed between control and mutant strains (****P < 0.0001, One-
way ANOVA with Dunnett's multiple comparisons test). The data for each time point represent
the mean value ± standard deviation.

Figure 1–figure supplement 7. Agar plates showing *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA*
strains following treatment with antibiotics at normalized concentrations. After treatment,
cells were washed and subjected to 10-fold serial dilutions, then plated on agar. Plates were
incubated for **(A)** 16 h, **(B)** 48 h, and **(C)** 72 h to assess colony formation over time. The panel
is a representative biological replicate. Consistent results were seen across all 3 biological
replicates.

Figure 1–figure supplement 8. Deletion of *crp* and *cyaA* reduces ampicillin and ofloxacin
persistence in the *hipA7* strain. Persistence levels were assessed by exposing *E. coli* K-12
MG1655 WT, *hipA7*, *hipA7Δcrp*, and *hipA7ΔcyaA* strains in the late stationary phase to the
specified antibiotics, followed by CFU quantification at designated time points. **(A)** Ampicillin
(200 μg/ml), **(B)** Ofloxacin (5 μg/ml) and **(C)** Gentamicin (50 μg/ml). n=4. Statistical significance
was observed between control and mutant strains (****P < 0.0001, One-way ANOVA with
Dunnett's multiple comparisons test). The data for each time point represent the mean
value ± standard deviation.

Figure 2–figure supplement 1. The pathway enrichment analysis comparing WT and Δ*crp*
strains during the early stationary phase cultures. The analysis was conducted using
MetaboAnalyst, with a threshold ratio (Δ*crp*/WT) set at ≤ 0.5 for downregulation and ≥ 2 for
upregulation. **(A)** Upregulated and **(B)** Downregulated pathways in the Δ*crp* strain compared to
WT during the early stationary growth phase (ESP).

Figure 2–figure supplement 2. The pathway enrichment analysis for the WT strain. The
analysis was conducted using MetaboAnalyst, with a threshold ratio (LSP/ESP) set at ≤ 0.5 for

downregulation and ≥ 2 for upregulation. **(A)** Upregulated and **(B)** Downregulated pathways of
the WT strain in the late stationary growth phase (LSP) compared to the WT strain in the early
stationary phase (ESP).

Figure 2–figure supplement 3. The pathway enrichment analysis for the Δ*crp* strain. The
analysis was conducted using MetaboAnalyst, with a threshold ratio (LSP/ESP) set at ≤ 0.5 for
downregulation and ≥ 2 for upregulation. **(A)** Upregulated and **(B)** Downregulated pathways of
the Δ*crp* strain in the late stationary growth phase (LSP) compared to the Δ*crp* strain in the
early stationary phase (ESP).

Figure 3–figure supplement 1. The MS analysis of proteins from both WT and Δ*crp* strains at
the late stationary phase. The proteomic data were subjected to unsupervised hierarchical
clustering. Each column in the figure represents a biological replica. n=3.

Figure 4–figure supplement 1. Persister levels of *E. coli* WT, Δ*crp*, and Δ*cyaA* cells with the
integrated mCherry expression system. Late stationary phase cultures (t=24 h) were
transferred to fresh media and treated with ampicillin (200 µg/mL), ofloxacin (5 µg/mL), and
gentamicin (50 µg/mL) for 20 hours. Subsequently, 1 mL of the treated culture underwent two
washes with 1X PBS to remove antibiotics. It was then serially diluted and plated on an agar
plate to count the CFUs. The levels of ofloxacin and gentamicin persisters in the mutant strains
were below the limit of detection. n=4. Statistical significance was observed between control
and mutant strains (***$P < 0.001$, ****$P < 0.0001$, Two-way ANOVA with Tukey's multiple
comparisons test). The data for each time point represent the mean value ± standard
deviation.

Figure 4–figure supplement 2. Non-growing cell levels in the *E. coli* strain carrying the Crp
expression system. (A, B) Flow cytometry histograms depict mCherry expression in
Δ*crp*+pUA66-*crp*, and Δ*crp*+pUA66 EV at early (t=5 h) and late (t=24 h) stationary phases,
respectively. Cells containing an IPTG-inducible mCherry expression system were cultivated
with IPTG. After washing and dilution of early and late stationary phase cells in IPTG-free fresh
media, fluorescence was tracked in non-growing and growing cells for 2.5 hours. The panel is a
representative biological replicate. Consistent results were seen across all 3 biological
replicates.

Figure 5–figure supplement 1. RSG staining control for bacterial metabolic activities.
Exponential phase (t=3 h) cells were stained with 1 µM RSG for 10 minutes at 37°C before
analyzing by flow cytometry. Unstained cells and cells treated with 20 µM CCCP + 1 µM RSG
were used as control. CCCP was expected to reduce cellular redox activities. The panel is a
representative biological replicate. Consistent results were seen across all 3 biological
replicates.

Figure 5–figure supplement 2. Intact (non-lysed) cell levels of *E. coli* WT, Δ*crp*, and Δ*cyaA* cells
with the integrated mCherry expression system. mCherry positive cells were diluted into fresh
media and treated with ampicillin (200 µg/mL) for 20 hours. Flow cytometry was used to
measure the red fluorescence of intact surviving cells at time points t=0, 1, 2, 4, 6 and 20 h. A
representative biological replicate is shown, with consistent results across all 3 replicates.

Figure 5–figure supplement 3. VBNC levels of *E. coli* WT, Δ*crp*, and Δ*cyaA* cells with the

integrated mCherry expression system. Flow cytometry was employed to quantify intact
surviving cells. Persister cells were quantified by plating the cells on agar media. Viable but
nonculturable (VBNC) cells were enumerated by subtracting persister levels from the intact cell
levels. n=4. Statistical significance was observed between control and mutant strains (****P <
0.0001, One-way ANOVA using Dunnett's multiple comparisons test). The data for each time
point represent the mean value ± standard deviation.

Figure 5–figure supplement 4. Cell counts of *E. coli* K-12 MG1655 WT and mutant strains using
flow cytometry at late stationary phase. Cells were diluted 100-fold into 1 mL of 1X PBS. n=4.
The data for each time point represent the mean value ± standard deviation.
1399
List of Supplementary Files

Supplementary File 1. MIC of antibiotics and concentrations of bactericidal antibiotics used in
persister assays.
Supplementary File 2. Analyzed metabolomics data.
Supplementary File 3. Analyzed proteomics data.
Supplementary File 4. Analyzed persister survival fraction data.
Supplemental File 5. The knockout strains generated using the *E. coli* K-12 MG1655
background in this study.
Supplemental File 6. Persister levels of *E. coli* K-12 MG1655 WT, Δ*crp*, and Δ*cyaA* strains in
late stationary phase.
Supplemental File 7. Persister levels of *E. coli* K-12 MG1655 (A) and BW25113 (B) WT, Δ*crp*,
and Δ*cyaA* strains in the exponential growth phase.
1412

List of Source Data

Figure 2 - Source Data 1: Normalized Metabolomics Data of Early and Late Stationary Phases
of Wild-Type and Mutant *crp* Across 3 Biological Replicates. Data Collected and Analyzed by
Metabolon Inc. (Morrisville, NC).
Figure 3 - Source Data 1: Normalized Proteomics Data of the Late Stationary Phase of Wild-
Type and Mutant *crp* Across 3 Biological Replicates. Data Collected by UT Health's Clinical and
Translational Proteomics Service Center (Houston, TX).
1420

A

B Upregulated pathways

C Early stationary phase

D Late stationary phase

Upregulated (red circle) and downregulated (blue circle) metabolites in the Δ*crp* strain compared to WT

A **Upregulated pathways**

- Anabolic metabolism
- Peptidoglycan metabolic processes
- Cell wall organization or biogenesis
- Cellular component organization or biogenesis
- Regulation of cell shape, cell cycle, and cell division

B **Downregulated pathways**

- Ribosome and translation factor activity
- Translation, and protein export
- Ribonucleoprotein, and translation factor activity
- RNA binding

- Tricarboxylic acid cycle
- Glycerol metabolism
- Pyruvate metabolism
- Glycolysis
- Energy metabolism
- Respiratory processes

- Periplasmic transporters
- Sugar transport
- Periplasm

A Early stationary phase

WT — Δcrp — ΔcyaA

t=0h, t=1h, t=2h, t=2.5h

Normalized cell count vs Red fluorescence

B Late stationary phase

WT — Δcrp — ΔcyaA

t=0h, t=1h, t=2h, t=2.5h

Normalized cell count vs Red fluorescence

C

WT — $t_d = 33\pm4$min, 51 ± 3min

Δcrp — $t_d = 72\pm16$min *, 32 ± 9min *

ΔcyaA — $t_d = 75\pm16$min *, 29 ± 9min *

Log_{10}(cells/mL) vs Time (h)