Immunity and bacterial recruitment in plant leaves are parallel processes whose link shapes sensitivity to temperature stress

Jisna Jose[1,9], Erik Teutloff[1], Simrat Naseem[1], Emanuel Barth[2,4], Rayko Halitschke[3,9], Manja Marz[2,5,6,7,8,9], and Matthew T. Agler[1,9*]

[1] Institute for Microbiology, Plant Microbiosis Group, Friedrich Schiller University Jena, Neugasse 23, Jena 07743, Germany.

[2] RNA Bioinformatics and High-throughput analysis, Faculty of Mathematics and Computer Science, Friedrich Schiller University Jena, Jena, Germany.

[3] Department of Biochemistry, Max-Planck Institute for Chemical Ecology, Hans-Knöll-Straße 8, Jena 07745, Germany.

[4] Bioinformatics Core Facility Jena, Friedrich Schiller University Jena, Leutragraben 1, 07743 Jena, Germany

[5] Leibniz Institute for Age Research, FLI, Beutenbergstraße 11, Jena 07743, Germany

[6] European Virus Bioinformatics Center, Friedrich Schiller University, Leutragraben 1, Jena 07743, Germany

[7] German Center for Integrative Biodiversity Research (iDiv), Puschstraße 4, Leipzig 04103, Germany

[8] Michael Stifel Center Jena, Friedrich Schiller University, Ernst-Abbe-Platz 2, Jena 07743, Germany

[9] Cluster of Excellence Balance of the Microverse, Friedrich Schiller University, Furstengraben 1, Jena 07743, Germany

*Corresponding author:

Matthew T. Agler

The Plant Microbiosis Lab, Institute of Microbiology

Friedrich Schiller University Jena

Neugasse 23

07743 Jena, Germany

E-mail: matthew.agler@uni-jena.de

Tel: +49 (0)3641 9 49980

Abstract

Rising global temperatures necessitate developing climate-resilient crops with better adaptability to
changing climates. Under elevated temperatures, plant immunity is downregulated, putting them at risk
of foliar pathogen attack. Manipulating plant defense hormones is one way to mitigate this detrimental
effect. However, it is unclear how plant immunity interacts with plant microbiome assembly and how
temperature will thus affect overall plant health and stability. We used chemical mutagenesis to identify
a phenotypically healthy genotype of *A. thaliana, "CLLF",* that compared to the wild type naturally
recruits an altered leaf bacteriome, including unusually high bacteria loads. Simultaneously, CLLF
hyperaccumulates salicylic acid (SA) and jasmonates, has constitutively upregulated systemic and
innate defenses, and has increased resistance to necrotrophic fungal and hemi-biotrophic bacterial
pathogens, indicating that pathogen immunity and non-pathogen recruitment function in parallel.
Growth of specific non-pathogenic leaf bacteria on SA as a carbon source suggests the same hormones
may even link the two processes. CLLF also showed high tolerance to heat stress in comparison to the
wild type, but SA-associated defense processes are not downregulated under heat. Synthetic community
(SynCom) experiments showed that when the taxonomic diversity of bacteria available to CLLF is
artificially reduced, resilience to heat stress is compromised, leading to dysbiosis, but this does not
occur with the full SynCom or in the wild type with any SynCom. Thus, the downregulation of defenses
in response to heat may contribute to avoidance of dysbiosis caused by some leaf bacteria, but full
bacteriome taxonomic diversity can restore balance.

Significance Statement

Plants are living ecosystems colonized by diverse microorganisms who strongly shape host health.
Understanding how balance arises in host-associated microbiomes is a key step to understanding how
to preserve, manage and possibly optimize these complex ecosystems, especially in a changing climate.
Using a random mutagenesis approach in a natural *A. thaliana* ecotype, we find that constitutively
upregulated defenses are associated with both tolerance to (a)biotic threats and healthy recruitment of
leaf bacteria, very likely in a direct manner. Thus, immunity and bacterial recruitment in leaves operate
in parallel. Synthetic community experiments show further that this link plays important roles in
shaping plant resilience to heat stress, an important consideration in developing plants more stable to
climate change.

 <u>Introduction</u>

Extreme conditions such as high temperature, drought, salinity, and high humidity due to climate
change may adversely affect agriculture and crop productivity. One reason is that changes in ambient
growth conditions that affect plant physiology and development, in turn, can alter plant-microbe
interactions, an inter-dependence known as the "disease triangle" (1). Altered plant-microbe
interactions carry the risk that balance in the microbiome may be lost, leading to dysbiosis, including
devastating plant diseases and loss of productivity. Thus, it is important to understand how plants
establish and maintain balance in the microbiome and how environmental factors will affect that
balance.

Leaf-colonizing bacteria are an important component of the microbiome with regard to plant health
because they protect plants but also must be regulated to prevent disease (2). Plants use a variety of
mechanisms to recruit bacteria and maintain balance. The phyllosphere (leaf) bacteriome is assembled
from both soil-born and air-born inoculums in either a deterministic or stochastic manner (3, 4). Plant
defenses, besides establishing immunity to pathogens, also plays critical roles in maintaining balance
in the bacteriome. For one, the interaction of commensal and opportunistic pathogenic bacteria with
immune components like MAMP-triggered immunity helps prevent dysbiosis by preventing
overgrowth of opportunistic pathogens (5–7). Additionally, secondary metabolites can contribute to
assembly via their toxicity (8) and likely by serving as resources for bacteria particularly adapted to use
them (9, 10). However, how plant immunity against (opportunistic) pathogens is more broadly linked
to the parallel process of bacteriome assembly in healthy leaves remains unclear.

The plant immune system is itself affected by the abiotic environment. Although the mechanisms
linking abiotic stressors to plant immunity are still underexplored, the effect of temperature is perhaps
best studied. For example, at moderately high temperatures (30°C vs. 23°C), the production of salicylic
acid (SA), a major plant defense hormone, is suppressed (11). Jasmonic acid (JA) biosynthesis is also
involved in temperature-dependent susceptibility of rice blast disease (12). Abscisic acid (ABA)
biosynthesis was not affected by heat stress, but ABA does negatively affect R gene-mediated immune
responses at high temperatures (13). Because of compromised immunity, these changes lead to
increased susceptibility to pathogens. Some microbes help plants mitigate heat stress by making them
more resilient (14, 15), suggesting that the leaf microbiome could offset some adverse effects of
increased temperature.

We hypothesized that the balance established by plant immune components during the natural
recruitment of leaf bacteriomes plays important roles in resilience to temperature stress. To investigate
these questions, we first studied how plant physiology shapes balance in the leaf bacteriome of a wild
A. thaliana genotype by generating a random mutant genotype that is healthy, but which naturally
assembles an alternatively balanced leaf bacteriome. We then studied immune signaling in the mutant

and responses to pathogens and temperature stress. Finally, we tested how the balanced bacteriome

compensates for altered immune responses to temperature.

A phenotypically distinct mutant displays an altered leaf bacteriome

Fig.1. A mutant *A. thaliana* NG2 genotype recruits a distinctive leaf bacteriome. A) Representative figures showing the phenotype of mutant created in *the A. thaliana* NG2 background "CLLF" (Curled and compressed Leaf with Late Flowering). B) Bacterial loads of CLLF and NG2 Wt were measured as colony-forming units (CFU) per gram of fresh leaf weight (n =12). C) Bacteria to host ratio of CLLF and NG2 Wt calculated using hamPCR amplicon sequencing data, the fraction of the total number of bacterial 16S rRNA gene reads to the sum of bacterial 16S rRNA gene and *A. thaliana* GI (*GIGANTEA*) reads per sample (n=12). D) Alpha diversity measures of CLLF and NG2 Wt leaf bacteriome based on 16S rRNA gene amplicon sequencing data grouped at the genus level. E) CCA plot based on the Bray-Curtis distance between samples constrained for genotype, generated using 16S rRNA gene amplicon sequencing data grouped at the genus level. 15.4% variance is explained by genotype (PERMANOVA with 999 permutations) p-value < 0.01. (CLLF; purple, NG2_Wt; green, ns; not significant, * p<0.05, ** p<0.01, *** p<0.001).

We previously established a collection of *Arabidopsis thaliana* ecotypes collected from Jena, Germany

(16). For this study, we generated a mutant plant library in ecotype NG2 using chemical mutagenesis

(see methods). In the M2 generation, we screened hundreds of mutant plants for higher bacterial load

than the wild type (WT). Approximately 5% of the mutant population had a higher bacterial load

(although this was only based on n=1 because every seed is a unique genotype), indicating altered

microbiota recruitment (Supplementary Fig.1). The mutant used further in this study, "CLLF", had the

highest bacterial load. In addition to that, it had a distinct phenotype, curled, and compressed leaves

with late flowering compared to the wild type (Fig 1A).

Whole leaf bacteriomes were characterized in the NG2 wild type (NG2 Wt), and the M4 generation of

CLLF (which should be mostly homozygous) grown in commercial soil supplemented with an extract

of natural soil. A plastic barrier was used to separate rosettes from soil (see methods) to ensure that

colonization differences were not due to increased CLLF leaf-soil direct contact. In 3-week-old plants,
bacterial loads in leaves of CLLF (colony-forming units) were approximately 100-fold higher compared
to the wild type. We also confirmed higher loads using the sensitive hamPCR method (17) (Fig.1B and
1C, $p<0.01$ and $p<0.05$, respectively). In addition, CLLF leaves recruited more diverse bacterial genera
than the wild type (Fig.1D, Chao1 $p < 0.01$) with a similar evenness (Fig. 1D, Shannon, and Simpson
indices). Furthermore, CLLF leaf bacteriome samples were more densely clustered and were separated
from NG2 wildtype in canonical correspondence analysis (CCA) of the Bray-Curtis distances between
samples (Fig.1E, 15.4% variation explained by the genotype, PERMANOVA, $p = 0.008$, 999
permutations). The distinct beta diversity patterns, higher alpha diversity, and high bacterial loads
together confirm an altered leaf microbiota recruitment in CLLF. Similar results were found without
the plastic barrier in three independent experiments (Supplemental Fig.2).

CLLF's altered recruitment is not dysbiotic

Fig.2. CLLF's microbiome is not dysbiotic and it has normal survival rates. A) Relative abundance of the top 12 Families (remaining families represented as "Others") in CLLF and NG2_Wt. Asterisks denote significant differences in abundance compared to the NG2 Wt. B) Differentially abundant bacterial families associated with genotypes. All families shown have a p-adjusted value <0.01 and data was normalized to plant-specific GI gene. C) Percentage of plants that survived from November 2022 to March 2023 in the garden (n=7). (CLLF; purple, NG2_Wt; green, ns; not significant, * p<0.05, ** p<0.01, *** p<0.001).

To check whether the healthy CLLF phenotype indicated a more taxonomically balanced bacteriome
than previously described "dysbiotic" mutants (over-colonization by one or a few opportunistic
pathogens), we looked closer at the taxonomic composition of the CLLF leaf bacteriome. Overall,
CLLF recruited similar bacterial families as the wild type (Fig. 2A). However, multiple families were
significantly more abundant in CLLF (p-value ≤ 0.01) (Fig. 2B). Differentially abundant taxa that were
significantly correlated to the genotype included members of most leaf-colonizing phyla. One-third of

the differentially abundant taxa were Betaproteobacteria (order Burkholderiales, families
Alcaligenaceae, Burkholderiaceae, Comamonadaceae, and Methylophilaceae), one-fourth of the
differentially abundant taxa were Alphaproteobacteria (Caulobacteraceae, Rhizobiaceae, and
Rhodobacteraceae), two families each belonging to Actinomycetota (Microbacteriaceae and
Pseudonocardiaceae) and Gammaproteobacteria (Moraxellaceae and Xathomonadaceae) and one
family belonging to Bacteroides (Spirosomaceae). Thus, the higher bacterial load of CLLF is not due
to the enrichment of a single or a few taxa, which should lead to dysbiosis, but rather an apparently
balanced enrichment of multiple families.

In support of balance in the CLLF bacteriome, plants were never observed to develop a disease
phenotype in the lab (Fig.1A). To test this further, we assessed CLLF's survival during overwintering
in nature in an outdoor garden experiment. CLLF had a similar survival rate as wild-type NG2 plants
(Fig.2C). Furthermore, leaf functions of CLLF were only slightly altered compared to wild type.
Specifically, the electron transport rate (ETR) and the vapor pressure differential were slightly higher,
and the stomatal conductance was slightly lower in CLLF compared to the wild type (p-value < 0.01)
(Supplementary Fig. 3). Together, these results suggest that the mutant phenotype of CLLF leads to the
recruitment of an alternate, but balanced, bacteriome.

Fig.3. CLLF shows upregulated defense hormone profiles and has a constitutively active immune system. A) Jasmonic acid (JA) biosynthesis precursors and derivatives measured in nanograms per fresh weight of leaf tissue (n=8). A schematic representation of the JA biosynthesis pathway is shown next to it (Highlighted in color are each precursor/derivative). B) Total salicylic acid (SA) levels in CLLF and NG2_Wt measured in nanograms per fresh weight of leaf tissue (n=8). C) Total abscisic acid (ABA) levels in CLLF and NG2_Wt measured in nanograms per fresh weight of leaf tissue (n=8) (CLLF; purple, NG2_Wt; green, ns; not significant, * p<0.05, ** p<0.01, *** p<0.001). D) Gene ontology (GO) enrichment analysis of differentially expressed genes in gnotobiotic CLLF compared to the NG2 Wt. The cut-off for the genes was set at log fold change > [2], base mean > 100, and p-value < 0.01.

To test whether the CLLF immune system is intact, we investigated defense hormone levels. Notably,

CLLF exhibited constitutively higher levels of defense hormones, with upregulated salicylic acid (SA)

(Fig.3B) and jasmonates (Fig.3A). CLLF plants grown in axenic conditions also showed higher levels

of SA and jasmonates (Supplementary Fig.4), indicating that the constitutive production of these

hormones is microbe-independent.

Next, we analyzed CLLF gene expression relative to NG2 Wt plants. In agreement with the

overproduction of defense hormones, gene expression profiles underscored CLLF's constitutively

active immune system. Genes involved in response to SA, systemic acquired resistance (SAR),

responses to bacterial molecules, and oomycetes were among the highest upregulated in CLLF (Fig.3D).

The strongest downregulated genes were mostly involved in glucosinolate-related processes.

We reasoned that an overactive immune system may still be dysfunctional, which could explain the

higher commensal bacterial loads. However, CLLF plants were more resistant to the bacterial pathogen

Pseudomonas syringae pv. syringae DC3000 (Supplementary Fig. 5C) and showed higher resilience

towards the fungal pathogen *Sclerotinia sclerotiorum* (Supplementary Fig. 5A and 5B) than the wild

type. Thus, CLLF appears able to mount a functional immune response. Therefore, the role of plant

defenses in maintaining plant immunity works in parallel with the recruitment of non-pathogenic leaf

bacterial communities.

Some leaf bacteria may directly utilize hormones as a resource for growth

Fig.4. Growth rate of representative strains from various phyla in minimal media supplemented with salicylic acid (SA) as carbon source. Control is the growth rate in minimal media without any carbon source, colored yellow. PC (positive control) is the growth rate in minimal media supplemented with sucrose and fructose, (11 mM sucrose + 11 mM glucose), colored green. SA is the growth rate in minimal media supplemented with salicylic acid (0.5 mM Salicylic acid) as a carbon source, colored purple. The growth rate is measured as optical density at 600nm (n=4).

High SA levels were previously linked to increased bacterial diversity in the phyllosphere (18), and SA

has been suggested to recruit specific taxa directly in the rhizosphere (9). Thus, we hypothesized that

plant hormones may simultaneously shape plant immunity and recruit leaf bacteriomes, explaining the

seemingly counterintuitive resistance and recruitment phenotypes of CLLF. To test this, we checked

whether strains representative of diverse CLLF phyla could utilize salicylic acid as a carbon source (Fig

4). We found that some bacteria belonging to Alphaproteobacteria (Rhizobium and Methylorubrum)

and Actinomycetes (Nocardioides and Brevibacterium) showed growth in minimal media supplemented

with 0.5mM salicylic acid. However, the Burkholderiales and Bacteroides we tested did not show

growth in SA *in vitro*. Thus, overproduction of SA and possibly other hormones in the phyllosphere at

least have the potential to contribute directly to the recruitment of specific bacterial taxa that can

metabolize them, as previously suggested.

High defense-related gene expression in CLLF does not decrease at elevated temperatures

A well-known effect of moderately high temperatures in plants is decreased expression of defense-

related genes, which causes increased pathogen susceptibility. During an accidental breakdown of plant

growth chambers in which temperatures of 34°C were reached, we observed that CLLF showed higher

tolerance to the extreme temperature than the wild type (Supplementary Fig. 5D), which could be caused

by increased SA signaling (19). To better understand this, we looked at what genes were differentially

regulated in CLLF with and without a controlled high temperature (30°C) treatment relative to NG2 Wt

plants (Supplementary Fig. 6). Overall, the gene expression profile of CLLF did not strongly change

due to heat treatment (Supplementary Fig. 6A). Specifically, very few genes and associated biological

processes were up- or down-regulated only in response heat treatment (Supplementary Fig. 6B). In

contrast, almost all processes differentially regulated in CLLF relative to NG2 Wt in normal conditions

were similarly regulated after heat treatment (Supplementary Fig. 6C). This included immune

activation-related processes, although a downregulation would typically be expected at elevated

temperatures (11). Thus, the constitutively active immune system of CLLF also does not suffer from

down-regulation of immune signaling in response to high temperature.

The balanced CLLF bacteriome is required to prevent dysbiosis at high temperatures

Because CLLF's activated immune system was linked to the alternative bacteriome balance and

downregulation of immune responses at high temperatures was inactive in CLLF, we decided to

investigate the role of bacteriome composition on balance during temperature stress. Therefore, we

used a semi-gnotobiotic synthetic community (SynCom) experiment (see methods). Plants were pre-

inoculated with either a full SynCom representative of the entire CLLF bacteriome (SynCom1) or

SynComs that lacked major taxonomic groups. This setup allowed us to evaluate whether significant

losses in taxonomic diversity would affect balance. SynCom2 lacked Burkholderiales

(Comamonadaceae and Oxalobacteriaceae), SynCom3 lacked Bacteroidota (Flavobacteriaceae and

Sphingobacteriaceae), and SynCom4 lacked both groups (Supplementary Fig 7 and Supplementary

Table 1). Notably, the bacteria remaining in SynCom4 include Actinomycetes and Alphaproteobacteria

that can utilize salicylic acid as a carbon source (Fig 4).

Plants not exposed to heat displayed normal, healthy phenotypes with all SynComs (Fig 5A and

Supplementary Fig. 8A). We next evaluated gene expression profiles of NG2 Wt and CLLF treated

with SynCom1 and the reduced SynCom4. As expected, compared to NG2 Wt with SynCom1, CLLF

with SynCom1 showed upregulation of various immune responses (Fig 5B), similar to the response

after growth in normal soil (Fig 3D). In response to SynCom4, NG2 Wt showed significantly increased

immune responses, phytoalexin biosynthetic pathways, and decreased photosynthesis-related genes,
while CLLF maintained high defense-related gene expression (Fig 5B and 5C). Thus, removing
Burkholderiales and Bacteroides in SynCom4 appeared to result in a disbalance that NG2 Wt addressed
with a defense response. CLLF was likely able to maintain balance due to already upregulated immune
responses.
Upon exposure to 30°C for 48-h, NG2 Wt plants showed a normal, healthy phenotype regardless of the
SynCom (Fig 5 and Supplementary Fig. 8B). Immune system-related biological processes that were
upregulated in NG2 Wt plants with SynCom4 under normal conditions decreased expression following
heat treatment. Thus, the NG2 Wt regulated immune genes as previously described due to heat, and this
appears to have been the appropriate response even with a highly reduced bacterial community, since
the plants remained healthy. In CLLF with SynCom4, on the other hand, we observed a strong effect of
heat stress (Fig. 5A and Supplementary Fig. 8C). Symptoms included deformed rosettes and chlorosis.
In these plants, immune responses, genes related to senescence, and low oxygen were extremely highly
upregulated, suggesting a potential growth-defense tradeoff. CLLF with SynCom1, on the other hand,
showed no signs of a dysbiotic phenotype under heat treatment (Supplementary Fig.8 and 9). Together,
these results suggest that with an incomplete bacteriome, proper immune system regulation in response
to temperature helps avoid dysbiosis. However, a complete bacteriome counteracted dysbiosis,
suggesting that leaf microbiome diversity can strongly contribute to host stability under abiotic stress
and contribute resilience to overcome altered immune signaling regulation. Notably, CLLF always
showed higher resistance than NG2 Wt to *Pseudomonas syringae pv. syringae* DC3000 (Pst DC3000),
even when inoculated with SynCom4 and after heat treatment (Supplementary Fig. 10). This further
supports that the mechanism of the observed growth-defense imbalance under temperature is linked to
microbiome assembly and is distinct and functions in parallel to the mechanisms of pathogen resistance.

Fig.5. Phenotype of SynCom inoculated plants with or without heat stress; CLLF develops dysbiosis at higher temperatures with a taxonomically reduced inoculum. A) Phenotype of SynCom4-inoculated plants at normal temperature (18/22°C) and after heat stress (treated at 30°C for 48h). Light green border- NG2Wt inoculated with SynCom4 at normal temperature, Yellow- NG2Wt inoculated with SynCom4 after heat stress, Pink- CLLF inoculated with SynCom4 at normal temperature, Blue- CLLF inoculated with SynCom4 after heat stress. **(**B. and C.) All up- (B.) or downregulated (C.) biological processes (BPs) in CLLF plants with SynCom1 (Diverse community) at normal temperature (Purple), NG2Wt and CLLF plants inoculated with SynCom4 at normal temperature (light green and pink, respectively), and NG2Wt and CLLF plants inoculated with SynCom4 with heat stress (yellow and blue, respectively), all in comparison to NG2Wt inoculated with SynCom1 at normal temperatures. All genes considered for assigning biological processes are subsetted based on p </= 0.01, basemean > 100, and foldchange > [2]. richFactor is the ratio of differentially expressed genes annotated to the term to the ratio of background genes annotated to the term. Data shown here are the top 10 commonly upregulated and downregulated biological processes, all the remaining BPs are added as a supplementary figure (Supplementary Fig. 9).

Discussion

The absence of key components of the immune system was previously shown to lead to dysbiosis - a breakdown of a healthy plant microbiome (5, 7). In these cases, disease phenotypes were characterized by high growth of specific taxa, suggesting that plant immune systems maintain balance by preventing the proliferation of opportunistic pathogens (2). Here, we identified CLLF, a mutant genotype with unusually high leaf bacterial loads when grown in natural soil. Detailed phenotyping revealed that the mutant has a constitutively active and apparently fully functional immune system despite high loads. Compared to the previously reported dysbiotic bacteriomes in immune-deficient mutants (5, 7), CLLF was healthy, robust, had a high abundance of diverse leaf-associated bacterial taxa, and no overgrowth of specific opportunistic pathogenic taxa. Thus, the constitutively active immune system of CLLF and associated high bacterial loads appear to represent an alternative balanced state of the leaf bacteriome. This finding suggests that although high bacterial loads indicate disease when opportunistic pathogens proliferate (20), balanced recruitment can stabilize higher loads. The findings also show that increased recruitment of non-pathogenic bacteria is compatible with and may even be directly linked to a fully functional and active immune system.

The observation that constitutively activated immunity was strongly correlated to leaf bacterial recruitment in CLLF could partly be explained by the ability of bacteria to use plant defense signals directly as resources for growth. It is already well-known that some bacteria can grow on SA directly (21) and SA was previously suggested to play positive direct roles in rhizosphere recruitment (9) and in promoting leaf bacterial diversity (18). SA is also likely available to leaf colonizers since it accumulates in extracellular compartments (e.g., apoplast) upon activation of immunity (22). In our tests, the growth of bacterial taxa used in the SynCom experiments on SA followed taxonomic boundaries: None of the Burkholderiales or Bacteroides tested showed growth in SA, while the tested alphaproteobacteria and actinobacteria did (SA consumers). Thus, a role for SA in the recruitment of these taxa is plausible.

Apart from SA, the CLLF mutant also accumulated jasmonates, including OPDA and JA, relative to NG2 Wt. Although this did not include the best-studied active form, JA-isoleucine, molecules such as OPDA play important roles in defense, thermotolerance, and regulation of other plant hormones (23, 24). JA species have also been shown to have effects on activity of specific plant-associated bacteria, including functioning as a chemoattractant and inducing the formation of biofilms in the rhizosphere (25, 26). Additionally, the aliphatic glucosinolate profile of CLLF was significantly altered compared to the wild type (data not shown here). Some bacteria can metabolize glucosinolates, and accordingly, we previously found that leaf glucosinolates function in bacterial recruitment, likely by acting as a carbon source for specific taxa (10, 27). This recruitment depends on the glucosinolate species, so the changed glucosinolate chemotype in CLLF likely also contributed to the altered leaf microbiome. Thus,

we hypothesize that the secondary metabolites linked to the plant immune system also function in direct

and specific recruitment processes for non-pathogenic leaf bacteria.

Alternative roles of plant hormones are not surprising since they are ancient signals in the green lineage

and, therefore, likely serve various purposes. SA-responsive NPR proteins, for example, appear to be

SA receptors in all plants, but conserved ancient functions shared by both bryophytes and tracheophytes

include light and temperature sensing, but not defense which probably evolved later (28). At any rate,

if defense signals help recruit leaf bacteria, forward genetics approaches such as mutant screens used in

this study could be a promising way to engineer plant microbiomes in a targeted way by screening for

changes in specific metabolites. However, to leverage such an approach, further research is needed to

determine how specific metabolites function in different contexts to recruit bacteria and how

microbiome balance is impacted by resource utilization.

The salicylic acid (SA) pathway is integral to plant resistance and immunity, playing major roles in

abiotic and biotic stress tolerance (28, 29). Down-regulation of SA biosynthesis at high temperatures

makes plants prone to infection by plant pathogens (29). Thus, both exogenous SA applications and

manipulating immune regulation to increase SA accumulation under higher temperatures have been

suggested as remedies (19, 30). However, it is unclear how altering the regulation of immunity at higher

temperatures will affect plant interactions with the broader microbiome. Here, the mutant CLLF

genotype showed many expected signs of constitutive and heat-insensitive SA-dependent immune

upregulation: Higher tolerance to plant pathogens under both conditions and higher tolerance to heat

stress than the wild-type. While this did not cause problems at normal temperatures, plants exposed to

a reduced microbiota showed extreme dysbiosis at higher temperatures. One explanation could be that

the upregulation of SA signalling under heat stress may have fuelled the overgrowth of SA consumers

that were not balanced by other colonizers, resulting in a growth-defense imbalance. Accordingly,

CLLF plants pre-inoculated with a fully diverse bacteriome did not show dysbiosis, indicating that a

full bacteriome diversity contributes resilience to altered immune signaling regulation.

In conclusion, our results suggest that alternative states of balance exist in the phyllosphere microbiome.

Specifically, plant immune activation alters the balance in the phyllosphere microbiome by altering

recruitment patterns without necessarily creating dysbiosis. We hypothesize that this occurs because

plant defense hormones, besides acting as defense signals, are also plant-bacterial communication

signals, driving colonization of the non-pathogenic bacteriome. However, as we observed, this implies

that hormone balance will play key roles in maintaining a stable microbiome under environmental

stressors. The microbiome of plants colonized in nature depends on both the environment and chance

colonization events (31), so not all taxa are always present. Given this risk, plants may have evolved to

prevent upregulation of defense signals under heat stress, at least in part, to mitigate the possibility of

imbalance caused by the proliferation of taxa that grow in response to defense signals. Thus, as

manipulating plants to better deal with multiple stressors increasingly finds use to combat changing

climates (32), it is important to consider how altered signaling patterns will affect interactions with the

diversity of the plant-associated bacteriome. In particular, strategies to ensure a full diversity is available

for colonization may help avoid unintended consequences.

 <u>Materials and Methods</u>

308 *EMS mutagenesis*

Method Fig.1. Workflow showing EMS mutagenesis and CLLF phenotyping. 1) NG2 wild-type seeds were treated with ethyl methane sulfonate. 2) Chemically mutagenized NG2 seeds were sown into plant growth trays with commercial soil. 3) We observed chlorosis (somatic mutagenesis) in ~ 1 in 100 plants indicating successful mutagenesis. Seeds were harvested in bulk from these plants 4) 1st generation of mutant plants were allowed to colonize from soil and air inoculums outside in the garden. Later we checked leaf bacterial loads of ~300 individual plants after moving it to individual pots with commercial soil supplemented with garden soil inoculum. 5) After detailed phenotyping, we harvested seeds from individual plants and created a mutant plant library. 6) Barrier experiment; we used a thin plastic barrier to separate seedlings from soil. We did detailed phenotyping of NG2 and stable lines of CLLF plants (M4 generation) and harvested leaves of 3-week-old plants for microbiome analysis.

A well-characterized *Arabidopsis thaliana* ecotype "NG2" (NASC ID N2110865; Je-1) previously
collected from Jena, Germany was used for this study (16). Approximately 5000 seeds of *A. thaliana*
NG2 seeds were incubated with 0.3% ethyl methane sulphonate (EMS, Sigma-Aldrich; M0880-1G)
overnight as previously described (33). After repeated washing with sterile Milli-Q water, the M1 seeds
were resuspended in 0.1% agarose. Seeds were vernalized for 5 days at 4°C and sowed into soil (2:1
Floraton 3 Floragard Potting soil and perlite supplemented with 25g/ 6L soil mix substral osmocote as
fertilizer) as batches of 100 plants per tray and moved to a growth cabinet set at 22°C day-16h/18°C
night- 8h cycle. Chlorotic spots were observed in roughly one in a hundred plants after 2.5 weeks,
indicating somatic mutagenesis. Once the plants started producing seeds, siliques were harvested in
bulk from each tray. Later, approximately 200 seeds collected from each tray were sown into a natural
garden soil after surface sterilization and vernalization. Two weeks after germination, approximately
300 healthy M2 seedlings were randomly selected and transferred into individual pots containing
commercial soil supplemented with garden soil inoculum (60g natural garden soil dissolved in 1L

autoclaved Milli-Q water, mixed with 3L commercial soil) and moved to growth shelves. The plants

were allowed to grow in controlled conditions (22°C day (16h) / 18°C night (8h) cycle) in the growth

shelf. Once the plants were 3.5 weeks old, 2 leaves were collected and used for colony-forming unit

(CFU) counting. The leaves were ground in sterile 1x PBS by bead beating (1400 rpm for 30s) and a

dilution series was prepared. This was plated onto R2 agar and incubated for 48 h before CFU counting.

Based on a bacterial load higher than the wild-type, siliques were harvested from individual plants (n=1

in the M2 generation, Method Fig.1). Bacterial loads were further tested by CFU counting in the

subsequent generations and for this study, one of the mutants with the highest bacterial load and a

unique phenotype (CLLF) was used for further experiments. All described experiments were carried

out with at least M4 generation seeds.

Plant growth conditions

For microbiome analysis of plants colonized from the soil, seeds of the *A. thaliana* NG2 wildtype

(hereafter NG2) and the mutant genotype CLLF were surface sterilized by washing for 2 min in 2%

bleach followed by 30s in 70% ethanol and 2 sterile water rinses. They were sown onto commercial

soil supplemented with garden soil inoculum (See method section 1) after vernalization in 0.1% agarose

at 4°C for 5 days. Later, a plastic wrap was used to cover individual pots, leaving a small hole in the

middle to allow the seedling to germinate. This acted as a barrier between the leaves and soil, preventing

detection of any soil microbes on leaves due to increased soil contact in CLLF (Barrier Experiment).

Plants were then allowed to grow at 22°C day (16h) / 18°C night (8h) for 3 weeks in a plant climate

chamber (PolyKlima, Freising, Germany) before leaf harvest. Each sample represents 4 leaves from a

single plant. No plastic barrier was used for replicate experiments in which leaves were used either for

total microbiome or phytohormone analysis. For phytohormone analysis, commercial soil without

garden soil inoculum was used. For gnotobiotic growth, surface sterilized and vernalized seeds were

sowed into Linsmaier and Skoog media (4.3g/L LS salt mix (Linsmaier & *Skoog* (1964)), 1% sucrose,

0.5% MES, 0.3% phytagel) solidified in sterile 24 well plates.

Plant DNA extraction

We used a high throughput DNA extraction method to isolate DNA from leaves. 2-6 leaves were taken

for DNA extraction depending on the weight and after washing away dust and debris with autoclaved

Milli-Q water, were frozen at -80°C. Later, frozen samples were ground in a bead beater at 1400rpm,

for 30s to 60s to make a fine homogenate using 3mm metal and 0.25-0.5 mm glass beads. DNA was

extracted in CTAB (2% (w/v) Cetylimethylammoniumbromid, 100mM Tris with pH 8.0, 20mM EDTA

with pH 8, 1,4M NaCl, 1% (w/v) Polyvinylpyroledon) buffer and then purified from the lysate using

Phenol:Chloroform:Isoamyl alcohol (25:24:1) and precipitated using 0.7 volume 2-propanol.

Precipitated DNA was then dissolved in 10mM-Tris buffer for the barrier experiment or nuclease-free

water for the remaining microbiome analysis.

16S rRNA gene amplicon sequencing

For amplicon library preparation, we used a modified version of the host-associated microbe PCR

(hamPCR) protocol (17). Kapa HiFi enzyme (Roche; 07958846001) was used for all PCR reactions.

Zymomix, nuclease-free water, and CTAB extraction buffer were used as internal controls to evaluate

contamination and sequencing depth. In the first tagging step, we used universal primers targeting the

V3-V4 region of 16S rDNA modified with a 5' overhang. In addition to that, we used primers targeting

host-specific single copy gene (GIGANTEA or GI gene) and blocking oligos to limit the amplification

of chloroplast 16S rDNA (34). Each 10 μL 1st PCR mix contained 1x Kapa Buffer, 0.3 mM Kapa

dNTPs, 0.08 μM of each of the forward and reverse 16S rDNA and GI primers, 0.25 μM of each of the

blocking oligos, 0.2μL Kapa HiFi DNA polymerase, and 50 to 100 ng template genomic DNA.

Thermocycling steps included initial denaturation at 95°C for 3min, denaturation at 98°C for 20s, two-

step annealing at 58°C for 30s and 55°C for 1min, extension at 72°C for 1min (cycle repeated 5x), and

final extension at 72°C for 2min. An enzymatic cleanup was done to remove primer dimers and

inactivate nucleotides by directly adding 0.5μL of Antarctic phosphatase and Exonuclease 1 (New

England Biolabs, Inc; M0293L, M0289L) and 1.22μL Antarctic phosphatase buffer (1x final

concentration) to the 1st PCR mix. It was then incubated at 37°C for 30 minutes followed by 80°C for

15 minutes to inactivate the enzyme activity. Next, in a second PCR barcoded primers targeted the 5'

overhangs to amplify 1st PCR products. The 20μL 2nd PCR mix contained 1x Kapa Buffer, 0.375 mM

Kapa dNTPs, 0.3 μM of each forward and reverse primer, 0.5μL Kapa HiFi DNA polymerase, and 5μL

1st PCR product. The PCR program included initial denaturation 95°C for 3min, denaturation at 98°C

for 20s, annealing at 60°C for 1min, extension at 72°C for 1min (repeated 35x), and final extension at

72°C for 2min. The barcoded PCR products from the second PCR were cleaned by magnetic separation

using Sera mag beads. Later, we estimated the concentration of each sample by measuring fluorescence

using PicoGreen (Quant-iT™ PicoGreen™). After adjusting the concentration, all samples were pooled

into one single library and concentrated using sera mag beads. The concentrated library was then loaded

into a 2% agarose gel and stained with Roti gel stain. The separated GI and 16S rDNA bands were cut

out of the gel and DNA was eluted from each of the gel pieces using the GeneJET gel extraction kit

(Thermo Scientific™; K0691). After measuring the concentration of gel-eluted DNA, a final library

was prepared with an adjusted concentration of 95% 16S rDNA and 5% GI. We quantified the final

library with a Qubit (Thermo Fisher Scientific, Inc). Then the library was denatured and loaded onto a

MiSeq lane spiked with 10% PhiX genomic DNA to ensure sequence diversity. Using conventional

Illumina sequencing primers, 600 cycles of Illumina sequencing were carried out to obtain 300 bp

sequences in both the forward and reverse directions.

Amplicon sequencing data processing

For all datasets, adapter sequences were first removed from reads using Cutadapt 3.5 and reads were

split into samples according to barcodes using a custom script. Quality filtering and clustering the data

into amplicon sequencing variants (ASVs) were performed with the dada2 (version 1.18.0) algorithm,

using only forward reads because of its higher quality. Taxonomy was assigned to the ASVs using dada2 with the Silva 16S rDNA database (version 138.1) supplemented with the A. thaliana GI gene sequence. We examined positive and negative controls from all data sets. Positive control (Zymomix) had an expected distribution of taxa and negative controls had very few reads (<60 reads) suggesting minimal contaminations, so these were not processed further. R packages Phyloseq (version 1.34.0), VEGAN (version 2.5-7), and DEseq2 (version 1.44.0), were used for downstream analysis. Host-derived reads were removed from the ASV tables by removing family "Mitochondria", order "Chloroplast" and Genus "Arabidopsis GI". In addition, all samples were normalized to GI reads before analysis so that diversity patterns reflect the true abundances of leaf bacteria. Plant GI reads were also used to normalize the total number of bacterial reads to get an estimate of the relative bacterial loads of each sample. Other packages such as ggplot2, reshape2, and ggpubr were used for statistical analysis and plotting data. Scripts for generating the main figures from the ASV tables and metadata will be made publicly available on Figshare prior to publication.

Assessing plant survival rate in a garden experiment

To compare survival rates of CLLF and NG2, we germinated seeds in a mix of 1:4 garden soil to commercial soil (as described above) under laboratory conditions. Plants were sown into 7 replicate trays, each divided into two halves, one for each genotype. Plants were thinned to 30 seedlings of each genotype per tray, which was considered as one replicate. The trays with seedlings were moved to the garden 7 days after germination (in Nov 2022), and the number of plants surviving in each tray was counted every week until plants started flowering (in March 2023).

Phytohormone analysis

For the phytohormone extraction, we used a high-throughput method previously published by Schäfer et al. (2016). We collected 100mg of leaves from 3-week-old lab-grown plants in a pre-weighed 2mL screw-cap tube with two metal beads each. Then we fast-froze the samples in liquid nitrogen and stored at -80°C until further processing. The frozen samples were ground into a fine powder using a tissue lyser (MiniG 1600, SPEX Sample Prep). For the extraction, samples were homogenized in 15:4:1 (v/v/v) methanol: H_2O: formic acid extraction buffer containing isotope-labeled phytohormone standards, centrifuged at high speed, and the supernatant was collected. The supernatant was then passed through a reversed-phase solid-phase extraction (SPE) column (HR-X, Machery-Nagel, 738530.025M). Methanol from the flow-through was then allowed to evaporate and the remaining samples were reconstituted in 1N formic acid. Reconstituted samples were then passed through a mixed-mode cation exchange SPE column (HR-XC, Machery-Nagel, 738540.025M), and the flow through was discarded. The column was washed once with 1N formic acid and eluted using an 80% aqueous methanol containing 0.2N formic acid. This eluted fraction contained abscisic acid (ABA), salicylic acid (SA), and jasmonates (JAs), which were analyzed by LC-MS/MS on a triple-quadrupole mass spectrometer (EVO-Q Elite, Bruker) using the chromatog. The chromatographic condition and

430 multiple-reaction-monitoring details for the compound detection are described in detail by Schäfer et
431 al. (35).

Pathogen infection assays

For the bacterial pathogen infection assay, CLLF and NG2-Wt were grown in gnotobiotic conditions in
LS media in square petri dishes. *Pseudomonas syringae pv syringae* DC3000 (Pst DC3000) was grown
in LB agar supplemented with rifampicillin for 48 h at 30°C. Colonies were then collected using a loop
and resuspended in 10 mM MgCl2. OD was measured using a UV spectrometer and adjusted to 0.02
OD. 0.02 % Silwet L-77 was freshly added to the bacterial suspension just before flood inoculation.
For flood inoculation, 40mL bacterial culture was poured into the petri dishes with plants. Plants
inoculated with 40mL of 10 mM MgCl2 plus 0.02 % Silwet L-77 were taken as controls. Disease
symptoms were measured after 24h.

For the fungal infection assay, plants grown in the semi-gnotobiotic (2-week-old seedlings grown in LS
media transplanted into commercial soil) system. *Sclerotinia sclerotiorum* (Ssc) was grown in Potato-
Dextrose agar (PDA) plates from a subculture for 4-6 days. We collected agar discs from the actively
growing region of Ssc plates and used this to inoculate liquid PDB (Potato-Dextrose Broth). Liquid
culture was then allowed to grow overnight at room temperature with shaking. We harvested the
mycelia the next day after passing it through an autoclaved nylon membrane. After washing them with
autoclaved milliQ water, they were resuspended in fresh PDB. This was then macerated to get small
hyphal fragments using a sterile macerator. Hyphal fragments were adjusted to 2×10^5 fragments/ml.
5µL of this hyphae suspension was used to inoculate leaves of 4-week-old plants. The leaf midrib was
avoided for easy measurement of the lesion and lesion size was measured at both 18 hours post-
inoculation (hpi) and 40hpi.

Synthetic community preparation

We used bacterial strains that were previously isolated from CLLF or NG2 leaves for preparing the
Synthetic Communities (SynComs) (Supplementary Table 1A). All strains were grown on R2 agar
plates. Depending on each strain's growth rate, it took 1 to 5 days to grow all strains. The bacteria were
then scraped off the plates and resuspended in 1x phosphate buffer saline (PBS). Optical density (OD)
was measured using a 1:10 dilution of each bacterial suspension in a plate reader. Since we observed
huge variation in the number of CFUs per OD of different bacteria, we decided to consider CFUs for
preparing SynComs. Therefore, an OD to CFU relationship of each bacterium was measured and this
was used to combine the bacterial suspensions into a single inoculum so that all strains would have
approximately same number of CFUs (~30 CFU per strain). A total of 45 strains belonging to diverse
plant-associated bacterial phyla constituted the SynCom1. The SynCom2 lacked strains belonging to
Comamonadaceae and Oxalobacteriaceae (Betaproteobacteria). SynCom3 lacked strains belonging to
Bacteroides and SynCom4 lacked Betaproteobacteria and Bacteroides. For each missing strain in

SynComs 2-4, we added the same amount of 1x PBS buffer to the SynCom stocks. In addition, aliquots

of all four SynComs were mixed with an equal amount of 40% glycerol for future analysis. To check

the SynCom compositions, 12µL from the glycerol stocks were later inoculated onto R2 agar plates and

grown for 5 days at 28°C. All cells were harvested from R2 plates by suspending into 1x PBS, and later,

all samples were centrifuged at 5000rpm in a table-top centrifuge and the pellet was resuspended in

CD1 solution of DNAeasy plant pro kit (Qiagen; 0142924730) along with 0.25-0.5 mm glass beads.

It was then bead-beated at 1400rpm for 30s. DNA was extracted using the kit following the suggested

protocol. The communities were analyzed using the 16S rRNA gene amplicon sequencing

pipeline as described above.

Semi-gnotobiotic plant growth conditions and stress treatments

Method Fig.2. Workflow showing SynCom experiment in a semi-gnotobiotic system. Gnotobiotic phase; Sterilized seeds of CLLF and NG2 plants were germinated in sterile LS media. 7-day old seedlings were inoculated with 5µL SynCom or 1X PBS. Plants were allowed to grow in LS media with or without the SynComs for 1 week at 22h day/8h night cycles. **Non-gnotobiotic phase;** 2-week-old pre-colonized plants are moved to commercial soil. After two weeks of growth (22h day/8h night cycles), half the of plants were challenged with temperature stress at 30°C for 48 h, followed by 2 days of recovery under nomal condition. Later half of plants from each treatment was challenged with PstDC3000. After 2 days, leaves were sampled for microbiome or RNAseq analysis.

We noted that in fully gnotobiotic, enclosed growth systems, plants tended to become stressed by humidity. Therefore, we developed a semi-gnotobiotic system in which plants are pre-colonized by a defined community in a gnotobiotic environment and then moved to a standard, non-sterile laboratory soil and grown in trays. When we tested the semi-gnotobiotic system with a fluorescently labeled isolate of *Stenotrophomonas* sp., leaf microbiomes were dominated by *Stenotrophomonas* even after several weeks in the soil, suggesting that this is an effective inoculation strategy (36). For this study, surface-sterilized and vernalized seeds of CLLF and NG2 were sown into LS media. 7-day-old seedlings were inoculated with 5µL inoculum of one of the four SynComs or 1x PBS buffer. This was enough inoculum to wet the leaves and roots. There were 48 replicates per genotype and SynCom. The plants were grown for 1 week in LS before being transplanted into commercial soil at 22°C day-16h/18°C night- 8h cycles. After 2 weeks of normal growth in soil, 16 plants from each genotype inoculated with one of the 4 SynCom or 1x PBS were given heat stress by moving into a growth chamber set at 30°C day (16h)

and night (8h) for 48h. The remaining plants were allowed to grow at normal 22°C day-16h/18°C night-

8h cycles. Later the plants that underwent heat stress were moved back into normal growth conditions
and all plants were grown for 2 days under 22°C day-16h/18°C night condition before treating with
Pseudomonas syringae pv. syrinage DC3000 (Pst DC3000).   Half of the plants from each treatment
were spray-inoculated with 0.02 OD cultures of Pst DC3000 suspended in sterile milliQ water
supplemented with 0.02% Silwet L-77 and the remaining half with sterile milliQ water. After 48 hours
post-inoculation, we measured leaf function using an LI-COR 600, and 2-3 leaves from at least 6 plants
were sampled for microbiome analysis from all the different treatments. 2-3 leaves from SynCom 1, 4,
and negative controls with and without heat stress were sampled for RNAseq analysis (Method Fig. 2).

RNA sequencing and data processing

All samples for RNAseq were fast-frozen using liquid nitrogen and stored at -80°C. The samples were
then ground to a fine powder in a bead beater using 2 metal beads. Finely ground samples were used
for RNA isolation using the RNAeasy Plant Mini Kit (Qiagen, ID: 74904). Total RNA was sent to
Eurofins genomics for rRNA depletion, cDNA synthesis, library preparation and sequencing. Raw data
received from Eurofins Genomics were then used for downstream processing as follows. Raw
sequencing reads were assessed for quality using FastQC (version
0.11.9; https://www.bioinformatics.babraham.ac.uk/projects/fastqc). Adaptor trimming, quality
filtering, and read preprocessing were performed using fastp (version 0.23.2) (37). In detail, 5′ and 3′
bases with a Phred quality score below 28 were cut and reads were removed if they had more than one
ambiguous base, an average quality score below 28, or a length of fewer than 15 bases. Processed reads
were aligned to the current *A. thaliana* genome (tair10.1) using Hisat2 (version 2.2.1) with standard
parameters (38). The aligned reads were sorted and indexed using SAMtools (version 1.11) (39). Read
counting was performed using featureCounts (version 2.0.1) (40) with the tair10.1 annotation as a
reference. Differential gene expression analysis was performed using DESeq2 (version 1.38.3) (41) and
comparisons having a false discovery rate (FDR) adjusted p-value <0.05 were deemed to be statistically
significant (42). Gene ontology (GO) enrichment was performed for each DE gene set using the R
package, clusterProfiler (version 4.10.0), and the GO category was assigned using the annotation data
package org.At.tair.db (version 3.18.0). The detailed scripts with step-by-step instructions were
uploaded to GitHub (https://github.com/Bioinformatics-Core-Facility-Jena/SE20231212_167).

Growth assay with salicylic acid

All strains belonging to Betaproteobacteria and Bacteroides and representative strains from the
remaining major taxonomic groups of the SynComs were used for growth assays. First, bacteria were
grown in R2 agar at 28°C for 4 days. Fresh colonies were then harvested from agar plates using a sterile
loop and resuspended in 1x PBS by vortexing and repeated pipetting. After measuring the optical
density (OD_{600}), the OD_{600} of all samples was adjusted to 0.02. 10 µL of this pre-adjusted culture was
used as an inoculum for measuring growth in 190 µL of M9 minimal media (M9 salts: 33.7mM
Na_2HPO_4, 22mM KH_2PO_4, 8.55mM NaCl, 9.35mM NH_4Cl, and 2 mM $MgSO_4*7H_20$, 0.2 mM $CaCl_2$, 1
µg/mL biotin, 1 µg/mL thiamine, and trace elements: 134 µM EDTA, 31 µM $FeCl_3-6H_2O$, 6.2 µM

ZnCl$_2$, 0.76 µM CuCl$_2$-2H$_2$O, 0.42 µM CoCl$_2$-2H$_2$O, 1.62 µM H$_3$BO$_3$, 0.081 µM MnCl$_2$-4H$_2$O)

supplemented either with 0.5mM Salicylic acid (SA), concentration adapted from S. L. Lebeis, *et al* (9),

11 mM Fructose plus 11 mM Sucrose (PC) or no additional carbon source (Control) for the growth

assay. All samples were then incubated at 28°C with shaking (220rpm) in 15-minute intervals in a plate

reader for 5 days. OD$_{600}$ was measured every 1.5 h.

Competing Interest Statement

The authors declare no competing interests.

Data Sharing Plan

Data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary
Materials. Raw sequencing data is available on NCBI-SRA (16S data: PRJNA1120601, RNAseq
data: XXX) and processed data with code to generate the main figures will be available on Figshare
before final publication. The detailed scripts with step-by-step instructions for RNA-seq analysis were
uploaded to GitHub (https://github.com/Bioinformatics-Core-Facility-Jena/SE20231212_167).

Funding Information

Carl Zeiss Foundation via Jena School for Microbial Communication (JJ, MTA)

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's
Excellence Strategy - EXC 2051 - Projektnummer 390713860 (JJ, MTA, MM)

Max Planck Society (RH)

The Ministry for Economics, Sciences and Digital Society of Thuringia (TMWWDG), under the
framework of the Landesprogramm ProDigital [DigLeben-5575/10-9]. (EB)
559
560   Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under CRC 1076
561   'AquaDiva', subproject A06. (MM)
562

563   <u>References</u>
564
1. McNew, George L., "The nature origin and evolution of parasitism" in *Plant Pathology: An*
Advanced Treatise, (Academic Press, 1960), pp. 19–69.

2. F. Entila, X. Han, A. Mine, P. Schulze-Lefert, K. Tsuda, Commensal lifestyle regulated by a
negative feedback loop between Arabidopsis ROS and the bacterial T2SS. *Nat. Commun.* **15**, 456
(2024).

3. L. Maignien, E. A. DeForce, M. E. Chafee, A. M. Eren, S. L. Simmons, Ecological Succession and
Stochastic Variation in the Assembly of Arabidopsis thaliana Phyllosphere Communities. *mBio* **5**
(2014).

4. K. M. Meyer, *et al.*, Plant neighborhood shapes diversity and reduces interspecific variation of
the phyllosphere microbiome. *ISME J.* **16**, 1376–1387 (2022).

5. T. Chen, *et al.*, A plant genetic network for preventing dysbiosis in the phyllosphere. *Nature* **580**,
653–657 (2020).

6. P. J. P. L. Teixeira, *et al.*, Specific modulation of the root immune system by a community of
commensal bacteria. *Proc. Natl. Acad. Sci.* **118**, e2100678118 (2021).

7. S. Pfeilmeier, *et al.*, The plant NADPH oxidase RBOHD is required for microbiota homeostasis in
leaves. *Nat. Microbiol.* **6**, 852–864 (2021).

8. C. J. Harbort, *et al.*, Root-Secreted Coumarins and the Microbiota Interact to Improve Iron
Nutrition in Arabidopsis. *Cell Host Microbe* **28**, 825-837.e6 (2020).

9. S. L. Lebeis, *et al.*, Salicylic acid modulates colonization of the root microbiome by specific
bacterial taxa. *Science* **349**, 860–864 (2015).

10. K. Unger, *et al.*, Beyond defense: Glucosinolate structural diversity shapes recruitment of a
metabolic network of leaf-associated bacteria. [Preprint] (2023). Available at:
http://biorxiv.org/lookup/doi/10.1101/2023.12.04.567830 [Accessed 13 May 2024].

11. B. Huot, *et al.*, Dual impact of elevated temperature on plant defence and bacterial virulence in
Arabidopsis. *Nat. Commun.* **8**, 1808 (2017).

12. J. Qiu, *et al.*, Warm temperature compromises JA-regulated basal resistance to enhance
Magnaporthe oryzae infection in rice. *Mol. Plant* **15**, 723–739 (2022).

13. H.-G. Mang, *et al.*, Abscisic Acid Deficiency Antagonizes High-Temperature Inhibition of Disease
Resistance through Enhancing Nuclear Accumulation of Resistance Proteins SNC1 and RPS4 in
Arabidopsis. *Plant Cell* **24**, 1271–1284 (2012).

14. A. Issa, *et al.*, Impacts of Paraburkholderia phytofirmans Strain PsJN on Tomato (Lycopersicon
esculentum L.) Under High Temperature. *Front. Plant Sci.* **9**, 1397 (2018).

15. R. J. Rodriguez, *et al.*, Stress tolerance in plants via habitat-adapted symbiosis. *ISME J.* **2**, 404–
416 (2008).

16. Mayer, Teresa, "Whose microbiome is it? Adaptive interactions of keystone species in plant
microbiomes.," Friedrich Schiller University of Jena, Jena, Germany. (2022).

17. D. S. Lundberg, *et al.*, Host-associated microbe PCR (hamPCR) enables convenient measurement
of both microbial load and community composition. *eLife* **10**, e66186 (2021).

18. S. A. Vincent, A. Ebertz, P. D. Spanu, P. F. Devlin, Salicylic Acid-Mediated Disturbance Increases
Bacterial Diversity in the Phyllosphere but Is Overcome by a Dominant Core Community. *Front.*
Microbiol. **13**, 809940 (2022).

19. L.-J. Wang, *et al.*, Salicylic acid alleviates decreases in photosynthesis under heat stress and
accelerates recovery in grapevine leaves. *BMC Plant Biol.* **10**, 34 (2010).

20. T. L. Karasov, *et al.*, The relationship between microbial population size and disease in the
Arabidopsis thaliana phyllosphere. [Preprint] (2019). Available at:
http://biorxiv.org/lookup/doi/10.1101/828814 [Accessed 13 May 2024].

21. K. Ogata, M. Ohsugi, M. Tomita, T. Tochikura, The Production of α-Ketoglutaric Acid from
Salicylic Acid by *Pseudomonas* sp.†. *Agric. Biol. Chem.* **34**, 364–369 (1970).

22. G.-H. Lim, *et al.*, Plasmodesmata Localizing Proteins Regulate Transport and Signaling during
Systemic Acquired Immunity in Plants. *Cell Host Microbe* **19**, 541–549 (2016).

23. W. Liu, S.-W. Park, 12-oxo-Phytodienoic Acid: A Fuse and/or Switch of Plant Growth and Defense
Responses? *Front. Plant Sci.* **12**, 724079 (2021).

24. I. Monte, *et al.*, An Ancient COI1-Independent Function for Reactive Electrophilic Oxylipins in
Thermotolerance. *Curr. Biol.* **30**, 962-971.e3 (2020).

25. M. Antunez-Lamas, *et al.*, Bacterial chemoattraction towards jasmonate plays a role in the entry
of *Dickeya dadantii* through wounded tissues. *Mol. Microbiol.* **74**, 662–671 (2009).

26. O. S. Kulkarni, *et al.*, Volatile methyl jasmonate from roots triggers host-beneficial soil
microbiome biofilms. *Nat. Chem. Biol.* **20**, 473–483 (2024).

27. C. S. Liou, *et al.*, A Metabolic Pathway for Activation of Dietary Glucosinolates by a Human Gut
Symbiont. *Cell* **180**, 717-728.e19 (2020).

28. H.-W. Jeon, *et al.*, Contrasting and conserved roles of NPR pathways in diverged land plant
lineages. [Preprint] (2022). Available at:
http://biorxiv.org/lookup/doi/10.1101/2022.07.19.500630 [Accessed 13 May 2024].

29. Y. Liu, *et al.*, Diverse Roles of the Salicylic Acid Receptors NPR1 and NPR3/NPR4 in Plant
Immunity. *Plant Cell* **32**, 4002–4016 (2020).

30. J. H. Kim, *et al.*, Increasing the resilience of plant immunity to a warming climate. *Nature* **607**,
339–344 (2022).

31. M. R. Wagner, *et al.*, Host genotype and age shape the leaf and root microbiomes of a wild
perennial plant. *Nat Commun* **7**, 12151 (2016).

32. B. N. Archibald, V. Zhong, J. A. N. Brophy, Policy makers, genetic engineers, and an engaged
public can work together to create climate-resilient plants. *PLOS Biol.* **21**, e3002208 (2023).

33. D. Weigel, J. Glazebrook, EMS Mutagenesis of *Arabidopsis* Seed. *Cold Spring Harb. Protoc.* **2006**,
pdb.prot4621 (2006).

34. T. Mayer, *et al.*, Obtaining deeper insights into microbiome diversity using a simple method to
block host and nontargets in amplicon sequencing. *Mol. Ecol. Resour.* **21**, 1952–1965 (2021).

35. M. Schäfer, C. Brütting, I. T. Baldwin, M. Kallenbach, High-throughput quantification of more
than 100 primary- and secondary-metabolites, and phytohormones by a single solid-phase
extraction based sample preparation with analysis by UHPLC–HESI–MS/MS. *Plant Methods* **12**,
30 (2016).

36. Teutloff, Erik, "Balance in plant bacterial recruitment affects resilience to fungal invasion of the
phyllosphere," Friedrich Schiller University of Jena, Jena, Germany. (2023).

37. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

38. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

39. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

40. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

41. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

42. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **57**, 289–300 (1995).

658