Systems Biology

The FBA Solution Space Kernel – Introduction and Illustrative Examples

Wynand S. Verwoerd[1,*] and Longfei Mao[2]

[1]Centre for Advanced Computational Solutions, Faculty of Ag & Life Sciences, Lincoln University, Lincoln 4647, New Zealand.

[2]Department of Pharmacy, College of Biology, Hunan University, Changsha, China.

*To whom correspondence should be addressed.

Abstract

Motivation: The solution space of an FBA-based model of cellular metabolism, can be characterized by extraction of a bounded, low dimensional kernel (the SSK) that facilitates perceiving it as a geometric object in multidimensional flux space. The aim is to produce an amenable description, intermediate between the single feasible extreme flux of FBA, and the intractable proliferation of extreme modes in conventional solution space descriptions. Fluxes that remain fixed are separated off while the focus of interest is put on the subset of variable fluxes that have a nonzero but finite range of values. For unbounded fluxes, a finite subrange that geometrically corresponds to the variable flux range is determined, and is supplemented by a limited set of rays that encapsulates their unbounded aspects. In this way the kernel emphasizes the realistic range of flux variation allowed in the interconnected biochemical network by e.g. limited nutrient uptake, an optimized objective and other model constraints.

Results: It is demonstrated how knowledge of the SSK and accompanying rays can be exploited to explore representative flux states of the metabolic network. A full presentation of the SSK approach was the subject of a research monograph (Verwoerd, 2022a). Moreover, since bioengineering interventions such as gene knockouts modify the solution space, new tools based on the SSK analysis are presented that predict the effects of such interventions on a target flux constructed to represent a desired metabolic output. A simple metabolic model is used first to demonstrate the special concepts and constructions needed to define and compute the SSK. The demonstration model is tweaked to produce typical behaviors of larger models, but with kernels in 1,2 or 3 dimensions that are explicitly displayed to visualize the concepts. General applicability to models where visualization is inaccessible, is illustrated by showing evaluation of potential bioengineering strategies for a genome scale model.

Availability: Results from an interactive software implementation freely available at doi:10.5281/zenodo.11483838.

Contact: Wynand.Verwoerd@lincoln.ac.nz

Supplementary information: Computational details and input data are available as accompanying documents at Bioinformatics online.

1 Introduction

A fundamental premise of the constraints-based modeling of cellular metabolism, is that stoichiometric, thermodynamic, and capacity constraints limit the vector of flux values for biochemical reactions in a cell, to a feasible region (the solution space) within the flux space created by taking each reaction flux as an independent axis.

Flux balance analysis (FBA) (Orth et al., 2010; B. O. Palsson, 2015) takes this one step further, assuming that the actual metabolic state of a cell is well represented by the flux vector that optimizes an appropriately chosen objective such as the biomass growth rate. With linear constraints and objective, solving for an optimal flux value is a standard linear programming (LP) (Jeter, 1986) problem.

However, even though realistic models remain extensively underdetermined even after fixing an objective, LP only delivers a single optimal flux value. In contrast, extreme pathway or elementary mode analysis (Papin *et al.*, 2004; Schilling *et al.*, 2000; Schuster *et al.*, 1999) gives an exact mathematical description of all feasible flux

values by finding a convex vector basis that completely spans the solution space.

Neither of these approaches gives a readily utilizable description of the flux or range of fluxes available to a cell. In the case of FBA, this is because the single flux it yields cannot be taken as typical; for the Simplex LP algorithm it is in fact by construction located at a vertex, i.e. at the very extreme of the solution space. For extreme pathways, the problem is that the sheer number of basis vectors becomes overwhelmingly large for realistic metabolic models.

A popular way to address this difficulty, is an elaboration of FBA called flux variability analysis (FVA) (Mahadevan & Schilling, 2003). This consists of using LP to find the minimum and maximum value that each reaction flux can achieve within the model constraints. This fixes a cuboid bounding box in the multidimensional flux space, within which the solution space lies. Numerical difficulties are often encountered in performing FVA, but a more fundamental limitation is that in high dimensional spaces the solution space polytope occupies a negligible fraction of the bounding box, rendering it uninformative. This point is further discussed below.

The solution space kernel (SSK) approach (Verwoerd, 2022a) steers a midway between the extremes and pitfalls described. It expands on FBA by specifying the range of feasible flux values rather than a single value. It characterizes this region in a way that is far more specific and informative than FVA, but in terms of far fewer parameters than the extreme pathway/elementary mode approach. In direct contrast to the FVA bounding box, every flux value that falls in the SSK is an actual feasible flux.

This goal is achieved by identifying a compact, low dimensional subspace of the SS from which most, if not all, feasible fluxes can be reached by the addition of a limited number of ray vectors that are also identified. This subspace is termed the *kernel* of the SS. The full SS is unbounded in some directions for virtually all metabolic models, but even so the kernel remains bounded. It is specified by a small number of constraints in far fewer dimensions than the full SS.

To enhance comprehension the SSK is treated as a geometrical object in multidimensional space with a shape described by a set of mutually orthogonal maximal chords and resulting aspect ratios. Even more concretely, a list of explicit flux vectors that specify the center and chosen periphery points of the SSK allows expected values and ranges for flux dependent attributes of cell metabolism, to be evaluated.

This paper clarifies these concepts by detailing the application to a simple metabolic network. No attempt is made here to explain the extensive computation that is needed to construct an SSK; more details are available in work referenced in the next section.

However, in broad outline the major stages are as follows:

1. Separating off all reaction fluxes that remain fixed over the entire SS.
2. Identifying directions in flux space for which the SS is unbounded and separating them off by defining ray vectors.
3. The remaining SS kernel constraints and flux variables are obtained by vector projection, introduction of capping constraints and other means.
4. The SSK extent and shape is delineated by finding a set of mutually orthogonal, maximal chords spanning the SSK.
5. The center and periphery of the compact kernel are determined and further characterizes the SSK location, size and shape in a reduced dimensional space.

The last stage allows an even more amenable approximation of the SSK for cases where it still has a sizable dimensionality N. A collection of representative periphery points is explicitly determined in this stage, and their convex hull delineates a central region that typically covers 80% or more of the full extent of the SSK. This is

termed the peripheral point polytope or PPP and has only of the order of $4N$ explicitly listed vertices that can be used as a "poor man's version" of extreme pathways, to specify the most physically significant central part of the SSK.

The solution space (SS) concept needs further discussion. Generally, in constraints based analysis this is considered to be the flux subspace that satisfies stoichiometry and other external constraints. For FVA the optimized objective is added as a further constraint, further reducing the subspace to what could be termed an objective subspace or OS. The distinction is ignored in most of what follows since the kernel analysis can be applied to either the SS or OS. However, for most models upper limits for individual fluxes are unknown and the SS is a simple unbounded hypercone with its apex at the flux space coordinate origin representing the zero flux solution. In this case just the apex point serves as the kernel, since all other feasible fluxes can be reached from it by adding a suitable ray vector.

For this the reason the SSKernel analysis mostly comes into its own for the OS, which has a more complex shape. A linear objective defines a hyperplane, and the OS is the intersection between this and the SS hypercone. This intersection generally retains some unbounded directions, but also acquires bounded facets. It is these bounded facets that embody the biologically significant flux constraints. Determining all feasible, bounded facets (FBF's) is one of the major computational challenges of the SSK calculation. Once found, the kernel is constructed as the hypervolume bordered by all FBF's, together with any additional capping constraints needed to partition it from the unbounded section of the OS.

This computational framework can also be exploited to characterize the dimensionality and shape of a simple cone SS more fully. To do this its minimal single point kernel is extended by capping the cone at a chosen radius. This allows the "cross-sectional" shape of the simple cone to be characterized by chords and aspect ratios in a similar way as is done for the OS.

In this sense the SS kernel approach can be usefully applied to both SS and OS, and to simplify terminology both concepts are henceforth referred to as the solution space.

Concepts touched upon in this brief description are further elucidated by the explicit examples that is the main subject of this article.

2   Methods

The computational processing is done by the software package, SSKernel (Verwoerd, 2024), readily available as a Github release. This implements an interactive interface that allows the user to perform the major stages as outlined in the previous section, adjusting relevant parameters for each subsequent stage as needed. The software is accompanied by a user manual that describes the concepts and calculations in some more detail than is possible here.

To process a model, it is specified in the format defined by the COBRA toolbox (Schellenberger et al., 2011) as a *.mat file or alternatively in SBML format.

An interactive run of the SSKernel software produces two sets of output. First, there is a 2-page PDF file that summarizes the main outcomes of the calculation such as stage by stage dimension reductions, SSK shape information etc. Secondly, the full numerical results including a constraint based specification of the SSK, explicit ray vectors and peripheral flux points are exported to an output file in Data Interchange Format (*.dif). Examples of the input file and two output files for the toy model discussed in section 3 are provided in the supplementary data.

Also included in the SSK software are tools for exploring the solution space and how it is changed by bioengineering interventions. These are used independently of the interactive interface for post-

processing, and takes the abovementioned DIF file as input. The realistic network example discussed in section 3.2 is based on processing the DIF output file of an SSK run on the red blood cell model iAB_RBC_283 (Bordbar, Jamshidi, & Palsson, 2011).

The calculation of the SSK entails many special concepts and algorithms for dealing with polytopes in high dimensional spaces, and a full exposition has been presented in a recent monograph (Verwoerd, 2022a). However, the exploratory tools are a subsequent development and the formal details of the new algorithms they use are presented in the Supplementary Information to this article.

3 Results

The results are presented in two main parts.

First, the main features of the SSK are illustrated based on a simple model system. This is a toy model formulated in a 10-dimensional flux space and allows SSK reductions to 2 or 3 dimensions so that they can be graphically displayed. The emphasis here is on transparency. Such a simple model does not demonstrate the full power of simplification that SSK analysis is capable of. For that, the reader is referred to the dedicated monograph (Verwoerd, 2022a) which discusses application to state-of-the-art models involving thousands of flux space dimensions.

Second, it is shown how the SSK and rays can be used to explore the solution space and its response to flux restrictions. Again, the toy model illustrates the concepts, and an application to a realistic metabolic model is also shown to demonstrate its practical usefulness.

3.1 Elucidating SSK Concepts

The toy network used here, originates from a textbook demonstration of extreme pathways (B. Ø. Palsson, 2006) and is shown in Figure . Various slight elaborations of the toy model are made below to reproduce a number of scenarios commonly encountered in models of real cellular metabolism.

Figure 1. Toy network that produces a product *E* and byproduct *byp* from six internal reactions with fluxes F1 to F6. F7 is an external input flux and F8 and F9 are external output fluxes.

3.1.1 Scenario 1: A simple unbounded cone solution space

The bare toy model is representative of realistic models that contain no objective. In the conventional description, metabolic states of this network are superpositions of its three distinct extreme pathways (B. Ø. Palsson, 2006), i.e. its SS is a 3-sided infinite cone in 9 flux dimensions.

The SSK analysis uses a default capping radius as mentioned above to reduce this to a finite kernel which is explicitly specified

by 6 constraints on 3 variables. The reduction from the 16 constraints on 9 variables of the original model merely hints at the large reductions in realistic models.

Figure 2 Shows a 3D plot of the SSK. The SSK algorithm finds 3 peripheral rays aligned along the 3 long edges of the cone, and

Figure 2. (a) The capped SSK for the toy model without objective. The main chords are shown as red arrows. (b) The periphery point polytope (PPP) superimposed on the SSK shows coverage of most but not all of its volume.

each yielding one capping plane visible at the top in the figure. These ray directions are identical to the extreme pathways listed in Figure 13.1 of (B. Ø. Palsson, 2006). However, that is generally not the case; for example, if the SS was a 5-sided cone in 3D, there would be 5 extreme pathways, but still only 3 peripheral rays, since they are constructed to be only a vectorially complete basis for the ray space.

The coordinate axes in Figure 2 represent the particular 3D subspace within the full 9D flux space, in which the model allows feasible metabolic fluxes. Specifically, they are by construction the directions of the main chords of the SSK, shown as red arrows in the figure. For the simple unbounded cone of this model, the arbitrary capping radius means that the actual chord lengths are not meaningful. However, their ratios are independent of the radius and characterize the shape of the cone. For this model, these aspect ratios are 1.2, 2.1 and 2.5 respectively. The lowest value reflects the fairly regular cross section, and the other values the quite sharp cone as seen in Figure 2.

3.1.2 Scenario 2: A fully bounded solution space

Next, the model is made slightly more realistic by introducing an upper limit for the input flux F7. FVA shows that this constrains all nine fluxes to finite, nonzero ranges. There are no fixed fluxes and the FVA bounding box is a cuboid in 9 dimensions.

With this SS already bounded, no rays or capping hyperplanes need to be calculated, and the SSK is in fact identical to the SS for this case. The SSK shape analysis remains applicable and the result is very similar to that shown in Figure 2, only the cone is now truncated by a single plane fixed by the upper limit on F7. So, the SSK is in this case given by just 4 constraints on 3 variables.

Provided that the F7 cutoff value has been set at a physically meaningful value, the 3 main chords do now indicate the range of metabolic flux variation allowed by the model along the 3 available orthogonal directions. Note how this is far more specific than the 9D FVA cuboid within which this truncated cone is contained.

The aspect ratios derived from the chords are 1.5, 1.8 and 2.8 indicating a similarly sharp cone, but with a slightly flattened cross section compared to Figure 2.

3.1.3 Scenario 3: Optimized objective in a bounded SS

Moving on to a case more appropriate to the purpose for which the SSK analysis is designed, we introduce an optimized objective into the model while keeping the input flux limitation. A biomass composition of 80% of metabolite *E* and 20% of the byproduct *byp* is assumed, and incorporated into the model by adding a further output flux F10 that exports these metabolites in a 4:1 ratio. The objective is taken as maximizing F10.

An FVA calculation to characterize the OS with F10 kept fixed to its optimized value, shows that a total of 5 fluxes become fixed, leaving a 5D cuboid as the bounding box for feasible metabolic states.

The OS is now bounded, so no rays or capping is required and the SSK is identical to the OS. The calculation algorithm shows that the SSK/OS is only 2-dimensional and is plotted in Figure 3.

Clearly this SSK is simply the intersection of the objective plane with the triangular cone of Figure 2 and as expected the aspect ratio of 1.4 is similar to that of the cone cross section.

Figure 3. The OS and SSK for the toy model with a restricted input flux and a fixed objective. A centering algorithm determines the green center point and a semi-random selection of periphery points shown in black. The main chords are shown as red arrows.

The figure also illustrates the outcome of the centering procedure incorporated in the SSK calculation. The centre is the green dot which also serves as the coordinate origin in the SSK specification of 3 constraints on 2 variables It also finds a series of periphery points shown as black dots. These points form a plausible sample of explicit feasible fluxes that encapsulates the range of variation of metabolic states allowed by the model. Their convex hull (the PPP), covers 100% of the SSK, but as shown in Figure 2(b) that is not generally the case.

3.1.4 Scenario 4: Comparing the SSK and FVA

In the first 3 scenarios, the FVA bounding box is in each case in a much higher dimensional space than the SSK and could not be drawn. To illuminate the relationship between the two constructs, we further restrict the model used in scenario 3 by keeping also F2 at a fixed value. This fixes F3 as well, reducing the bounding box from 5 to 3 dimensions. Removal of the F2 degree of freedom also reduces the SSK to one dimension. The result is shown in Figure 4.

Unlike the previous figures, for this case not just the SSK subspace is shown but rather the 3D subspace that contains the variable fluxes F4, F5 and F6 taken as its coordinate axes.. The FVA bounding box accurately depicts the range of variation along each axis, but globally gives a very poor rendering of the actual extent of feasible

flux solutions. Only the SSK shows this exactly, namely as just a single line, the diagonal of the cuboid box. This only occupies a negligible fraction of the FVA box volume, so for example random sampling from the cuboid is unlikely to discover a single feasible flux combination.

Such dimensional discrepancy between the bounding cuboid and

Figure 4. The 1D linear SSK (red line) and 3D FVA bounding box (in colour) for the toy model, with an optimized objective but constrained to leave only F4, F5 and F6 variable, while other fluxes assume fixed non-zero values.

the SSK was also noted in the previous scenarios and is a universal feature of realistic metabolic models. There the FVA overestimation is far worse, because even where the bounding box occupies the same dimensions as the SSK, the fraction of the box that the SSK occupies is still negligible. This is because of its typical high aspect ratio and the scaling effects in high dimensions mentioned in the introduction and discussed in more detail in (Verwoerd, 2022a).

In the discussion so far, the FVA limits were used as a familiar indication of flux ranges to discover fixed fluxes, but we note that the Hop, Skip and Jump algorithm incorporated in the SSK calculation performs this task faster by orders of magnitude and also more reliably. These advantages are not manifest in the small toy model.

3.1.5 Scenario 5: SS partially bounded; tangent capping.

The next scenario is chosen to resemble a realistic solution space as closely as possible within the size limitations of the toy model.

This is brought about by putting a lower limit on F2 and allowing F3 to be reversible. The same objective and input flux limitation as in scenario 3 is used. Again there are 5 fixed fluxes and 5 variable ones, and the feasible solutions are restricted to a 2D plane within this 5D subspace. Unlike the previous scenarios this *reduced solution space* or RSS (reduced by the exclusion of fixed fluxes) is not bounded any more. So there are ray directions in the solution space – in fact only a single ray, the green arrow shown in Figure .

Despite the RSS being unbounded, some of its boundary facets remain bounded, emphasized as thick lines in the figure. Along with the 2 bounded facets (FBF's), there are also 2 unbounded ones. To achieve a bounded kernel, the SSK procedure introduces a single *capping hyperplane*, the dotted line BC, that partitions the RSS.

Geometrically, any flux value in the unbounded section can be reached from at least one flux point in the bounded section (the SSK) by adding a multiple of the ray vector.

This is guaranteed by having chosen the capping line such that it does not bisect any of the bounded facets. The capping radius (the perpendicular distance from the origin to the capping line) is chosen to allow the capping line to intercept any bounded facet at most in a single point. This procedure is denoted as *tangential capping*.

To grasp the biochemical meaning of the capping, note that the ray vector in fact represents the single cycle created in the network of Figure when reaction 3 was made reversible. That is shown by

the calculated 10D ray vector {0, 0.57735, -0.57735, 0, 0.57735, 0, 0, 0, 0, 0} i.e., only fluxes F2, F3 and F5 participate equally.

Such a cycle has no net consumption/production of metabolites, so it can support arbitrarily large flux values without violating limits such as the maximum uptake imposed in the scenario under discus-

Figure 5. The RSS (light blue shading) and SSK (darker shading) for the toy model with a single reversible reaction. The RSS has two bounded facets, AB and AD. The other two parallel facets remain unbounded and leaves a single ray direction (green arrow). The SSK is bounded by the capping line BC, perpendicular to the ray and only intersecting the bounded facets tangentially at the single point B. The inscribed circle is shown in a lighter shade, but the (green) centroid of the chosen periphery points (black dots) is taken as the coordinate origin and the axes are aligned with the main chords shown as dashed red arrows.

sion. This means that partitioning off the unbounded section of the RSS by capping simply excludes non-physical cyclical fluxes. The remaining solutions that define the SSK, embody the physically meaningful flux variation as limited by the model constraints.

Note that this partitioning is not unique. The line BD would be a minimal capping line. The key feature is that a viable SSK needs to include all bounded facets (FBF's), which is why FBF's are considered to encompass the physically meaningful constraints that result from the way that individual flux constraints combine across a network. In higher dimensions minimal tangent capping is hard to compute so merely capping all ray directions orthogonally is considered a sufficient strategy.

3.1.6   Scenario 6: Ray space and coincidence capping.

Figure 6. The toy model RSS (unbounded 2D shaded region) and 1D SSK (the line AB) for scenario 6. There are two peripheral ray directions shown as green arrows. The coordinate system of scenario 5 is chosen to facilitate comparison.

This case is similar to scenario 5, but reaction 6 is now also allowed to be reversed. The result, shown in Figure 6 , is that only the single bounded facet AB of scenario 5 survives. The RSS remains 2D, but there are now two peripheral ray directions, aligned with the two unbounded facets. Intermediate rays can be formed by linear combination of these two rays, i.e. they span the *ray space*.

It would be possible to define a 2D SSK for this case by taking a capping line perpendicular to each peripheral ray to form a triangle with AB as one side. However, a better alternative is to notice that there exists a ray that is perpendicular to line AB. Suppose that this ray is used to define a capping constraint, resulting in a capping line parallel to AB. Adjusting the capping radius until the capping line first intersects the FBF (i.e. line AB itself), that only happens when the two lines coincide. At that point, the entire shaded region of the RSS is separated off and only the single, one dimensional SSK defined by line AB remains.

This is an example of *coincidence capping*. In contrast to tangent capping, it gives an SSK of lower dimensions than the RSS.

Regarding the biochemical interpretation, inspection of the two 10D numerical ray vectors shows that they represent fluxes that involve (F2, F3, F5) and (F4, F5, F6) respectively. These are easily recognizable from Figure as overlapping but independent cyclical paths formed when F3 and F6 are reversed.

Rays are, however, not necessarily associated with cycles as happens for the examples shown; e.g., if the toy network contained a linear network path that does not involve the constrained uptake flux F2 or the metabolites that make up the objective, this might also have no upper limit on its flux and thus constitute a ray.

3.1.7   Scenario 7: Coincidence capping of prismatic rays

The RSS in scenario's 5 and 6 are examples of facetted cones, and the rays are described as *conical rays*. However, for the SSK calculation several additional types of ray are distinguished (Verwoerd, 2022a) and the most common type, typically about half of all rays for realistic metabolic models, are prismatic rays. This type can be induced into the toy model by inserting an additional unidirectional reaction into the network, which only involves some newly added metabolites. That would add an additional flux axis F11 to the flux space. If the same constraints are used as in scenario 3, the OS shown in Figure 3 is modified by the addition of a vertical axis, perpendicular to the plane of the paper. The assumed direction of the reaction limits feasible fluxes to positive values. So instead of the 2D triangle shown in the figure, the OS would now be a semi-infinite triangular prism stretching vertically to positive infinity, and bordered by one horizontal constraint plane (the plane of the paper, constraining F11 to positive values) and three vertical constraint planes.

This modification adds a single ray direction to the OS, which is for obvious reasons described as a *prismatic ray*. The distinguishing feature of a prismatic ray is that it is antiparallel to one constraint vector (the one defining the horizontal plane in this example) and orthogonal to all others. Prismatic rays are particularly easy to deal with as they can be coincidence capped. In this case, that would reduce the 3D OS back to the 2D triangular SSK that is shown in Figure 3.

In a similar way, if the constraints of scenarios 5 or 6 are applied to the 11D extension of the toy model, the RSS in both cases become three dimensional semi-infinite prisms, partially unbounded in both horizontal and vertical directions. They each acquires an additional prismatic ray as a result. Once more, this can be coincidence capped, and then tangent or coincidence capping in the horizontal plane respectively produces the same 2D or 1D SSK's as were illustrated by Figures 5 and 6.

3.1.8   Discussion.

The toy model scenarios that were sketched, reproduce in isolation some of the behaviors encountered in full scale metabolic models. But in a realistic model, multiple instances are usually combined, such as the simultaneous occurrence of multiple ray types and capping methods along different dimensions.

Moreover, in a realistic higher dimensional RSS there are hierarchies of facet levels, similar to the boundary planes, edges and vertices of a 3D polytope.

For realistic models the boundaries formed by the original model constraint hyperplanes are invariably unbounded, and bounded facets only occur at much higher facet levels. This requires an extensive computational framework for their detection and utilization to construct appropriate tangent capping. An exponential proliferation of facets at higher levels makes this challenging, but it turns out that with appropriate algorithms it is possible to reduce most models to a fairly low dimensional SSK with a moderately sized accompanying set of ray vectors that suffice to specify the FBA solution space. That is, for example, the case for the red blood cell model discussed in the next section.

3.2   Exploring and Manipulating the SS.

Knowledge of the SSK provides insight into the location, extent and shape of the solution space and can plausibly be exploited to direct cellular metabolism along desired channels. This section introduces some new computational tools for that and demonstrates their use.

Although realistic models are not amenable to visualization, the SSK periphery point calculation yields a representative sample of feasible metabolic states that similarly serves to display the range of variation allowed by the model.

The key idea pursued here is to resample a particular subregion of the SS that corresponds to a metabolic intervention of interest, such as manipulating nutrient availability, gene regulation, etc. Inspection of the resulting new sample yields information about the cellular responses predicted by the model.

Resampling a section of the SSK could be done by mere interpolation of the SSK periphery point set, but the *HyperSampler* function provided in the SSKernel software goes beyond that by also using the ray basis to extrapolate sample points located in SS regions outside the SSK. Execution of the combined interpolation and extrapolation is not entirely straightforward, and details of the algorithm used is provided in the Supplementary Information of this article.

Given a target metabolite or flux on which some bioengineering effort focusses, a further function *TargetVariation* tabulates the mean value and range of variation of the target, observed in the new sample, for each of a chosen list of interventions such as gene knockouts. Examples are shown and interpreted below.

3.2.1   The toy model with *cof* as a target metabolite

Consider the toy model as modified in scenario 5 above, and for illustration suppose that we wish to influence production of the metabolite *cof* (see Figure) by an external intervention. Two tweaks of the model are required. First, in the model as stated the stoichiometry guarantees net zero production of *cof* (as appropriate for its intended role as a cofactor). Increasing the coefficient of the metabolite for reactions 3 and 4 from 1.0 to 2.0 provides for an excess production. Secondly, a sink reaction to absorb this excess is added to the model and its flux defined as the target.

The effects on the target flux of various interventions, chosen to reflect typical cases and including some single gene knockouts, are shown in Table 1.

Table 1.   Variation of target flux in response to flux assignments

ASSIGN		SAMPLE			
Flux	Value	Count	Mean	Lower	Upper
SSK	---	23	0.265	0	∞
F3, F5	0, ≥0.8	12	1.05	0.8	1.5
F3	0.	18	0.869	0	1.5
F6	0.	5	0.621	0.5	∞
F2	1.	20	0.396	0.	1.0
F8	0.	23	0.265	0.	∞
F3, F4	0.5, 0.5	1	0.	0.	0.
F5	0.	9	0.	0.	0.
F7	0.	0	---	---	---

For comparison the first line shows values for the SSK.

Starting at the bottom of the table, blocking the input flux F7 makes the model infeasible as indicated by 0 sample points. Blocking F5 does allow a range of feasible points, but the target flux remains fixed to zero. Fixing F3 and F4 to 0.5 units each, restricts feasibility to a single flux point still with zero target flux value. In contrast, blocking F8 has no discernible effect: the sample count and target flux values remain the same as for the SSK. Fixing F2 to 1.0 produces almost as many sample points, but now the maximum target flux is reduced to a finite value while the mean value is increased. Blocking either F6 or F3 respectively restricts the lower and upper target flux limits, boosting the mean value in both cases. The strongest boosting effect is observed by simultaneously blocking F3 and restraining F5 to a range of values above 0.8 units.

All of these behaviors are easily interpreted by graphically displaying the SS and various flux contours as in Figure 7. The SSK is the same as in Figure , only reoriented for better display, and the unbounded SS now stretches to infinity along the vertical direction. The actual sample points generated are shown as coloured dots for a few of the single fixed flux cases included in the table. Note how these extend beyond the SSK where appropriate, and cover the full range of variation fairly uniformly.

Figure 7: a) SSK with sampled periphery points shown as black dots, and contours of fixed target flux values as dashed lines. b) Fixed flux lines (dotted) and resampled points for some flux assignments: F3 = 0. (green), F2 = 1. (cyan), F5 = 0. (red) and F6 = 0 (blue).

The case of assigning a value range is not explicitly shown, but for example resampling F2 ≤ 1 produces a sample corresponding to the cyan, blue and some of the red points, and in addition some points on the lower right SSK boundary. The points represent all finite SS boundaries but may also include a few interior points inherited from the SSK sample.

Comparing a fixed flux line in Figure 7b with the target flux contours in Figure 7a, the value range shown in Table 1 for the corresponding case is easily reconciled. Such graphic interpretation is not possible for realistic models, but the table still conveys the relevant information.

3.2.2 Resampling a realistic model

The red blood cell (RBC) metabolic model (Bordbar *et al.*, 2011) included as a demonstration example with the SSKernel software is used to illustrate that resampling is viable also for realistic models. This model has 469 fluxes, of which 408 are found to remain fixed leaving 61 fluxes that vary across the SS, but restricted to a 8-dimensional bounded SSK and with 4 ray directions.

Simply inspecting points sampled from the SS can reveal aspects of the metabolism. For example, it is observed that the flux for the water transport sink reaction EX_h2O_e varies between -6.3 and 4.8 units, allowing both import and export of water. What would happen if either of these is ruled out or if no net water transport is allowed?

This is investigated by resampling from the subregions where for that particular reaction respectively either $F \leq 0$, ≥ 0 or the intersection of hyperplane $F = 0$ with the SS. In the latter case, for example, all changes of the remaining mean fluxes from their values over the SSK, are plotted in Figure 8. There is quite a large contrast in the

Figure 8: Mean value changes from SSK values for the variable fluxes, in response to requiring net zero water exchange in RBC model.

observed responses. For reference, the seven reactions where the magnitude of the change exceeded 0.1, are labeled EX_ade_e, EX_adn_e, EX_bilglcur_e, EX_fum_e, EX_h2o2_e, EX_hco3_e and EX_man_e. The biochemical significance of this observation, if any, could perhaps be further explored by sampling from SS subregions that focus attention on subsets of these particular fluxes.

3.2.3 Maximizing RBC oxygen production

To show an example of how choosing a target metabolite could suggest a bioengineering target, consider that the function of the red blood cell is to transport oxygen. Is it possible to enhance this function by knocking out any single gene? To address that question, we identify the oxygen export reaction EX_O2_e in the existing model as the target flux. The broader SSK analysis discussed above has already narrowed down the field from 469 fluxes to the 61 for which variability is compatible with the RBC model constraints.

Table 2. Oxygen export flux for selected single gene knockouts.

ASSIGN		SAMPLE			
Flux	Value	Count	Mean	Lower	Upper
SSK	---	64	2.61	0.	5.
ADEt	0.	64	3.13	0.	5.
UREAt	0.	10	2.77	0.	5.
SBTR_D2	0.	14	2.76	0.	5.
H2Ot	0.	64	2.75	2.32	3.14
EX_H2O_e	0.	64	2.75	2.32	3.14
NH4t3r	0.	17	2.68	0.	5.
PYRt2	0.	5	2.21	0.	5.
O2t	0.	2	0.	0.	0.
L_LACt2r	0.	0	---	---	---

The *TargetVariation* function illustrated above for the toy model, provides a table of predicted outcomes for a systematic trial of each of the 61 knockouts. An extract of the result is shown in Table 2.

There are again some cases that produce an infeasible model or restrict the target flux to zero. Most knockouts leave the range of the target flux unchanged, while either increasing or reducing the mean target flux relative to the SSK value. Two cases stand out, however:

knocking out either the H2Ot or EX-H2o_e reaction restricts the target EX_O2 flux to a fairly narrow, positive value range. Although there are other knockouts such as ADEt that produce a higher average value, this is arguably a more potent intervention since it guarantees a substantial oxygen export flux.

This result is not claimed to be a profound biochemical insight, but clearly demonstrates that interrogation of the SSK using the resampling tools can be useful to identify bioengineering interventions for further study.

For this realistic but still rather small model, it is practical to use a brute force approach of testing all available gene knockouts since less than 10 seconds of computing time was required to trial all 61 cases. For larger and more complex models, this may well become intractable. However, since the method merely requires a list of candidate interventions as input, the application of biochemical knowledge to set up such a list should be able to reduce the task to manageable proportions.

Funding

This work was supported by provision of publication charges by Hunan University and is gratefully acknowledged by the authors.
Conflict of Interest: none declared.

References

Bordbar, A., Jamshidi, N., & Palsson, B. Ø. (2011) iAB-RBC-283: A proteomically derived knowledge-base of erythrocyte metabolism that can be used to simulate its physiological and patho-physiological states. *BMC Systems Biology*, **5**. doi:10.1186/1752-0509-5-110

Jeter, M. W. (1986) *Mathematical Programming : an Introduction to Optimization*, 1 st. New York: Taylor & Francis, Routledge. doi:10.1201/9780203749333

Mahadevan, R., & Schilling, C. H. (2003) The effects of alternate optimal solutions in constraint-based genome-scale metabolic models. *Metab Eng*, **5**, 264–276. doi:10.1016/j.ymben.2003.09.002

Orth, J. D., Thiele, I., Palsson, B. Ø., *et al.* (2010) What is flux balance analysis? *Nat Biotechnol*, **28**, 245–248. doi:10.1038/nbt.1614

Palsson, B. O. (2015) *Systems biology: constraint based reconstruction and analysis*. Cambridge, UK: Cambridge University Press. doi:10.1017/CBO9781139854610

Palsson, B. Ø. (2006) *Systems biology properties reconstructed networks*. Cambridge, UK: Cambridge University Press. Retrieved from https://www.cambridge.org/nz/academic/subjects/life-sciences/genomics-bioinformatics-and-systems-biology/systems-biology-properties-reconstructed-networks?format=HB&isbn=9780521859035

Papin, J., Stelling, J., Price, N., *et al.* (2004) Comparison of network-based pathway analysis methods. *Trends Biotechnol*, **22**, 400–405. doi:10.1016/j.tibtech.2004.06.010

Schellenberger, J., Que, R., Fleming, R. M. T., *et al.* (2011) Quantitative prediction of cellular metabolism with constraint-based models: the COBRA Toolbox v2.0. *Nature Protocols*, **6**, 1290–1307. doi:10.1038/nprot.2011.308

Schilling, C. H., Letscher, D., & Palsson, B. O. (2000) Theory for the systemic definition of metabolic pathways and their use in interpreting metabolic function from a pathway-oriented perspective. *Journal of Theoretical Biology*, **203**, 229–248. doi:10.1006/jtbi.2000.1073

Schuster, S., Dandekar, T., & Fell, D. A. (1999, February 1) Detection of elementary flux modes in biochemical networks: A promising tool for pathway analysis and metabolic engineering. *Trends in Biotechnology*. Elsevier Ltd. doi:10.1016/S0167-7799(98)01290-6

Verwoerd, W. S. (2022a) *The Compact Kernel of a Metabolic Flux Balance Solution Space*. Singapore: World Scientific. doi:10.1142/12821

Verwoerd, W. S. (2024) SSKernel; Mathematica implementation. GitHub.com. doi:10.5281/zenodo.11483838