Genetic analysis of *Mycobacterium abscessus* reveals genomic diversity linked to global plasmid distribution

Kensuke Ohse[1], Atsushi Yoshida[2,3,4], Keisuke Kamada[3,5], Hironobu Kitazawa[6,7], Yusuke Ito[8], Takayo Shoji[9], Kenichiro Watanabe[10], Hiroshi Koganemaru[2], Ken Kikuchi[3] and Masashi Toyoda[1,11, *]

[1]Research Team for aging science (Vascular Medicine), Tokyo Metropolitan Institute for Geriatrics and Gerontology (TMIG), Tokyo, Japan

[2]Department of Infectious Diseases, TMIG, Tokyo, Japan

[3]Department of Infectious Diseases, Tokyo Women's Medical University, Tokyo, Japan

[4]Department of Genetic Diagnosis and Laboratory Medicine, Dokkyo Medical University, Tochigi, Japan

[5]Department of Mycobacterium Reference and Research, The Research Institute of Tuberculosis, Japan Anti-Tuberculosis Association, Tokyo, Japan

[6]Department of Pediatrics, Nagoya University Graduate School of Medicine, Aichi, Japan

[7]Department of Hematology and Oncology, Children's Medical Center, Japanese Red Cross Aichi Medical Center Nagoya First Hospital, Aichi, Japan

[8]Department of Pediatric Infectious Diseases, Hyogo Prefectural Amagasaki General Medical Center, Hyogo, Japan

[9]Department of Infectious Diseases, Shizuoka Children's Hospital, Shizuoka, Japan

[10]Department of Hematology and Oncology, Shizuoka Children's Hospital, Shizuoka, Japan

[11]Lead contact

*Correspondence:mtoyoda@tmig.or.jp

SUMMARY

Genetic epidemiological analysis of mobile genetic elements, such as plasmids, has rarely been carried out in *Mycobacterium abscessus,* regardless of its usefulness in speculating the past contact between bacteria. In this study, through whole genome sequencing analysis of clinical isolates of *M. abscessus* sequentially collected from the same patient, we identified a single nucleotide variant that could cause antibiotic resistance, and three plasmids (pMAB625-1, pMAB625-2 and pMAB625-3) not found in the type strain. We investigated the distribution of plasmids previously identified in *M. abscessus,* including pMAB625 plasmids, in 462 clinical isolates worldwide *in silico*. pMAB625-3 was detected in the largest number of isolates. Furthermore, phylogenetic tree analysis revealed that these plasmids transferred beyond regions and subspecies and acquired unique mutations. These results indicate that transmission of plasmids increases the genomic diversity in *M. abscessus,* and plasmid epidemiology is useful for estimating the past contact between bacteria.

KEYWORDS

Mycobacterium abscessus; Whole genome sequencing; Plasmid; Horizontal gene transfer

	INTRODUCTION

In recent years, the threat of antimicrobial-resistant bacteria has been increasing worldwide, and they could
be responsible for 1.91 million deaths per year by 2050[1]. Hence, research into the mechanisms and routes
of transmission is becoming increasingly important to develop effective countermeasures against
antimicrobial-resistant bacteria. The global incidence of non-tuberculous mycobacterial infections has been
on the rise, presenting a significant challenge due to their increasing prevalence and innate drug resistance,
particularly in *Mycobacterium abscessus* infections[2]. *M. abscessus* is not limited to pulmonary infections
but also serves as the causative agent for various extrapulmonary infections, including bloodstream, skin,
soft tissue, surgical site, and peritoneal dialysis-related infections[3]. Furthermore, it demonstrates resistance
to a wide range of antimicrobial agents.
45	
46	The route of infection of *M. abscessus* remains controversial. It has been assumed that *M. abscessus* is
47	transmitted from the environment, such as soil and water systems, and person-to-person transmission does
48	not occur. However, a few cases of suspected person-to-person transmission have been reported[4–6]. In
49	addition, the existence of "dominant clones", which were isolated in almost all regions and phylogenetically
50	close[6,7], makes the source and transmission routes of the bacteria mysterious[8].
51	
Whole genome sequencing (WGS) analysis and phylogenetic tree analysis of the core genome, which is
the genomic region shared by all isolates, are often conducted to predict the origin and transmission route
of the isolates[9]. However, the accessory genome, which is shared by a certain percentage but not all of the
isolates such as plasmids, prophage, and genomic islands, is made up of approximately 30% of the typical
number of all genes of *M. abscessus*[10]. Therefore, the analysis of the pan-genome, which includes both the
core genome and accessory genome, was performed to compare the genetic relation of all gene sets of *M.*
abscessus[8,11]. Although the pan-genome analysis has the advantage of comparing whole bacterial
genomes, it has the disadvantage of analyzing accessory genomes as gene units, resulting in the loss of
information on the units of mobile genetic elements. Therefore, epidemiological analysis of mobile genetic
element units, such as individual plasmids, may more accurately predict the origin and route of infection of
bacteria.
63	
In this study, we performed the WGS analysis of the clinical isolates that acquired elevated minimum
inhibitory concentrations (MICs) for carbapenem antibiotics in a case of disseminated infection caused by
M. abscessus subsp. *abscessus* (ABS) during antimicrobial therapy. We detected one mutation potentially
responsible for antimicrobial resistance (AMR) and identified three plasmids (pMAB625-1, pMAB625-2 and
pMAB625-3) that type strain did not harbor. Therefore, we analyzed the distribution of plasmids identified
in *M. abscessus* before, including pMAB625 plasmids and found that pMAB625-3 was the most widely
spreading plasmid among the clinical isolates of *M. abscessus* worldwide. Thus, we investigated whether
predicting the past contact between isolates is possible by analyzing single nucleotide variants (SNVs) and
structural variants (SVs) of the plasmids.
73	

RESULTS

The clinical isolates had decreased susceptibility to carbapenem antibiotics compared to the type
strain

We sequentially isolated five ABS strains from one patient who had osteomyelitis over six months (Table
1). We first compared the growth rate and antimicrobial susceptibility of these clinical isolates with the type
strain (ATCC 19977). The growth rates of Isolate57625 and Isolate57626 were significantly slower than the
type strain (Figure S1). Antimicrobial susceptibility testing showed that the MIC of Isolate57626 against IPM,
which binds the penicillin-binding proteins (PBPs) and inhibits the bacterial cell wall synthesis, was higher
than that of the type strain (Table 1). The MICs of Isolate57625 and Isolate57626 against meropenem
(MEPM), which has the same mechanism of action as IPM and improved stability, were higher than that of
the type strain (Table 1). However, there were no differences in MIC between ATCC 19977 and two isolates
against the other antibiotics. These results indicated the decreased susceptibility of the isolates to
carbapenem antibiotics compared to the type strain.

WGS of the clinical isolates

To explore the factors determining the difference in growth rate and antimicrobial susceptibility between the
type strain and the clinical isolates, we performed WGS using short-read sequencer and Nanopore long-
read sequencer. We analyzed the *de novo* assembled chromosome sequences and found that two large
insertions (13 kbp and 8 kbp) and a deletion (16 kbp) existed in the sequence of Isolate57625 (Figure S2A,
B). The coding sequences of phage integrase and transposase were present near the three large SVs
(Figure S3), indicating that these regions were prophage and transposon. We also analyzed the SNVs in
the chromosome (Table S3), however, we could not identify the cause of the phenotypic differences.
Phylogenetic analysis showed that the isolates were genetically close to the type strain (Figure S2C).
Furthermore, to analyze the difference in the chromosome sequence between five clinical isolates, we
analyzed SVs and SNVs between the isolates (Figure S4). This analysis suggested one mutation located
at the putative *PbpA*, which was associated with the synthesis of the bacterial cell wall[12] and the target of
carbapenem antibiotics[13], may be responsible for the reduced susceptibility of Isolate57626 to IPM
compared to Isolate57625.

The isolates had three plasmids that encode various virulence factors

WGS analysis also revealed that the clinical isolates have three plasmids not found in the type strain
(named pMAB625-1, pMAB625-2 and pMAB625-3, respectively) (Table 2). Thus, we next focused on the
pMAB625 plasmids and compared the pMAB625 plasmid sequences with other plasmids registered to
PLSDB[14] previously (Table 3). A taxonomic keyword search for "*abscessus*" returned 20 plasmids. To
investigate the homology between these plasmids, we calculated ANI between them and pMAB625. No
plasmid had a high ANI for pMAB625-1, however, the ANI between pMAB625-2 and pGD21-2 was 97.4%
with some SVs. Furthermore, the ANI for pMAB625-3 against pGD69A-1 and pGD42-1 was 100%. These
results indicated that pMAB625-2 and pMAB625-3 were retained in the other clinical isolates of *M.*
abscessus. Next, we performed qPCR using primers specific to each plasmid to verify the copy numbers
of pMAB625 plasmids in the clinical isolates. This analysis showed that the copy number of pMAB625-1
and pMAB625-2 was less than one per bacterium, and that of pMAB625-3 was more than five per bacterium
in the isolates (Figure S5). Therefore, the clinical isolates were considered to be a heterogeneous
population in terms of plasmid possession.

We then investigated the function of the genes encoded on pMAB625 plasmids using homology search.
Plasmids encode various factors that influence host characteristics. Therefore, we listed the lost and
acquired genes in Isolate57625 compared to the type strain to investigate the factors affecting antimicrobial
susceptibility (Table S4). Of note, 480 candidate open reading frames (ORFs) were acquired and 37
candidate ORFs were lost in Isolate57625 compared to the type strain. We then searched the AMR genes
from several databases using Blast+ based on the amino acid sequences of the acquired candidate ORFs.
This analysis showed that the coding sequence (CDS) named Isolate57625_54550 had a similarity to *qacA*,
a subunit of the qac multidrug efflux pump[15], (identity, 31 %; *e*-value, 1.69e-10; coverage, 16.1%) and that
the CDS named Isolate57625_54800 had a similarity to *MexL*, a specific repressor of *MexJK* multidrug
efflux system[16] (identity, 33%; *e*-value, 5.54e-4; coverage, 34.4%). However, the identity and coverage
were not sufficiently high.

To gain further insight into the effect of gene acquisition, we searched the homology of the acquired
candidate ORFs using the mycobacterial gene database of Mycobrowser[17]. This analysis detected 37
mycobacterial proteins with amino acid homology to candidate ORFs of the isolates (identity > 30 %, *e*-
value < 1e-50) (Table S5). Notably, genes associated with the ESX secretion system were highly enriched

(10/37, 27%). ESX secretion systems are encoded in various bacteria on their chromosomes and plasmids.
It has been reported that the ESX secretion system (also known as the type VII secretion system) is involved
in bacterial virulence through several mechanisms, including evasion of host immune system[18,19], and
biofilm formation[20]. We further searched whether other ORFs of the components of the ESX secretion
system were encoded near the loci of the detected ORFs of the components of the ESX secretion system
using blast+. We detected several components of the ESX secretion system in pMAB625-1 and pMAB625-
2 (Figure 1A). Core genes that construct membrane pores (*eccB, eccC, eccD, mycP,* and *eccE*) were
present in these gene clusters. The type strain and clinical isolates have ESX-3 and ESX-4 loci on their
chromosomes; therefore, the isolates acquired two additional ESX loci by the plasmid acquisition. To predict
the function of these genes, we performed phylogenetic tree analysis using known amino acid sequences
of the ESX secretion system components in chromosomes and plasmids. It was previously reported that
plasmid-encoded ESX loci can be classified into 4 major clusters[21]. Importantly, our analysis revealed that
plasmid-encoded ESX loci were classified into 5 major clusters (ESX-P cluster 1-5) (Figure 1B). Notably,
the newly classified ESX-P cluster 5 contained the ESX loci of pMAB625-1 and pMAB625-2. As shown in
the results of the plasmid homology search (Table 3), the ESX loci of pGD21-2 and pMAB625-2 were
genetically close. The ESX loci of pMAB625-1 were genetically close to that of pMFLV01, which was
harboured by *Mycobacterium gilvum* PYR GCK. It was previously reported that pMFLV01 was enriched in
biofilm formed in household water purifiers by metagenomic analysis[22]. Biofilm formation is one of the drug
resistance mechanisms in mycobacterium and acts as a barrier against antibiotics. We also detected the
toxin-antitoxin (TA) system components, YefM/YoeB and VapB5/VapC5 in pMAB625-1. The TA system is
present in various bacteria and archaea and encoded on their chromosomes and plasmids. It contributes
to plasmid persistence, inhibiting bacteriophage propagation, and increases antimicrobial tolerance by
inducing dormancy[23]. These results indicated that the clinical isolates acquired factors that could contribute
to bacterial virulence through plasmid transfer.

Gene expression analysis on the pMAB625 plasmids in the clinical isolates

We found that the various genes associated with bacterial virulence were encoded on the pMAB625
plasmid. However, it was unclear whether genes encoded on these plasmids were expressed in the isolates
and to what extent gene expression on the plasmids influenced the expression of genes on the
chromosome. To address this question, we analyzed the global gene expression in the type strain and the
isolates using RNA-seq analysis (Figure 1C and Table S6). The expression levels of the genes on plasmids
were low to moderate (below the 75th percentile of the distribution of expression levels of all genes that
were expressed in the isolates and the type strain). Notably, the expression levels of the most genes (19/27
genes) associated with the ESX secretion system, which were encoded on the pMAB625-1 and pMAB625-
2, were moderate (below the 75th percentile and above the 25th percentile) (Figure 1C). In addition, the
genes associated with the TA system and the mycobacterial membrane protein large (MMPL) family
transporter, which were also encoded on the pMAB625-1 and pMAB625-3, respectively, were expressed
at moderate levels. These results indicated that the numerous genes associated with bacterial virulence
encoded on the plasmids were expressed at moderate levels in the clinical isolates, despite the low number
of copies per bacterium for the pMAB625-1 and pMAB625-2. Next, in the clinical isolates, 393 genes were
highly expressed and 29 genes were low expressed compared to the type strain. Of these differentially
expressed genes, 87.2% of genes (368 highly expressed genes) were acquired genes by SVs or plasmids,
and 12.8% of the genes (25 highly expressed genes and 29 low expressed genes) were present on the
chromosomes of both the isolates and the type strain (Figure 1C). These results indicated that the gene
expression on the plasmid could directly or indirectly affect the expression on the chromosome but to a
lesser extent.

Figure 1. Phylogenetic and gene expression analysis of the genes related to the ESX secretion system

(A) The loci of the ESX secretion system are encoded on pMAB625-1 and pMAB625-2. CDSs of the ESX secretion system were searched using Blast+. The core components that construct membrane pore (*eccB, eccC, eccD, mycP,* and *eccE*) are contained in this operon.

(B) Phylogenetic tree analysis of various ESX loci encoded on chromosomes and plasmids. The amino acid sequence of the genes that construct the membrane pore were concatenated and the phylogenetic tree was created by NGphylogeny.fr[24] using the PhyML+SMS inference method. The classification of each cluster was followed as previously reported[21].

(C) RNA-seq analysis of the type strain and the isolates. The x-axis represents the mean log expression levels, and the y-axis of the top plot represents the log fold change. The bottom dot plot and box plot represent the distribution of the expression levels of each gene. The green dots represent the data of genes that the isolates acquired by SVs and plasmid acquisition. The red dots represent the data of ESX secretion system genes encoded on the plasmids.

pMAB625 plasmids were globally distributed in the clinical isolates of M. abscessus

So far, we have found that the isolates carried three plasmids that the type strain did not, that these plasmids encoded the genes that could contribute to bacterial virulence, and that these genes also expressed in the

isolates. However, we had analyzed the strains sequentially isolated from only one patient and it was uncertain whether the plasmid acquisition occurred extensively in clinical isolates of mycobacterium. We therefore investigated whether other clinical isolates of mycobacterium carry the pMAB625 plasmids. We first performed PCR using the genomic DNA of the clinical isolates collected in Japan[25] with primers specific to each pMAB625 plasmid. We detected pMAB625-3 in two isolates other than Isolate57625, although we did not detect pMAB625-1 and pMAB625-2 in any isolates other than Isolate57625 (Figure 2A). Both the isolates in which pMAB625-3 was detected were *M. abscessus*. We then verified the plasmid possession by mapping the read data from the short-read sequencer to the pMAB625 plasmid sequences. We used the read data of *M. abscessus* clinical isolates in Japan[7] obtained from the NCBI database in addition to our original data. Out of 125 clinical isolates, pMAB625-2 was detected in four isolates (3.2%) and pMAB625-3 was detected in 16 isolates (12.8%) (Figure 2B). These isolates were collected from the four prefectures in Japan (Tokyo, Shizuoka, Aomori, and Okinawa) (Figure 2C). We also mapped the read data of the other mycobacterium species (three of *Mycobacterium iranicum*, 20 of *Mycobacterium fortuitum*, and 25 of *Mycobacterium chelonae*), however, no pMAB625 plasmid was detected in these isolates (data not shown).

Figure 2. The distribution of pMAB625 plasmid in the clinical isolates in Japan
(A) PCR analysis of the possession of pMAB625 plasmid in the clinical isolates which our group isolated in Japan. PCR was performed using the primers specific to each plasmid and *HrpA*. The color bar represents the species of each isolate. The primer sets of *HrpA* were different depending on the species.

220   (B) Mapping the read data of the clinical isolates in Japan to Isolate57625 genome sequence. The bar
221   plot represents the depth of coverage to chromosome and each plasmid and the dot plot represents the
222   percentage of covered genome region.
223   (C) Geographical map of Japan. The prefectures in which the pMAB625 plasmids were detected are
224   represented by color.
225

Moreover, to investigate whether clinical isolates in other areas of East Asia possess pMAB625 plasmids,
we mapped the available read data about the clinical isolates from Taiwan[7] and China[26]. pMAB625-2 was
detected in the isolates from China (2/69, 2.9%) and pMAB625-3 was detected in the isolates from Taiwan
(8/98, 8.2%), and China (5/69, 7.2%) (Figure S6). Two plasmids were detected in ABS and *M. abscessus*
subsp. *massiliense* (MAS) but not in *M. abscessus* subsp. *bolletii* (BOL). In addition, we analyzed the clinical
isolates in Spain, the UK, and the US to determine whether these isolates carried the pMAB625 plasmids
(Table 4). pMAB625-2 plasmid was detected in isolates from the UK (2/30, 6.7%), and the US (2/110, 1.8%).
pMAB625-3 plasmid was detected in the isolates from Spain (9/30, 30%), the UK (1/30, 3.3%), and the US
(14/110, 12.7%). Overall, pMAB625-1 was detected in one isolate (1/462, 0.2%), pMAB625-2 was detected
in 10 isolates (10/462, 2.2%), and pMAB625-3 was detected in 53 isolates (53/462, 11.5%).

As previous analyses have shown that the pMAB625 plasmids are retained in many clinical isolates, we
investigated to what extent other plasmids are retained in the clinical isolates. First, we searched the
plasmid sequence in PLSDB as the taxonomic name for "*abscessus"* and got 20 plasmid sequences (Table
3). We removed three plasmids (pGD42-1, pGD69A-1 and FDAARGOS_1640_plasmid) because they were
identical to other plasmids with no SV. We selected 20 plasmids shown in Table 3 as "TRUE" for "USE for
Reference" together with three pMAB625 plasmids. Then, we mapped the read data of the clinical isolates
to these plasmid sequences. Except for the pMAB625 plasmid, three plasmids (pGD69A-2, pGD22-2, and
pGD42-2) were detected in more than 10 isolates (Table 4). Seven plasmids (pGD21-1, pGD21-2, pGD22-
1, pGD25-3, pJCM30620_2, BRA100 and 50594_plasmid1) were not detected in our dataset. Statistical
analysis showed regional differences in the frequency of appearance of these plasmids (Table 4). pGD69A-
2 was detected more frequently in the US (9/110, 8.2%) than in Japan (0/125, p=0.014). pGD22-2 was
detected more frequently in Spain (3/40, 13.3%) than in Japan (0/125, p=0.018) and Taiwan (0/98, p=0.036).
pGD42-2 was detected more frequently in Spain (4/30, 13.3%) than in Japan (0/125, p=0.017). pGD42-2
was also detected more frequently in the US (12/110, 10.9%) than in Japan (0/125, p=0.001), China (0/69,
p=0.045), and Taiwan (1/98, 1.0%, p=0.041). pGD25-1 was detected more frequently in the UK (4/30,
13.3%) than in Japan (0/125, p=0.018) and in Taiwan (0/98, p=0.036). Moreover, we performed a
phylogenetic tree analysis of the chromosomes to determine whether these plasmids are horizontally
transferred or whether the same strains harboring the plasmids are spreading (Figure 3). The phylogenetic
tree analysis showed that pGD22-2 and pGD42-2 were only detected in the isolates belonging to Clade756.
In contrast, pMAB625-3 was detected mainly in the isolates belonging to the four clades (Clade595, 674,
791 and 813). pGD25-1 and pGD69A-2 were detected mainly in the isolates belonging to Clade678 and
Clade813. pMAB625-2 and pGD19 were also detected in the isolates belonging to several clades. These
results indicated that some plasmids, such as pMAB625-3, were spread across clades and others were
restricted to the specific clade.

Figure 3. The relation between the phylogenetic distance of host bacterial chromosomes and plasmid distribution in clinical isolates worldwide

The phylogenetic tree was created by Gubbins[27] using chromosome sequences and visualized by ggtree[28]. The read data of the clinical isolates collected worldwide were mapped to pMAB625 and known plasmid sequences to verify the plasmid distribution. Tree-scale represents the number of SNVs per genome.

Phylogenetic and structural variant analysis of pMAB625 plasmids enabled the strict classification of isolates and prediction of past inter-subspecies contact

So far, we acquired the sequence information for pMAB625-3, which is retained in 53 different clinical
isolates, and for pMAB625-2, which is retained in 10 different clinical isolates. Therefore, we examined the
genetic relation between the plasmids and investigated whether it could predict host origin and route of
transmission. At first, we performed a phylogenetic tree analysis of 53 sequences of pMAB625-3 (Figure
4A). Forty-two sequences (covered by a yellow box in Figure 4A) belonged to the same clade of the
sequence of pMAB625-3 in Isolate57625. However, 11 sequences belonged to the different clades. In
particular, the pMAB625-3 sequence of SRR7800511 (US, Clade595) was genetically distant from other
pMAB625-3 sequences in the isolates belonging to Clace595. The pMAB625-3 sequence of DRR317373
(Japan, Clade791) was also genetically distant from other pMAB625-3 sequences in the isolates belonging
to Clade791. These results suggested that plasmid epidemiology, when combined with core-genome
analysis, could accurately predict the classification of clinical isolates. In addition, the verification of the
mapping results also showed that pMAB625-3 sequences in the six isolates (DRR317373, Japan;
SRR18969469, China; SRR7800518, US; SRR7800703, US; SRR7800409, US; SRR7800511, US) were
heterogeneous because the unmutated reads and mutated ones were mixed (Figure 4B). Thus, we ran
freebays with the parameter "--ploidy 2" to detect SNVs in these mixed reads. Interestingly, the variant
hotspot region was located at the CDS of the MMPL family transporter, which was the most reliably highly
expressed gene in Isolate57625 and the other four isolates in the RNA-seq analysis. The variant pattern in
the MMPL family transporter was classified into three types (Figure 4B). Four sequences (DRR317373,
Japan; SRR7800518, US; SRR7800703, US; SRR7800409 US) were classified as Type A and one
sequence each was classified as Type B (SRR18969469, China) and Type C (SRR7800511, US), the other
47 sequences had no mutation. There were three missense mutations and 14 or 15 silent mutations in Type
A, 20 missense mutations and 43 silent mutations in Type B, and 47 missense mutations and 126 silent
mutations in Type C (Table S7). Surprisingly, despite many mutations, no mutation was responsible for the
frameshift and nonsense mutations. It was therefore speculated that the mutant MMPL family transporters
were being translated into full-length proteins in these isolates. Thus, these data indicated that pMAB625-
3 was spread across strains and regions, each acquiring unique mutations and some clinical isolates
retaining pMAB625-3 expressed mutated MMPL family transporters.

Figure 4 Phylogenetic tree and SNVs analysis of pMAB625-3

(A) Phylogenetic tree of the sequences of pMAB625-3 detected in the 53 isolates. The phylogenetic tree was created using the maximum-likelihood method using iQ-TREE2 based on the core-SNVs data. The sequences belonging to the clade covered by yellow color were identical, although the sequence of SRR18969469 was heterogeneous. The classification of "Clade (Chromosome)" is based on the clade in

the phylogenetic tree indicated in Figure 3. pMAB625-3 sequences of the isolates covered by yellow box
were identical except for SRR18969469, whose sequence of MMPL family transporter was mixed.

(B) The classification of the pMAB625 sequences based on the SNVs. The mapping results of the reads
data from each isolate to pMAB625-3 were also visualized by IGV. The mutation hotspot was located at
the MMPL family transporter (surrounded by a black line square). The mutation types of the MMPL
transporters were classified into three types, and all mutations at this region were missense or silent.

Then, we classified pMAB625-2 sequences in terms of SVs. The sequences of pMAB625-2 were classified
into four types (Figure 5A). Assuming that Type I, which was the sequence of pMAB625-2 in Isolate57625,
was the base sequence, Type II and III had the large SVs at Region B, and Type II but not Type III had at
Region C. In contrast, Type IV had large SVs at Region A and C but not at Region B. Type I, Type III, and
Type IV were detected in ABS, and Type II was detected in MAS, except for ERR3198514. In addition,
Type III was detected only in the isolates collected in China, and Type IV was detected only in the isolates
collected in the US. All pMAB625-2 sequences retained the ESX locus. These results indicated that
pMAB625-2 had been horizontally transferred between subspecies and had acquired the SVs unique to
each subspecies and region. Next, we analyzed the relation between the phylogenetic distance of hosts'
chromosome sequences and pMAB625-2 type (Figure 5B). The distribution of the pMAB625-2 type was
almost consistent with the phylogenetic relation of hosts' chromosomes. However, the pMAB625-2 type of
Isolate57625 and ERR3198415 were different from those of the other isolates belonging to the same clade
of phylogenetic tree of hosts' chromosomes. Interestingly, Type II was detected in four MAS isolates but
also in ERR3198415 whose subspecies was ABS. This result indicated the possibility of direct or indirect
contact between ERR3198415 and the other four MAS isolates in the past, especially with ERR3198440
isolated from the same region.

Finally, we investigated the distribution in our dataset regarding the plasmids identified in other NTMs
except for *M. abscessus* to verify whether plasmids transferred between *M. abscessus* and other NTM
species. Recently, Wetzstein *et al*. reported clinical and genomic features in Europe about *Mycobacterium*
avium complex (MAC)[29], the most frequently detected NTM globally[30,31]. In the report, they investigated the
distribution of the known 152 plasmids registered in PLSDB and showed that plasmids identified in *M.*
abscessus, including pGD69B-1 (identical to pGD69A-1) and pGD42-1, which were identical to pMAB625-
3, were not detected in their dataset. Therefore, we investigated the distribution of these 152 plasmids in
our dataset using our methods in this study. As a result, we detected the same plasmids indicated in Table
3 but did not other plasmids identified in other NTM except for pMN23 (accession number: NC_010604.1)
and pMFLV03 (accession number: NC_009341.1) in two and one isolates, respectively (Table S8). pMN23
was almost identical to pMAB23 (ANI: 99.9%), so pMN23 was detected redundantly in ATCC 19977 and
SRR7800408, which had pMAB23. Interestingly, pMFLV03 was detected in ERR3198408 (covered
genome region: 78.7%, depth of coverage: 73.1x). pMN23 had originally identified in *Mycobacterium*
marinum[32] and pMFLV03 had identified in *M. gilvum* PYR-GCK (according to the GenBank registry
information), indicating that these plasmids could be present across NTM species. These results suggested
that plasmid distribution is strictly different from *M. abscessus* and other NTMs except for very few cases.

Figure 5. Classification of pMAB625-2 by structural variants and relation with the phylogenetic distance of host bacterial chromosomes

(A) The mapping results of the reads data from each isolate to pMAB625-2 were visualized using IGV. Each sequence was classified into 4 types by the pattern of structural variants (surrounded by black line squares).

(B) Relation between the phylogenetic distance and pMAB625-2 distribution.

DISCUSSION

Plasmids are mobile genetic elements that can move between different bacterial strains and alter host characteristics, such as virulence, and antimicrobial susceptibility. Plasmids are horizontally transferable among bacteria by transformation and conjugation. Thus, plasmid epidemiology, in conjunction with core-

genome analysis, may help to infer past contact between bacteria. In this study, we showed that the
multidrug-resistant clinical isolates of ABS isolated from a patient with osteomyelitis had three plasmids
(pMAB625-1, pMAB625-2 and pMAB625-3) that the type strain did not harbor. Additionally, analysis of the
clinical isolates of *M. abscessus* identified worldwide indicated that pMAB625 plasmids were horizontally
transferred between regions and subspecies and that it is sometimes possible to predict past contact
between bacteria by tracing the phylogenetic relation of plasmids. Dedrick *et al*. reported that pGD69-1
(pGD69A-1), identical to pMAB625-3, is the same as the other five plasmids[33]. Lewin *et al*. identified pMabs-
09-13 through WGS of the clinical isolates of *M. abscessus* within cystic fibrosis patients and reported that
pMabs-09-13 exhibits an identity of 99.86% to pGD42-1, pGD69A-1, and pGD69B-1[34]. These reports
suggested that pMAB625-3 is spreading among many other isolates of *M. abscessus*, but the detailed
distribution of pMAB625 plasmids in the world and other subspecies has remained unclear. Our study
shows that pMAB625-3 is the plasmid spreading beyond regions and subspecies in the largest number of
clinical isolates of *M. abscessus*. This allowed us to investigate the possibility of speculating the precise
classification of bacteria and the past contact between them. The phylogenetic tree analysis of pMAB625-
3 based on the SNVs of plasmids (Figure 4A) indicated that the SNV pattern was correlated with the
phylogenetic relation of the hosts' chromosome sequences; however, a unique variant (SRR7800511, US)
different from the other sequences in the clade was found. This indicates that it is possible to predict the
bacterial contact using the distribution of the unique variant like SRR7800511. In the case of pMAB625-2,
it is actually indicated that the past contact between the different subspecies is predictable based on the
relation between the phylogenetic tree of the hosts' chromosomes and the distribution of pMAB625-2 type.

As important findings, we found that 2.2% (10/462) and 11.5% (53/462) of the clinical isolates worldwide
had pMAB625-2, and pMAB625-3, respectively (Table 4). This result provides an epidemiological insight
into the spread of plasmids in *M. abscessus* beyond regions and strains. Other than person-to-person
transmission of *M.abscessus*, this global spread of pMAB625 plasmids may be mainly due to increased
global logistical activity and human mobility with associated interference of human and microbial habitats[35].
The habitat of *M. abscessus* is soil and water systems, and it is also possible that increased flooding due
to recent climate change may have increased the chance of contact between the different strains. Analysis
of plasmid distribution revealed that some plasmids (pMAB625-2, pMAB625-3, pGD19, pGD25-1 and
pGD69A-2) have been widely spread between strains, regions and subspecies, while others (pGD22-2 and
pGD42-2) have been restricted to specific strains and regions (Figure 3). Therefore, it is speculated that
regarding pGD22-2 and pGD42-2, the identical isolates retaining these plasmids had been spread within
or between the regions, whereas, regarding pMAB625-2, pMAB625-3, pGD19, pGD25-1 and p69A-2,
plasmids themselves had been spread between different strains, subspecies and region through horizontal
transfer. It is unknown what factors determine these differences in plasmid distribution. It is possible that
the factors encoded on the plasmid or the periods from the emergence were different among the plasmids.
In addition, we also showed the plasmid distribution was different from *M. abscessus* and MAC (Table S8),
indicating that incompatibility also exists in the plasmids present in NTM, strictly distinguishing *M.*
abscessus from other NTMs except for very few cases, such as pMN23 and pMFLV03.

We identified various factors associated with bacterial virulence and multidrug resistance on the pMAB625
plasmids. The components of the ESX secretion system, the MMPL family transporter, and the components
of the TA system were encoded on the plasmids, and these genes were highly expressed in the clinical
isolates compared to the type strain. The ESX secretion system is involved in the export of the various
pathogenic proteins and metal ions associated with proliferation, survival in the cytosol of macrophage, and
conjugation. It has also been reported that ESX-1 components are required for sliding motility and biofilm
formation in *M. avium*[20]. Therefore, given the close genetic distance between ESX-1 and ESX-P cluster 5,
which include the components of the ESX secretion system on the pMAB625-1 and pMAB625-2 (Figure
1B), the components of ESX-P cluster 5 may be involved in biofilm formation. In addition, MMPL family

transporters can transport antibiotics across the cell membrane. It has been reported that the increased
expression of MMPL5 is associated with decreased susceptibility to bedaquiline, and clofazimine in *M.*
tuberculosis[36,37], bedaquiline, and clofazimine in *M. intracellulare*[38], and thiactazone derivative in *M.*
abscessus[39]. The function of the MMPL family transporter encoded on the pMAB625-3 is unknown;
however, the presence of mutational hotspots (Figure 4B) indicates that this MMPL transporter may be
compatible with a variety of substrates, including antimicrobial agents. Therefore, our results suggest that
Isolate57625 enhanced defence mechanisms against antibiotics through the acquisition of the ESX
secretion system and transporters, accompanied by changes in biofilm barriers, and transport of antibiotics.
These results suggest that pMAB625 plasmids appeared to be a cause of the low susceptibility to
carbapenem antibiotics in the clinical isolates compared with the type strain; however, further studies will
be essential to determine whether these plasmids are the main cause of AMR.

In conclusion, we reported the distribution of plasmids in *M. abscessus* and found that pMAB625-3 is the
most widely distributed plasmid in the clinical isolates of *M. abscessus* worldwide. In addition, phylogenetic
tree analysis and mutational analysis of the pMAB625 plasmids partially predict past bacterial contact.
These findings provide a new perspective on the acquisition of genomic diversity, the origin and
transmission route of *M. abscessus.* These findings will also lead to the development of effective
countermeasures to identify pathways of bacterial transmission and prevent serious bacterial outbreaks.

Limitations of the study

Although it is assumed that phylogenetically close plasmids with similar SNV and SV patterns were spread
via bacterial contact, it cannot be excluded that they may have acquired identical mutation patterns in their
respective clinical isolates without bacterial contact. To clarify this, it is necessary to calculate the mutation
rate of the plasmid and compare the mutation acquisition period with the interval of appearance in the
different strains. In addition, future studies are required to verify whether the pMAB625 plasmids induce
multidrug resistance in *M. abscessus* by plasmid transfection.

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by
the lead contact, Masashi Toyoda (mtoyoda@tmig.or.jp).

Materials availability
This study did not generate new unique reagents.

Data and code availability
• All raw sequence data generated in this study have been deposited in the DDBJ database at
https://www.ddbj.nig.ac.jp/ under the BioProject accession number PRJDB16220. The complete
genome sequences of the isolates are available in the DDBJ database. They can be accessed
using the accession numbers, AP028613, AP028614, AP028615, and AP028616 for Isolate57625;
AP028617, AP028618, AP028619, and AP028620 for Isolate57629; AP028621, AP028622,
AP028623, and AP028624 for Isolate57630; AP028625, AP028626, AP028627, and AP028628 for
Isolate57596; AP028629, AP028630, AP028631, and AP028632 for Isolate57626, for the
chromosome and three plasmids in each isolate.
• The analyzed data of RNA-seq have been deposited in the Genomic Expression Archive at
https://www.ddbj.nig.ac.jp/gea/ under the accession number E-GEAD-827.

• All data reported in this paper will be shared by the lead contact upon request.
• This paper does not report original code.

• Any additional information required to reanalyze the data reported in this paper is available from
the lead contact upon request.
457

ACKNOWLEDGMENTS

459 The authors thank the staff of the microbiology laboratory of Dokkyo Medical University Hospital for
460 performing identification and drug susceptibility tests, and Dr. Yoshishige Masuda and Dr. Takashi
461 Inamatsu at TMIG for supervising in clinical practice. Computations were partially performed on the NIG
462 supercomputer at ROIS National Institute of Genetics. This work was supported by Japan Society for the
463 Promotion of Science (JSPS) KAKENHI Grant Number JP19K08938.

AUTHOR CONTRIBUTIONS

466 Conceptualization,K.O., A.Y.; Formal analysis, K.O.; Methodology, K.O., A.Y., and Hiroshi Koganemaru;
467 Data curation, K.O.; Investigation,K.O., Keisuke Kamada, and Ken Kikutchi ; Validation, K.O.; Writing-
468 original draft, K.O. and A.Y.; Writing-review & editing, K.O., Y.A. and M.T.; Funding acquisition, A.Y. and
469 Ken Kikuchi.; Patient care and resources, Hironobu Kitazawa, Y.I., T.S. and K.W.; Supervision, M.T.

DECLARATION OF INTERESTS

472 The authors declare no competing interests.

SUPPLEMENTAL INFORMATION

Document S1. Figures S1–S6
Table S1. Primer sequences used in this study
Table S2. Genbank accession numbers of the genome and plasmid sequences in this study
Table S3. Gene lists on chromosome which were acquired, lost and mutated in Isolate57625
compared with ATCC19977
Table S4. Gene lists which were acquired, lost and mutated in Isolate57625 compared with
ATCC19977
Table S5. Homology search results of genes on pMAB625 plasmids against genes of
Mycobacterium
Table S6. The results of RNA-seq analysis
Table S7. The mutation type of MMPL family transporter
Table S8. The analysis of distribution regarding NTM plasmids in the clinical isolates of *M.*
abscessus

 FIGURE TITLES AND LEGENDS

Figure 1. Phylogenetic and gene expression analysis of the genes related to the ESX secretion
system
(A) The loci of the ESX secretion system are encoded on pMAB625-1 and pMAB625-2. CDSs of the ESX
secretion system were searched using Blast+. The core components that construct membrane pore
(*eccB, eccC, eccD, mycP,* and *eccE*) are contained in this operon.
(B) Phylogenetic tree analysis of various ESX loci encoded on chromosomes and plasmids. The amino
acid sequence of the genes that construct the membrane pore were concatenated and the phylogenetic
tree was created by NGphylogeny.fr[24] using the PhyML+SMS inference method. The classification of
each cluster was followed as previously reported[21].
(C) RNA-seq analysis of the type strain and the isolates. The x-axis represents the mean log expression
levels, and the y-axis of the top plot represents the log fold change. The bottom dot plot and box plot
represent the distribution of the expression levels of each gene. The green dots represent the data of
genes that the isolates acquired by SVs and plasmid acquisition. The red dots represent the data of ESX
secretion system genes encoded on the plasmids.
518
Figure 2. The distribution of pMAB625 plasmid in the clinical isolates in Japan
(A) PCR analysis of the possession of pMAB625 plasmid in the clinical isolates which our group isolated
in Japan. PCR was performed using the primers specific to each plasmid and *HrpA*. The color bar
represents the species of each isolate. The primer sets of *HrpA* were different depending on the species.
(B) Mapping the read data of the clinical isolates in Japan to Isolate57625 genome sequence. The bar
plot represents the depth of coverage to chromosome and each plasmid and the dot plot represents the
percentage of covered genome region.
(C) Geographical map of Japan. The prefectures in which the pMAB625 plasmids were detected are
represented by color.
528
Figure 3. The relation between the phylogenetic distance of host bacterial chromosomes and
plasmid distribution in clinical isolates worldwide

The phylogenetic tree was created by Gubbins[27] using chromosome sequences and visualized by
ggtree[28]. The read data of the clinical isolates collected worldwide were mapped to pMAB625 and known
plasmid sequences to verify the plasmid distribution. Tree-scale represents the number of SNVs per
genome.

Figure 4 Phylogenetic tree and SNVs analysis of pMAB625-3

(A) Phylogenetic tree of the sequences of pMAB625-3 detected in the 53 isolates. The phylogenetic tree
was created using the maximum-likelihood method using iQ-TREE2 based on the core-SNVs data. The
sequences belonging to the clade covered by yellow color were identical, although the sequence of
SRR18969469 was heterogeneous. The classification of "Clade (Chromosome)" is based on the clade in
the phylogenetic tree indicated in Figure 3. pMAB625-3 sequences of the isolates covered by yellow box
were identical except for SRR18969469, whose sequence of MMPL family transporter was mixed.

(B) The classification of the pMAB625 sequences based on the SNVs. The mapping results of the reads
data from each isolate to pMAB625-3 were also visualized by IGV. The mutation hotspot was located at
the MMPL family transporter (surrounded by a black line square). The mutation types of the MMPL
transporters were classified into three types, and all mutations at this region were missense or silent.

Figure 5. Classification of pMAB625-2 by structural variants and relation with the phylogenetic
distance of host bacterial chromosomes

(A) The mapping results of the reads data from each isolate to pMAB625-2 were visualized using IGV.
Each sequence was classified into 4 types by the pattern of structural variants (surrounded by black line
squares).

(B) Relation between the phylogenetic distance and pMAB625-2 distribution.

Table 1. The isolation time point of the clinical isolates and antimicrobial susceptibility of the type strain and the isolates

Isolates	Source	Period from the first isolation (month)	MICa (µg/mL)														
			AMK	TOB	IPM	FRPM	LVFX	MFLX	AZM	CAM 4day	CAM 14day	ST	DOXY	MEPM	LZD	CLF	STFX
ATCC 19977	-	-	8	8	16	>64	32	>8	64	2	>64	>152/8	>16	64	32	1	2
Isolate57625	catheter	0	8	8	16b	>64	16	>8	64	2	>64	>152/8	>16	>64	32	1	2
Isolate57629	catheter	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Isolate57630	bone biopsy	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Isolate57596	heel	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Isolate57626	malleolus	6	8	8	>64c	>64	32	>8	64	2	>64	>152/8	>16	>64	32	1	2

[a]MICs were determined as the minimum concentration at which the bacteria did not grow. No antimicrobial susceptibility testing was conducted on the isolates other than Isolate57625 and Isolate57626, representing hyphens in the table. The test was repeated three times and all MICs were confirmed to be the same between the experiments with the exception of imipenem for the isolates. AMK, Amikacin; TOB, Tobramycin; IPM, Imipenem; FRPM, Faropenem; LVFX, Levofloxacin; MFLX, Moxifloxacin; AZM, Azithromycin; CAM, Clarithromycin; ST, Trimethoprim/sulfamethoxazole; DOXY, Doxycycline; MEPM, Meropenem; LNZ, Linezolid; CLF, Clofazimine; STFX, Sitafloxacin
[b]For Isolate57625, the MICs against IPM in the three experiments were 64, 16, and 16 µg/mL, respectively.
[c]For Isolate57626, the MICs against IPM in the three experimenst were >64, 16, and >64 µg/mL, respectively.

Table 2. Genome assembly and the annotation statics of the Isolate57625

Contig	Form	Length(bp)	GC contents (%)	Number of CDS	Number of rRNA	Number of tRNA	Number of tmRNA
Chromosome	circular	5,072,346	64.1	5,034	3	48	1
pMAB625-1	linear	202,398	64.7	247	0	29	0
pMAB625-2	linear	142,689	62.2	171	0	0	0
pMAB625-3	circular	24,988	65.6	30	0	0	0

Table 3. Homology search result between plasmids identified in *M. abscessus*

Plasmid	Accession no.	Topology	Length (bp)	Homologous plasmid (ANI(%)>99)	Homologous plasmid (97< ANI(%) <99)	Use for "Reference"
pMAB625-1	AP028614.1	linear	202,398	-	-	TRUE
pMAB625-2	AP028615.1	linear	142,689	-	pGD21-2b	TRUE
pMAB625-3	AP028616.1	circular	24,988	pGD69A-1, pGD42-1	-	TRUE
pMAB23	NC_010394.1	circular	23,319	pMabS_GZ002a, FDAARGOS_1604_plasmid	-	TRUE
pMabS_GZ002	NZ_CP034180.1	linear	15,203	pMAB23, FDAARGOS_1604_plasmid	-	TRUE

pGD19	NZ_CP063329.1	circular	18,605	-	-	TRUE
pGD21-1	NZ_CP065285.1	circular	112,633	-	-	TRUE
pGD21-2	NZ_CP065286.1	linear	155,609	-	pMAB625-2[b]	TRUE
pGD22-1	NZ_CP063325.1	circular	19,694	-	-	TRUE
pGD22-2	NZ_CP063326.1	circular	18,117	pGD42-2[a]	-	TRUE
pGD25-1	NZ_CP063321.1	circular	31,413	-	-	TRUE
pGD25-2	NZ_CP063322.1	circular	27,424	-	-	TRUE
pGD25-3	NZ_CP063323.1	circular	23,599	-	-	TRUE
pGD42-1	NZ_CP065281.1	circular	24,993	pMAB625-3, pGD69A-1	-	FALSE
pGD42-2	NZ_CP065282.1	circular	9,547	pGD22-2	pGD69A-2	TRUE
pGD69A-1	NZ_CP065270.1	circular	25,000	pMAB625-3, pGD42-1	-	FALSE
pGD69A-2	NZ_CP065271.1	circular	18,611	-	pGD42-2[a]	TRUE
pJCM30620_1	NZ_AP022622.1	linear	119,389	50594_plasmid2[a]	-	TRUE
pJCM30620_2	NZ_AP022623.1	linear	38,763	50594_plasmid1[b]	-	TRUE
BRA100	NC_017908.2	circular	56,265	-	-	TRUE
FDAARGOS_1604_plasmid	NZ_CP085920.1	circular	23,319	pMAB23, pMabS_GZ002[a]	-	FALSE
50594_plasmid1	NC_021278.1	circular	172,814	pJCM30620_2[a]	-	TRUE
50594_plasmid2	NC_021279.1	circular	97,240	pJCM30620_1[b]	-	TRUE

572   [a]One Structural Variant (SV) exists in the sequence.

573   [b]More than two SVs exist in the sequence.

575 **Table 4. pMAB625 and known plasmids distribution in the clinical isolates worldwide**

	Number (%) of isolates retaining plasmid							
Region	Unknown (Type strain)	Japan	China	Taiwan	US	Spain	UK	total[a]
Analyzed isolates	3	125	69	98	110	30	30	462
pMAB625-1	0	1 (0.8)	0	0	0	0	0	1 (0.2)
pMAB625-2	0	4 (3.2)	2 (2.9)	0	2 (1.8)	0	2 (6.7)	10 (2.2)
pMAB625-3	0	16 (12.8)	5 (7.2)	8 (8.2)	14 (12.7)	9 (30.0)	1 (3.3)	53 (11.5)
pMAB23	1 (33.3)	0	0	0	1 (0.9)	0	0	1 (0.2)
pMabS_GZ002	1 (33.3)	0	0	0	1 (0.9)	0	0	1 (0.2)

pGD69A-2	0	0	0	1 (1.0)	9 (8.2)[b]	2 (6.7)	3 (10.0)	15(3.2)
pGD19	0	0	4 (5.8)	3 (3.1)	0	1 (3.3)	1 (3.3)	9(1.9)
pJCM30620_1	0	0	1 (1.4)	0	0	0	1 (3.3)	2(0.4)
50594_plasmid2	0	0	1 (1.4)	0	0	0	1 (3.3)	2(0.4)
pGD22-2	0	0	0	0	6 (5.5)	4 (13.3)[c]	1 (3.3)	11(2.4)
pGD42-2	0	0	0	1 (1.0)	12 (10.9)[d]	4 (13.3)[b]	1 (3.3)	18(3.9)
pGD25-1	0	0	0	0	3 (2.7)	0	4 (13.3)[c]	7(1.5)
pGD25-2	0	0	0	0	1 (0.9)	0	2 (6.7)	3(0.6)

Fisher's exact tests were performed to verify regional differences in the frequency of appearance of each
plasmid. Corrections for the multiple comparisons were performed using Holm's method.

[a]Total number excludes the number of the type strains.

[b]$p<0.05$, compared with Japan

[c]$p<0.05$, compared with Japan and Taiwan

[d]$p<0.05$, compared with Japan, China and Taiwan

REFERENCES

1.   Naghavi, M., Vollset, S.E., Ikuta, K.S., Swetschinski, L.R., Gray, A.P., Wool, E.E., Aguilar, G.R.,
Mestrovic, T., Smith, G., Han, C., et al. (2024). Global burden of bacterial antimicrobial resistance 1990–
2021: a systematic analysis with forecasts to 2050. The Lancet *404*, 1199–1226.
https://doi.org/10.1016/S0140-6736(24)01867-1.

2.   Johansen, M.D., Herrmann, J.-L., and Kremer, L. (2020). Non-tuberculous mycobacteria and the
rise of Mycobacterium abscessus. Nat Rev Microbiol *18*, 392–407. https://doi.org/10.1038/s41579-020-
0331-1.

3.   Cristancho-Rojas, C., Varley, C.D., Lara, S.C., Kherabi, Y., Henkle, E., and Winthrop, K.L. (2024).
Epidemiology of *Mycobacterium abscessus*. Clinical Microbiology and Infection *30*, 712–717.
https://doi.org/10.1016/j.cmi.2023.08.035.

4.   Bryant, J.M., Grogono, D.M., Greaves, D., Foweraker, J., Roddick, I., Inns, T., Reacher, M.,
Haworth, C.S., Curran, M.D., Harris, S.R., et al. (2013). Whole-genome sequencing to identify
transmission of Mycobacterium abscessus between patients with cystic fibrosis: a retrospective cohort
study. Lancet *381*, 1551–1560. https://doi.org/10.1016/S0140-6736(13)60632-7.

5.   Lipworth, S., Hough, N., Weston, N., Muller-Pebody, B., Phin, N., Myers, R., Chapman, S., Flight,
W., Alexander, E., Smith, E.G., et al. (2021). Epidemiology of Mycobacterium abscessus in England: an
observational study. Lancet Microbe *2*, e498–e507. https://doi.org/10.1016/S2666-5247(21)00128-2.

6.   Bryant, J.M., Grogono, D.M., Rodriguez-Rincon, D., Everall, I., Brown, K.P., Moreno, P., Verma,
D., Hill, E., Drijkoningen, J., Gilligan, P., et al. (2016). Emergence and spread of a human-transmissible
multidrug-resistant nontuberculous mycobacterium. Science *354*, 751–757.
https://doi.org/10.1126/science.aaf8156.

7.   Yoshida, M., Chien, J.-Y., Morimoto, K., Kinjo, T., Aono, A., Murase, Y., Fujiwara, K., Morishige,
Y., Nagano, H., Jou, R., et al. (2022). Molecular Epidemiological Characteristics of Mycobacterium
abscessus Complex Derived from Non-Cystic Fibrosis Patients in Japan and Taiwan. Microbiol Spectr *10*,
e00571-22. https://doi.org/10.1128/spectrum.00571-22.

8.   Davidson, R.M. (2018). A Closer Look at the Genomic Variation of Geographically Diverse
Mycobacterium abscessus Clones That Cause Human Infection and Disease. Front Microbiol *9*, 2988.
https://doi.org/10.3389/fmicb.2018.02988.

9.   Zhang, M., Adroub, S., Ummels, R., Asaad, M., Song, L., Pain, A., Bitter, W., Guan, Q., and
Abdallah, A.M. (2024). Comprehensive pan-genome analysis of Mycobacterium marinum: insights into
genomic diversity, evolution, and pathogenicity. Sci Rep *14*, 27723. https://doi.org/10.1038/s41598-024-
75228-0.

10.   Choo, S.W., Wee, W.Y., Ngeow, Y.F., Mitchell, W., Tan, J.L., Wong, G.J., Zhao, Y., and Xiao, J.
(2014). Genomic reconnaissance of clinical isolates of emerging human pathogen Mycobacterium
abscessus reveals high evolutionary potential. Sci Rep *4*, 4061. https://doi.org/10.1038/srep04061.

11.   Bronson, R.A., Gupta, C., Manson, A.L., Nguyen, J.A., Bahadirli-Talbott, A., Parrish, N.M., Earl,
A.M., and Cohen, K.A. (2021). Global phylogenomic analyses of Mycobacterium abscessus provide
context for non cystic fibrosis infections and the evolution of antibiotic resistance. Nat Commun *12*, 5145.
https://doi.org/10.1038/s41467-021-25484-9.

12.   Sauvage, E., Kerff, F., Terrak, M., Ayala, J.A., and Charlier, P. (2008). The penicillin-binding
proteins: structure and role in peptidoglycan biosynthesis. FEMS Microbiol Rev *32*, 234–258.
https://doi.org/10.1111/j.1574-6976.2008.00105.x.

13.   Hellinger, W.C., and Brewer, N.S. (1991). Imipenem. Mayo Clin Proc *66*, 1074–1081.
https://doi.org/10.1016/s0025-6196(12)61732-7.

14.   Schmartz, G.P., Hartung, A., Hirsch, P., Kern, F., Fehlmann, T., Müller, R., and Keller, A. (2022).
PLSDB: advancing a comprehensive database of bacterial plasmids. Nucleic Acids Research *50*, D273–
D278. https://doi.org/10.1093/nar/gkab1111.

15.   Paulsen, I.T., Brown, M.H., Littlejohn, T.G., Mitchell, B.A., and Skurray, R.A. (1996). Multidrug
resistance proteins QacA and QacB from Staphylococcus aureus: membrane topology and identification
of residues involved in substrate specificity. Proceedings of the National Academy of Sciences *93*, 3630–
3635. https://doi.org/10.1073/pnas.93.8.3630.

16.   Chuanchuen, R., Gaynor, J.B., Karkhoff-Schweizer, R., and Schweizer, H.P. (2005). Molecular
Characterization of MexL, the Transcriptional Repressor of the mexJK Multidrug Efflux Operon in
Pseudomonas aeruginosa. Antimicrob Agents Chemother *49*, 1844–1851.
https://doi.org/10.1128/AAC.49.5.1844-1851.2005.

17.   Kapopoulou, A., Lew, J.M., and Cole, S.T. (2011). The MycoBrowser portal: a comprehensive
and manually annotated resource for mycobacterial genomes. Tuberculosis (Edinb) *91*, 8–13.
https://doi.org/10.1016/j.tube.2010.09.006.

18.   Osman, M.M., Shanahan, J.K., Chu, F., Takaki, K.K., Pinckert, M.L., Pagán, A.J., Brosch, R.,
Conrad, W.H., and Ramakrishnan, L. (2022). The C terminus of the mycobacterium ESX-1 secretion
system substrate ESAT-6 is required for phagosomal membrane damage and virulence. Proceedings of
the National Academy of Sciences *119*, e2122161119. https://doi.org/10.1073/pnas.2122161119.

19.   Laencina, L., Dubois, V., Le Moigne, V., Viljoen, A., Majlessi, L., Pritchard, J., Bernut, A., Piel, L.,
Roux, A.-L., Gaillard, J.-L., et al. (2018). Identification of genes required for Mycobacterium abscessus
growth in vivo with a prominent role of the ESX-4 locus. Proceedings of the National Academy of
Sciences *115*, E1002–E1011. https://doi.org/10.1073/pnas.1713195115.

20.   Lai, L.-Y., Lin, T.-L., Chen, Y.-Y., Hsieh, P.-F., and Wang, J.-T. (2018). Role of the
Mycobacterium marinum ESX-1 Secretion System in Sliding Motility and Biofilm Formation. Frontiers in
Microbiology *9*, 1160.

21.   Dumas, E., Christina Boritsch, E., Vandenbogaert, M., Rodríguez de la Vega, R.C., Thiberge, J.-
M., Caro, V., Gaillard, J.-L., Heym, B., Girard-Misguich, F., Brosch, R., et al. (2016). Mycobacterial Pan-
Genome Analysis Suggests Important Role of Plasmids in the Radiation of Type VII Secretion Systems.
Genome Biology and Evolution *8*, 387–402. https://doi.org/10.1093/gbe/evw001.

22.   Zhou, Z., Xu, L., Zhu, L., Liu, Y., Shuai, X., Lin, Z., and Chen, H. (2021). Metagenomic analysis of
microbiota and antibiotic resistome in household activated carbon drinking water purifiers. Environment
International *148*, 106394. https://doi.org/10.1016/j.envint.2021.106394.

23.   Harms, A., Brodersen, D.E., Mitarai, N., and Gerdes, K. (2018). Toxins, Targets, and Triggers: An
Overview of Toxin-Antitoxin Biology. Molecular Cell *70*, 768–784.
https://doi.org/10.1016/j.molcel.2018.01.003.

24.   Lemoine, F., Correia, D., Lefort, V., Doppelt-Azeroual, O., Mareuil, F., Cohen-Boulakia, S., and
Gascuel, O. (2019). NGPhylogeny.fr: new generation phylogenetic services for non-specialists. Nucleic
Acids Research *47*, W260–W265. https://doi.org/10.1093/nar/gkz303.

25.   Kamada, K., Yoshida, A., Iguchi, S., Arai, Y., Uzawa, Y., Konno, S., Shimojima, M., and Kikuchi,
K. (2021). Nationwide surveillance of antimicrobial susceptibility of 509 rapidly growing mycobacteria
strains isolated from clinical specimens in Japan. Sci Rep *11*, 12208. https://doi.org/10.1038/s41598-021-
91757-4.

26.   Jin, P., Dai, J., Guo, Y., Wang, X., Lu, J., Zhu, Y., and Yu, F. (2022). Genomic Analysis of
Mycobacterium abscessus Complex Isolates from Patients with Pulmonary Infection in China. Microbiol
Spectr *10*, e00118-22. https://doi.org/10.1128/spectrum.00118-22.

27.   Croucher, N.J., Page, A.J., Connor, T.R., Delaney, A.J., Keane, J.A., Bentley, S.D., Parkhill, J.,
and Harris, S.R. (2015). Rapid phylogenetic analysis of large samples of recombinant bacterial whole
genome sequences using Gubbins. Nucleic Acids Research *43*, e15. https://doi.org/10.1093/nar/gku1196.

28.   Yu, G., Smith, D.K., Zhu, H., Guan, Y., and Lam, T.T.-Y. (2017). ggtree: an r package for
visualization and annotation of phylogenetic trees with their covariates and other associated data.
Methods in Ecology and Evolution *8*, 28–36. https://doi.org/10.1111/2041-210X.12628.

29.   Wetzstein, N., Diricks, M., Anton, T.B., Andres, S., Kuhns, M., Kohl, T.A., Schwarz, C., Lewin, A.,
Kehrmann, J., Kahl, B.C., et al. (2024). Clinical and genomic features of Mycobacterium avium complex: a
multi-national European study. Genome Med *16*, 86. https://doi.org/10.1186/s13073-024-01359-8.

30.   Namkoong, H., Kurashima, A., Morimoto, K., Hoshino, Y., Hasegawa, N., Ato, M., and Mitarai, S.
(2016). Epidemiology of Pulmonary Nontuberculous Mycobacterial Disease, Japan. Emerging Infectious
Diseases *22*, 1116. https://doi.org/10.3201/eid2206.151086.

31.   Hoefsloot, W., Ingen, J. van, Andrejak, C., Ängeby, K., Bauriaud, R., Bemer, P., Beylis, N.,
Boeree, M.J., Cacho, J., Chihota, V., et al. (2013). The geographic diversity of nontuberculous
mycobacteria isolated from pulmonary samples: an NTM-NET collaborative study. European Respiratory
Journal *42*, 1604–1613. https://doi.org/10.1183/09031936.00149212.

32.   Stinear, T.P., Seemann, T., Harrison, P.F., Jenkin, G.A., Davies, J.K., Johnson, P.D.R., Abdellah,
Z., Arrowsmith, C., Chillingworth, T., Churcher, C., et al. (2008). Insights from the complete genome
sequence of Mycobacterium marinum on the evolution of Mycobacterium tuberculosis. Genome Res *18*,
729–741. https://doi.org/10.1101/gr.075069.107.

33.   Dedrick, R.M., Aull, H.G., Jacobs-Sera, D., Garlena, R.A., Russell, D.A., Smith, B.E.,
Mahalingam, V., Abad, L., Gauthier, C.H., and Hatfull, G.F. (2021). The Prophage and Plasmid Mobilome
as a Likely Driver of Mycobacterium abscessus Diversity. mBio *12*, e03441-20.
https://doi.org/10.1128/mBio.03441-20.

34.   Lewin, A., Kamal, E., Semmler, T., Winter, K., Kaiser, S., Schäfer, H., Mao, L., Eschenhagen, P.,
Grehn, C., Bender, J., et al. (2021). Genetic diversification of persistent Mycobacterium abscessus within
cystic fibrosis patients. Virulence *12*, 2415–2429. https://doi.org/10.1080/21505594.2021.1959808.

35.   Castañeda-Barba, S., Top, E.M., and Stalder, T. (2024). Plasmids, a molecular cornerstone of
antimicrobial resistance in the One Health era. Nat Rev Microbiol *22*, 18–32.
https://doi.org/10.1038/s41579-023-00926-x.

36.   Andries, K., Villellas, C., Coeck, N., Thys, K., Gevers, T., Vranckx, L., Lounis, N., Jong, B.C. de,
and Koul, A. (2014). Acquired Resistance of Mycobacterium tuberculosis to Bedaquiline. PLOS ONE *9*,
e102135. https://doi.org/10.1371/journal.pone.0102135.

37.   Hartkoorn, R.C., Uplekar, S., and Cole, S.T. (2014). Cross-Resistance between Clofazimine and
Bedaquiline through Upregulation of MmpL5 in Mycobacterium tuberculosis. Antimicrob Agents
Chemother *58*, 2979–2981. https://doi.org/10.1128/AAC.00037-14.

38.   Alexander, D.C., Vasireddy, R., Vasireddy, S., Philley, J.V., Brown-Elliott, B.A., Perry, B.J.,
Griffith, D.E., Benwill, J.L., Cameron, A.D.S., and Wallace, R.J. (2017). Emergence of mmpT5 Variants
during Bedaquiline Treatment of Mycobacterium intracellulare Lung Disease. Journal of Clinical
Microbiology *55*, 574–584. https://doi.org/10.1128/jcm.02087-16.

39.   Halloum, I., Viljoen, A., Khanna, V., Craig, D., Bouchier, C., Brosch, R., Coxon, G., and Kremer,
L. (2017). Resistance to Thiacetazone Derivatives Active against Mycobacterium abscessus Involves
Mutations in the MmpL5 Transcriptional Repressor MAB_4384. Antimicrob Agents Chemother *61*,
e02509-16. https://doi.org/10.1128/AAC.02509-16.

40.   De Coster, W., D'Hert, S., Schultz, D.T., Cruts, M., and Van Broeckhoven, C. (2018). NanoPack:
visualizing and processing long-read sequencing data. Bioinformatics *34*, 2666–2669.
https://doi.org/10.1093/bioinformatics/bty149.

41.   Wick, R.R., Judd, L.M., Gorrie, C.L., and Holt, K.E. (2017). Completing bacterial genome
assemblies with multiplex MinION sequencing. Microb Genom *3*, e000132.
https://doi.org/10.1099/mgen.0.000132.

42.   Wang, J.R., Holt, J., McMillan, L., and Jones, C.D. (2018). FMLRC: Hybrid long read error
correction using an FM-index. BMC Bioinformatics *19*, 50. https://doi.org/10.1186/s12859-018-2051-3.

43.   Kolmogorov, M., Yuan, J., Lin, Y., and Pevzner, P.A. (2019). Assembly of long, error-prone reads
using repeat graphs. Nat Biotechnol *37*, 540–546. https://doi.org/10.1038/s41587-019-0072-8.

44.   Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A., Zeng,
Q., Wortman, J., Young, S.K., et al. (2014). Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. PLOS ONE *9*, e112963.
https://doi.org/10.1371/journal.pone.0112963.

45.   Gurevich, A., Saveliev, V., Vyahhi, N., and Tesler, G. (2013). QUAST: quality assessment tool for
genome assemblies. Bioinformatics *29*, 1072–1075. https://doi.org/10.1093/bioinformatics/btt086.

46.   Wick, R.R., Schultz, M.B., Zobel, J., and Holt, K.E. (2015). Bandage: interactive visualization of
de novo genome assemblies. Bioinformatics *31*, 3350–3352.
https://doi.org/10.1093/bioinformatics/btv383.

47.   Tanizawa, Y., Fujisawa, T., and Nakamura, Y. (2018). DFAST: a flexible prokaryotic genome
annotation pipeline for faster genome publication. Bioinformatics *34*, 1037–1039.
https://doi.org/10.1093/bioinformatics/btx713.

48.   Tanizawa, Y., Fujisawa, T., Kaminuma, E., Nakamura, Y., and Arita, M. (2016). DFAST and
DAGA: web-based integrated genome annotation tools and resources. Bioscience of Microbiota, Food
and Health *35*, 173–184. https://doi.org/10.12938/bmfh.16-003.

49.   Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S. (2018). High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun *9*,
5114. https://doi.org/10.1038/s41467-018-07641-9.

50.   Darling, A.C.E., Mau, B., Blattner, F.R., and Perna, N.T. (2004). Mauve: Multiple Alignment of
Conserved Genomic Sequence With Rearrangements. Genome Res *14*, 1394–1403.
https://doi.org/10.1101/gr.2289704.

51.   Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
Preprint at arXiv, https://doi.org/10.48550/arXiv.1303.3997 https://doi.org/10.48550/arXiv.1303.3997.

52.   Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and BCFtools.
Gigascience *10*, giab008. https://doi.org/10.1093/gigascience/giab008.

53.   Garrison, E., and Marth, G. (2012). Haplotype-based variant detection from short-read
sequencing. Preprint at arXiv, https://doi.org/10.48550/arXiv.1207.3907.

54.   Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., and
Ruden, D.M. (2012). A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w [1118]; iso-2; iso-3. Fly *6*,
80–92. https://doi.org/10.4161/fly.19695.

55.   Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24–26.
https://doi.org/10.1038/nbt.1754.

56.   Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*, 3094–
3100. https://doi.org/10.1093/bioinformatics/bty191.

57.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

58.   Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq—a Python framework to work with high-
throughput sequencing data. Bioinformatics *31*, 166–169. https://doi.org/10.1093/bioinformatics/btu638.

59.   Sun, J., Nishiyama, T., Shimizu, K., and Kadota, K. (2013). TCC: an R package for comparing tag
count data with robust normalization strategies. BMC Bioinformatics *14*, 219.
https://doi.org/10.1186/1471-2105-14-219.

60.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,
and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Molecular Biology and Evolution *37*, 1530–1534. https://doi.org/10.1093/molbev/msaa015.

61.   Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic Acids Research *49*, W293–W296.
https://doi.org/10.1093/nar/gkab301.

62.   Chen, S. (2023). Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp. iMeta *2*, e107. https://doi.org/10.1002/imt2.107.

63.   Shen, W., Le, S., Li, Y., and Hu, F. (2016). SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. PLoS ONE *11*, e0163962. https://doi.org/10.1371/journal.pone.0163962.

64.   Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics *26*, 841–842. https://doi.org/10.1093/bioinformatics/btq033.

65.   Clinical & Laboratory Standards Institute (2018). Susceptibility Testing of Mycobacteria, Nocardia
spp., and Other Aerobic Actinomycetes, M24Ed3E, (CLSI Publication).

66.   Bouso, J.M., and Planet, P.J. (2019). Complete nontuberculous mycobacteria whole genomes
using an optimized DNA extraction protocol for long-read sequencing. BMC Genomics *20*, 793.
https://doi.org/10.1186/s12864-019-6134-y.

67.   Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden,
T.L. (2009). BLAST+: architecture and applications. BMC Bioinformatics *10*, 421.
https://doi.org/10.1186/1471-2105-10-421.

68.   Feldgarden, M., Brover, V., Haft, D.H., Prasad, A.B., Slotta, D.J., Tolstoy, I., Tyson, G.H., Zhao,
S., Hsu, C.-H., McDermott, P.F., et al. (2019). Validating the AMRFinder Tool and Resistance Gene
Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates.
Antimicrob Agents Chemother *63*, e00483-19. https://doi.org/10.1128/AAC.00483-19.

69.   Jia, B., Raphenya, A.R., Alcock, B., Waglechner, N., Guo, P., Tsang, K.K., Lago, B.A., Dave,
B.M., Pereira, S., Sharma, A.N., et al. (2017). CARD 2017: expansion and model-centric curation of the

comprehensive antibiotic resistance database. Nucleic Acids Research *45*, D566–D573.
https://doi.org/10.1093/nar/gkw1004.

70.   Zankari, E., Hasman, H., Cosentino, S., Vestergaard, M., Rasmussen, S., Lund, O., Aarestrup,
F.M., and Larsen, M.V. (2012). Identification of acquired antimicrobial resistance genes. Journal of
Antimicrobial Chemotherapy *67*, 2640–2644. https://doi.org/10.1093/jac/dks261.

71.   Gupta, S.K., Padmanabhan, B.R., Diene, S.M., Lopez-Rojas, R., Kempf, M., Landraud, L., and
Rolain, J.-M. (2014). ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic Resistance Genes in
Bacterial Genomes. Antimicrobial Agents and Chemotherapy *58*, 212–220.
https://doi.org/10.1128/AAC.01310-13.

72.   Chen, L., Zheng, D., Liu, B., Yang, J., and Jin, Q. (2016). VFDB 2016: hierarchical and refined
dataset for big data analysis—10 years on. Nucleic Acids Research *44*, D694–D697.
https://doi.org/10.1093/nar/gkv1239.

73.   Carattoli, A., Zankari, E., García-Fernández, A., Voldby Larsen, M., Lund, O., Villa, L., Møller
Aarestrup, F., and Hasman, H. (2014). In Silico Detection and Typing of Plasmids using PlasmidFinder
and Plasmid Multilocus Sequence Typing. Antimicrobial Agents and Chemotherapy *58*, 3895–3903.
https://doi.org/10.1128/AAC.02412-14.

74.   Ingle, D.J., Valcanis, M., Kuzevski, A., Tauschek, M., Inouye, M., Stinear, T., Levine, M.M.,
Robins-Browne, R.M., and Holt, K.E. (2016). In silico serotyping of E. coli from short read data identifies
limited novel O-loci but extensive diversity of O:H serotype combinations within and between pathogenic
lineages. Microb Genom *2*, e000064. https://doi.org/10.1099/mgen.0.000064.

75.   Doster, E., Lakin, S.M., Dean, C.J., Wolfe, C., Young, J.G., Boucher, C., Belk, K.E., Noyes, N.R.,
and Morley, P.S. (2020). MEGARes 2.0: a database for classification of antimicrobial drug, biocide and
metal resistance determinants in metagenomic sequence data. Nucleic Acids Research *48*, D561–D569.
https://doi.org/10.1093/nar/gkz1010.

76.   Sun, J., Nishiyama, T., Shimizu, K., and Kadota, K. (2013). TCC: an R package for comparing tag
count data with robust normalization strategies. BMC Bioinformatics *14*, 219.
https://doi.org/10.1186/1471-2105-14-219.

77.   García-Alcalde, F., Okonechnikov, K., Carbonell, J., Cruz, L.M., Götz, S., Tarazona, S., Dopazo,
J., Meyer, T.F., and Conesa, A. (2012). Qualimap: evaluating next-generation sequencing alignment data.
Bioinformatics *28*, 2678–2679. https://doi.org/10.1093/bioinformatics/bts503.

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
M. abscessus subsp. *abscessus* type strain, ATCC 19977	ATCC	19977
M. abscessus subsp. *massiliense* type strain, JCM 15300	Japan Collection of Microorganisms (JCM), RIKEN BRC	15300
M. abscessus subsp. *bolletii* type strain, JCM 15297	JCM, RIKEN BRC	15297
Isolate57625, Isolate57629, Isolate57630, Isolate57596, and Isolate57626	This study	N/A
Chemicals, peptides, and recombinant proteins		
Trypticase soy agar with 5% sheep blood	Beckton Dickinson	Cat.# 251239
BBL™ Mueller Hinton II Broth (Cation-Adjusted)	Beckton Dickinson	Cat.# 212322
Middlebrook 7H9 Medium	Sigma-Adlrich	Cat.# M0178
2% OGAWA medium	Kyokuto Pharmaceutical Industrial	Cat.# 551-08026
Lysozyme	Sigma-Aldrich	Cat.# L4919
Critical commercial assays		
BrothMIC RGM	Kyokuto Pharmaceutical Industrial	Cat.# 551-10087
EZ-Beads	Promega	Cat.# AMR76813M
Wizard^R HMW DNA Extraction Kit	Promega	Cat.# A2920
Qubit™ BR DNA Quantitation Kit	Thermo Fisher Scientific	Cat.# Q32850
Ion Xpress™ Plus Fragment Library Kit	Thermo Fisher Scientific	Cat.# 4471252
Ion Xpress™ Barcode Adapters 1-16 kit	Thermo Fisher Scientific	Cat.# 4471250
TaqMan™ Quantitation Kit	Thermo Fisher Scientific	Cat.# 4468802
Ion PI™ Chip kit v3	Thermo Fisher Scientific	Cat.# A26711
UltraPure™ DNase/RNase-free Distilled Water	Thermo Fisher Scientific	Cat.# 10977015
SUPERase-In™ RNase Inhibitor	Thermo Fisher Scientific	Cat.# AM2696
mirVana™ miRNA Isolation Kit	Thermo Fisher Scientific	Cat.# AM1561
IonXpress™ RNA-Seq Barcode 1-16 Kit	Thermo Fisher Scientific	Cat.# 4475485
Ion PI™ Hi-Q™ OT2 200 Kit v2	Thermo Fisher Scientific	Cat.# A26434
Ion Total RNA-Seq Kit v2	Thermo Fisher Scientific	Cat.# 4479789
PowerUp™ SYBR™ Green Master Mix	Thermo Fisher Scientific	Cat.# A25741
KOD One^R PCR Master Mix	TOYOBO	Cat.#LMM-101
Short Read Eliminator Kit	PacBio	Cat.# 102-208-300
Ligation Sequencing Kit	Oxford Nanopore Technologies	Cat.# SQK-LSK109
Native Barcoding Expansion 1-12	Oxford Nanopore Technologies	Cat.# EXP-NBD104
Flow cell (R9.4.1)	Oxford Nanopore Technologies	Cat.# FLO-MIN106

NEBNext[R] rRNA Depletion Kit (Bacteria)	NEB	Cat.# E7850
Deposited data		
All raw sequence data generated in this study	DDBJ	PRJDB16220
Complete genome sequences (Chromosome, pMAB625-1, pMAB625-2 and pMAB625-3) of the Isolate57625	DDBJ	AP028613 AP028614 AP028615 AP028616
The analyzed RNA-seq data	Genomic Expression Archive (GEA)	E-GEAD-827
Oligonucleotides		
Forward primer for validation of large deletion in Figure S1 for Deletion_1, CAAGAACTGATGGCTGAAATCCTAATG	This study	N/A
Reverse primer for validation of large deletion in Figure S1 for Deletion_1, AAAGTGATGCTTGCTCTGCCAGTC	This study	N/A
Forward primer for validation of large deletion in Figure S1 for Insertion_1, GTTAGATATGTGCAGGAGTTCCGAGTG	This study	N/A
Reverse primer for validation of large deletion in Figure S1 for Insertion_1, ACCAATACTGCCGAATTCTCTAAAAGC	This study	N/A
Forward primer for validation of large deletion in Figure S1 for Insertion_2, GTGACATGATCTTGCTGCTCAAAAG	This study	N/A
Reverse primer for validation of large deletion in Figure S1 for Insertion_2, GCTATCTGTCGCAGTACTTCGTGACC	This study	N/A
Forward Primers for plasmid copy number analysis in Figure S5 for *HrpA* (MAB), GGATCTGGTCCACAAGACCTACAG	This study	N/A
Reverse Primers for plasmid copy number analysis in Figure S5 for *HrpA* (MAB), AAAGTGATGCTTGCTCTGCCAGTC	This study	N/A
Forward Primers for plasmid copy number analysis in Figure S5 for *HrpA* (no MAB), ACCCTGGTGAAGTTGATGACCGAC	This study	N/A
Reverse Primers for plasmid copy number analysis in Figure S5 for *HrpA* (no MAB), AGGAGGAAGTCGATGTTGAGGCTG	This study	N/A
Forward Primers for plasmid copy number analysis in Figure S5 for pMAB625-1(LOCUS_52450), TCTACTACGCAGACCCTGCACATATC	This study	N/A
Reverse Primers for plasmid copy number analysis in Figure S5 for pMAB625-1(LOCUS_52450), AACTGCTGCCACATCGCTTTCTG	This study	N/A
Forward Primers for plasmid copy number analysis in Figure S5 for pMAB625-2(LOCUS_53740), GACCGATCACGTCATCATTCCTAATC	This study	N/A
Reverse Primers for plasmid copy number analysis in Figure S5 for pMAB625-2(LOCUS_53740), GAACCTCAGCCTCAATATGCTCCTC	This study	N/A
Forward Primers for plasmid copy number analysis in Figure S5 for pMAB625-3(LOCUS_54770), CGACCAGGTACACATCCTGACCAC	This study	N/A
Reverse Primers for plasmid copy number analysis in Figure S5 for pMAB625-3(LOCUS_54770), GAGTTGGCTGATTCCGTCTTGGAG	This study	N/A
Software and algorithms		
R (ver. 4.3.1)	R Core Team (2024)	https://www.R-project.org/
MinKNOW (ver.21.06.0)	Oxford Nanopore Technologies	https://nanoporetech.com/
Guppy (ver. 5.0.7)	Oxford Nanopore Technologies	https://nanoporetech.com/
NanoPlot (ver. 1.38.0)	De Coster et al.[40]	https://github.com/wdecoster/NanoPlot

Porechop (ver. 0.2.4)	Wick et al[41]	https://github.com/rrwick/Porechop
Filtlong (ver. 0.2.1)	Wick	https://github.com/rrwick/Filtlong
Fmlrc (ver. 1.0.0)	Wang et al.[42]	https://github.com/holtjma/fmlrc
Flye (ver. 2.9-b1768)	Kolmogorov et al.[43]	https://github.com/mikolmogorov/Flye
Pilon (ver. 1.24.0)	Walker et al.[44]	https://github.com/broadinstitute/pilon
QUAST (ver. 5.0.2)	Gurevich et al.[45]	https://quast.sourceforge.net/
Bandage (ver. 0.8.1)	Wick et al.[46]	https://rrwick.github.io/Bandage/
DFAST (ver.1.5.0)	Tanizawa et al.[47,48]	https://dfast.ddbj.nig.ac.jp/
FastANI (ver.1.34)	Jain et al.[49]	https://github.com/ParBLiSS/FastANI
Mauve (ver. 2.4.0)	Darling et al.[50]	https://darlinglab.org/mauve/mauve.html
Abricate (ver.1.0.1)	Seeman. T	https://github.com/tseemann/abricate
BWA-MEM (ver. 0.7.17-r1188)	Li H.[51]	https://github.com/lh3/bwa
Samtools (ver.1.12)	Danecek et al.[52]	https://github.com/samtools/samtools
Freebayes (ver. 1.3.5)	Garrison et al.[53]	https://github.com/freebayes/freebayes
SnpEff (ver. 5.2)	Cingolani et al.[54]	https://pcingola.github.io/SnpEff/
IGV (ver.2.17.4)	Robinson et al.[55]	https://igv.org/
Minimap2 (ver. 2.21-r1071)	Li H.[56]	https://github.com/lh3/minimap2
Bowtie2 (ver. 2.5.4)	Langmead and Salzberg[57]	https://github.com/BenLangmead/bowtie2
HTseq (ver. 1.99.2)	Anders et al.[58]	https://github.com/htseq/htseq
R package, TCC (ver. 1.26.0)	Sun et al.[59]	https://bioconductor.org/packages/release/bioc/html/TCC.html
Snippy (ver. 4.6.0)	Seeman. T	https://github.com/tseemann/snippy
IQ-TREE (ver.2)	Minh et al.[60]	https://www.iqtree.org
iTOL ver.6.7	Letunic and Bork[61]	https://itol.embl.de
NGphylogeny.fr	Lemoine et al.[24]	https://ngphylogeny.fr
Fastp (ver. 0.23.4)	Chen[62]	https://github.com/OpenGene/fastp
Seqkit ver.2.4.0	Shen et al.[63]	https://github.com/shenwei356/seqkit
Bedtools ver. 2.30.0	Quinlan and Hall[64]	https://github.com/arq5x/bedtools2
Gubbins ver.3.3.1	Croucher[27]	https://github.com/nickjcroucher/gubbins

| R package, ggtree (ver 3.10.1) | Yu et al.[28] | https://bioconductor.org/packages/release/bioc/html/ggtree.html |
| R package, RVAideMemoire (ver. 0.9-83-7) | Herve M. | https://github.com/cran/RVAideMemoire |

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Clinical isolates and type strain

834
Ethical approval for this study was obtained from the Ethics Committee of Dokkyo Medical University
(approval number 28045). Regarding Isolate57625, Isolate57629, Isolate57630, Isolate57596, and
Isolate57626, specimens were obtained from a patient with osteomyelitis at the five time points in six
months indicated in Table 1. Each strain was isolated from the specimen at Shizuoka Children's Hospital
for diagnosis. The other clinical isolates used for PCR or WGS were stored at Tokyo Metropolitan Institute
for Geriatrics and Gerontology or Tokyo Women's Medical University or Dokkyo Medical University[25]. The
type strain of ABS, ATCC 19977, was purchased from ATCC (Manassas, VA, US). The type strain of MAS,
JCM 15300, and the type strain of BOL, JCM 15297, were provided by Japan Collection of Microorganisms,
RIKEN BRC which is participating in the National BioResource Project of the MEXT, Japan.

METHOD DETAILS

Growth curve

846
The type strain and the five clinical isolates were inoculated onto trypticase soy agar with 5% sheep blood
(Beckton Dickinson, Franklin Lakes, NJ, US) and incubated at 30 °C for five days. Next, some colonies
were inoculated into BBL^{TM} Mueller Hinton II Broth (Cation-Adjusted) (Beckton Dickinson) and incubated
overnight at 30°C. These culture media were measured for absorbance (OD_{600}), from which they were
prepared to 0.1 by medium. Seventy microliter of the adjusted culture solution was inoculated into 7 mL of
BBL^{TM} Mueller Hinton II Broth (Cation-Adjusted) and incubated at 30 °C for eight days with agitation (200
rpm). OD_{600} was measured on days one to eight after inoculation. Statistical analysis was performed using
R ver. 4.3.1.

Antimicrobial susceptibility testing

856
The type strain and the clinical isolates were subcultured on 2% OGAWA medium (Kyokuto Pharmaceutical
Industrial, Tokyo, Japan) at 30 °C for five days. Antimicrobial susceptibility testing was performed using
BrothMIC RGM (Kyokuto Pharmaceutical Industrial) following CLSI M24 3rd ed[65]. MICs were determined
as the minimum concentration at which bacteria had not grown on day four (all antibiotics) and day 14
(clarithromycin). The test was repeated three times and all MICs were confirmed to be the same between
the experiments, except for imipenem (IPM) for the isolates. For IPM, the MICs of the type strain were
confirmed to be the same between all three experiments, while for the MICs of Isolate57625 and
Isolate57626, they were confirmed to be the same between two out of three experiments, as indicated in
Table 1.

Mycobacterial DNA isolation and whole-genome sequencing

The type strain and the clinical isolates of *M. abscessus* were stored in frozen stock containing skim milk
and 5% (v/v) glycerol. Each strain was inoculated from frozen stocks into Middlebrook 7H9 Medium (Sigma-
Aldrich, St. Louis, MO, US) containing 10% (v/v) oleic acid-albumin-dextrose-catalase and 0.2% (v/v)
glycerol and cultured at 30 °C for six days with agitation (200 rpm). Bacterial genomic DNA for short-read
sequencing was extracted as previously reported[66]. The quality of the extracted DNA was confirmed by
agarose gel electrophoresis, and the concentration was measured using Qubit^{TM} BR DNA Quantitation Kit
(Thermo Fisher Scientific, Waltham, MA, US). Genomic DNA was fragmented, and the sequencing library
was prepared using the Ion Xpress^{TM} Plus Fragment Library Kit and the Ion Xpress^{TM} Barcode Adapters 1-
16 kit (Thermo Fisher Scientific) according to the manufacturer's instructions. The quantity of sequencing
library was assessed using the Ion Library TaqMan^{TM} Quantitation Kit (Thermo Fisher Scientific). Emulsion
PCR for sequence template synthesis was performed with the Ion PI^{TM} Hi-Q^{TM} OT2 200 Kit v2 using the

Ion One Touch™ II System and the Ion One Touch™ II ES (Thermo Fisher Scientific). Sequencing was
performed using the Ion PI™ Chip kit v3 and the Ion Proton™ System (Thermo Fisher Scientific). Data were
collected using the Torrent Suite v5.0.5 software (Thermo Fisher Scientific).

High-molecular-weight genomic DNA for long-read sequencing was extracted as follows. The bacterial
pellet was collected by centrifugation and homogenized using EZ-beads (Promega, Madison, WI, US) and
Shakeman 6 (Bio Medical Science, Tokyo, Japan). The homogenized pellet was lysed with Lysozyme
(Sigma-Aldrich) at 37 °C for one hour. Next, DNA was extracted using Wizard® HMW DNA Extraction Kit
(Promega) following the manufacturer's instructions. Fragmented DNA in the extracted DNA was removed
using the Short Read Eliminator Kit (PacBio, Menlo Park, CA, US). The sequencing library for long-read
sequencing was prepared using the Ligation Sequencing Kit (SQK-LSK109, Oxford Nanopore
Technologies, Oxford, UK) and the Native Barcoding Expansion 1-12 (EXP-NBD104, Oxford Nanopore
Technologies). In brief, the barcode sequences for multiplex sequencing were attached to the ends of the
HMW genomic DNA. In addition, the sequencing adapters were ligated to the end of HMW genomic DNA.
The prepared DNA was loaded into a flow cell (FLO-MIN106, Oxford Nanopore Technologies) attached to
MinION Mk1B (Oxford Nanopore Technologies). Sequencing data were acquired using MinKNOW
ver.21.06.0 (Oxford Nanopore Technologies), and base calling was performed using Guppy ver. 5.0.7
(Oxford Nanopore Technologies) with high accuracy mode using the National Institute of Genomics
supercomputer system (NIG, Shizuoka, Japan). Data quality was confirmed using NanoPlot ver. 1.38.0[40]
and adapter sequences were trimmed using Porechop ver. 0.2.4[41]. Raw reads shorter than 1000 bp
(Isolate57625, Isolate57629, Isolate57630, Isolate57596) or 5000 bp (Isolate57626) were removed using
Filtlong ver. 0.2.1 (https://github.com/rrwick/Filtlong). Filtered reads were corrected using Fmlrc ver. 1.0.0[42]
with the data of short-read sequencing acquired by the Ion Proton™ Sequencer. *De novo* assembly was
performed using Flye ver. 2.9-b1768[43] with corrected long-reads. Assembled genome sequences were
polished using Pilon ver. 1.24.0[44], evaluated using QUAST ver. 5.0.2[45], and visualized using Bandage ver.
0.8.1[46]. Genome annotation was performed using DFAST ver.1.5.0[47,48].

Plasmid sequence analysis

Known plasmid sequences were downloaded from PLSDB[14] by the taxonomic search for "*abscessus*".
Average nucleotide identity (ANI) was calculated using FastANI ver.1.34[49] and structural variants (SVs)
were analyzed using Mauve ver. 2.4.0[50].

Homology search was performed by blast+[67] using the amino acid sequence as the query or by Abricate
ver.1.0.1 (http://github.com/tseemann/abricate) using the nucleotide sequences as the query. NCBI
AMRFinderPlus[68], CARD[69], Resfinder[70], ARG-ANNOT[71], VFDB[72], PlasmidFinder[73], EcOH[74], and
MEGARes 2.00[75] were used as the database for the search of AMR genes. Mycobrowser[17] was used as
the database for the homology search of the mycobacterium genes.

SNVs and SVs analysis
Short-read data acquired by the Ion Proton™ Sequencer were mapped using BWA-MEM ver. 0.7.17-
r1188[51] against the reference sequence (CU458896.1, CU458745.1). Output BAM files were sorted using
samtools ver.1.12[52]. Valiant calling was performed using Freebayes ver. 1.3.5[53], and variants were
annotated using SnpEff ver. 5.2[54] with annotated genome sequence. The positions of single nucleotide
variants (SNVs) were visualized using IGV[55] (Broad Institute, Cambridge, MA, US). Assembled genome
sequences were aligned using Mauve ver. 2.4.0[50] for SVs analysis. Mapping the short- and long-read to

the assembled genome sequence using minimap2 ver. 2.21-r1071[56] was performed for the validation of
assembly.

PCR for the validation of SVs was performed using KOD One® PCR Master Mix (TOYOBO, Osaka, Japan)
and the primers indicated in Table S1. Amplified DNA was confirmed using agarose gel electrophoresis
following staining with SYBR™ Safe Gel Stain (Thermo Fisher Scientific).

RNA-seq analysis

Total RNA was extracted as follows: The bacterial pellet was collected by centrifugation and suspended in
UltraPure™ DNase/RNase-free Distilled Water (Thermo Fisher Scientific) and homogenized using EZ-
beads (Promega) and Shakeman 6 (Bio Medical Science). SUPERase-In™ RNase Inhibitor (Thermo Fisher
Scientific) was added to the homogenized pellet. Next, the homogenized pellet was lysed by Lysozyme
(Sigma-Aldrich) at 37 °C for an hour. Total RNA was purified by mirVana™ miRNA Isolation Kit (Thermo
Fisher Scientific). The size and quality of total RNA were assessed using the Agilent 2100 Bioanalyzer
(Agilent Technologies, Santa Clara, CA, US) and the RNA 6000 Nano Kit (Agilent Technologies). rRNA
was depleted using the NEBNext® rRNA Depletion Kit (Bacteria) (NEB, Ipswich, MA, US) from 500 ng of
total RNA. The whole transcriptome library was prepared using the Ion Total RNA-Seq Kit v2 (Thermo
Fisher Scientific) as follows: First, the rRNA-depleted RNA was fragmented by RNase III at 37 °C for 10
minutes and purified using the Magnetic Beads Cleanup Module. Next, adapter sequences were ligated
and reverse transcription was performed. The whole transcriptome library was amplified and barcoded
using the IonXpress™ RNA-Seq Barcode 1-16 Kit (Thermo Fisher Scientific). The quantity of the amplified
whole transcriptome library was assessed using the Ion Library TaqMan™ Quantitation Kit (Thermo Fisher
Scientific). Emulsion PCR for sequence template synthesis was performed using the Ion PI™ Hi-Q™ OT2
200 Kit v2 with the Ion One Touch™ II System and the Ion One Touch™ II ES (Thermo Fisher Scientific).
Sequencing was performed using the Ion Proton™ System with Ion PI™ Chip kit v3 (Thermo Fisher
Scientific). Data was collected using the Torrent Suite ver. 5.0.5 software (Thermo Fisher Scientific).
Sequence reads were mapped using Bowtie2 ver. 2.5.4[57] against the assembled genome sequence of
Isolate57625. Mapped reads were counted using HTseq ver. 1.99.2[58]. Merged transcript count data were
normalized using TCC ver. 1.26.0[76] and differentially expressed genes (DEG) were detected under false
discovery rate (FDR) < 0.01.

Phylogenetic tree analysis
The genome sequences of the various clinical isolates shown in Figure S2C for phylogenetic tree analysis
were obtained from the NCBI database and are listed in Table S2. Core-genome SNVs were detected using
Snippy ver. 4.6.0 (https://github.com/tseemann/snippy) with each assembled genome sequence and the
type strain (ATCC 19977, CU458896.1) as the reference. The concatenated core-genome SNVs were also
aligned using the snippy-core script. The phylogenetic tree was constructed using IQ-TREE ver.2[60] using
the maximum likelihood criterion and rendered using iTOL ver.6.7[61]. Phylogenetic tree analysis of pMAB625
sequences was performed by the same method. For the phylogenetic tree analysis regarding the
components of the ESX secretion system, the amino acid sequences of ESX loci (*eccB*, *eccC*, *eccD*, *mycP*,
and *eccE*) were concatenated and used as input for NGphylogeny.fr[24] using the PhyML+SMS inference
method.
966
PCR for plasmid detection and copy number analysis
PCR was performed using KOD One® PCR Master Mix (TOYOBO) and the primers indicated in Table S1.
Amplified DNA was confirmed using agarose gel electrophoresis. The plasmid copy number was analyzed
by qPCR using PowerUp™ SYBR™ Green Master Mix (Thermo Fisher Scientific). qPCR was performed
using the QuantStudio™ 5 Real-Time PCR System (Thermo Fisher Scientific). Five nanogram of genomic

DNA was used for a 20 µL reaction volume of qPCR. The plasmid copy number is shown as a relative value
normalized using the copy number of *HrpA*, which is present in one copy on the chromosome.
974
Mapping the read data of various clinical isolates to the genome sequence of Isolate57625

Raw read data of *M. abscessus* clinical isolate in Japan, Taiwan[7], China[26], the US, the United Kingdom
(UK) and Spain were obtained from the NCBI sequence reads archive (SRA). Raw read data were trimmed
using fastp ver. 0.23.4[62] and reduced using seqkit ver.2.4.0[63] to 200 Mb for the Ion Proton™ single-end
data and 100 Mb for the Illumina pair-end data to equalize the depth of coverage among isolates. Reads
were mapped using BWA-MEM ver. 0.7.17-r1188 against the chromosome and plasmid sequences of
Isolate57625 and other plasmids which were hit using PLSDB by the taxonomic searching of "*abscessus*"
with removing the redundant plasmid sequences (ANI (%)>97, without large SV (>8 kbp)). The depth of
coverage was calculated using Qualimap ver. 2.2.2-dev[77]. The covered genome region was calculated
using bedtools ver. 2.30.0[64]. We considered the isolate whose depth of coverage was higher than 10x and
the covered genome region was higher than 75% as the plasmid-harboring isolate. The 152 sequences for
the analysis of the distribution of plasmids identified in other NTM and previously reported[29] were listed in
Table S2. For the phylogenetic tree analysis, the aligned assumed chromosome sequences using Snippy-
core script were used as input for Gubbins ver.3.3.1[27]. The phylogenetic tree was rendered using ggtree
ver 3.10.1[28].

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis in this study were performed using R ver. 4.3.1. Statistical significance in Figure S1 were
calculated using one-way ANOVA followed by Dunnett's test and data were represents the mean of the
three independent experiments and the error bar represents the standard deviation. Statistical analysis of
the number of clinical isolates retaining plasmid in Table 4 was performed using Fisher's exact test with
Holm's method for correction of multiple comparisons using RVAideMemoire ver. 0.9-83-7
(https://github.com/cran/RVAideMemoire).