Dimeric assembly of F$_1$-like ATPase for the gliding motility of

Mycoplasma

Takuma Toyonaga[1,2], Takayuki Kato[3], Akihiro Kawamoto[3], Tomoko Miyata[4,5], Keisuke

Kawakami[6], Junso Fujita[4,5,7], Tasuku Hamaguchi[8], Keiichi Namba[4,5], Makoto Miyata[1,2]

[1]Graduate School of Science, Osaka Metropolitan University, 3-3-138 Sugimoto,

Sumiyoshi-ku, Osaka 558-8585, Japan

[2]The OMU Advanced Research Institute for Natural Science and Technology, Osaka

Metropolitan University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

[3]Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka

565-0871, Japan

[4]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita,

Osaka 565-0871, Japan

[5]JEOL YOKOGUSHI Research Alliance Laboratories, Osaka University, 1-3

Yamadaoka, Suita, Osaka 565-0871, Japan

[6]Biostructual Mechanism Laboratory, RIKEN, SPring-8 Center, 1-1-1, Kouto, Sayo,

Hyogo 679-5148, Japan

[7]Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita,

Osaka 565-0871, Japan

[8]Institute of Multidisciplinary Research for Advanced Materials, Tohoku University,

2-1-1 Katahira, Aoba-ku, Sendai 980-8577, Japan

Corresponding author: Makoto Miyata; Email: miyata@omu.ac.jp

Present address: Takuma Toyonaga: Institute of Multidisciplinary Research for

Advanced Materials, Tohoku University, 2-1-1 Katahira, Aoba-ku, Sendai 980-8577,

Japan

Abstract

Rotary ATPases, including F_1F_O-, V_1V_O- and A_1A_O-ATPases, are molecular motors that

exhibit rotational movements for energy conversion[1]. In the gliding bacterium,

Mycoplasma mobile, a dimeric F_1-like ATPase forms a chain structure within the cell[2],

which is proposed to drive the gliding motility[2–4]. However, the mechanisms of force

generation and transmission remain unclear. We determined the electron

cryomicroscopy (cryo-EM) structure of the dimeric F_1-like ATPase complex. The

structure revealed an assembly distinct from those of dimeric F_1F_O-ATPases[5]. The

F_1-like ATPase unit associated by two subunits, GliD and GliE was named G_1-ATPase

as a new R_1 domain of rotary ATPases. G_1-β subunit, a homologue of the F_1-ATPase

catalytic subunit[6], exhibited a specific N-terminal region that incorporates the glycolytic

enzyme, phosphoglycerate kinase into the complex. Structural features of the ATPase

displayed strong similarities to F_1-ATPase, suggesting a rotation based on the rotary

catalytic mechanism[1,7–10]. Overall, the cryo-EM structure provides insights into the

mechanism through which G_1-ATPase drives the *Mycoplasma* gliding motility.

Introduction

Rotary ATPases are molecular rotary motors that link ATP hydrolysis or synthesis with

ion transport via rotational motion in biological membranes[1]. These motors comprise

two common domains: soluble R_1, which is responsible for ATP hydrolysis and

synthesis, and membrane-embedded R_O, which is responsible for ion transport. Rotary

ATPase family defined by R_1R_O includes F_1F_O-, V_1V_O- and A_1A_O-ATPases[1]. Notably,

F_1F_O-ATPase, also known as F_1F_O ATP synthase is conserved across most eukaryotes,

archaea, and bacteria, and responsible for ATP synthesis and the maintenance of

membrane potential[5]. An F_1-like ATPase gene cluster, referred to as Type 2 ATPase, was

reported in four *Mycoplasma* species, *M. mobile*, *M. pulmonis*, *M. agassizii*, and *M.*

testudineum, in addition to the genuine F_1F_O-ATPase gene cluster, referred to as Type 1

ATPase[11,12] (Extended Data Fig. 1). The presence of F_1-ATPase α and β subunit

homologues (MMOBs 1660 and 1670 in *M. mobile*) in the Type 2 ATPase gene cluster

suggests a shared evolutionary origin in the F_1 domain. Type 2 ATPase is proposed to

drive the gliding motility observed in *M. mobile* and *M. pulmonis*[2–4,13,14]. *M. mobile*, a

fish pathogen, glides at speeds of up to 4 µm per second on a solid surface covered with

sialylated oligosaccharides found on host cell surfaces[15] (Fig. 1a, Extended Data Fig. 2).

In our proposed model for gliding, filamentous leg structures on the mycoplasma cell

surface, which are approximately 50 nm, propel the cell body by catching and pulling

sialylated oligosaccharides on the solid surface. Inside the cell, motors composed of

Type 2 ATPase and phosphoglycerate kinase (PGK), which converts ADP to ATP

through glycolysis, are linked near the cell membrane to form 'internal chains'[2,6]. The

motor is formed by dimerised F_1-like ATPases with unique bridge structures and is

referred to as a twin motor[2,16]. However, the mechanisms of force generation and

transmission remain unclear. This study aimed to elucidate the detailed structure of the

twin motor using cryo-EM single-particle analysis (SPA). Notably, the structural

features had a strong homology to those of F_1-ATPase, suggesting a rotary catalytic

mechanism analogous to that of F_1-ATPase[1,7–10] for the gliding motility of *M. mobile*.

Results

Protein identification using the cryo-EM structure of twin motor

Cryo-EM SPA was performed for twin motors isolated from *M. mobile* cells using

images collected on the epoxidised graphene grid (EG-grid)[17] (Extended Data Table 1).

The 3D structure of the twin motor was determined at a resolution of 3.2 Å (Fig. 1b,

Extended Data Figs. 3, 4). Local refinement improved the resolution of the regions,

each containing the F_1-like ATPase unit, to 3.1 Å (Extended Data Figs. 3b, 4). The

density maps enabled the construction of the atomic model for most regions of the twin

motor (Fig. 1c). The modelled sequence regions of each subunit are summarised in

Extended Data Table 2. The twin motor exhibited a rectangular structure with

dimensions of approximately $350 \times 250 \times 150$ Å and at least 24 polypeptide chains with

a total mass of approximately 1.5 MDa. Notably, the two F_1-like ATPases were inclined

to each other by approximately 5° and formed a dimer with pseudo-two-fold rotational

symmetry and several accessories (Fig. 1c, d). The F_1-like ATPase comprised a

hexameric ring of two alternating subunits and a central shaft subunit, identified as

MMOBs 1660, 1670, and 1630, respectively. The conserved ATPase structure of the R_1

domain in a unique motor indicates that the F_1-like ATPase clarified here is a new R_1

domain. In the present study, the ATPase unit was designated G_1-ATPase, with

assignment of MMOBs 1660, 1670, and 1630 to α, β, and γ, respectively, according to

the subunit names of F_1-ATPase[10]. In the hexameric ring, nucleotide-derived densities

were observed at five of the six subunit interfaces, despite the isolation and analysis of

samples under nucleotide free condition, indicating the presence of endogenous

nucleotides as reported in F_1-ATPase[18] (Fig. 1c, e). The subunit attached to α, forming a

'bridge' between two $\alpha_3\beta_3\gamma$ subcomplexes, was identified as MMOB1620 and named

GliD in the present study (Fig. 1c). The three densities protruding from the G_1-ATPase

dimer were identified as MMOB4530, which was previously annotated as PGK. Due to

the low resolution of the PGK1–3 regions at the edge of the map, atomic models were

constructed for only 204, 203 and 203 aa of the total length of 511 aa, including 60, 72

and 98 aa alanine truncations, respectively (Extended Data Table 2, Extended Data Fig.

5). The denoised 3D map revealed additional distinct densities at six locations,

including the PGK1–3 regions (Extended Data Fig. 6a). These six densities exhibited

correlation coefficients ranging from 0.37 to 0.48 with the crystal structure of PGK

from *Staphylococcus aureus* (PDB ID: 4DG5) (Extended Data Fig. 6b, c). Although the

values are low, these assignments are consistent with the stoichiometry previously

estimated via SDS-PAGE[2], suggesting these maps are PGK molecules. A density bound

only to one of the two $\alpha_3\beta_3\gamma$ subcomplexes was identified as MMOB3660, which was

newly detected via SDS-PAGE of the twin motor and named GliE as a G_1-ATPase

subunit in the present study (Fig. 1b, c, Extended Data Fig. 7). This small protein (112

aa), characterised by a β-sandwich, was nestled between the ATPase hexamer and other

subunits, breaking the rotational symmetry of the complex (Extended Data Fig. 8).

Although MMOB1640, encoded in the Type 2 ATPase gene cluster, was detected via

SDS-PAGE of the twin motor (Extended Data Fig. 7), the protein density was not

observed from the overall 3D map.

Structural features of G_1-ATPase αβγ

G_1-β and G_1-α exhibited high structural similarity to the β and α subunits of the

F_1-ATPase from the thermophilic bacterium, *Bacillus* PS3 (PDB ID: 8HH5), based on

the superimposition of their Cα atoms with the root mean squared deviation (RMSD)

values of 1.95 Å and 2.79 Å, respectively (Fig. 1g, i). Both G_1-β and G_1-α had three

domains: crown, nucleotide-binding domain (NBD), and C-terminal domain (CTD) (Fig.

1f, h). The NBDs of G_1-β and G_1-α comprise a phosphate-binding motif of P-loop[19],

consisting of the amino acid sequences GGAGVGKT and GDRGTGKT, respectively.

The G_1-β-CTD contained a DELSEED loop, which is important for torque transmission

in the F_1-ATPase β subunit[20], with an amino acid sequence of DALSEDD (Fig. 1f). A

remarkable difference between G_1-β and the F_1-ATPase β subunit was the presence of

an extended N-terminal region consisting of the N-terminal domain (NTD) and linker

(Fig. 1f, Extended Data Fig. 9a). This region comprised approximately 100 residues

located within the extra N-terminal 299 residues of G_1-β[6,12]. Notably, the linker hangs

outside the G_1-ATPase along the crown and NBD of G_1-β (Extended Data Fig. 9b).

G_1-β-NTDs, consisting of 4 to 6 helices, were oriented in various directions among G_1-β,

depending on their position in the complex (Extended Data Fig. 9c, d). Further,

G_1-β-NTD interacted with PGK (Fig. 2a, b). In addition, at the interface of the two

G_1-ATPase monomers, the N-terminal regions of G_1-β3 and G_1-β6 interacted with the

NBDs of G_1-α6 and G_1-α3 of another G_1-ATPase monomer, respectively (Fig. 2a, c).

These interactions suggest that the β-specific N-terminal region plays a role in the

incorporation of PGK into the twin motor and stabilisation of the ATPase dimer. A

difference between G_1-α and the F_1-ATPase α subunit is the absence of the H1 helix,

which interacts with the peripheral stalk[21] (Extended Data Fig. 10). This absence is

consistent with the lack of a gene homologous to the peripheral stalk in the Type 2

ATPase gene cluster[12]. G_1-γ, which is composed of a coiled coil and an α/β fold typical

of the F_1-ATPase γ subunit, was superimposed on the γ subunit of the *Bacillus* PS3

F_1-ATPase with an RMSD value of 5.79 Å for Cα atoms due to a lower sequence

conservation than that of the catalytic subunits[15] (Fig. 1j, k). The folded C-terminal

α-helix, which is 25 Å longer than the F_1-ATPase γ subunit, protrudes from the

N-terminal side of the hexameric ring and interacts with the crowns of G_1-$\alpha\beta$ (Fig. 1k,

Extended Data Fig. 11a, b). The coiled coil of G_1-γ hydrophobically interacted with the

CTDs of G_1-$\alpha\beta$ in the hexameric ring (Extended Data Fig. 11c), and in the F_1-ATPase,

this interaction is involved in axle hold and torque transmission[22].

Accessories of ATPase dimer

Generally, PGK is bifurcated into two domains interconnected by an α-helix: the NTD,

which binds to 3-phosphoglycerate (3-PG) or 1,3-bisphosphoglycerate (1,3-BPG), and

the CTD, which binds to Mg-ADP or Mg-ATP[23] (Extended Data Fig. 12a). The

twin-motor PGK structure, comprising the CTD and a segment of the α-helix, exhibited

a high structural similarity to the PGK from *S. aureus* (PDB ID: 4DG5), with an RMSD

value of 3.12 Å for Cα atoms (Extended Data Fig. 12b, c). The β loop of the PGK-CTD

invades the depression of the folded β-NTD and interacts with the H2 helix and the

H2–H3 loop of the β-NTD (Fig. 2d). The H4–β loop and H5 helix of the PGK-CTD also

interacted with the H5 and H4 helices of the β-NTD, respectively. Nucleotide-derived

densities were not found in the PGK-CTD of the twin motor, despite conservation of the

residues important for nucleotide binding in PGK[24] (Extended Data Figs. 12d–f, 13).

None of the twin-motor PGK-NTDs were modelled due to the poor densities of the

reconstruction. However, the binding and catalytic residues for 3-PG or 1,3-BPG in

PGK were conserved in terms of amino acid sequence, suggesting that the twin-motor

PGK possesses enzymatic activity (Extended Data Fig. 13). As the binding and catalytic

residues were not conserved in another PGK homologue, MMOB4490, encoded in the

M. mobile genome, the twin-motor PGK is expected to function as the original PGK in

glycolysis (Extended Data Fig. 13). The GliD (GliD1,3,4,6) bridge components formed

dimers at the interface of two G_1-ATPase monomers, whereas GliD2,5 at the twin-motor

end existed as monomers (Fig. 2a, e, f). GliD contained a globular domain and an

N-terminal long α-helix (H2), which was only visualised in the dimer. The globular

domain interacted with the H20–H25 region at the C-terminus of α (Fig. 2g, h). The two

H2 helices in the GliD dimer aligned parallel to each other across the H20–H25 region

of α, GliE, β-NTD, and β-CTD (Fig. 2f). These two α-helices are reminiscent of the

dynein buttress that connects the AAA+ ring and the tubulin-binding domain[25],

suggesting a regulatory role for structural changes induced by the ATPase activity of the

G_1-ATPases.

Structural information suggesting a catalytic mechanism analogous to that of

F_1-ATPase

The crown region of the hexameric ring exhibited approximately six-fold rotational

symmetry, whereas the CTD region was asymmetric (Extended Data Fig. 11a). This

observation suggests a structural change with a pivot point between the crown and CTD.

β1–3 adopted three distinct conformations, which appear to be the transition states of

CTD flexion motion towards γ, with a positional difference of 11 Å for the DELSEED

loop (Fig. 3a). These conformations were named $β_C$ (closed), $β_{HO}$ (half-open), and $β_O$

(open), according to the conformation names of F_1-ATPase[8]. The conformations were

arranged clockwise when viewed from the C-terminus of the hexameric ring (Fig. 3b).

These conformations resemble the states referred to as *ATP-waiting* or *step-waiting*,

which occurs during the catalytic reaction of *Bacillus* PS3 F_1-ATPase; the latter occurs

after ATP binding[8] (Extended Data Table 3). In α, the region corresponding to the

DELSEED loop exhibited only a 5 Å difference in position relative to the γ side,

whereas the H20–H25 region interacting with GliD exhibited a difference of 9 Å

(Extended Data Fig. 14). These conformations were named $α_C$ (close), $α_{HO}$ (half-open),

and $α_O$ (open) based on the H20–H25 positions viewed from the γ side. If this difference

in α represents a conformational change associated with ATP hydrolysis of the twin

motor, force transfer may occur from the $α_3β_3γ$ subcomplex of G_1-ATPase to GliD. The

catalytic sites formed at the α–β interface of $α_Cβ_C$ and $α_{HO}β_{HO}$ contained Mg-ATP and

Mg-ADP + P_i, respectively, whereas no nucleotide density was found at the catalytic

site of $α_Oβ_O$ (Fig. 3b–d, Extended Data Fig. 15). The nucleotide-binding states were also

shared between the two G_1-ATPase monomers according to the pseudo-two-fold

rotational symmetry. The residues of F_1-ATPase involved in ATP hydrolysis, including

those in the arginine finger, were conserved in G_1-αβ, suggesting a shared ATP

hydrolysis pathway between F_1-ATPase and G_1-ATPase (Fig. 3d). Mg-ATP was

observed at the remaining three interfaces, hereafter referred to as the β–α interfaces

(Fig. 3b, Extended Data Fig. 15). At the β–α interface, α-K179 of G_1-ATPase

corresponded to the position of conserved residue Q200 of the F_1-ATPase α subunit

(Extended Data Fig. 16a, b). Compared with the α–β interface, α-K179 is charge

inverted relative to the glutamate essential for the catalytic activity (E190 of the

F_1-ATPase β subunit), suggesting that the β–α interface, akin to F_1-ATPase, does not

exhibit ATP hydrolysis activity (Extended Data Fig. 16c). The conformational sequence

pattern of three β subunits with bound nucleotides was identical to that of the

ATP-waiting state of F_1-ATPase[8] (Fig. 3b). These data suggest that Mg-ATP binding to

the catalytic site of $\alpha_O\beta_O$ promotes the catalytic reaction of the G_1-ATPase based on the

catalytic mechanism analogous to that of F_1-ATPase, as discussed below.

Twin motors in the chain

The atomic model of the twin motor comprising G_1-ATPase dimer and PGK was fitted

into the previously reconstructed density of the internal chain, obtained using negative

staining electron microscopy (EM) SPA[2] (Fig. 4a). To circumvent misfitting of the

long-axis orientation of the twin motors due to the low-resolution density map (29.7 Å)

of the chain structure, regions with asymmetric structures, such as the NTDs of β3,6,

GliE, and PGK3, were excluded from the fitted model. At the interface between

twin-motor units, GliDs that project along the long axis of the twin motor faced each

other, suggesting their role as a linker for chain formation. However, the 3D map of the

internal chain revealed a remaining density at the interface that cannot be filled by

GliDs, which may correspond to the approximately 100 residues comprising an

N-terminal region and loops that have not been modelled.

Discussion

The major difference between the G_1-ATPase dimer and the dimeric form of eukaryotic

mitochondrial F_1F_O-ATPases[5] is the interaction between the R_1 domains through GliD

and the N-terminal region of β. Although the GliD dimer does not have a

transmembrane segment, the two long α-helices at the N-terminus are reminiscent of the

structure of a peripheral stalk. This observation is consistent with the hypothesis that the

MMOB1620 gene encoding GliD is derived through gene fusion from the b and δ

subunits, which comprise the peripheral stalk[15]. The GliD-mediated dimer and chain

formation may allow G_1-ATPases to work cooperatively. Consistently, changes in the

distance between twin motors have been associated with the ATP hydrolysis reaction,

suggesting force transfer between the twin motors through GliD[3,4]. The connections

between $\alpha_3\beta_3$ catalytic hexamers in the G_1-ATPase dimer may be related to the lower

ATPase activity than that of the F_1-ATPase, with a maximum turnover rate of $0.18\ \text{s}^{-1}$

per β molecule and a Km of $74\ \mu\text{M}^2$. The twin-motor PGK may regulate the assembly

and activity of ATPases, as observed for 6-phosphofructo-1-kinase and aldolase,

glycolytic enzymes that bind to V_1V_O-ATPase[26,27]. Further, PGK molecules protruding

from the transverse axis of the twin motor may be involved in the sheet formation of the

internal chain, according to negative staining EM[16]. The weak density of PGK in the

cryo-EM map can be explained by its flexibility, which was identified using high-speed

atomic force microscopy[2]. The flexibility of the PGK molecule may suggest that the

chain sheet serves as a cytoskeleton to withstand mechanical stress in the mycoplasma

cell, which does not have a cell wall. The GliE subunit, which is bound to only one side

of the dimeric G_1-ATPase, creates asymmetry in the twin motor structure and may

contribute to the unidirectional nature of the gliding motility. However, this protein may

also bind to another side of the dimeric G_1-ATPase in the cell but dislodge during

purification.

The three distinct forms of the catalytic subunit β and their corresponding nucleotide

binding suggest a rotary catalytic mechanism analogous to that of F_1-ATPase[8] (Fig. 4b).

In this rotary catalytic mechanism, $\alpha_O\beta_O$ transitions to $\alpha_C\beta_C$, and the DELSEED loop of

β in this transition approaches γ after ATP binding at the catalytic site. The other αβ

dimers in the hexameric ring also operate cooperatively, with $\alpha_C\beta_C$ transitioning to

$\alpha_{HO}\beta_{HO}$, associated with ATP hydrolysis and $\alpha_{HO}\beta_{HO}$ transitioning to $\alpha_O\beta_O$, associated

with the release of ADP and phosphate. This one-step conformational change of the

hexameric ring prompts a 120° rotation of γ. Each $\alpha\beta$ dimer sequentially hydrolyses

ATP, and through a three-step process, the hexameric ring structure and position of γ

revert to their initial states. The pseudo-two-fold rotational symmetry structure of the

twin motor suggests that the two G_1-ATPase monomers operate in a synchronised

manner, maintaining the structural symmetry. Throughout the catalytic reaction,

significant changes in the positioning of the two G_1-ATPase monomers within the twin

motor may be hindered due to the presence of the N-terminal regions of β bridging the

two monomers and a buttress-like structure composed of two long α-helices of the GliD

dimer (Fig. 2c, f).

For cell gliding, the rotary motions of the twin motors must be converted into a linear

motion. This conversion mechanism may occur due to the membrane protein Gli521,

which consists of a hook and rod, thereby functioning as a crank[28] (Fig. 4c). Upon the

conversion of the rotary motion to linear motion, Gli521 pulls the leg protein, Gli349,

which is bound to sialylated oligosaccharides on the host cell surface, thereby propelling

the mycoplasma cell forward[29]. The presence of a globular domain in G_1-γ, similar to

that in the F_1-ATPase γ subunit, suggests a potential interaction with a

membrane-embedded F_O-like domain. MMOB1610 and MMOB1650, encoded in the

Type 2 ATPase gene cluster, possess 12 and 2 transmembrane segments, respectively[15].

These two proteins are potential interaction partners for the globular domain of γ at the

cell membrane, transmitting the rotary motion to Gli521 on the cell surface.

The roles of rotary ATPases and their relatives, such as the flagellar type III protein

export apparatus, which exhibit a hexameric ring with a central shaft, involve protein

and ion transport[30,31]. We hypothesised that F_1F_O-ATPase has acquired the role and

mechanism of the twin motor through an incidental contact with an adhesin on the

membrane during the evolution of mycoplasma from a non-motile to a motile state[2,32].

This primitive gliding motility would have conferred a survival advantage. Another

F_1-like ATPase, referred to as Type 3 ATPase, is also found in mycoplasma, and this

F_1-like ATPase may drive the system that cleaves host antibodies[12,33]. Notably, Type 2

and 3 ATPases transition from F_1-ATPase to drive mycoplasma-specific systems.

Understanding these unique ATPases will contribute to investigations into the working

principles and evolution of rotary ATPases.

References

1. Stewart, A. G., Sobti, M., Harvey, R. P. & Stock, D. Rotary ATPases. *Bioarchitecture* **3**, 2–12 (2013).

2. Toyonaga, T. *et al.* Chained structure of dimeric F1-like ATPase in *Mycoplasma mobile* gliding machinery. *mBio* **12**, e0141421 (2021).

3. Nishikawa, M. S. *et al.* Refined mechanism of *Mycoplasma mobile* gliding based on structure, ATPase activity, and sialic acid binding of machinery. *mBio* **10**, e02846-19 (2019).

4. Kobayashi, K. *et al.* Movements of *Mycoplasma mobile* gliding machinery detected by high-speed atomic force microscopy. *mBio* **12**, e0004021 (2021).

5. Kühlbrandt, W. Structure and mechanisms of F-Type ATP synthases. *Annu Rev Biochem* **88**, 515–549 (2019).

6. Nakane, D. & Miyata, M. Cytoskeletal 'jellyfish' structure of *Mycoplasma mobile*. *Proc Natl Acad Sci U S A* **104**, 19518–19523 (2007).

7. Noji, H. & Ueno, H. How Does F1-ATPase Generate Torque?: Analysis From Cryo-Electron Microscopy and Rotational Catalysis of Thermophilic F1. *Front Microbiol* **13**, 904084 (2022).

8.   Nakano, A., Kishikawa, J. ichi, Mitsuoka, K. & Yokoyama, K. Mechanism of
ATP hydrolysis dependent rotation of bacterial ATP synthase. *Nat Commun* **14**,
4090 (2023).

9.   Noji, H., Yasuda, R., Yoshida, M. & Kinosita, K. Direct observation of the
rotation of F1-ATPase. *Nature* **386**, 299–302 (1997).

10. Abrahams, J. P., Leslie, A. G. W., Lutter, R. & Walker, J. E. Structure at 2.8 Â
resolution of F1-ATPase from bovine heart mitochondria. *Nature* **370**, 621–628
(1994).

11. Tulum, I., Kimura, K. & Miyata, M. Identification and sequence analyses of the
gliding machinery proteins from *Mycoplasma mobile*. *Scientific Reports 2020*
10:1 **10**, 1–11 (2020).

12. Béven, L. *et al.* Specific evolution of F1-like ATPases in mycoplasmas. *PLoS*
One **7**, e38793 (2012).

13. Bredt, W. & Radestock, U. Gliding motility of *Mycoplasma pulmonis*. *J Bacteriol*
130, 937 (1977).

14. Rosengarten, R. & Kirchhoff, H. Gliding motility of Mycoplasma sp. nov. strain
163K. *J Bacteriol* **169**, 1891–1898 (1987).

15. Miyata, M. & Hamaguchi, T. Prospects for the gliding mechanism of
Mycoplasma mobile. *Curr Opin Microbiol* **29**, 15–21 (2016).

16. Fukushima, M., Toyonaga, T., Tahara, Y. O., Nakane, D. & Miyata, M. Internal
structure of *Mycoplasma mobile* gliding machinery analyzed by negative staining
electron tomography. *Biophys Physicobiol* e210015 (2024).

17. Fujita, J. *et al.* Epoxidized graphene grid for highly efficient high-resolution
cryoEM structural analysis. *Sci Rep* **13**, 2279 (2023).

18. Leimgruber, R. M. & Senior, A. E. Removal of 'tightly bound' nucleotides from
phosphorylating submitochondrial particles. *Journal of Biological Chemistry* **251**,
7110–7113 (1976).

19. Menz, R. I., Walker, J. E. & Leslie, A. G. W. Structure of bovine mitochondrial
F(1)-ATPase with nucleotide bound to all three catalytic sites: implications for
the mechanism of rotary catalysis. *Cell* **106**, 331–341 (2001).

20. Tanigawara, M. *et al.* Role of the DELSEED Loop in Torque Transmission of
F1-ATPase. *Biophys J* **103**, 970–978 (2012).

21. Sobti, M. *et al.* Cryo-EM structures of the autoinhibited *E. coli* ATP synthase in
three rotational states. *Elife* **5**, e21598 (2016).

22. Kohori, A. *et al.* Torque generation in F1-ATPase devoid of the entire
amino-terminal helix of the rotor that fills half of the stator orifice. *Biophys J* **101**,
188–195 (2011).

23. Bowler, M. W. Conformational dynamics in phosphoglycerate kinase, an open
and shut case? *FEBS Lett* **587**, 1878–1883 (2013).

24. Rojas-Pirela, M. *et al.* Phosphoglycerate kinase: structural aspects and functions,
with special emphasis on the enzyme from Kinetoplastea. *Open Biol* **10**, 200302
(2020).

25. Rao, L., Berger, F., Nicholas, M. P. & Gennerich, A. Molecular mechanism of
cytoplasmic dynein tension sensing. *Nature Communications 2019 10:1* **10**, 1–17
(2019).

26. Lu, M., Ammar, D., Ives, H., Albrecht, F. & Gluck, S. L. Physical Interaction
between Aldolase and Vacuolar H+-ATPase Is Essential for the Assembly and
Activity of the Proton Pump. *Journal of Biological Chemistry* **282**, 24495–24503
(2007).

27. Chan, C. Y., Dominguez, D. & Parra, K. J. Regulation of Vacuolar H+-ATPase
(V-ATPase) Reassembly by Glycolysis Flow in 6-Phosphofructo-1-kinase
(PFK-1)-deficient Yeast Cells. *J Biol Chem* **291**, 15820–15829 (2016).

28. Nonaka, T., Adan-Kubo, J. & Miyata, M. Triskelion structure of the Gli521
protein, involved in the gliding mechanism of *Mycoplasma mobile*. *J Bacteriol*
192, 636–642 (2010).

29. Hamaguchi, T., Kawakami, M., Furukawa, H. & Miyata, M. Identification of
novel protein domain for sialyloligosaccharide binding essential to *Mycoplasma*
mobile gliding. *FEMS Microbiol Lett* **366**, fnz016 (2019).

30. Ibuki, T. *et al.* Common architecture of the flagellar type III protein export
apparatus and F- and V-type ATPases. *Nat Struct Mol Biol* **18**, 277–282 (2011).

31. Majewski, D. D. *et al.* Cryo-EM structure of the homohexameric T3SS
ATPase-central stalk complex reveals rotary ATPase-like asymmetry. *Nat*
Commun **10**, 626 (2019).

32. Miyata, M. *et al.* Tree of motility - A proposed history of motility systems in the

tree of life. *Genes. Cell* **25**, 6–21 (2020).

33. Nottelet, P. *et al.* The mycoplasma surface proteins MIB and MIP promote the dissociation of the antibody-antigen interaction. *Sci Adv* **7**, eabf2403 (2021).

34. Aluotto, B. B., Wittler, R., Williams, C. & Faber, J. Standardized bacteriologic techniques for the characterization of Mycoplasma species1, 2. *Int J Syst Evol Microbiol* **20**, 35–58 (1970).

35. Miyata, M. *et al.* Gliding mutants of *Mycoplasma mobile*: relationships between motility and cell morphology, cell adhesion and microcolony formation. *Microbiology (Reading)* **146 (Pt 6)**, 1311–1320 (2000).

36. Uenoyama, A., Seto, S., Nakane, D. & Miyata, M. Regions on Gli349 and Gli521 protein molecules directly involved in movements of *Mycoplasma mobile* gliding machinery, suggested by use of inhibitory antibodies and mutants. *J Bacteriol* **191**, 1982–1985 (2009).

37. Mizutani, M., Tulum, I., Kinosita, Y., Nishizaka, T. & Miyata, M. Detailed analyses of stall force generation in *Mycoplasma mobile* gliding. *Biophys J* **114**, 1411–1419 (2018).

38. Hellman, U., Wernstedt, C., Góñez, J. & Heldin, C. H. Improvement of an 'In-Gel' digestion procedure for the micropreparation of internal protein fragments for amino acid sequencing. *Anal Biochem* **224**, 451–455 (1995).

39. Fujita, J. *et al.* Structures of a FtsZ single protofilament and a double-helical tube in complex with a monobody. *Nat Commun* **14**, 4073 (2023).

40. Mastronarde, D. N. Automated electron microscope tomography using robust prediction of specimen movements. *J Struct Biol* **152**, 36–51 (2005).

41. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017).

42. Zivanov, J. *et al.* A Bayesian approach to single-particle electron cryo-tomography in RELION-4.0. *Elife* **11**, e83724 (2022).

43. Bepler, T., Kelley, K., Noble, A. J. & Berger, B. Topaz-Denoise: general deep denoising models for cryoEM and cryoET. *Nat Commun* **11**, 5208 (2020).

44. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

412   45.   Šali, A. & Blundell, T. L. Comparative protein modelling by satisfaction of
413   spatial restraints. *J Mol Biol* **234**, 779–815 (1993).

414   46.   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory
415   research and analysis. *J Comput Chem* **25**, 1605–1612 (2004).

416   47.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development
417   of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

418   48.   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and
419   crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018).

420   49.   Liebschner, D. *et al.* Macromolecular structure determination using X-rays,
421   neutrons and electrons: recent developments in Phenix. *Acta Crystallogr D Struct*
422   *Biol* **75**, 861–877 (2019).

423   50.   Pintilie, G. *et al.* Measurement of atom resolvability in cryo-EM maps with
424   Q-scores. *Nat Methods* **17**, 328–334 (2020).

425   51.   Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM
426   single-particle structure refinement and map calculation using Servalcat. *Acta*
427   *Crystallogr D Struct Biol* **77**, 1282–1291 (2021).

428   52.   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
429   educators, and developers. *Protein Science* **30**, 70–82 (2021).

Figure legends

Fig. 1 Structure of the twin motor featuring dimeric F_1-like ATPase.

(**a**) Gliding machinery of *Mycoplasma mobile*. Overall view of the gliding machinery

(top) and part of the internal chain (bottom). (**b**) Overall structure of the twin motor. The

illustration is drawn in the upper left. Scale bar = 25 Å. (**c**) Atomic model of the twin

motor. Black triangles represent nucleotides. Maps in (b) and (c) are contoured at 0.55

in UCSF ChimeraX. (**d**) Dimeric F_1-like ATPase in the twin motor. Dotted lines indicate

the axis of each ATPase. (**e**) Density of Mg-ATP bound to G_1-ATPase. The dotted region

in (c) is shown. (**f, h, j**) Subunits constituting F_1-like ATPase ($\alpha_3\beta_3\gamma$ subcomplex of

G_1-ATPase). (**g, i, k**) Overlay of subunits of G_1-ATPase and *Bacillus* PS3 F_1-ATPase.

The subunits of the *Bacillus* PS3 F$_1$-ATPase, αβ (PDB ID: 8HH5) and γ (PDB ID:

8HH2), are coloured purple.

Fig. 2 Interaction between subunits.

(**a**) N- and C-terminal regions of the twin motor. The twin motor is viewed from the top

of the illustration. Arrows indicate the interface between G$_1$-ATPase monomers. Maps

are contoured at 0.55 in UCSF ChimeraX. (**b**) Structure of β and PGK. (**c**) Interaction

between the N-terminal region of β and α-NBD at the interface between G$_1$-ATPase

monomers. The dotted region in (a) is shown. (**d**) Interaction of PGK with the β-NTD.

The dotted region in (a) is shown. (**e**) Structure of the α and GliD monomers. (**f**) GliD

dimer at the interface between G$_1$-ATPase monomers. The figure corresponds to the

dotted region in the illustration. (**g, h**) Interaction between GliD and α-CTD. Each

corresponds to the dotted regions in (f).

Fig. 3 Structural conformations and nucleotide binding of β.

(**a**) Differences in the conformations of the three β. Superposed three βs in the

G$_1$-ATPase. The dotted lines indicate the position of the DELSEED loop of each β. (**b**)

Nucleotide binding pattern of dimeric G$_1$-ATPase and F$_1$-ATPase. The catalytic site is

circled by black dotted lines. (**c**) αβ dimer in each conformation in the hexamer. (**d**)

Comparison of the catalytic sites. Each catalytic site corresponds to the dot square

above. The catalytic sites are superimposed on the corresponding catalytic sites of

F$_1$-ATPase (PDB ID: 8HHA), which are coloured purple.

Fig. 4 Structure and motion of the gliding machinery.

(**a**) Fitting of the atomic model of the twin-motor into the EM map of the motor chain.

The right panel indicates the interface between GliDs. (**b**) Possible rotational catalytic

mechanism of G_1-ATPase. The 120° rotation of γ coupled with the conformational

change in the catalytic hexamer is explained, based on the tri-site mechanism of

F_1-ATPase[8]. γ is indicated by yellow arrows, the direction of which indicates the

conformational tilt. (**c**) A possible explanation for the conversion mechanism from

rotational to linear motion. Dotted arrows indicate unknown force transduction in the

cell membrane. Double-headed arrows indicate the changes in the distance between

twin motors. Gli349 and Gli521 are depicted in red and green, respectively.

Unidentified protein regions in the gliding machinery are coloured purple.

Methods

Optical microscopy

The cells of the *M. mobile* mutant strain (gli521 [P476R]), which can glide like the wild

type strain but binds sialylated oligosaccharides more tightly, were cultured, observed

using phase-contrast microscopy, and recorded as previously described[34–37]. The

recorded video was analysed using ImageJ software v1.54d (https://imagej.nih.gov/ij/).

Isolation of the twin motor

The twin motor was isolated from the *M. mobile* mutant strain (gli521 [P476R]), as

previously described[2]. The twin motor sample with phosphate-buffered saline (PBS)

consisting of 8.1 mM Na_2HPO_4, 1.5 mM KH_2PO_4 (pH 7.3), 2.7 mM KCl and

137 mM NaCl, and 1 mM $MgCl_2$ was concentrated to 1 mg/mL using a 100K Vivaspin

concentrator (Merck, Germany) with 0.1% (w/v) CHAPS to prevent membrane

adsorption. The concentrated twin-motor was subjected to 12.5% SDS-PAGE and

stained using Coomassie brilliant blue R-250. The bands of the component proteins

were identified using peptide mass fingerprinting, as previously described[38].

Cryo-EM grid preparation and data acquisition

An EG-grid was prepared as previously described[39]. A 2.6 µL sample solution was

applied on the grid, automatically blotted from both sides with filter paper at 4 °C for 2

s at 100% humidity, and vitrificated using the semi-automated vitrification device,

Vitrobot Mark IV (Thermo Fisher Scientific, USA). Cryo-EM imaging was performed

using a CRYO ARM 300 (JEOL, Japan) operated at 300 kV with a K3 direct electron

detector camera (AMETEK, Gatan, USA). A total of 7,350 movies were recorded using

SerialEM software[40] with a total dose of approximately 80 electrons $Å^{-2}$ for 40 frames,

an exposure time of 3.3 s per movie, and a nominal defocus range of -0.8 to -1.8 µm.

The nominal magnification was 60,000×, corresponding to 0.87 Å per pixel.

Cryo-EM image processing

The image processing steps for SPA are summarised in Extended Data Figs. 3 and 4.

Image processing was performed using cryoSPARC v3.3.2[41], unless otherwise stated.

Movies were aligned using patch-based motion correction, and the contrast transfer

function (CTF) parameters were estimated using patch CTF estimation. Micrographs

with a CTF fit resolution worse than 8 Å were removed. Particles were automatically

picked up using a Blob Picker, extracted with ×4 binning, and 2D-classified. The

obtained 2D averaged images were used as templates for the Template Picker to pick up

good particles. After two rounds of 2D classification, the selected good particles were

subjected to local motion correction and re-extracted with the original pixel size. The

particles were subjected to Ab-initio reconstruction using C1 symmetry enforced with a

final resolution limit of 12 Å and heterogeneous reconstruction with 3 classes.

Particles in class 2 were subjected to homogeneous reconstruction using C1 symmetry,

resulting in a map at a resolution of 3.5 Å. The final map was generated via local and

global CTF refinement, and then non-uniform refinement. To improve the local

resolution, the final map and particles were subjected to particle subtraction and local

refinement using C1 symmetry to generate half-divided maps, each containing a

monomeric ATPase. The whole and two local maps were subjected to mask creation and

postprocessing in RELION 4.0[42], resulting in resolutions of 3.2, 3.1, and 3.1 Å,

respectively. Local resolution estimation was also performed in RELION 4.0. The

post-processed map was denoised using Topaz software v0.2.5 with the trained model

Unet-3d-10a[43]. RELION 4.0 was used to calculate the Fourier shell correlation (FSC)

curve between two half-maps and estimate the local resolution.

Modelling

Initial models for α, β, γ, GliD, and GliE were generated using AlphaFold2[44], whereas

the homology model for PGK was generated using Modeller v10.4[45] with the atomic

model of PGK (PDB ID: 4DG5) from *S. aureus*. The generated models were fitted to

the corresponding densities of the EM map as rigid bodies using UCSF Chimera v1.15[46].

The fitted models were manually inspected and adjusted using COOT v0.9.8.1[47] and

refined using the *phenix.real_space_refine* program in PHENIX v1.19[48,49] with

reference model restraints using atomic models of F_1-ATPase (PDB ID: 5IK2) from

Caldalaklibacillus thermarum and PGK (PDB ID: 3ZLB) from *Streptococcus*

pneumoniae. Manual adjustment using COOT and refinement using PHENIX were

repeated until model parameters were no longer improved. For the modelling of PGK3,

the atomic model of PGK1 was fitted to the PGK3 map and then manually inspected

using COOT. FSC curves between the map and model were calculated using PHENIX.

The refined model was also evaluated via comprehensive validation in PHENIX and the

Q-score[50]. To identify the nucleotide binding sites of G_1-ATPases, F_o-F_c maps were

generated using Servalcat[51]. Superposition of models and fitting of models to the map

were performed using UCSF ChimeraX v1.2.5[52]. All structural figures were obtained

using UCSF ChimeraX.

Data availability

548   The cryo-EM dataset for the twin motor has been deposited in the Electron Microscopy

549   Public Image Archive (EMPIAR) under accession code: EMPIAR-12340. The cryo-EM

550   map is available on the Electron Microscopy Data Bank (EMDB) under accession code:

551   EMD-60718. The coordinates have been deposited in the Protein Data Bank (PDB)

552   under accession code: 9IO5.

Acknowledgments

We thank Tomomi Shimonaka at Osaka Metropolitan University and Fumiaki Makino at

Osaka University for their technical assistance. This manuscript was edited using

Copilot, an AI tool. This study was supported by Grants-in-Aid for Scientific Research

(A) (JP17H01544), a JST CREST grant (JPMJCR19S5) to M.M., and Research Support

Project for Life Science and Drug Discovery (BINDS) from AMED under Grant

Number, JP22am121003, to K.N., and JEOL YOKOGUSHI Research Alliance

Laboratories of Osaka University to K.N.

Author contributions

T.T. and M.M. conceived the strategy. M.M. and K.N. supervised the research. T.T.

prepared the cryo-EM samples and carried out the biochemical analyses. J.F. prepared

EG-grid for cryo-EM observation. A.K. and T.M. performed cryo-EM. T.T. carried out

3D reconstruction and model building. T.K., T.H., A.K., T.M., K.K., and K.N.

participated in the discussion regarding the structural analyses. T.T. and M.M. wrote the

manuscript with input from all authors.

Competing interests

572 The authors have declared no competing interest.

a Internal chain

Sialylated oligosaccharides
Host cell surface
Leg

Bridge
Twin motor

b GliE GliD β PGK α

PGK3 GliD4 β6 PGK6
GliD3 α4 γ2 PGK
GliE α3
PGK1 β2 α5
β1 GliD5
GliD2 α2
α1
γ1 β3 α6 β5 PGK2
PGK5 β1 PGK4
GliD1 GliD6

c PGK3 GliD4 PGK6
GliD3 β6 α4
GliE α3 γ2
PGK1 β2 β4
α2 α5
GliD2 β1 GliD5
γ1 α1
β3 α6
PGK5 β5 PGK2
GliD1 GliD6 PGK4

Side view
90°

PGK1 α3 β3 β4 α5
β2 α2 β6 α4 PGK6
PGK2
GliD2 GliE GliD5
γ1 GliD3 GliD4 γ2
PGK3

d F₁-like ATPase   F₁-like ATPase
β2 α3 β3 β6 α4 β4
α2 α5
γ1 γ2
~5°

e β1-E484
Mg β1-R485
β1-K458 β1-R487
α1-R344
ATP

f G₁-β2
Crown Linker
NTD
NBD
P-loop N
CTD C
DELSEED loop (DALSEDD)

g G₁-β2 vs. *Bacillus* PS3 F₁-β
N
N
C
C

h G₁-α2
Crown N
NBD
P-loop
CTD
C

i G₁-α2 vs. *Bacillus* PS3 F₁-α
N N
C
C

j G₁-γ1
C
N
α/β fold

k G₁-γ1 vs. *Bacillus* PS3 F₁-γ
Extended helix
C
C
N N

Fig. 1

Fig. 2

Fig. 3

Fig. 4