Comprehensive profiling of transcriptional regulation in cartilage reveals pathogenesis of osteoarthritis

Wen Tian[1#], Shan-Shan Dong[1#], Feng Jiang[1#], Chang-Yi He[1], Chen Wang[1], Shou-Ye Hu[2], Ruo-Han Hao[1], Hui-Miao Song[1], Hui-Wu Gao[1], Jun-Qi Zhang[1], Ke An[1], Dong-Li Zhu[1], Zhi Yang[2], Yan Guo[1*], Tie-Lin Yang[1*]

[1]Key Laboratory of Biomedical Information Engineering of Ministry of Education, Biomedical Informatics & Genomics Center, School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, Shaanxi 710049, P.R. China

[2]Honghui Hospital, Xi'an Jiaotong University, Xi'an, Shaanxi 710054, P.R. China

[#]These authors contributed equally

[*]Correspondence authors:

Tie-Lin Yang, Ph.D., Yan Guo, Ph.D.

E-MAIL: yangtielin@xjtu.edu.cn (T.-L.Y.), guoyan253@xjtu.edu.cn (Y.G.)

Conflict of Interest: The authors declare that they have no competing interests.

Abstract

Cartilage damage is a leading cause of osteoarthritis (OA) etiology, however, the underlying mechanism governing gene expression regulation in this progress is poorly understood. Here, we described a comprehensive profiling of transcriptional regulation of 234 primary human cartilage samples. Based on speculated chondrocyte subtype proportions, we demonstrated the heterogeneity of cell composition contributed to the variance of gene expression across samples. We identified 3,232 significant expression quantitative trait loci (eQTLs) for 2,999 genes. Cell type-interaction eQTL (ci-eQTL) analyses revealed the regulation relationship of 81 eQTL-gene pairs showing cell type dependency. Further, by integrating with genome-wide association studies (GWASs) data of OA, we nominated 55 candidate effector genes, of which, 26 were novel genes that have not been reported by previous studies. Finally, we established a multi-level functional variants-deciphering pipeline by integrating the cartilage-specific ATAC-seq data, and identified 112 causal eQTLs that display allele-specific open chromatin (ASoC), as well as 410 transcription factor binding-disruption (TBD) eQTLs. We experimentally verified that the T allele at rs11750646, which was located within a GWAS-eQTL co-localized region, increased the AR binding affinity to an open chromatin region and promoted the expression of an OA-related gene *PIK3R1*. Altogether, our findings provide new insights into the unique regulatory landscape of cartilage and elucidate potential mechanisms underlying the OA pathogenesis.

Key word: Osteoarthritis; Cartilage; eQTL; Fine-mapping

Introduction

Osteoarthritis (OA) is the most prevalent chronic joint disease, which causes pain, disability and loss of function, and affects over 500 million people worldwide[1]. Knee is the leading site of OA, followed by hand and hip[1,2]. In the process of OA, articular cartilage is both the primary victim and betrayer[2]. Figuring out the role of articular cartilage in OA and its specific mechanisms, especially underlying gene expression regulation, is critical to developing efficient therapies.

Genome-wide association studies (GWASs) have reported more than 100 susceptibility loci for OA[3-9], however, the majority of SNPs map to noncoding regions and the downstream effector genes in articular cartilage have not been well clarified. Expression quantitative trait loci (eQTLs) data provide a valuable tool for screening effector genes for noncoding SNPs. The first eQTL regulation landscape of OA articular cartilage has been constructed from 99 European patients[10], and identified a total of 1,453 genes with eQTLs, which allows the nomination of four effector genes of OA GWAS loci. However, the power to detect mechanistically informative expression signals rely on sufficient numbers of samples in relevant tissues[11], a larger sample size of articular cartilage is needed to obtain more statistically significant eQTLs. Furthermore, given the discrepancy of minor allele frequencies (MAF) in different populations, a cartilage eQTL resource from non-European population will provide additional information to elucidate the gene expression roles of GWAS variants.

Previous studies have shown that some of the eQTL effects are cell type-specific[12-16].

Chondrocytes have several subtypes[17-19], but their genetic regulatory heterogeneity

remains unknown. As an effort to solve this issue, an approach based on the in silico

speculated cell subtype proportions from bulk RNA-seq data has been developed to

reveal the cell type dependency of eQTL effects[20] and has been broadly used for bulk

RNA-seq data from several tissues[21-24].

In this study, we performed comprehensive characterization for transcriptional

regulation of knee cartilage samples from 204 OA patients (Fig. 1). Based on the

deconvoluted chondrocyte subtypes proportions from bulk RNA-seq data, we

evaluated the contribution of cell composition to the overall gene expression variance

across samples and its contribution to the differential expression between damaged

and intact cartilage samples. We not only identified *cis*-eQTLs in this key tissue, but

also evaluated the dependence of the *cis*-eQTLs effect on the chondrocyte subtypes.

Next, we nominated the effector genes for OA GWAS loci and annotated their

potential function using multiple lines of evidence. Finally, leveraging the

cartilage-specific epigenomic data and our multi-level functional variants-deciphering

pipeline, we conducted functional fine-mapping to reveal candidate causal variants

that linked both to OA risk and gene expression regulation through transcription

factors (TFs) binding site disruption or chromatin accessibility alteration. In all, our

study represented a valuable eQTL resource in cartilage, and provide new insights

into how genetics contributes to the pathogenesis of OA.

Results

Contribution of chondrocyte composition to the expression variance of cartilage

samples

We generated 235 qualified knee cartilage RNA sequencing (RNA-seq) data from 204

Chinese Han individuals undergoing joint replacement for OA with extensive quality

controls (Fig. 1 and Supplementary Tables 1,2). We obtained a median of 14,809

expressed protein coding genes, 3,407 long non-coding RNA (lncRNA) genes and

1,915 genes of other types (expressed gene was defined as TPM $\geq$ 0.1, Supplementary

Fig. 1a). The most of protein coding genes showed high expressions (TPM $\geq$ 10),

while the majority of non-coding genes had low expressions (TPM $<$ 1)

(Supplementary Fig. 1b).

Using a liner mixed model-based method[25], we compared the expression between the

paired damaged and intact cartilage samples from 31 donors. We observed 334 genes

significantly high expressed in damaged cartilage, and 715 genes significantly high

expressed in intact cartilage (FDR $<$ 0.05, no fold change cutoff; Fig. 2a and

Supplementary Table 4). Gene set enrichment analyses showed that the genes

expressed lower in damaged cartilage were significantly enriched in the cell markers

of chondrocyte subtypes, effector chondrocytes (EC) and regulatory chondrocytes

(RegC) (Fig. 2b,c). Cell type composition differences have been proven to be a vital

factor for driving gene expression variance in bulk RNA-seq data[26]. Cartilage consists

of multiple subtypes of chondrocyte[17-19]. To evaluate the contribution of chondrocyte

composition to the overall variance of gene expression, we deconvoluted our bulk

gene expression data (204 from damaged cartilage and 31 from intact cartilage) into

chondrocyte subtype proportions based on the expression signature of each subtype

extracted from scRNA-seq data[19] (Fig. 2d,e and Supplementary Table 5). By

comparing the speculated cell proportions in the paired damaged and intact cartilage

samples, our results showed that the damaged cartilage samples had significantly

lower proportions in EC and RegC, and significantly higher proportion in

prehypertrophic chondrocytes (preHTC) (Fig. 2f), which were consistent with the

findings from scRNA-seq data[19] and enrichment analyses.

Next, for each gene, we estimated the proportion of expression variance that could be

explained by each chondrocyte subtype. After adjusting multiple common biological

factors and artificial factors, the variance could be largely explained by proportion of

chondrocyte subtypes (Fig. 2g). In combination, the median of variance explained by

all six chondrocyte subtypes was 21.3% (Fig. 2h), indicating the great impacts of

chondrocytes composition heterogeneity on the expression variance.

***Cis*-eQTL mapping**

Regular cis-eQTL

To identify *cis*-eQTLs in cartilage, we generated genome-wide genotype data from

peripheral blood samples of all participants (Supplementary Tables 1,3). The genotype

data showed no significant stratification (Supplementary Fig. 2a), and the minor allele

frequencies (MAF) were strongly correlated with that of Asians but weaker correlated

with that of Europeans in the 1000 Genomes Project (Supplementary Fig. 2b.c). We

then performed *cis*-eQTL mapping by integrating the chondrocyte subtype

proportions as covariates to correct for the impact of cell type heterogeneity. Totally,

we identified 2,999 genes (eGenes) with at least one genome variant (eVariant)

significantly associated with its expression at 5% false discovery rate (FDR) level

(Fig. 3a and Supplementary Table 6), including 2,449 (81.7%) protein coding genes,

385 (12.8%) lncRNAs and 165 (5.5%) of other types (Fig. 3b and Supplementary

Table 7). Compared with non-eGenes, eGenes showed significantly higher expression

($P < 2.2 \times 10^{-16}$; Wilcoxon rank sum test; Fig. 3c) and lower expression variation

coefficient ($P < 2.2 \times 10^{-16}$; Wilcoxon rank sum test; Fig. 3c).

Conditional analysis

To identify independent eQTL signals, we conducted conditional analysis and found

that 216 of 2,999 eGenes had more than one independent eQTL signals (Fig. 3d and

Supplementary Table 8). Importantly, 98 eGenes (45.4%) had at least one newly

identified independent signal which could not be detected by regular eQTL analysis

(Fig. 3e). For example, rs623263 showed significant association with the *SNAP91*

expression only when conditional on the effect of rs145051363, which tagged the

other independent signal (Fig. 3f). We compared the characteristics of the newly

identified independent lead eVariants (the most significant eVariant for each eGene)

by conditional analysis with those significant lead eVariants by regular eQTL analysis.

In general, the newly identified lead eVariants showed significantly lower MAF ($P =$

1.7×10^{-4}; Wilcoxon rank sum test; Supplementary Fig. 3a) and longer distance to

transcription start sites (TSS) of the target genes ($P = 2.6 \times 10^{-2}$; Wilcoxon rank sum

test; Supplementary Fig. 3b), but no difference in effect size for the target genes ($P =$

0.404; Wilcoxon rank sum test; Supplementary Fig. 3c). Down sampling analysis

showed that conditional analysis with our larger sample size could help uncover more

independent association signals (Supplementary Fig. 4).

For the 3,232 unique independent lead eVariants, the largest part (1,606, 49.7%) was

located in the intron regions, followed by the intergenic regions (747, 23.1%),

promoter regions (607, 18.8%), and exon regions (272, 8.4%) (Fig. 3g). Lead

eVariants were almost symmetrically concentrated on both sides of the TSS of their

target genes (Fig. 3h). In total, 1,978 (61.2%) lead eVariants were located within ± 50

Kb from the TSS. However, there were also 754 lead eVariants (23.3%) located > 100

Kb from the TSS of the target genes, suggesting the long-range regulation models

between eVariants and their target genes.

To assess eQTL reproducibility, we compared our results with a previously published

cartilage study involving approximately 90 European individuals[10]. Despite the high

direction consistency of regulatory effect of the shared variant-gene pairs, our data

also uncovered a large number of pairs that were specifically significant in our study

(Supplementary Fig. 5). This was partly attributable to the larger sample size of our

study (Supplementary Fig. 6). Besides, the difference in MAF between the Chinese

and European populations was also a considerable factor. Among the 236,922 specific

eVariants identified in our study, 13,568 (5.76%) and 4,300 (1.83%) with a MAF <

0.01 or < 0.001 in the 1000 Genomes Project European population, respectively

(Supplementary Fig. 7). This meant that if the minimum minor allele count was set to

10, as used in this study, it would take about 1,000 and 10,000 samples to calculate

their effects on the expression of *cis*-genes, respectively, which exceed the maximum

sample size of any single tissue in the GTEx project (Muscle-Skeletal, 803 samples,

v8)[27], posing a great challenge in terms of sample collection and highlighting the

values of eQTL mapping in different populations.

Characteristics of eVariants

We assessed the functional characteristics of the identified eVariants using

epigenomic data from human knee cartilage[28] and human mesenchymal stem cells

(hMSCs) derived chondrocytes[29]. As expected, eVariants were significantly enriched

in open chromatin regions with ATAC-seq peaks (Fig. 3i). Significant enrichments

were also observed in multiple active histone markers (H3K4me3, H3K27ac,

H3K4me1, and H3K36me3) (Fig. 3i). Specifically, we found that the eVariants

exhibited stronger enrichment in super-enhancers (SEs) than that in typical enhancers

(TEs) (Fig. 3j), suggesting a deeper implication of SEs in genetic regulation of gene

expression.

Genetically regulatory heterogeneity among chondrocyte subtypes

Genetic regulation of gene expression exhibited a broad cellular specificity[12-16]. To

explore whether there was regulatory heterogeneity among chondrocytes subtypes, we

performed cell type-interaction eQTL analyses (ci-eQTL), from which the possible

driver or confirmable non-driver cell types for a single variant-gene regulatory pair

would be revealed (Fig. 4a; Methods). After applying to three widely distributed

chondrocyte subtypes (EC, preHTC, and RegC) in our deconvolution results (Fig.

2d,e), we identified 29, 23 and 29 genes with significant ci-eQTLs for EC, preHTC

and RegC, respectively (Supplementary Tables 9-11). Of which, most of variant-gene

regulatory pairs were aligned to the driver model (Fig. 4b).

We estimated the sharing and specificity of all 81 significant cell type-interaction

variant-gene pairs using multivariate adaptive shrinkage (mash)[30] analysis. Totally, we

obtained 69 variant-gene pairs with local false sign rate (LFSR) < 0.05 in at least one

cell type (Supplementary Table 12). We clustered the 69 pairs into 11 clusters

according to their model type and mash posterior effect size (Fig. 4c), and observed

similar pattern for EC and RegC, opposite pattern for preHTC, which were consistent

with the correlation between their proportions that speculated from our samples

(Supplementary Fig. 8). We found 7, 6 and 11 cell type-specific pairs that aligned to

driver model for preHTC, RegC and EC, respectively. As examples, the expression of

the *IGFBP2* showed a significantly positive correlation with the count of T allele at

rs12620516 (Fig. 4d). Consistently, with the proportion of preHTC increased, T allele

carriers exhibited significantly higher *IGFBP2* expression, but no such pattern was

observed in the EC and RegC (Fig. 4e). The *IGFBP2* was specifically expressed in

preHTC in the scRNA-seq data of knee cartilage[19] (Supplementary Fig. 9), which

could be the cause for cell type-specific eQTL effects[23]. Similarly, RegC and EC were

the diver cell types of the regulatory pairs rs2356553~*MAST2* (Fig. 4f,g) and

rs1408829~*TOR3A* (Fig. 4h,i), respectively. These findings demonstrate the

genetically regulatory heterogeneity of gene expression among chondrocytes

subtypes.

eQTL-GWAS integration nominates risk genes for OA

To date, GWASs have successfully discovered numerous genetic loci associated with

OA, yet their underlying gene expression regulatory mechanisms remain elusive. Here,

we performed co-localization analysis to integrate our cartilage eQTL data (regular,

conditional and ci-eQTL results) and 28 publicly available GWAS summary statistics

of OA from six studies (Supplementary Table 13)[4-9]. In total, we identified 55 genes

for which the corresponding eQTL signals showed strong evidence (PPH4 $\geq$ 0.7) that

shared a common causal variant with GWAS signals (Fig. 5a, Supplementary Fig. 10

and Supplementary Table 14). Specifically, our eQTL data supplemented 26 new

genes to the candidate OA risk gene list (gene symbols were shown in red in Fig. 5b),

which could not be nominated from eQTL data of any tissues in the GTEx project and

previous cartilage eQTL data (Fig. 5b and Supplementary Figs. 11-12), revealing the

sufficient power of our relatively larger sample size in cartilage.

Our eQTL data linked OA GWAS signals to 14 functionally known OA related genes

(Fig. 5b and Supplementary Table 15), including five genes showed differential

expression between paired damaged and intact cartilage in our data or abnormal

cartilage/chondrocytes related phenotypes in gene knockout mice (Fig. 5b and

Supplementary Table 16). For example, *COLGALT2* showed a significant lower

expression in our damaged cartilage samples ($\log_2$FC = -1.61, *P* = 6.19 × 10^{-4},

adjusted *P* = 0.023; Supplementary Table 15). Previous study has reported that

COLGALT2 initiates collagen glycosylation to organize and stabilizes collagen triple

helix[31]. Particularly, our data nominated four functionally known genes (*IGF1R*,

PIK3R1, *SERPINF1* and *WNT3*) that could not be nominated by eQTL data of GTEx

tissues and previously reported cartilage tissue. For example, *IGF1R* is located on the

cartilage cell membrane and could be activated by IGF1, and further inhibit cartilage

matrix degradation and maintain the homeostasis of articular cartilage[32,33].

To reveal biological processes involved for more functionally unknown effector genes,

we generated gene sets to annotate their potential functions based on a time series

expression data of chondrocytes differentiation[34]. Briefly, we classified all

differentially expressed genes into 12 clusters to shown the overall tendency during

the chondrocyte differentiation (Supplementary Fig. 13a and Supplementary Table 17),

and a total of 23 co-localized genes belonged to these gene clusters (Fig. 5b). We

performed gene ontology (GO) enrichment analyses to reveal the primary function of

each cluster, and identified four clusters (cluster 3, 6, 7 and 12) enriched in biological

processes related to chondrocytes differentiation (Supplementary Fig. 13b),

suggesting the potential function of genes involved in these cluster. Especially, five

newly functionally unknown effector genes (*FILIP1*, *OGFOD2*, *MMP23B*,

CTD-2287O16.5 and *SLC44A2*) belonged to these four clusters (Fig 5b and

Supplementary Fig. 13a). In addition, for another seven new effectors genes (*ALAS1*,

NRXN2, *NSFP1*, *VAC14*, *ABCA9*, *MARVELD2* and *SLC35E2B*) belonged to three

clusters (cluster 1, 9 and 10), although the top enriched biological processes of the

gene clusters they belonged to were not directly associated with cartilage formation,

their expressions were gradually decreased during chondrocyte differentiation

(Supplementary Fig. 13a), suggesting their potential roles involved in OA. Taken

together, leveraging our tissue match eQTL data, we provide a high confidence OA

risk gene list for future research.

Functional fine-mapping reveal candidate causal variants underlying both OA

risk and cartilage gene expression

To fill the mechanistic gap between the eVariants and their target genes expression,

we established a multi-level functional variants-deciphering pipeline to evaluate

whether they act by altering the chromatin accessibility of regulatory elements or

disrupting the TF binding to the regulatory elements.

Allele-specific open chromatin (ASoC) eVariants

We first explored the impact of eVariants on the chromatin accessibility (Fig. 6a).

Using the read coverage of heterozygous sites called from knee cartilage ATAC-seq

data[35,36] (Supplementary Fig. 14; Methods), we identified 1,170 allele-specific open

chromatin (ASoC) variants showing significant imbalance between the two alleles

(nominal $P \leq 0.05$) (Supplementary Fig. 15 and Supplementary Table 18), which was

limited by the low heterozygosity rate of variants in the ATAC-seq data sets. A total of

112 eVariants were annotated as ASoC variants and they associated to a total of the

expression of 129 eGenes (Fig. 6b and Supplementary Table 18).

ASoC annotation of eVariants allowed us to interpret the underlying mechanism of

OA GWAS loci. For example, the eVariant rs3810154 was associated with the

expression of *SLC44A2* (Fig. 6c), which was assigned as an OA risk gene by

co-localization analyses (Fig. 6d). Intriguingly, rs3810154 was located in an open

region downstream *SLC44A2* and the local chromatin region around the G allele at

rs3810154 showed significantly higher accessibility than that of A allele (44 reads for

G allele; 20 reads for A; $P = 3.69 \times 10^{-3}$; binomial test; Fig. 6e), suggesting that

rs3810154 regulates *SLC44A2* expression by altering the chromatin accessibility. Our

soft-clustering and functional enrichment analyses revealed that *SLC44A2* belonged to

a gene cluster with increased expression during chondrocytes differentiation (Fig. 6f,

Supplementary Fig. 13), indicating its potential important role in cartilage

development (Supplementary Fig. 13b).

TF binding-disruption (TBD) eVariants

We and others had revealed the impacts of genetic variants on disrupting the TF

binding on DNA and the consequent gene expression alteration[37-42]. As shown in Fig.

3i, eVariants were significantly enriched in the H3K4me3-defined promoters and

H3K27ac-definded enhancers, which load TF binding motifs. We further investigated

the effect of these eVariants on the TF binding. Due to the lack of ChIP-seq data of

TFs in cartilage tissues, we established an allele-sensitive TF footprint analysis

method, which considered the TF binding specificity to both the reference allele and

alternative allele of each variant in motif scan step (Fig. 7a; Methods), to speculate the

binding sites for 441 expressed TFs (median TPM $\geq$ 1) in our damaged cartilage

samples (Supplementary Fig. 16a and Supplementary Table 19). We identified 222 to

39,416 non-overlapped unique binding sites for each TF (Supplementary Fig. 16b and

Supplementary Table 19). Comparing to the traditional footprint analysis, our allele

sensitive method could detect a total of 22,627 new footprints for all 673 motifs used

for analysis (Supplementary Fig. 17). To validate accuracy of the footprint results, we

used the CTCF ChIP-seq data from cartilage samples, and the high validation rates

were achieved for all five motifs of CTCF (Supplementary Fig. 18), indicating the

reliability of our method.

Next, we used a TF motif information content-based method to investigate the

potential motif disruption effects of eVariants (Fig. 7a; Methods). A total of 1087

variants were defined as candidate TBD variants (Supplementary Table 20). Further,

we performed a TF expression-interaction eQTL analysis (TFi-eQTL) to assess

whether the TF for which the motif was disrupted by a variant was the true mediator

of the eQTL effect (Fig. 7a; Methods). Briefly, we constructed a linear-mixed

regression model with an interaction term of genotype dosage and TF expression to

detect the effect of the eVariant on the co-expression of TF and target gene. As a result,

we identified 410 TBD eVariants that might regulate the expression of 369 eGenes

through the disruption of binding sites of 231 TFs with a nominal significance

threshold (Fig. 7b and Supplementary Table 21). These results allowed us to explain

the underlying regulation mechanisms of candidate causal eVariants on seven

co-localized effector genes (*PIK3R1*, *GNL3*, *hsa-mir-8072*, *TRIOBP*, *SLC44A2*,

RNF212 and *PCCB*) (Supplementary Table 20). As a representative example,

rs11750646 was located within a co-localized signal region of hip OA and cartilage

eQTL (Fig. 7c). The T allele at rs11750646 significantly increased the expression of

PIK3R1 ($P = 4.53 \times 10^{-8}$; Fig. 7d). It was potential to disrupt the motifs of five TFs

(AR, FOXA2, FOXA3, FOXM1 and FOXO1) that left footprints in an open region

downstream the *PIK3R1* (Fig. 7e,f and Supplementary Table 20). TFi-eQTL results

showed that rs11750646 most significantly altered the co-expression relationship

between *AR* and *PIK3R1* ($P = 2.81 \times 10^{-4}$; Fig. 7g and Supplementary Table 21),

suggesting that *AR* was the true regulator affected by this SNP.

We further performed functional experiments to validate our hypothesis derived from

our data analyses. Using human chondrocyte-like cell line SW1353, a commonly used

cell model for OA *in vitro*, the dual-luciferase reporter assay revealed that

rs11750646-T contained sequence showed significant enhancement ability on target

gene *PIK3R1*, while rs11750646-C contained sequence showed slightly inhibiting

ability for target gene expression (Fig. 7h,i). Further, in line with in silico motif

analysis results, our ChIP-PCR followed by Sanger sequencing results proved that AR

preferentially bound to the T allele at rs11750646 *in vivo* in SW1353 cells (Fig. 7j).

Knockdown of AR significantly repressed *PIK3R1* expression (Fig. 7k).

Dual-luciferase reporter assay showed that the gene expression-promoting role of

rs11750646-T contained fragment was significantly abolished after AR knockdown,

while rs11750646-C contained fragment had a relatively weaker change (Fig. 7l). The

PIK3R1 encoded the regulatory subunit 1 of phosphoinositide-3-kinase (PI3K), which

have been reported to regulate the chondrocyte apoptosis through PI3K/Akt pathway

and thus involved in development of OA[43]. Collectively, our integrated computational

and functional results suggested that AR could allele-preferentially bind the protective

T allele (Beta = -0.0521 in GWAS) at rs11750646 to reinforce *PI3KR1* expression and

thus associated with OA (Fig. 7m). In addition, it was noteworthy that the

binding-preferred allele for AR at this SNP was T, which is an alternative allele that is

not presented in human reference genome sequence version hg19 (Fig. 7f) and this

AR footprint only identified with our allele-sensitive footprint analysis method,

demonstrating its great value in uncovering biological mechanisms of genomic

variants.

Discussion

In this study, we presented the largest sample size of human knee cartilage reported to

date. We found the chondrocyte subtype composition heterogeneity vastly contributed

to the expression variance across samples, which also partially explained the

differential expression between damaged and intact cartilage samples. We reported the

eQTLs for cartilage gene and investigated their cell type dependency. By integrating

with OA GWAS data and epigenetic data of cartilage samples, we extended the list of

OA risk genes and prioritized the potential causal variants linked both to the OA

susceptibility and gene expression regulation.

For the high-throughput bulk RNA-seq generated expression data, the reported

expression levels are the average number of all cell types in each sample. Thus, cell

composition is an important factor that may contribute to the expression variance of

heterogeneous tissues. Although chondrocytes cell type composition of cartilage

tissue is relatively simple when compared to other tissues, our results still illustrated

the difference of chondrocyte subtypes proportion explained a notable part of variance

of gene expression across samples. Importantly, we revealed that the differentially

expressed genes between damaged and intact cartilage samples using bulk RNA-seq

data significantly enriched in the marker gene list of cell types (EC and RegC) with

significant composition difference between damaged and intact cartilage. According

396 to previous study, the main distribution area of EC and RegC is the cartilage

397 superficial zone[19], which suffered abrasion during the development of OA, so it is

398 hard to say whether the cell type composition change is the cause of OA. Our results

399 will prompt the need to consider the non-negligible impact of cell composition on the

400 differential expression analysis and the interpretation of results.

402 As an efficient approach, eQTL analysis has been widely used to investigate the

403 genetic impact on the gene expression of human tissues or cell types. To date, the

404 GTEx project has generated eQTL data resources for 49 different human tissues. For

405 some easily accessible samples, such as blood, a considerable number of eQTL

406 studies have been reported and obtained a great number of samples. All of these have

407 greatly facilitated the discovery of disease risk genes. As for OA research, Julia et al.

408 published the only eQTL so far for articular cartilage, which is the main tissue related

409 to OA[10]. In our present study, we doubled the sample size and reported a larger

410 number of significant eQTLs for cartilage samples, and significantly complemented

411 the previous research. We also successfully distinguished the independent eQTL

412 signal from each other for 216 genes and further reported the capacity of conditional

413 analysis to uncover the eQTL signals that have been missed in the regular eQTL

414 analysis, indicating a new value for conditional analysis.

416 Genetic regulatory heterogeneity among chondrocyte subtypes remains poorly

417 understood. By introducing the interaction item of genome dosage and cell proportion

proxy to the regression model, it enabled the cell type dependency test for eQTLs

without using single cell expression data or bulk expression data from sorted cell of

single type[20]. An increasing number of studies based on this strategy have revealed

specifically genetic regulation effect on gene expression between different cell

types[20,21,44]. In this study, we indeed demonstrated the genetic regulatory

heterogeneity between three chondrocyte subtypes, much like the heterogeneity

reported in immune cell subpopulations[45]. We believe it would stimulate future

research in this aspect.

Integration of eQTL and GWAS is a powerful method for understanding the

molecular mechanism influencing complex traits. The tissue non-matched eQTL data

from the GTEx project have broadly been used for co-localization analyses to reveal

the gene expression regulation role of OA GWAS loci[5,6], but the tissue specific

genetic regulation effect may introduce inevitable false nomination of true effector

genes. While efforts have been made to nominate effector genes using cartilage

expression data derived annotation information, limitations still remained challenging

to overcome. For example, Hollander et al. reported the transcriptome-wide allelic

imbalance data in OA cartilage to link the GWAS variants to gene expression[46], but

only variants in exons could be evaluated with this method. Further, existing OA

studies have limited the nomination of risk genes using colocalization analysis

because of their relatively small sample size. Here, by integrating our cartilage eQTL

data and OA GWAS data, we nominated 26 novel OA risk genes that could not be

found by other studies. Our data presented in this study helped to make a big step in

OA pathogenic mechanism analysis.

We also made efforts to fill the mechanism gap between genomic variants and gene

expression. We first revealed that some eVariants were involved in the alteration of

chromatin accessibility. As a prerequisite for genomic regulatory functions, open

chromatin allowed the binding and assembling of transcription machines, including

TFs, co-factors and RNA polymerase. Increasing number of studies have reported the

chromatin accessibility quantitative trait loci (caQTLs)[47-50], through which could link

the GWAS loci to chromatin accessibility regulation and further to gene expression

alteration. However, there are no such resources for cartilage samples. As a

compromise, we generated an allele-specific open chromatin SNP set using the

publicly available ATAC-seq data from cartilage samples. Although it limited the

detectable number of variants that affect chromatin accessibility, we still found the

variant rs3810154 linked both to OA susceptibility and *SLC44A2* expression via the

alteration of chromatin accessibility. We believe future caQTL data in cartilage would

promote the discovery of variants that regulate gene expression through the regulation

of chromatin accessibility.

Second, we established an allele-sensitive TF footprint analysis method and then

identified footprints for 441 expressed TFs in cartilage samples. Here, we verified one

example. The variant rs11750646, associating both with OA susceptibility and

PIK3R1 expression, had a disruption effect on the DNA binding of AR. Notably, this

footprint of AR was only identified when using the alternative allele-replaced genome

sequence for footprint analysis, it was due to the strong conservation of the alternative

allele T in the motif. The allele T is the minor allele at rs11750646, the frequency in

our samples was 0.1866, and its frequency in 1000 genome European samples was

0.3598, indicating it is a very common allele, emphasizing the great value of our

allele sensitive footprint analysis method. Although more footprints were identified,

our result still underestimated the value of allele sensitive footprint analysis method

because in the current study, we only considered the variants genotyped in our

samples, which only represented a small part of the variants across whole genome

sequence.

In summary, we performed comprehensive profiling for transcriptional regulation in

knee cartilage from OA patients, reported the largest cartilage eQTL dataset to date,

and nominated new risk genes for OA. We revealed potential causal variants for both

OA risk and gene regulation based on epigenetic feature resources identified from

cartilage specific data. Our study provides a valuable resource to elucidate regulatory

mechanisms for genes underlying OA etiology.

Methods

Sample collection

All patients clinically diagnosed with OA and required total knee replacement surgery

from 2019–2022 in Xi'an Honghui Hospital were eligible for this study after signing

the informed consent. During total knee replacement surgery, we isolated the knee

damaged or/and intact cartilage with scalpels after the knee joint was removed from

the donors and froze it in liquid nitrogen rapidly followed by transferred to a -80□

refrigerator for storage. Besides, we collected 2-3 ml peripheral blood from the

participants. The summary of clinical information for donors, including age, sex was

shown in Supplementary Table S1.

DNA isolation, genotyping and imputation

DNA was extracted from whole blood using FlexGen Blood DNA Kit (CWBIO)

according to the manufacturer's instructions. DNA purity and concentration were

measured by NanoDrop, and DNA integrity was assessed using gel electrophoresis.

Samples were genotyped using Illumina Infinium Asian Screening Array-24 v1.0

(ASA) containing 659,184 variants in total. TOP strand-based genotypes were

reported in plink format using GenomeStudio software (Illumina) and then converted

to reference strand-based genotypes using PLINK (v1.9). Subsequently, we performed

initial quality control at variants and sample level following the pipelines used by the

Genotype-Tissue Expression (GTEx) consortium[27] and that introduced in ref,[51] with a

little modification. Briefly, at variants level, keeping variants with consistent genome

position and allele with that of 1000 Genomes Project, with genotype missing rate

across all samples < 0.02, MAF > 0.01 and the Hardy–Weinberg Equilibrium $P > 1 \times$

10^{-6}. At sample level, keeping samples with genotype missing rate across all variants

< 0.02, with matched sex tag between clinical record and genotype-imputed result,

with heterozygosity within 3 standard deviations from the mean heterozygosity.

Genotype imputation was performed using IMPUTE2 (v2.3.2)[52] with 1000 Genome

Project genotype data (phase 3) as reference panel. To increase the imputation

accuracy, we changed the default values of the parameter -k and –buffer to 100 and

300, respectively. After imputation, we performed another round of quality control,

keeping variants with quality score > 0.3, genotype missing rate across all samples <

0.05, MAF > 0.05 and the Hardy–Weinberg Equilibrium $P > 1 \times 10^{-6}$, keeping samples

with genotype missing rate across all variants < 0.05.

We performed principal component analysis (PCA) using smartpca[53] to detect the

stratification of population.

Cartilage RNA isolation, sequencing and data processing

Cartilage RNA was extracted with TRIzol. RNA integrity (RIN) was assessed using

the Fragment Analyzer 5400 (Agilent Technologies). Samples with RIN ≥ 5.6 were

used for sequencing library preparation using NEBNext® Ultra™ RNA Library Prep

Kit for Illumina® (NEB) according to manufacturer's instructions. RNA sequencing

was performed on an Illumina Novaseq 6000 platform and 150 bp paired-end reads

were generated.

Clean RNA-seq reads were mapped to human reference genome hg19 (only contained

autosomes, sex chromosomes and mitochondrial chromosomes) with STAR aligner

(v2.7.9a)[54] based on the GENCODE v19 (ref.[55]) annotation, the parameters setting

was same as the GTEx v8 pipeline. Gene-level read counts and TPM values were

generated using RNA-SeQC (v2.3.5)[56] with default parameters. The read mapping

rate and base mismatch rate were calculated from the output of SAMtools (v1.9)[57]

stats sub-command. The proportions of genomic origin of sequencing reads were

produced using QualiMap (v2.2.2-dev)[58]. Samples are needed to meet the following

metrics as used by the GTEx consortium[27]: read mapping rate $\geq$ 0.2, base mismatch

rate $\leq$ 0.01, intergenic mapping rate $\leq$ 0.3, and rRNA mapping rate $\leq$ 0.3. In this study,

all samples passed this mapping quality control step. Then, we conducted expression

outlier filtration with the method described in ref.[59]. Briefly, we calculated the

pairwise expression correlation coefficients using the log transformed read count of

all genes, let the r_{ij} denote the expression correlation coefficient between sample i and

sample j, we calculated

$$\bar{r}_i = \sum_j r_{ij}/n$$

544   the average correlation coefficient of sample i with all others of the total n samples.

545   Lower $\bar{r}_i$ represented a lower quality. Then we calculated

$$D_i = (\bar{r}_i - \bar{\bar{r}})/median(|\bar{r}_i - \bar{\bar{r}}|)$$

to provide a sense of distance from the grand correlation mean $\bar{\bar{r}}$. Samples with $D <$

-5 were considered as outliers and removed. Finally, samples that passed both

genotype and RNA-seq quality control were used for the following analysis, which

contained a total of 245 samples, of which, 204 were damaged cartilage samples and

31 were intact cartilage samples.

Differential expression analyses

We performed differential expression analyses using the differential expression for

repeated measures (dream) R package from variancePartition[25]. For the comparison

between damaged and intact cartilage, we produced the gene-level read counts matrix

for 31 donors from whom we obtained paired damaged and intact cartilage samples.

Genes with count per million (CPM) value < 0.1 in more than half samples were

removed, then gene-level read counts were normalized with trimmed means of m

values (TMM) method applied by edgeR (v3.40.2)[60] and voom-transformed with

voomWithDreamWeights function from variancePartition R package. Besides

cartilage type (damaged or intact), donor ID, sex, age, first two principal components

of genotype, RNA integrity number (RIN), and sequencing batch were also included

in the differential expression analysis model to correct the effects of technical and

non-cartilage type-related biological factors. FDR ≤ 0.05 was considered as the

threshold for the identification of differentially expressed genes.

Gene set enrichment analysis

Gene set enrichment analysis was performed for differentially expressed genes

between paired damaged and intact cartilage samples using GSEA software[61,62]. The

marker gene list of seven chondrocyte subtypes were calculated using FindAllMarkers

function from R package Seurat (v4.1.3)[63] with the scRNA-seq data and cell labels

from ref.[19].The enrichment *P*-values were calculated based on 100,000 times

permutation.

Deconvolution of bulk RNA-seq data into chondrocyte subtype composition

Processed single-cell RNA-seq data of knee cartilage[19] was downloaded from the

GEO database (GSE104782). UMI counts matrix where the columns were named by

cell type (HomC, HTC, RegC, PreHTC, FC, EC and ProC) and the rows were named

by gene symbols was used as input for CIBERSORTx[64] web server to produce a

chondrocyte subtype signature matrix. Then the signature matrix was applied to the

expression matrix of the bulk RNA-seq expression data to speculated estimate the

proportions of chondrocyte subtypes using the CIBERSORTx web server. In bulk

RNA-seq expression matrix, the columns were named by sample ID, rows were

named by gene symbols, and genes with the same symbols were removed. S-mode

was turned on for CIBERSORTx to correct the technical bias between single-cell and

bulk RNA-seq data.

Variation partition of gene expression

To get the contributor of gene expression variation at sample level, we applied

variancePartition (v1.28.4)[65], which is based on linear-mixed regression model, to

measure the impacts of biological and technical confounders on gene expression

variance, including donor ID, sex, age, cartilage type (damaged or intact), genetic

background (first two principal component of genotype), speculated proportions of six

chondrocyte subtypes (HomC, HTC, RegC, PreHTC, FC, EC, ProC was excluded

because of the extremely low fraction across all samples), RNA integrity number

(RIN), and sequencing batch. Genes with count per million (CPM) value < 0.1 in

more than half samples were removed, then gene-level read counts were normalized

with trimmed means of m values (TMM) method applied by edgeR (v3.40.2)[60] and

voom-transformed with voomWithDreamWeights function from variancePartition R

package. Participants identity (represented with donor ID) and sequencing batch were

modeled as random effects and the others confounders were modeled as fixed effects.

Cis-eQTL mapping

Expression data preprocessing

We followed the *cis*-eQTL mapping pipeline from the GTEx consortium[27]. Expression

data from damaged cartilage samples were used for eQTL analysis. Genes with TPM >

0.1 and read count ≥ 6 in at least 20% samples were selected. Further, genes on sex

and mitochondrial chromosomes were removed. Filtered expression data were

normalized using TMM to correct the library size difference between samples, then

the rank-based inverse normal transformation (RINT) was applied for each gene

across all samples, this method could increase the analysis power when test the

significance of correlation with large sample size than other normalization methods

did[66].

Covariates file generation

PEER factors[67] were calculated to estimate the hidden confounders of gene expression.

Following the advice from the GTEx consortium, 30 PEER factors were used for

covariates as our sample size is between 150 and 250. Integrated with sex, age, the

first two principal components of genotype and the proportions of seven chondrocyte

subtypes, we produced a covariate file used for *cis*-eQTL mapping.

Regular cis-eQTL mapping

All variants within 1 Mb of transcription start site (TSS) of each gene were used for

regular *cis*-eQTL mapping with linear regression model implemented by FastQTL

(v2.0)[68] and nominal *P*-value was calculated for each variant-gene pair. Beta

distribution-adjusted empirical *P*-values (from adoptive permutation results by setting

--permute 1000 10000 for FastQTL) of top variant of each gene were used for

multiple testing correction (FDR estimation) using the method described by Storey

and Tibshirani[69], which was implemented in R package qvalue, and FDR ≤ 0.05 was

used as a threshold to identify genes with significant eQTL (eGenes). To assess the

association significance of variants with eGenes, first, we got the mean of Beta

distribution-adjusted empirical *P*-values of two genes closest to the 0.05 FDR

threshold (one > 0.05 and the other one ≤ 0.05) as genome-wide empirical *P*-value

threshold (p_t), then the nominal *P*-value threshold for each gene was calculated using

the two beta distribution parameters of the minimum P-value distribution obtained

from the adoptive permutation procedure for the gene based on the genome-wide

empirical P-value ($F^{-1}(p_t)$, F^{-1} denote the inverse cumulative distribution of beta

distribution). For each gene, variants with a nominal P-value below the gene-level

nominal P-value threshold were considered statistically significant (eVariants).

Conditional analysis

To get the independent *cis*-eQTL signal for each eGene, a two-step conditional

analysis (a forward stepwise regression followed by a backward selection) was

performed using cis_independent mode of tensorQTL[70] with the regular *cis*-eQTL

analysis result as input, the lead variant for each independent eQTL signal was

reported. For visualization of each post-conditional independent eQTL signal of single

gene, we integrated the genotype dosage of the others lead variants of independent

signal into the covariates file and re-conducted the *cis*-eQTL analysis with FastQTL

for the gene to get a new P-value for each variant.

Epigenetic data processing

ATAC-seq data

The ATAC-seq data of knee cartilage samples from 16 donors were downloaded from

GEO database (GSE108301)[36]. The reads were mapped to human reference genome

hg19 with Bowtie2 (v2.3.5)[71] for each sample, reads with mapping quality < 30 or

duplicated were filtered. Peak calling was performed with MACS2 (v2.2.7.1)[72,73] as

described before[38]. Peaks presented only in one sample were removed and the others

were merged to produce a consistent peak set.

Histone modification ChIP-seq data

ChIP-seq data for four histone modifications (H3K4me1, H3K4me3, H3K27ac, and

H3K27me3) of adult knee cartilage samples were downloaded from from GEO

database (GSE111850)[28]. The reads were mapped to human reference genome hg19

with Bowtie2 (v2.3.5) for each sample, reads with mapping quality < 30 or duplicated

were removed. Then the bam files from three biological replications of each

modification type were merged to a single bam file. Peak calling was performed with

MACS2 (v2.2.7.1), for H3K4me1 and H3K27me3, the broad peaks were called with

--broad-cutoff set to 0.05, for H3K4me3 and H3K27ac, the narrow peak were called

with -q set to 0.05.

Peak files, which were generated by ENCODE project[29] using

pseudo-replications-based method, for six histone modification (H3K27ac,

H3K27me3, H3K36me3, H3K4me1, H3K4me3, and H3K9me3) of hMSCs derived

chondrocytes were download from GEO database under the access number

GSE188124, GSE187652, GSE187044, GSE187055, GSE187641, and GSE187471,

respectively. Then the genome coordinates were transformed from hg38 to hg19 using

liftOver with hg38ToHg19.over.chain.gz annotation file. The corresponding bam files

were downloaded from ENCODE portal. The alignment coordinates were transformed

to hg19 using CrossMap[74] with hg38ToHg19.over.chain.gz annotation file and then

the bam files for replication of each modification type were merged.

Super-enhancers identification

We performed the super-enhancers identification based on the H3K27ac ChIP-seq

data using ROSE2 with default settings. The H3K27ac peaks used for indicating the

enhancer position and the merged bam files used for enhancer signal strength

estimation.

eVariants enrichment analyses

All lead variants of conditional independent eQTL signals were LD-pruned with a

threshold $r^2 < 0.8$ to produce a query set. Then for each query variant, we extracted at

most 100 random variants from the SNPsnap dataset[75] as a background set, which

accounted for MAF deviation ($\pm$ 0.05), distance to gene deviation ($\pm$ 0.5), gene

density deviation ($\pm$ 0.5) and LD buddy deviation ($\pm$ 0.5). Then, the epigenetic

enrichment analysis was performed using fisher's exact test.

Cell type-interaction eQTL (ci-eQTL) analysis

As described in ref.[23], to measure the contribution of a single chondrocyte subtype to

eQTL effect, the significant variant-gene pairs in regular *cis*-eQTL result were used

for ci-eQTL analysis. We modeled following two linear-mixed regression models with

R package lme4 (v1.1-30)[76]:

$$E \sim Beta_g \times G + covariates \#(1)$$

$$E \sim Beta_i \times G{:}C + Beta_g \times G + Beta_c \times C + covariates \#(2)$$

where E denotes the RINT normalized gene expression value. G denotes the genotype

dosage. C denotes the proportion of specific chondrocyte subtype (ranging from 0 to

1). *Beta* denotes the effect size of each item. Covariates included in the models were

the same as that used in regular *cis*-eQTL analysis. Sex was modeled as random effect

and the other covariates were modeled as fixed effect.

The difference of those two models was compared with analysis of variance (ANOVA)

implemented by R package pbkrtest (v0.5.1)[77] and a χ^2 P-value was calculated to

indicate the statistical significance. We extracted the lead variant with the smallest χ^2

P-value for each tested gene and corrected the χ^2 P-value with Benjamini–Hochberg

FDR control method[78]. Genes for which the lead variants with FDR < 0.05 and

P-value $< 5 \times 10^{-5}$ of genotype dosage and cell proportion interaction item were

considered as statistically significant cell type-interaction eGenes. We divided the

significant variant-gene pairs into driver and non-driver model based on the effect

direction of $Beta_g$ from formula (1) and $Beta_i$ from formula (2). The same effect

direction meant that the eQTL effect increased as the cell fraction increased (driver

model), suggesting the eQTL effect may come from the cell type used for ci-eQTL

analysis (Fig. 4a). In contrast, the opposite effect direction meant that the eQTL effect

decreased as the cell fraction increased (non-driver model), suggesting the eQTL

effect didn't come from the cell type used for ci-eQTL analysis (Fig. 4a). The cell

proportions of EC, preHTC and RegC were used for ci-eQTL analyses because of

their broader distribution than that of the other four subtypes (Figs. 2d,e).

Sharing and specificity estimation of the ci-eQTL effect

To assess whether the cell type-interaction variant-gene pairs were shared or specific

across multiple chondrocytes subtypes (EC, preHTC and RegC). We applied the

multivariate adaptive shrinkage method (mash)[30], which was implemented in R

package mashr (v0.2.57), based on the workflow used for the GTEx data

(https://github.com/stephenslab/gtexresults). The effect size of interaction item of

genotype dosage and cell proportion ($Beta_i$) and the standard error (s.e.$_{Betai}$ calculated

form formula 2) were used as inputs for mashr. We extracted the variant with the

smallest P-value across all three cell types, and all tested variants for each gene and

produced two matrices which saved their effect sizes and standard errors for the gene

in three cell types, respectively. In addition, two matrices with the same format for a

200,000 randomly selected variant-gene pairs set also be produced. A mash model

was built using the exchange effects mode to estimate the prior. The model was then

applied to all significant variant-gene pairs for each cell type to get the posterior mash

effect sizes and the local false sign rate (LFSR). Pairs with LFSR < 0.5 were

considered statistically significant.

Co-localization analyses

We downloaded a total of 28 OA GWAS summary data from six published studies[4-9]

(Supplementary Table 13). For each GWAS summary data, we produced a list of

200-Kb signal window centered on each genome-wide significant variant ($P < 5 \times$

10^{-8}). The final signal region list for co-localization analysis was produced after

merging the overlapping signal windows. Then, the GWAS association P-value, MAF

and LD structure (calculated from UK biobank samples) of variants within each

region were integrated for further co-localization analysis.

To get *cis*-eQTL signal regions for co-localization analysis, for eQTL signal of each

gene from our regular *cis*-eQTL analysis, we merged the 200-Kb signal windows

centered on each significant eVariant of genome-wide significant eGenes or 200-Kb

signal windows centered on the top variant which with P-value < 0.05 for genes that

did not reach genome-wide significance. Then, the eQTL association P-value, MAF

(calculated from our samples) and LD structure (calculated from our samples) of

variants within each region were integrated for co-localization analysis. For *cis*-eQTL

signal from conditional analysis and ci-eQTL analysis, the information of variants

within 1-Mb windows of TSS were used for co-localization analysis.

Co-localization analysis was performed using R package coloc (v5.1.0.1)[79], which

implemented a Bayesian test to assess whether two association signals shared a

consistent causal variant by reporting the posterior probabilities of following five

hypotheses: H0, No association with either trait; H1, Association with trait 1, not with

trait 2; H2, Association with trait 2, not with trait 1; H3, Association with trait 1 and

trait 2, two independent SNPs; H4, Association with trait 1 and trait 2, one shared SNP.

The results were denoted as PPH0, PPH1, PPH2, PPH3, PPH4, higher PPH4

suggested higher probability that a single variant effecting both two traits. After

preparation of input files, we performed co-localization analysis using coloc.abf

function with default setting. Genes of which the eQTL signal showed PPH4 ≥ 0.7

with GWAS signal were called co-localized genes and considered as candidate OA

risk genes.

To verify that our eQTL data have the capacity to nominate more risk genes for OA

GWAS signals, we performed co-localization analysis using the eQTL signals of

co-localized genes from all tissues in the GTEx project v8. We also verified that our

cartilage eQTL data could nominate more OA risk genes than that using cartilage

eQTL data from ref. [10].

Processing of time-series gene expression data of chondrocytes differentiation

Soft-clustering of time-series expression data

To assess the functional roles of colocalized genes, we performed clustering and

enrichment analysis on a chondrogenic differentiation time series RNA-seq data

(GSE128554)[34]. We supposed the genes with similar expression change patterns have

similar functions. First, the gene-level read counts and TPM values of each sample

were obtained with the same method used for quantification of genes of our cartilage

samples. Then, pairwise differentially expressed gene identification was performed

using R package DESeq2 (v1.28.1)[80]. All genes with $P \leq 0.05$ in at least one

comparison were used for clustering analysis.

We performed soft-clustering of all differentially expressed genes based on the mean

TPM values at each time point using R package Mfuzz (v2.48.0)[81]. This method

extracted the patterns of pre-specified n clusters and then assigned a membership

score μ_{ij} ($0 < \mu_{ij} < 1$) to gene i to indicate the degree of membership of this gene for

cluster j. We chose to cluster the genes into 12 classes by visual observation. Finally,

we considered genes with membership score ≥ 0.6 as final members for each cluster.

GO enrichment analyses

GO enrichment was performed by R package clusterProfiler (v3.16.4)[82] with default

setting. GO terms with P-value ≤ 0.5 were considered statistically significant

TF binding-disruption analysis

Allele-sensitive TF footprint analysis

Footprint analysis was limited to the expressed TFs which were defined as the genes

with at least one DNA binding motif included in JASPAR 2022 core non-redundant

set[83] or HOCOMOCO v11 (ref.[84]) database and with a median TPM ≥ 1 across all

damaged cartilage samples.

TF footprint analysis involved a step to search TF motifs in the ATAC signal valleys

using sequence-match-based method. We supposed there was a possible scenario

where the reference genome-stored allele of a SNP may strongly disrupt a TF motif

but the alternative allele may help to form a sequence context suitable for TF binding.

To account for this, we produced an alternative allele--replaced genome sequence

based on hg19 using SNPsplit (v0.6.0)[85] by replacing the reference allele to the

alternative allele of all variants presented in genotype data of our samples. Both the

original reference genome sequence and newly generated allele-replaced genome

sequence were used for TF footprint analysis.

We performed TF footprint analysis using TOBIAS (v0.13.3)[86] with default settings,

TOBIAS was specially developed to account for the ATAC-seq data specific bias. All

16 bam files from ATAC-seq pipeline described above were merged to produce a

single input to increase the detection power. The search regions were limited to the

peaks presented in at least two samples. All motifs corresponding to the expressed

TFs were used for motif search. The results reported by TOBIAS were the genome

coordinates of the motifs and the speculated binding status.

TF binding-disruption evaluation

To investigate the potential impacts of eVariants on the TF binding, we applied a TF

motif information content-based method. Briefly, for each motif of expressed TF, we

calculated the information content (IC) of each position based on the position

frequency matrix (PFM). For each position of the motif, the total possible IC was

fixed

$$IC_{total} = log_2 n = log_2 4 = 2$$

832   where *n* denote the number of all possible base could be presented in this position,

833   thus *n* is 4 for any TF motif. Then, let the $P(N)$ denote the probability of a certain

834   base in the position, we calculated the sum of the uncertainty of all bases (A, C, G, T):

$$U = -\sum_{N=A}^{T} P(N) \times log_2 P(N)$$

835   Finally, the final information of the position was calculated:

$$IC_{final} = IC_{total} - U$$

The variants located in a position that with $IC_{final} \geq 0.5$ for a certain motif, and the

$|P(ref) - P(alt)| \geq 0.3$ were considered as candidate TF binding-disruption

variants.

TF expression-interaction (TFi-eQTL) eQTL analysis

To verify whether the TF for which the motif was disrupted by a variant was the true

mediator of the eQTL effect. we performed TFi-eQTL analysis that was similar to the

ci-eQTL analysis. Briefly, we modeled following two linear-mixed regression models

with lme4 (v1.1-30)[76]:

$$E_{target} \sim Beta_g \times G + covariate \#(3)$$

$$E_{target} \sim Beta_{tfi} \times G{:}E_{tf} + Beta_g \times G + Beta_{tf} \times E_{tf} + covariates \#(4)$$

where E_{target} denote the RINT normalized expression value of target gene. E_{tf}

denote the RINT normalized expression of TF. *Beta* denotes the effect size of each

item. Covariates included in the models were the same as that used in the regular

eQTL analysis plus the expression value of TF. Sex was modeled as random effect

and the other covariates were modeled as fixed effect. The difference of those two

models was compared with analysis of variance (ANOVA) implemented by R

package pbkrtest (v0.5.1)[77] and a χ^2 P-value was calculated to indicate the statistical

significance. For each TF-target pair for test, the motif and variant with smallest

P-value was retained.

855 Identification of allele-specific open chromatin (ASoC) SNPs

We adopted the pipeline described in ref.[35] to identify the variants that showed

imbalanced chromatin accessibility with a little modification(Supplementary Fig. 13).

Briefly, BAM files from 16 ATAC-seq data of cartilage samples[36] were calibrated by

WASP[87] and used as inputs for GATK (v4.2.6.1)[88]. Following the GATK Best Practice

recommendation, HaplotypeCaller sub-command was used for variants calling based

on human hg19 genome sequence and corresponding dbSNP version 138 annotation.

Then, the construction and application of InDels and SNPs recalibration model were

performed in tandem using VariantRecalibrator and ApplyVQSR sub-command,

respectively. In addition, the databases used for VariantRecalibrator as true training set

were hg19 versions of Mills and 1000G gold standard InDels list (priority = 12),

HapMap v3.3 and 1000G_omni v2.5 (priority = 12), the database used as non-true

training set was 1000G phase 1 high confidence SNP list (priority = 10) and dbSNP

138 (priority = 2), the database used as known set was dbSNP 138 (priority = 2).

Heterozygous variants with tranche level > 99.0% and located within ATAC-seq peaks

were retained. Due to the low read coverage in individual samples, we added up the

read depth corresponding to each allele across all heterozygous samples after

checking the consistency of allelic effect direction (Supplementary Fig. 14). Finally,

we did the binomial test for variant sites with minimum read depth count (DP) ≥ 20

using R function binom.test with a null hypothesis that the two alleles showed the

same effect on chromatin accessibility (namely the parameter p was set to 0.5). FDR

control was applied using R function p.adjust.

Cell culture

SW1353 cell line was obtained from Zeye (Shanghai, China). Cells were cultured in

DMEM medium supplemented with 10% fetal bovine serum (Biological Industries,

Israel), penicillin (100 U/ml) and streptomycin (0.1 mg/ml) in 5% CO_2 at 37℃

incubator.

Transfection and dual-luciferase reporter assays

A 876 bp putative open chromatin region containing rs11750646 (C alle or T alle),

and a 1500 bp fragment surrounding *PI3KR1* transcription start site were inserted into

pGL3 Basic vector (Promega, USA) (Supplementary Table 21). The luciferase

reporter plasmid and phRLTK containing Renilla luciferase were co-transfected into

cells using Lipo8000 Transfection Reagent (Beyotime, China). After 48 h, the relative

luciferase signal was computed as the ratio of firefly luciferase activity to Renilla

signal, and relative activity was normalized by pGL3-promoter plasmid. All data

came from four replicate wells, and statistical analyses were performed by a

two-tailed *t* test.

shRNA expression plasmid constructs and shRNA knockdown

For short hairpin RNA (shRNA) knockdown of *AR*, two independent oligonucleotides

targeting *AR* were cloned into linearized miR30 backbone. The shRNA and negative

control (NC) sequences are shown (Supplementary Table 22). A total of 2.5 μg of

each plasmid were independently transfected into 80% confluence of SW1353 cells

using Lipo8000 Transfection Reagent (Beyotime, China) according to the

manufacturer's instruction. After 48 hours of transfection, total RNA was isolated to

detect the mRNA expression by quantitative real-time PCR (qRT-PCR). Moreover, *AR*

shRNA plasmids were independently co-transfected with the expression plasmids

including rs11750646 (C alle or T alle) with *PI3KR1* promoter used in the luciferase

reporter assay. The transfection and measuring of luciferase activity are the same as

indicated in dual-luciferase reporter assay section. All data came from at least three

replicate wells, and statistical analyses were performed by a two-tailed *t* test.

Cell line RNA isolation, reverse transcription, and quantitative real-time PCR

Total RNA was extracted using RNA fast200 Kit (Fastagen, China). 1 μg of total RNA

was then reverse transcribed to cDNA using SuperScripts II First-Strand cDNA

Synthesis Kit (Invitrogen, Waltham, MA). qRT-PCR was conducted with the

QuantiTect SYBR® Green PCR Kit (Qiagen, Germany) on Bio-Rad instrument. The

$2^{-\Delta\Delta CT}$ method was used for data analysis. The qPCR primers are listed in

Supplementary Table 22.

Chromatin immunoprecipitation assay

Chromatin immunoprecipitation was performed using Simple-ChIP® Enzymatic

Chromatin IP Kit (Magnetic Beads) (Cell Signaling Technology, USA). Pre-cleared

chromatin was immunoprecipitated with monoclonal antibodies against AR

(ab108341, Abcam, USA), normal rabbit IgG or antibody against acetyl Histone H3

(included in the kit). The total cell chromatin extract was used as input. Protein-DNA

crosslinks were precipitated by using ChIP-Grade Protein G Magnetic Beads. And

DNA is purified using DNA purification spin columns (included in the kit). The target

DNA fragments were analyzed by Sanger sequencing of PCR fragments harboring

SNP (Supplementary Table 22).

Data visualization

ChIP-seq signals were visualized using R package Gviz (v1.32.0)[89]. Heatmap was

produced using R package ComplexHeatmap (v2.4.3)[90]. Other results were visualized

using R package ggplot2 (v3.3.6).

Data availability

Raw RNA-seq data reported in this paper have been deposited in the Genome

Sequence Archive (GSA)[91] in National Genomics Data Center[92], China National

Center for Bioinformation / Beijing Institute of Genomics, Chinese Academy of

Sciences   (GSA-Human: HRA006224)   that   are   accessible   at

https://ngdc.cncb.ac.cn/gsa-human with reasonable request. Individual-level genotype

data are available upon reasonable request: T.-L.Y. (yangtielin@xjtu.edu.cn). Access

to individual-level genotypes from samples recruited within mainland China is subject

to the policies and approvals from the Human Genetic Resource Administration,

Ministry of Science and Technology of the People's Republic of China. All

differentially expressed genes, lead variants of eQTL analyses (including regular

eQTL analysis, conditional analysis, ci-eQTL analysis, and TFi-eQTL analyses),

GWAS-eQTL co-localization, and allele-specific open chromatin results are presented

as Supplementary Tables.

Processed knee cartilage single-cell RNA data can download from supplementary file

at https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE104782. Knee cartilage

eQTL   summary   data   in   ref.[10]   is   available   at

https://personal.broadinstitute.org/ryank/Southam_FunGen_eQTL_LowGradeCartilag

e.FastQTL_perm_nom_info.ForMSK-KP_16Jan2021.zip   for   low-grade   (intact)

cartilage   and

https://personal.broadinstitute.org/ryank/Southam_FunGen_eQTL_HighGradeCartila

ge.FastQTL_perm_nom_info.ForMSK-KP_16Jan2021.zip for high-grade (damaged)

cartilage. The GTEx v8 eQTL summary data which contain associations of all tested

variant-gene pairs is available at

https://console.cloud.google.com/storage/browser/gtex-resources. Position Frequency

Matrices (PFMs) of JASPAR 2022 core non-redundant motif set of vertebrates is

available at

https://jaspar.genereg.net/download/data/2022/CORE/JASPAR2022_CORE_vertebrat

es_non-redundant_pfms_jaspar.txt. Position Frequency Matrices (PFMs) of

HOCOMOCO v11 motif set is available at

https://hocomoco11.autosome.org/final_bundle/hocomoco11/core/HUMAN/mono/H

OCOMOCOv11_core_HUMAN_mono_jaspar_format.txt. 1000 Genomes Project

Phase 3 haplotype reference is available at

https://mathgen.stats.ox.ac.uk/impute/1000GP_Phase3.tgz. Download links of GWAS

summary data of osteoarthritis are presented in Supplementary Table 13.

Code availability

Code used in this study is available at https://github.com/tianwen0003/OA-eQTL. The

GTEx consortium modified version of FastQTL and corresponding auxiliary scripts

were available at https://github.com/francois-a/fastqtl. Scripts used for gene-level

filtration and PEER factor calculation are available at

https://github.com/broadinstitute/gtex-pipeline/archive/refs/tags/gtex_v8.tar.gz.

Pipeline code for mashR is available at https://github.com/stephenslab/gtexresults.

Additional code may be available upon reasonable request from the corresponding

authors.

Acknowledgements

We thank the patients who donated material for these studies. This work was

supported by the National Natural Science Foundation of China (32170616, 32370653,

and 82170896); Science Fund for Distinguished Young Scholars of Shaanxi Province

(2021JC-02); Innovation Capability Support Program of Shaanxi Province

(2022TD-44); Key Research and Development Project of Shaanxi Province

(2022GXLH-01-22, 2023-YBSF-180), and the Fundamental Research Funds for the

Central Universities. This study was also supported by the High-Performance

Computing Platform and Instrument Analysis Center of Xi'an Jiaotong University.

Author contributions

Conceptualization: T.-L.Y. and Y.G.; methodology: W.T., F.J., C.W., R.-H.H. and

S.-S.D.; data analysis: W.T. and C.-Y.H.; experiment: C.W. and K.A.; visualization:

W.T. and C.W.; resource: S.-Y.H., H.-M.S., H.-W.G.and Z.Y.; public data collection:

W.T. and C.W.; writing-original draft: W.T. and C.W.; writing-review and editing:

T.-L.Y. and Y.G.; project administration and supervision: T.-L.Y. and Y.G.

Competing interests

The authors declare no competing interests.

1000

References

1001

1. Long, H. *et al.* Prevalence Trends of Site-Specific Osteoarthritis From 1990 to 2019: Findings
From the Global Burden of Disease Study 2019. *Arthritis & Rheumatology* **74**, 1172-1183
(2022).

2. Hunter, D.J. & Bierma-Zeinstra, S. Osteoarthritis. *The Lancet* **393**, 1745-1759 (2019).

3. Cibrián Uhalte, E., Wilkinson, J.M., Southam, L. & Zeggini, E. Pathways to understanding the
genomic aetiology of osteoarthritis. *Human Molecular Genetics* **26**, R193-R201 (2017).

4. Zengini, E. *et al.* Genome-wide analyses using UK Biobank data provide insights into the
genetic architecture of osteoarthritis. *Nature Genetics* **50**, 549-558 (2018).

5. Tachmazidou, I. *et al.* Identification of new therapeutic targets for osteoarthritis through
genome-wide analyses of UK Biobank data. *Nat Genet* **51**, 230-236 (2019).

6. Boer, C.G. *et al.* Deciphering osteoarthritis genetics across 826,690 individuals from 9
populations. *Cell* **184**, 4784-4818 e17 (2021).

7. Boer, C.G. *et al.* Genome-wide association of phenotypes based on clustering patterns of hand
osteoarthritis identify WNT9A as novel osteoarthritis gene. *Annals of the rheumatic diseases* **80**,
367-375 (2021).

8. Donertas, H.M., Fabian, D.K., Valenzuela, M.F., Partridge, L. & Thornton, J.M. Common
genetic associations between age-related diseases. *Nat Aging* **1**, 400-412 (2021).

9. Jiang, L., Zheng, Z., Fang, H. & Yang, J. A generalized linear mixed model association tool for
biobank-scale data. *Nat Genet* **53**, 1616-1621 (2021).

10. Steinberg, J. *et al.* A molecular quantitative trait locus map for osteoarthritis. *Nat Commun* **12**,
1309 (2021).

11. Gamazon, E.R. *et al.* Using an atlas of gene regulation across 44 human tissues to inform
complex disease- and trait-associated variation. *Nature Genetics* **50**, 956-967 (2018).

12. Bryois, J. *et al.* Cell-type-specific cis-eQTLs in eight human brain cell types identify novel risk
genes for psychiatric and neurological disorders. *Nature Neuroscience* **25**, 1104-1112 (2022).

13. Yazar, S. *et al.* Single-cell eQTL mapping identifies cell type-specific genetic control of
autoimmune disease. *Science* **376**, eabf3041 (2022).

14. van der Wijst, M.G.P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs
and co-expression QTLs. *Nat Genet* **50**, 493-497 (2018).

15. Schmiedel, B.J. *et al.* Impact of Genetic Polymorphisms on Human Immune Cell Gene
Expression. *Cell* **175**, 1701-1715 e16 (2018).

16. Ota, M. *et al.* Dynamic landscape of immune cell-specific gene regulation in immune-mediated
diseases. *Cell* **184**, 3006-3021 e17 (2021).

17. Dreier, R. Hypertrophic differentiation of chondrocytes in osteoarthritis: the developmental
aspect of degenerative joint disorders. *Arthritis Research & Therapy* **12**, 216 (2010).

18. Prein, C. *et al.* Structural and mechanical properties of the proliferative zone of the developing
murine growth plate cartilage assessed by atomic force microscopy. *Matrix Biology* **50**, 1-15
(2016).

19. Ji, Q. *et al.* Single-cell RNA-seq analysis reveals the progression of human osteoarthritis. *Ann
Rheum Dis* **78**, 100-110 (2019).

20. Westra, H.J. *et al.* Cell Specific eQTL Analysis without Sorting Cells. *PLoS Genet* **11**, e1005223
(2015).

21.   Kim-Hellmuth, S. *et al.* Cell type-specific genetic regulation of gene expression across human
1045   tissues. *Science* **369**(2020).
1046   22.   Elorbany, R. *et al.* Single-cell sequencing reveals lineage-specific dynamic genetic regulation of
1047   gene expression during human cardiomyocyte differentiation. *PLoS Genet* **18**, e1009666 (2022).
1048   23.   Sheng, X. *et al.* Mapping the genetic architecture of human traits to cell types in the kidney
1049   identifies mechanisms of disease and potential treatments. *Nature Genetics* **53**, 1322-1333
1050   (2021).
1051   24.   Patel, D. *et al.* Cell-type-specific expression quantitative trait loci associated with Alzheimer
1052   disease in blood and brain tissue. *Transl Psychiatry* **11**, 250 (2021).
1053   25.   Hoffman, G.E. & Roussos, P. Dream: powerful differential expression analysis for repeated
1054   measures designs. *Bioinformatics* **37**, 192-201 (2021).
1055   26.   Sturm, G., List, M. & Zhang, J.D. Tissue heterogeneity is prevalent in gene expression studies.
1056   *NAR Genomics and Bioinformatics* **3**(2021).
1057   27.   Consortium, G.T. The GTEx Consortium atlas of genetic regulatory effects across human tissues.
1058   *Science* **369**, 1318-1330 (2020).
1059   28.   Ferguson, G.B. *et al.* Mapping molecular landmarks of human skeletal ontogeny and pluripotent
1060   stem cell-derived articular chondrocytes. *Nature Communications* **9**, 3634 (2018).
1061   29.   Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome. *Nature*
1062   **489**, 57 (2012).
1063   30.   Urbut, S.M., Wang, G., Carbonetto, P. & Stephens, M. Flexible statistical methods for
1064   estimating and testing effects in genomic studies with multiple conditions. *Nat Genet* **51**,
1065   187-195 (2019).
1066   31.   Bann, J.G., Peyton, D.H. & Bächinger, H.P. Sweet is stable: glycosylation stabilizes collagen.
1067   *FEBS Letters* **473**, 237-240 (2000).
1068   32.   Middleton, J., Manthey, A. & Tyler, J. Insulin-like growth factor (IGF) receptor, IGF-I,
1069   interleukin-1 beta (IL-1 beta), and IL-6 mRNA expression in osteoarthritic and normal human
1070   cartilage. *Journal of Histochemistry & Cytochemistry* **44**, 133-141 (1996).
1071   33.   Tanaka, N. *et al.* The attenuation of insulin-like growth factor signaling may be responsible for
1072   relative reduction in matrix synthesis in degenerated areas of osteoarthritic cartilage. *BMC
1073   Musculoskeletal Disorders* **22**, 231 (2021).
1074   34.   Vail, D.J., Somoza, R.A., Caplan, A.I. & Khalil, A.M. Transcriptome dynamics of long
1075   noncoding RNAs and transcription factors demarcate human neonatal, adult, and human
1076   mesenchymal stem cell-derived engineered cartilage. *Journal of Tissue Engineering and
1077   Regenerative Medicine* **14**, 29-44 (2020).
1078   35.   Zhang, S. *et al.* Allele-specific open chromatin in human iPSC neurons elucidates functional
1079   disease variants. *Science* **369**, 561-565 (2020).
1080   36.   Liu, Y. *et al.* Chromatin accessibility landscape of articular knee cartilage reveals aberrant
1081   enhancer regulation in osteoarthritis. *Scientific Reports* **8**, 15499 (2018).
1082   37.   Wang, N.N. *et al.* Enhancer variants on chromosome 2p14 regulating SPRED2 and ACTR2 act
1083   as a signal amplifier to protect against rheumatoid arthritis. *Am J Hum Genet* **110**, 625-637
1084   (2023).
1085   38.   Wang, C. *et al.* Lineage-selective super enhancers mediate core regulatory circuitry during
1086   adipogenic and osteogenic differentiation of human mesenchymal stem cells. *Cell Death &
1087   Disease* **13**, 866 (2022).

39.   Gao, P. *et al.* Biology and Clinical Implications of the 19q13 Aggressive Prostate Cancer
Susceptibility Locus. *Cell* **174**, 576-589.e18 (2018).
40.   Chen, X.F. *et al.* An Osteoporosis Risk SNP at 1p36.12 Acts as an Allele-Specific Enhancer to
Modulate LINC00339 Expression via Long-Range Loop Formation. *Am J Hum Genet* **102**,
776-793 (2018).
41.   Dong, S.S. *et al.* An Intronic Risk SNP rs12454712 for Central Obesity Acts As an
Allele-Specific Enhancer To Regulate BCL2 Expression. *Diabetes* **70**, 1679-1688 (2021).
42.   Thynn, H.N. *et al.* An Allele-Specific Functional SNP Associated with Two Systemic
Autoimmune Diseases Modulates IRF5 Expression by Long-Range Chromatin Loop Formation.
J Invest Dermatol **140**, 348-360 e11 (2020).
43.   Fan, Z. *et al.* MiR-155 promotes interleukin-1β-induced chondrocyte apoptosis and catabolic
activity by targeting PIK3R1-mediated PI3K/Akt pathway. *Journal of Cellular and Molecular*
Medicine **24**, 8441-8451 (2020).
44.   Donovan, M.K.R., D'Antonio-Chronowska, A., D'Antonio, M. & Frazer, K.A. Cellular
deconvolution of GTEx tissues powers discovery of disease and cell-type associated regulatory
variants. *Nat Commun* **11**, 955 (2020).
45.   Aguirre-Gamboa, R. *et al.* Deconvolution of bulk blood eQTL effects into immune cell
subpopulations. *BMC Bioinformatics* **21**, 243 (2020).
46.   den Hollander, W. *et al.* Annotating Transcriptional Effects of Genetic Variants in
Disease-Relevant Tissue: Transcriptome-Wide Allelic Imbalance in Osteoarthritic Cartilage.
Arthritis & Rheumatology **71**, 561-570 (2019).
47.   Kosoy, R. *et al.* Genetics of the human microglia regulome refines Alzheimer's disease risk loci.
Nat Genet **54**, 1145-1154 (2022).
48.   Liang, D. *et al.* Cell-type-specific effects of genetic variation on chromatin accessibility during
human neuronal differentiation. *Nature Neuroscience* **24**, 941-953 (2021).
49.   Alasoo, K. *et al.* Shared genetic effects on chromatin and gene expression indicate a role for
enhancer priming in immune response. *Nat Genet* **50**, 424-431 (2018).
50.   Currin, K.W. *et al.* Genetic effects on liver chromatin accessibility identify disease regulatory
variants. *The American Journal of Human Genetics* **108**, 1169-1189 (2021).
51.   Wang, T. *et al.* A pipeline for RNA-seq based eQTL analysis with automated quality control
procedures. *BMC Bioinformatics* **22**, 403 (2021).
52.   Howie, B.N., Donnelly, P. & Marchini, J. A flexible and accurate genotype imputation method
for the next generation of genome-wide association studies. *PLoS Genet* **5**, e1000529 (2009).
53.   Price, A.L. *et al.* Principal components analysis corrects for stratification in genome-wide
association studies. *Nature Genetics* **38**, 904-909 (2006).
54.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
55.   Harrow, J. *et al.* GENCODE: the reference human genome annotation for The ENCODE Project.
Genome research **22**, 1760-1774 (2012).
56.   Graubert, A., Aguet, F., Ravi, A., Ardlie, K.G. & Getz, G. RNA-SeQC 2: efficient RNA-seq
quality control and quantification for large cohorts. *Bioinformatics* **37**, 3048-3050 (2021).
57.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079
(2009).
58.   Okonechnikov, K., Conesa, A. & García-Alcalde, F. Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data. *Bioinformatics* **32**, 292-294 (2015).

59. Wright, F.A. *et al.* Heritability and genomics of gene expression in peripheral blood. *Nature*
Genetics **46**, 430-437 (2014).
60. McCarthy, D.J., Chen, Y. & Smyth, G.K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research* **40**,
4288-4297 (2012).
61. Subramanian, A. *et al.* GSEA: Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-50 (2005).
62. Mootha, V.K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are
coordinately downregulated in human diabetes. *Nature Genetics* **34**, 267-273 (2003).
63. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).
64. Newman, A.M. *et al.* Determining cell type abundance and expression from bulk tissues with
digital cytometry. *Nature Biotechnology* **37**, 773-782 (2019).
65. Hoffman, G.E. & Schadt, E.E. variancePartition: interpreting drivers of variation in complex
gene expression studies. *BMC Bioinformatics* **17**, 483 (2016).
66. Bishara, A.J. & Hittner, J.B. Testing the significance of a correlation with nonnormal data:
comparison of Pearson, Spearman, transformation, and resampling approaches. *Psychol*
Methods **17**, 399-417 (2012).
67. Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of
expression residuals (PEER) to obtain increased power and interpretability of gene expression
analyses. *Nat Protoc* **7**, 500-7 (2012).
68. Ongen, H., Buil, A., Brown, A.A., Dermitzakis, E.T. & Delaneau, O. Fast and efficient QTL
mapper for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479-85 (2016).
69. Storey, J.D. & Tibshirani, R. Statistical significance for genomewide studies. *Proceedings of the*
National Academy of Sciences **100**, 9440-9445 (2003).
70. Taylor-Weiner, A. *et al.* Scaling computational genomics to millions of individuals with GPUs.
Genome Biology **20**, 228 (2019).
71. Langmead, B. & Salzberg, S.L. Bowtie2: Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357-9 (2012).
72. Zhang, Y. *et al.* MACS: Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137
(2008).
73. Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X.S. Identifying ChIP-seq enrichment using MACS.
Nature Protocols **7**, 1728-1740 (2012).
74. Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between genome assemblies.
Bioinformatics **30**, 1006-1007 (2013).
75. Pers, T.H., Timshel, P. & Hirschhorn, J.N. SNPsnap: a Web-based tool for identification and
annotation of matched SNPs. *Bioinformatics* **31**, 418-20 (2015).
76. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67**, 1 - 48 (2015).
77. Halekoh, U. & Højsgaard, S. A Kenward-Roger Approximation and Parametric Bootstrap
Methods for Tests in Linear Mixed Models – The R Package pbkrtest. *Journal of Statistical*
Software **59**, 1 - 32 (2014).
78. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*

57, 289-300 (1995).

79.   Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association

studies using summary statistics. *PLoS Genet* **10**, e1004383 (2014).

80.   Love, M.I., Huber, W. & Anders, S. DESeq2: Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

81.   Kumar, L. & Futschik, M.E. Mfuzz: a software package for soft clustering of microarray data.

Bioinformation **2**, 5 (2007).

82.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology* **16**,

284-287 (2012).

83.   Castro-Mondragon, J.A. *et al.* JASPAR 2022: the 9th release of the open-access database of

transcription factor binding profiles. *Nucleic Acids Res* **50**, D165-D173 (2022).

84.   Kulakovskiy, I.V. *et al.* HOCOMOCO: towards a complete collection of transcription factor

binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Research*

46, D252-D259 (2017).

85.   Krueger, F. & Andrews, S. SNPsplit: Allele-specific splitting of alignments between genomes

with known SNP genotypes [version 2; peer review: 3 approved]. *F1000Research* **5**(2016).

86.   Bentsen, M. *et al.* ATAC-seq footprinting unravels kinetics of transcription factor binding during

zygotic genome activation. *Nat Commun* **11**, 4267 (2020).

87.   van de Geijn, B., McVicker, G., Gilad, Y. & Pritchard, J.K. WASP: allele-specific software for

robust molecular quantitative trait locus discovery. *Nat Methods* **12**, 1061-3 (2015).

88.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Research* **20**, 1297-1303 (2010).

89.   Hahne, F. & Ivanek, R. Visualizing Genomic Data Using Gviz and Bioconductor. in *Statistical*

Genomics: Methods and Protocols (eds. Mathé, E. & Davis, S.) 335-351 (Springer New York,

New York, NY, 2016).

90.   Gu, Z. Complex heatmap visualization. *iMeta* **1**, e43 (2022).

91.   Chen, T. *et al.* The Genome Sequence Archive Family: Toward Explosive Data Growth and

Diverse Data Types. *Genomics, Proteomics & Bioinformatics* **19**, 578-583 (2021).

92.   Members, C.-N. & Partners. Database Resources of the National Genomics Data Center, China

National Center for Bioinformation in 2023. *Nucleic Acids Research* **51**, D18-D28 (2023).

1207

1208

Figure Legends

Figure 1. Overview of the study

We collected damaged and intact knee cartilage samples for bulk RNA-seq and blood

samples for DNA genotyping. We then deconvoluted bulk gene expression data into

chondrocyte subtype proportions based on single-cell RNA-seq data of knee cartilage

samples. We performed *cis*-eQTL mapping and followed by conditional analysis to

identify independent eQTL signals for each eGene. We also performed cell

type-interaction eQTL analyses to reveal the genetically regulatory heterogeneity of

gene expression between chondrocyte subtypes. By integrating our eQTL data and

OA GWAS summary data, we nominated risk genes for OA and annotated their

function using multiple lines of evidences. Finally, using our established multi-level

functional variants-deciphering pipeline, we performed functional fine-mapping based

on cartilage epigenetic features that called from cartilage ATAC-seq data to nominate

the candidate causal eVariants which may regulate gene expression through alteration

of chromatin accessibility and TF binding.

1224

Figure 2. Expression analyses results

a, The differentially expressed genes (DEGs) (FDR $\leq$ 0.05, no fold change cutoff)

between paired damaged and intact cartilage samples when not corrected the impact

of cell composition heterogeneity. The x axis is log-converted fold change (damaged

vs intact) of expression; The y axis is statistical significance in $-\log_{10}$(adjusted

P-value). **b,c**, Gene set enrichment analyses (GSEA) result showed the genes

expressed higher in intact cartilage samples ($n = 715$ as showed in Fig. f) enriched in

EC (b) and RegC markers (c). The FDR q-value was calculated based on the *P*-values

obtained from 100,000 permutations. **d**, Chondrocytes subtype proportions speculated

by CIBERSORTx per sample. Each column is a different sample. The y axis is the

accumulated cell proportions. **e**, Distribution of proportions of each chondrocyte

subtype across all damaged and intact cartilage samples. The y axis is the sample

density; The x axis is the cell proportion. **f**, Comparison of speculated chondrocytes

subtype proportions between paired damaged and intact cartilage sample. The

P-values were calculated using paired Student's t-test. The x axis is different cell type;

The y axis is the cell proportion. **g**, Distribution of gene expression variance explained

per gene by different confounders. The x axis is different factors; The y axis is

percentage (%) of total variance explained. Biological factors are colored in light blue;

artificial factors are colored in orange; chondrocyte subtype proportions are colored in

brown. **h**, Distribution of gene expression variance explained per gene by all

chondrocyte subtype proportions. The y axis is percentage (%) of total variance

explained.

1247

Figure 3. *cis*-eQTL results

a, Manhattan plot shows the association strength of lead variants of all eQTL signals

of all tested gene ($n = 18,756$). The y axis is the statistical significance in

$-\log_{10}$(*P*-value). **b**, Biotype distribution of eGenes. **c**, Comparison of expression and

variable coefficients distribution of significant eGenes and non-eGenes. The x axis is

log-converted mean expression (top panel) and variable coefficients (bottom panel).

P-value was calculated using Wilcoxon rank sum test. **d** Number of genes with certain

number of independent signals in conditional analysis. **e**, Distribution of genes with or

without newly identified independent signal (not significant in regular *cis*-eQTL

analysis) after conditional analysis. **f**, *SNAP91* has two independent eQTL signals

tagged by rs145051363 (middle) and rs623263 (bottom), respectively. In regular

cis-eQTL analysis, only the rs145051363 tagged signal is significant, the rs623263

tagged signal is not significant (top). **g**, Genome distribution of lead eVariants of all

independent eQTL signals. **h**, Distance of lead variants of all independent eQTL

signals from the transcription start site (TSS) of the associated eGenes and

corresponding nominal statistical significance in -$\log_{10}$(*P*-value). **i**, Enrichment of

LD-pruned ($r^2 < 0.8$) lead eVariants in different epigenetic features. Data are showed

as odds ratio (OR) ± 95% confidence intervals. The utilized epigenetic feature data

are from both original adult human knee cartilage samples and hMSCs derived

chondrocytes. ATAC-seq data is absent from hMSCs derived chondrocytes.

H3K36me3 and H3K9me3 ChIP-seq data are absent from adult human knee cartilage.

j, Enrichment of LD-pruned ($r^2 < 0.8$) lead eVariants in H3K27ac-defined SEs and

TEs from both adult human knee cartilage and hMSCs derived chondrocytes.

1271

Figure 4. Genetically regulatory heterogeneity between chondrocyte subtypes

a, Schematic outline of ci-eQTL analysis. For significant variant-gene pairs in regular

cis-eQTL analysis, an interaction term of genotype dosage and cell type proportion is

modeled in the linear-mixed model to test the cell type dependency of certain eQTL

effect. Driver model means the eQTL effect emerging with increase of cell proportion.

Non-driver model means that eQTL effect disappearing with increase of cell

proportion. **b**, Number of significant genes discovered in ci-eQTL analyses that

aligned to driver and non-driver model for three chondrocyte subtypes. **c**, Heatmap

shows the detailed mash results of all significant ci-eQTL results. The color depth

represents the mash-reported posterior effect size estimates of interaction item of

genotype dosage and cell proportion. Red represents the diver model; Blue represents

non-Driver model. Each row is a variant-gene pair. The rows and columns were

clustered based on the similarity. **d**, Association between the genotypes of variant

rs12620516 and *IGFBP2* gene expression. The y axis is the normalized expression of

IGFBP2 in damaged cartilage samples; the x axis is the genotype at the rs12620516.

The whiskers extend to the 5th and 95th percentiles; the outliers are represented by

the dots. The *P*-value was reported by FastQTL nominal procedure. **e**, ci-eQTL results

of rs12620516~*IGFBP2* pair across three different chondrocyte subtypes. The y axis

is the normalized expression of *IGFBP2* in damaged cartilage samples; the x axis is

the cell proportion. Each point represents one sample and colored by genotype types.

The *P*-values indicate the statistical significance of interaction term in linear-mixed

regression model. LFSR was estimated by mash. **f**, Association between the genotypes

of variant rs2356553 and *MAST2* gene expression. **g**, ci-eQTL results of

rs2356553~*MAST2* pair across three different chondrocyte subtypes. The y axis is the

normalized expression of *MAST2* in damaged cartilage samples; the x axis is the cell

proportion. Each data point represents one sample and colored by genotype types. The

P-values indicate the statistical significance of interaction term in linear-mixed

regression model. LFSR was estimated by mash. **h**, Association between the

genotypes of variant rs1408829 and *TOR3A* gene expression. **i**, ci-eQTL results of

rs1408829~*TOR3A* pair across three different chondrocyte subtypes. The y axis is the

normalized expression of *TOR3A* in damaged cartilage samples; the x axis is the cell

proportion. Each data point represents one sample and colored by genotype types. The

P-values indicate the statistical significance of interaction term in linear-mixed

regression model. LFSR was estimated by mash.

1306

Figure 5. Co-localization analysis nominate risk genes for osteoarthritis

a, Co-localization result of OA GWAS loci with cartilage eQTLs. The numbers next

to the triangles represent the highest value of probability of a single shared variant for

both GWAS and eQTL signal (PPH4) for each type of OA site (see more details in

Supplementary Table 13). Genes nominated with PPH4 ≥ 0.7 in at least one GWAS

data were showed. **b**, Function annotation of co-localized genes from multiple lines of

evidences.

1314

Figure 6. Allele-specific open chromatin (ASoC) variant links both to OA risk

and cartilage gene expression

a, The schematic diagram of an allele specific open chromatin variant. **b**, Number of

eVariant that is ASoC variant and eGenes associated with ASoC eVaraints under

different statistical significance threshold. **c**, Association between the genotypes of

variant rs3810154 and *SLC44A2* gene expression. The y axis is the normalized

expression of *SLC44A2* in damaged cartilage samples; the x axis is the genotype at the

rs3810154. The whiskers extend to the 5th and 95th percentiles; Each dot is a

different sample. The *P*-value was reported by FastQTL nominal procedure. **d**,

Co-localization of OA GWAS locus and eQTL signal associated to *SLC44A2*. The y

axis is the statistical significance of OA GWAS and cartilage eQTL for *SLC44A2* gene

in -$\log_{10}$(*P*-value). The x axis is the chromosome coordinate. **e**, Variant rs3810154

located within an open chromatin region marked by an ATAC-seq peak downstream

the *SLC44A2* and altered the local chromatin accessibility. The sequencing reads with

G allele piled up a significantly higher peak than that piled up by A allele-contained

reads. **f**, *SLC44A2* expression at each time point during the chondrocyte

differentiation. The y axis is the expression value in TPM; The x axis is the time point.

Each time point has three biological replications; The data are showed as mean ± sd.

1333

Figure 7. TF binding-disruption (TBD) variants links both to OA risk and

cartilage gene expression

a, Schematic diagram for TF binding-disruption analysis module in our multiple-level

functional variants-deciphering pipeline. Three main steps included. For the

allele-sensitive TF footprint analysis step, we introduced both the original reference

genome sequence and the allele-replaced genome sequence corresponding to the

signal valley regions as inputs for TF motif search. In this example scenario, only the

alternative allele-replaced genome sequence matched to the TF motif. **b**, The number

of eVariant that is TBD variant, eGenes associated with TBD eVaraints and TFs

whose binding sites were affected under different statistical significance threshold. **c**,

Co-localization of OA GWAS locus and eQTL signal associated to *PIK3R1*. The y

axis is the statistical significance of OA GWAS and cartilage eQTL for *PIK3R1* gene

in -$\log_{10}$(*P*-value). The x axis is the chromosome coordinate. **d**, Association between

the genotypes of variant rs11750646 and *PIK3R1* gene expression. The y axis is the

normalized expression of *PIK3R1* in damaged cartilage samples; the x axis is the

genotype at the rs11750646. The whiskers extend to the 5th and 95th percentiles;

Each dot is a different sample. The *P*-value was reported by FastQTL nominal

procedure. **e**, rs11750646 overlapped with footprints of AR, FOXA2, FOXA3,

FOXM1 and FOXO1, that were identified in an open chromatin region marked by an

ATAC peak downstream the *PIK3R1*. **f**, Reference allele C at rs11750646 disrupts the

motif (ID: ANDR_HUMAN.H11MO.0.A) of AR. The motif coordinate was shown on

the top. The symbol (+) represents the genome sequence shown is positive strand. **g**,

TFi-eQTL analysis reveal rs11750646 modified the regulatory role of *AR* on *PIK3R1*

expression. The y axis is the normalized expression of *PIK3R1* in damaged cartilage

samples; The x axis is the normalized expression of *AR* in damaged cartilage samples.

The *P*-value indicates the statistical significance of interaction term of TF expression

and target expression in linear-mixed regression model. **h**, Schematic diagram of dual

luciferase reporter vector. **i**, Detection of activity effect on open chromatin region

containing rs11750646 (T and C allele) compared with pGL3-*PIK3R1*-promoter in

SW1353 cells by dual-luciferase reporter assay. Luciferase signal was computed as

the ratio of firefly luciferase activity to Renilla signal and relative activity was

normalized by pGL3-*PIK3R1*-promoter. *P*-values are calculated using two-tailed

Student's *t*-test. **j**, AR preferred the binding to the T allele at rs11750646 confirmed by

ChIP followed by Sanger sequencing in SW1353cells. **k**, Effect of *AR* knockdown on

PIK3R1 expression in SW1353 cells. Two independent shRNAs (shRNA1 and

shRNA2) and shRNA-NC (negative control) were used. RT-qPCR analysis of gene

expression normalized to *GAPDH* expression level. *P*-values are calculated using

two-tailed Student's *t*-test. **l**, Dual-luciferase reporter assay containing rs11750646-C

or -T allele plasmids co-transfected with *AR* shRNAs in SW1353 cells. The

shRNA-NC is used as the negative control. Luciferase signals are normalized to

Renilla signals. *P*-values are calculated using two-tailed Student's *t*-test. **m**, The OA

risk allele C at rs11750646 disrupts binding of AR to a regulatory element, resulting

in decreased *PIK3R1* expression and further increasing the risk of osteoarthritis.

Fig. 1: Overview of the study

Fig. 2: Expression analyses results

Fig. 3: *cis*-eQTL results

Fig. 4: Genetically regulatory heterogeneity between chondrocyte subtypes

Fig. 5: Co-localization analysis nominate candidate risk genes for osteoarthritis

Fig. 6: Allele-specific open chromatin (ASoC) variant links both to OA risk and cartilage gene expression

Fig. 7: TF binding-disruption (TBD) variants links both to OA risk and cartilage gene expression