Genomic and morphological characterization of *Trichuris incognita n. sp.*, a Human-
Infecting Trichuris species

Max Alexander Bär [1,2], **Nadège Akissi Kouamé** [3,4], **Sadikou Touré** [4], **Jean Tenena**
Coulibaly [1,2,3,4], **Pierre Henri Hermann Schneeberger** [† 1,2], **Jennifer Keiser** [† 1,2]

[†] These authors contributed equally as senior authors

[1] Department of Medical Parasitology and Infection Biology, Swiss Tropical and Public Health
Institute, Allschwil, Switzerland

[2] University of Basel, Basel, Switzerland

[3] Unité de Formation et de Recherche Biosciences, Université Felix Houphouët-Boigny,
Abidjan, Cote d'Ivoire

[4] Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Cote d'Ivoire

Abstract

Background: Trichuriasis is a neglected tropical disease that affects as many as 500 million
individuals and can cause significant morbidity. For decades, it was thought to be caused by
one species of whipworm, *Trichuris trichiura*. Significant differences in response rates to the
best available anthelmintic treatment for trichuriasis — a combination of albendazole and
ivermectin — exist across clinical trial reports yet the underlying reasons are unclear.

Methods: Combining long- and short- read sequencing approaches, we assembled a high-
quality reference genome of *T. incognita n. sp.* isolated in an interventional study conducted
in Côte d'Ivoire. The species tree of the *Trichuris* genus was constructed using 12434
orthologous groups. We sequenced a total of 747 individual worms which were used to confirm
the phylogenetic placement and investigate patterns of adaptation through comparative
genomic analysis.

Findings: Here, we present and characterize a human-infecting *Trichuris* species that we
named *Trichuris incognita n. sp*. Comparative genomic analysis of genes suspected to confer
resistance to either albendazole or ivermectin in helminths revealed a high number of beta-
tubulin (TBB) orthologs, present in the whole population of *T. incognita n. sp.*, compared to
the canonical *T. trichiura* species, but these genes were not associated to a resistant
phenotype. We conducted a genome wide association study (GWAS) comparing 179
albendazole-ivermectin sensitive to 542 drug non-sensitive worms, which did not conclusively
show an adaptation to drug pressure within the same species.

Interpretation: Our results demonstrate that trichuriasis can be caused by multiple species of
whipworm and that the differences in response rates, may be a result of species responding
differently to drug treatment, as opposed to an establishment of resistance. This discovery,
coupled with the high tolerability of *T. incognita n. sp.* to albendazole-ivermectin marks a
significant shift in how we understand and approach whipworm infections.

Funding: European Research Council (ERC) Nr. 101019223.

Research in context Panel:

Evidence before this study

The recommended treatment with the benzimidazoles albendazole or mebendazole shows consistently low efficacy against *T. trichiura*, and therefore combination chemotherapy of benzimidazoles and ivermectin has been recommended. Recent clinical trials with albendazole-ivermectin, showed especially low cure rates against what was presumed to be *T. trichiura* in Côte d'Ivoire, in contrast to higher efficacy observed in Laos and Pemba, Tanzania. Amplicon sequencing data pointed towards underlying genomic differences and potentially a new species of human infecting *Trichuris* in Côte d'Ivoire.

Added value of this study

We identified and described a morphologically indistinguishable, novel causative agent of trichuriasis that doesn't respond to conventional drug treatment: *Trichuris incognita n. sp*. We provide a high-quality reference genome of this novel species and sequencing data of 747 individual isolates. A genome wide association study did not conclusively show adaptation to drug pressure. This study demonstrates that trichuriasis can be caused by multiple species of whipworm and that the differences in response rates, may be a result of species responding differently to drug treatment, as opposed to an establishment of resistance.

Implications of all the available evidence

The current whole body of evidence calls for new effective treatments against *T. incognita n. sp*. Due to the morphological indistinguishableness, molecular diagnostic tools need to be implemented to identify species level differences as opposed to the traditional Kato Katz method. The geographical distribution of *T. incognita* as well as longitudinal changes of species distributions will need to be mapped, in addition to better understanding of the new species pathology and transmission dynamics.

Introduction

Soil-transmitted helminthiasis is the most prevalent neglected tropical disease (NTD) worldwide. Over one billion people, particularly in impoverished nations, may develop debilitating chronic infections due to one of the five most common soil-transmitted helminths (STHs): *Ascaris lumbricoides*, *Trichuris trichiura*, the two hookworm species, *Necator americanus* and *Ancylostoma duodenale*, and *Strongyloides stercoralis*.(1) Despite their genetic diversity, the first four parasites are classified under the umbrella of STHs by the World Health Organization (WHO) and are treated with benzimidazoles, the cornerstone of large-scale deworming efforts through preventative chemotherapy (PC) since 1984 or ivermectin for strongyloidiasis.(2-5)

With more than 3.3 billion tablets of albendazole donated and distributed up until August 2020 and 5 billion treatments of ivermectin for onchocerciasis until January 2025, concern regarding anthelmintic resistance extending to humans, is rising. Reports of decreased efficacy and documented resistance to the same drugs within other helminths in livestock support projections suggesting the emergence of resistance within the next decade.(6, 7) However, no molecular evidence of resistance has yet been presented in human STHs. Notably, albendazole and ivermectin monotherapies show consistently low efficacy against *T. trichiura*, possibly due to differences in cuticle structure, mucosal embedding, or drug efflux, though these remain unproven.(8)

To circumvent potential resistance and improve efficacy against *T. trichiura*, the use of a combination therapy of albendazole and ivermectin has been taken forward. Recent clinical trials with albendazole-ivermectin, revealed especially low cure rates against what was presumed to be *T. trichiura* in Côte d'Ivoire, in contrast to higher efficacy observed in Laos and Pemba, Tanzania.(9) Amplicon sequencing data, normally not recorded in clinical trials,

pointed towards underlying genomic differences and potentially a new species of human
infecting *Trichuris* in Côte d'Ivoire, which may have been observed previously by S. Nissen *et*
al. already in 2012.(10, 11)

Despite the clinical relevance of these findings in addition to harboring information on potential
immunotherapeutics,(12) genomic resources for whipworms and molecular adaptation to
anthelmintic pressure in *Trichuris* remain limited.(12-14) Currently, only the genomes of three
Trichuris species are publicly available on WormBase ParaSite, *T. suis*,(15) *T. muris(16)* and
T. trichiura(16).

We present a new species of *Trichuris*, *Trichuris incognita n. sp.*, naturally infecting humans
in Côte d'Ivoire. *T. incognita n. sp.* occurs where drug treatment fails and is characterized by
a high number of TBB orthologs. Phylogenetic analysis based on the species tree of 12434
orthologous groups of genes, places the isolated worms within a new, monophyletic clade
which is genetically closer to *T. suis* than to *T. trichiura*. We elucidate the genome of this new
species, leveraging whole-genome-sequencing (WGS) data to produce a high quality
reference genome of this species. Finally, we provide the first ever dataset of human infecting
helminths comparing pre- and post-treatment WGS data. We investigate the whole genome
sequencing data of 747 isolates in context of resistance by looking at duplication events across
the species tree and conducting a GWAS.

Methods

Expulsion study design fieldwork

This trial was conducted in the Dabou and Jacqueville districts of Côte d'Ivoire. This area was
selected, because communities with high endemicity were identified as part of a previous trial
on *T. trichiura*.(9) All community members aged 6-12 who were positive for *T. trichiura* in
duplicate Kato-Katz smears, with an infection intensity of at least 200 eggs per gram of stool
(EPG), were eligible for trial inclusion. Excluded from the trial were individuals with reported
major systemic illnesses or chronic disease, known allergy to study medications and having
received anthelmintic treatment with ivermectin in the past two weeks.

This trial was conducted in accordance with the protocol, International Conference on
Harmonization Good Clinical Practice E6 (R2) (ICH-GCP) and the current version of the
Helsinki Declaration. Parents or guardians of participating children provided written informed
consent. Children provided written assent. All authors take responsibility for the accuracy and
completeness of the data and the fidelity of the trial to the protocol, which is available together
with the statistical analysis plan in appendix 1. The protocol was approved by independent
ethics committees in Côte d'Ivoire (reference numbers 156-22/MSHPCMU/CNESVS-kp and
ECCI00918) and Switzerland (reference number AO_2022_00028).

Treatment and sample collection from humans

Guardians/caregivers were invited to participate in an information session. The research team
explained the purpose and procedures of the study, as well as potential benefits and risks of
participation. The timeline of the procedures is summarized in Figure 1a. For initial diagnosis,
participants were asked to provide one stool sample from which duplicate Kato-Katz thick
smears using 41·7 mg of stool were prepared and assessed under a light microscope for the
identification of *T. trichiura*, *A. lumbricoides*, and hookworm ova by laboratory technicians.
10% of all Kato-Katz slides were randomly chosen for subsequent quality control by picking
every tenth slide of all slides read by each laboratory technician on the respective day. All
eligible participants were treated with albendazole-ivermectin at day 0. Albendazole, the
product of Glaxo Smith Kline (Zentel®), was a single tablet of 400 mg. Ivermectin, the product

of Merck (Stromectol®), was given at a dose of 200 µg/kg. On day 8 a single oral dose of 20 mg/kg of oxantal pamoate was administered. Oxantel pamoate tablets (500 mg) were provided by the University of Basel.

From day 1 or 2 and 8 onwards, 24h-stool was collected, based on previously reported expulsion dynamics and weighed for 2-3 days.(17, 18) Expelled worms were counted, washed twice with distilled water, once with sterile PBS and then stored in pure ethanol at 4°C. A picture of each worm in ethanol was taken and is provided on online (https://github.com/max-baer).

Sample collection from pigs

To obtain *T. suis* worms, a list of all pig keepers was drawn by the help of local village chief. Pig keepers were gathered in formal meetings where the purpose of the study was explained. Pig keepers were asked to give written informed consent to participate in the study. Fecal samples were then collected directly from the rectum using gloves lubricated with glycerin, by a veterinarian. The samples were analyzed using the Mini Flotac technique.(19) 2 pigs that tested positive for *T. suis* roaming freely in Tiagba were sacrificed and worms were collected directly from the cecum and stored in either pure ethanol at 4 °C or mashed and stored in RNA-later at -20 °C.

Defining a holotype for *T. incognita n. sp.* and depositing a sample at the natural history museum in Basel

A female specimen of *T. incognita n. sp.* isolated from *Homo sapiens* in the town of Tiagba in the district des lagunes in Côte d'Ivoire has been chosen as the holotype and was archived at the natural history museum in Basel under the number NMB-79 a in absolute ethanol together with a male paratype. All authors of this publication have been registered as the discoverers of this new species. This work and the nomenclatural acts it contains for the new species have been registered in ZooBank: LSID: 34409100-BF94-4A0D-9088-D28215D231F6.

DNA extraction and high-throughput sequencing

Worms stored in absolute ethanol were photographed and used for sequencing experiments. The posterior end of the worms was severed and DNA was extracted using a DNeasy Blood & Tissue Kit (Qiagen, Cat: 69504) according to the manufacturer's protocol from the anterior part of the worm. Sequencing libraries for illumina were prepared using the NEBNext® Ultra™ II FS DNA Library Prep Kit (New England Biolabs, Cat: E7805L) according to the manufacturer's protocol after which DNA concentration and library size was determined (HS NGS Fragment Analyzer). Libraries were sequenced using 150 bp paired-end chemistry at an average 20X coverage on an Illumina NovaSeq6000 platform with an S4 flow cell at the Genomics Facility Basel as part of the Department of Biosystems Science and Engineering. Sequencing libraries for Nanopore were prepared using the Native Barcoding Kit 24 V14 (Oxford Nanopore Technologies, Cat: SQK-NBD114.2) from a single whole female worm and sequenced on a PromethION flow cell and PromethION 2 device.

PCR and Nanopore sequencing experiment of the TBB gene

Two primers were selected surrounding the regions containing known mutations (Phe168, Glu198, Phe200), observed variable position (Ser194) and predicted truncation site. The primer sequences were AAAGAGACCGGACATTCGC for the forward primer and TGAATTGCCTGGTTTCTAGAATG for the reverse primer and the amplified region was 402 base pairs. Genomic DNA from one sequenced worm was used as input and the OneTaq® One-Step RT-PCR Kit was used to amplify the region during 40 cycles with an annealing

temperature of 56°C for 30 seconds and an extension temperature of 68°C for 1 minute
Sequencing libraries for Nanopore were prepared using the native barcoding kit 24 V14
(Oxford Nanopore Technologies, Cat: SQK-NBD114.2), and sequenced on a Flongle flow cell
and the MinION Mk1c device. The dorado basecaller (version 0.7.2, model
400bps_hac@v5.0.0) was used for duplex basecalling. Reads were filtered by alignment
score (AS>600) and length (>390 BP) and aligned using minimap2 to the reference TBB
sequence. Variants were called using bcftools and coverage plots were created in R.

Pipeline for raw read processing and mitochondrial genome assembly

For all sequencing, assembly and annotation pipelines Nextflow (version 23.04.1 build 5866)
was used with Java (version 11.0.3).(20). Raw reads were first processed using Trimmomatic
(version 0.39).(21) for PE reads, adapters were removed and N-bases trimmed. Next, reads
were channeled to getorganelle (version 1.7.7.0) to reconstruct the mitochondrial genome.(22)
Runs which resulted in a complete mitochondrial genome were further processed by flipping
in cases where the reverse complement was generated and adjustment to COX1 as starting
gene. Reference and newly constructed mitochondrial genomes were annotated and
visualized using MitoZ (version 3.6).(23) Coding genes were then extracted and aligned as
amino acid and nucleic acid sequences using MAFFT (version 7.490).(24) All reconstructed
mitochondrial genomes and the aligned MAFFT files used as input for phylogenetic inference
are provided online.

Phylogenetic analysis

BEAST2 (version 2.7.5) was used for all phylogenetic inference with a pure birth process (Yule
process).(25) Aligned concatenated nucleic acid sequences from all assembled and reference
mitochondrial genomes were treated as homochronous. The JC69 substitution model with
gamma-distributed rate heterogeneity was used (JC69 + Γ_4) and a strict molecular clock was
assumed. Trees and clock models were linked and all model parameters were estimated
jointly. A Markov Chain Monte Carlo was run for each analysis. Tracer (version 1.7.2) was
used to assess convergence and effective sample size. The percentage of samples discarded
as burn-in was at least 10%. For the amino acid inferred species tree, 2 assembled sequences
of each subclade clade with a posterior distribution over 0.95 were selected at random. The
mito REV substitution model was used and all genes were partitioned separately, allowing for
different rates for each gene. Effective sample size (ESS) was at least 900 for each of the
inferred parameters. The intraspecies phylogeny was inferred using a TN96 + Γ_4 model. Each
gene was partitioned separately and codon positions 1 and 2 were partitioned separately to 3.
The mean mutation rates of the genes and there codon positions are provided in the
supporting information and show a faster mutation rate for codon position 3 in all cases.

Pipeline for Hybrid-genome assembly and gene prediction

Nextflow (version 23.04.1 build 5866) was used with Java (version 11.0.3). Nanopore raw
reads from a single worm were channeled into chopper (version 0.6.0) where reads below 5kb
and low quality reads were filtered out and adapters trimmed.(26, 27) Trimmed reads were
channeled into Kraken2 (version 2.1.1) and reads with a contamination ratio >0.5 were filtered
out with a custom R (version 4.3.0) script, resulting in a 50X coverage.(28) Illumina reads of
the same worm at a 65X coverage were channeled into Kraken2 where reads of impurities
were identified and filtered out. Illumina and Nanopore channels were combined to run a hybrid
de-novo genome assembly using MaSuRCA (version 4.0.9) which allowed for paired end
illumina reads and nanopore reads.(29) A total of 1564 contigs were assembled and the raw
assembly is provided on github. The BUSCO showed a completeness of 64.3% of complete
BUSCOs, 11.2% fragmented and 24.5% missing and the N50 was 0.26 Mb. The outputted

contigs were channeled into RagTag (version 2.1.0) for correction and scaffolding using the
T. trichiura (PRJEB535) reference genome.(30) The corrected scaffolds were then assigned
to one of three linkage groups from *T. muris* using Liftoff (version 1.6.3),(31) as described in
literature.(32) The genome was frozen for quality assessment using BUSCO (version 5.1.2
with Metazoa_obd10 database)(33) and QUAST (version 5.0.2).(34) At this point all available
reference whole genomes of *Trichuris* species and *Trichinella spiralis* were channeled into the
pipeline to run a de-novo gene prediction on all reference genomes and reduce
methodological bias from gene prediction software. Repeat regions were masked using
RepeatMasker (version 4.1.4) before channeling the genome to BRAKER (version 3.0.6,
Braker2 pipeline using the Metazoa protein database from OrthoDB 11).(35, 36) Gene
prediction statistics including exon and intron count, mean length and median length were
gathered using a custom bash script.

Phylogenetic orthology inference, species tree, orthologous group functional
annotation and GO term enrichment analysis

Orthofinder (version 2.5.5) was used to infer orthology and the species tree of the braker2
predicted genes.(37) 81425 out of 92202 (88.3%) genes were classified into 12434
orthologous groups with a mean group size of 6.5 and median group size of 5. Gene ontology
of orthologous groups of interest was inferred using InterProScan (version 5.63-95.0). A list of
resistance associated genes was downloaded from WormBase and blasted against all
orthologous groups using an E-value threshold of $1e^{-50}$ to identify orthologous groups of
resistance associated genes. In SI Table 4 a list of all genes with matching orthologous groups
and Gene IDs from WormBase ParaSite is provided. Species tree with heat map was
visualized using iTOL (version 6.8.1).(38) Duplicated, gained, lost and retained genes across
the species tree were inferred using OMA standalone (version 2.6.0) and pyHam (version
1.1.11).(39, 40) Massively expanded gene families were defined as ≥7 copies in *Trichuris*
incognita n. sp. and ≤4 in all other species and a list is provided in the supporting information
together with the functional annotation. The orthology of predicted transcripts was inferred
analogously. A custom R-script was used to visualize the Venn diagram and identify
transcripts shared by *T. incognita n. sp.*, *T. trichiura* and *T. muris.* SI Table 3, a list of these
transcripts with their functional annotation is provided. Protein structures were predicted using
Alphafold (version 2.2.0, template date cutoff was set to 2021-09-01) and visualized with
PyMOL (version 2.5). GO terms were extracted from the InterProScan annotated genes file of
T. incognita n. sp. (background GO terms) and of *T. incognita n. sp.* genes shared with *T.*
trichiura but not with *T. suis* or *T. muris* (GO terms of interest). A fishers exact test was
performed on the GO term counts of each GO term in the intersection compared to the
background GO terms. All gene lists of *T. incognita n. sp.* and the InterProScan annotations
are provided on github.

Pipeline for variant calling

Nextflow (version 23.04.1 build 5866) was used with Java (version 11.0.3). Variants were
called by adapting the analytical pipeline in by Doyle *et al.* into Nextflow .(32) Raw reads were
first processed using Trimmomatic (version 0.39).(21) BWA (version 0.7.17) and SAMtools
(version 1.14) were used to index the assembled reference genome, and align, sort and filter
raw reads.(41) Flagstat and Kraken2 (version 2.1.1) were used as QC tools. The coverage
ratio was used to classify the gender of each worm by analyzing the coverage ratio of the
largest X-chromosome scaffold to the largest scaffold on chromosome 2. Next, the standard
GATK (version 4.2.6.1) pipeline was followed for variant calling, base quality score
recalibration (BQSR) and genotyping. First, duplicates were marked using the
MarkDuplicatesSpark option in GATK. Basecalling was done in a first round to obtain a subset

of very likely SNPs for base quality score recalibration. Plots were generated for QUAL, DP,
QD, FS, MQ, MQRankSum, SQR, ReadPosRankSum. Heterozygosity was set to 0.015 and
indel heterozygosity to 0.01.(32) Initial hard filtering of the variants for BQSR was done in a
way to exclude 5% of SNVs with the following lowest quality metrics: QUAL < 194 || QD < 10.3
|| MQ < 43.61 || FS > 9.9 || SOR > 2.30 || MQRankSum < -1.269 || ReadPosRankSum < -
1.174 for SNPs and QUAL < 101 || QD < 5.3 || FS > 11.4 || ReadPosRankSum < -1.336 for
indels. Next, BQSR was conducted before generating a merged gVCF which was used for
genotyping and then filtered for SNVs using the following hard filter expression: QUAL < 35 ||
QD < 1.3 || MQ < 28.76 || FS > 50.5 || SOR > 2.30 || MQRankSum < -4.261 ||
ReadPosRankSum < -1.421 for SNPs and QUAL < 32 || QD < 0.5 || FS > 30.1 ||
ReadPosRankSum < -1.836 for indels. Indels were filtered out and the variants were further
filtered using the genotype filter expression DP < 3 and a minor allele frequency of 0.02 which
resulted in a total of 6'282'017 SNPs. Next, genes in the VCF file were annotated using
BCFtools (version 1.15) and BEDTools (version 2.30.0) and the braker2 produced gff file
containing all genes of *T. incognita n. sp.*. Using SnpEff (version 5.2c) a total of 111'994
missense SNPs leading to a different amino acid on the genes of *T. incognita n. sp.* . This set
of SNPs as well as the non-filtered set was used as an input in the GWAS.

Pipeline for genome wide association study

Plink (version 1.9) was used to conduct the genome wide association study.(42) First, the
principal components were analyzed, as presented in SI Figure ? to assess population
stratification and correct any falsely assigned sexes. Next, the phenotype data (SI Table 8) on
treatment one (T1) or treatment 2 (T2) was added using the "--pheno" option as well as the
worm sex using the "--update-sex" option. Next, the complete set of variants, as well as the
subset of variants leading to missense mutations on predicted genes were filtered using a
minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the
Hardy-Weinberg Equilibrium at a significance level of p=0.001 resulting in 1254247 variants
over all, and 56273 missense variants on predicted genes, which were used for association
testing. The GWAS was performed a total of 24 times and the results of each run are presented
to be as unbiased as possible. Each GWAS for both variant sets was performed once for both
sexes together, and once for each sex individually, to account for variation associated to worm
sex. Next, each GWAS was performed once with population stratification and once without
population stratification as the PC1 and PC2 did not show any significant association to
treatment, however, did show association to some villages in both male and female
populations. Finally, the GWAS was ran once with a logistical model and once with a linear
model resulting in 24 runs of the GWAS. The GWAS' were ran using linear and logistic models
with the "--assoc" or "--logistic" option respectively, with both sexes, only females and only
males. Next, a principal component analysis was conducted to account for population
stratification. The option "--pca" was used on the file containing all filtered variants. As an
example for the subset of variants on coding genes, which is representative for all runs, the
analysis of the principal components revealed an eigenvalue of 52.7 for the first principal
component and 3.1 for the second, indicating that the consideration of only the first principal
component will account for most population stratification. Next, the logistic and linear models
were run again under consideration of the first principal component by using the options "--
covar" and "--covar-number 1" and the "--linear" or "--logistic" option for the linear or logistic
model respectively for both sexes, only females and only males respectively on both sets of
variants. To visualize the results and generate the QQ-plots the R package qqman (version
0.1.9) was used.

Data availability

Raw sequencing data of 747 samples together with the associated metadata and the whole genome of *T. incognita* (TaxID: 3388467) is available under the Sequencing Read Archive, BioProject "PRJNA1282940". 551 assembled mitochondrial genomes including *T. incognita n. sp.* and *T. suis*, as well as reference mitochondrial genomes and the *de novo* assembled *Trichuris incognita n. sp.* reference genome are available under the BioProject number and on GitHub https://github.com/max-baer. The publicly available mitochondrial genomes used for comparative analysis are available from ENA and NCBI. There are no restrictions on data availability.

Code availability

Custom code to analyze data and reproduce the figures presented is available at https://github.com/max-baer. Where available, all input scripts and log files are provided including nextflow log files documenting a complete run, BEAST2 XML input and log files.

Results

Trichuris expulsion study confirms a new species of whipworm infecting humans, responding poorly to conventional albendazole-ivermectin treatment

We screened 670 children in 7 villages, namely Akakro, Ahouya, N'Doumikro, Tiagba, Bekpou and Teffredji in southern Côte d'Ivoire for trichuriasis. 243 participants were enrolled in the study and received a single dose albendazole-ivermectin combination treatment (T1) after which stool was collected daily and screened for worms for up to four days. Subsequently, 8 days after T1, the same individuals received a single oral dose of the trichuricidal drug oxantel-pamoate (T2) after which stool was collected for 2 consecutive days, as presented in Figure 1a. We isolated 271 worms after T1, indicating partial efficacy of the combination treatment, with an expulsion peak on days 2 and 3 and 827 worms after T2 with expulsion peaking on day 1 and 2 after treatment. A total of 1098 *Trichuris* worms were collected, of which 747 provided enough DNA for genomic analyses. The female to male ratio of the worm sex was 3.42:1 and 1.6:1 during T1 and T2 respectively. A significant correlation ($R^2 = 0.34$; $P = 0.0061$, SI Figure 1) was observed between the number of expelled worms and the mean egg count per day, yielding a fecundity of 14307 eggs per female worm per day, as presented in SI Figure 1. Figure 1b shows near complete expulsion of all worms as the median EPG dropped from 1524 to 24 at the follow-up time point on day 12. To assess the genetic relatedness of human worms with *T. suis*, 25 individual worms from 2 swine hosts were collected directly from the caecum. A picture of each worm which was used to identify the worm sex is provided on github.

Figure 1: Timeline of expulsion study including the number of expelled worms and the female/male distribution of the sequenced worms

a Timeline of expulsion study including the number of expelled worms and the female/male distribution. T1: albendazole-ivermectin combination treatment, standard-of-care. T2: Oxantel-pamoate treatment. Representative pictures of an expelled female and male worm are provided on the top right and bottom right respectively. Further morphological data is provided in Figure 3. **b** EPG before T1 and after T2.

To phylogenetically place these new isolates of whole worms from humans into context of the *Trichuris* genus, an analysis on the mitogenome was conducted as presented in Figure 2. The phylogenetic analysis was done using the coding regions of the mitochondrial genomes of 535 individual worms that yielded a complete mitogenome and is presented in SI Figure 2. The reconstructed mitogenomes contained 13 protein coding genes (*cox*1-3, *nad*1-6, *nad*4L, *atp*6, *atp*8 and *co*B). Concatenated nucleic acid sequences from all assembled and reference mitochondrial genomes showed that the *Trichuris* species encountered in Côte d'Ivoire forms a separate clade from the canonical human infective *T. trichiura*, closer to *T. suis*, while the phylogenetic placement of the 16 sequenced *T. suis* fall within the expected clade of *T. suis* reference mitochondrial genomes. Previously reported sequences of a *Trichuris* species, isolated from a colobus monkey in Spain by Rivero et al.,{Rivero, 2020 #19} show the closest genomic relationship with the species from Côte d'Ivoire. Finally, the phylogenetic tree of the *T. incognita n. sp.* clade was plotted in SI Figure 3 to see if any of the clades within the species correlates to treatment, worm sex, host sex, host, coinfection with other worms, however no correlation could be identified. This species was named *Trichuris incognita n. sp.*

Figure 2: Phylogenetic placement of expelled *Trichuris* worms from Côte d'Ivoire

379 Bayesian inferred phylogenetic tree based on the amino acid sequences of genes in the mitochondrial
380 genomes of 535 *Trichuris incognita n. sp.* worms and available reference mitochondrial sequences,
381 using *Trichinella spiralis* as an outgroup. Timescale indicates numbers of substitutions per site.

Morphological description of *T. incognita n. sp.*

This parasite has a threadlike structure, filiform. Both female and male can be divided into a
thin hair-like anterior part and broader handle-like posterior section. The posterior section of
the female worms is slightly ventrally incurved compared to the strong ventral incurvature of
the male, adopting a Fibonacci spiral-like structure. The overall length of the worms (displayed
in Figure 1a) is 45 mm and 49 mm and the ratio of anterior to posterior is 2.5 and 3.1 for female
and male worms respectively. An analysis of a subset of five male and five female worms are
within literature reported min-max values for *T. trichiura* lengths as shown in SI Table 1.(43)
The main organs of the male posterior end are the intestine, testis and ejaculatory duct, which
run in parallel along the long axis of the body (Figure 3a). At the height of the ending of the
testis, the ejaculatory duct and intestine join, forming the cloaca, which opens at the posterior
end of the male body. The distal cloacal tube contains the spicule. The spicule is surrounded
by a shiny spicule sheath (Figure 3b, 3c) with a granular surface structure. The female
posterior starts at the esophagus-intestinal junction, where the vulva is located (Figure 3d).
The vagina shows thick walls, which connect back into the oviduct (Figure 3e). The cuticle of
the anterior structure shows two patterns, one, which is striated with transverse grooves on
one side and the other, a tuberculate band (Figure 3g). Within the anterior part is the
stichosome, comprised of a row of stichocytes surrounding the esophagus which starts at the
mouth (Figure 3i), and continues to the posterior part of the worm merging into the intestine
(Figure 3d) and marking the division between anterior and posterior section. The ova present
in the oviduct contained two polar plugs, were 68 μM long and 29 μM wide (Figure 3f) placing
them closer to a larger observed egg population of two distinct ova groups of *T. trichiura* in
literature.(44)

Figure 3: Morphology of *T. incognita n. sp.*

Figure 3: Morphology of *T. incognita n. sp.* the same scale was used for a,b,d,e,g and h, and a different
one for c, f and i and is provided in the bottom right corner. **a** Posterior end showing male organs
including testis, ejaculatory duct, intestine, union of intestine and ejaculatory duct, cloacal tube and
spicule with a ruptured sheath. **b** Male spicule with an intact sheath partially covering the spicule. **c** 40X
magnification of the shiny spicular sheath showing a surface structure. **d** Morphology showing female
reproductive organs at the intersection between anterior and posterior part of the worm including
stichosome, esophagus-intestinal junction, intestine, vagina and vulva. **e** Enlarged oviduct filled with *T.*
incognita n. sp. ova. **f** 40X magnification on ova showing polar plugs. The ova are 68 μM long and 29
μM wide in this image. **g** Anterior showing one row of stichocytes and 2 types of cuticle patterns. The
left side showing transverse grooves and the right side the tuberculate band. **h** Anterior ending leading
up to the mouth. **i** 40X magnification of the mouth.

***T. incognita n. sp.* hybrid *de novo* whole genome assembly and comparative genomic**
analyses

The *de novo* assembled genome of *T. incognita n. sp.,* sequenced at a coverage of 50X
Nanopore and 65X Illumina reads, resulted in a length of 84.26 Mb, compared to 80.57 Mb of
T. trichiura, 63.84 Mb, 73.46 Mb and 70.45 Mb of *T. suis* and 111.83 Mb of *T. muris.* The *T.*

incognita n. sp. genome consisting of 387 contigs with an N50 of 11.78 Mb is presented in
Table 1. The genome completeness amounted to 75.4% of whole, duplicated, and fragmented
single copy orthologues as determined by BUSCO. GC content was 43%. Genes were
predicted *de novo* for all available reference whole genomes using the braker2 pipeline and
the metazoa database, resulting in the prediction of 13739 genes for *T. incognita n. sp.* An
exploratory analysis of these genes in context to the *in silico* predicted genes of *T. trichiura*,
T. suis and *T. muris* presented in SI Table 2 and SI Table 5 identified one family of each
serpins (OG0000305) and chymotrypsin-like serine proteases (OG0010319) that are shared
only by *T. trichiura* and *T. incognita n. sp.* (SI Figure 4) additionally to a family of tetraspanins
(OG0009162) also shared by *T. muris.*(45, 46)

Table 1: Whole genome assembly metrics

	Trichuris incognita n. sp.	*Trichuris trichiura*	*Trichuris suis*	*Trichuris suis*	*Trichuris suis*	*Trichuris muris*	*Trichinella spiralis*
Accession	NA	PRJEB535	PRJNA208416	PRJNA208415	PRJNA179528	PRJEB126	PRJNA12603
Genome Size [Mb]	84.26	80.57	70.45	73.46	63.84	111.83	63.51
Number of contigs	387	113	1131	1470	306	803	6819
N50 [Mb]	11.78	11.3	0.45	0.5	1.32	28.94	6.37
GC percentage [%]	42.8	42.3	43.5	43.6	43.31	44.6	33.9
Complete single BUSCOs [%]	61.3	64.3	65.6	65.4	65.2	61.5	64.8
Duplicated BUSCOs [%]	2.7	1.8	0.4	0.7	0.4	3.9	3.8
Fragmented BUSCOs [%]	11.4	10	10	9.7	9.9	10.3	7
Missing BUSCOs [%]	24.6	23.9	24	24.2	24.5	24.3	24.4
Number of genes	13739	11809	11224	11884	9755	20537	13254
Total exon number	82285	71304	69897	71543	66000	93618	103944
Mean exon length [bp]	255	240	235	239	222	281	193
Median exon length [bp]	152	150	150	151	148	164	136
Total intron number	67462	58645	58015	58973	55584	71798	88078
Mean intron length [bp]	267	291	274	278	277	315	187
Median intron length [bp]	57	57	56	57	56	57	74
Total coding sequence [Mb]	20.98	17.14	16.45	17.07	14.63	26.23	20.03
Sequencing technique	Illumina + Nanopore	Pacbio	Illumina	Illumina	Pacbio	Pacbio + Illumina	Whole genome shotgun*

434 *bacterial artificial chromosome (BAC)

The species tree confirms the existence of a new species of *Trichuris*, and reveals highly duplicated genes associated to microtubule dynamics and mitotic processes

The species tree of the *Trichuris* genus presented in Figure 4a was inferred using species tree
inference from all genes (STAG)(47, 48) with 12434 orthologous groups of the braker2
predicted genes in the publicly available whole genomes of *Trichuris* species, *T. incognita n.*
sp. and *Trichinella spiralis*. The species tree shows the same topology as for the mitogenome
inferred phylogeny.

We further inferred the gene orthology of all genes predicted in the publicly available whole
genomes of *Trichuris* species, *T. incognita n. sp.* and *Trichinella spiralis* to identify duplication
events of genes that have been associated with resistance to either albendazole or ivermectin
in helminths. These resistance-associated genes include TBB for albendazole or glutamate-
gated chloride channel subunits (*glc*-1, *glc*-2, *glc*-3, *avr*-14, *avr*-15), voltage gated chloride ion
channels (*clh*-3), sodium:potassium:chloride symporter (*nkcc*-1), bestrophin chloride channel
(*best*-19), p-glycoprotein ABC transporters (*pgp*-1, *pgp*-2, *pgp*-9, *pgp*-11), and transcription
factor *cky*-1, for ivermectin respectively.(14, 49) Amongst these genes, one orthologuous
group (OG0000671) of the TBB gene showed 9 orthologs of beta tubulin gene in *T. incognita*
n. sp. and only 1 ortholog in all *T. suis* and *T. trichiura* reference species as presented in
Figure 4a. One gene was fragmented into 2 sections, g475 and g476, and g5082 was also

found to be fragmented. Intriguingly, the α-tubulin gene was only present as a single ortholog. SI Figure 5 shows the 7 predicted TBB genes from the braker 2 pipeline. None of the genes show previously encountered Phe168, Glu198 or Phe200 mutations. Position 194, showed the existence of 3 different variants carrying either Ile, Met or a polar Asn. Reference sequences of *T. trichiura*, and *T. suis* also show different variants at this position. *In silico*, a truncation was predicted in 3 of the 7 TBB genes as shown in SI Figure 5, which warrants further transcriptomic verification. To further investigate, if the variants at position 194 were reproducible in an independent Nanopore sequencing experiment of a single isolate, a region was amplified that was present in all genes and contained the predicted variants (primer sequences are provided in SI Figure 6). As shown in SI Figure 6, the different variants at position 194 were confirmed and the coverage ratio from the IGV software indicated that 8% of all reads carried an Asn, 37% carried an Ile, and 55% a Met, indicating that out of all orthologs of the TBB gene, one ortholog is present in the Asn form. Notably, these variants are very close to the predicted binding site of albendazole and are at variable sites compared to reference sequences.

The gene tree of the orthologous group of TBB, OG0000671, shown in Figure 4c gives insight into the history of these duplication events. The gene tree shows a similar distance in number of substitutions per site from *T. suis* gene 4606 to *T. incognita n. sp.* gene 34, compared to the distance from *T. incognita n. sp.* gene 472 to *T. incognita n. sp.* gene 34. This indicates that interspecies genetic variation is similar to intraspecies genetic variation for at least 2 clades of TBB genes. Clade one, including gene 7506, 529, 472, 460, 463, and clade two, including gene 34. Genes 475, 476, and 5082 were not considered, as they were found to be fragmented.

Figure 4: Duplications across the species tree show a high number of β-tubulin gene orthologs.

a Species tree indicating gene duplication, gains, losses and retention as pie charts across the tree. Heat map of ortholog numbers for all genes present in orthologues groups of genes, which have previously been associated to resistance and the α-tubulin gene. **b** Coverage ratio compared between worm sex (n=747), treatment (n=747) and from *T. incognita n. sp.* (n=747) to *T. suis* (n=26) from Côte d'Ivoire and other species. Three *T. suis* genomes and one *T. trichiura* genome were used to calculate the average coverage in the species containing only one gene in OG0000671. *** signifies a p-value < 0.001 from a two tailed t-test. **c** TBB gene tree of OG0000671, timescale indicates numbers of substitutions per site.

We then sought to conduct an unbiased investigation of gene families which are as expanded
or more expanded compared to the β-tubulin gene in *T. incognita n. sp.* and identified gene
families associated with microtubule function and mitotic processes. Massively expanded
gene families as in ≥7 copies per gene in *T. incognita n. sp.* and ≤4 in all other species were
investigated and a list of gene families and their annotation is provided in the supporting
information (SI Table 6). From 81 gene families, 68 could be associated to DNA transposable
elements and retrotransposons, 6 remained unknown and without blast hits and 7 could be
assigned a functionality. Apart from the TBB gene, other genes related to microtubule function
and flexibility were found within these 7 assignable orthologous groups. Amongst these highly
duplicated genes is the motor protein kinesin and MAP65/Ase1, which promotes microtubule
flexibility and prevents microtubule severing by cross-linking.(50, 51) Finally, Condensin,
Transcription factor (TF) IIIC, and a haspin like kinase, were identified and are involved in
chromatid condensation, organization and cohesion respectively.

To extrapolate the TBB duplication events from the assembled genome of one single worm to
the population, we leveraged the ratio of the average coverage of illumina reads of the
amplified TBB region compared to the longest autosomal contig present in the genome of all
747 worms as shown in Figure 4b. Intriguingly, the gene ortholog count was reflected in the
average overall coverage showing a significant difference between worm sex (p-value <
0.001) but not showing any significance with regard to treatment. On average 11 copies were
present in female worms and 8 copies in male worms. To compare the coverage ratios against
species, where only one ortholog was identified, the raw sequencing data of 26 *T. suis* worms
isolated in Côte d'Ivoire was used as a comparison, where, intriguingly, only 1-2 copies of the
TBB was present. Thus, one unique feature about *T. incognita n. sp.* is the high number of the
TBB gene orthologs. The coverage ratios were further compared to coverage ratios from
publicly available raw sequencing data that was downloaded for *T. suis* (PRJNA20815,
PRJNA20816, PRJNA179528) and *T. trichiura* (PRJNA304165), the same region of TBB was
used and the coverage ratio was calculated again from the amplified TBB region to the
autosomal region indicating one ortholog of the TBB gene.

This work and the nomenclatural acts it contains for the new species have been registered in
ZooBank: LSID: 34409100-BF94-4A0D-9088-D28215D231F6. Furthermore, a female
holotype has been archived at the natural history museum in Basel under the number NMB-
79 a.

Genome wide association study does not provide substantial evidence of adaptation to
drug treatment

To investigate potential SNPs associated with resistance, a genome wide association study
(GWAS) was conducted with sequencing data of 747 individual worms, 721 of which passed
the filtering, comparing the populations isolated after T1 and T2 respectively comprising of
179 drug sensitive (T1) and 542 drug non-sensitive (T2) isolates. Investigation of the PCA
presented in SI Figure 7, showed that worm-sex contributed the most to population
stratification. As a result, the GWAS was conducted once for both sexes pooled, and for male
and female worms separately. In the PCA of the subpopulation, significant population
stratification was associated with the village of isolation. However, no association was found
to T1 or T2 in the subpopulation, justifying the consideration of population stratification only
on a worm sex level. Both a logistical and linear model was used to investigate 1254246
overall mutations and a subset of 56273 missense mutations on *in-silico* predicted genes and
a genome wide significance level of 3.99E-8 and 8.89E-7 respectively using the Bonferroni
correction at α=0.05 on the combined sample cohort of male and female worms and on each
sex individually resulting in a total of 24 different runs presented in Figure 5 and SI Figures 8-

15. QQ-plots showed consistent artificial inflation when using combined data of male and female worms, as seen in Figure 5a (right half of the panel). Furthermore, population stratification occasionally resulted in increased deviations from expected and observed values, for the probability of association to treatment outcome. When running only male or female isolates, the artificial inflation was mitigated (Figure 5b, 5c on the right). In total, 29 SNPs were identified with genome wide significance in all the 24 GWAS' combined and are presented with their sequence in SI Table 7. Using a linear model without population stratification on missense mutations on coding genes (Figure 5), a total of 3 genes carried SNPs with genome wide significance in the male population: g207, a muscle M-line assembly protein unc-89 homolog; g1478, a FSA C domain containing protein; and g2562, an integrase catalytic domain-containing protein. The GWAS with a linear model without population stratification on all SNPs yielded 22 hits on 10 genes, including a voltage-gated chloride ion channel.. No further missense SNPs were identified with a genome wide significance. The QQ-plots were most consistent in the logistic model without population stratification as shown in SI Figure 10. Using this model, no SNPs with genome wide significance were identified.

SNPs on expanded gene families were also investigated. However, no genome wide significant SNPs were found in any of the 24 GWAS'. The strongest association of a missense SNP associated to drug treatment on the TBB gene was the Asn194Ile mutation with a p-value of p = 0.12 which is far below the genome wide significance level.

Figure 5: Genome wide association study of worms expelled after albendazole-ivermectin treatment (T1) versus worms expelled after oxantel-pamoate treatment (T2).

GWAS displaying p-values of missense SNPs associated to treatment outcome (T1 vs T2) using a linear model without population stratification in PLINK. PCA is provided in the SI. A total of 24 GWAS' were performed as described in the methodology section, the results of all of them are provided in the SI. The linear model without population stratification yielded the most hits while maintaining stable QQ plots for both sexes individually. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the Bonferroni corrected significance of 8.89E-7 at α=0.05 and 56273 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138 controls. **c**

569   Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS
570   hits, from each of the 24 analyses, the sequence, the best blast hit, the location and functional
571   annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

Discussion:

Our study presents an in-depth characterization of *T. incognita n. sp.* found to be less
susceptible to albendazole-ivermectin treatment and morphologically indistinguishable to *T.*
trichiura. A genome wide association study and the investigation of genes previously
associated with resistance, did not conclusively show adaptation to drug pressure within the
same species of *Trichuris*.

The existence of a species of *Trichuris*, phylogenetically distinct from the canonical human-
infective species, has been proposed in recent studies involving amplicon sequencing by
Rahman *et al.*,Venkatesan *et al.*, and already in 2012 by Nissen *et al.*(11, 52, 53) Adopting
the genetic species concept, our data confirm the existence of this species at the whole
genome level, using the species tree constructed from 12434 orthologous groups, and 535
mitogenomes of 747 individually isolated and sequenced worms, all falling into the same
clade. The whole-genome assembly metrics align with established *Trichuris* genomes,
underscoring the robustness of our dataset.(15, 16, 32) However, structural variations might
be missed, as the *T. muris* genome was used as a reference for scaffolding, similar to what
was done in literature.(16) The prediction of genes in the publicly available *T. trichiura*, *T. suis*,
T. muris and *Trichinella spiralis* genomes, and the newly assemble *T. incognita n. sp.* genome,
using the same reference database, allowed the exploration of orthologous gene groups and
duplication events across the species tree. The identification of a family of serpins and
chemotryphsin-like serine proteases uniquely present in human infecting *Trichuris* may be of
particular interest, as these proteins, which were identified to be upregulated in the stichosome
of *T. suis*, have gained particular interest in the field of immunomodulatory diseases in general
and vaccine development for *T. spiralis*.(15, 54, 55)

Our investigation of duplication events across the species tree revealed a unique characteristic
of *T. incognita n. sp.*: A high number of TBB gene orthologs. Duplication events are
encountered across the tree of life and constitute an important mechanism to provide new
genetic material. In eukaryotes, including *C. elegans*, gene duplication rates are estimated to
be as frequent as single-nucleotide polymorphisms. However, they have never been
investigated within the *Trichuris* genus.(56, 57) Drawing upon previous studies linking gene
duplication to drug adaptation across diverse parasitic organisms, and deducing modes of
action of drugs from observations of duplication events, we sought to investigate gene
duplication events in connection to drug insensitivity.(58, 59) In total, 7 variants of the β-tubulin
gene were identified in the whole genome, providing substantial redundancy that may allow
for adaptation through sub-functionalization and increased gene expression. This contrasts
with the low number of TBB orthologs in *T. trichiura*. It is of note that *T. trichiura* does not
respond well to single benzimidazole treatment. Leveraging coverage ratios, duplication
events could be confirmed in the whole population for each individual worm. Notably, while β-
tubulin exhibited significant duplication, α-tubulin did not. Furthermore, massively expanded
gene families are associated with microtubule dynamics and mitotic processes. Kinesin acts
as intracellular shuttle and can de-polymerize microtubules. It has been shown to bind to
negatively charged residues E410, D417 and E421 on the β-tubulin subunit.(60) MAP65/Ase1,
a microtubule-associated protein is reported to promote microtubule flexibility and prevent
microtubule severing by cross-linking.(50, 51) These findings suggest that the duplications
observed are associated to repairing and protecting microtubule functionality which is
intriguing considering the mode of action of albendazole, inhibiting microtubule formation.(61,

62) Further, highly duplicated genes including Condensin, Transcription factor (TF) IIIC and a haspin like kinase were associated to chromatid condensation, organization and cohesion respectively. While these genes have not been associated to benzimidazole resistance, they are linked to mitotic processes which in turn are connected to microtubule function with regards to the mitotic spindle.(63-65) *T. muris* is the only other organism with a high number of TBB orthologs and it has been shown that albendazole or mebendazole had an $IC_{50} > 200$ µg/mL in *T. muris*.(66) However, comparing the number of TBB orthologs of the population obtained after T1 to the population obtained after T2, showed no significant difference indicating that duplications are not the result of a recent adaptation nor an adaptation to drug pressure. Assuming an average mutation rate of 2×10^{-8} point mutations per gene per generation in *C. elegans* as an estimate, the duplication events would be dated back several centuries. These findings support the hypothesis that the observed lower cure rates in Côte d'Ivoire are not a result of resistance establishment, but potentially a result of different response rates of different *Trichuris* species. Finally, the functional significance of different TBB genes remains unknown and is subject for future studies.

Patterns of gene duplication, gain, and loss of function events in *T. incognita n. sp.* compared to within *T. suis* suggest a more ancient and divergent evolutionary lineage. Despite the close genetic relationship of *T. incognita n. sp.* to *T. suis*, our phylogenetic analysis demonstrates clear separation between *T. incognita n. sp.* and *T. suis* isolates from swine hosts in Côte d'Ivoire. While zoonosis events of trichuriasis from *T. suis* to *T. incognita n. sp.* have been reported,(67) these findings suggest that there may have been a zoonotic event sometime in the past between the human and swine host that was successful, but could not be observed currently. However, the occurrence of closely related *Trichuris* isolates from the colobus monkeys (*Colobus guereza kikuyensis*) raises the question, if these might belong to the same species of *Trichuris* and have zoonotic potential similar to *T. trichiura* species, which are found both within primates and humans.(32) Furthermore, J. Rivero *et al.* identified *Trichuris* species, which were closely related to the one found in the colobus monkey and may be the same species as *T. incognita n. sp.* based on ITS sequences or *cox*1 and *co*b genes.(43) If *T. incognita n. sp.* were able to infect non-human hosts, which is the case for *T. trichiura,* this would provide a refugia from drug pressure, which might prevent the establishment of anthelmintic resistance.(68)

With the unique experimental setup of the expulsion study, we were able to categorize helminths into drug sensitive and non-sensitive and thus enabling a genome wide association study (GWAS) to investigate the potential establishment of resistance to albendazole-ivermectin. MDA has been ongoing in Côte d'Ivoire for multiple decades,(4) which is well within the timeframe of a possible resistance establishment in STH through PC.(7) The most surprising finding of the GWAS may be, that there is no conclusive evidence to show a clear adaptation to drug treatment by comparing drug sensitive and non-sensitive *T. incognita* worms. Depending on the set of SNPs analyzed, regression model chosen for the GWAS, logistic vs linear, and the inclusion of principal components to account for population stratification or not, SNPs of genome wide significance with acceptable QQ-plots were only observed using a linear model in the male population without considering population stratification. Two hits are worth mentioning, as they can be found on genes that are to some extent in context of the mode of action of albendazole or ivermectin: a homolog of the muscle M-line assembly protein unc-89 found in *C. elegans* and a voltage gated chloride ion channel. The first one is a structural component of the muscle M line which regulates Ca^{2+} signaling and is involved in preventing the degradation of microtubule severing protein mei-1 by binding mel-26, which may be interesting regarding albendazoles mode of action.(69) The latter is interesting regarding the glutamate gated chloride ion channel binding activity of ivermectin,

leading to increased hyperpolarization in the neuromuscular junction. (70) However, while these hits will be subject of future investigation, there is no clear indication of resistance establishment in the analysis conducted. This is in contrast to reported quantitative trait loci (QTL) associated with ivermectin resistance in *H. contortus* and *C. elegans* found on the cky-1 gene,(14) and transmembrane proteins such as the chloride ion channel avr-15,(71) all of which did not show an association with genome wide significance in this study. Importantly, our experimental design also has limitations, such as the potential failure of capturing of multi-loci resistance mechanisms against a combination treatment, or gene expression related mechanisms, which might be visible in transcriptomic experiments.

The expulsion study also allowed us to gain insights into the fecundity and expulsion dynamics of *T. incognita*. Following the administration of albendazole and ivermectin during T1, only 271 out of 1098 worms were expelled, which is in-line with previously observed low cure rates in Côte d'Ivoire.(9, 72) The female to male ratio is comparable to recent reports for *T. trichiura*.(17) The fecundity of around 14,300 eggs per female per day of *T. incognita n. sp.* matches literature reported values of approximately 16,700 for *T. trichiura*.(17) Overall, these results suggest that the biology of *T. incognita n. sp.* in terms of fecundity is comparable to *T. trichiura*. Also morphologically, *T. incognita n. sp.* and *T. trichiura* were comparable. The presented morphological characteristics of *T. incognita n. sp.* do not enable a morphological distinction to *T. trichiura* indicating the limits of the morphological species concept for the *Trichuris* genus.(43) The morphological indistinguishability of *T. incognita n. sp.*, *T. trichiura* and *T. suis* worms may be the reason why this species has remained undescribed for many decades and emphasizes the need of diagnostic tools able to differentiate between these worms to gain further insight into the transmission dynamics and prevalence of this new species, as it responds differently to drug treatment. Intriguingly in a study from Rahman *et al.*, analyzing a sample cohort from 2019 in similar regions, both *T. trichiura* and *T. incognita* were detected, where in our study exclusively *T. incognita* was observed in 2022.(53) Longitudinal studies investigating a potential connection between species distribution and drug efficacies will be of particular interest.

This study demonstrates that human trichuriasis can be caused by multiple species of whipworm, and that the differences in response rates may be a result of species responding differently to drug treatment, as opposed to an establishment of resistance. While the current WHO guideline for treatment of *T. trichiura* (combination therapy with (ALB/IVM) should still stand the current findings show the necessity to incorporate molecular diagnostics into clinical trials and to develop a new treatment against *T. incognita n. sp.*

Acknowledgements

We would like to acknowledge Dr. John Gilleard and Dr. Abhinaya Venkatesan from the Faculty of Veterinary Medicine, Host-Parasite Interactions Program, University of Calgary, AB T2N1N4, Canada for the fruitful discussions on *Trichuris* phylogeny. We are grateful to the European Research Council for financial support (Nr. 101019223).

References:

1.   James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. The Lancet. 2018;392(10159):1789-858.

2.   Montresor A, Mupfasoni D, Mikhailov A, Mwinzi P, Lucianez A, Jamsheed M, et al. The
global progress of soil-transmitted helminthiases control in 2020 and World Health
Organization targets for 2030. PLoS Negl Trop Dis. 2020;14(8):e0008505.
3.   World Health O, Onchocerciasis Control Programme in the Volta River Basin A. Drug
development. Ouagadougou: Onchocerciasis Control Programme in the Volta River Basin
Area; 1976 1976. Contract No.: OCP/STAC/4.2.
4.   Bland J, Viedma C, Davies P, Ozorio P. This wormy world [full issue]. World health.
1984(March 1984):2-31.
5.   Ogoussan KT, Hopkins A. Mectizan((R)) procurement and delivery for onchocerciasis
mass drug administration programmes. Acta Trop. 2011;120 Suppl 1:S173-6.
6.   Rose Vineer H, Morgan ER, Hertzberg H, Bartley DJ, Bosco A, Charlier J, et al.
Increasing importance of anthelmintic resistance in European livestock: creation and meta-
analysis of an open database. Parasite. 2020;27:69.
7.   Coffeng LE, Stolk WA, de Vlas SJ. Predicting the risk and speed of drug resistance
emerging in soil-transmitted helminths during preventive chemotherapy. Nat Commun.
2024;15(1):1099.
8.   Hansen TV, Nejsum P, Friis C, Olsen A, Thamsborg SM. Trichuris suis and
Oesophagostomum dentatum show different sensitivity and accumulation of fenbendazole,
albendazole and levamisole in vitro. PLoS Negl Trop Dis. 2014;8(4):e2752.
9.   Hürlimann E, Keller L, Patel C, Welsche S, Hattendorf J, Ali SM, et al. Efficacy and
safety of co-administered ivermectin and albendazole in school-aged children and adults
infected with Trichuris trichiura in Côte d'Ivoire, Laos, and Pemba Island, Tanzania: a double-
blind, parallel-group, phase 3, randomised controlled trial. The Lancet Infectious Diseases.
2022;22(1):123-35.
10.   Venkatesan A, Chen R, Bar M, Schneeberger PHH, Reimer B, Hurlimann E, et al.
Trichuriasis in Human Patients from Cote d'Ivoire Caused by Novel Trichuris incognita Species
with Low Sensitivity to Albendazole/Ivermectin Combination Treatment. Emerg Infect Dis.
2025;31(1):104-14.
11.   Nissen S, Al-Jubury A, Hansen TV, Olsen A, Christensen H, Thamsborg SM, et al.
Genetic analysis of Trichuris suis and Trichuris trichiura recovered from humans and pigs in a
sympatric setting in Uganda. Vet Parasitol. 2012;188(1-2):68-77.
12.   Ghedin E. Panning for molecular gold in whipworm genomes. Nat Genet.
2014;46(7):661-3.
13.   Doyle SR. Improving helminth genome resources in the post-genomic era. Trends
Parasitol. 2022;38(10):831-40.
14.   Doyle SR, Laing R, Bartley D, Morrison A, Holroyd N, Maitland K, et al. Genomic
landscape of drug response reveals novel mediators of anthelmintic resistance. Preprint.
2022.
15.   Jex AR, Nejsum P, Schwarz EM, Hu L, Young ND, Hall RS, et al. Genome and
transcriptome of the porcine whipworm Trichuris suis. Nat Genet. 2014;46(7):701-6.
16.   Foth BJ, Tsai IJ, Reid AJ, Bancroft AJ, Nichol S, Tracey A, et al. Whipworm genome
and dual-species transcriptome analyses provide molecular insights into an intimate host-
parasite interaction. Nat Genet. 2014;46(7):693-700.
17.   Hansen EP, Tejedor AM, Thamsborg SM, Alstrup Hansen TV, Dahlerup JF, Nejsum
P. Faecal egg counts and expulsion dynamics of the whipworm, Trichuris trichiura following
self-infection. J Helminthol. 2016;90(3):298-302.
18.   Bundy DA, Thompson DE, Cooper ES, Blanchard J. Rate of expulsion of Trichuris
trichiura with multiple and single dose regimens of albendazole. Trans R Soc Trop Med Hyg.
1985;79(5):641-4.
19.   Cringoli G, Maurelli MP, Levecke B, Bosco A, Vercruysse J, Utzinger J, et al. The Mini-
FLOTAC technique for the diagnosis of helminth and protozoan infections in humans and
animals. Nat Protoc. 2017;12(9):1723-32.
20.   Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow
enables reproducible computational workflows. Nat Biotechnol. 2017;35(4):316-9.

21. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30(15):2114-20.
22. Jin JJ, Yu WB, Yang JB, Song Y, dePamphilis CW, Yi TS, et al. GetOrganelle: a fast
and versatile toolkit for accurate de novo assembly of organelle genomes. Genome Biol.
2020;21(1):241.
23. Meng G, Li Y, Yang C, Liu S. MitoZ: a toolkit for animal mitochondrial genome
assembly, annotation and visualization. Nucleic Acids Res. 2019;47(11):e63.
24. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80.
25. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchene S, Fourment M, Gavryushkina
A, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLoS
Comput Biol. 2019;15(4):e1006650.
26. De Coster W, Rademakers R. NanoPack2: population-scale evaluation of long-read
sequencing data. Bioinformatics. 2023;39(5).
27. Sun J, Li R, Chen C, Sigwart JD, Kocot KM. Benchmarking Oxford Nanopore read
assemblers for high-quality molluscan genomes. Philos Trans R Soc Lond B Biol Sci.
2021;376(1825):20200160.
28. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome
Biol. 2019;20(1):257.
29. Zimin AV, Marcais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA
genome assembler. Bioinformatics. 2013;29(21):2669-77.
30. Alonge M, Lebeigle L, Kirsche M, Jenike K, Ou S, Aganezov S, et al. Automated
assembly scaffolding using RagTag elevates a new tomato system for high-throughput
genome editing. Genome Biol. 2022;23(1):258.
31. Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations.
Bioinformatics. 2021;37(12):1639-43.
32. Doyle SR, Søe MJ, Nejsum P, Betson M, Cooper PJ, Peng L, et al. Population
genomics of ancient and modern Trichuris trichiura. Preprint. 2021.
33. Manni M, Berkeley MR, Seppey M, Simao FA, Zdobnov EM. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for
Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021;38(10):4647-54.
34. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome
assembly evaluation with QUAST-LG. Bioinformatics. 2018;34(13):i142-i50.
35. Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, et al.
OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. Nucleic
Acids Res. 2023;51(D1):D445-D51.
36. Bruna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein
database. NAR Genom Bioinform. 2021;3(1):lqaa108.
37. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019;20(1):238.
38. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res. 2021;49(W1):W293-W6.
39. Altenhoff AM, Levy J, Zarowiecki M, Tomiczek B, Warwick Vesztrocy A, Dalquen DA,
et al. OMA standalone: orthology inference among public and custom genomes and
transcriptomes. Genome Res. 2019;29(7):1152-63.
40. Train CM, Pignatelli M, Altenhoff A, Dessimoz C. iHam and pyHam: visualizing and
processing hierarchical orthologous groups. Bioinformatics. 2019;35(14):2504-6.
41. Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler
transform. Bioinformatics. 2010;26(5):589-95.
42. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a
tool set for whole-genome association and population-based linkage analyses. Am J Hum
Genet. 2007;81(3):559-75.

43.   Rivero J, Cutillas C, Callejon R. Trichuris trichiura (Linnaeus, 1771) From Human and
Non-human Primates: Morphology, Biometry, Host Specificity, Molecular Characterization,
and Phylogeny. Front Vet Sci. 2020;7:626120.
44.   Yoshikawa H, Yamada M, Matsumoto Y, Yoshida Y. Variations in egg size ofTrichuris
trichiura. Parasitology Research. 1989;75(8):649-54.
45.   Tran MH, Pearson MS, Bethony JM, Smyth DJ, Jones MK, Duke M, et al. Tetraspanins
on the surface of Schistosoma mansoni are protective antigens against schistosomiasis. Nat
Med. 2006;12(7):835-40.
46.   Susa KJ, Kruse AC, Blacklow SC. Tetraspanins: structure, dynamics, and principles of
partner-protein recognition. Trends Cell Biol. 2023.
47.   Emms DM, Kelly S. STAG: Species Tree Inference from All Genes. 2018.
48.   Emms DM, Kelly S. STRIDE: Species Tree Root Inference from Gene Duplication
Events. Molecular Biology and Evolution. 2017;34(12):3267-78.
49.   Laing R, Doyle SR, McIntyre J, Maitland K, Morrison A, Bartley DJ, et al.
Transcriptomic analyses implicate neuronal plasticity and chloride homeostasis in ivermectin
resistance and response to treatment in a parasitic nematode. PLoS Pathog.
2022;18(6):e1010545.
50.   Burkart GM, Dixit R. Microtubule bundling by MAP65-1 protects against severing by
inhibiting the binding of katanin. Mol Biol Cell. 2019;30(13):1587-97.
51.   Portran D, Zoccoler M, Gaillard J, Stoppin-Mellet V, Neumann E, Arnal I, et al.
MAP65/Ase1 promote microtubule flexibility. Mol Biol Cell. 2013;24(12):1964-73.
52.   Venkatesan A, Chen R, Baer M, Schneeberger PHH, Reimer B, Hürlimann E, et al.
2024.
53.   Rahman N, Bär MA, Dommann J, Hürlimann E, Coulibaly JT, Ali S, et al. Nanopore-
based analysis unravels the genetic landscape and phylogenetic placement of human-
infecting Trichuris species in Côte d'Ivoire, Tanzania, Uganda, and Laos.
bioRxiv. 2024:2024.07.31.605962.
54.   Mkaouar H, Mariaule V, Rhimi S, Hernandez J, Kriaa A, Jablaoui A, et al. Gut
Serpinome: Emerging Evidence in IBD. Int J Mol Sci. 2021;22(11).
55.   Xu J, Bai X, Wang LB, Shi HN, van der Giessen JWB, Boireau P, et al. Influence of
adjuvant formulation on inducing immune response in mice immunized with a recombinant
serpin from Trichinella spiralis. Parasite Immunol. 2017;39(7).
56.   Lynch M, Conery JS. The Evolutionary Fate and Consequences of Duplicate Genes.
Science. 2000;290(5494):1151-5.
57.   Kuzmin E, Taylor JS, Boone C. Retention of duplicated genes in evolution. Trends
Genet. 2022;38(1):59-72.
58.   Kondrashov FA. Gene duplication as a mechanism of genomic adaptation to a
changing environment. Proc Biol Sci. 2012;279(1749):5048-57.
59.   Zajac N, Zoller S, Seppala K, Moi D, Dessimoz C, Jokela J, et al. Gene Duplication
and Gain in the Trematode Atriophallophorus winterbourni Contributes to Adaptation to
Parasitism. Genome Biol Evol. 2021;13(3).
60.   Cushion TD, Leca I, Keays DA. MAPping tubulin mutations. Front Cell Dev Biol.
2023;11:1136699.
61.   Enos A, Coles GC. Effect of benzimidazole drugs on tubulin in benzimidazole resistant
and susceptible strains of Caenorhabditis elegans. Int J Parasitol. 1990;20(2):161-7.
62.   Dawson PJ, Gutteridge WE, Gull K. A comparison of the interaction of anthelmintic
benzimidazoles with tubulin isolated from mammalian tissue and the parasitic nematode
Ascaridia galli. Biochem Pharmacol. 1984;33(7):1069-74.
63.   Hagstrom KA, Meyer BJ. Condensin and cohesin: more than chromosome compactor
and glue. Nat Rev Genet. 2003;4(7):520-34.
64.   Stutzman AV, Liang AS, Beilinson V, Ikegami K. Transcription-independent TFIIIC-
bound sites cluster near heterochromatin boundaries within lamina-associated domains in C.
elegans. Epigenetics Chromatin. 2020;13(1):1.
65.   Dai J, Kateneva AV, Higgins JM. Studies of haspin-depleted cells reveal that spindle-
pole integrity in mitosis requires chromosome cohesion. J Cell Sci. 2009;122(Pt 22):4168-76.

66.   Keiser J, Tritten L, Adelfio R, Vargas M. Effect of combinations of marketed human anthelmintic drugs against Trichuris muris in vitro and in vivo. Parasites & Vectors. 2012;5(1):292.

67.   Nejsum P, Betson M, Bendall RP, Thamsborg SM, Stothard JR. Assessing the zoonotic potential of Ascaris suum and Trichuris suis: looking to the future from an analysis of the past. J Helminthol. 2012;86(2):148-55.

68.   Vercruysse J, Albonico M, Behnke JM, Kotze AC, Prichard RK, McCarthy JS, et al. Is anthelmintic resistance a concern for the control of human soil-transmitted helminths? Int J Parasitol Drugs Drug Resist. 2011;1(1):14-27.

69.   Wilson KJ, Qadota H, Mains PE, Benian GM. UNC-89 (obscurin) binds to MEL-26, a BTB-domain protein, and affects the function of MEI-1 (katanin) in striated muscle of Caenorhabditis elegans. Mol Biol Cell. 2012;23(14):2623-34.

70.   El-Saber Batiha G, Alqahtani A, Ilesanmi OB, Saati AA, El-Mleeh A, Hetta HF, et al. Avermectin Derivatives, Pharmacokinetics, Therapeutic and Toxic Dosages, Mechanism of Action, and Their Biological Effects. Pharmaceuticals (Basel). 2020;13(8).

71.   Dent JA, Smith MM, Vassilatis DK, Avery L. The genetics of ivermectin resistance in Caenorhabditis elegans. Proceedings of the National Academy of Sciences. 2000;97(6):2674-9.

72.   Sprecher VP, Coulibaly JT, Hurlimann E, Hattendorf J, Keiser J. Efficacy and Safety of Moxidectin-Albendazole and Ivermectin-Albendazole Combination Therapy Compared to Albendazole Monotherapy in Adolescents and Adults Infected with Trichuris trichiura: A Randomized, Controlled Superiority Trial. Clin Infect Dis. 2023;77(9):1294-302.

Supporting Information

900   **Figures**

902   SI Figure 1: Correlation between observed expelled worms and average egg count per day.
903   Error bars indicate standard error with no error bar present if there was only one value.

SI Figure 3: Phylogenetic tree based on mitochondrial sequences from 535 isolated *T.*
incognita n. sp. worms from human hosts. The circle on branches indicates the posterior
distribution. The outmost 4 rings represent binary metadata. A filled red tile indicates that the
worm was isolated after T2 where the empty tile indicates isolation after T1. The dark violet
tile indicates the worm sex and the light violet tile the host sex while purple indicates if a
coinfection with other worms was observed. The middle ring indicates two hosts: A host with
the ID number "531 – 1250" in light brown and a host with the ID number "498 – 1173" in dark
brown, showing that the worms isolated from the two different hosts are scattered throughout
the phylogenetic tree. The inner circle indicates the village from which the worm was isolated.

SI Figure 4: Tetraspanin, Serpin and Chymotrypsin-like serine protease from analyzing orthologous groups in Venn diagram sections shared by *T. incognita n. sp.*, *T. trichiura* (Serpin and Chymotrypsin-like serine protease) and *T. muris* (Tetraspanin).

SI Figure 5: Beta tubulin genes found in *T. incognita n. sp.* compared to *T. suis* and *T. trichiura*, predicted from the braker2 pipeline. Black indicates positions carrying known mutations (Phe168, Glu198, Phe200), observed variable position (Ile194). A truncation was predicted *in silico* on 3 of the 7 genes. Gene IDs from top to bottom: g34, g460, g463, g472, g475, g529 and g7506.

SI Figure 6: Coverage and variants from an independent PCR and Nanopore genome sequencing experiment of the regions containing known mutations of TBB and the *in silico* predicted truncation. Blue line indicates coverage, red lines the called variants using bcftools. Gray indicating positions carrying known mutations (Phe168, Glu198, Phe200), observed

variable position (Ile194) and predicted truncation indicating that the truncation was not
observed from the amplified genomic sequence.

Forward primer sequence: AAAGAGACCGGACATTTCGC.

Reversed primer sequence: TGAATTGCCTGGTTTCTAGAATG.

SI Figure 7: Summary of the PCA of all missense mutations on coding genes for the GWAS
showing no population structure associated to treatment. a-c PCA plots of all missense
mutations showing that worm sex has the biggest impact on population stratification. d-i PCA
plots of only male worms showed significant population stratification by Tukey multiple
comparisons of means in PC1 between the villages Tiagba-Akakro (p=4.96 E-14) and Bekpou-

 Akakro (p=2.79 E-05) and in PC2 between the villages Tiagba-Akakro (p=0.0032). No
association to the treatment (ANOVA p > 0.05) was found in PC1 or PC2. j-o PCA plots of
only female worms showed significant population stratification by Tukey multiple comparisons
of means in PC1 between the villages Tiagba-Ahouya (p=0.0009) and Ahouya-Akakro
(p=0.0006) and in PC2 between the villages Ahouya-Akakro (p=0.027). No association to the
treatment (ANOVA p > 0.05) was found in PC1 or PC2.

SI Figure 8: GWAS displaying p-values of missense SNPs on predicted genes associated to
treatment outcome (T1 vs T1) using a linear model without population stratification in PLINK.
Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of
0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001.
The red line indicates the Bonferroni corrected significance of 8.89E-7 at α=0.05 and 56273
mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs
found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data
of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive
and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-
sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug
non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast
hit, the location and functional annotation, where possible, is provided in the SI Table 7
"hits_gwas_all.xlsx"

SI Figure 9: GWAS displaying p-values of missense SNPs on predicted genes associated to treatment outcome (T1 vs T1) using a linear model with population stratification in PLINK. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the Bonferroni corrected significance of 8.89E-7 at α=0.05 and 56273 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast hit, the location and functional annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

SI Figure 10: GWAS displaying p-values of missense SNPs on predicted genes associated to
treatment outcome (T1 vs T1) using a logistic model without population stratification in PLINK.
Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of
0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001.
The red line indicates the Bonferroni corrected significance of 8.89E-7 at α=0.05 and 56273
mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs
found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data
of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive
and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-
sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug
non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast
hit, the location and functional annotation, where possible, is provided in the SI Table 7
"hits_gwas_all.xlsx"

SI Figure 11: GWAS displaying p-values of missense SNPs on predicted genes associated to
treatment outcome (T1 vs T1) using a logistic model with population stratification in PLINK.
Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of
0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001.
The red line indicates the Bonferroni corrected significance of 8.89E-7 at α=0.05 and 56273
mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs
found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data
of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive
and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-
sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug
non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast
hit, the location and functional annotation, where possible, is provided in the SI Table 7
"hits_gwas_all.xlsx"

1012

1013

SI Figure 12: GWAS displaying p-values of all SNPs associated to treatment outcome (T1 vs T1) using a linear model without population stratification in PLINK. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the Bonferroni corrected significance of 3.99 E-8 at α=0.05 and 1254247 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast hit, the location and functional annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

1026

SI Figure 13: GWAS displaying p-values of all SNPs associated to treatment outcome (T1 vs T1) using a linear model with population stratification in PLINK. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the Bonferroni corrected significance of 3.99 E-8 at α=0.05 and 1254247 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138

controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast hit, the location and functional annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

SI Figure 14: GWAS displaying p-values of all SNPs associated to treatment outcome (T1 vs T1) using a logistic model without population stratification in PLINK. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the Bonferroni corrected significance of 3.99 E-8 at α=0.05 and 1254247 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **b** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138 controls. **c** Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast hit, the location and functional annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

SI Figure 15: GWAS displaying p-values of all SNPs associated to treatment outcome (T1 vs T1) using a logistic model with population stratification in PLINK. Variants were filtered at a minor allele frequency of 0.05, a missing genotype frequency of 0.05, and a deviation from the Hardy-Weinberg Equilibrium at a significance level of p=0.001. The red line indicates the

Bonferroni corrected significance of 3.99 E-8 at α=0.05 and 1254247 mutations. The scaffolds are separated by black/grey colors. The green dots indicate all SNPs found on the specific gene of interest carrying a SNP with genome wide significance. **a** and **b** Data of both sexes. 247 males, 471 females, and 3 ambiguous 542 cases as in drug non-sensitive and 179 controls. **c** and **d** Only female SNPs. 0 males, 471 females, 333 cases as in drug non-sensitive and 138 controls. **e** and **f** Only male SNPs. 247 males, 0 females, 208 cases as in drug non-sensitive and 39 controls. All GWAS hits, from each analysis, the sequence, the best blast hit, the location and functional annotation, where possible, is provided in the SI Table 7 "hits_gwas_all.xlsx"

Tables

Sample ID	Sex	Total length	Posterior length	Posterior width	Anterior length	Anterior width
995	F	36	12	1.2	24	0.09
787	F	48	15	0.8	33	0.15
66	F	44	13	0.6	31	0.12
618	F	45	11	0.7	34	0.15
368	F	40	12	0.6	28	0.11
Mean		**42.39**	**12.53**	**0.75**	**29.77**	**0.12**
SD		**4.67**	**1.52**	**0.25**	**4.06**	**0.03**
Sample ID	**Sex**					
9	M	36	11	1	25	0.2
965	M	40	17	1.1	23	0.25
960	M	41	14	0.5	27	0.15
271	M	38	13	0.6	25	0.07
17	M	41	18	0.5	23	0.1
Mean		**39.15**	**14.37**	**0.70**	**24.55**	**0.14**
SD		**2.17**	**2.88**	**0.29**	**1.67**	**0.07**

Females	Total length (Min-Max)	Posterior length (Min-Max)	Posterior width (Min-Max)	Anterior length (Min-Max)	Anterior width (Min-Max)
T. incognita n. sp. (Present study)	36-45	11-15	0.6-1.2	24-34	0.09-0.15
T. trichiura from J. Rivero *et al.*	20-49	6-16	0.38-0.9	13-33	0.09-0.2
Males	**Total length (Min-Max)**	**Posterior length (Min-Max)**	**Posterior width (Min-Max)**	**Anterior length (Min-Max)**	**Anterior width (Min-Max)**
T. incognita n. sp. (Present study)	36-41	11-15	0.5-1	23-27	0.07-0.25
T. trichiura from J. Rivero *et al.*	25-49	7-20	0.4-0.7	18-29	0.09-0.31

SI Table 1: Five randomly sampled male and female worms (top) and literature reported values (bottom). Total length, posterior length, posterior width, anterior length and anterior width are provided, including the geometrical mean and standard deviation (top) or Min-Max values (bottom).

Venn diagram

Category	GO term	p-value	Description
Cellular component	GO:0005615	3.10E-21	extracellular space
	GO:0000786	4.02E-17	nucleosome
Molecular function	GO:0003676	1.29E-52	nucleic acid binding
	GO:0004830	1.21E-33	tryptophan-tRNA ligase activity
	GO:0030527	2.78E-18	structural constituent of chromatin
	GO:0003723	6.72E-18	RNA binding
Biological process	GO:0006436	1.21E-33	tryptophanyl-tRNA aminoacylation
	GO:0044571	3.17E-19	[2Fe-2S] cluster assembly
	GO:0051259	3.17E-19	protein complex oligomerization

1076

SI Table 2. Venn diagram of orthologous groups from predicted transcripts, where 175 were
exclusively shared between T. incognita n. sp. and T. trichiura and their GO enrichment
analysis. Availability of a *Trichuris* genome, closely related to *T. suis* but from a species
naturally infecting humans, poses a unique opportunity to conduct an exploratory analysis of
genes that might be crucial for the adaptation to a human host. We looked into the occurrence
of orthologous groups throughout the genomes of *T. incognita n. sp.*, *T. suis*, *T. trichiura* and
T. muris as shown in Table 2. Leveraging the close genetic relationship with *T. suis*, predicted
transcripts were investigated that are shared with *T. trichiura* but do not occur in *T. suis* or *T.*
muris. A GO term enrichment analysis was performed on this subset of genes using the
predicted genes in *T. incognita n. sp.* as a background gene set and the full list of significant
terms is provided in SI Table 3. A list of GO enriched terms provided in Table 2 indicate
extracellular space to be the most significantly enriched term in the category of cellular
components along with the nucleosom. Nucleic acid binding, tryptophan-tRNA ligase activity,
structural constituent of chromatin and RNA binding were the most significant in molecular
functions and finally tryptophanyl-tRNA aminoacylation, [2Fe-2S] cluster assembly and protein
complex oligomerization in biological process. Finally, considering that the anterior section is
in a most intimate relationship with the host, we searched for genes within this gene-set which
Jex *et. al* identified to be upregulated in the stichosome and potentially have
immunomodulatory effects like galactins, serpins, venom allergen–like proteins, apyrase or
calreticulin and chymotrypsin-like serine proteases.(15) We identified one family of each
serpins (OG0000305) and chymotrypsin-like serine proteases (OG0010319) that are shared
only by *T. trichiura* and *T. incognita n. sp.* (SI Figure 4). Additionally, we identified, a family of
tetraspanins (OG0009162) which are investigated as vaccine candidates in
schistosomiasis.(45, 46)

"Gene_list_venn_intersection.xlsx"

SI Table 3. Gene list of cross section in Venn diagram.

"Gene_list_resistance_associated_literature.xlsx"

SI Table 4. Gene list of resistance associated genes towards albendazole or ivermectin in
helminths.

"Orthologous_group_list_duplications.xlsx"

SI Table 5. GO term enrichment analysis results

"GO-Term_enrichment_analysis.xlsx"

SI Table 6. Highly duplicated genes.

"hits_gwas_all.xlsx"

SI Table 7

"Main_Phenotype_File_sex.txt"

SI Table 8. Worm ID's with sex for PLINK input and phenotype of drug sensitive or non
sensitive: Sex code ('1' = male, '2' = female, '0' = unknown), Phenotype value ('1' = control, '2'
= case, '-9'/'0'/non-numeric = missing data if case/control)

1116