A snapshot of the Physcomitrella N-terminome reveals N-terminal methylation of organellar proteins

Sebastian N.W. Hoernstein[1], Andreas Schlosser[2], Kathrin Fiedler[3,*], Gabor L. Igloi[3], Daniel Lang[1,4], Ralf Reski[1,5]

[1] Plant Biotechnology, Faculty of Biology, University of Freiburg, Schaenzlestr. 1, 79104 Freiburg, Germany

[2] Rudolf Virchow Center for Experimental Biomedicine, University of Würzburg, Josef-Schneider-Str. 2, 97080 Würzburg, Germany

[3] Institute of Biology III, University of Freiburg, Schaenzlestr. 1, 79104 Freiburg, Germany

[4] Bundeswehr Institute of Microbiology, Microbial Genomics and Bioforensics, Neuherbergstr. 11, 80937 Munich, Germany

[5] Signalling Research Centres BIOSS and CIBSS, Schaenzlestr., 79104 Freiburg, Germany

* Present address: Lonza, Hochbergerstr. 60A, 4057 Basel, Switzerland

Corresponding author: ralf.reski@biologie.uni-freiburg.de

ORCID IDs

Sebastian Hoernstein	https://orcid.org/0000-0002-2095-689X
Andreas Schlosser	https://orcid.org/0000-0003-0612-9932
Kathrin Fiedler	https://orcid.org/0009-0006-6265-0836
Gabor Igloi	https://orcid.org/0000-0002-4490-8892
Daniel Lang	https://orcid.org/0000-0002-2166-0716
Ralf Reski	https://orcid.org/0000-0002-5496-6711

Abstract

Post- or co-translational N-terminal modifications of proteins influence their half-life as well as mediating protein sorting to organelles *via* cleavable N-terminal sequences that are recognized by the respective translocation machinery. Here, we provide an overview on the current modification state of the N-termini of over 4500 proteins from the model moss Physcomitrella (*Physcomitrium patens*) using a compilation of 24 N-terminomics datasets. Our data reveal distinct proteoforms and modification states and confirm predicted targeting peptide cleavage sites of 1144 proteins localized to plastids and the thylakoid lumen, to mitochondria, and to the secretory pathway. Additionally, we uncover extended N-terminal methylation of mitochondrial proteins. Moreover, we identified PpNTM1 (*P. patens* alpha N-terminal protein methyltransferase 1) as a promising candidate for protein methylation in plastids, mitochondria and the cytosol. These data can now be used to optimize computational targeting predictors, for customized protein fusions and their targeted localization in biotechnology, and they provide novel insights into potential dual targeting of proteins.

Key message:

Analysis of the N-terminome of Physcomitrella reveals N-terminal monomethylation of nuclear encoded, mitochondria-localized proteins.

Key words:

AARE, N-terminus, organelle, proteases, protein methylation, protein modification, protein targeting

Introduction

Following translation at the ribosome, the N-terminus of a protein is subjected to a plethora of modifications among which are proteolytic processing and the addition of moieties such as acetyl, methyl or other functional groups (Fortelny et al. 2015; Meinnel and Giglione 2008). In turn, the N-terminus enables subcellular targeting and is a determinant of protein half-life (Varshavsky 1996; Kunze and Berger 2015; Armenteros et al. 2019; Varshavsky 2019). Modifications of the N-terminus are introduced in a co- or a posttranslational manner with co-translational acetylation and methionine-excision being among the most abundant modifications in eukaryotes (Ree et al. 2018; Giglione and Meinnel 2021). In plants, however, N-terminal acetylation occurs also in a posttranslational manner on plastid stromal proteins after import and cleavage of their targeting peptide (Giglione and Meinnel 2021). In contrast to proteolytic trimming, amino acids can also be added to the apparent N-terminus of a protein in a ribosome-independent manner (Varshavsky 1996; Tasaki 2012) as part of the N-degron pathway for targeted proteolysis. Various methods such as COFRADIC (Staes et al. 2011), TAILS (Kleifeld et al. 2010) or HUNTER (Demir et al. 2022) have been established and permit the characterization of proteases, high-throughput degradomics and profiling of N-terminal acetylation. In turn, N-terminomics data are available from public databases such as TopFIND (https://topfind.clip.msl.ubc.ca/) for various organisms including human, mouse, yeast, and Arabidopsis.

In contrast, almost no N-terminomics data were available for the model plant Physcomitrella (*Physcomitrium patens*; Lueth and Reski, 2023). This moss is a versatile model system for evo-devo studies (Horst et al. 2016), plant physiology (Decker et al. 2017; Wiedemann et al. 2018) and evolution of metabolic pathways (Renault et al. 2017) due to its interesting evolutionary position at the early divergence of land plants (Rensing et al. 2008). It has further proven to be a valuable system for proteomics and proteogenomics research due to its easy and axenic culture conditions enabling highly reproducible and even GMP-compliant culture conditions (Sarnighausen et al. 2004; Heintz et al. 2006; Mueller et al. 2014; Hoernstein et al. 2016, 2018; Fesenko et al. 2019, 2021). Besides broad application in basic research, Physcomitrella is employed as a production platform for recombinant biopharmaceuticals in GMP-compliant bioreactors (Decker and Reski 2020; Ruiz-Molina et al. 2021; Tschongov et al. 2024). Furthermore, genomic and transcriptomic resources are well established and publicly

available (Goodstein et al. 2012; Lang et al. 2018; Perroud et al. 2018; Fernandez-Pozo et al. 2020; Bi et al. 2024).

Here, we provide a first snapshot of the N-terminome of the moss Physcomitrella with a focus on the cleavage of N-terminal targeting sequences, N-terminal acetylation and N-terminal monomethylation. The data was compiled using 24 datasets from various experimental setups and subsequent N-terminal peptide enrichment using a modified TAILS approach. We reveal apparent N-terminal methylation not only of plastid and cytosolic proteins but also of mitochondrial proteins. Further, we provide a list of confirmed targeting peptide cleavage sites along with a candidate list of proteins which are dually targeted to plastids and mitochondria as well as to mitochondria and the cytosol. With this, we provide a resource for basic research as it contains information about translation of splice variants as well as posttranslational and posttranscriptional processing of proteins. Moreover, targeting of recombinant proteins to plastids of *Nicotiana benthamiana* (MacLean et al. 2007) or the extracellular space in Physcomitrella (Schaaf et al. 2005) enabled high yields of the desired recombinant product. Consequently, our present data provide also a comprehensive resource for further tailored recombinant protein production and targeting in Physcomitrella.

Results and Discussion

Overview of identified N-termini

The present data provide a qualitative overview of the N-terminome of the moss Physcomitrella using a compilation of 24 datasets from N-terminal peptide enrichments from different tissues, treatments and different sample processing protocols. The datasets were obtained during method establishment for various purposes not related to the analysis performed in this study and hence the data was only assessed qualitatively and will not allow any cross-sample comparison. A table providing details about the sample type, employed tissue and other experimental parameters is available from Supplemental Table S1.

Enrichment of N-terminal peptides was performed as described in Hoernstein et al. (2018) with modifications. Free amino groups in the protein sample were blocked by reductive dimethylation according to Kleifeld et al. (2010) and depletion of internal peptides after proteolysis was performed according to McDonald and Beynon (2006). MS measurements

were performed on an LTQ-Orbitrap Velos Pro (ThermoScientific) and raw data were processed and searched with *Mascot* (Matrix Science). All database search results were loaded in *Scaffold5*[TM] (V5.0.1, https://www.proteomesoftware.com/) software and proteins were accepted with a *ProteinProphet*[TM] (Nesvizhskii et al. 2003) probability of at least 99% and a minimum of 1 identified peptide. Peptides were accepted at a *PeptideProphet*[TM] (Keller et al. 2002) probability of at least 95% and a *Mascot* ion score of a least 40 (Supplemental Table S2). Using these settings, from a total of 24 datasets we identified 11,533 protein N-termini using 32,213 spectra corresponding to 4517 proteins (3920 protein groups) with a decoy FDR (false discovery rate) of 0.4% at the protein level and 0.08% at the peptide level (Supplemental Tables S2, S3). Approximately 20% of the identified N-termini represented either the initiator methionine (start index 1, Figure 1A) or the subsequent amino acid after cleavage of the initiator methionine (start index 2). For approx. 40% a start index between 2 and 100 was identified, indicating proteolytic processing and cleavage of subcellular targeting sequences. A single experimentally determined N-terminus was observed in approx. 70% of all cases whereas for approx. 30% of the identified proteins two or more N-termini were observed (Figure 1B). This is in strong contrast to previous findings from proteins in Physcomitrella bioreactor supernatants (Hoernstein et al. 2018) where at least 2 distinct N-termini were observed for approx. 80% of all identified proteins. Further, we analyzed the presence of N-terminal modifications with a focus on N-terminal acetylation, mono-methylation and presence of pyro-glutamate (pyroGlu) at the N-terminus. The latter modification can occur spontaneously or *via* enzymatic catalysis on N-terminal glutamine residues (Schilling et al. 2008). Since pyro-glutamate formation can also occur following proteolysis during sample processing (Purwaha et al. 2014), only peptides where the preceding P1 amino acid did not match the specificity of the employed protease (e.g., no peptides with K or R as preceding amino acid in the case of trypsin digests) were considered here.

Approximately 25% of all identified N-termini were acetylated, 71% were identified without any modification and 4% were identified being either methylated or having N-terminal pyroGlu (Figure 1C). The actual level of N-terminal pyroGlu occurrence is likely higher, but due to specificity ambiguity with the experimentally employed proteases this cannot be further analyzed. At the protein level, we found approximately 76% of the nuclear-encoded proteins having either the retained or the cleaved initiator methionine (1436 protein groups, Supplemental Table S3) to be N-terminally acetylated (1097 protein groups, Supplemental

Table S3). This degree is slightly below the estimated degree of around 90% of N-terminal acetylated proteins in plants (Bienvenut et al. 2012; Linster and Wirtz 2018).

Figure 1 Overview of identified N-termini, N-terminal modifications and identified cleavage sites of targeting peptides. (**A**) Frequency of identified N-terminal positions per identified protein accession. The start index represents the position number of the identified N-terminal amino acid in the corresponding protein model. (**B**) Frequency of the number of identified N-termini per protein. (**C**) Bar chart depicting the distribution of identified N-terminal modifications. Peptides bearing N-terminal pyro-glutamate (pyroGlu) were only counted if the preceding amino acid did not match the specificity of the applied protease (e.g., peptides identified with K|R in the P1 position were rejected in the case of trypsin digests).

Post-import trimming of plastid proteins

Cleavable N-terminal sequences are required for sub- and extracellular targeting of nuclear-encoded proteins. Their cleavage *via* specific proteases after translocation across a respective organellar membrane generates a new N-terminus that represents either the final N-terminus of the translocated protein or a new site for further proteolytic processing by organellar proteases. For Physcomitrella, a total of 8681 cleavable N-terminal targeting sequences are predicted (Supplemental Figure S1) and here we compared our experimentally observed N-termini to these predictions allowing a tolerance window of ±5 amino acids around a predicted targeting peptide cleavage site (Figure 2A). In the following, a difference of 0 indicates agreement of an observed N-terminal amino acid with a predicted cleavage site (predicted P1 amino acid). Within this range we confirm the predicted cleavage sites of 748 plastid targeting signals (cTP), of 57 thylakoid luminal targeting signals (luTP), of 154 mitochondrial presequences (mTP), and of 185 secretory signal peptides (SP) using our present N-terminomics data. This data is compiled in Supplemental Table S4. Among the confirmed

plastid proteins, we find approx. 42% to be N-terminally acetylated (317 protein isoforms, Supplemental Table S4). Apparently, N-termini identified around a plastid transit peptide cleavage site only matched in approx. 47% the predicted cleavage site exactly (Figure 2A). This percentage is strikingly higher in all other cases with almost 70% for thylakoid luminal transit peptides (Figure 2B) and around 65% for mitochondrial pre-sequences and secretory signal peptides (Figures 2C, D). Further, approx. 43% of the N-termini of plastid proteins within the chosen difference window deviate by 1-5 amino acids upstream of the predicted cleavage site with decreasing frequency. This effect is less apparent for luminal targeting sequences (approx. 33%), mitochondrial pre-sequences (approx. 22%) and secretory signal peptides (approx. 32%). Consequently, the distribution of differences between predicted and observed plastid transit cleavage site (Figure 2A) indicates a successive proteolytic post-processing pattern of plastid proteins after cleavage of their transit peptide. A similar scenario with multiple cleavage sites around predicted plastid transit peptide cleavage sites has also been observed in Arabidopsis (Bienvenut et al. 2012; Rowland et al. 2015).

Figure 2. Comparison of experimentally observed N-termini with predicted organellar targeting peptide cleavage sites. Depicted are frequencies of identified N-termini around a predicted targeting peptide cleavage site. A difference of 0 indicates an identified N-terminal amino acid corresponds to the P1' amino acid of a predicted cleavage site. Cleavages of plastid (**A**), thylakoid lumen (**B**), mitochondrial (**C**) and secretory (**D**) targeting signals were predicted with *TargetP2.0*. All data are available from Supplemental Table S4. (**E**) Bar chart depicting the distribution of plastid protein isoforms with confirmed cleavage of a plastid targeting peptide and their identified N-terminal modifications. Percentages are related to the total number of identified proteins with a cleaved N-terminal plastid targeting peptide (748) within a window of ±5 amino acids around a predicted cleavage site. All data are available from Supplemental Table S4. Frequency of identified N-termini around a predicted plastid transit peptide cleavage site being either acetylated (**F**), unmodified (free, (**G**)) or monomethylated (**H**). Cum. [%]: cumulative percentage (red points).

N-terminal modifications of plastid and mitochondrial proteins

Among the proteins with identified plastid transit peptide cleavage site, we found 42% protein isoforms with an acetylated N-terminus and almost 10% with a monomethylated N-terminus (Figure 2E). Strikingly, plastid N-termini being acetylated and non-modified (free) both show this successive cleavage pattern whereas monomethylated N-termini do not show this pattern (Figures 2F-H).

This raises the question whether this processing occurs on both, acetylated and free N-termini, or whether only free N-termini are processed and subsequently acetylated. One explanation would be that N^α-acetylation of plastid proteins is incomplete and affects only a fraction of each protein isoform. Effectively, many N-termini of plastid proteins were identified in this study in a dual state, being acetylated and free (e.g., Pp3c18_19140V3.1, Pp3c15_7750V3.4; Supplemental Table S4). In this case, N^α-acetylation would prevent N-terminal trimming, whereas the fraction with an unmodified N-terminus would be proteolytically processed to different levels and subsequently acetylated. This scenario may be supported by the fact that both, N^α-acetylated and free N-termini share a similar preference of N-terminal amino acids (Figure 3) with alanine and serine being the most prominent ones. Apparently, the relative amino acid frequency of the N-terminally acetylated plastid proteins is strikingly similar to the relative frequency in Arabidopsis (Huesgen et al. 2013). Although the plastid protease inventory is under active investigation (van Wijk 2015; Meinnel and Giglione 2022), a specific protease for such N-terminal trimming has not yet been identified. Aminopeptidases

identified in Arabidopsis were recently proposed to confer also trimming functions (Rowland et al. 2015; Meinnel and Giglione 2022), but their activity was investigated on released plastid transit peptides in conjunction with pre-sequence proteases (Teixeira et al. 2017), but not on the protein N-terminus of the corresponding protein.

Figure 3 Sequence logos of identified N-termini with different modification states of plastid and mitochondrial proteins. "Dif" indicates the position difference upstream of a predicted plastid or mitochondrial transit peptide cleavage site. Transit peptide cleavage sites were predicted with *TargetP2.0*. The prediction data are available from Supplemental Table S4. "n" represents the total number of non-redundant sequences. Sequences were aligned at the identified N-terminal amino acid.

On the other hand, a cleavage of an N^α-acetylated amino acid may also be possible, although until now no protease has clearly proven activity on acetylated N-termini of intact proteins. However, acylamino acid-releasing enzyme (AARE), a bifunctional serine protease (Tsunasawa et al. 1975; Fujino et al. 2000; Shimizu et al. 2003; Nakai et al. 2012; Hoernstein et al. 2023), has a proven activity on N^α-acetylated oligopeptides and activity on intact proteins is repeatedly considered (Tsunasawa et al. 1975; Arfin and Bradshaw 1988; Adibekian et al. 2011). Moreover, one moss (Physcomitrella) AARE isoform and the Arabidopsis AARE are localized not only to the cytoplasm but also to plastids and mitochondria (Hoernstein et al. 2023). This renders AARE an interesting novel candidate for plastid protein processing, especially since there is a strong substrate preference of this protease towards Ac-Ala (Hoernstein et al. 2023; Yamauchi et al. 2003), which is the most frequent N^α-acetylated amino acid of plastid proteins observed here (Figure 3).

We also identified several mitochondrial proteins being N-terminally acetylated around a predicted transit peptide cleavage site (Supplemental Tables S4, S5). However, despite well-known N^α-acetylation of plastid proteins, this modification has not been identified yet to a similar extent on mitochondrial proteins and its apparent presence is not clear (Giglione and Meinnel 2021). The N-terminally acetylated amino acids were A, T, S and in all cases, they were preceded by a methionine which may also indicate alternative translation initiation. Consequently, we considered this not to be an as yet undiscovered modification of mitochondrial proteins, but rather as a co-translational modification of a shorter, e.g. cytoplasmic, isoform derived from alternative translation initiation or from alternative splicing. In Physcomitrella, both mechanisms are known to target protein isoforms to distinct subcellular localizations (Kiessling et al. 2004; Hoernstein et al. 2023). Apart from these two scenarios, dual targeting of proteins to plastids and mitochondria *via* ambiguous targeting signals is also considerable. In this case, the N-terminally acetylated protein would represent the plastid-localized variant. Hence, we investigated those N-terminally acetylated potentially mitochondria-localized proteins with a focus on alternative translation initiation sites or splice variants that would give rise to shorter, possibly cytoplasmic, protein isoforms (Supplemental Table S5). Subcellular targeting predictions were performed with *Localizer* (Sperschneider et al. 2017) and the presence of ambiguous targeting signals was predicted with *ATP2* (Fuß et al. 2013). In two cases (Pp3c13_17110V3.1, Pp3c21_2600V3.1) potential dual targeting to plastids and mitochondria was predicted by *Localizer* and *ATP2* but alternative translation

initiation from the downstream methionine was also likely (Supplemental Table S5). In most other cases, we found either a potential alternative translation initiation site (Pp3c15_21480V3.1, Pp3c4_3210V3.1) or alternative splice variants (Pp3c22_8300V3.1, Pp3c7_24050V3.2) that facilitate translation of a shorter open reading frame. For one protein (Pp3c9_14150V3.1, Pp3c9_14150V3.2) the situation remains unclear. Nevertheless, the present data indicate that the observed N-terminally acetylated proteins are localized in plastids or the cytoplasm rather than in mitochondria.

Besides acetylation, we also observed N-terminal methylation on plastid proteins and on mitochondrial proteins. Strikingly, monomethylation on plastid proteins was also identified predominantly on N-terminal alanine and serine but also on N-terminal methionine (Figure 3). The apparent absence of a successive cleavage pattern similar to that observed for free or acetylated N-termini may indicate a stabilizing effect of monomethylation on the modified protein. This modification is found in eukaryotes and prokaryotes (Stock et al. 1987) but is poorly investigated in plants. It has been proven for the small subunit of Rubisco (RbcS) in pea, spinach, barley and corn (Grimm et al. 1997). Accordingly, we found the N-terminal methionine of RbcS to be monomethylated at its N-terminal methionine (after transit peptide removal, Pp3c12_19890V3.4, Supplemental Figure S2) in Physcomitrella suggesting evolutionary conservation of this modification on RbcS.

Apart from those plastid proteins, we also identify 49 mitochondrial proteins being monomethylated at their N-terminus, matching the predicted presequence cleavage site (Supplemental Table S4), including cytochrome C subunit 5B (COX5B, Pp3c19_11870V3.1, Figure S3). A strong overrepresentation of serine as N-terminally monomethylated amino acid was observed (Figure 3), whereas alanine and serine were equally frequent on non-modified N-termini of mitochondrial proteins. This specificity of methylated proteins in plastids and mitochondria resembles only partially the specificity of human NTM1A (NAC with transmembrane motif1; UNIPROT: Q9BV86) (Schaner Tooley et al. 2010; Wu et al. 2015) which methylates N-terminal alanine and serine when followed by a proline and a lysine. In our data, proline and lysine were not frequently observed as subsequent amino acids (Figure 3). Intriguingly, the observed amino acid frequency of methylated plastid and mitochondrial proteins rather resembles the situation in yeast (Chen et al. 2021). Despite RbcS, N-terminal methylation of plant proteins was reported only on cytosolic and plastid ribosomal subunits

and histones (Carroll et al. 2008; Webb et al. 2010). Also, here the N-terminal amino acid sequence (Webb et al. 2010) share almost no homology with the methylated N-termini observed in our data. Nevertheless, we identified the cytosolic ribosomal subunit RPL19 to be methylated at its mature N-terminus (Pp3c18_14440V3.1, Figure S4), but also other likely cytosolic proteins (Supplemental Table S3).

Lastly, we also identified several proteins to be methylated at their N-terminus after cleavage of a predicted secretory signal peptide (Supplemental Table S4). Here, most of them seemed to be false positive identifications due to isotope peak errors. Hence, they were not further considered.

The knowledge of protein N-terminal methylation especially in plants is scarce and until now the methylation of RbcS was regarded as an exception (Grimm et al. 1997; Petkowski et al. 2013). In contrast, our data reveal that this modification affects several plastid and mitochondrial proteins with similar specificity of the methylating enzyme. In humans, two N-terminal methyltransferases are known so far, NTM1 and NTM2 (Schaner Tooley et al. 2010; Petkowski et al. 2013). To investigate whether homologues in Physcomitrella do exist, we used the sequence of human NTM1 (UNIPROT: Q9BV86) as query for a *BlastP* search (Altschul 1997) against all Physcomitrella V3.3 protein models (Lang et al. 2018) using the Phytozome database (https://phytozome-next.jgi.doe.gov/) (Goodstein et al. 2012) and identified a single protein (Pp3c22_8670V3.1; identity 36%, alignment length 74 amino acids) sharing the protein family annotations as human NTM1A (InterPro: Alpha-N-methyltransferase NTM1 (IPR008576); S-adenosyl-L-methionine-dependent methyltransferase (IPR029063). In a reciprocal *BlastP* search using Pp3c22_8670V3.1 (hereafter referred to as PpNTM1) as a query, human NTM1 appeared as best Blast hit confirming the orthology. A full *InterPro* search against all Physcomitrella V3.3 protein sequences (Lang et al. 2018) did not reveal any further hits with this protein family annotation. In agreement, PpNTM1 was also the only Blast hit when using the sequence of NTM2 (UNIPROT: Q5VVY1), a human homolog of NTM1, as Blast query. In Physcomitrella, PpNTM1 is expressed in all major tissues at moderate levels (Supplemental Figure S6). Interestingly, PpNTM1 is predicted via *TargetP2.0* (Armenteros et al. 2019) to localize to mitochondria, whereas plastid localization is predicted by *Localizer* (Sperschneider et al. 2017) (Supplemental Table S6). We also found only a single Arabidopsis homologue (AT5G44450.1) which is predicted to localize to plastids by both predictors. Yet, it

is not fully clear whether PpNTM1 is the responsible methyltransferase and whether it might be dually targeted to plastids and mitochondria, at least in Physcomitrella. Further, a potential alternative translation initiation site might be at M^{56} (Kozak Similarity Score ≥ 0.7 and < 0.8; Gleason et al. 2022a, b) which would not interfere with the predicted domain structure and enable cytosolic localization. However, our present data do not provide any support for predicted transit peptide cleavages or alternative translation initiation for this protein. Hence, further research is required to investigate function and localization of PpNTM1.

Conclusion

In the present study we used a compilation of 24 proteomic datasets obtained from various experiments to provide first insights into the N-terminome of the model plant Physcomitrella. We found that the percentage of N-terminal acetylation of cytosolic proteins appears slightly lower than the estimated percentage in Arabidopsis. Our data allow the confirmation of hundreds of predicted targeting peptide cleavage sites localizing proteins to plastids. These data can now be used to optimize computational targeting predictors, for customized protein fusions and their targeted localization in biotechnology, and it provides novel insights into potential dual targeting of proteins. Moreover, we reveal N-terminal monomethylation as a yet unknown modification of mitochondrial proteins. The function and impact of this modification remains to be further analyzed, but we propose PpNTM1 as a promising candidate for protein methylation in plastids, mitochondria and the cytosol.

Methods

Cell culture

For all experiments, Physcomitrella WT (new species name: *Physcomitrium patens* (Hedw.) Mitt.; Medina et al. 2019) ecotype "Gransden 2004" available from the international Moss Stock Centre (IMSC, www.moss-stock-center.org, #40001) was used. Cultivation was performed using Knop medium (Reski and Abel 1985) containing 250 mg/l KH_2PO_4, 250 mg/l KCl, 250 mg/l $MgSO_4$ x 7 H_2O, 1,000 mg/l $Ca(NO_3)_2$ x 4 H_2O and 12.5 mg/l $FeSO_4$ x 7 H_2O (pH 5.8). According to Egener et al. (2002) and Schween et al. (2003) 10 mL of a microelement solution 309 mg/l H_3BO_3, 845 mg/l $MnSO_4$ x 1 H_2O, 431 mg/l $ZnSO_4$ x 7 H_2O, 41.5 mg/l KI, 12.1 mg/l Na_2MoO_4 x 2 H_2O, 1.25 mg/l $CoSO_4$ x 5 H_2O, 1.46 $Co(NO_3)_2$ x 6 H_2O) was added per liter of medium. Gametophores were either cultivated on plates containing 12 g Agar per liter liquid medium or on hydroponic ring cultures as described in (Erxleben et al. 2012; Hoernstein et al. 2016). Hydroponic gametophore cultures were started from protonema culture that was dispersed weekly using an Ultra Turrax (IKA, Staufen, Germany) at 18,000 rpm for 90 s. All cultivation was done at 25°C in a day/night cycle of 16 h light with a light intensity of 70 µmol/sm^2 and 8 h dark.

Enrichment of nuclei from gametophores

Eighteen g fresh weight (FW) gametophores were harvested from hydroponic culture and chopped in buffer I containing 1 M 2-Methyl-2,4-pentandiol, 10 mM HEPES pH 7.5, 10 mM KCl 10 mM DTT, 0.1% PVP40, 0.1% PPI (P9599, Sigma-Aldrich) according to Nelson et al. (1994) using a custom 4 razorblade chopping device. The homogenate was successively filtered through a 40 µm and a 20 µm sieve and the flow-through was centrifuged for 30 min at 300 x g at 2°C. The supernatant was discarded, and the pellets were carefully dissolved in buffer II containing 110 mM KCl, 15 mM HEPES, pH 7.5, 5mM DTT and 0.1% PPI. The enriched nuclei were further purified using three-step Percoll-gradients (100%/60%/30%, 17-0891-01, GE Healthcare, Solingen, Germany) modified after Marienfeld et al. (1989). The Percoll-gradients were centrifuged at 200 x g at 2°C for 30 min. The interface between 100% and 60% was recovered as well as the pellet at the top of the gradient attached to the tube wall. Both fractions were strongly enriched in nuclei and thus pooled for further experiments. The samples were combined and washed with buffer II and centrifuged again for 10 min at 300 x g at 2°C. The pellet containing enriched nuclei was stored at -20°C until further use.

Sequential protein extraction from enriched nuclei

Pellets containing enriched nuclei were dissolved in 400 µL 50 mM Tris-HCl, pH7.6, 1% PPI (P9599, Sigma-Aldrich) and sonicated (Sonopuls HD2070, Bandelin, Berlin, Germany) 3 times for 20 s with an amplitude between 60 and 90%. 50 µL DNAse buffer and 50 µL DNase I (EN0521, Thermo Scientific, Waltham, USA) were added and the samples were incubated for 1 h at 37°C. After centrifugation at 20,000 x g for 20 min at 4°C the supernatant (Tris extract) was recovered and directly precipitated with 5 vol ice-cold acetone containing 0.2% DTT overnight at -20°C. The remaining pellet was dissolved in 400 µL 50 mM Tris-HCl, pH 7.6, 2% Triton X-100, 1% PPI, and again sonicated 3 times as before. Again, the sample was centrifuged, and the supernatant (Triton extract) was also acetone-precipitated overnight. The remaining pellet was dissolved in 50 mM Tris-HCl, pH 7.6, 4% SDS, 1% PPI, 50 mM DTT and incubated at 95°C for 10 min. The sample was centrifuged, and the supernatant was acetone-precipitated overnight. All acetone precipitations were centrifuged at 20,000 x g at 0°C for 15 min. The supernatant was discarded, and the remaining protein pellet was washed for 1 h with 1 volume ice-cold acetone without DTT. The centrifugation step was repeated, and the supernatant was discarded afterwards. The remaining protein pellets were air dried and stored at -20°C for further experiments.

Sequential protein extraction from gametophores

One to two g FW of gametophores were ground in liquid nitrogen for 10-15 min. The fine powder was dissolved in Tris buffer containing 40 mM Tris-HCl, pH 7.6, 0.5% PVPP and 1% PPI. The homogenate was sonicated for 15 min and afterwards centrifuged at 20,000 x g at 4°C for 30 min. The supernatant (Tris-extract) was recovered. The remaining pellet containing cell debris was dissolved in 40 mM Tris-HCl pH 7.6, 2% Triton X-100, 1% PPI and again sonicated for 15 min. Again, centrifugation was performed at 20,000 x g at 4°C for 30 min and the supernatant (Triton-extract) was recovered. Protein concentrations of the extracts were directly determined via the Bradford assay (Bradford 1976) and aliquots corresponding to 100 µg protein were precipitated with acetone containing DTT as described before.

Enrichment of N-terminal peptides

The dimethylation reaction was carried out according to Kleifeld et al. (2010) with some modifications. Protein pellets were dissolved in 100 mM HEPES-NaOH pH pH 7.5, 0.2% SDS. Reduction of cysteine residues was carried out using *Reducing Agent* (NP0009, Life

Technologies™, Carlsbad, USA) 1:10 at 95°C for 10 min or *Bond-Breaker®* (77720, Thermo Scientific) 1:100 at 28°C for 30 min. Alkylation was performed at a final concentration of 100 mM iodoacetamide for 20 min at RT. The dimethylation reaction was carried out by adding 2 µL of a 4% formaldehyde solution (Formaldehyde ^{13}C, d$_2$ solution, 596388, Sigma-Aldrich or Formaldehyde-D2, DLM-805-PK, Cambridge Isotope Laboratories Inc.) and 2 µL of a 500 mM NaCNBH$_3$ solution per 100 µL sample at 37°C for 4 h. The same volumes of formaldehyde and NaCNBH$_3$ were added again to the sample and the reaction was carried out overnight at 37°C. The dimethylation reaction was stopped by adding 2 µL of a 4% NH$_4$OH solution per 100 µL sample for 1 h at 37°C. Afterwards the samples were precipitated as described before using acetone without DTT for at least 3 h at -20°C. The final enrichment was modified after McDonald and Beynon (2006).

SDS-based enrichment

The dried protein pellets were dissolved in binding buffer according to McDonald and Beynon (2006) containing 20 mM NaH$_2$PO$_4$, 150 mM NaCl pH 7.5 with 0.2% SDS and in solution digest using either Trypsin (V5280, Promega, Madison, USA), GluC (90054, Thermo Scientific) or Chymotrypsin (V1062, Promega) was performed at an enzyme-to-substrate ratio of 1:25 for 4 h at 37°C (Trypsin, GluC) or 25°C (Chymotrypsin). Then the ratio was increased to 1:20 and the reaction was carried out overnight. Enrichment of N-terminal labeled peptides was carried out using 200 µL NHS-Sepharose slurry (17-0906-01, GE Healthcare, Solingen, Germany) per 100 µg protein. The slurry was centrifuged for 30 sec at 200 x g. The supernatant was discarded and 400 µL ice-cold 1 mM HCl was added. The slurry was centrifuged again, and the supernatant was discarded. Afterwards the Sepharose was washed with 1 mL binding buffer without SDS. The samples were applied to the prepared Sepharose and incubated for 4 h at RT. The Sepharose was again centrifuged, and the supernatant was transferred to a new tube containing freshly prepared Sepharose. The used Sepharose was washed with 20 µL binding buffer and the supernatant was added also to the freshly prepared Sepharose. The enrichment reaction was carried out overnight at 4-8°C. The enriched peptides were desalted using 200 µl C18 *StageTips* (SP301, Thermo Scientific) that were supplemented with an additional layer of *Empore^{TM} SPE Disk* C18 material (66883-U Sigma-Aldrich). The tips were washed prior to use with 100 µl 0.1% TFA and subsequently with 100 µl 80% ACN, 0.1% TFA. The tips were again equilibrated with 100 µl 0.1% TFA and the samples were loaded afterwards. The remaining

Sepharose was washed with 50 µl binding buffer and the supernatant was also transferred to the tip. The tips were washed with 100 µl binding buffer and the retained peptides were eluted with 300 µl 80% ACN, 0.1% TFA. The eluate was vacuum dried and the samples were stored at -20°C until further analysis.

RapiGest-based enrichment

The dried protein pellets were dissolved in 50 mM HEPES-KOH, 0.1% *Rapi*Gest surfactant (RPG, 18600186, Waters, Milford Massachusetts, USA). Proteolytic digest was performed as described before. After digestion, the RPG surfactant was cleaved by acidifying the sample to pH 2 using TFA as recommended by the manufacturer. The cleavage was performed at 37°C for 45 min. Insoluble RPG remnants were removed by centrifugation at 13000 rpm for 10 min at RT. The peptide-containing supernatant was subjected to solid phase extraction using SampliQ C18 cartridges (1 ml, 100 mg, 5982-1111, Agilent, Santa Clara, USA). Prior to the extraction the cartridge was washed successively with 1 ml 0.1% TFA,0.1% TFA in 80% ACN, 0.1% TFA. Then the peptide solution was applied and washed with1 ml 0.1% TFA. The peptides were eluted in 600 µl 0.1% TFA in 80% ACN and vacuum dried. The dried peptides were stored at -20°C until further use. For enrichment of the N-terminal peptides, the dried and purified peptides were dissolved in binding buffer as described in before without SDS and the enrichment using NHS-Sepharose was performed accordingly. Finally, the enriched N-terminal peptides were desalted again using SampliQ C18 cartridges as described before. The dried peptides were stored at -20°C until mass spectrometric analysis.

Mass spectrometry

NanoLC-MS/MS analyses were performed on an LTQ-Orbitrap Velos Pro (ThermoScientific) equipped with an EASY-Spray Ion Source and coupled to an EASY-nLC 1000 (Thermo Scientific). Peptides were loaded on a trapping column (2 cm x 75 µm ID. PepMap C_{18} 3 µm particles, 100 Å pore size, Dionex, Thermo Scientific) and separated either on a 25 cm EASY-Spray column (25 cm x 75 µm ID, PepMap C_{18} 2 µm particles,100 Å pore size) with a 30 min linear gradient from 3% to 30% ACN (V5280, Promochem) and 0.1% FA (56302, Thermo Scientific), or on a 50 cm EASY-Spray column (50 cm x 75 µm ID, PepMap C_{18} 2 µm particles, 100 Å pore size, Dionex, Thermo Scientific) with a 360 min linear gradient from 3% to 30% ACN and 0.1% FA in the case of in solution digested proteins such as the enriched N-terminal peptides. MS scans were acquired in the Orbitrap analyzer with a resolution of 30,000 at m/z 400, MS/MS scans were

acquired in the Orbitrap analyzer with a resolution of 7500 at m/z 400 using HCD fragmentation with 30% normalized collision energy. A TOP5 or TOP10 data-dependent MS/MS method was used. Dynamic exclusion was applied with a repeat count of 1 and an exclusion duration of 30 sec or 2 min in the case of long gradients. Singly charged precursors were excluded from selection. Minimum signal threshold for precursor selection was set to 50,000. Predictive AGC was used with a target value of 10^6 for MS scans and $5*10^4$ for MS/MS scans. Lock mass option was applied for internal calibration using background ions from protonated decamethylcyclopentasiloxane (m/z 371.10124). ETD fragmentation was performed with 35% normalized collision energy. A TOP5 data dependent MS/MS method was used. Dynamic exclusion was applied with a repeat count of 1 and an exclusion duration of 30 seconds. Singly charged precursors were excluded from selection. Minimum signal threshold for precursor selection was set to 75,000. Predictive AGC was used with AGC target a value of 10^6 for MS scans and $5*10^4$ for MS/MS scans. ETD activation time was set to 250 ms for doubly, 166 ms for triply and 125 ms for quadruple charged precursors, AGC target was set to 200,000 for fluoranthene. Lock mass option was applied for internal calibration in all runs using background ions from iron(III) citrate (m/z 263.956311).

Raw data processing and database search

Raw data were processed with *Mascot Distiller* (V2.8.3.0, https://www.matrixscience.com/) and database searches were performed using *Mascot Server* (V2.7.0, https://www.matrixscience.com) against a database containing all V3 Physcomitrella protein models (Lang et al. 2018) as well as their reversed sequences as decoys. In parallel, a search was performed against a database containing the sequences of known contaminants, such as keratin etc. (269 entries, available on request). For all samples semi-specific protease specificities were chosen and in the case of tryptic digests the specificity was set to semi-ArgC. Variable modifications were Gln >pyro Glu (N term Q) -17.026549 Da, oxidation (M) +15.994915 Da, acetyl (N-term) +42.010565 Da, $^{13}C,D_2$ dimethyl (N-term) +34.063117 Da, hybrid-methylation (N-term) +31.047208 ($^{13}CD_2CH_2$) Da and deamidation of asparagine (N) +0.984016 Da. Fixed modifications were carbamidomethyl (C) +57.021464 Da and $^{13}C,d_2$ dimethyl (K) +34.063117 Da. In the case of samples dimethylated with D_2 formaldehyde, the dimethylation (K, N-term) mass shift was +32.056407 Da and hybrid-methylation (N-term) was set to +30.043854 Da ($C_2D_2H_2$). A precursor mass tolerance of ±8 ppm and a fragment mass

tolerance of ±0.02 Da was specified. Search results were loaded in *Scaffold5^{TM}* software (V5.0.1, https://www.proteomesoftware.com/) using the high mass accuracy scoring and independent sample grouping method.

Computational analysis

The presence of cleavable N-terminal targeting signals (plastid, mitochondria, secretory) was performed with *TargetP2.0* (Armenteros et al. 2019) and in selected cases with *Localizer* (Sperschneider et al. 2017). Ambiguous targeting to plastids and mitochondria in Physcomitrella was predicted with *ATP2* (Fuss et al. 2013). Potential alternative translation initiation was predicted with TIS (https://www.tispredictor.com/) (Gleason et al. 2022a, b). All plots and tables were created using custom PERL scripts and *R* (R Core Team 2022).

Data availability

The mass spectrometry proteomics data have been deposited to the *ProteomeXchange* Consortium via the *PRIDE* partner repository (Deutsch et al. 2023; Perez-Riverol et al. 2022) with the dataset identifier PXD052824 and 10.6019/PXD052824.

Author contributions

R.R., G.L.I. and A.S. designed research and acquired funding. D.L. supervised research and helped analyzing the data. S.N.W.H. and K.F. performed experiments. S.N.W.H. analyzed the data and wrote the manuscript together with R.R. All authors approved the final version of the manuscript.

Conflict of Interest Statement

All authors declare no conflict of interest.

Acknowledgements

Funding by Deutsche Forschungsgemeinschaft (DFG IG9/8-1 to G.L.I., A. S. and R.R.) and under Germany's Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984) and by Wissenschaftliche Gesellschaft Freiburg is gratefully acknowledged. We thank Christine Glockner and Stephanie Lamer for technical assistance and Anne Katrin Prowse for proof reading.

References

Adibekian A, Martin BR, Wang C, Hsu KL, Bachovchin DA, Niessen S, Cravatt BF (2011) Click-generated triazole ureas as ultrapotent *in vivo*–active serine hydrolase inhibitors. Nat Chem Biol 7:469-478. https://doi.org/10.1038/nchembio.579

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25:3389–3402. https://doi.org/10.1093/nar/25.17.3389

Arfin SM, Bradshaw RA (1988) Cotranslational processing and protein turnover in eukaryotic cells. Biochemistry 27, 7979-7984. https://doi.org/10.1021/bi00421a001

Armenteros JJA, Salvatore M, Emanuelsson O, Winther, O Von Heijne G, Elofsson A Nielsen H (2019) Detecting sequence signals in targeting peptides using deep learning. Life Sci Alliance 2:e201900429. https://doi.org/10.26508/lsa.201900429

Ashton NW, Grimsley NH, Cove DJ (1979) Analysis of gametophytic development in the moss, *Physcomitrella patens*, using auxin and cytokinin resistant mutants. Planta 144:427-435. https://doi.org/10.1007/BF00380118

Bi G, Zhao S, Yao J, Wang H, Zhao M, Sun Y, Hou X, Haas FB, Varshney D, Prigge M, Rensing SA, Jiao Y, Ma Y, Yan J, Dai J (2024) Near telomere-to-telomere genome of the model plant *Physcomitrium patens*. Nat Plants 10:327–343. https://doi.org/10.1038/s41477-023-01614-7

Bienvenut WV, Sumpton D, Martinez A, Lilla S, Espagne C, Meinnel T, Giglione C (2012) Comparative large scale characterization of plant versus mammal proteins reveals similar and idiosyncratic N-α-acetylation features. Mol Cell Proteomics 11.6. https://doi.org/10.1074/mcp.M111.015131

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72, 248-254. https://doi.org/10.1016/0003-2697(76)90527-3

Carroll AJ, Heazlewood JL, Ito J, Millar AH (2008) Analysis of the Arabidopsis cytosolic ribosome proteome provides detailed insights into its components and their post-translational modification. Mol Cell Proteomics 7:347-369. https://doi.org/10.1074/mcp.M700052-MCP200

Chen P, Paschoal Sobreira TJ, Hall MC, Hazbun TR (2021) Discovering the N-terminal methylome by repurposing of proteomic datasets. J Proteome Res 20:4231-4247. https://doi.org/10.1021/acs.jproteome.1c00009

Decker EL, Reski R (2020) Mosses in biotechnology. Curr Opin Biotech 61:21–27. https://doi.org/10.1016/j.copbio.2019.09.021

Decker EL, Alder A, Hunn S, Ferguson J, Lehtonen MT, Scheler B, Kerres KL, Wiedemann G, Safavi-Rizi V, Nordzieke S, Balakrishna A, Baz L, Avalos J, Valkonen JPT, Reski R, Al-Babili S (2017) Strigolactone biosynthesis is evolutionarily conserved, regulated by phosphate

starvation and contributes to resistance against phytopathogenic fungi in a moss, *Physcomitrella patens*. New Phytol 216:455-468. https://doi.org/10.1111/nph.14506

Demir F, Perrar A, Mantz M, Huesgen PF (2022) Sensitive plant N-terminome profiling with HUNTER. In Plant Proteases and Plant Cell Death: Methods and Protocols 139-158. New York, NY: Springer US. https://doi.org/10.1007/978-1-0716-2079-3_12

Deutsch EW, Bandeira N, Perez-Riverol Y, Sharma V, Carver JJ, Mendoza L, Kundu DJ, Wang S, Bandla C, Kamatchinathan S, Hewapathirana S, Pullman BS, Wertz J, Sun Z, Kawano S, Okuda S, Watanabe Y, MacLean B, MacCoss MJ, Zhu Y, Ishihama Y, Vizcaíno JA (2023) The ProteomeXchange consortium at 10 years: 2023 update. Nucleic Acids Res 51:D1539-D1548. https://doi.org/10.1093/nar/gkac1040

Erxleben A, Gessler A, Vervliet-Scheebaum M, Reski R (2012) Metabolite profiling of the moss *Physcomitrella patens* reveals evolutionary conservation of osmoprotective substances. Plant Cell Rep 31:427-436. https://doi.org/10.1007/s00299-011-1177-9

Fernandez-Pozo N, Haas FB, Meyberg R, Ullrich KK, Hiss M, Perroud P-F, Hanke S, Kratz V, Powell AF, Vesty EF, Daum CG, Zane M, Lipzen A, Sreedasyam A, Grimwood J, Coates JC, Barry K, Schmutz J, Mueller LA, Rensing SA (2020) PEATmoss (Physcomitrella Expression Atlas Tool): a unified gene expression atlas for the model plant *Physcomitrella patens*. Plant J 102:165–177. https://doi.org/10.1111/tpj.14607

Fesenko I, Kirov I, Kniazev A, Khazigaleeva R, Lazarev V, Kharlampieva D, Grafskaia E, Zgoda V, Butenko I, Arapidi G, Mamaeva A, Ivanov V, Govorun V (2019) Distinct types of short open reading frames are translated in plant cells. Genome Res 29:1464-1477. https://doi.org/10.1101/gr.253302.119

Fesenko I, Shabalina SA, Mamaeva A, Knyazev A, Glushkevich A, Lyapina I, Ziganshin R, Kovalchuk S, Kharlampieva D, Lazarev V, Taliansky M, Koonin EV (2021) A vast pool of lineage-specific microproteins encoded by long non-coding RNAs in plants. Nucleic Acids Res 49:10328-10346. https://doi.org/10.1093/nar/gkab816

Fortelny N, Yang S, Pavlidis P, Lange PF, Overall CM (2015) Proteome TopFIND 3.0 with TopFINDer and PathFINDer: database and analysis tools for the association of protein termini to pre-and post-translational events. Nucleic Acids Res 43:D290-D297. https://doi.org/10.1093/nar/gku1012

Fujino T, Kojima M, Beppu M, Kikugawa K, Yasuda H, Takahashi K (2000) Identification of the cleavage sites of oxidized protein that are susceptible to oxidized protein hydrolase (OPH) in the primary and tertiary structures of the protein. J Biochem 127:1087-1093. https://doi.org/10.1093/oxfordjournals.jbchem.a022702

Fuss J, Liegmann O, Krause K, Rensing SA (2013) Green targeting predictor and ambiguous targeting predictor 2: the pitfalls of plant protein targeting prediction and of transient protein expression in heterologous systems. New Phytol 200:1022-1033. https://doi.org/10.1111/nph.12433

Giglione C, Meinnel T (2021) Evolution-driven versatility of N terminal acetylation in photoautotrophs. Trends Plant Sci 26:375-391. https://doi.org/10.1016/j.tplants.2020.11.012

Gleason AC, Ghadge G, Chen J, Sonobe Y, Roos RP (2022a) Machine learning predicts translation initiation sites in neurologic diseases with nucleotide repeat expansions. PLOS One 17:e0256411. https://doi.org/10.1371/journal.pone.0256411

Gleason AC, Ghadge G, Sonobe Y, Roos RP (2022b) Kozak similarity score algorithm identifies alternative translation initiation codons implicated in cancers. Int J Mol Sci 23:10564. https://doi.org/10.3390/ijms231810564

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Therese Mitros, Dirks W, Hellsten U, Putnam N, Rokhsar DS (2012) Phytozome: a comparative platform for green plant genomics. Nucleic Acids Res 40:D1178-D1186. https://doi.org/10.1093/nar/gkr944

Grimm R, Grimm M, Eckerskorn C, Pohlmeyer K, Röhl T, Soll J (1997) Postimport methylation of the small subunit of ribulose-1, 5-bisphosphate carboxylase in chloroplasts. FEBS Lett 408:350-354. https://doi.org/10.1016/S0014-5793(97)00462-6

Heintz D, Erxleben A, High AA, Wurtz V, Reski R, Van Dorsselaer A, Sarnighausen E (2006) Rapid alteration of the phosphoproteome in the moss *Physcomitrella patens* after cytokinin treatment. J Prot Res 5:2283-2293. https://doi.org/10.1021/pr060152e

Hoernstein SNW, Mueller SJ, Fiedler K, Schuelke M, Vanselow JT, Schuessele C, Lang D, Nitschke R, Igloi GL, Schlosser A, Reski, R. (2016) Identification of targets and interaction partners of arginyl-tRNA protein transferase in the moss *Physcomitrella patens*. Mol Cell Proteomics 15:1808-1822. https://doi.org/10.1074/mcp.M115.057190

Hoernstein SNW, Fode B, Wiedemann G, Lang D, Niederkrueger H, Berg B, Schaaf A, Frischmuth T, Schlosser A, Decker EL, Reski R (2018) Host cell proteome of *Physcomitrella patens* harbors proteases and protease inhibitors under bioproduction conditions. J Proteome Res 17:3749–3760. https://doi.org/10.1021/acs.jprot eome.8b00423

Hoernstein SNW, Özdemir B, van Gessel N, Miniera AA, Rogalla von Bieberstein B, Nilges L, Schweikert Farinha J, Komoll R, Glauz S, Weckerle T, Scherzinger F, Rodriguez-Franco M, Müller-Schüssele SJ, Reski R. (2023) A deeply conserved protease, acylamino acid-releasing enzyme (AARE), acts in ageing in Physcomitrella and Arabidopsis. Comm Biol 6:61. https://doi.org/10.1038/s42003-023-04428-7

Horst NA, Katz A, Pereman I, Decker EL, Ohad N, Reski R (2016) A single homeobox gene triggers phase transition, embryogenesis and asexual reproduction. Nat Plants 2:15209. https://doi.org/10.1038/nplants.2015.209

Huesgen PF, Alami M, Lange PF, Foster LJ, Schröder WP, Overall CM, Green, BR (2013) Proteomic amino-termini profiling reveals targeting information for protein import into complex plastids. PLOS One 8:e74483. https://doi.org/10.1371/journal.pone.0074483

Keller A, Nesvizhskii AI, Kolker E, Aebersold R (2002) Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. Anal Chem 74:5383-5392. https://doi.org/10.1021/ac025747h

Kiessling J, Martin A, Gremillon L, Rensing SA, Nick P, Sarnighausen E, Reski R (2004) Dual targeting of plastid division protein FtsZ to chloroplasts and the cytoplasm. EMBO Rep 5:889-894. https://doi.org/10.1038/sj.embor.7400238

Kleifeld O, Doucet A, auf dem Keller U, Prudova A, Schilling O, Kainthan RK, Starr AE, Leonard Foster J, Kizhakkedathu JN, Overall CM. (2010) Isotopic labeling of terminal amines in complex samples identifies protein N-termini and protease cleavage products. Nat Biotech 28:281-288. https://doi.org/10.1038/nbt.1611

Kunze M, Berger J (2015) The similarity between N-terminal targeting signals for protein import into different organelles and its evolutionary relevance. Front Physiol 6:159761. https://doi.org/10.3389/fphys.2015.00259

Lang D, Ullrich KK, Murat F, Fuchs J, Jenkins J, Haas FB, Piednoel M, Gundlach H, Van Bel M, Meyberg R, Vives C, Morata J, Symeonidi A, Hiss M, Muchero W, Kamisugi Y, Saleh O, Blanc G, Decker EL, van Gessel N, Grimwood J, Hayes RD, Graham SW, Gunter LE, McDaniel S, Hoernstein SNW, Larsson A, Li F-W, Phillips J, Ranjan P, Rokshar DS, Rothfels CJ, Schneider L, Shu S, Stevenson DW, Thümmler F, Tillich M, Villarreal A. JC, Widiez T, Wong GK-S, Wymore A, Zhang Y, Zimmer AD, Quatrano RS, Mayer KFX, Goodstein D, Casacuberta JM, Vandepoele K, Reski R, Cuming AC, Tuskan J, Maumus F, Salse J, Schmutz J, Rensing SA (2018) The *Physcomitrella patens* chromosome-scale assembly reveals moss genome structure and evolution. Plant J 93:515–533. https://doi.org/10.1111/TPJ.13801

Lueth VM, Reski R (2023). Mosses. Curr Biol 33:R1175–R1181. https://doi.org/10.1016/j.cub.2023.09.042

Linster E, Wirtz M (2018) N-terminal acetylation: an essential protein modification emerges as an important regulator of stress responses. J Ex Bot 69:4555-4568. https://doi.org/10.1093/jxb/ery241

Maclean J, Koekemoer M, Olivier AJ, Stewart D, Hitzeroth II, Rademacher T, Fischer R, Williamson AL, Rybicki EP (2007). Optimization of human papillomavirus type 16 (HPV-16) L1 expression in plants: comparison of the suitability of different HPV-16 L1 gene variants and different cell-compartment localization. J Gen Virol 88, 1460-1469. https://doi.org/10.1099/vir.0.82718-0

Marienfeld JR, Reski R, Friese C, Abel WO (1989) Isolation of nuclear, chloroplast and mitochondrial DNA from the moss *Physcomitrella patens*. Plant Sci 61:235-244. https://doi.org/10.1016/0168-9452(89)90230-6

McDonald L, Beynon RJ (2006) Positional proteomics: preparation of amino-terminal peptides as a strategy for proteome simplification and characterization. Nat Protoc 1:1790-1798. https://doi.org/10.1038/nprot.2006.317

Medina R, Johnson MG, Liu Y, Wickett NJ, Shaw AJ, Gofnet B (2019) Phylogenomic delineation of Physcomitrium (Bryophyta: Funariaceae) based on targeted sequencing of nuclear exons and their fanking regions rejects the retention of Physcomitrella, Physcomitridium and Aphanorrhegma. J Syst Evol 57:404–417. https://doi.org/10.1111/JSE.12516

Meinnel T, Giglione C (2008) Tools for analyzing and predicting N-terminal protein modifications. Proteomics 8:626-649. https://doi.org/10.1002/pmic.200700592

Meinnel T, Giglione C (2022) N-terminal modifications, the associated processing machinery, and their evolution in plastid-containing organisms. J Exp Bot 73:6013-6033. https://doi.org/10.1093/jxb/erac290

Mueller SJ., Lang D, Hoernstein SNW, Lang EG, Schuessele C, Schmidt A, Fluck M, Leisibach D, Niegl C, Zimmer AD, Schlosser A, Reski R (2014) Quantitative analysis of the mitochondrial and plastid proteomes of the moss *Physcomitrella patens* reveals protein macrocompartmentation and microcompartmentation. Plant Physiol 164:2081-2095. https://doi.org/10.1104/pp.114.235754

Nakai A, Yamauchi Y, Sumi S, Tanaka K (2012) Role of acylamino acid-releasing enzyme/oxidized protein hydrolase in sustaining homeostasis of the cytoplasmic antioxidative system. Planta 236:427-436. https://doi.org/10.1007/s00425-012-1614-1

Nelson D, Salamini F, Bartels D (1994) Abscisic acid promotes novel DNA-binding activity to a desiccation-related promoter of *Craterostigma plantagineum*. Plant J 5:451-458. https://doi.org/10.1046/j.1365-313X.1994.05040451.x

Nesvizhskii AI, Keller A, Kolker E, Aebersold R (2003) A statistical model for identifying proteins by tandem mass spectrometry. Anal Chem 75:4646-4658. https://doi.org/10.1021/ac0341261

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaíno JA (2022) The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res 50:D543-D552. https://doi.org/10.1093/nar/gkab1038

Perroud P-F, Haas FB, Hiss M, Ullrich KK, Alboresi A, Amirebrahimi M, Barry K, Bassi R, Bonhomme S, Chen H, Coates JC, Fujita T, Guyon-Debast A, Lang D, Lin J, Lipzen A, Nogué F, Oliver MJ, Ponce de León I, Quatrano RS, Rameau C, Reiss B, Reski R, Ricca M, Saidi Y, Sun N, Szövenyi P, Sreedasyam A, Grimwood J, Stacey G, Schmutz J, Rensing SA (2018) The *Physcomitrella patens* gene atlas project: large-scale RNA-seq based expression data. Plant J 95:168–182. https://doi.org/10.1111/tpj.13940

Petkowski JJ, Bonsignore LA, Tooley JG, Wilkey DW, Merchant ML, Macara IG, Schaner Tooley, CE (2013) NRMT2 is an N-terminal monomethylase that primes for its homologue NRMT1. Biochem J 456:453-462. https://doi.org/10.1042/BJ20131163

Purwaha P, Silva LP, Hawke DH, WeinsteinJN, Lorenzi PL (2014). An artifact in LC-MS/MS measurement of glutamine and glutamic acid: in-source cyclization to pyroglutamic acid. Anal Chem 86:5633-5637. https://doi.org/10.1021/ac501451v

R Core Team (2022) R: A Language and environment for statistical computing. R Foundation for Statistical Computing. https://www.R-project.org

Ree R, Varland S, Arnesen T. (2018) Spotlight on protein N-terminal acetylation. Exp Mol Med 50:1-13. https://doi.org/10.1038/s12276-018-0116-z

Renault H, Alber A, Horst NA, Basilio Lopes A, Fich EA, Kriegshauser L, Wiedemann G, Ullmann P, Herrgott L, Erhardt M, Pineau E, Ehlting J, Schmitt M, Rose JKC, Reski R, Werck-Reichhart D (2017) A phenol-enriched cuticle is ancestral to lignin evolution in land plants. Nat Comm 8: 14713. https://doi.org/10.1038/ncomms14713

Rensing SA, Lang D, Zimmer AD, Terry A, Salamov A, Shapiro H, Nishiyama T, Perroud PF, Lindquist EA, Kamisugi Y, Tanahashi T, Sakakibara K, Fujita T, Oishi K, Shin-I T, Kuroki Y, Toyoda A, Suzuki Y, Hashimoto SI, Yamaguchi K, Sugano S, Kohara Y, Fujiyama A, Ashton N, Anterola A, Aoki S, Ashton N, Barbazuk WB, Barker E, Bennetzen JL, Blankenship R, Cho SH, Dutcher SK, Estelle M, Fawcett JA, Gundlach H, Hanada K, Heyl A, Hicks KA, Hughes J, Lohr M, Mayer K, Melkozernov A, Murata T, Nelson DR, Pils B, Prigge M, Reiss B, Renner T, Rombauts S, Rushton PJ, Sanderfoot A, Schween G, Shiu SH, Stueber K, Theodoulou FL, Tu H, Van de Peer Y, Verrier PJ, Waters E, Wood A, Yang L, Cove D, Cuming AC, Hasebe M, Lucas S, Mishler BD, Reski R, Grigoriev IV, Quatrano RS, Boore JL (2008) The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants. Science 319:64-69. https://doi.org/10.1126/science.1150646

Reski R, Abel WO (1985) Induction of budding on chloronemata and caulonemata of the moss, *Physcomitrella patens*, using isopentenyladenine. Planta 165:354–358. https://doi.org/10.1007/BF00392232

Rowland E, Kim J, Bhuiyan NH, van Wijk KJ (2015) The Arabidopsis chloroplast stromal N-terminome: complexities of amino-terminal protein maturation and stability. Plant Physiol 169:1881-1896. https://doi.org/10.1104/pp.15.01214

Ruiz-Molina N, Parsons J, Müller M, Hoernstein SNW, Bohlender LL, Pumple S, Zipfel PF, Häffner K, Reski R, Decker EL (2022) A synthetic protein as efficient multitarget regulator against complement over-activation. Comm Biol 5:152. https://doi.org/10.1038/s42003-022-03094-5

Sarnighausen E, Wurtz V, Heintz D, van Dorsselaer A, Reski R (2004) Mapping of the *Physcomitrella patens* proteome. Phytochem 65:1589-1607. https://doi.org/10.1016/j.phytochem.2004.04.028

Schaaf A, Tintelnot S, Baur A, Reski R, Gorr G, Decker EL (2005) Use of endogenous signal sequences for transient production and efficient secretion by moss (*Physcomitrella patens*) cells. BMC Biotech 5:30. https://doi.org/10.1186/1472-6750-5-30

Schaner Tooley CE, Petkowski JJ, Muratore-Schroeder TL, Balsbaugh JL, Shabanowitz J, Sabat M, Minor D, Hunt DF, Macara IG (2010) NRMT is an α-N-methyltransferase that methylates RCC1 and retinoblastoma protein. Nature 466:1125-1128. https://doi.org/10.1038/nature09343

Schilling S, Wasternack C, Demuth HU (2008) Glutaminyl cyclases from animals and plants: a case of functionally convergent protein evolution. Biol Chem 389:983–991. https://doi.org/10.1515/BC.2008.111

Schween G, Hohe A, Koprivova A, Reski R (2003) Effects of nutrients, cell density and culture techniques on protoplast regeneration and early protonema development in a moss,

Physcomitrella patens. J Plant Physiol 160:209–212. https://doi.org/10.1078/0176-1617-00855

Shimizu K, Fujino T, Ando K, Hayakawa M, Yasuda H, Kikugawa K (2003) Overexpression of oxidized protein hydrolase protects COS-7 cells from oxidative stress-induced inhibition of cell growth and survival. Biochem Biophys Res Comm 304:766-771. https://doi.org/10.1016/S0006-291X(03)00657-0

Sperschneider J, Catanzariti AM, DeBoer K, Petre B, Gardiner DM, Singh KB, Dodds, PN, Taylor, JM (2017) LOCALIZER: subcellular localization prediction of both plant and effector proteins in the plant cell. Sci Rep 7:44598. https://doi.org/10.1038/srep44598

Staes A, Impens F, Van Damme P, Ruttens B, Goethals M, Demol H, Timmerman E, Vandekerckhove J, Gevaert K (2011) Selecting protein N-terminal peptides by combined fractional diagonal chromatography. Nat Protoc 6:1130-1141. https://doi.org/10.1038/nprot.2011.355

Stock A, Clarke S, Clarke C, Stock J (1987) N-terminal methylation of proteins: structure, function and specificity. FEBS Lett 220:8-14. https://doi.org/10.1016/0014-5793(87)80866-9

Tasaki T, Sriram SM, Park KS, Kwon YT (2012) The N-end rule pathway. Annu Rev Biochem 81:261-289. https://doi.org/10.1146/annurev-biochem-051710-093308

Teixeira PF, Kmiec B, Branca RM, Murcha MW, Byzia A, Ivanova A, Whelan J, Drag M, Lehtiö J, Glaser E. (2017) A multi-step peptidolytic cascade for amino acid recovery in chloroplasts. Nat Chem Biol 13:15-17. https://doi.org/10.1038/nchembio.2227

Tschongov T, Konwar S, Busc, A, Sievert C, Hartmann A, Noris M, Gastoldi S, Aiello S, Schaaf A Panse J, Zipfel PF, Dabrowska-Schlepp P, Häffner K (2024) Moss-produced human complement factor H with modified glycans has an extended half-life and improved biological activity. Front Immun 15:1383123. https://doi.org/10.3389/fimmu.2024.1383123

Tsunasawa S, Narita K, Ogata K (1975) Purification and properties of acylamino acid-releasing enzyme from rat liver. J Biochem 77:89-102. https://doi.org/10.1093/oxfordjournals.jbchem.a130722

van Wijk KJ. (2015) Protein maturation and proteolysis in plant plastids, mitochondria, and peroxisomes. Annu Rev Plant Biol 66:75-111. https://doi.org/10.1146/annurev-arplant-043014-115547

Varshavsky A (1996) The N-end rule: functions, mysteries, uses. Proc Natl Acad Sci USA 93:12142-12149. https://doi.org/10.1073/pnas.93.22.12142

Varshavsky A (2019) N-degron and C-degron pathways of protein degradation. Proc Natl Acad Sci USA 116:358-366. https://doi.org/10.1073/pnas.181659611

Webb KJ, Lipson RS, Al-Hadid Q, Whitelegge JP, Clarke SG (2010) Identification of protein N-terminal methyltransferases in yeast and humans. Biochem 49:5225-5235. https://doi.org/10.1021/bi100428x

Wiedemann G, van Gessel N, Köchl F, Hunn L, Schulze K, Maloukh L, Nogué F, Decker EL, Hartung F, Reski, R (2018) RecQ helicases function in development, DNA repair, and gene targeting in *Physcomitrella patens*. Plant Cell 30:717-736. https://doi.org/10.1105/tpc.17.00632

Wu R, Yue Y, Zheng X, Li H (2015) Molecular basis for histone N-terminal methylation by NRMT1. Gene Dev, 29:2337-2342. https://doi.org/10.1101/gad.270926.115

Yamauchi Y, Ejiri Y, Toyoda Y, Tanaka K (2003) Identification and biochemical characterization of plant acylamino acid–releasing enzyme. J Biochem 134:251-257. https://doi.org/10.1093/jb/mvg138

Supplementary information

Figure S1 Overview on the numbers of cleavable targeting sequences in Physcomitrella predicted by *TargetP2.0*

Figure S2 HCD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of RbcS

Figure S3 HCD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of COX5B

Figure S4 ETD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of RPL19

Figure S5 Expression levels of PpNTM1 in different tissues and cell culture types of Physcomitrella

Table S1 Overview on the performed experiments, employed culture type and sample processing

Table S2 Protein identification summary on all experiments exported from *Scaffold5* software

Table S3 Overview on identified N-terminal peptides

Table S4 Comparison of identified N-terminal positions with predicted transit peptide cleavage sites

Table S5 Overview on proteins possibly dually targeted to mitochondria and the cytosol or to plastids

Table S6 Overview on subcellular predictions for PpNTM1

Figure S1 Overview on the numbers of cleavable targeting sequences in Physcomitrella predicted by *TargetP2.0*. Predictions were done on a non-redundant protein isoform list (49,589 entries) of all Physcomitrella V3.3 protein models (Lang et al. 2018). luTP: thylakoid luminal targeting peptide; mTP: mitochondrial targeting peptide; cTP: plastid targeting peptide; SP: (secretory) signal peptide; noTP: no targeting peptide predicted.

(A)

(B)

Figure S2 HCD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of RbcS (Pp3c12_19890V3.4). (A) HCD fragment mass spectrum of the peptide MQVWNPIGMTKFE. A mass shift of +31 indicates the hybrid modification of a post-translationally incorporated methyl group and the further addition methyl group from the preformed reductive methylation during sample preparation ($^{13}CD_2CH_2$, +31.047208). **(B)** Fragment mass error distribution of the b- and y-ion series.

(A)

(B)

Figure S3 HCD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of COX5B (Pp3c19_11870V3.1). (**A**) HCD fragment mass spectrum of the peptide SGHAASGQLDEFGIATGAER. A mass shift of +31 indicates the hybrid modification of a post-translationally incorporated methyl group and the further addition methyl group from the preformed reductive methylation during sample preparation ($^{13}CD_2CH_2$, +31.047208). (**B**) Fragment mass error distribution of the b- and y-ion series.

(A)

(B)

Figure S4 ETD fragment mass spectrum and fragment mass error distribution of the identified N-terminal peptide of RPL19 (Pp3c18_14440V3.1). (A) ETD fragment mass spectrum of the peptide GKQISEIKDFLLTAR. A mass shift of +30 indicates the hybrid modification of a post-translationally incorporated methyl group and the further addition methyl group from the preformed reductive methylation during sample preparation ($C_2D_2H_2$, +30.043854 Da). **(B)** Fragment mass error distribution of the b- and y-ion series.

Figure S5 Expression levels of PpNTM1 in different tissues and cell culture types of Physcomitrella. Expression levels are represented as mean fpkm (fragments per kilobase million) values with standard deviation. Single datapoints represent biological replicates. All replicate data were downloaded from *PEATmoss* (https://peatmoss.plantcode.cup.uni-freiburg.de/) and sampling and methods are described in Perroud et al. (2018) and Fernandez-Pozo et al. (2020). Abbreviations: B = BCD medium (Ashton et al. 1979); K = Knop medium (Reski and Abel 1985); A =ammonium tartrate; lq = liquid; sl = solid; Germ. = germinating.