Cyclical MinD membrane affinity differences are not necessary for MinD gradient

formation in *Bacillus subtilis*

Laura C. Bohorquez[1, 4], Henrik Strahl[2, #], Davide Marenduzzo[3], Martin J. Thiele[2, 5], Frank

Bürmann[2, 6], Leendert W. Hamoen[1, #]

[1] The Bacterial Cell Biology, Swammerdam Institute for Life Sciences, University of

Amsterdam, Amsterdam, The Netherlands.

[2] The Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle

upon Tyne, United Kingdom.

[3] SUPA, School of Physics and Astronomy, The University of Edinburgh, Edinburgh, Scotland,

United Kingdom.

[#]   For   correspondence: Henrik   Strahl   (h.strahl@ncl.ac.uk),   Leendert   Hamoen

(l.w.hamoen@uva.nl)

[4] Present address: FIND, Geneva, Switzerland

[5] Present address: Carl Zeiss Meditec AG, Jena, Germany

[6] Present address: MRC Laboratory of Molecular Biology, Structural Studies Division,

Cambridge, UK

Running title: Effect of MinD membrane affinity on gradient formation

ABSTRACT

Proteins can diffuse micrometers in seconds, yet bacterial cells are able to maintain stable

protein gradients. The best studied bacterial protein gradient is the Min system of

Escherichia coli. In rod-shaped bacteria the MinCD proteins prevent formation of minicells

by inhibiting FtsZ polymerization close to the cell poles. In *E. coli* these proteins oscillate

between cell poles within a minute, resulting in an increased MinCD concentration at the

poles. This oscillation is caused by the interaction between MinD and the protein MinE,

which form an ATP-driven reaction-diffusion system, whereby the ATPase MinD cycles

between a monomeric cytosolic and a dimeric membrane attached states. *Bacillus subtilis*

also has MinCD, but lacks MinE. In this case MinCD form a static gradient that requires the

transmembrane protein MinJ, located at cell poles and cell division sites. A recent reaction-

diffusion model was successful in recreating the MinD gradient in *B. subtilis*, assuming that

MinD cycles between cytosol and membrane, like in *E. coli*. Here we show that the

monomeric and dimeric states of *B. subtilis* MinD have comparable membrane affinities,

that MinD interacts with MinJ as a dimer, and that MinJ is not required for membrane

localization of MinD. Based on these new findings we tested different models, using kinetic

Monte Carlo simulations, and found that a difference in diffusion rate between the

monomer and dimer, rather than a difference in membrane affinity, is important for *B.*

subtilis MinCD gradient formation.

IMPORTANCE (150 max)

Proteins can diffuse micrometers in seconds, yet bacterial cells are able to maintain stable

protein gradients. One of the best studied examples is the membrane associated MinD

protein gradient in *Escherichia coli*. This oscillating gradient requires cycling of MinD

between a monomeric cytosolic and a dimeric membrane attached state. *Bacillus subtilis*

also has a Min system, but in this case MinD forms a static gradient. Mathematical models

have been successful in recreating the MinD gradient in *B. subtilis*, using an *E. coli*-like

membrane attachment cycle of MinD. Here we show that, in contrast to the *E. coli* situation,

the monomeric and dimeric state of *B. subtilis* MinD have comparable membrane affinities.

Using this and other information, we tested Monte Carlo simulations and found that a

difference in diffusion rate between MinD monomer and dimer, rather than a difference in

membrane affinity, is important for MinD gradient formation in *B. subtilis*.

INTRODUCTION

Bacterial cells are capable of positioning proteins at midcell and cell poles, and can form protein gradients without the support of specific membrane compartments or guidance by cytoskeleton elements, as found in eukaryotic cells. A well-known example is the formation of a MinCD protein gradient along the cell axis of rod-shaped *E. coli* and *B. subtilis* cells. The formation of these protein gradients is remarkable considering that the diffusion of proteins in cells is in the order of μm square per second, whereas these bacteria are only a few μm in length (1).

The Min system of *E. coli* is the best studied gradient system, and comprises the membrane associated MinD and MinE proteins that form a reaction-diffusion couple resulting in a MinCD gradient that oscillates between cell poles in a matter of seconds (2, 3). This system has been reconstructed on artificial membranes resulting in dynamic wave patterns (4), and extensively simulated to understand the reaction parameters (e.g. (5-11)). *B. subtilis* also forms a MinD gradient that decreases in concentration from cell poles and nascent cell division sites towards the middle of the cell. However, unlike *E. coli*, this gradient does not oscillate (12-14). In this study, we investigated how the MinCD gradient is formed in *B. subtilis*.

The core of the Min system consists of the protein couple MinC and MinD that prevent aberrant polymerization of the key cell division protein FtsZ close to newly formed septa or cell poles (14-16). MinC inhibits polymerization of FtsZ by direct protein-protein interactions and needs to bind to the Walker A-type ATPase MinD for its activity (17-20). In case of *E. coli* MinD, ATP binding leads to dimerization of MinD, which induces a conformational change in the C-terminal region, thereby exposing an amphipathic helix that

functions as a membrane binding domain (18, 21). The MinD dimer recruits MinC to the cell

membrane and stimulates its activity (21-23). The *E. coli* Min system comprises a third

protein, the peripheral membrane protein MinE, that interacts with MinD, thereby

displacing MinC and stimulating the ATPase activity of MinD, which ultimately triggers the

release of MinD from the membrane (22, 24-27). The interaction with MinD causes a

conformational change in MinE that stimulates its membrane affinity (24). These reciprocal

interactions of MinD and MinE represent a natural reaction-diffusion system, resulting in

the oscillation of MinCD proteins between cell poles with a periodicity of approximately 50

seconds (2-4). Through this oscillation, the average concentration of MinC is higher close to

cell poles and lower at midcell, thereby inhibiting polymerization of FtsZ close to cell poles

and favoring Z-ring formation at midcell (28).

The MinCD proteins of *E. coli* and *B. subtilis* are highly conserved. However, *B.*

subtilis does not encode a MinE homologue and instead requires the proteins DivIVA and

MinJ for the proper polar and septal localization of MinCD (29, 30). DivIVA is a general

scaffold protein conserved in Gram-positive bacteria that associates with strong negatively

curved membrane areas at the base of division septa, resulting in polarly located DivIVA

clusters after division is completed (31, 32). The integral membrane protein MinJ interacts

with DivIVA and shows the same localization pattern. MinJ also interacts with MinD, and

inactivation of MinJ eliminates a MinCD gradient, resulting in filamentous cells and minicell

formation due to uncontrolled activity of MinC (29, 30). Since DivIVA is recruited to midcell

in the early stages of septum synthesis when a strong concave membrane area is formed

(31), the Min proteins are already present when septum synthesis is ongoing, and their

primary activity it to inhibit Z-ring formation next to nascent division sites (14).

The ATP-driven cycle between membrane association and dissociation of MinD is

critical to the formation of the oscillating MinD gradient in *E. coli*. Mathematical modelling

has shown that such an ATP-driven reversible membrane attachment cycle can also explain

the formation of a MinD gradient in *B. subtilis* (33, 34). However, the C-terminal

amphipathic alpha helix of *B. subtilis* MinD has a much stronger affinity for the cell

membrane compared to its *E. coli* counterpart, and when fused to GFP can recruit this

protein to the cell membrane, which the *E. coli* MinD C-terminal membrane anchor is not

capable of (35). In fact, a MinD point mutant that prevents binding of ATP and maintains the

protein in its monomeric state is still recruited to the *B. subtilis* cell membrane (36). It is

therefore unclear whether dimerization-dependent changes in the membrane affinity of the

MinD are crucial for the formation of a *B. subtilis* MinCD gradient.

In this study, we analyzed and manipulated the membrane affinity of *B. subtilis*

MinD, and combined *in vivo* experiments with *in silico* Monte Carlo simulations to assess the

role of membrane affinity in the formation of a MinCD gradient. These experiments

demonstrate that, unlike in *E. coli*, a difference in membrane affinity between MinD

monomer and dimer is not critical for the creation of a MinCD gradient. In addition, we

examined how differences in MinD membrane affinity affect the recruitment of MinC to the

membrane, a process that is critical for the FtsZ-regulating activity of the Min-system.

RESULTS

Effect of dimerization on the localization of *B. subtilis* MinD

We began by examining how much the membrane affinity of monomeric *B. subtilis* MinD

differs from the dimeric form by constructing MinD variants that are fixed in one of these

conformations. Dimerization of MinD and other members of the MinD/ParA protein family

requires binding of ATP. The conserved lysine in the Walker A-type ATPase domain, amino

acid position 16 in the protein (Fig. S1), forms hydrogen bonds with the phosphate groups of

the nucleotide, as revealed in the crystal structure of *Pyrococcus furiosus* MinD (37).

Exchanging this residue into an alanine (K16A exchange) prevents ATP binding and

formation of dimers in several Walker motif-containing proteins, including *E. coli* MinD and

RecA, and *B. subtilis* ParA/Soj (26, 38-40). The conserved glycine at position 12 makes

contacts with the γ-phosphate of ATP across the dimer interface. It has been shown that

changing this residue into a valine in *B. subtilis* ParA and *E. coli* MinD causes a steric clash in

the active site, thereby preventing dimerization while retaining ATP binding (38, 40). These

two monomer-forcing amino acid exchanges (K16A and G12V) were introduced into *B.*

subtilis MinD, which was fused at the N-terminus with GFP to follow its localization in the

cell. It has been shown that a GFP-MinD fusion retains its biological activity (13), and the

fusion protein was expressed at levels that resulted in normal sized cells and the absence of

minicells (Fig. 1A). To prevent possible localization artefacts due to the weak dimerization

characteristics of GFP (41), we used a monomeric GFP (mGFP) variant. The different MinD

fusion proteins were expressed from the ectopic *amyE* locus, under control of the xylose-

inducible *Pxyl* promoter (42), in a strain lacking wild type *minD* (Δ*minD*), as well as in a strain

lacking *minD* and *minC* (Δ*minCD*). Western blot analysis showed that the different MinD-

variants were stable and expressed at levels comparable to that of wild type MinD (Fig. S2).

Induction of mGFP-MinD with 0.1 % xylose reduced minicell formation in a Δ*minD*

background to wild type levels (from 14.7 % to 0.07 %, n > 300). However, the mGFP-MinD-

K16A and -G12V variants were unable to prevent formation of minicells (11.2 % and 9.5 %,

respectively, n > 600).

Fig. 1B shows the effect of the K16A exchange on mGFP-MinD localization. In line

with previous results (36), preventing the binding of ATP abolishes the mGFP-MinD

concentration gradient, which becomes clearer in longitudinal fluorescence intensity

profiles (Fig. 1, right panels). The fluorescence peaks visible at midcell are caused by the

presence of two septal cell membranes layers, as indicated by the fluorescence intensity

profiles of a wild type cell stained with the fluorescent membrane dye FM 5-95 (Fig. 1G).

The monomeric ATP binding G12V variant shows the same absence of a protein

gradient as the K16A variant (Fig. 1C). In contrast to *E. coli* MinD (26), both monomeric *B.*

subtilis MinD mutants still bind to the cell membrane. To measure the membrane

association *in vivo*, transverse fluorescence intensity profiles were collected (Fig. 1, middle

panels), and the membrane affinities were estimated based on the ratio between valley

(cytoplasmic) and peak (membrane) signals. This confirmed that the different monomeric

mutants have the same membrane affinity as wild type MinD (Fig. 1F). As controls we used

the fluorescence membrane dye FM5-95 and cells expressing cytosolic GFP (Fig. 1E).

It has been shown that exchanging a conserved aspartic acid residue in the ATPase

domains of *B. subtilis* ParA and *E. coli* MinD blocks ATP hydrolysis and traps the proteins in a

dimeric state (38, 40). When this aspartic acid residue, located at amino acid position 40 in

the *B. subtilis* MinD sequence (Fig. S1), was replaced by an alanine in the mGFP-MinD fusion,

expression of this variant in a Δ*minD* strain resulted in highly filamentous cells with clusters

of mGFP-MinD-D40A formed along the membrane (Fig. 1D, left picture). Introduction of a

minC deletion restored cell division (Fig. 1D, right picture), indicating that the locked MinD

dimer causes hyper activation of MinC. Interestingly, in this Δ*minCD* background the clusters

disappeared and mGFP-MinD-D40A showed a clear polar and septal accumulation, although

the polar and septal signal intensities were approximately 60-70 % compared to that of wild

type cells (Fig. S3B), and the longitudinal protein gradient was clearly reduced (Fig. 1D).

Quantification of the membrane affinity showed that the membrane affinity of the D40A

variant is only slightly higher compared to the monomeric mutants (Fig. 1F).

Localization of trapped MinD dimer depends on MinJ

The polar and septal localization of the D40A trapped dimer-variant of MinD presumably

relies on the interaction with MinJ. It has been reported that the absence of MinJ results in

detachment of MinD from the membrane (29, 34). However, when we introduced the Δ*minJ*

in our Δ*minCD* deletion strain background we found that all MinD-GFP variants remained

attached to the membrane (Fig. S4), and the septal and polar localization of wild type and

the trapped dimer MinD disappeared. These results show that while MinJ is not required for

membrane attachment of MinD *per se,* it is required for polar and septal localization of

MinD through interactions specifically with the dimeric form of MinD.

Recruitment of MinC by the different MinD mutants

Several biochemical studies with different bacterial species, including *B. subtilis*, have shown

that MinC and MinD dimers can assemble into long polymers (43-46). However, the

assembly of large MinCD multimers has not been observed *in vivo*, and a genetic study

indicated that the formation of such polymers are not necessary for the function of the Min

system in *E. coli* (47). The D40A MinD variant showed a clustered membrane localization

pattern that resolved into a smooth membrane pattern when *minC* was deleted (Fig. 1D).

This change in localization pattern might be related to the formation of higher-order MinCD

assemblies. To confirm this, we introduced an active mCherry-MinC fusion to see whether

mGFP-MinD and mCherry-MinC would colocalize when the former is fixed in a dimeric state.

First, we examined the localization of MinC in the presence of monomeric K16A and G12V

MinD variants. As shown in Fig. 2B and C, the mCherry-MinC signal was dispersed

throughout the cytosol in these strains, indicting a lack of interaction with monomeric MinD

variants, in line with a previous report (36). In case of the mGFP-MinD-D40A variant, the

mCherry-MinC expression was only induced for 1 h to prevent excessive filamentation. As

shown in Fig. 2D, the mCherry-MinC signal showed foci along the membrane that often

colocalized with mGFP-MinD-D40A foci, confirming that MinD dimerization is crucial for

MinC interaction, and that both proteins can indeed form large assemblies in the cell,

presumably in the form of MinCD copolymers.

The colocalization of the MinD-D40A and MinC fluorescent fusion proteins enabled

us to investigate whether MinD dimerization is sufficient for MinC activity or whether

localization at the cell membrane is also required. To test this, the membrane binding

capacity of MinD was impaired by replacing isoleucine 260, located in the C-terminal

membrane targeting amphipathic alpha helix, with a glutamate residue (Fig. 2E) (48). Next,

we introduced this mutation in the mGFP-MinD-D40A mutant to create a trapped MinD

dimer that was no longer able to bind to the membrane. Expression of this cytosolic MinD

variant did not result in cell filamentation (Fig. 2F), indicating that interaction with MinD

dimers alone is not sufficient to inhibit FtsZ polymerization. Rather, MinC needs to be

localized and enriched to the membrane surface to be effective, which is not surprising

since Z-ring formation occurs at the membrane periphery.

Weak membrane interaction is important for MinD gradient formation

The absence of a clear difference in membrane affinity between monomeric and dimeric

MinD raised the question whether a reversable membrane binding is important for the

formation a protein gradient. To examine this, we tried to increase the membrane affinity of

MinD by adding an extra copy of its C-terminal membrane targeting amphipathic helix. The

extra amphipathic helix was connected to the original helix by a short 11 amino acid long

flexible linker (Fig. 3A). First, we tested whether such tandem amphipathic helix (2xAH)

increases the membrane affinity by fusing it to mGFP. As shown in Fig. 3, fusion of this

tandem helix to mGFP increased the membrane association of the latter considerably when

compared to a single MinD amphipathic helix.

Next, we replaced the membrane targeting amphipathic helix of mGFP-MinD by the

tandem amphipathic helix. Western blot analysis showed that this new C-terminus did not

affect the protein levels to a significant degree (Fig. S2). As shown in Fig. 4B, the tandem

helix notably increased the fluorescent membrane signal all along the cell membrane (Fig.

4D and S5), thereby reducing the longitudinal fluorescence gradients.

To confirm that the tandem amphipathic helix increased the membrane affinity of

MinD, we tested its sensitivity to membrane depolarization. Previously, it has been shown

that binding of the MinD amphipathic helix to the cell membrane requires the presence of

the membrane potential (48). Dissipation of the proton motive force by the specific proton-

ionophore carbonyl cyanide m-chlorophenylhydrazone (CCCP) did not affect the membrane

localization of the tandem amphipathic helix fusion, whereas CCCP quickly disturbed the

membrane attachment of wild type MinD (Fig. S6).

240   We also tested what happens with the MinD gradient when its membrane affinity is

241 reduced. For this, the well-studied membrane binding amphipathic helix 2 from the

242 Hepatitis C virus protein NS4B was chosen (49, 50), which is hardly able to attach mGFP to

243 the cell membrane (Fig. 3). Replacing the wild type membrane targeting helix of MinD by

244 the weak NS4B amphipathic helix reduced the membrane association of mGFP-MinD (Fig.

245 4D), as well as its septal and polar localization (Fig. 4C and Fig. S5). Interestingly, this mutant

246 still displayed a longitudinal protein gradient. Together these data suggests that a reversible

247 membrane association is important for the formation of a MinD gradient along the cell axis.

249 **Effect of MinD membrane affinity on MinC recruitment and activity**

250 The results shown in Fig. 2 indicated that the attachment of the MinD dimer to the

251 membrane is essential for the activity of MinC. When this attachment was strengthened by

252 using the tandem amphipathic helix, the respective strain regularly featured additional

253 septa resulting in minicells (Fig. 4B, red arrows), although the frequency was lower

254 compared to a Δ*minD* mutant (Fig. 5A). The strain also exhibited an elongated cell

255 phenotype that was comparable to that of the *minD* deletion strain (Fig. 5B). As shown in

256 Fig. 6A-B, the mGFP-MinD(2xAH) variant is capable of recruiting mCherry-MinC along the

257 whole cell membrane, which likely explains the elongated cell phenotype.

258   Weakening the membrane affinity of mGFP-MinD, by replacing the wild type

259 amphipathic helix with the NS4B amphipathic helix, also resulted in occasional minicell

260 formation, but again not as numerous as in a *minD* deletion strain (Fig. 5A). The cell length

261 distribution of the strain expressing the mGFP-MinD(NS4B-AH) variant was closer the wild

262 type strain (Fig. 5B). When mCherry-MinC was co-expressed in this strain background, also

263 the MinC fusion protein showed a clear fluorescent gradient along the cell axis (Fig. 6C).

Thus, a reversible association of MinD with the cell membrane seems to be critical to

prevent misplacing the FtsZ inhibiting activity of MinC along the cell axis. Both the mGFP-

MinD(2xAH) and mGFP-MinD(NS4B-AH) variants were, at least partially, able to prevent

minicell formation, presumably because they are still recruited to cell poles and cell division

sites through interaction with MinJ.

Membrane gradient formation in kinetic Monte Carlo simulations

The different experiments indicated that the formation of a MinCD gradient requires the

dynamic switching between monomer and dimer states of MinD (Fig. 1), and a reversible

MinD membrane interaction (Fig. 4). So far, *in silico* simulations of the *B. subtilis* Min

gradient have assumed a strong difference in membrane interaction between the

monomeric and dimeric state of MinD, which is based on properties of the *E. coli* Min

system (33, 34). However, our data suggests that for *B. subtilis* these affinities do not differ

much (Fig. 1). To investigate more closely whether a difference in membrane affinity

between monomer and dimer is required for the formation of a MinD gradient in *B. subtilis*,

we resorted to kinetic Monte-Carlo simulations. The Monte-Carlo algorithm attempts to

move one molecule at a time by a small amount in a three-dimensional space. The new

interaction energy experienced by the particle is then calculated and compared with its old

energy, and the move is subsequently accepted or rejected according to the standard

Metropolis test (51, 52). This procedure is used to guide the system to the correct

Boltzmann distribution in equilibrium. It has been shown that this Monte-Carlo dynamics

corresponds well with Brownian or molecular dynamics (53, 54), and can be used to

examine mechanisms underlying spatiotemporal protein gradients in cells, provided that the

moves are small enough and that the acceptance probability remains high (31).

MinD monomers were presented as spheres (hydrodynamic radius 2.5 nm) that

diffuse within and along a spherocylindrical cell geometry. The following rules were

implemented: (i) Diffusion along the membrane is 10-fold slower that through the

cytoplasm, and (ii) the dimer, due to its larger diameter, diffuses 2-fold slower compared to

the monomer in both environments, emanating from Stokes-Einstein law and a

hydrodynamic radius for the dimer which is double that of the monomer. (iii) Based on the

data from Fig. 2, we started out with a 10 % stronger membrane affinity for the dimer. (iv)

The membrane dwell time of monomers and dimers is 0.01 - 0.5 sec and during the

simulations approximately 70 % of MinD molecules reside at the membrane, approaching

the valley to peak values observed experimentally (Fig. 1F). (v) Dimerization and

monomerization are modelled as chemical reactions between a monomer and a dimer

state. Baseline rates are set such that the two states are in a 1:1 ratio, based on the

longitudinal fluorescence intensity profile in Fig. 2A, and the assumption that dimers are

located at the polar regions and monomers along the lateral wall. In our baseline model,

dimerization at the pole, membrane and in the cytoplasm is equally likely, and is modelled

in a simple way as a reaction turning a monomeric to a dimeric MinD at constant rate. (vi)

The transition from dimer to monomer, stimulated by ATP hydrolysis, occurs stochastically

with a half-life of approximately 1/sec, which is based on information from *E. coli* MinD (8).

(vii) MinD dimers that come in close proximity of MinJ, represented as the surface area at

the polar caps, will remain attached to MinJ for some time, so that approximately 25 % of

MinD dimers is associated with MinJ, reminiscent of the septal localization of the D40A

variant (Fig. S3). Simulation details are described in the material and methods.

When we applied these rules and ran the simulation, a strong accumulation of MinD

at the poles was observed, but no gradient (Fig. 7A). In this simulation, the chance that

MinD can form a dimer is the same for monomers in the cytoplasm and those that are

attached to the membrane. However, it is likely that dimer formation in the latter situation

is more likely since monomers attached to the membrane by their C-terminal amphipathic

helix will diffuse slower and therefore spend more time in close proximity and, importantly,

they have the same spatial orientation, which will facilitate dimerization. However, when we

repeated the simulation, and implemented that membrane attached monomers have a 2-

fold higher chance of forming dimers than monomers in the cytoplasm, the localization

pattern did not change (Fig. 7B).

Since MinJ interacts with MinD dimers, there is the possibility that MinJ actually

stimulates MinD dimerization. Interestingly, when this property was added to the

simulation, a clear MinD gradient emerged along the cell length, together with a strong

polar localization signal (Fig. 7C), a pattern that is in good agreement with the longitudinal

fluorescence intensity profiles observed *in vivo* (Fig. 1A). This pattern did not change when

we included the condition that only membrane attached dimers are able to revert to

monomers (not shown), as suggested by the stimulated ATPase activity of *E. coli* MinD in the

presence of liposomes (27). When we performed the simulation of Fig. 7C but now with the

assumption that MinJ only stimulates dimerization and does not bind MinD dimers, a MinD

gradient is still formed, although the strong accumulation at the polar caps was abolished

(Fig. 7D). Subsequently, we repeated the simulation of Fig. 7C under conditions whereby

there was no difference in the diffusion rate of monomer and dimer (Fig. 7E). In this case

the MinD gradient was completely lost, showing that the differential diffusion of monomers

and dimers is necessary to create a steady-state gradient.

Next, we tested the role of the membrane affinity in gradient formation. First, the

simulation of Fig. 7C was repeated without a difference between the membrane affinity of

the MinD monomer and dimer (Fig. 7F). Interestingly, under these conditions the MinD

protein gradient is still being formed. When we repeated the simulation of Fig. 7C and kept

the 10 % difference in membrane affinity between MinD monomer and dimer, but made

their membrane affinities stronger, the gradient was reduced (Fig. 7G), in good agreement

with our *in vivo* observation with a MinD variant containing a tandem membrane anchor

(Fig. 4B). Lastly, we tested what happens when the simulation of Fig. 7C was repeated with a

weaker membrane affinity of MinD. As shown in Fig. 7H, a clear gradient is formed, again in

line with what was found when the membrane targeting amphipathic helix of MinD was

replaced with the weaker amphipathic helix 2 from Hepatitis C virus protein NS4B (Fig. 4C).

In conclusion, our kinetic Monte Carlo simulations suggests that the formation of a

MinD gradient requires: (i) a reversible and not too strong interaction of MinD with the

membrane, (ii) a clear difference between the diffusion rates of monomer and dimer, (iii) an

ATPase driven dimer-monomer cycle, and (iv) the stimulation of dimerization at the cell

poles and division sites. Importantly, a difference in membrane affinity between monomer

and dimer is not required.

DISCUSSION

Protein gradients in bacteria

Due to their small size, proteins diffuse rapidly throughout bacterial cells, in a matter of

seconds, yet stable protein gradients can be found in a large variety of bacteria. What these

systems have in common, including the MinD gradient, is a considerably reduced diffusion

rate of the proteins involved, which allows stable protein concentration gradients to form.

The viscosity of the cell membrane is roughly 100 times larger compared to the cytoplasm

(55, 56), and the Min system uses attachment to the cell membrane to slow down diffusion.

Another approach to reduce diffusion is to bind to DNA, which occurs in the ParABS system

involved in chromosome segregation in the asymmetric bacterium *Caulobacter crescentus*

(57). This system consists of the ParA ATPase, which belongs to the same ATPase family as

MinD (58) and ParB that binds to *parS* DNA binding sites. The ParA dimers associate with

DNA nonspecifically whereby ParB stimulates ATP hydrolysis of ParA dimers, resulting in the

dissociation from DNA. The localized dissociation and subsequent increased diffusion rate

results in a ParA concentration gradient away from the *parS* locus (59). The cyanobacterium

Synechococcus elongatus uses the ParA-like protein McdA to position carboxysomes along

the longitudinal axis. In this system, the protein McdB, which interacts with carboxysomes,

stimulates McdA ATPase activity and its release from DNA, resulting in an oscillation of

McdA over the nucleoid (60). Recently it was described that *Staphylococcus aureus* trigger

factor (TF) forms a gradient by interacting with the cell division protein FtsK, resulting in an

increased TF concentration towards the septal region (61). How TF reduces its diffusion in *S.*

aureus cells is still unclear but might be a consequence of the fact that the chaperone

directly interacts with ribosomes (62), which diffuse much slower due to their size and

association with mRNA and nascent polypeptide chains.

Previous MinD gradient modelling studies

As mentioned in the introduction, there are many mathematical models describing the

oscillating MinD gradient of *E. coli* but so far only two mathematical models focused on the

MinD gradient of *B. subtilis* (33, 34). Both models make use of the reversible interaction of

MinD with the cell membrane. The first study, almost 20 years ago, modelled the

localization of DivIVA and MinD by a reaction-diffusion system in one spatial dimension (33).

At that time neither MinJ nor the localization mechanism of DivIVA were known. In this

early study both MinD and DivIVA were assumed to cycle between the membrane and the

cytosol, and the membrane attachment of DivIVA would depend on MinD, whereas the

membrane attachment of the latter would be stabilized at the cell poles by DivIVA.

However, we now know that DivIVA binds to the membrane by itself, and there is no

interaction between MinD and DivIVA (31).

The ATPase stimulated monomer-dimer cycle of MinD was not included in this early

model, but it was used in a recent study that employed a minimal reaction-diffusion system

in three-dimensional cellular space (34). In this model *B. subtilis* MinD cycles between the

cell membrane and cytosol, whereby the membrane detached MinD is in the ADP-bound

state and can only rebind after exchanging ADP for ATP, like in *E. coli*. By assuming that the

membrane recruitment rate is higher at the poles and septa due to the presence of MinJ or

due to a kinetic increase of diffusional impacts with the membrane due to the curved

geometry at poles, a steep MinD gradient could be simulated (34).

New data on MinD localization

Here, we have shown that monomeric MinD, both in its ATP-bound and unbound state, has

a membrane affinity that is comparable to that of the dimeric form of MinD. This shows a

clear difference with the *E. coli* situation, where dimerization of MinD causes a

conformational change of the C-terminal region enabling the amphipathic helix to insert into

the lipid bilayer (Hu & Lutkenhaus 2003). Apparently, the C-terminal amphipathic helix of *B.*

subtilis MinD is continuously exposed and capable to bind the membrane, thus increasing

affinity only by an additive effect upon dimerization.

We have also shown that MinJ is not required for the membrane recruitment of

MinD. We have no explanation for the differences between our findings and that of the

previous studies in which it has been shown that MinJ is needed for the membrane

localization of MinD (29, 34). However, since the membrane-targeting amphipathic alpha

helix of *B. subtilis* MinD is by itself sufficient to recruit GFP to the cell membrane, even in *E.*

coli, which does not contain a MinJ homologue (35, 48), it is difficult to envision how the

absence of MinJ would bring MinD to the cytosol.

In another study, it has been suggested that the C-terminal amphipathic helix of

MinD by itself is sufficient to recruit GFP to cell division sites, even in the absence of either

MinJ or DivIVA (63). However, as indicated in Fig. 1, the presence of two perpendicular-

observed membranes that are formed during septum synthesis gives a higher fluorescence

membrane signal, which might have been mistakenly interpreted as septal localization. In

addition, in our hands a GFP fusion with the C-terminal amphipathic helix of *B. subtilis* MinD

shows a clear fluorescence membrane stain but no accumulation at division sites (Fig. 3B).

Kinetic Monte Carlo model

Our *in vivo* findings warranted a new mathematical model to study the formation of the *B.*

subtilis MinD gradient. The kinetic Monte Carlo model showed that a membrane

attachment-detachment cycle is not required to form a protein gradient, but that a dynamic

ATP-driven cycle between monomer and dimer is crucial, coupled to the differences in

diffusion rates. We have shown that the MinD dimer associates with MinJ, but this by itself,

even when the interaction was reversible, was not enough to create a gradient. We ascribe

this to the fact that both monomers and dimers bind to the membrane with small difference

in affinities in our model, whereas in (34) the monomer state is not membrane-associated.

To establish a gradient in our simulations, we needed to include the requirement that MinJ

stimulates and/or stabilizes MinD dimerization, but this MinJ activity remains speculative. It

should be stressed that MinJ is not crucial for MinD dimerization, since a *minJ* mutant shows

a strong filamentous phenotype that can be suppressed by a *minC* deletion, indicating that

MinD dimerization, required for MinC activity, also occurs in the absence of MinJ.

Although a difference in membrane affinity between monomer and dimer is not

necessary to establish a gradient, the modelling shows that a transient and reversible

membrane binding of MinD is necessary. This was also confirmed by the reduced formation

of a gradient when the membrane affinity of MinD was increased by using a tandem C-

terminal amphipathic helix (Fig. 4). This can be explained by considering the emergent

length scale over which the gradient develops. As in reaction-diffusion models for

morphogenesis, this length scale is given by the square root of a diffusion coefficient over a

reaction rate. Our simulations suggest that a possible formula is given by $l \sim \sqrt{\frac{D_d - D_m}{k_m}}$, with

D_d and D_m the dimer and monomer diffusion (the relevant rate is that on the membrane),

and k_m the rate of conversion between dimers and monomers. Increasing the interaction

with the membrane leads to a decrease in diffusivity, and concomitantly a reduction in the

length scale over which the gradient can be observed.

In our modelling, MinJ is simply represented by the polar cap area. FRAP studies

have shown that the recruitment of MinJ to cell poles and division sites is quite dynamic

with a fluorescence recovery rate close to a minute (34). However, the fluorescence

recovery rate of MinD is approximate 8-fold higher (34). Because of this large difference in

diffusion rates, we have ignored the diffusion of MinJ in our modelling. We do not foresee

different results if we were to include MinJ diffusion in our Monte Carlo simulations,

provided that in steady state MinJ is on average localized at the poles.

Another aspect that we have not included in the simulations is MinD polymerization.

It has been shown that purified *E. coli* MinD can form large polymers in the presence of ATP

(64), although this was not found for purified *B. subtilis* MinD (46). FRAP experiments with

B. subtilis shows a fluorescence recover rate of seconds for MinD at division sites,

suggesting that if such polymers exist, they are not very stable. Nevertheless, any

multimerization of MinD dimer will further increase the diffusion difference between

monomer and dimer, thus contributing to the formation of a protein gradient.

In conclusion, we have shown here that a dynamic cyclic change in membrane

affinity of MinD is not required for the formation of a concentration gradient in the cell.

Instead, our modelling suggests that stimulation of MinD-dimerization at cell poles and cell

division sites is needed. It will be interesting to see whether future experiments proves this

prediction right or wrong.

MATERIAL AND METHODS

Bacterial strains, growth conditions and media

Luria-Bertani (LB) agar was used for the routine selection and maintenance of both *B.*

subtilis and *E. coli* strains. For *B. subtilis*, cells were grown in LB medium or Spizizen minimal

salt medium supplemented with 0.05 % yeast extract (SMMY). SMMY consists of 2 g/l

$(NH_4)_2SO_4$, 14 g/l K_2HPO_4, 6 g/l KH_2PO_4, 1 g/l sodium citrate, 2 g/l $MgSO_4$, 5 g/l fructose, 2 g/l

tryptophan, 0.2 g/l casamino acids and 2.2 g/l ammonium ferric citrate. All strains were

grown at 37 °C. The supplements were added as required: 100 µg/ml ampicillin, 5 µg/ml

chloramphenicol, 5 µg/ml kanamycin, 100 µg/ml spectinomycin, 10 µg/ml tetracycline and 1

µg/ml erythromycin. The *B. subtilis* strains used are listed in Table S1. The mutant strains

kindly provided by other labs were transformed into our laboratory strain to ensure that all

strains were isogenic.

Strain constructions

Molecular cloning, PCR and transformations were carried out by standard techniques.

Plasmids and oligonucleotides used in this study are listed in Table S2 and S3, respectively.

The *minC*, *minD*, *minCD* and *minJ* gene deletions were introduced from existing strains listed

in Table S1 by standard natural transformation (65). The *aprE*-integration vector containing

mCherry under an IPTG-inducible promoter (P*spac*) was constructed as follows. The

spectinomycin marker of pAPNC213 (66) was replaced with a chloramphenicol resistance

cassette (*cat*) using In-Fusion cloning (Clonetech). For this aim pAPNC123 was PCR-linearized

with oligonucleotide pair HS05/HS06 and the *cat* cassette was amplified from pSG2 (67)

with oligonucleotide pair HS07/HS08. These PCR products were fused together with In-

Fusion cloning, resulting in pAPNC*cat,* this plasmid was verified by sequencing. Next, the

mCherry encoding gene was amplified using oligonucleotide pair HS437/HS438 and plasmid

template pSS153 (68) as template. SalI and BamHI restriction sites were inserted into the

primers. The mCherry PCR product and the pAPNC*cat* were digested with SalI and BamHI

and ligated. The resulting plasmid was verified by sequencing and named pHJS112.

The IPTG-inducible mCherry-*minC* fusion was constructed as follows. A PCR fragment

containing *minC* was amplified with oligonucleotide pair LB1/LB2 and genomic DNA of the

wild type strain 168 as template. BamHI and EcoRI restriction sites were inserted into the

primers. The PCR product and the *aprE*-integration vector pHJS112 were digested with

BamHI and EcoRI restriction enzymes and ligated. The resulting plasmid pLB11 was verified

by sequencing and transformed into *B. subtilis* competent cells, resulting in strain LB31. The

aprE integration was verified by PCR, amplifying the genomic DNA with oligonucleotide pair

HS508/HS509 and sequencing.

The *amyE*-integration vector pSG1730 containing GFP fused to *minD* under a xylose-

inducible promoter (12) was used to exchange the *minD* residue glycine 12 to valine (G12V)

and aspartic acid 40 to alanine (D40A), by incorporation of the corresponding mutation

using the QuickChange site-directed mutagenesis reaction (Stratagene) and the

oligonucleotide pairs HS320/HS321 and HS322/HS323, respectively. The resulting plasmids

were verified by sequencing and named pHJS115 and pHJS116, respectively. The plasmid

pHJS113 containing GFP fused to *B. subtilis minD* coding sequence with the lysine 16 to

alanine (K16A) exchange under a xylose-inducible promoter was reported previously (48).

Plasmid pSG1729 (42), containing GFP under a xylose-inducible promoter, was used

to generate a construct with GFP fused to the native *minD* membrane targeting amphipathic

helix (KGMMAKIKSFFG). This domain was amplified from *B. subtilis* wild type strain 168 with

oligonucleotide pair FB136/FB149. EcoRI and BamHI restriction sites were inserted into the

primers. The PCR product and the plasmid pSG1729 were digested with EcoRI and BamHI

restriction enzymes and ligated. The new plasmid was verified by sequencing and named

pHJS117. Plasmid pHJS117 was used to generate a construct with GFP fused to two *minD*

amphipathic helices in tandem with a short linker of glycines in between (2xAH). This

tandem membrane targeting sequence was created by amplification with oligonucleotides

FB138/FB149 and pHJS117 as template. The primer FB138 carries a long overhang that

generates an extra MinD membrane targeting amphipathic helix fused to the existing

membrane targeting domain (KGMMAKIKSFFGVRS**GG**VLEEQNKGMMAKIKSFFG). EcoRI and

BamHI restriction sites were inserted into the primers. The PCR product and the plasmid

pSG1729 were digested with *Eco*RI and *Bam*HI restriction enzymes and ligated. The new

plasmid was verified by sequencing and named pHJS119.

Another construct with GFP fused to the NS4B amphipathic helix 2 from Hepatitis C

virus (NS4B-AH) was generated as follows. Plasmid pSG1729 was amplified with

oligonucleotides FB141/FB149. The primer FB141 carries a long overhang that generates the

NS4B-AH anchor (TVTQLLRRLHQWI). EcoRI and BamHI restriction sites were inserted into

the primers. The PCR product and the plasmid pSG1729 were digested with EcoRI and

BamHI restriction enzymes and ligated. The new plasmid was verified by sequencing and

named pHJS121.

We constructed two different *amyE*-integration plasmids with GFP fused to *minD*

with different membrane targeting sequences. The native *minD* amphipathic helix was

either replaced with two *minD* amphipathic helices in tandem with a short linker of glycines

in between (MinD2xAH) or with the Hepatitis C virus NS4B amphipathic helix 2 (MinDNS4B-

AH). The *minD* gene was amplified from the *B. subtilis* genome (strain 168) with

oligonucleotides pairs FB135/FB138 and FB135/FB141, respectively. BamHI and EcoRI

restriction sites were inserted into the primers. The primers FB138 and FB141 carry a long

overhang that generates the 2xAH and the NS4B-AH, respectively. The digested PCR

fragments were ligated with BamHI/EcoRI-linearized pSG1729, containing GFP under a

xylose-inducible promoter. The new plasmids were verified by sequencing and named

pHJS123 and pHJS125, respectively.

Plasmids pSG1730, pHJS113, pHJS115, pHJS116, pHJS123 and pHJS125 were used as

template for the QuickChange site-directed mutagenesis reaction (Stratagene) with

oligonucleotide pair HS410/HS411 to introduce the A206K exchange in the GFP coding

sequence to reduce protein dimerization (mGFP). The resulting plasmids were verified by

sequencing and named pLB21, pLB22, pLB23, pLB24, pLB49 and pLB50 respectively.

Plasmids pLB21 and pLB24 were used to further exchange the *minD* isoleucine 260 to

glutamic acid (I260E) by QuickChange site-directed mutagenesis reaction (Stratagene) with

oligonucleotide pair HS205/HS206. The resulting plasmids were verified by sequencing and

named pLB71 and pLB72, respectively. These constructs were integrated at the *amyE* locus

in different genetic backgrounds.

Western blot analysis

To detect the cellular levels of the different GFP-MinD variants the strains were induced

with 0.1 % xylose for three doublings. 2 ml of cultures, with comparable ODs were

centrifuged and the cell pellets flash frozen in liquid nitrogen. The pellets were then

resuspended in 200 µl of buffer containing 10 mM Tris-Hcl pH 8, 100 mM NaCl and 1 %

TritonX-100. The cells were lysed with 100 µg/ml lysozyme during 15 min incubation at

room temperature. Cell debris were spun down (20 min Eppendorf centrifuge, 14000 rpm,

4°C) and 10 µl of samples were loaded onto a 10 % SDS-PAGE followed by separation by

electrophoresis at 150 V for 60 min. The proteins were then transferred to a PVDF

membrane (Thermo Scientific) with transfer buffer (14.4 g/l glycine, 3 g/l Tris base and 15 %

methanol) using mini-wet electroblotting system (Bio-Rad) at 55 V for 130 min. The

transferred membrane was then blocked in blocking buffer (1 % skimmed milk powder in

TBST buffer 50 mM Tris, 150 mM NaCl and 0.05 % Tween-20) overnight and rinsed twice for

5 min in TBST. The proteins were probed with a 1:2000 dilution of rabbit anti-GFP primary

antibody (Invitrogen) in 10 ml TBST incubated 1 h, the membrane was washed 3x in TBST

and incubated in a 1:20 000 dilution of secondary antibody goat anti rabbit IR dye (Li-cor

Biosciences) in 10 ml TBST during 1 h. The immunoblot was developed by using an Odyssey®

imaging system (Li-cor Biosciences).

Microscopy

Microscopy was performed on an inverted fluorescence Nikon Eclipse Ti microscope

equipped with a CFI Plan Apochromat DM 100x oil objective, an Intensilight HG 130bW lamp

and C11440-22CU Hamamatsu ORCA camera. Images were analysed using ImageJ v.1.48d5

(National Institutes of Health) software and the plugin ObjectJ (69). To measure the

transverse fluorescence intensity profiles (FIP), line-scans perpendicular to the cell axis were

marked. The polar gradient areas, marked in red in figures, were manually assigned by

means of visual inspection of the fluorescence gradient of the FIPs. The membrane affinity

was estimated as the ratio valley/peaks of the transverse FIP, using an average of at least 30

cells per data set.

For microscopy, cells were grown to exponential phase by diluting 1:100 a starter

culture grown overnight in liquid medium (SMMY) into fresh medium, and allowing at least

three doublings before mounting on microscopy slides covered with a thin film of 1 %

agarose. 0.1 % xylose and 0.01 mM IPTG were used for the induction since these

concentrations complemented a Δ*minCD* deletion strain, preventing minicell formation.

Membranes were stained with the fluorescent membrane dye FM-95 (0.5 µg/ml final

concentration), incubated for 5 min prior to mounting onto microscope slides covered with

a thin film of 1 % agarose. Dissipation of the proton motive force (pmf) was carried out by

the addition of 100 µM carbonyl cyanide m-chlorophenylhydrazone (CCCP) for 10 min. CCCP

was dissolved in DMSO (0.1 % final concentration of DMSO). As control, cells were

incubated with 0.1 % DMSO.

Kinetic Monte Carlo simulations

To model the formation of a steady state gradient in *B. subtilis* wild-type cells and mutants,

kinetic Monte-Carlo simulation were performed. Kinetic Monte Carlo is a stochastic

algorithm which is known to be equivalent to molecular dynamics for sufficiently small local

moves (53, 54). The system simulated consisted in a collection of 1000 MinD proteins

enclosed inside a spherocylinder (aspect ratio 4, sizes 80 by 20 in simulation units)

modelling a *B. subtilis* cell. MinD proteins can be in one of two states, either monomeric (M)

or dimeric (D). The M and D states have both affinities for the membrane (the surface of the

spherocylinder). To model this, we introduce an attractive potential, a square well with

range 1 and strength e_M and e_D for the M and D state respectively. Given our experimental

results that the dimer is only weakly stickier than the monomer, we ran our simulations with

e_D = 1.1e_M=1 k_BT. Additionally, the D state, but not the M state, may interact with the caps

(poles) in the spherocylinders, with an extra interaction again represented by a square well,

with range 1 and strength e_C. This interaction simulates binding between dimeric MinD and

MinJ. Various cases were considered but results are reported, for simplicity for two cases

only: e_C=0 (no additional polar interaction), or e_C = 0.7 e_M. A third parameter which varies

between M and D is the diffusion coefficient: for simplicity, unless otherwise stated, we

assume this to scale inversely with molecular weight, so that the diffusivity of the dimer, D,

is half that of the monomer, M.

Key to the formation of the gradient is the fact that the reaction between the M and

D states of MinD is not in thermodynamic equilibrium, as it is linked to ATP hydrolysis. In our

model, D turns into M, due to ATP hydrolysis, at a constant rate k_{ATP}, in all cases. We

consider instead three cases for the dimerization reaction. In our baseline model the

conversion of M into D also occurred at the same rate in the cytoplasm, at the membrane

and at the pole. We have set this rate such that the M and D state are on average

comparable in steady state, as suggested by our experiments. We also considered the case

where dimerization at the membrane occurs 2-fold more often at the membrane, and a

third case where polar localization (through MinJ interaction) strongly stimulates

dimerization, so that the M to D transition occurs at a rate $k_{pole} \gg k_{ATP}$ at poles. When

considering polar stimulation, dimerization occurs at a rate k_{ATP}/10 in the cytosol; otherwise,

rates of dimerization and monomerization in the cytoplasm are equal. The diffusion of

monomeric MinD in the cytosol was set to 0.0017 in simulation units (corresponding to

approximately 1 micron square per second, see below), where the hydrolysis rate k_{ATP} was

set to 10^{-5} (corresponding to 1/s). When considering polar stimulation, we set k_{pole} such that

dimerization at the pole occurs with unit probability during a timestep (we have verified

though that the results are qualitatively similar down to at least a value of k_{pole} equal to 100

k_{ATP}). More details on the parameters used in the model are given in the caption of Fig. 7.

To simulate the monomer (K16A) and dimer (D40A) mutants, we used another

version of the same model, in which all particles were of type M or D respectively, and the

rate of interconversion between the two was set to 0.

Simulation units may be mapped to physical units by noting that 1 space and time

units in simulations can be made to correspond to 0.08 microns and 10 microseconds,

respectively.

ACKNOWLEDGEMENTS

We would like to thank especially Norbert Vischer for analysis and quantification of

fluorescence images and developing ObjectJ plugins, members of the Bacterial Cell Biology

group (University of Amsterdam) for insightful discussions, Dirk-Jan Scheffers for critical

reading of the manuscript, and Marc Bramkamp for prior communications. This research

was funded by EU Marie Curie ITN grants ATP-BCT (020496-2) and AMBER (317338), Marie

Curie CIG grant DIVANTI (618452) and STW Vici grant (12128).

REFERENCES
648
1. Kumar M, Mommer MS, Sourjik V. 2010. Mobility of cytoplasmic, membrane, and
DNA-binding proteins in *Escherichia coli*. Biophys J 98:552-9.
2. Hu Z, Lutkenhaus J. 1999. Topological regulation of cell division in *Escherichia coli*
involves rapid pole to pole oscillation of the division inhibitor MinC under the control
of MinD and MinE. Mol Microbiol 34:82-90.
3. Raskin DM, de Boer PA. 1999. Rapid pole-to-pole oscillation of a protein required for
directing division to the middle of *Escherichia coli*. Proc Natl Acad Sci USA 96:4971-6.
4. Loose M, Fischer-Friedrich E, Ries J, Kruse K, Schwille P. 2008. Spatial regulators for
bacterial cell division self-organize into surface waves in vitro. Science 320:789-92.
5. Howard M, Rutenberg AD, de Vet S. 2001. Dynamic compartmentalization of
bacteria: accurate division in *E. coli*. Phys Rev Lett 87:278102.
6. Meinhardt H, de Boer PA. 2001. Pattern formation in *Escherichia coli*: a model for the
pole-to-pole oscillations of Min proteins and the localization of the division site. Proc
Natl Acad Sci U S A 98:14202-7.
7. Kruse K. 2002. A dynamic model for determining the middle of *Escherichia coli*.
Biophys J 82:618-27.
8. Huang KC, Meir Y, Wingreen NS. 2003. Dynamic structures in *Escherichia coli*:
spontaneous formation of MinE rings and MinD polar zones. Proc Natl Acad Sci U S A
100:12724-8.
9. Di Ventura B, Sourjik V. 2011. Self-organized partitioning of dynamically localized
proteins in bacterial cell division. Mol Syst Biol 7:457.
10. Wurthner L, Brauns F, Pawlik G, Halatek J, Kerssemakers J, Dekker C, Frey E. 2022.
Bridging scales in a multiscale pattern-forming system. Proc Natl Acad Sci U S A
119:e2206888119.
11. Glock P, Brauns F, Halatek J, Frey E, Schwille P. 2019. Design of biochemical pattern
forming systems from minimal motifs. Elife 8.
12. Marston AL, Thomaides HB, Edwards DH, Sharpe ME, Errington J. 1998. Polar
localization of the MinD protein of *Bacillus subtilis* and its role in selection of the
mid-cell division site. Genes Dev 12:3419-30.
13. Marston AL, Errington J. 1999. Selection of the midcell division site in *Bacillus subtilis*
through MinD-dependent polar localization and activation of MinC. Mol Microbiol
33:84-96.
14. Gregory JA, Becker EC, Pogliano K. 2008. *Bacillus subtilis* MinC destabilizes FtsZ-rings
at new cell poles and contributes to the timing of cell division. Genes Dev 22:3475-
88.
15. Levin PA, Shim JJ, Grossman AD. 1998. Effect of minCD on FtsZ ring position and
polar septation in *Bacillus subtilis*. J Bacteriol 180:6048-51.
16. Bi E, Lutkenhaus J. 1993. Cell division inhibitors SulA and MinCD prevent formation
of the FtsZ ring. J Bacteriol 175:1118-25.
17. de Boer PA, Crossley RE, Rothfield LI. 1989. A division inhibitor and a topological
specificity factor coded for by the minicell locus determine proper placement of the
division septum in *E. coli*. Cell 56:641-9.
18. de Boer PA, Crossley RE, Hand AR, Rothfield LI. 1991. The MinD protein is a
membrane ATPase required for the correct placement of the *Escherichia coli* division
site. EMBO J 10:4371-80.

19. Hu Z, Lutkenhaus J. 2000. Analysis of MinC reveals two independent domains
involved in interaction with MinD and FtsZ. J Bacteriol 182:3965-71.
20. Cordell SC, Anderson RE, Lowe J. 2001. Crystal structure of the bacterial cell division
inhibitor MinC. EMBO J 20:2454-61.
21. Hu Z, Lutkenhaus J. 2003. A conserved sequence at the C-terminus of MinD is
required for binding to the membrane and targeting MinC to the septum. Mol
Microbiol 47:345-55.
22. Hu Z, Saez C, Lutkenhaus J. 2003. Recruitment of MinC, an inhibitor of Z-ring
formation, to the membrane in *Escherichia coli*: role of MinD and MinE. J Bacteriol
185:196-203.
23. de Boer PA, Crossley RE, Rothfield LI. 1992. Roles of MinC and MinD in the site-
specific septation block mediated by the MinCDE system of *Escherichia coli*. J
Bacteriol 174:63-70.
24. Park KT, Villar MT, Artigues A, Lutkenhaus J. 2017. MinE conformational dynamics
regulate membrane binding, MinD interaction, and Min oscillation. Proc Natl Acad
Sci U S A 114:7497-7504.
25. Hu Z, Lutkenhaus J. 2001. Topological regulation of cell division in E. coli.
spatiotemporal oscillation of MinD requires stimulation of its ATPase by MinE and
phospholipid. Mol Cell 7:1337-43.
26. Hu Z, Gogol EP, Lutkenhaus J. 2002. Dynamic assembly of MinD on phospholipid
vesicles regulated by ATP and MinE. Proc Natl Acad Sci USA 99:6761-6.
27. Lackner LL, Raskin DM, de Boer PA. 2003. ATP-dependent interactions between
Escherichia coli Min proteins and the phospholipid membrane *in vitro*. J Bacteriol
185:735-49.
28. Rowlett VW, Margolin W. 2013. The bacterial Min system. Curr Biol 23:R553-6.
29. Bramkamp M, Emmins R, Weston L, Donovan C, Daniel RA, Errington J. 2008. A novel
component of the division-site selection system of *Bacillus subtilis* and a new mode
of action for the division inhibitor MinCD. Mol Microbiol 70:1556-69.
30. Patrick JE, Kearns DB. 2008. MinJ (YvjD) is a topological determinant of cell division in
Bacillus subtilis. Mol Microbiol 70:1166-79.
31. Lenarcic R, Halbedel S, Visser L, Shaw M, Wu LJ, Errington J, Marenduzzo D, Hamoen
LW. 2009. Localisation of DivIVA by targeting to negatively curved membranes.
EMBO J 28:2272-2282.
32. Ramamurthi KS, Losick R. 2009. Negative membrane curvature as a cue for
subcellular localization of a bacterial protein. Proc Natl Acad Sci USA 106:13541-5.
33. Howard M. 2004. A mechanism for polar protein localization in bacteria. J Mol Biol
335:655-63.
34. Feddersen H, Wurthner L, Frey E, Bramkamp M. 2021. Dynamics of the *Bacillus
subtilis* Min System. mBio 12.
35. Szeto TH, Rowland SL, Habrukowich CL, King GF. 2003. The MinD membrane
targeting sequence is a transplantable lipid-binding helix. J Biol Chem 278:40050-6.
36. Karoui ME, Errington J. 2001. Isolation and characterization of topological specificity
mutants of minD in Bacillus subtilis. Mol Microbiol 42:1211-21.
37. Hayashi I, Oyama T, Morikawa K. 2001. Structural and functional studies of MinD
ATPase: implications for the molecular recognition of the bacterial cell division
apparatus. EMBO J 20:1819-28.

38. Wu W, Park KT, Holyoak T, Lutkenhaus J. 2011. Determination of the structure of the
MinD-ATP complex reveals the orientation of MinD on the membrane and the
relative location of the binding sites for MinE and MinC. Mol Microbiol 79:1515-28.
39. Story RM, Steitz TA. 1992. Structure of the recA protein-ADP complex. Nature
355:374-6.
40. Leonard TA, Butler PJ, Lowe J. 2005. Bacterial chromosome segregation: structure
and DNA binding of the Soj dimer - a conserved biological switch. Embo J.
41. Landgraf D, Okumus B, Chien P, Baker TA, Paulsson J. 2012. Segregation of molecules
at cell division reveals native protein localization. Nat Methods 9:480-2.
42. Lewis PJ, Marston AL. 1999. GFP vectors for controlled expression and dual labelling
of protein fusions in *Bacillus subtilis*. Gene 227:101-10.
43. Ghosal D, Trambaiolo D, Amos LA, Lowe J. 2014. MinCD cell division proteins form
alternating copolymeric cytomotive filaments. Nat Commun 5:5341.
44. Conti J, Viola MG, Camberg JL. 2015. The bacterial cell division regulators MinD and
MinC form polymers in the presence of nucleotide. FEBS Lett 589:201-6.
45. Huang H, Wang P, Bian L, Osawa M, Erickson HP, Chen Y. 2018. The cell division
protein MinD from *Pseudomonas aeruginosa* dominates the assembly of the MinC-
MinD copolymers. J Biol Chem 293:7786-7795.
46. Wang N, Zhang T, Du S, Zhou Y, Chen Y. 2022. How Do MinC-D Copolymers Act on Z-
Ring Localization Regulation? A New Model of *Bacillus subtilis* Min System. Front
Microbiol 13:841171.
47. Park KT, Du S, Lutkenhaus J. 2015. MinC/MinD copolymers are not required for Min
function. Mol Microbiol 98:895-909.
48. Strahl H, Hamoen LW. 2010. Membrane potential is important for bacterial cell
division. Proc Natl Acad Sci USA 107:12281-6.
49. Gouttenoire J, Montserret R, Kennel A, Penin F, Moradpour D. 2009. An amphipathic
alpha-helix at the C terminus of hepatitis C virus nonstructural protein 4B mediates
membrane association. J Virol 83:11378-84.
50. Adams DW, Wu LJ, Errington J. 2015. Nucleoid occlusion protein Noc recruits DNA to
the bacterial cell membrane. EMBO J 34:491-501.
51. Allen MP, Tildesley DJ. 1987. Computer simulations of liquids.
52. Binder K, Heermann DW. 2002. Monte Carlo Simulation in Statistical Physics: An
Introduction, 4th ed, vol 80.
53. Whitelam S, Geissler PL. 2007. Avoiding unphysical kinetic traps in Monte Carlo
simulations of strongly attractive particles. J Chem Phys 127:154101.
54. Sanz E, Marenduzzo D. 2010. Dynamic Monte Carlo versus Brownian dynamics: A
comparison for self-diffusion and crystallization in colloidal fluids. J Chem Phys
132:194102.
55. Mika JT, Thompson AJ, Dent MR, Brooks NJ, Michiels J, Hofkens J, Kuimova MK. 2016.
Measuring the Viscosity of the *Escherichia coli* Plasma Membrane Using Molecular
Rotors. Biophys J 111:1528-1540.
56. Mullineaux CW, Nenninger A, Ray N, Robinson C. 2006. Diffusion of green
fluorescent protein in three cell environments in *Escherichia coli*. J Bacteriol
188:3442-8.
57. Ptacin JL, Lee SF, Garner EC, Toro E, Eckart M, Comolli LR, Moerner WE, Shapiro L.
2010. A spindle-like apparatus guides bacterial chromosome segregation. Nat Cell
Biol 12:791-8.

58. Leipe DD, Wolf YI, Koonin EV, Aravind L. 2002. Classification and evolution of P-loop
GTPases and related ATPases. J Mol Biol 317:41-72.
59. Surovtsev IV, Lim HC, Jacobs-Wagner C. 2016. The Slow Mobility of the ParA
Partitioning Protein Underlies Its Steady-State Patterning in Caulobacter. Biophys J
110:2790-2799.
60. MacCready JS, Hakim P, Young EJ, Hu L, Liu J, Osteryoung KW, Vecchiarelli AG, Ducat
DC. 2018. Protein gradients on the nucleoid position the carbon-fixing organelles of
cyanobacteria. Elife 7.
61. Veiga H, Jousselin A, Schaper S, Saraiva BM, Marques LB, Reed P, Wilton J, Pereira
PM, Filipe SR, Pinho MG. 2023. Cell division protein FtsK coordinates bacterial
chromosome segregation and daughter cell separation in *Staphylococcus aureus*.
EMBO J doi:10.15252/embj.2022112140:e112140.
62. Deeng J, Chan KY, van der Sluis EO, Berninghausen O, Han W, Gumbart J, Schulten K,
Beatrix B, Beckmann R. 2016. Dynamic Behavior of Trigger Factor on the Ribosome. J
Mol Biol 428:3588-602.
63. Ishikawa K, Matsuoka S, Hara H, Matsumoto K. 2017. Septal membrane localization
by C-terminal amphipathic alpha-helices of MinD in *Bacillus subtilis* mutant cells
lacking MinJ or DivIVA. Genes Genet Syst 92:81-98.
64. Suefuji K, Valluzzi R, RayChaudhuri D. 2002. Dynamic assembly of MinD into filament
bundles modulated by ATP, phospholipids, and MinE. Proc Natl Acad Sci U S A
99:16776-81.
65. Hamoen LW, Smits WK, de Jong A, Holsappel S, Kuipers OP. 2002. Improving the
predictive value of the competence transcription factor (ComK) binding site in
Bacillus subtilis using a genomic approach. Nucleic Acids Res 30:5517-28.
66. Morimoto T, Loh PC, Hirai T, Asai K, Kobayashi K, Moriya S, Ogasawara N. 2002. Six
GTP-binding proteins of the Era/Obg family are essential for cell growth in *Bacillus*
subtilis. Microbiology 148:3539-52.
67. Fort P, Errington J. 1985. Nucleotide sequence and complementation analysis of a
polycistronic sporulation operon, *spoVA*, in *Bacillus subtilis*. J Gen Microbiol
131:1091-105.
68. Syvertsson S, Wang B, Staal J, Gao Y, Kort R, Hamoen LW. 2021. Different Resource
Allocation in a *Bacillus subtilis* Population Displaying Bimodal Motility. J Bacteriol
203:e0003721.
69. Vischer NO, Verheul J, Postma M, van den Berg van Saparoea B, Galli E, Natale P,
Gerdes K, Luirink J, Vollmer W, Vicente M, den Blaauwen T. 2015. Cell age dependent
concentration of *Escherichia coli* divisome proteins analyzed with ImageJ and
ObjectJ. Front Microbiol 6:586.
824
825

Fig. 1. Localization of monomeric and dimeric MinD variants

Cellular localization of (A) wild type MinD, (B) MinD K16A (apo monomer), (C) MinD G12V

(ATP-bound monomer), and (D) MinD D40A (ATP-bound dimer, ATP hydrolysis deficient).

Localization was monitored by an N-terminal mGFP fusion. Fusion proteins were expressed

at the ectopic *amyE* locus, under control of a xylose-inducible promoter in either a Δ*minD* or

Δ*minCD* background. Fluorescence images (left panels) and corresponding phase contrast

images (inset) are shown in the left panels. Some minicells are indicated with red arrows.

Scale bar is 2 μm. Middle panels show the transverse fluorescence intensity profiles (FIP)

calculated using an average of at least 30 cells (Δ*minCD* background) per data set. Right

panels depict the longitudinal fluorescence intensity profiles (FIP) using the Δ*minCD*

background. The manually shaded red areas highlight the polar gradients. Additional

examples for wild type MinD and the D40A variants are shown in Fig. S3. (E) Transversal

fluorescence intensity profile (FIP) of exponentially growing wild-type cells stained with

fluorescence membrane dye FM5-95, and wild-type cells expressing GFP are shown as

controls. (F) Membrane affinities estimated from the valley/peak ratios shown in the middle

panels of (A-D) and controls (E). (G) Longitudinal fluorescence intensity profile (FIP) along

the exponentially growing wild-type cells stained with the fluorescence membrane dye

FM5-95. Strains used in (A): LB249 and LB305, (B): LB250 and LB306, (C): LB251 and LB307,

(D): LB252 and LB308, (E): LB609 and (G) 168.

Fig. 2. Membrane recruitment of MinC by MinD variants

Fluorescence microscopy images of cells expressing different mGFP–MinD variants (cyan) and mCherry–MinC (red). Corresponding phase contrast images shown in the insets. (A)

Wild type MinD, (B) MinD K16A (apo monomer), (C) MinD G12V (ATP-bound monomer), (D)

MinD D40A (ATP-bound fixed dimer), (E) MinD I260E (non-membrane binding), (F) MinD

D40A, I260E (non-membrane binding ATP-bound fixed dimer). Right panels show the

transverse fluorescence intensity profiles (FIP) averaged over at least 30 cells. White arrows

in (D) highlights colocalization. Scale bar is 2 μm. mGFP–MinD variants were expressed in a

Δ*minCD* background strain from the ectopic *amyE* locus under a xylose-inducible promoter

(0.1 % xylose), and mCherry–MinC was expressed from the ectopic *aprE* locus under an

IPTG-inducible promoter (0.01 mM IPTG). Strains used: (A) LB318, (B) LB319, (C) LB320, (D)

LB321. (E) LB643 and (F) LB644.

Fig. 3. Membrane association of different amphipathic helices

(A) Schematic presentation of the tandem amphipathic helix and the weak amphipathic helix from Hepatitis C virus protein NS4B$_{241-253}$. (B) Fluorescence microscopy images and transverse fluorescence intensity profiles of cells expressing the different amphipathic helix

864 sequences fused to the C-terminus of GFP. A strain expressing cytoplasmic GFP was included

865 for comparison. Scale bar is 2 μm. Strains used: FBB043 (GFP-AH$_{MinD}$), FBB05 (GFP-AH$_{2x}$),

866 FBB046 (GFP-AH$_{NS4B}$), LB609 (GFP). (C) Average membrane affinities calculated from the

867 transverse fluorescence intensity profiles ($n > 30$).

Fig. 4. Membrane affinity affects MinD gradient

Fluorescence microscopy and fluorescence intensity profiles (right panels) of cells

expressing mGFP-MinD containing either the native amphipathic helix (wt) (A), the tandem

MinD amphipathic helix (2xAH) (B) or the weak Hepatitis C virus protein NS4B amphipathic

helix (NS4B-AH) (C). The manually shaded red areas highlight the polar gradients. Additional

examples are shown in Fig. S5. (D) Relative membrane affinities of the different mGFP-MinD

variants calculated from transverse fluorescence intensity profiles ($n > 30$). The MinD

variants were expressed in a Δ*minD* background from the ectopic *amyE* locus under control

of a xylose-inducible promoter (0.1 % xylose). Phase contrast images are shown as insets.

Some double septa are indicated with red arrows. Scale bar is 2 μm. Red areas in the

intensity profiles highlight the polar gradients. Strains used in (A) LB249, in (B) LB507, in (C)

LB508.

Fig. 5. Functionality of MinD with different membrane affinities

(A) Minicell formation in cells expressing mGFP-MinD with different membrane affinities.

The MinD variants were expressed in a Δ*minD* background from the ectopic *amyE* locus

under regulation of a xylose-inducible promoter (*n* > 300). (B) Cell length distributions of the

different strains (*n* > 300). Strains used: 1901 (Δ*minD*), LB249 (wild type MinD), LB507

(2xAH) and LB508 (NS4B-AH).

Fig. 6. Increased MinD membrane affinity affects MinC recruitment

Fluorescence microscopy images of cells expressing mGFP-MinD (cyan) and mCherry-MinC (red), with either the native membrane anchor (A), the tandem amphipathic helix (B), or the weak Hepatitis C virus protein NS4B derived amphipathic helix (C). Corresponding phase contrast images are shown in insets. Scale bar is 2 μm. Transverse fluorescence intensity profiles are shown in the right panels ($n > 30$). Right panels depict the longitudinal fluorescence intensity profiles (FIP) with manually shaded red areas to highlight the polar gradients. Additional examples are shown in Fig. S7. The mGFP-MinD variants were expressed at the ectopic *amyE* locus from a xylose-inducible promoter (0.1 % xylose), and mCherry-MinC was expressed at the ectopic *aprE* locus regulated by an IPTG-inducible promoter (0.01 mM IPTG) in a Δ*minCD* background strain. Strains used: (A) LB318, (B)

900   LB584, (C) LB559.

902
903
Fig. 7. Kinetic Monte Carlo simulations of MinD localization

Whole-cell kinetic Monte-Carlo simulations employing the following conditions. (A) Start
situation whereby (i) diffusion along the membrane is 10-fold slower compared to
cytoplasm, (ii) MinD dimer diffuses 2-fold slower compared to the monomer in both
environments, (iii) 10 % stronger membrane affinity for dimer, (iv) membrane dwell time of
monomers and dimers on average 1.4-4.5 sec, (v) dimerization and monomerization rates
such that dimers and monomers are approximately in a 1:1 ratio, (vi) transition from dimer
to monomer occurs stochastically with a half-life of approximately 1/sec, and (vii) MinD
dimers in close proximity of polar regions (peak MinJ concentration) will remain attached
for some time, so that approximately 25 % of MinD dimers is associated with the polar caps,
representing MinJ. (B) Same as simulation A but membrane attached monomers have a 2-
fold higher chance of forming dimers compared to cytoplasmic monomers. (C) Same as
simulation A but MinJ also stimulates MinD dimerization. (D) Same as simulation C but MinD
dimers are not retained by MinJ. (E) Same as simulation C but diffusion rates of monomer

and dimer are the same. (F) Same as simulation C but the membrane affinity of MinD

monomers and dimers is the same. (G) Same as simulation C but membrane affinity of MinD

is stronger (such that the diffusion is a further 10-fold slower on the membrane). (H) Same

as simulation C but membrane affinity of MinD is 2-fold weaker. Graphs indicate the average

lateral projection of MinD in simulated cells.