Title: **A human-ACE2 knock-in mouse model for SARS-CoV-2 infection recapitulates**
respiratory disorders but avoids neurological disease associated with the transgenic K18-
hACE2 model.

Running title: B.1 SARS-CoV-2 infection of a new hACE-2 Knock-in mice

Anna Pons-Grífols[1], Ferran Tarrés-Freixas[1,2,3,4], Mònica Pérez[2,3], Eva Riveira-Muñoz[1], Dàlia
Raïch-Regué[1], Daniel Pérez-Zsolt[1], Jordana Muñoz-Basagoiti[1], Barbara Tondelli[5], Edwards
Pradenas[1], Nuria Izquierdo-Useros[1,6,7,], Sara Capdevila[8], Júlia Vergara-Alert[2,3], Victor Urrea[1],
Jorge Carrillo[1,6,7], Ester Ballana[1,6], Stephen Forrow[5], Bonaventura Clotet[1,4], Joaquim
Segalés[2,9], Benjamin Trinité[1,#], Julià Blanco[1,4,6,7,#].

[1]IrsiCaixa, Can Ruti Campus, Badalona, Spain.
[2]Unitat mixta d'investigació IRTA-UAB en Sanitat Animal, Centre de Recerca en Sanitat
Animal (CReSA), Campus de la Universitat Autònoma de Barcelona (UAB), Bellaterra,
Spain.
[3]IRTA, Programa de Sanitat Animal, Centre de Recerca en Sanitat Animal (CReSA), Campus
de la Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain.
[4]University of Vic-Central University of Catalonia (UVic-UCC), Vic, Spain.
[5]Mouse Mutant Core Facility (PBA2B2), Institute for Research in Biomedicine (IRB), Parc
Científic de Barcelona, Barcelona, Spain.
[6]Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain.
[7]CIBER Infectious Diseases (CIBERINFEC), Carlos III Health Institute, Madrid, Spain.
[8]Comparative Medicine and Bioimage Centre of Catalonia (CMCiB), Germans Trias i Pujol
Research Institute (IGTP), Badalona, Spain.
[9]Departament de Sanitat i Anatomia Animals, Facultat de Veterinària, Universitat
Autònoma de Barcelona (UAB), Campus de la UAB, Bellaterra, Spain.

 # Corresponding authors

Benjamin Trinité: btrinite@irsicaixa.es

Julià Blanco: jblanco@irsicaixa.es

JS, BT and JB are co-senior authors

Abstract Word count: 242
Text word count: 5984
33

Abstract

Animal models have been instrumental in elucidating the pathogenesis of SARS-CoV-2

infection and testing COVID-19 vaccines and therapeutics. Wild-type (WT) mice are not

susceptible to many SARS-CoV-2 variants, therefore transgenic K18-hACE2 mice have

emerged as a standard model system. However, this model is characterized by severe

disease, particularly associated with neuroinfection, which leads to early humane endpoint

euthanasia. Here, we established a novel knock-in (KI) mouse model by inserting the original

K18-hACE2 transgene into the collagen COL1A1 locus using a recombinase mediated

cassette exchange (RMCE) system. Once the Col1a1-K18-hACE2 mouse colony was

established, animals were challenged with a B.1 SARS-CoV-2 (D614G) isolate and were

monitored for up to 14 days. Col1a1-K18-hACE2 mice exhibited an initial weight loss similar

to the K18-hACE2 transgenic model but did not develop evident neurologic clinical signs.

The majority of Col1a1-K18-hACE2 mice did not reach the preestablished humane endpoint,

showing progressive weight gain after 9 days post-infection (dpi). Importantly, despite this

apparent milder pathogenicity of the virus in this mouse model compared to the K18-hACE2

transgenic model, high levels of viral RNA were detected in lungs, oropharyngeal swab, and

nasal turbinate. Remaining lesions and inflammation in lungs were still observed after 14

dpi. In contrast, although low level viral RNA could be detected in a minority of Col1a1-K18-

hACE2 animals, no brain lesions were observed at any timepoint. Overall, Col1a1-K18-

hACE2 mice constitute a new model for investigating SARS-CoV-2 pathogenesis and

treatments, with potential implications for studying long-term COVID-19 sequelae.

Importance

K18-hACE2 mice express high levels of the human protein ACE2, the receptor for SARS-CoV-2, and can therefore be infected by this virus. These animals have been crucial to understand viral pathogenesis and to test COVID-19 vaccines and antiviral drugs. However, K18-hACE2 often die after infection with initial SARS-CoV-2 variants likely due to a massive brain infection that does not occur in humans. Here, we used a technology known as knock-in that allows for the targeted insertion of a gene into a mouse and we have generated a new hACE2-mouse. We have characterized this new animal model demonstrating that the virus replicates in the respiratory tract, damaging lung tissue and causing inflammation. In contrast to K18-hACE2 mice, only limited or no brain infection could be detected, and most animals recovered from infection with remaining lung lesions. This new model could be instrumental for the study of specific disease aspects such as post-COVID condition, sequelae, and susceptibility to reinfection.

Keywords: animal model, lung inflammation, lung damage, neuroinvasion, post covid condition.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is the etiologic agent of the coronavirus disease 2019 (COVID-19). The COVID-19 pandemic fueled an unprecedented collaborative research effort to develop therapeutics and vaccines. Animal models that recapitulate key clinical and pathological features of COVID-19 have played a pivotal role in testing novel vaccines, antivirals and other treatments (1).

The spike glycoprotein of SARS-CoV-2 uses the angiotensin-converting enzyme 2 (ACE2) as receptor to enter and infect host cells (2). ACE2 is expressed on the surface of cells of different organs such as lungs, intestines, kidneys, and heart. However, the spike glycoprotein from ancestral SARS-CoV-2 strains does not efficiently bind to mouse ACE2 (mACE2), rendering wild-type (WT) mice refractory to infection. Therefore, several strategies were used to develop mouse models susceptible to infection. These strategies include viral adaptation (3) or the introduction of the human ACE2 (hACE2) receptor via viral vectors and transgenic approaches (4–8).

Both K18-hACE2 transgenic mice and Golden Syrian hamsters (GSH) became reference animal models for investigating SARS-CoV-2 pathogenesis in vivo. They mainly differ in pathogenesis: GSH, which are naturally susceptible to SARS-CoV2 infection, recapitulate a milder disease phenotype (1). The K18-hACE2 mouse model was originally developed for the study of SARS-CoV, which also targets hACE2. It was created by random insertion of multiple copies of the *hACE2* gene under the control of the human cytokeratin 18 gene promoter (*KRT18* or K18). This promoter allows for high-level expression and is specific to epithelial cells, including those in the airways (9, 10). This approach allows the infection of

mice by SARS-CoV-2, while maintaining the activity of the mACE2 receptor. Infection of K18-

hACE2 mice with pre-Omicron variants of SARS-CoV-2 results in progressive weight-loss and

strong clinical signs by 3-5 days post-infection (dpi), a severity which often requires

euthanasia by 5-7 dpi. The lethality in this model is dose- and SARS-CoV-2 variant-

dependent (10, 11), and has been associated with neuroinvasion and extensive brain

infection (9, 12–14). Neuroinvasion in K18-hACE2 transgenic mice has been linked to the

aleatory gene insertion and the high number of inserted gene copies, around 8 in the

commercial *B6.Cg-Tg(K18-ACE2)2Prlmn/J-K18-hACE2 mouse* (10), leading to altered

expression levels and tissue distribution of the receptor (15). Specifically, a higher number

of gene copies correlates with a worse disease prognosis and encephalitis in this model,

following SARS-CoV infection (16). These neurological lesions leading to death are not

consistent with the pathology of COVID-19 in humans, in which infection of the central

nervous system can be detected post-mortem in severe COVID-19 cases, although to a

lower extent and unrelated to the severity of brain lesions (17, 18). Therefore, new mouse

models that better recapitulate the human disease, without lethal viral neuroinvasion,

could complement other existing animal models (hamster or NHP) (1).

Knock-in (KI) models are characterized by the targeted insertion of a defined gene copy

number in a specific locus, therefore allowing for a more accurate and predictable

expression of the transgene. In this work, we used a recombinase-based approach to insert

the original K18-hACE2 transgene into the collagen type I alpha chain (COL1A1) locus to

generate a Col1a1-K18-hACE2 KI mouse. To characterize this new model, mice were

challenged with SARS-CoV-2 B.1 (D614G) isolate and pathogenicity was compared with the

well-characterized K18-hACE2 transgenic mice. After viral challenge of Col1a1-K18-hACE2

mice, we confirmed viral replication and histological lesions in the lungs, but minimal viral

neuroinvasion.

Results

Assessment of hACE2 expression in the Col1a1-K18-hACE2 mice

We generated a new hACE2 KI mouse model through addition of the original K18-hACE2

transgene, kindly supplied by Paul McCray (9) at the COL1A1 locus using a recombinase

mediated system (**Figure 1A**) (19). Traditional transgenic techniques rely on the random

genomic insertion of an unpredictable number of transgene copies. This frequently results

in a highly variable level and pattern of transgene expression, due to cis acting regulatory

elements at the site of insertion. A single copy of a transgene, inserted into the well

characterised Col1A1 locus, is expected to result in a more predictable level and pattern of

expression of the transgene sequence. To confirm the correct expression of the transgene

in this KI model, we initially quantified *hACE2* transcripts relative to the house-keeping gene

GAPDH by RT-qPCR in RNA samples extracted from twelve different tissues. Expression of

the hACE2 mRNA in K18-hACE2 mice was similar among most tested tissues except in

muscle and heart, which showed lower values (**Figure 1B).** In contrast, hACE2 mRNA

expression in Col1a1-K18-hACE2 mice was more heterogenous, with the lowest expression

observed in the brain and liver. When comparing the relative expression of hACE2 mRNA

between both models, we observed higher expression in the pancreas, lymph nodes and

muscle of Col1a1-K18-hACE2 mice. Conversely, in the kidney, brain and liver, hACE2 mRNA

expression was higher in the K18-hACE2 mouse model. Minor differences were observed in

the rest of the analyzed tissues (**Figure 1C**). To complement mRNA expression data, protein

expression was assessed by Wester-blot (WB) in 6 different tissues using a monoclonal

antibody specific for hACE2 (20) and a polyclonal antibody recognizing both hACE2 and

mACE2. GAPDH was used as control, and tissues from Col1a1-K18-hACE2, K18-hACE2 and

WT mice were analyzed. Data showed undetectable protein expression in muscle, and

higher hACE2 expression in lung, pancreas and spleen of Col1a1-K18-hACE2 mice compared

to K18-hACE2 animals **(Figure 1D)**. Conversely, as observed by RT-qPCR, a higher hACE2

expression was detected in K18-hACE2 brain and liver compared to Col1a1-K18-hACE2

samples **(Figure 1D)**.

Survival of Col1a1-K18-hACE2 mice following SARS-CoV-2 B.1 infection.

Col1a1-K18-hACE2 mice (n=27) and K18-hACE2 mice (n=4) were challenged intranasally

with a SARS-CoV-2 B.1 (D614G) isolate. Uninfected KI animals, used as a control, received

an intranasal dose of PBS (n=5). Animal weight and clinical signs were monitored daily for

14 days after infection. The endpoints to collect samples were set at 3-, 7-, and 14-days

post-infection (dpi, n=8 Col1a1-K18-hACE2 mice per endpoint) or upon fulfilment of

humane endpoint criteria (**Figure 2A**).

Consistent with previous studies in transgenic animals with the B.1 SARS-CoV-2 variant,

weight-loss could be observed in both animal models starting around 3 dpi (**Figure 2B**) (13).

In K18-hACE2 transgenic animals, the decrease in weight was associated with the

progressive appearance of severe clinical signs: decrease in mobility, dyspnea, and

neurological effects **(Supplementary Table 1),** fulfilling the humane endpoint criteria and

leading to euthanasia of all K18-hACE animals by 6 or 7dpi (**Figure 2C**). In contrast, while

Col1a1-K18-hACE2 mice underwent weight loss, they did not show severe clinical signs

apart from a slight decrease in activity. Three Col1a1-K18-hACE2 animals had to be

euthanized (exitus) exclusively due to their weight loss exceeding the 20% limit (humane

endpoint) at 8, 9 and 10 dpi. Convalescent animals started to regain weight after 9 dpi,

reaching 90% of their initial weight by 14 dpi (**Figure 2B/C**). These results indicate that B.1

SARS-CoV-2 infection is significantly less pathogenic in Col1a1-K18-hACE2 mice than in K18-

hACE2 transgenic mice (**Figure 2C**).

SARS-CoV-2 B.1 replicates in the respiratory tract of Col1a1-K18-hACE2 KI mice.

To fully characterize the kinetics of viral replication in the respiratory tract, samples from

oropharyngeal swab, lung, and nasal turbinate were collected at 3, 7, and 14 dpi or at the

humane endpoint (exitus) in all Col1a1-K18-hACE2 mice (n=32). Samples from K18-hACE2

transgenic mice obtained at euthanasia (6 and 7 dpi) were used as reference (n=4). Viral

RNA analysis in oropharyngeal swab, lung, and nasal turbinate revealed widespread

infection in both K18-hACE2 and Col1a1-K18-hACE2 mice in all mentioned tissues across

two independent experiments (**Figure 3A**). VL in the nasal turbinate were similar in both

animal models, with some viral RNA was still detected at 14 dpi in Col1a1-K18-hACE2 mice.

RNA levels in oropharyngeal swabs and lung peaked at 3 dpi in Col1a1-K18-hACE2 mice and

tended to decay overtime up to 14 dpi. To further confirm viral clearance, titration of

replication-competent virus was performed in lung samples. Similar to the VL kinetics, viral

titers peaked at 3 dpi, followed by a quick and significant decrease (**Figure 3B,** 3 dpi vs 7 dpi:

p = 0.00178), concomitant with the elicitation of neutralizing antibodies **(Supplementary**

Figure 1). Col1a1-K18-hACE2 animals euthanized at the humane endpoint at 8-10 dpi still

displayed high viral RNA in lungs, which could explain their more pronounced weight-loss

and early fulfilment of endpoint criteria. However, high titers of neutralizing antibodies in

these animals probably prevented the recovery of infectious viruses **(Figure 3B and**

Supplementary Fig 1).

To characterize histopathological changes, formalin-fixed lung samples from both animal

models were analyzed by histology and immunohistochemistry (IHC) to score the presence

and extension of lesions and SARS-CoV-2 nucleoprotein antigen (NP), respectively.

Histological lesions were semi-quantified as none, mild, moderate, or severe (score 0-3),

and the detection of viral antigen as lack, low, moderate, or high (score 0-3). In Col1a1-K18-

hACE2 KI mice, histopathological analysis showed development of bronchointerstitial

pneumonia characterized by multifocal increased thickness of interalveolar wall, presence

of macrophage-like cells into alveoli surrounding bronchi and bronchiole, and hyperplasia

of type II pneumocytes. These lesions evolved from mild and multifocal by 3 dpi to moderate

in most of the animals by 14 dpi, also with increasing lymphoplasmacytic infiltration until

the end of the study. The extension of hyperplasia of type II pneumocytes was barely

detected at 3 dpi, mild by 7 dpi and moderate by 14 dpi. Therefore, an increase in lung lesion

average was observed over time **(Figure 3C and D).** Animals reaching humane endpoint

displayed areas of severe bronchointerstitial pneumonia (**Supplementary Figure 2**). IHC

detection of SARS-CoV-2 NP in the lungs of Col1a1-K18-hACE2 KI mice was consistent with

the VL analyses, showing the highest score early upon infection (3 dpi). Antigen detection

decreased with time showing apparent clearance by 14 dpi **(Figure 3C and D).** However,

animals reaching humane endpoint at 8,9 and 10 dpi showed mild to moderate SARS-CoV-

2 antigen staining mainly located in bronchiolar epithelium from different bronchiole

(**Supplementary Figure 2**).

To assess infection-driven inflammation, we analyzed the levels of IL-6 and IFNγ cytokines,

involved in inflammation and immune activation; and IP-10, MCP-1 and MIP-1β

chemokines, primarily involved in the recruitment of immune cells. Cytokine and chemokine

levels were measured by Luminex in lung samples from both animal models at the indicated

timepoints. While all markers were increased in infected animals, compared to uninfected

controls, IP-10 and IL-6 peaked at 3 dpi and significantly decreased by 14 dpi in Col1a1-K18-

hACE2 KI mice (**Figure 4**). In contrast, IFNγ, MCP-1, and MIP1β peaked at 7 dpi and decayed

afterwards, although no statistical differences were observed among the analyzed

timepoints. Overall, levels of inflammatory mediators at 7 dpi in Col1a1-K18-hACE2 KI mice

were comparable to K18-hACE2 transgenic mice at the time of euthanasia (6 and 7 dpi).

Residual levels of IFNγ and MIP1β were observed at 14dpi in some Col1a1-K18-hACE2 mice,

probably associated with the remaining tissue lesions at this time point, since no viral

antigen could be detected **(Figure 4. C-D).**

SARS-CoV-2 B.1 replication in other tissues

To explore the extent of viral infection in both animal models, viral loads (VL, cell-free viral

RNA) were quantified in a large set of samples (brain, muscle, intestine, liver, kidney,

pancreas, salivary glands, lymph nodes and spleen) collected from a subset of animals

(n=20). In both models, VL were mostly undetected in most tissues except for low levels

found in the heart and salivary glands (**Figure 5**). The analysis of brain samples showed high

VL exclusively in K18-hACE2 transgenic animals. However, the initial screening of brain

tissue from Col1a1-K18-hACE2 mice showed low but detectable VL in one animal at 3 dpi

and two animals at 7 dpi. In sharp contrast, euthanized animals, which showed the highest

weight loss and lung VL, lacked detectable VL in the brain (**Figure 5**).

Col1a1-K18-hACE2 KI mice do not develop massive brain infection.

To further investigate the level of viral infection in the brain (**Figure 5**), we performed a

longitudinal virological and histological analysis of brain samples in all animals (n=32).

Considering the full set of data, VL was detected at low levels in 3 of 8 Col1a1-K18-hACE2

mice at 3dpi and 2 out of 8 Col1a1-K18-hACE2 KI mice at 7 dpi. No VL was found in the

additional animal that reached the humane endpoint. In contrast, all K18-hACE2 brain

samples showed significantly higher VL at euthanasia (6-7dpi) (**Figure 6A**). However, no

infective virus could be found in any Col1a1-K18-hACE2 mice at any timepoint by viral

titration, while all K18-hACE2 presented high viral titers at 6-7dpi (**Figure 6B**).

For histological analysis, formalin-fixed brain samples were scored in a blinded fashion as

described above. Histopathological analysis of the brain in the Col1a1-K18-hACE2 model

showed no lesions at any of the analyzed timepoints nor those animals that required

humane endpoint (**Supplementary Figure 2**). In contrast, brain samples from K18-hACE2

infected animals showed lesion scores between 1 and 2 **(Figure 6C,Left).** Furthermore, no

detection of SARS-CoV-2 NP antigen in the brain of Col1a1-K18-hACE2 mice was observed

by IHC at any timepoint **(Figure 6C)** or at the humane endpoint **(Supplementary Figure 1);**

while K18-hACE2 mice showed a high density of antigen diffused throughout the tissue

(score 3) at euthanasia **(Figure 6C, Right).** Brain changes in the transgenic model were

consistent with the widely reported pathology induced by initial SARS-CoV-2 variants (12,

14, 21) and were observed alongside the detection of viral antigen in the brain (**Figure 6D**).

Discussion

In this study, we characterized a novel hACE2 KI murine model by evaluating the course of infection with the B.1 D614G SARS-CoV-2 variant with a direct comparison to the well-described K18-hACE2 transgenic mouse model, in which this variant is known to be highly pathogenic. The most relevant findings included the susceptibility of the KI model to SARS-CoV-2 infection and the reduced lethality imputed to the reduced viral replication and lesions in the brain.

The development of this KI model was started at the beginning of the COVID-19 pandemic as an alternative to the K18-hACE2 transgenic mouse model. Even though the transgenic model had already been established and used in previous SARS-CoV studies (9), the availability of these animals was limited at first, as colonies needed to be restarted from cryopreserved embryos. In this context, several laboratories developed new hACE2 murine models for SARS-CoV-2 infection. Expression of the hACE2 transgene was achieved through various methods, including directed insertion using CRISPR/Cas9 technology (20) using different promoters such as Hepatocyte nuclear factor-3/forkhead homologue 4 (Hfh4) promoter (5) or the mouse ACE2 promoter (6), and by infecting mice with replication-defective adenovirus encoding hACE2 (3). In this work, we followed a different strategy wherein a single copy of the transgene was inserted into the COL1A1 locus, but under the K18 promoter as the original transgenic K18-hACE2 model. The selection of this locus was based on its known ability to provide reliable and ubiquitous expression of inserted sequences (19). Additionally, RMCE allowed the insertion of a single transgene copy which should better mimic physiological expression, potentially impacting lethality through

reduction of the ectopic expression and modification of tissue distribution of the hACE2

receptor. A recent study identifying a correlation between lower hACE2 copy number and

reduced pathogenicity further supports our initial hypothesis (22).

When compared with K18-hACE2 transgenic mice, B.1 infection of the new Col1a1-K18-

hACE2 KI mouse model resulted in much less severe clinical signs with better responsiveness

to stimuli, better physical appearance and no neurological signs. These observations were

consistent with a lower viral replication in brain of Col1a1-K18-hACE2 KI mice, as

summarized in **Supplementary Figure 3**. Focusing on the respiratory tract, viral replication

was observed in both models, although VL and antigen detection were in general lower at

7 dpi in Col1a1-K18-hACE2 KI mice. In contrast, lung tissue lesions and inflammation were

fully comparable in both models at this timepoint (**Supplementary Figure 3)**.

To explain the different pathogenicity between each model, hACE2 expression was analyzed

in different tissues. At mRNA level, expression in the lungs and nasal turbinates was

comparable or higher in Col1a1-K18 hACE2 mice, while the brain and liver showed lower

hACE2 expression, as reported for other transgenic approaches (22, 23). Expression of

hACE2 protein showed additional differences; it confirmed lower expression in brain and

liver and higher expression in pancreas of Col1A1-K18-hACE2 animals, but revealed higher

expression in lung and spleen. In general, brain hACE2 expression in this new KI model

seems to be closer to the physiological situation in humans, which showed low expression

in the CNS, based on several techniques including transcriptomics and IHC (24). Among the

clinical parameters analyzed in Col1a1-K18-hACE2 KI infected mice, we found that initial

weight-loss was similar to that of the K18-hACE mice but was not accompanied by the

increased severity of clinical signs that characterizes infection in these transgenic mice

(neurological signs, lack of responsiveness, and poor general appearance) (13). Consistent

with this observation, infection of Col1a1-K18-hACE2 KI mice led to partial recovery in

weight by 14 dpi in most animals. Importantly, the survival rate of Col1a1-K18-hACE2 KI

mice was over 70%, with three euthanized animals exclusively due to a weight loss of more

than 20%, which was one of the humane endpoints in our study. These data are in line with

other hACE2 KI mouse models in which substantial viral replication within the upper and

lower respiratory tracts with limited spread to extrapulmonary organs has been described

(8, 22, 25).

Among the tissues analyzed in Col1a1-K18-hACE2 KI infected mice, we found significant

replication of SARS CoV-2 in the respiratory tract, mainly in the lungs. Despite viral clearance

in Col1a1-K18-hACE2 KI mice which was confirmed by viral titration and IHC, lung lesions

were similar between both models by 7dpi and remained after 14 dpi in KI animals

recovering from the infection. The impact of viral replication on lung inflammation in

Col1a1-K18-hACE2 was confirmed and found to be comparable to the inflammation profile

reported in K18-hACE2 transgenic mice (21, 26). Infection of Col1a1-K18-hACE2 KI mice

induced high levels of IFNγ, IL-6, IP-10, MCP-1 and MIP-1β cytokines. The longitudinal

analysis showed early impact on IL-6 and IP-10 levels (3 dpi) and a clear trend to

normalization by 14 dpi, with some residual inflammation remaining. This is consistent with

the presence of lesions at this timepoint, and the immune cell infiltration described by 21

dpi in K18-hACE2 transgenic mice surviving infection (21).

The main difference between the Col1a1-K18-hACE2 KI and the K18-hACE2 transgenic

models was found in neuroinvasion. Residual levels of viral RNA but no replication

competent virus or viral proteins were detected in the brains of Col1a1-K18-hACE2 KI mice.

Furthermore, no detectable brain lesions, such as the multifocal lymphoplasmacytic

meningoencephalitis reported in K18-hACE2 transgenic mice (14) were detected by

histology, and no SARS-CoV-2 nucleoprotein could be evidenced by IHC. Importantly, the

three animals that reached humane endpoints did not present any evidence of brain viral

replication but showed enhanced lung infection. Considering that the expression of hACE2

is lower in brain and higher in lung in Col1a1-K18-hACE2 compared to K18-hACE2, it seems

that the targeted insertion of the hACE2 transgene can be a crucial factor leading to lower

levels of brain infection and neurological clinical signs, as well as enhanced respiratory

affection in this new model. However, different studies have shown that neuroinvasion is

only partially dependent on the expression of hACE2 receptor (12), and that both delivery

route and viral dose can play a role in the magnitude of the neuroinvasion (23, 27). In our

case, to exclusively analyze the effect of receptor expression, we kept the intranasal

administration route and TCID$_{50}$ dose established and characterized in our own and others'

studies to ensure comparability of the results (13, 28).

Our study is a preliminary characterization of a new animal model and has consequently

several limitations. The first limitation was the age of the animals used. The Col1a1-K18-

hACE2 mice ranged in age from 5 to 11 months, depending on availability at the time of the

study. However, although pathogenicity of SARS-CoV-2 infection has the potential to be

higher in older animals, we could not detect significant age differences between euthanized

and convalescent animals (mean+/-SD age of 9.1+/-2.0 and 8.0+/-2.1 months, respectively).

Second, the expression of hACE2 has been analyzed at both the mRNA and protein levels in

bulk tissues, but a further characterization in specific cell types will be necessary to link

expression levels with pathology. Moreover, we have focused on the characterization of

one viral variant and a single dose. The B.1 variant was selected for its demonstrated

severity in K18-hACE2 mice and its inability to use mACE2 as an entry receptor, unlike

B.1.351/Beta and Omicron BA1.1 variants (13, 15). While ancestral and earlier variants

(B.1.1.7/Alpha, B.1.351/Beta B.1.617.2/Delta) were highly pathogenic in K18-hACE2 mice,

Omicron BA.1.1 caused a milder infection with no weight loss nor neuroinvasion, therefore

reducing lethality in this model (13). However, more recent Omicron subvariants such as

BQ.1.1, BA.5 and XBB.1.5 appear to be regaining pathogenicity in K18-hACE2 mice (29–31).

Increased lung infection, pro-inflammatory cytokines, and lung pathology are observed with

these variants, with data on neuroinvasion and neurovirulence on the latter (31). Similarly,

lower omicron variants has been described in KI mice (25, 32). Nevertheless, for the B.1

SARS-CoV-2 variant the Col1a1-K18-hACE2 model shows similarity with GSH in viral

dynamics and pathology, with minimal neuroinvasion. This new model, however, has the

upside of a wider reagent availability, scarce in GSH (1), and easier animal facility allowance.

In summary, several mouse models for the evaluation of antivirals and vaccines have been

developed to date, with disease phenotypes ranging from mild to severe COVID-19-like

condition. None of these models, however, can fully recapitulate all aspects of the disease

as it occurs in humans. The development and further characterization of new animal

models, like the Col1a1-K18-hACE2 model, may overcome some of these limitations and

provide valuable tools to study certain aspects of COVID-19. Survival of Col1a1-K18-hACE2

KI mice after infection with a highly pathogenic variant could provide a model which better

mimics human disease progression and thus could be instrumental for the study of specific

disease aspects such as the consequences of inflammation (including loss of taste or smell)

post-COVID conditions (PCC) or sequelae, long-term effects of drug therapies, and

susceptibility to reinfection.

Materials and methods

Creation of the Knock-in Model

A new KI mouse model, **B6;129S-Col1a1^{tm1(K18-hACE2)Irb}/Irsi** (Col1a1-K18-hACE2), was created

by inserting the original K18-hACE2 transgene into the collagen COL1A1 locus using a

recombinase-mediated cassette exchange (RMCE) FLP-FRT system in KH2 cells (19). The

actual insertion site lies approximately 0.3 kb downstream of the 3'UTR end of COL1A1. As

such, the inserted transgene remains identical to the original designed one (9) and the

hACE2 cDNA is under the control of the K18 promoter rather than COL1A1. Specifically, the

plasmid   PGK-ATG-Frt   was   digested   with   EcoRV,   and   the   sequence

ATCAGACGTCGCTAGCGGCGCGCCGGTACTAGT was inserted to create a multi-cloning site.

The plasmid containing hACE2 under the control of the K18 promoter (K18-hACE2), supplied

by Paul McCray, was digested with HpaI and XbaI enzymes. The resulting transgene

fragment was isolated and purified, and then cloned into the EcoRV-NheI sites of the

modified PGK-ATG-Frt plasmid to generate the targeting vector. The targeting vector was

then co-transfected with an Flp expression construct into KH2 cells by electroporation. The

cells were then placed under hygromycin selection for 9 days, after which drug resistant

colonies were picked, expanded, and screened for the presence of the targeted transgene.

Once confirmed, the modified embryonic stem cells were injected into mouse blastocysts.

These injected mouse blastocysts were then placed into recipient foster females via embryo

transfer techniques.

Biosafety Approval and Virus Isolation

The execution of SARS-CoV-2 experiments was approved by the biologic biosafety

committee of Germans Trias i Pujol Research Institute (IGTP) and performed at the Biosafety

Level 3 laboratory (BSL-3) of the Comparative Medicine and Bioimage Centre (CSB-20-015-

M8; CMCiB, Badalona, Spain).

The B.1 SARS-CoV-2 isolate used in this study was isolated from nasopharyngeal swabs of

hospitalized patients in Spain as described elsewhere (33, 34). Briefly, viruses were

propagated in Vero E6 cells (CRL-1586; ATCC, Virginia, VA, United States) for two passages

and recovered by supernatant collection. The sequence of the SARS-CoV-2 variant tested is

deposited at the GISAID Repository with accession IDs EPI_ISL_510689. EPI_ISL_510689 was

the first SARS-CoV-2 virus isolated in Catalonia in March 2020 and, compared to the Wuhan/

Hu-1/2019 (WH1) strain, this isolate had the S protein mutations D614G, which is associated

with the B.1 lineage, and R682L. Viral stocks were titrated on Vero E6 cells to use equivalent

TCID50/mL using the Reed-Muench method and sequential 1/10 dilutions of the viral stocks

as described previously (34).

Animal Procedures and Study Design

All animal procedures were approved by the Committee on the Ethics of Animal

Experimentation of the IGTP and were authorized by the *Generalitat de Catalunya* (code:

11222). All animal experiments followed the principles of animal welfare and the 3Rs. All

experiments and sample processing were performed inside the BSL-3 facility. Col1a1-K18-

hACE2 hemizygous KI mice were produced and bred at Parc Científic de Barcelona (PCB) by

pairing hemizygous males for Tg(K18-ACE2)2Prlmn (or K18-hACE2) with non-carrier

C57Bl6/J females. Animals used as control, B6.Cg-Tg(K18-ACE2)2Prlmn/J (or K18-hACE2)

hemizygous transgenic mice (034860, Jackson Immunoresearch, West Grove, PA, United

States) were bred at CMCiB in the Specific Pathogen Free Area (SPF) by pairing hemizygous

K18-hACE2 males with non-carrier C57Bl6/J females. The genotype of the offspring

regarding both K18-hACE2 and Col1a1-K18-hACE2 mice was determined by qPCR at the

IGTP's Genomics Platform from tail samples. Both animal models were kept in the BSL-3

facility during the whole experiment including acclimatization period. The housing

conditions in the BSL-3 room were maintained as follows: a temperature of 22±2ºC,

humidity levels between 30-70%, 20 ACH, a 12h dark/light cycle, and access to food and

water *ad libitum.*

A total of 36 adult mice aged 5-11 months were used in this experiment, consisting of 32 KI

Col1a1-K18-hACE2 mice and 4 transgenic K18-hACE2 mice. All groups were sex balanced.

The infections were performed in two separate experiments between January and June

2022. Mice were anaesthetized with isoflurane (FDG9623; Baxter, Deerfield, IL, USA) and

infected with a B.1 isolate (27 Col1a1-K18-hACE2 and 4 K18-hACE2 mice). The uninfected

control group received PBS (5 Col1a1-K18-hACE2).

Infection was performed using 1,000 $TCID_{50}$ of B.1 SARS-CoV-2 isolate in 50 µl of PBS (25

µL/nostril), or PBS only (25 µL/nostril) for the control group. All mice fully recovered from

the challenge and anaesthesia procedures. Following the challenge, body weight and clinical

signs were monitored daily. Eight animals per group were euthanized at days 3, 7, 14 dpi or

upon fulfilment of human endpoint, for viral RNA quantification and histological analyses

(**Supplementary Table 2**). The endpoint criteria for animal welfare were established based

on a body weight loss of more than to 20% of the initial body weight and/or the display of

moderate to severe clinical signs (including neurological signs), in accordance with previous

studies (12–14). The evaluated clinical signs included respiration, physical appearance, lack

of responsiveness and neurological signs, that were scored from 0-2 depending on severity

(**Supplementary Table 1**). Euthanasia was performed under deep isoflurane anaesthesia by

whole blood extraction via cardiac puncture and cervical dislocation. Oropharyngeal swab,

lung, brain, and nasal turbinate were collected for viral and hACE2 RNA quantification,

histological and IHC analyses. For the latter techniques, tissues were fixed by immersion in

10% buffered formalin. An additional set of 9 tissues was collected to further characterize

the KI model both for viral RNA quantification and hACE2 receptor expression. These tissues

included muscle, intestine, liver, kidney, lymph nodes, spleen, heart, pancreas, and salivary

glands.

Tissue sampling and processing

The collected tissues were processed as described by Tarrés-Freixas et al. 2022 (13). Briefly,

approximately 100 mg of each tissue was collected in 1.5 mL Sarstedt tubes (72607;

Sarstedt, Nümbrecht, Germany) containing 500 µL of DMEM medium (11995065;

ThermoFisher Scientific) supplemented with 1% penicillin–streptomycin (10378016;

ThermoFisher Scientific). A 1.5 mm Tungsten bead (69997; QIAGEN, Hilden, Germany) was

added to each tube, and samples were homogenized twice at 25 Hz for 30 s using a

TissueLyser II (85300; QIAGEN) before being centrifuged for 2 min at 2,000 × g. Supernatants

were then stored at −80°C until the analysis of hACE2 expression and VL by RT-qPCR.

hACE2 mRNA Expression

RNA tissue extraction was performed by using the Viral RNA/ Pathogen Nucleic Acid Isolation kit (A42352; ThermoFisher Scientific), optimized for use with a KingFisher instrument (5400610; ThermoFisher Scientific), following the manufacturer's instructions. The expression of hACE2 receptor in tissue was detected by RT-qPCR with a predefined TaqMan assay with a ACE2 target (Hs01085333_m1, ThermoFisher Scientific). Mouse *gapdh* gene expression was measured in duplicate for each sample using TaqMan®gene expression assay (Mm99999915_g1; ThermoFisher Scientific) as amplification control. Data was graphed both as the delta-Ct values of hACE2 minus GAPDH for each tissue, and the relative expression of the receptor (delta-delta Ct value) of Col1a1-K18-hACE2 minus K18-hACE2 tissues.

ACE2 Protein Expression

ACE2 protein expression was assessed by WB. Tissue samples were obtained from uninfected Col1a1-K18-hACE2, K18-hACE2 and WT mice and flash frozen. Frozen sections of each tissue were sliced by cryostat processing (Leica CM1950). and homogenized in 100uL of RIPA buffer for 15 min on ice, followed by centrifugation at 1000g for 10 min. Supernatant was then isolated, and total protein concentration was quantified by Pierce™ BCA Protein Assay (23225; ThermoFisher Scientific) following manufacturer's instructions.

For WB, 3 µg of lysate total protein (as quantified Pierce™ BCA Protein Assay (23225; ThermoFisher Scientific) were run in NuPAGE Bis-Tris 4-12% acrylamide gels (NP0321; ThermoFisher Scientific). BlueStar pre-stained protein ladder was used as a molecular weight marker (523005; NIPPON Genetics, Tokyo, Japan). Proteins were blotted onto PVDF

membrane (Ref. 1704156; Bio-Rad) using the Trans-Blot Turbo Transfer System (Ref.

1704150 Bio-Rad) and membranes were blocked with EveryBlot Blocking Buffer (12010020;

Bio-Rad). Primary antibodies used were: Anti-hACE2 specific monoclonal (1:1000;

AB108209, Abcam) (20), Anti-ACE2 polyclonal (1:1000 AF933; R&D Systems) and Anti-

GAPDH (1:5000; Ab9485; Abcam). Secondary antibodies were: donkey anti-Rabbit (1:10000;

Ref. 711-036-152; Jackson Immunoresearch) or a donkey anti-Goat (1:10000; Ref. 705-035-

147; Jackson Immunoresearch). Membranes were developed with SuperSignal™ West

Femto Maximum Sensitivity Substrate (Ref. 34094; ThermoFisher Scientific) according to

the manufacturer's instructions and acquired with ChemiDOC^{TM} MP Imaging System

(12003154; Bio-Rad). Images were processed and merged with ImageLab 6.0.1 software

(Bio-Rad).

SARS-CoV-2 PCR Detection and Viral Load Quantification

Viral RNA was quantified by RT-qPCR in both the standard set of samples (oropharyngeal

swab, lung, brain, and nasal turbinate), and the extended set (muscle, intestine, liver,

kidney, pancreas, salivary glands, lymph nodes and spleen).

RNA extraction was performed by using the Viral RNA/ Pathogen Nucleic Acid Isolation kit

(A42352; ThermoFisher Scientific), optimized for use with a KingFisher instrument

(5400610; ThermoFisher Scientific), following the manufacturer's instructions. PCR

amplification was based on the 2019-Novel Coronavirus Real-Time RT-PCR Diagnostic Panel

guidelines and protocol developed by the American Center for Disease Control and

Prevention (CDC-006-00019, v.07). Briefly, a 20 μL PCR reaction was set up containing 5 μL

of RNA, 1.5 μL of N2 primers and probe (2019-nCov CDC EUA Kit; catalog number 10006770;

Integrated DNA Technologies, Coralville, IA, USA) and 10 µl of GoTaq 1-Step RT-qPCR

(Promega, Madison, WI, USA). Thermal cycling was performed at 50°C for 15 min for reverse

transcription, followed by 95°C for 2 min and then 45 cycles of 95°C for 10 s, 56°C for 15 s

and 72°C for 30 s in the Applied Biosystems 7,500 or QuantStudio5 Real-Time PCR

instruments (ThermoFisher Scientific). For absolute quantification, a standard curve was

built using 1/5 serial dilutions of a SARS-CoV-2 plasmid (2019-nCoV_N_Positive Control;

catalog number 10006625, 200 copies/µL, Integrated DNA Technologies), which was run in

parallel with all PCR determinations. Viral RNA from each sample was quantified in

triplicate, and the mean viral RNA concentration (in copies/mL) was extrapolated from the

standard curve and corrected by the corresponding dilution factor. Mouse *gapdh* gene

expression was measured in duplicate for each sample using TaqMan®gene expression

assay (Mm99999915_g1; ThermoFisher Scientific) as amplification control.

Viral Titration

Lung tissues were evaluated for the presence of replicative virus by titration in Vero E6 cells

as previously described (14, 34, 35). Briefly, after tissue homogenization, each sample

underwent sequential 10-fold dilutions in duplicate, transferred onto a monolayer of Vero

E6 cell in a 96-well plate, and incubated at 37°C and 5% CO_2. Plates were monitored daily

under a microscope, and at 5 dpi, wells were evaluated for the presence of cytopathic

effects. The amount of infectious virus was calculated by determining the TCID50 using the

Reed–Muench method.

Histological and Immunohistochemical Analyses

Tissue samples were recovered at the designated endpoint (3,7,14 dpi or humane endpoint) and fixed by immersion in 10% buffered formalin. Lung, nasal turbinate, and brain samples were routinely processed for histological examination, with haematoxylin & eosin-stained slides examined under an optical microscope in a blinded fashion. Brain was cut into two coronal sections, including cerebellum/pons and hemispheres at thalamus level. A semi-quantitative approach based on the amount of inflammation (none, mild, moderate, or severe) was used to assess the damage caused by SARS-CoV-2 infection in mice, following a previously published scoring system (14, 36). Additionally, an IHC technique was employed to detect SARS-CoV-2 NP antigen in nasal turbinate, lung, and brain sections from all animals, using a rabbit monoclonal antibody (40143-R019, Sino Biological, Beijing, China) at a 1:15,000 dilution. The amount of viral antigen in tissues was semi-quantitatively scored in a blinded fashion (low, moderate, and high amount, or lack of antigen detection) (14, 36).

Cytokine Quantification

To assess the viral-driven inflammation in lung in both animal models, the levels of IP-10, IL-6, IFNγ, MCP-1 and MIP-1β cytokines were analyzed by Luminex in tissue extracts. Lung samples were processed as described above and stored at -80ºC until analysis. In Col1a1-K18-hACE2 mice, cytokines were analyzed at 3,7,14 dpi and those euthanized by humane endpoint criteria (exitus). Uninfected Col1a1-K18-hACE2 and K18-hACE2 infected animals were used as reference groups.

Cytokines were measured by Luminex xMAP technology and analyzed with xPONENT 3.1 software (Luminex Corporation) using the MCYTOMAG-70 kit, according to the

manufacturers' protocol with minor modifications. Briefly, after cytokine staining, samples

underwent an overnight incubation on a rocking shaker at 4°C, using 2% PFA to ensure

complete inactivation of any remaining SARS-CoV-2 particles; a fixation that does not alter

cytokine quantification (37). Before plate acquisition, the PFA was washed away and

replaced with sheath fluid.

Pseudovirus generation and neutralization assay

To assess the generation of neutralizing antibodies upon SARS-CoV-2 B.1 infection, we used

a previously described pseudovirus-based neutralization assay (38). Briefly, in Nunc 96-well

cell culture plates (Thermo Fisher Scientific), 200 TCID50 of a luciferase-reporter HIV-based

pseudovirus bearing the BA.1 SARS-CoV-2 spike were preincubated with four-fold serial

dilutions (1/60 - 1/61,440) of the heat-inactivated (56ºC for 30 minutes) serum samples, for

1 hour at 37ºC. Then 1×10^4 HEK293T/hACE2 cells treated with DEAE-Dextran (Sigma-

Aldrich) were added. Results were read after 48 hours using the EnSight Multimode Plate

Reader and BriteLite Plus Luciferase reagent (PerkinElmer, USA). The values were

normalized, and the ID50 (reciprocal dilution inhibiting 50% of the infection)

were calculated as described (39).

Statistical Analyses

All figures were generated using GraphPad Prism 9.0.0. Statistical analyses were performed

using R v4.3. Survival Rates were estimated with Kaplan-Meier curves and compared with

the Log-rank test. Datasets with an abundance of data below the limit of detection, like VL,

were analyzed using the Peto-Peto Left-censored samples test with correction for multiple

comparisons. Histopathological, IHC scores and viral titrations were compared using an

Independence Asymptotic Generalized Pearson Chi-Squared Test for ordinal data. Cytokine

titers were compared by a Kruskal Wallis, with pairwise comparisons conducted using

Conover's non-parametric test.

Acknowledgements

The authors would like to acknowledge Jorge Díaz, Yaiza Rosales-Salgado, Rosa María

Ampudia-Carrasco, Sergi Sunyé-Casas and Mireia Martínez from the CMCiB for their

essential help in the BSL3 facility and the K18-hACE2 mouse colony. We also thank Marisa

Larramona from Parc Científic de Barcelona for her invaluable help with the knock-in mouse

colony. We thank the Dormeur Fondation for their financial support for the acquisition of

the QuantStudio-5 real-time PCR system.

Author Contributions

AP-G, FT-F, BT, and JB conceived and designed the experiments. AP-G, FT-F, and BT

performed the animal procedures. AP-G, FT-F, MP, ER-M, DR-R, DP-Z, JM-B, EP, JS, BT

performed the analytical experiments. AP-G, B, JS, JV-A, VU and JB analyzed and interpreted

the data. SF, BTondelli established and provided the Knock-in mouse colony. SC established

communication between animal facilities and contributed to the veterinary report

verification to allow the knock-in animal shipment to the CMCiB facility. DP-Z, JM-B, DR-R,

NI-U provided key reagents. AP-G, BT, and JB wrote the paper. All authors contributed to

the article and approved the submitted version.

Competing Interests /Funding

A.P-G was supported by a predoctoral grant from Generalitat de Catalunya and Fons Social

Europeu (2022 FI_B 00698). This study was funded by Grifols, the Departament de Salut of

the Generalitat de Catalunya (grant nos. SLD016 to J.B. and SLD015 to J.C.), the Spanish

Health Institute Carlos III, CERCA Programme/Generalitat de Catalunya 2021 SGR 00452,

and the crowdfunding initiatives #joemcorono, BonPreu/Esclat, and Correos. NI-U is

supported by the Spanish Ministry of Science and Innovation (grants PID2023-147498OB-

I00, PID2020-117145RB-I00 and 10.13039/501100011033, Spain), EU HORIZON-HLTH-

2021CORONA-01 (grant 101046118, European Union) and by institutional funding of

Pharma Mar, Grifols, HIPRA, and Amassence. J. Blanco has received institutional funding

from Grífols, Nesapor Europe, HIPRA and MSD. Unrelated to the submitted work, J.B. and

J.C. are founders and shareholders of AlbaJuna Therapeutics, SL. B.C. is founder and

shareholder of AlbaJuna Therapeutics, SL, and AELIX Therapeutics, SL.

The funders had no role in study design, data collection and interpretation, or the decision

to submit the work for publication

Data availability

The data supporting the findings of this study are documented within the paper and are

available from the corresponding authors upon request.

References

1.   Muñoz-Fontela C, Dowling WE, Funnell SGP, Gsell PS, Riveros-Balta AX, Albrecht RA,
Andersen H, Baric RS, Carroll MW, Cavaleri M, Qin C, Crozier I, Dallmeier K, de Waal
L, de Wit E, Delang L, Dohm E, Duprex WP, Falzarano D, Finch CL, Frieman MB,
Graham BS, Gralinski LE, Guilfoyle K, Haagmans BL, Hamilton GA, Hartman AL,
Herfst S, Kaptein SJF, Klimstra WB, Knezevic I, Krause PR, Kuhn JH, Le Grand R, Lewis
MG, Liu WC, Maisonnasse P, McElroy AK, Munster V, Oreshkova N, Rasmussen AL,
Rocha-Pereira J, Rockx B, Rodríguez E, Rogers TF, Salguero FJ, Schotsaert M,
Stittelaar KJ, Thibaut HJ, Tseng C Te, Vergara-Alert J, Beer M, Brasel T, Chan JFW,
García-Sastre A, Neyts J, Perlman S, Reed DS, Richt JA, Roy CJ, Segalés J, Vasan SS,
Henao-Restrepo AM, Barouch DH. 2020. Animal models for COVID-19. Nature
586:509–515.

2.   Jackson CB, Farzan M, Chen B, Choe H. 2021. Mechanisms of SARS-CoV-2 entry into
cells. Nat Rev Mol Cell Biol 2021 231 23:3–20.

3.   Hassan AO, Case JB, Winkler ES, Thackray LB, Kafai NM, Bailey AL, McCune BT, Fox
JM, Chen RE, Alsoussi WB, Turner JS, Schmitz AJ, Lei T, Shrihari S, Keeler SP,
Fremont DH, Greco S, McCray PB, Perlman S, Holtzman MJ, Ellebedy AH, Diamond
MS. 2020. A SARS-CoV-2 Infection Model in Mice Demonstrates Protection by
Neutralizing Antibodies. Cell 182:744-753.e4.

4.   Tseng C-TK, Huang C, Newman P, Wang N, Narayanan K, Watts DM, Makino S,
Packard MM, Zaki SR, Chan T, Peters CJ. 2007. Severe Acute Respiratory Syndrome
Coronavirus Infection of Mice Transgenic for the Human Angiotensin-Converting
Enzyme 2 Virus Receptor. J Virol 81:1162–1173.

5.   Jiang R Di, Liu MQ, Chen Y, Shan C, Zhou YW, Shen XR, Li Q, Zhang L, Zhu Y, Si HR,
Wang Q, Min J, Wang X, Zhang W, Li B, Zhang HJ, Baric RS, Zhou P, Yang X Lou, Shi
ZL. 2020. Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing Human
Angiotensin-Converting Enzyme 2. Cell 182:50-58.e8.

6.   Bao L, Deng W, Huang B, Gao H, Liu J, Ren L, Wei Q, Yu P, Xu Y, Qi F, Qu Y, Li F, Lv Q,
Wang W, Xue J, Gong S, Liu M, Wang G, Wang S, Song Z, Zhao L, Liu P, Zhao L, Ye F,
Wang H, Zhou W, Zhu N, Zhen W, Yu H, Zhang X, Guo L, Chen L, Wang C, Wang Y,
Wang X, Xiao Y, Sun Q, Liu H, Zhu F, Ma C, Yan L, Yang M, Han J, Xu W, Tan W, Peng
X, Jin Q, Wu G, Qin C. 2020. The pathogenicity of SARS-CoV-2 in hACE2 transgenic
mice. Nat 2020 5837818 583:830–833.

7.   Zhou B, Thao TTN, Hoffmann D, Taddeo A, Ebert N, Labroussaa F, Pohlmann A, King
J, Steiner S, Kelly JN, Portmann J, Halwe NJ, Ulrich L, Trüeb BS, Fan X, Hoffmann B,
Wang L, Thomann L, Lin X, Stalder H, Pozzi B, de Brot S, Jiang N, Cui D, Hossain J,
Wilson MM, Keller MW, Stark TJ, Barnes JR, Dijkman R, Jores J, Benarafa C,
Wentworth DE, Thiel V, Beer M. 2021. SARS-CoV-2 spike D614G change enhances
replication and transmission. Nature 592:122–127.

8.   Winkler ES, Chen RE, Alam F, Yildiz S, Case JB, Uccellini MB, Holtzman MJ, Garcia-Sastre A, Schotsaert M, Diamond MS. 2022. SARS-CoV-2 Causes Lung Infection without Severe Disease in Human ACE2 Knock-In Mice. J Virol 96:e0151121.

9.   McCray PB, Pewe L, Wohlford-Lenane C, Hickey M, Manzel L, Shi L, Netland J, Jia HP, Halabi C, Sigmund CD, Meyerholz DK, Kirby P, Look DC, Perlman S. 2007. Lethal Infection of K18- hACE2 Mice Infected with Severe Acute Respiratory Syndrome Coronavirus . J Virol 81:813–821.

10. Zheng J, Wong LYR, Li K, Verma AK, Ortiz ME, Wohlford-Lenane C, Leidinger MR, Knudson CM, Meyerholz DK, McCray PB, Perlman S. 2021. COVID-19 treatments and pathogenesis including anosmia in K18-hACE2 mice. Nature 589:603–607.

11. Winkler ES, Bailey AL, Kafai NM, Nair S, McCune BT, Yu J, Fox JM, Chen RE, Earnest JT, Keeler SP, Ritter JH, Kang LI, Dort S, Robichaud A, Head R, Holtzman MJ, Diamond MS. 2020. SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung inflammation and impaired function. Nat Immunol 2020 2111 21:1327–1335.

12. Carossino M, Kenney D, O'Connell AK, Montanaro P, Tseng AE, Gertje HP, Grosz KA, Ericsson M, Huber BR, Kurnick SA, Subramaniam S, Kirkland TA, Walker JR, Francis KP, Klose AD, Paragas N, Bosmann M, Saeed M, Balasuriya UBR, Douam F, Crossland NA. 2022. Fatal Neurodissemination and SARS-CoV-2 Tropism in K18-hACE2 Mice Is Only Partially Dependent on hACE2 Expression. Viruses 14:535.

13. Tarrés-Freixas F, Trinité B, Pons-Grífols A, Romero-Durana M, Riveira-Muñoz E, Ávila-Nieto C, Pérez M, Garcia-Vidal E, Perez-Zsolt D, Muñoz-Basagoiti J, Raïch-Regué D, Izquierdo-Useros N, Andrés C, Antón A, Pumarola T, Blanco I, Noguera-Julián M, Guallar V, Lepore R, Valencia A, Urrea V, Vergara-Alert J, Clotet B, Ballana E, Carrillo J, Segalés J, Blanco J. 2022. Heterogeneous Infectivity and Pathogenesis of SARS-CoV-2 Variants Beta, Delta and Omicron in Transgenic K18-hACE2 and Wildtype Mice. Front Microbiol 13:840757.

14. Vidal E, López-Figueroa C, Rodon J, Pérez M, Brustolin M, Cantero G, Guallar V, Izquierdo-Useros N, Carrillo J, Blanco J, Clotet B, Vergara-Alert J, Segalés J, E V, C L-F, J R, M P, M B, G C, V G, N I-U, J C, J B, B C, J V-A, J S. 2021. Chronological brain lesions after SARS-CoV-2 infection in hACE2-transgenic mice. Vet Pathol 59:613–626.

15. Shuai H, Chan JF-W, Yuen TT-T, Yoon C, Hu J-C, Wen L, Hu B, Yang D, Wang Y, Hou Y, Huang X, Chai Y, Chan CC-S, Poon VK-M, Lu L, Zhang R-Q, Chan W-M, Ip JD, Chu AW-H, Hu Y-F, Cai J-P, Chan K-H, Zhou J, Sridhar S, Zhang B-Z, Yuan S, Zhang AJ, Huang J-D, To KK-W, Yuen K-Y, Chu H. 2021. Emerging SARS-CoV-2 variants expand species tropism to murines. EBioMedicine 73:103643.

16. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. 2008. Severe Acute Respiratory Syndrome Coronavirus Infection Causes Neuronal Death in the Absence of Encephalitis in Mice Transgenic for Human ACE2. J Virol 82:7264–7275.

17. Meinhardt J, Streit S, Dittmayer C, Manitius R v., Radbruch H, Heppner FL. 2023. The
neurobiology of SARS-CoV-2 infection. Nat Rev Neurosci 2023 251 25:30–42.

18. Matschke J, Lütgehetmann M, Hagel C, Sperhake JP, Schröder AS, Edler C,
Mushumba H, Fitzek A, Allweiss L, Dandri M, Dottermusch M, Heinemann A,
Pfefferle S, Schwabenland M, Sumner Magruder D, Bonn S, Prinz M, Gerloff C,
Püschel K, Krasemann S, Aepfelbacher M, Glatzel M. 2020. Neuropathology of
patients with COVID-19 in Germany: a post-mortem case series. Lancet Neurol
19:919–929.

19. Beard C, Hochedlinger K, Plath K, Wutz A, Jaenisch R. 2006. Efficient method to
generate single-copy transgenic mice by site-specific integration in embryonic stem
cells. genesis 44:23–28.

20. Sun SH, Chen Q, Gu HJ, Yang G, Wang YX, Huang XY, Liu SS, Zhang NN, Li XF, Xiong R,
Guo Y, Deng YQ, Huang WJ, Liu Q, Liu QM, Shen YL, Zhou Y, Yang X, Zhao TY, Fan CF,
Zhou Y Sen, Qin CF, Wang YC. 2020. A Mouse Model of SARS-CoV-2 Infection and
Pathogenesis. Cell Host Microbe 28:124-133.e4.

21. Yinda CK, Port JR, Bushmaker T, Owusu IO, Purushotham JN, Avanzato VA, Fischer
RJ, Schulz JE, Holbrook MG, Hebner MJ, Rosenke R, Thomas T, Marzi A, Best SM, de
Wit E, Shaia C, van Doremalen N, Munster VJ. 2021. K18-hACE2 mice develop
respiratory disease resembling severe COVID-19. PLOS Pathog 17:e1009195.

22. Bradshaw CM, Georgieva T, Tankersley TN, Taylor-Doyle T, Johnson L, Uhrlaub JL,
Besselsen D, Nikolich JŽ. 2024. Cutting Edge: Characterization of Low Copy Number
Human Angiotensin-Converting Enzyme 2–Transgenic Mice as an Improved Model
of SARS-CoV-2 Infection. J Immunol 212:523–528.

23. Dong W, Mead H, Tian L, Park J-G, Garcia JI, Jaramillo S, Barr T, Kollath DS, Coyne
VK, Stone NE, Jones A, Zhang J, Li A, Wang L-S, Milanes-Yearsley M, Torrelles JB,
Martinez-Sobrido L, Keim PS, Barker BM, Caligiuri MA, Yu J. 2022. The K18-Human
ACE2 Transgenic Mouse Model Recapitulates Non-severe and Severe COVID-19 in
Response to an Infectious Dose of the SARS-CoV-2 Virus. J Virol 96:e0096421.

24. Hikmet F, Méar L, Edvinsson Å, Micke P, Uhlén M, Lindskog C. 2020. The protein
expression profile of ACE2 in human tissues. Mol Syst Biol 16:e9610.

25. Barut GT, Halwe NJ, Taddeo A, Kelly JN, Schön J, Ebert N, Ulrich L, Devisme C,
Steiner S, Trüeb BS, Hoffmann B, Veiga IB, Leborgne NGF, Moreira EA, Breithaupt A,
Wylezich C, Höper D, Wernike K, Godel A, Thomann L, Flück V, Stalder H, Brügger M,
Esteves BIO, Zumkehr B, Beilleau G, Kratzel A, Schmied K, Ochsenbein S, Lang RM,
Wider M, Machahua C, Dorn P, Marti TM, Funke-Chambour M, Rauch A, Widera M,
Ciesek S, Dijkman R, Hoffmann D, Alves MP, Benarafa C, Beer M, Thiel V. 2022. The
spike gene is a major determinant for the SARS-CoV-2 Omicron-BA.1 phenotype.
Nat Commun 13:5929.

26. Oladunni FS, Park J-G, Pino PA, Gonzalez O, Akhter A, Allué-Guardia A, Olmo-

Fontánez A, Gautam S, Garcia-Vilanova A, Ye C, Chiem K, Headley C, Dwivedi V, Parodi LM, Alfson KJ, Staples HM, Schami A, Garcia JI, Whigham A, Platt RN, Gazi M, Martinez J, Chuba C, Earley S, Rodriguez OH, Mdaki SD, Kavelish KN, Escalona R, Hallam CRA, Christie C, Patterson JL, Anderson TJC, Carrion R, Dick EJ, Hall-Ursone S, Schlesinger LS, Alvarez X, Kaushal D, Giavedoni LD, Turner J, Martinez-Sobrido L, Torrelles JB. 2020. Lethality of SARS-CoV-2 infection in K18 human angiotensin-converting enzyme 2 transgenic mice. Nat Commun 11:6122.

27. Fumagalli V, Ravà M, Marotta D, Di Lucia P, Laura C, Sala E, Grillo M, Bono E, Giustini L, Perucchini C, Mainetti M, Sessa A, Garcia-Manteiga JM, Donnici L, Manganaro L, Delbue S, Broccoli V, De Francesco R, D'Adamo P, Kuka M, Guidotti LG, Iannacone M. 2022. Administration of aerosolized SARS-CoV-2 to K18-hACE2 mice uncouples respiratory infection from fatal neuroinvasion. Sci Immunol 7:eabl9929.

28. Radvak P, Kwon HJ, Kosikova M, Ortega-Rodriguez U, Xiang R, Phue JN, Shen RF, Rozzelle J, Kapoor N, Rabara T, Fairman J, Xie H. 2021. SARS-CoV-2 B.1.1.7 (alpha) and B.1.351 (beta) variants induce pathogenic patterns in K18-hACE2 transgenic mice distinct from early strains. Nat Commun 12:1–15.

29. Case JB, Scheaffer SM, Darling TL, Bricker TL, Adams LJ, Harastani HH, Trende R, Sanapala S, Fremont DH, Boon ACM, Diamond MS. 2023. Characterization of the SARS-CoV-2 BA.5.5 and BQ.1.1 Omicron variants in mice and hamsters. J Virol 97:e0062823.

30. Halfmann PJ, Iida S, Iwatsuki-Horimoto K, Maemura T, Kiso M, Scheaffer SM, Darling TL, Joshi A, Loeber S, Singh G, Foster SL, Ying B, Case JB, Chong Z, Whitener B, Moliva J, Floyd K, Ujie M, Nakajima N, Ito M, Wright R, Uraki R, Warang P, Gagne M, Li R, Sakai-Tagawa Y, Liu Y, Larson D, Osorio JE, Hernandez-Ortiz JP, Henry AR, Ciuoderis K, Florek KR, Patel M, Odle A, Wong L-YR, Bateman AC, Wang Z, Edara V-V, Chong Z, Franks J, Jeevan T, Fabrizio T, DeBeauchamp J, Kercher L, Seiler P, Gonzalez-Reiche AS, Sordillo EM, Chang LA, van Bakel H, Simon V, Consortium Mount Sinai Pathogen Surveillance (PSP) study group, Douek DC, Sullivan NJ, Thackray LB, Ueki H, Yamayoshi S, Imai M, Perlman S, Webby RJ, Seder RA, Suthar MS, García-Sastre A, Schotsaert M, Suzuki T, Boon ACM, Diamond MS, Kawaoka Y. 2022. SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. Nature 603:687–692.

31. Stewart R, Yan K, Ellis SA, Bishop CR, Dumenil T, Tang B, Nguyen W, Larcher T, Parry R, Sng JDJ, Khromykh AA, Sullivan RKP, Lor M, Meunier FA, Rawle DJ, Suhrbier A. 2023. SARS-CoV-2 omicron BA.5 and XBB variants have increased neurotropic potential over BA.1 in K18-hACE2 mice and human brain organoids. Front Microbiol 14:1320856.

32. Zhang Z, Zhou L, Liu Q, Zheng Y, Tan X, Huang Z, Guo M, Wang X, Chen X, Liang S, Li W, Song K, Yan K, Li J, Li Q, Zhang Y, Yang S, Cai Z, Dai M, Xian Q, Shi Z-L, Xu K, Lan K, Chen Y. 2024. The lethal K18-hACE2 knock-in mouse model mimicking the severe

pneumonia of COVID-19 is practicable for antiviral development. Emerg Microbes
Infect 13:2353302.

33. Perez-Zsolt D, Muñoz-Basagoiti J, Rodon J, Elousa M, Raïch-Regué D, Risco C, Sachse
M, Pino M, Gumber S, Paiardini M, Chojnacki J, Erkizia I, Muñiz X, Ballana E, Riveira-
Muñoz E, Noguera M, Paredes R, Trinité B, Tarrés-Freixas F, Blanco I, Guallar V,
Carrillo J, Blanco J, Telenti A, Heyn H, Segalés J, Clotet B, Martinez-Picado J, Vergara-
Alert J, Izquierdo-Useros N. 2021. Siglec-1 on dendritic cells mediates SARS-CoV-2
trans -infection of target cells while on macrophages triggers proinflammatory
responses. bioRxiv. Cold Spring Harbor Laboratory
https://doi.org/10.1101/2021.05.11.443572.

34. Rodon J, Muñoz-Basagoiti J, Perez-Zsolt D, Noguera-Julian M, Paredes R, Mateu L,
Quiñones C, Perez C, Erkizia I, Blanco I, Valencia A, Guallar V, Carrillo J, Blanco J,
Segalés J, Clotet B, Vergara-Alert J, Izquierdo-Useros N. 2021. Identification of
Plitidepsin as Potent Inhibitor of SARS-CoV-2-Induced Cytopathic Effect After a Drug
Repurposing Screen. Front Pharmacol 12:646676.

35. Haagmans BL, van den Brand JMA, Raj VS, Volz A, Wohlsein P, Smits SL, Schipper D,
Bestebroer TM, Okba N, Fux R, Bensaid A, Solanes Foz D, Kuiken T, Baumgärtner W,
Segalés J, Sutter G, Osterhaus ADME. 2016. An orthopoxvirus-based vaccine
reduces virus excretion after MERS-CoV infection in dromedary camels. Science (80-
) 351:77–81.

36. Brustolin M, Rodon J, Rodríguez de la Concepción ML, Ávila-Nieto C, Cantero G,
Pérez M, Te N, Noguera-Julián M, Guallar V, Valencia A, Roca N, Izquierdo-Useros N,
Blanco J, Clotet B, Bensaid A, Carrillo J, Vergara-Alert J, Segalés J. 2021. Protection
against reinfection with D614- or G614-SARS-CoV-2 isolates in golden Syrian
hamster. Emerg Microbes Infect 10:797.

37. Dowall SD, Graham VA, Tipton TRW, Hewson R. 2009. Multiplex cytokine profiling
with highly pathogenic material: use of formalin solution in luminex analysis. J
Immunol Methods 348:30–35.

38. Pradenas E, Trinité B, Urrea V, Marfil S, Ávila-Nieto C, Rodríguez de la Concepción
ML, Tarrés-Freixas F, Pérez-Yanes S, Rovirosa C, Ainsua-Enrich E, Rodon J, Vergara-
Alert J, Segalés J, Guallar V, Valencia A, Izquierdo-Useros N, Paredes R, Mateu L,
Chamorro A, Massanella M, Carrillo J, Clotet B, Blanco J. 2021. Stable neutralizing
antibody levels 6 months after mild and severe COVID-19 episodes. Med 2:313-
320.e4.

39. Trinité B, Tarrés-Freixas F, Rodon J, Pradenas E, Urrea V, Marfil S, Rodríguez de la
Concepción ML, Ávila-Nieto C, Aguilar-Gurrieri C, Barajas A, Ortiz R, Paredes R,
Mateu L, Valencia A, Guallar V, Ruiz L, Grau E, Massanella M, Puig J, Chamorro A,
Izquierdo-Useros N, Segalés J, Clotet B, Carrillo J, Vergara-Alert J, Blanco J. 2021.
SARS-CoV-2 infection elicits a rapid neutralizing antibody response that correlates
with disease severity. Sci Rep 11:2608.

Figure Legends

Figure 1. Expression of hACE2 in the Col1a1-K18-hACE2 KI model. (A) Schematic

representation of the insertion strategy. The original K18-hACE2 transgene was inserted

into the collagen COL1A1 locus using a recombinase mediated cassette exchange (RMCE)

FLP-FRT system in KH2 cells via blastocyst injection. pPGK-ATG-frt plasmid: vector backbone

with Ampicillin resistant gene (Amp), transcription start site (ATG) and Flippase recognition

target (Frt). Hygro: Hygromicin resistance gene. pCAGGS-FlpE: expression plasmid for FLPe

recombinase expression. (B) Relative quantification of hACE2 receptor expression to GAPDH

expression in uninfected Col1a1-K18-hACE2 ((blue empty dot, n=3) and K18-hACE2 males

(black empty square, n=3). Delta Ct values are inversely shown to facilitate interpretation.

The lower the absolute number, the higher the relative expression. Solid line and bars

represent mean and SEM. (C) Relative comparison of hACE2 receptor expression in Col1a1-

K18-hACE2 versus K18-hACE2 mice (n=3 each, males). Higher receptor expression in Col1a1-

K18-hACE2 (negative values in Y axis) is marked by blue bars, and higher expression in K18-

hACE2 (positive values in Y axis) is marked in black bars. (D) Western Blot analysis of hACE2,

mACE2 and GAPDH in representative tissues. hACE2 signal was obtained using a specific

monoclonal antibody (top panels); mACE2 and hACE2 signal was obtained with a cross-

reactive polyclonal antibody (middle panel), and anti-GAPDH was analyzed as reference

(bottom panels). Molecular weight marker is shown on the right.

Figure 2. Experimental setting and progression of SARS-CoV-2 infection in Col1a1-K18-

hACE2 and K18-hACE2 mouse models. (A) Schematic representation of the experimental

setting. Knock-in Col1a1-K18-hACE2 (n=27) and transgenic K18-hACE2 mice (n=4) were

intranasally challenged with a 1000 TCID$_{50}$ dose of a B.1 SARS-CoV-2 isolate. A Col1a1-K18-

hACE2 uninfected control group (n=5) was challenged with PBS. Mice were monitored for

weight-loss and clinical signs for 14 days post infection (dpi). Euthanasia was performed at

3, 7, and 14 dpi or upon fulfilment of humane endpoint criteria, for sample and tissue

collection (n=8 per timepoint). Infections were performed in two separate experiments

between January and June 2022. Created with Biorender.com. (B) Relative body weight

follow-up referred to day 0. Col1a1-K18-hACE2 uninfected (blue empty dot), Col1a1-K18-

hACE2 infected (blue dot), K18-hACE2 infected (black square). Solid lines and bars represent

mean±SD. (C) Survival (Kaplan-Meier). All K18-hACE2 infected animals (n=4) had to be

euthanized due to endpoint criteria by 7dpi, and only three infected Col1a1-K18-hACE2 at

8, 9, and 10 dpi. No uninfected Col1a1-K18-hACE2 had to be euthanized under this criterion.

Col1a1-K18-hACE2 uninfected (blue dashed line), Col1a1-K18-hACE2 infected (blue line),

K18-hACE2 infected (black line). Statistical differences were identified using a Log-rank

(Mantel-Cox) test (<0.0001), followed by individual comparisons (**p < 0.005; ****p <

0.0001)

Figure 3. Progression of SARS-CoV-2 infection in Col1a1-K18-hACE2 KI mice. Col1a1-K18-

hACE2 (blue circles, *n=27*) and K18-hACE2+ mice (black squares, *n=4*) were inoculated with

1000 TCID$_{50}$ of a SARS-CoV-2 B.1 isolate (full shapes) or uninfected (empty shapes of each

colour, *n=5*). Animals were euthanized at 3dpi (n=8), 7dpi (n=8), 14 dpi (n=8) or upon

fulfilment of humane endpoint criteria **(n=3, euthanized at 8,9 and 10 dpi)**. (A) SARS-CoV-2

viral RNA loads (copies/mL) of oropharyngeal swab, lung, and nasal turbinate samples.

Dashed line represents limit of detection, established by 2SD of uninfected animals.

Statistical differences were identified using a Peto-Peto Left censored test (*p < 0.05,

**p<0.005). (B) Viral titration of replicative virus ($TCID_{50}$/mL) in lung samples from B.1

infected mice at different endpoints in Vero E6 cells on day 5 of culture. Titers were

compared using an Independence Asymptotic Generalized Pearson Chi-Squared Test for

ordinal data (*p < 0.05, **p<0.005). (C) Lung histopathological scoring of broncho-

interstitial pneumonia (left) and SARS-CoV-2 NP immunohistochemical (IHC) scoring (right)

in both models at 3, 7, 14 dpi and endpoint. Statistical differences were identified using an

Independence Asymptotic Generalized Pearson Chi-Squared Test for ordinal data (*p <

0.05). (D) Representative lung histology and IHC pictures of both models at 3, 7, 14 dpi and

endpoint. Images show low-power magnification bars (200 μm).

Figure 4. Inflammatory response in the lung. The concentration of Inflammatory cytokines

in lung extracts in the knock-in and transgenic models at 3,7,14 dpi or endpoint are shown.

Col1a1-K18-hACE2 infected (blue full circles, *n=23*), uninfected (blue empty circles, n=3) and

K18-hACE2 infected mice (black full squares, *n=4*). The bar shows Median with interquartile

range. The limit of detection for each cytokine is indicated by a dotted line. Statistical

differences were identified using a Kruskal Wallis, and a Conover's nonparametric all-pairs

comparison test (*p < 0.05, **p<0.005).

Figure 5. Tissue Tropism of SARS-CoV-2 in Col1a1-K18-hACE2 and K18-hACE2 Mice. SARS-

CoV-2 viral RNA loads (copies/mL) were analyzed by RT-qPCR in Col1a1-K18-hACE2 mice

tissues collected at 3,7 and 14 dpi (n=3 per timepoint) or upon fulfilment of humane

endpoint criteria (n=1 at 8 dpi, n=1 at 9 dpi). K18-hACE2 tissues were collected at endpoint

at 6 (n=1) and 7 dpi (n=1). Solid lines show the mean ±SD. Dashed line represents limit of

detection, established by 2SD of uninfected animals.

Figure 6. Progression of SARS-CoV-2 infection in brain from the Col1a1-K18-hACE2 KI mice.

Col1a1-K18-hACE2 (blue circles, *n=27*) and K18-hACE2+ mice (black squares, *n=4*) were

inoculated with 1000 TCID$_{50}$ of a B.1 SARS-CoV-2 isolate (full shapes) or uninfected (empty

shapes, *n=5*) and followed until 14 dpi. Samples were collected at 3, 7, 14 dpi and endpoint.

(A) SARS-CoV-2 viral RNA loads (copies/mL) in brain extracts. Dashed line represents limit

of detection, established by 2SD of uninfected animals. Statistical differences were

identified using a Peto-Peto Left-censored samples test with correction for multiple

comparisons. Solid line and bars represent mean and SD. (B) Viral titration of replicative

virus (TCID$_{50}$/mL) in brain samples from B.1 infected mice at different endpoints in Vero E6

cells on day 5 of culture. Solid line and bars represent mean and SD. (C) Brain

histopathological scoring of multifocal lymphoplasmacytic meningo-encephalitis in brain

(Left) and SARS-CoV-2 NP IHC scoring (right) in brain of both models. Statistical differences

were identified using an Independence Asymptotic Generalized Pearson Chi-Squared Test

for ordinal data (*p < 0.05; **p < 0.01). (D) Representative images of brain Histology and

IHC images of both hACE2+ mouse models. Images show low-power magnification (top;

bars: 600μm) and medium-power magnification (bottom; bars: 300μm). Col1a1-K18-hACE2

KI mice samples shown were collected at 7 dpi, while K18-hACE2 samples were collected at

humane endpoint (6-7dpi).

A

B

hACE2 Receptor Expression

ΔCt value (hACE2 - GAPDH)

Pancreas, Lymph nodes, Salivary, Spleen, Intestine, Muscle, Nasal Turbinate, Lung, Kidney, Heart, Brain, Liver

○ Col1a1-K18-hACE2   □ K18-hACE2

C

Relative hACE2 Receptor Expression

ΔΔCt value (Col1a1-K18 vs K18)

Higher expression in K18-hACE2

Higher expression in Col1a1-K18-hACE2

Pancreas, Lymph nodes, Salivary, Spleen, Intestine, Muscle, Nasal Turbinate, Lung, Kidney, Heart, Brain, Liver

D

	Lung			Brain				Muscle			Pancreas				Liver			Spleen		
	K18	KI	WT	K18	KI	WT		K18	KI	WT	K18	KI	WT		K18	KI	WT	K18	KI	WT
moACE2																				
pACE2																				
GAPDH																				

125 kDa
125 kDa
93 kDa
42 kDa
31 kDa

A

B