Title: **A human-ACE2 knock-in mouse model for SARS-CoV-2 infection recapitulates respiratory disorders but avoids neurological disease associated with the transgenic K18-hACE2 model.**

Running title: B.1 SARS-CoV-2 infection of a new hACE-2 Knock-in mice

Anna Pons-Grífols[1], Ferran Tarrés-Freixas[1,2,3,8], Mònica Pérez[2,3], Eva Riveira-Muñoz[1], Dàlia Raïch-Regué[1], Daniel Pérez-Zsolt[1], Jordana Muñoz-Basagoiti[1], Barbara Tondelli[4], Nuria Izquierdo-Useros[1,5,6], Sara Capdevila[7], Júlia Vergara-Alert[2,3], Victor Urrea[1], Jorge Carrillo[1,5,6], Ester Ballana[1,5], Stephen Forrow[4], Bonaventura Clotet[1,8], Joaquim Segalés[2,9], Benjamin Trinité[1,#], Julià Blanco[1,5,6,8,#].

[1]IrsiCaixa, Can Ruti Campus, Badalona, Spain.
[2]Unitat mixta d'investigació IRTA-UAB en Sanitat Animal, Centre de Recerca en Sanitat Animal (CReSA), Campus de la Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain.
[3]IRTA, Programa de Sanitat Animal, Centre de Recerca en Sanitat Animal (CReSA), Campus de la Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain.
[4]Mouse Mutant Core Facility (PBA2B2), Institute for Research in Biomedicine (IRB), Parc Científic de Barcelona, Barcelona, Spain.
[5]Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain.
[6] CIBER Infectious Diseases (CIBERINFEC), Carlos III Health Institute, Madrid, Spain.
[7]Comparative Medicine and Bioimage Centre of Catalonia (CMCiB), Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain.
[8] University of Vic-Central University of Catalonia (UVic-UCC), Vic, Spain.
[9]Departament de Sanitat i Anatomia Animals, Facultat de Veterinària, Universitat Autònoma de Barcelona (UAB), Campus de la UAB, Bellaterra, Spain.

 # Corresponding authors

Benjamin Trinité: btrinite@irsicaixa.es

Julià Blanco: jblanco@irsicaixa.es

JS, BT and JB are co-senior authors

33   Abstract Word count: 245
34   Text word count: 4955
35

Abstract

Animal models have been instrumental in elucidating the pathogenesis of SARS-CoV-2 infection and testing COVID-19 vaccines and therapeutics. Wild-type (WT) mice are not susceptible to many SARS-CoV-2 variants, therefore transgenic K18-hACE2 mice have emerged as a standard model system. However, this model is characterized by severe disease, particularly associated with neuroinfection, which leads to early humane endpoint euthanasia. Here, we established a novel knock-in (KI) mouse model by inserting the original K18-hACE2 transgene into the collagen COL1A1 locus using a recombinase mediated cassette exchange (RMCE) system. Once the Col1a1-K18-hACE2 mouse colony was established, animals were challenged with a B.1 SARS-CoV-2 (D614G) isolate and were monitored for up to 14 days. Col1a1-K18-hACE2 mice exhibited an initial weight loss similar to the K18-hACE2 transgenic model but did not develop evident neurologic clinical signs. The majority of Col1a1-K18-hACE2 mice did not reach the preestablished humane endpoint, showing progressive weight gain after 9 days post-infection (dpi). Importantly, despite this apparent milder pathogenicity compared to the K18-hACE2 transgenic model, high levels of viral RNA were detected in lungs, oropharyngeal swab, and nasal turbinate. Conversely, in sharp contrast to K18-hACE2 transgenic mice, no viral replication was detected in the brains of Col1a1-K18-hACE2 animals at any timepoint, explaining the reduced severity of clinical signs. At 14 dpi, while infection was cleared in the lungs, increased lesions and residual inflammation were detected. Overall, Col1a1-K18-hACE2 mice constitute a new model for investigating SARS-CoV-2 pathogenesis and treatments, with potential implications for studying long-term COVID-19 sequelae.

Importance

K18-hACE2 mice express high levels of the human protein ACE-2, the receptor for SARS-

CoV-2, and therefore are infected by this virus. These animals have been crucial to

understand viral pathogenesis and to test COVID-19 vaccines and antiviral drugs.

However, K18-hACE2 rapidly die after infection with initial SARS-CoV-2 variants due to

a massive brain infection that does not occur in humans. Here, we used a technology

known as knock-in that allows for the targeted insertion of a gene into a mouse and we

have generated a new hACE2-mouse. We have characterized this new animal model

demonstrating that the virus replicates in the respiratory tract, damaging and inflaming

the lungs; however, in contrast to K18-hACE2 mice, no brain infection was observed,

and most animals recovered from infection. This new model could be instrumental for

the study of specific disease aspects such as post-COVID condition, sequelae, and

susceptibility to reinfection.

Keywords: animal model, lung inflammation, lung damage, neuroinvasion, post covid

condition.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is the etiologic agent of the coronavirus disease 2019 (COVID-19). The COVID-19 pandemic fuelled an unprecedented collaborative research effort to develop therapeutics and vaccines. Animal models that recapitulate key clinical and pathological features of COVID-19 have played a pivotal role in testing novel vaccines, antivirals and other treatments (1).

The spike glycoprotein of SARS-CoV-2 uses the angiotensin-converting enzyme 2 (ACE2) as receptor to enter and infect host cells (2). ACE2 is expressed on the surface of cells of different organs such as lungs, intestine, kidneys, and heart. However, the spike glycoprotein from ancestral SARS-CoV-2 strains does not efficiently bind to mouse ACE2 (mACE2), rendering wild-type (WT) mice refractory to infection. Therefore, several strategies were used to develop mouse models susceptible to infection. These strategies include viral adaptation (3) or the introduction of the human ACE2 (hACE2) receptor via viral vectors and transgenic approaches (4–8).

Both K18-hACE2 transgenic mice and Golden Syrian hamsters (GSH) became reference animal models for investigating SARS-CoV-2 pathogenesis in vivo. They mainly differ in pathogenesis: GSH, which are naturally susceptible to SARS-CoV2 infection, recapitulate a milder disease phenotype (1). The transgenic mouse model was originally developed for the study of SARS-CoV, which also targets human ACE2. It was created by random insertion of multiple copies of the *hACE2* gene under the control of the human cytokeratin 18 gene promoter (*KRT18* or K18). This promoter allows for high-level expression and is specific to epithelial cells, including those in the airways (9, 10). This approach allows the infection of

mice by SARS-CoV-2, while maintaining the activity of the mACE2 receptor. Infection of K18-

hACE2 mice with pre-Omicron variants of SARS-CoV-2 courses with progressive weight-loss

and strong clinical signs by 3-5 days post-infection (dpi), a severity which requires

euthanasia by 5-7 dpi. The lethality in this model is dose- and SARS-CoV-2 variant-

dependent (10, 11), and has been associated with neuroinvasion and extensive brain

infection (McCray et al., 2007; Carossino et al., 2022; Tarrés-Freixas et al., 2022; Vidal et al.,

2022). Neuroinvasion in K18-hACE2 transgenic mice has been linked to the aleatory gene

insertion and the high number of inserted gene copies, around 8 in the commercial *B6.Cg-*

Tg(K18-ACE2)2Prlmn/J-K18-hACE2 mouse (https://www.jax.org/strain/034860), leading to

altered expression levels and tissue distribution of the receptor (15). Specifically, a higher

number of gene copies correlates with a worse disease prognosis and encephalitis in this

model, following SARS-CoV infection (16). These neurological lesions are not consistent with

the pathology of COVID-19 in humans, where infection of the central nervous system was

not observed at such level (17). Therefore, new animal models that better recapitulate the

human disease, without viral neuroinvasion, should be developed.

Knock-in (KI) models are characterized by the targeted insertion of a defined gene copy

number in a specific locus, therefore providing a more predictable expression of the

transgene. In this work, we used a recombinase-based approach to insert the original K18-

hACE2 transgene into the collagen type I alpha chain (COL1A1) locus to generate a Col1a1-

K18-hACE2 KI mouse. To characterize this new model, mice were challenged with a SARS-

CoV-2 B.1 (D614G) isolate and pathogenicity was compared with the well-characterized

K18-hACE2 transgenic mice. After viral challenge of a Col1a1-K18-hACE2 mice, we

confirmed viral replication and lesions in the lungs, but absence of viral neuroinvasion.

Results

Characterization of hACE2 expression in the Col1a1-K18-hACE2 mice

We generated a new hACE2 KI mouse model through addition of the original K18-hACE2

transgene, kindly supplied by Paul McCray (9) at the COL1A1 locus using a recombinase

mediated system (**Figure 1A**) (18). Traditional transgenic techniques rely on the random

genomic insertion of an unpredictable number of transgene copies. This frequently results

in a highly variable level and pattern of transgene expression, due to cis acting regulatory

elements at the site of insertion. A single copy of a transgene, inserted into the well

characterised Col1A1 locus, is expected to result in a more predictable level and pattern of

expression of the transgene sequence. To confirm the correct expression of the transgene

in this KI model, we quantified *hACE2* transcripts relative to the house-keeping gene GAPDH

by RT-qPCR in RNA samples extracted from twelve different tissues. Additionally, the same

analysis was performed in K18-hACE2 mice to compare both models. Expression of the

hACE2 transgene in K18-hACE2 mice was similar among most tested tissues except in

muscle and heart, which showed lower values (**Figure 1B).** In contrast, hACE2 expression in

Col1a1-K18-hACE2 mice was more heterogenous, with the lowest expression observed in

the brain and liver. When comparing the relative expression of hACE2 between both

models, we observed higher expression in the pancreas, lymph nodes and muscle of Col1a1-

K18-hACE2 mice. Conversely, in the kidney, brain and liver, expression was higher in the

K18-hACE2 mouse model. Minor differences were observed in the rest of the analyzed

tissues   (**Figure   1C**).

Survival of Col1a1-K18-hACE2 mice following SARS-CoV-2 B.1 infection.

Col1a1-K18-hACE2 mice (n=27) and K18-hACE2 mice (n=4) were challenged intranasally

with a SARS-CoV-2 B.1 (D614G) isolate. Uninfected KI animals, used as a control, received

an intranasal dose of PBS (n=5). Animal weight and clinical signs were monitored daily for

14 days after infection. The endpoints to collect samples were set at 3-, 7-, and 14-days

post-infection (dpi, n=8 Col1a1-K18-hACE2 mice per endpoint) or upon fulfilment of

humane endpoint criteria (**Figure 2A**).

Both animal models started losing weight around 3 dpi, consistent with previous studies

with B.1 isolates in transgenic animals (**Figure 2B**) (13). In K18-hACE2 transgenic animals,

the decrease in weight was associated with a dramatic increase of clinical signs: mobility,

dyspnoea, and neurological effects (Supplementary Table 1). This severity fulfilled the

humane endpoint criteria, leading to euthanasia of all K18-hACE animals by 6 or 7dpi (**Figure**

2C). In contrast, while Col1a1-K18-hACE2 mice underwent weight loss, they did not show

severe clinical signs apart from a slight decrease in activity. Three Col1a1-K18-hACE2

animals had to be euthanized (exitus) due exclusively to a weight-loss exceeding the 20%

limit (humane endpoint) at 8, 9 and 10 dpi. But interestingly, all other KI animals started to

regain weight after 9 dpi, reaching 90% of their initial weight by 14 dpi (**Figure 2B/C**). These

results indicate that B.1 SARS-CoV-2 infection in Col1a1-K18-hACE2 mice is significantly less

lethal compared to k18-hACE2 transgenic mice (**Figure 2C**).

To explore the extent of viral infection in both animal models, viral loads (VL, cell-free viral

RNA) were quantified in a large set of samples (oropharyngeal swabs, lung, brain, nasal

turbinate, muscle, intestine, liver, kidney, pancreas, salivary glands, lymph nodes and

spleen) collected from a subset of animals (n=20). Respiratory tract tissues showed

detectable VL in both models, while the brain showed high VL exclusively in K18-ACE2

transgenic samples. In both models, VL were mostly undetected in all other tissues except

for low levels found in the heart and salivary glands (**Supplementary Figure 1**).

B.1 replicates in the respiratory tract of Col1a1-K18-hACE2 KI mice.

To fully characterize the kinetics of viral replication in the respiratory tract, samples from

oropharyngeal swab, lung, and nasal turbinate were collected at 3, 7, and 14 dpi or at the

humane endpoint (exitus) in all Col1a1-K18-hACE2 mice. Samples from K18-hACE2

transgenic mice obtained at euthanasia (6 and 7 dpi) were used as references. Virological

analysis in oropharyngeal swab, lung, and nasal turbinate revealed widespread infection in

both K18-hACE2 and Col1a1-K18-hACE2 mice in all mentioned tissues across two

independent experiments (**Figure 3A**). VL in the nasal turbinate were similar in both animal

models, with some viral RNA still detected at 14 dpi in Col1a1-K18-hACE2 mice. RNA levels

in oropharyngeal swabs and lung peaked at 3 dpi in Col1a1-K18-hACE2 mice and tended to

decay overtime up to 14 dpi. Col1a1-K18-hACE2 animals euthanized at the humane

endpoint at 8-10 dpi still displayed high viral RNA in lungs, which could explain their more

pronounced weight-loss and early fulfilment of endpoint criteria. To further confirm viral

clearance, titration of replication-competent virus was performed in lung samples. Similar

to the VL kinetics, viral titers peaked at 3 dpi, followed by a quick and significant decrease

(**Figure 3B,** 3 dpi vs 7 dpi: p = 0.00178).

To further characterize viral pathogenesis, formalin-fixed lung samples from both animal

models were analyzed by histology and immunohistochemistry (IHC) to score the presence

of lesions and SARS-CoV-2 antigen (nucleoprotein, NP), respectively. Histological lesions

were semi-quantified as none, mild, moderate, or severe (score 0-3), and the detection of

viral antigen as lack, low, moderate, or high (score 0-3). In Col1a1-K18-hACE2 KI mice,

histopathological analysis showed an increase in lung lesion average overtime, reaching a

maximum score of two at 14 dpi in most animals **(Figure 3C and D).** IHC detection of SARS-

CoV-2 NP in the lungs of Col1a1-K18-hACE2 KI mice was consistent with the VL analyses,

showing the highest score early upon infection (3 dpi). Antigen detection decreased with

time showing apparent clearance by 14 dpi **(Figure 3C and D).**

To assess infection-driven inflammation, the levels of IP-10, IL-6, IFNγ, MCP-1 and MIP-1β

were measured by Luminex in lung samples from both animal models at the indicated

timepoints. While all cytokines were increased in infected animals, compared to uninfected

controls, IP-10 and IL-6 showed a longitudinal dynamic comparable to viral load and IHC

data in Col1a1-K18-hACE2 KI mice, peaking at 3 dpi and significantly decreasing by 14 dpi

(**Figure 4**). In contrast, IFNγ, MCP-1, and MIPβ peaked at 7 dpi and decayed afterwards,

although no statistical differences were observed among the analyzed timepoints. Overall,

levels of inflammatory mediators at 7 dpi in Col1a1-K18-hACE2 KI mice were comparable to

K18-hACE2 transgenic mice at the time of euthanasia.

Col1a1-K18-hACE2 KI mice do not develop brain infection.

To investigate the discrepancy in VL observed in the brain (**Supplementary Figure 1**) and

the differences in survival and clinical signs observed between each model (**Figure 2C**), we

performed a longitudinal virological and histological analysis of brain samples. VL were

detected at low levels in only 2 out of 8 Col1a1-K18-hACE2 KI mice analyzed at 7 dpi and

importantly, no VL was found in animals reaching the humane endpoint. In contrast, all K18-

hACE2 brain samples showed high VL at euthanasia (6-7dpi) (**Figure 5A**). Similar results were

obtained when formalin-fixed brain samples were analyzed for histopathology and IHC,

scored in a blinded fashion as described above. Histopathological analysis of the brain in the

Col1a1-K18-hACE2 model showed no lesions at any of the analyzed timepoints. In contrast,

brain samples from K18-hACE2 infected animals showed lesion scores of 2 and 1 **(Figure 5B**

and C). Furthermore, no detection of SARS-CoV-2 NP in the brain of Col1a1-K18-hACE2 mice

was observed by IHC at any timepoint or at the humane endpoint, while K18-hACE2 mice

showed a high density of antigen diffused throughout the tissue (score 3) at euthanasia

(Figure 5C). Brain lesions in the transgenic model were characterized as multifocal

lymphoplasmacytic meningoencephalitis and were observed alongside the detection of

viral antigen in the brain (**Figure 5C**).

Discussion

In this study, we characterized a novel hACE2 KI murine model by evaluating the course of infection with the B.1 D614G SARS-CoV-2 variant with a direct comparison to the well-described K18-hACE2 transgenic mouse model, in which this variant is known to be highly pathogenic. The most relevant findings included the susceptibility of the KI model to SARS-CoV-2 infection and the reduced lethality imputed to the absence of viral replication and lesions in the brain.

The development of this KI model was started at the beginning of the COVID-19 pandemic as an alternative to the K18-hACE2 transgenic mouse model. Even though the transgenic model had already been established and used in previous SARS-CoV studies (9), the availability of these animals was limited at first, as colonies needed to be restarted from cryopreserved embryos. In this context, several laboratories developed new hACE2 murine models for SARS-CoV-2 infection. Expression of the hACE2 transgene was achieved through various methods, including directed insertion using CRISPR/Cas9 technology (19) using different promoters such as Hepatocyte nuclear factor-3/forkhead homologue 4 (Hfh4) promoter (20) or the mouse ACE2 promoter (6), and by infecting mice with replication-defective adenovirus encoding hACE2 (3). In this work, we followed a different strategy wherein a single copy of the transgene was inserted into the COL1A1 locus, but under the K18 promoter as the original transgenic K18-hACE2 model. The selection of this locus was based on its known ability to provide reliable and ubiquitous expression of inserted sequences (18). Additionally, RMCE allowed the insertion of a single transgene copy which should better mimic physiological expression, potentially impacting lethality through

reduction of the ectopic expression and modification of tissue distribution of the hACE2

receptor. A recent study identifying a correlation between lower hACE2 copy number and

reduced pathology further supports our initial hypothesis (21).

When comparing with K18-hACE2 transgenic mice, B.1 infection of the new Col1a1-K18-

hACE2 KI mouse model resulted in much less severe clinical signs with better responsiveness

and physical appearance and no neurological sign. These observations were consistent with

a lack of detectable viral replication in brain of Col1a1-K18-hACE2 KI mice, as summarised

in **Figure 6**. Focusing on the respiratory tract, viral replication was observed in both models,

although VL and antigen detection were in general lower at 7 dpi in Col1a1-K18-hACE2 KI

mice. In contrast, lung tissue lesions and inflammation were fully comparable in both

models at this timepoint (**Figure 6**).

To explain the different behaviour between each model, hACE2 expression was analyzed in

different tissues. While expression in the lungs and nasal turbinates was comparable, the

brain and liver showed lower expression of hACE2 in Col1a1-K18-hACE2 KI mice. This finding

better mimics the physiological distribution in humans, where hACE2 expression in the brain

is low (22). Expression of mACE2 was not analyzed, as mice are not susceptible to the B.1

variant through the murine receptor, and no changes in their levels or distribution were

expected compared to WT mice (15, 23). Among the clinical parameters analyzed in Col1a1-

K18-hACE2 KI infected mice, we found that initial weight-loss was similar to that of the K18-

hACE mice but was not accompanied by the increased severity of clinical signs that

characterizes infection in these transgenic mice (neurological signs, lack of responsiveness,

and poor general appearance) (13). Consistent with this observation, infection of Col1a1-

K18-hACE2 KI mice coursed with a partial recovery in weight by 14 dpi in most animals.

Importantly, the survival rate of Col1a1-K18-hACE2 KI mice was over 70%, with the three

deaths exclusively due to a weight loss of more than 20%, which was one of the humane

endpoints in our study.

Among the tissues analyzed in Col1a1-K18-hACE2 KI infected mice, we found significant

replication of SARS CoV-2 in the respiratory tract, mainly in the lungs. Despite viral clearance

in Col1a1-K18-hACE2 KI mice which was confirmed by viral titration and IHC, lung lesions

were similar between both models by 7dpi and remained after 14 dpi in KI animals

recovering from the infection. The impact of viral replication in lungs, on local expression of

inflammatory markers, was also confirmed and found to be comparable to the inflammation

profile reported in K18-hACE2 transgenic mice (24, 25). Infection of Col1a1-K18-hACE2 KI

mice induced higher levels of IFN-g, IL-6, IP-10, MCP-1 and MIP-1b cytokines. The

longitudinal analysis showed early impact on IL-6 and IP-10 levels (3 dpi) and a clear trend

to normalization by 14 dpi, with some residual inflammation remaining. This is consistent

with the presence of lesions at this timepoint, and the immune cell infiltration described by

21 dpi in K18-hACE2 transgenic mice surviving infection (25).

The main difference between the Col1a1-K18-hACE2 KI and the K18-hACE2 transgenic

models was found in neuroinvasion. Only residual levels of viral RNA were detected in the

brains of Col1a1-K18-hACE2 KI mice. Furthermore, no detectable brain lesions, such as the

multifocal lymphoplasmacytic meningoencephalitis reported in K18-hACE2 transgenic mice

(14) were detected by histology. A complete absence of SARS-CoV-2 nucleoprotein in the

brain was evidenced by IHC. Importantly, the three animals that reached humane endpoints

did not present any evidence of brain viral replication but showed enhanced lung infection.

These observations are consistent with the lower level of hACE2 expression in the brain of

Col1a1-K18-hACE2 KI compared to the K18-hACE2 transgenic animals. Therefore, the

targeted insertion of the hACE2 transgene seems to be a crucial factor leading to lower

levels of brain infection and neurological clinical signs. However, different studies have

shown that neuroinvasion is only partially dependent on the expression of hACE2 receptor

(12), and that both delivery route and viral dose can also play a role in the magnitude of the

neuroinvasion (22, 26). In our case, to exclusively analyze the effect of receptor expression,

we kept the intranasal administration route and $TCID_{50}$ dose established and characterized

in our own and others' studies to ensure comparability of the results (13).

Our study is a preliminary characterization of a new animal model and has consequently

several limitations. First, the expression of hACE-2 in different tissues has only been

analyzed at the mRNA level, with no data on immunohistochemistry to confirm the

presence of the protein. Moreover, we have focused on the characterization of one viral

variant and a single dose. The B.1 (D614G) variant was selected for its demonstrated

severity in K18-hACE2 mice and its inability to use mACE2 as an entry receptor, unlike

B.1.351/Beta and BA1.1/Omicron variants (13, 15). While ancestral and earlier variants

(B.1.1.7/Alpha, B.1.351/Beta B.1.617.2/Delta) were highly pathogenic in K18-hACE2 mice,

BA.1.1/Omicron caused a milder infection with no weight loss nor neuroinvasion, therefore

reducing lethality in this model (13). However, more recent Omicron subvariants such as

BQ.1.1, BA.5 and XBB.1.5 appear to be regaining pathogenicity in K18-hACE2 mice (27–29).

Increased lung infection, pro-inflammatory cytokines, and lung pathology are observed with

these variants, with data on neuroinvasion and neurovirulence on the latter (29). Further

characterization of the latest variants in the KI model would provide relevant insights into

the full impact of the KI insertion of the transgene and confirm the suitability of this model

for the study of both ancestral and emerging variants. Nevertheless, for the B.1 SARS-CoV-

2 variant it shows similarity with GSH in viral dynamics and pathology, with no

neuroinvasion. This new model, however, has the upside of a wider reagent availability,

scarce in GSH (1), and easier animal facility allowance.

In summary, several mouse models for the evaluation of antivirals and vaccines have been

developed to date, with disease phenotypes ranging from mild to severe COVID-19-like

condition. None of these models, however, can fully recapitulate all aspects of the disease

as it occurs in humans. The development and further characterization of new animal

models, like the Col1a1-K18-hACE2 model, may overcome some of these limitations and

provide valuable tools to study certain aspects of COVID-19. Survival of Col1a1-K18-hACE2

KI mice after infection with a highly pathogenic variant could provide a model which better

mimics human disease progression, and thus could be instrumental for the study of specific

disease aspects such as post-COVID condition (PCC), sequelae, long-term effects of drug

therapies, and susceptibility to reinfection.

Materials and methods
Creation of the Knock-in Model

A new KI mouse model, **B6;129S-Col1a1$^{tm1(K18-hACE2)Irb}$/Irsi** (Col1a1-K18-hACE2), was created

by inserting the original K18-hACE2 transgene into the collagen COL1A1 locus using a

recombinase-mediated cassette exchange (RMCE) FLP-FRT system in KH2 cells (18). The

actual insertion site lies approximately 0.3 kb downstream of the 3'UTR end of COL1A1. As

such, the inserted transgene remains identical to the original designed one (9) and the

hACE2 cDNA is under the control of the K18 promoter rather than COL1A1. Specifically, the

plasmid PGK-ATG-Frt was digested with EcoRV, and the sequence

ATCAGACGTCGCTAGCGGCGCGCCGGTACTAGT was inserted to create a multi-cloning site.

The plasmid containing hACE2 under the control of the K18 promoter (K18-hACE2), supplied

by Paul McCray, was digested with HpaI and XbaI enzymes. The resulting transgene

fragment was isolated and purified, and then cloned into the EcoRV-NheI sites of the

modified PGK-ATG-Frt plasmid to generate the targeting vector. The targeting vector was

then co-transfected with an Flp expression construct into KH2 cells by electroporation. The

cells were then placed under hygromycin selection for 9 days, after which drug resistant

colonies were picked, expanded, and screened for the presence of the targeted transgene.

Once confirmed, the modified embryonic stem cells were injected into mouse blastocysts.

These injected mouse blastocysts were then placed into recipient foster females via embryo

transfer techniques.

Biosafety Approval and Virus Isolation

The execution of SARS-CoV-2 experiments was approved by the biologic biosafety

committee of Germans Trias i Pujol Research Institute (IGTP) and performed at the Biosafety

Level 3 laboratory (BSL-3) of the Comparative Medicine and Bioimage Centre (CSB-20-015-

M8; CMCiB, Badalona, Spain).

The B.1 SARS-CoV-2 isolate used in this study was isolated from nasopharyngeal swabs of

hospitalized patients in Spain as described elsewhere (30, 31). Briefly, viruses were

propagated in Vero E6 cells (CRL-1586; ATCC, Virginia, VA, United States) for two passages

and recovered by supernatant collection. The sequence of the SARS-CoV-2 variant tested is

deposited at the GISAID Repository with accession IDs EPI_ISL_510689. EPI_ISL_510689 was

the first SARS-CoV-2 virus isolated in Catalonia in March 2020 and, compared to the Wuhan/

Hu-1/2019 (WH1) strain, this isolate had the S protein mutations D614G, which is associated

with the B.1 lineage, and R682L. Viral stocks were titrated on Vero E6 cells to use equivalent

TCID50/mL using the Reed-Muench method and sequential 1/10 dilutions of the viral stocks

as described previously (31).

Animal Procedures and Study Design

All animal procedures were approved by the Committee on the Ethics of Animal

Experimentation of the IGTP and were authorized by the *Generalitat de Catalunya* (code:

11222). All animal experiments followed the principles of animal welfare and the 3Rs. All

experiments and sample processing were performed inside the BSL-3 facility. Col1a1-K18-

hACE2 hemizygous KI mice were produced and bred at Parc Científic de Barcelona (PCB) by

pairing hemizygous males for Tg(K18-ACE2)2Prlmn (or K18-hACE2) with non-carrier

C57Bl6/J females. Animals used as control, B6.Cg-Tg(K18-ACE2)2Prlmn/J (or K18-hACE2)

hemizygous transgenic mice (034860, Jackson Immunoresearch, West Grove, PA, United

States) were bred at CMCiB in the Specific Pathogen Free Area (SPF) by pairing hemizygous

K18-hACE2 males with non-carrier C57Bl6/J females. The genotype of the offspring

regarding both K18-hACE2 and Col1a1-K18-hACE2 mice was determined by qPCR at the

IGTP's Genomics Platform from tail samples. Both animal models were kept in the BSL-3

facility during the whole experiment including acclimatization period. The housing

conditions in the BSL-3 room were maintained as follows: a temperature of 22±2ºC,

humidity levels between 30-70%, 20 ACH, a 12h dark/light cycle, and access to food and

water *ad libitum.*

A total of 36 adult mice aged 5-11 months were used in this experiment, consisting of 32 KI

Col1a1-K18-hACE2 mice and 4 transgenic K18-hACE2 mice. All groups were sex balanced.

The infections were performed in two separate experiments between January and June

2022. Mice were anaesthetized with isoflurane (FDG9623; Baxter, Deerfield, IL, USA) and

infected with a B.1 isolate (27 Col1a1-K18-hACE2 and 4 K18-hACE2 mice). The uninfected

control group received PBS (5 Col1a1-K18-hACE2).

Infection was performed using 1,000 $TCID_{50}$ of B.1 SARS-CoV-2 isolate in 50 µl of PBS (25

µl/nostril), or PBS only (25 µl/nostril) for the control group. All mice fully recovered from

the challenge and anaesthesia procedures. Following the challenge, body weight and clinical

signs were monitored daily. Eight animals per group were euthanized at days 3, 7, 14 dpi or

upon fulfilment of human endpoint, for viral RNA quantification and histopathological

analyses (**Supplementary Table 2**). The endpoint criteria for animal welfare were

established based on a body weight loss superior to 20% of the initial body weight and/or

the display of moderate to severe clinical signs (including neurological signs), in accordance

with previous studies (12–14). Categories evaluated included respiration, physical

appearance, lack of responsiveness and neurological changes, that were scored from 0-2

depending on severity. Euthanasia was performed under deep isoflurane anaesthesia by

whole blood extraction via cardiac puncture and was confirmed by cervical dislocation.

Oropharyngeal swab, lung, brain, and nasal turbinate were collected for viral RNA

quantification, histological and ICH analyses. For the latter techniques, tissues were fixed

by immersion in 10% buffered formalin. An additional set of 9 tissues was collected to

further characterize the KI model both for viral RNA quantification and hACE2 receptor

expression. These tissues included muscle, intestine, liver, kidney, lymph nodes, spleen,

heart, pancreas, and salivary glands.

hACE2 Receptor Tissue Expression

Several samples from both KI and transgenic models were collected and processed to

analyze potential differences in hACE2 receptor expression. These samples included

oropharyngeal swab, lung, brain, nasal turbinate, muscle, intestine, liver, kidney, pancreas,

salivary glands, lymph nodes, and spleen. Tissues were processed with a TissueLyser II

(85,300; QIAGEN) for further VL quantification, and transgene expression was analyzed by

RT-qPCR analysis.

SARS-CoV-2 PCR Detection and Viral Load Quantification

Viral RNA was quantified by RT-PCR in both the standard set of samples (oropharyngeal

swab, lung, brain, and nasal turbinate), and the extended set (muscle, intestine, liver,

kidney, pancreas, salivary glands, lymph nodes and spleen). The collected tissues were

processed, and VL determined as described by Tarrés-Freixas et al. 2022 (13). Briefly,

approximately 100 mg of each tissue was collected in 1.5 mL Sarstedt tubes (72,607;

Sarstedt, Nümbrecht, Germany) containing 500 µl of DMEM medium (11,995,065;

ThermoFisher Scientific) supplemented with 1% penicillin–streptomycin (10,378,016;

ThermoFisher Scientific, Waltham, MA, United States). A 1.5 mm Tungsten bead (69,997;

QIAGEN, Hilden, Germany) was added to each tube, and samples were homogenized twice

at 25 Hz for 30 s using a TissueLyser II (85,300; QIAGEN) before being centrifuged for 2 min

at 2,000 × g. Supernatants were then stored at −80°C until analysis.

RNA extraction was performed by using the Viral RNA/ Pathogen Nucleic Acid Isolation kit

(A42352, ThermoFisher Scientific), optimized for use with a KingFisher instrument

(5,400,610; ThermoFisher Scientific), following the manufacturer's instructions. PCR

amplification was based on the 2019-Novel Coronavirus Real-Time RT-PCR Diagnostic Panel

guidelines and protocol developed by the American Center for Disease Control and

Prevention (CDC-006-00019, v.07). Briefly, a 20 μL PCR reaction was set up containing 5 μl

of RNA, 1.5 μL of N2 primers and probe (2019-nCov CDC EUA Kit, catalogue number

10,006,770, Integrated DNA Technologies, Coralville, IA, USA) and 10 μl of GoTaq 1-Step RT-

qPCR (Promega, Madison, WI, USA). Thermal cycling was performed at 50°C for 15 min for

reverse transcription, followed by 95°C for 2 min and then 45 cycles of 95°C for 10 s, 56°C

for 15 s and 72°C for 30 s in the Applied Biosystems 7,500 or QuantStudio5 Real-Time PCR

instruments (ThermoFisher Scientific). For absolute quantification, a standard curve was

built using 1/5 serial dilutions of a SARS-CoV-2 plasmid (2019-nCoV_N_Positive Control,

catalog number 10006625, 200 copies/μL, Integrated DNA Technologies), which was run in

parallel with all PCR determinations. Viral RNA from each sample was quantified in

triplicate, and the mean viral RNA concentration (in copies/mL) was extrapolated from the

standard curve and corrected by the corresponding dilution factor. Mouse *gapdh* gene

expression was measured in duplicate for each sample using TaqMan®gene expression

assay (Mm99999915_g1; ThermoFisher Scientific) as amplification control.

Viral Titration

Lung tissues were evaluated for the presence of replicative virus by titration in Vero E6 cells

as previously described (14, 31, 32). Briefly, after tissue homogenization, each sample

underwent sequential 10-fold dilutions in duplicate, transferred onto a monolayer of Vero

E6 cell in a 96-well plate, and incubated at 37°C and 5% CO_2. Plates were monitored daily

under a microscope, and at 5 dpi, wells were evaluated for the presence of cytopathic

effects. The amount of infectious virus was calculated by determining the TCID50 using the

Reed–Muench method.

Histopathological and Immunohistochemical Analyses

Tissue samples were recovered at the designated endpoint (3,7,14 dpi or exitus) and fixed

by immersion in 10% buffered formalin. Lung, nasal turbinate, and brain samples were

routinely processed for histopathological examination, with haematoxylin & eosin-stained

slides examined under an optical microscope in a blinded fashion. A semi-quantitative

approach based on the amount of inflammation (none, mild, moderate, or severe) was used

to assess the damage caused by SARS-CoV-2 infection in mice, following a previously

published scoring system (14, 33). Additionally, an IHC technique was employed to detect

SARS-CoV-2 NP antigen in nasal turbinate, lung, and brain sections from all animals, using a

rabbit monoclonal antibody (40143-R019, Sino Biological, Beijing, China) at a 1:15,000

dilution. The amount of viral antigen in tissues was semi-quantitatively scored in a blinded

fashion (low, moderate, and high amount, or lack of antigen detection) (14, 33).

463 **Cytokine quantification**

464 To assess the viral-driven inflammation in lung in both animal models, the levels of IP-10,

465 IL-6, IFNγ, MCP-1 and MIP-1β cytokines were analyzed by Luminex in tissue extracts. Lung

466 samples were processed as stated in the VL quantification section and stored at -80ºC until

467 analysis. In Col1a1-K18-hACE2 mice, cytokines were analyzed at 3,7,14 dpi and those

468 euthanized by humane endpoint criteria (exitus). Uninfected Col1a1-K18-hACE2 and K18-

469 hACE2 infected animals were used as reference groups.

470 Cytokines were measured by Luminex xMAP technology and analyzed with xPONENT 3.1

471 software (Luminex Corporation) using the MCYTOMAG-70 kit, according to the

472 manufacturers' protocol with minor modifications. Briefly, after staining the desired

473 cytokines, samples underwent an overnight incubation on a rocking shaker at 4°C, using 2%

474 PFA to ensure complete inactivation of any remaining SARS-CoV-2 particles; a fixation that

475 does not alter cytokine quantification (34). Before plate acquisition, the PFA was washed

476 away and replaced with sheath fluid.

Statistical Analyses

All figures were generated using GraphPad Prism 9.0.0. Statistical analyses were performed

using R v4.3. Survival Rates were estimated with Kaplan-Meier curves and compared with

the Log-rank test. Datasets with an abundance of data below the limit of detection, like VL,

were analyzed using the Peto-Peto Left-censored samples test with correction for multiple

comparisons. Histopathological, IHC scores and viral titrations were compared using an

Independence Asymptotic Generalized Pearson Chi-Squared Test for ordinal data. Cytokine

titers were compared by a Kruskal Wallis, with pairwise comparisons conducted using

Conover's non-parametric test.

Acknowledgements

The authors would like to acknowledge Jorge Díaz, Yaiza Rosales-Salgado, Rosa María

Ampudia-Carrasco, Sergi Sunyé-Casas and Mireia Martínez from the CMCiB for their

essential help in the BSL3 facility and the K18-hACE2 mouse colony. We also thank Marisa

Larramona from Parc Científic de Barcelona for her unvaluable help with the knock-in

mouse colony. We thank the Dormeur Fondation for their financial support for the

acquisition of the QuantStudio-5 real-time PCR system.

Author Contributions

AP-G, FT-F, BTrinité, and JB conceived and designed the experiments. AP-G, FT-F, and

BTrinité performed the animal procedures. AP-G, FT-F, MP, ER-M, DR-R, DP-Z, JM-B, JS, BT

performed the analytical experiments. AP-G, BTrinité, JS, JV-A, VU and JB analyzed and

interpreted the data. SF, BTondelli established and provided the Knock-in mouse colony. SC

established communication between animal facilities and contributed to the veterinary

report verification to allow the knock-in animal shipment to the CMCiB facility. DP-Z, JM-B,

DR-R, NI-U provided key reagents. AP-G, BT, and JB wrote the paper. All authors contributed

to the article and approved the submitted version.

Competing Interests /Funding

A.P-G was supported by a predoctoral grant from Generalitat de Catalunya and Fons Social

Europeu (2022 FI_B 00698). This study was funded by Grifols, the Departament de Salut of

the Generalitat de Catalunya (grant nos. SLD016 to J.B. and SLD015 to J.C.), the Spanish

Health Institute Carlos III, CERCA Programme/Generalitat de Catalunya 2021 SGR 00452,

and the crowdfunding initiatives #joemcorono, BonPreu/Esclat, and Correos. NI-U is

supported by the Spanish Ministry of Science and Innovation (grant PID2020-117145RB-I00

and 10.13039/501100011033, Spain), EU HORIZON-HLTH-2021CORONA-01 (grant

101046118, European Union) and by institutional funding of Pharma Mar, Grifols, HIPRA,

and Amassence. J. Blanco has received institutional funding from Grífols, Nesapor Europe,

HIPRA and MSD. Unrelated to the submitted work, J.B. and J.C. are founders and

shareholders of AlbaJuna Therapeutics, SL. B.C. is founder and shareholder of AlbaJuna

Therapeutics, SL, and AELIX Therapeutics, SL.

The funders had no role in study design, data collection and interpretation, or the decision

to submit the work for publication

Data availability

The data supporting the findings of this study are documented within the paper and are

available from the corresponding authors upon request.

References

1. Muñoz-Fontela C, Dowling WE, Funnell SGP, Gsell PS, Riveros-Balta AX, Albrecht RA,
Andersen H, Baric RS, Carroll MW, Cavaleri M, Qin C, Crozier I, Dallmeier K, de Waal
L, de Wit E, Delang L, Dohm E, Duprex WP, Falzarano D, Finch CL, Frieman MB,
Graham BS, Gralinski LE, Guilfoyle K, Haagmans BL, Hamilton GA, Hartman AL,
Herfst S, Kaptein SJF, Klimstra WB, Knezevic I, Krause PR, Kuhn JH, Le Grand R, Lewis
MG, Liu WC, Maisonnasse P, McElroy AK, Munster V, Oreshkova N, Rasmussen AL,
Rocha-Pereira J, Rockx B, Rodríguez E, Rogers TF, Salguero FJ, Schotsaert M,
Stittelaar KJ, Thibaut HJ, Tseng C Te, Vergara-Alert J, Beer M, Brasel T, Chan JFW,
García-Sastre A, Neyts J, Perlman S, Reed DS, Richt JA, Roy CJ, Segalés J, Vasan SS,
Henao-Restrepo AM, Barouch DH. 2020. Animal models for COVID-19. Nature
586:509–515.

2. Jackson CB, Farzan M, Chen B, Choe H. 2021. Mechanisms of SARS-CoV-2 entry into
cells. Nat Rev Mol Cell Biol 2021 231 23:3–20.

3. Hassan AO, Case JB, Winkler ES, Thackray LB, Kafai NM, Bailey AL, McCune BT, Fox
JM, Chen RE, Alsoussi WB, Turner JS, Schmitz AJ, Lei T, Shrihari S, Keeler SP,
Fremont DH, Greco S, McCray PB, Perlman S, Holtzman MJ, Ellebedy AH, Diamond
MS. 2020. A SARS-CoV-2 Infection Model in Mice Demonstrates Protection by
Neutralizing Antibodies. Cell 182:744-753.e4.

4. Tseng C-TK, Huang C, Newman P, Wang N, Narayanan K, Watts DM, Makino S,
Packard MM, Zaki SR, Chan T, Peters CJ. 2007. Severe Acute Respiratory Syndrome
Coronavirus Infection of Mice Transgenic for the Human Angiotensin-Converting
Enzyme 2 Virus Receptor. J Virol 81:1162–1173.

5. Jiang R Di, Liu MQ, Chen Y, Shan C, Zhou YW, Shen XR, Li Q, Zhang L, Zhu Y, Si HR,
Wang Q, Min J, Wang X, Zhang W, Li B, Zhang HJ, Baric RS, Zhou P, Yang X Lou, Shi
ZL. 2020. Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing Human
Angiotensin-Converting Enzyme 2. Cell 182:50-58.e8.

6. Bao L, Deng W, Huang B, Gao H, Liu J, Ren L, Wei Q, Yu P, Xu Y, Qi F, Qu Y, Li F, Lv Q,
Wang W, Xue J, Gong S, Liu M, Wang G, Wang S, Song Z, Zhao L, Liu P, Zhao L, Ye F,
Wang H, Zhou W, Zhu N, Zhen W, Yu H, Zhang X, Guo L, Chen L, Wang C, Wang Y,
Wang X, Xiao Y, Sun Q, Liu H, Zhu F, Ma C, Yan L, Yang M, Han J, Xu W, Tan W, Peng
X, Jin Q, Wu G, Qin C. 2020. The pathogenicity of SARS-CoV-2 in hACE2 transgenic
mice. Nat 2020 5837818 583:830–833.

7. Zhou B, Thao TTN, Hoffmann D, Taddeo A, Ebert N, Labroussaa F, Pohlmann A, King
J, Steiner S, Kelly JN, Portmann J, Halwe NJ, Ulrich L, Trüeb BS, Fan X, Hoffmann B,
Wang L, Thomann L, Lin X, Stalder H, Pozzi B, de Brot S, Jiang N, Cui D, Hossain J,
Wilson MM, Keller MW, Stark TJ, Barnes JR, Dijkman R, Jores J, Benarafa C,
Wentworth DE, Thiel V, Beer M. 2021. SARS-CoV-2 spike D614G change enhances
replication and transmission. Nature 592:122–127.

8.   Winkler ES, Chen RE, Alam F, Yildiz S, Case JB, Uccellini MB, Holtzman MJ, Garcia-
Sastre A, Schotsaert M, Diamond MS. 2022. SARS-CoV-2 Causes Lung Infection
without Severe Disease in Human ACE2 Knock-In Mice. J Virol 96.

9.   McCray PB, Pewe L, Wohlford-Lenane C, Hickey M, Manzel L, Shi L, Netland J, Jia HP,
Halabi C, Sigmund CD, Meyerholz DK, Kirby P, Look DC, Perlman S. 2007. Lethal
Infection of K18- hACE2 Mice Infected with Severe Acute Respiratory Syndrome
Coronavirus . J Virol 81:813–821.

10.   Zheng J, Wong LYR, Li K, Verma AK, Ortiz ME, Wohlford-Lenane C, Leidinger MR,
Knudson CM, Meyerholz DK, McCray PB, Perlman S. 2021. COVID-19 treatments
and pathogenesis including anosmia in K18-hACE2 mice. Nature 589:603–607.

11.   Winkler ES, Bailey AL, Kafai NM, Nair S, McCune BT, Yu J, Fox JM, Chen RE, Earnest
JT, Keeler SP, Ritter JH, Kang LI, Dort S, Robichaud A, Head R, Holtzman MJ,
Diamond MS. 2020. SARS-CoV-2 infection of human ACE2-transgenic mice causes
severe lung inflammation and impaired function. Nat Immunol 2020 2111 21:1327–
1335.

12.   Carossino M, Kenney D, O'connell AK, Montanaro P, Tseng AE, Gertje HP, Grosz KA,
Ericsson M, Huber BR, Kurnick SA, Subramaniam S, Kirkland TA, Walker JR, Francis
KP, Klose AD, Paragas N, Bosmann M, Saeed M, Balasuriya UBR, Douam F, Crossland
NA. 2022. Fatal Neurodissemination and SARS-CoV-2 Tropism in K18-hACE2 Mice Is
Only Partially Dependent on hACE2 Expression. Viruses 2022:535.

13.   Tarrés-Freixas F, Trinité B, Pons-Grífols A, Romero-Durana M, Riveira-Muñoz E,
Ávila-Nieto C, Pérez M, Garcia-Vidal E, Perez-Zsolt D, Muñoz-Basagoiti J, Raïch-
Regué D, Izquierdo-Useros N, Andrés C, Antón A, Pumarola T, Blanco I, Noguera-
Julián M, Guallar V, Lepore R, Valencia A, Urrea V, Vergara-Alert J, Clotet B, Ballana
E, Carrillo J, Segalés J, Blanco J. 2022. Heterogeneous Infectivity and Pathogenesis of
SARS-CoV-2 Variants Beta, Delta and Omicron in Transgenic K18-hACE2 and
Wildtype Mice. Front Microbiol 13.

14.   Vidal E, López-Figueroa C, Rodon J, Pérez M, Brustolin M, Cantero G, Guallar V,
Izquierdo-Useros N, Carrillo J, Blanco J, Clotet B, Vergara-Alert J, Segalés J. 2022.
Chronological brain lesions after SARS-CoV-2 infection in hACE2-transgenic mice.
Vet Pathol 59:613–626.

15.   Shuai H, Chan JF-W, Yuen TT-T, Yoon C, Hu J-C, Wen L, Hu B, Yang D, Wang Y, Hou Y,
Huang X, Chai Y, Chan CC-S, Poon VK-M, Lu L, Zhang R-Q, Chan W-M, Ip JD, Chu AW-
H, Hu Y-F, Cai J-P, Chan K-H, Zhou J, Sridhar S, Zhang B-Z, Yuan S, Zhang AJ, Huang J-
D, To KK-W, Yuen K-Y, Chu H. 2021. Emerging SARS-CoV-2 variants expand species
tropism to murines. EBioMedicine 73:103643.

16.   Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. 2008. Severe Acute
Respiratory Syndrome Coronavirus Infection Causes Neuronal Death in the Absence
of Encephalitis in Mice Transgenic for Human ACE2. J Virol 82:7264–7275.

17. Meinhardt J, Streit S, Dittmayer C, Manitius R v., Radbruch H, Heppner FL. 2023. The
neurobiology of SARS-CoV-2 infection. Nat Rev Neurosci 2023 251 25:30–42.

18. Beard C, Hochedlinger K, Plath K, Wutz A, Jaenisch R. 2006. Efficient method to
generate single-copy transgenic mice by site-specific integration in embryonic stem
cells. Genesis 44:23–28.

19. Sun SH, Chen Q, Gu HJ, Yang G, Wang YX, Huang XY, Liu SS, Zhang NN, Li XF, Xiong R,
Guo Y, Deng YQ, Huang WJ, Liu Q, Liu QM, Shen YL, Zhou Y, Yang X, Zhao TY, Fan CF,
Zhou Y Sen, Qin CF, Wang YC. 2020. A Mouse Model of SARS-CoV-2 Infection and
Pathogenesis. Cell Host Microbe 28:124-133.e4.

20. Jiang R Di, Liu MQ, Chen Y, Shan C, Zhou YW, Shen XR, Li Q, Zhang L, Zhu Y, Si HR,
Wang Q, Min J, Wang X, Zhang W, Li B, Zhang HJ, Baric RS, Zhou P, Yang X Lou, Shi
ZL. 2020. Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing Human
Angiotensin-Converting Enzyme 2. Cell 182:50-58.e8.

21. Bradshaw CM, Georgieva T, Tankersley TN, Taylor- T, Johnson L, Uhrlaub JL,
Besselsen D, Janko Ž. 2023. Characterization of low copy number human
angiotensin- converting enzyme 2 (hACE2) -transgenic mice as an improved model
of SARS-CoV-2 infection 2.

22. Dong W, Mead H, Tian L, Park J-G, Garcia JI, Jaramillo S, Barr T, Kollath DS, Coyne
VK, Stone NE, Jones A, Zhang J, Li A, Wang L-S, Milanes-Yearsley M, Torrelles JB,
Martinez-Sobrido L, Keim PS, Barker BM, Caligiuri MA, Yu J. 2022. The K18-Human
ACE2 Transgenic Mouse Model Recapitulates Non-severe and Severe COVID-19 in
Response to an Infectious Dose of the SARS-CoV-2 Virus. J Virol 96.

23. Gembardt F, Sterner-Kock A, Imboden H, Spalteholz M, Reibitz F, Schultheiss HP,
Siems WE, Walther T. 2005. Organ-specific distribution of ACE2 mRNA and
correlating peptidase activity in rodents. Peptides 26:1270–1277.

24. Oladunni FS, Park JG, Pino PA, Gonzalez O, Akhter A, Allué-Guardia A, Olmo-
Fontánez A, Gautam S, Garcia-Vilanova A, Ye C, Chiem K, Headley C, Dwivedi V,
Parodi LM, Alfson KJ, Staples HM, Schami A, Garcia JI, Whigham A, Platt RN, Gazi M,
Martinez J, Chuba C, Earley S, Rodriguez OH, Mdaki SD, Kavelish KN, Escalona R,
Hallam CRA, Christie C, Patterson JL, Anderson TJC, Carrion R, Dick EJ, Hall-Ursone S,
Schlesinger LS, Alvarez X, Kaushal D, Giavedoni LD, Turner J, Martinez-Sobrido L,
Torrelles JB. 2020. Lethality of SARS-CoV-2 infection in K18 human angiotensin-
converting enzyme 2 transgenic mice. Nat Commun 11.

25. Yinda CK, Port JR, Bushmaker T, Owusu IO, Purushotham JN, Avanzato VA, Fischer
RJ, Schulz JE, Holbrook MG, Hebner MJ, Rosenke R, Thomas T, Marzi A, Best SM, de
Wit E, Shaia C, van Doremalen N, Munster VJ. 2021. K18-hACE2 mice develop
respiratory disease resembling severe COVID-19. PLOS Pathog 17:e1009195.

26. Fumagalli V, Ravà M, Marotta D, Di Lucia P, Laura C, Sala E, Grillo M, Bono E,
Giustini L, Perucchini C, Mainetti M, Sessa A, Garcia-Manteiga JM, Donnici L,

Manganaro L, Delbue S, Broccoli V, De Francesco R, D'Adamo P, Kuka M, Guidotti
LG, Iannacone M. 2022. Administration of aerosolized SARS-CoV-2 to K18-hACE2
mice uncouples respiratory infection from fatal neuroinvasion. Sci Immunol 7.

27. Case JB, Scheaffer SM, Darling TL, Bricker TL, Adams LJ, Harastani H, Trende R,
Sanapala S, Fremont DH, Boon ACM, Diamond MS. 2023. Characterization of the
SARS-CoV-2 BA.5.5 and BQ.1.1 Omicron Variants in Mice and Hamsters. bioRxiv
Prepr Serv Biol 2023.04.28.538747.

28. Halfmann PJ, Iida S, Iwatsuki-Horimoto K, Maemura T, Kiso M, Scheaffer SM, Darling
TL, Joshi A, Loeber S, Singh G, Foster SL, Ying B, Case JB, Chong Z, Whitener B,
Moliva J, Floyd K, Ujie M, Nakajima N, Ito M, Wright R, Uraki R, Warang P, Gagne M,
Li R, Sakai-Tagawa Y, Liu Y, Larson D, Osorio JE, Hernandez-Ortiz JP, Henry AR,
Ciuoderis K, Florek KR, Patel M, Odle A, Wong LYR, Bateman AC, Wang Z, Edara VV,
Chong Z, Franks J, Jeevan T, Fabrizio T, DeBeauchamp J, Kercher L, Seiler P,
Gonzalez-Reiche AS, Sordillo EM, Chang LA, van Bakel H, Simon V, Alburquerque B,
Alshammary H, Amoako AA, Aslam S, Banu R, Cognigni C, Espinoza-Moraga M,
Farrugia K, van de Guchte A, Khalil Z, Laporte M, Mena I, Paniz-Mondolfi AE,
Polanco J, Rooker A, Sominsky LA, Douek DC, Sullivan NJ, Thackray LB, Ueki H,
Yamayoshi S, Imai M, Perlman S, Webby RJ, Seder RA, Suthar MS, García-Sastre A,
Schotsaert M, Suzuki T, Boon ACM, Diamond MS, Kawaoka Y. 2022. SARS-CoV-2
Omicron virus causes attenuated disease in mice and hamsters. Nat 2022 6037902
603:687–692.

29. Stewart R, Yan K, Ellis SA, Bishop CR, Dumenil T, Tang B, Nguyen W, Larcher T, Parry
R, Sng JDJ, Khromykh AA, Sullivan RKP, Lor M, Meunier FA, Rawle DJ, Suhrbier A.
2023. SARS-CoV-2 omicron BA.5 and XBB variants have increased neurotropic
potential over BA.1 in K18-hACE2 mice and human brain organoids. Front Microbiol
14.

30. Perez-Zsolt D, Muñoz-Basagoiti J, Rodon J, Elousa M, Raïch-Regué D, Risco C, Sachse
M, Pino M, Gumber S, Paiardini M, Chojnacki J, Erkizia I, Muñiz X, Ballana E, Riveira-
Muñoz E, Noguera M, Paredes R, Trinité B, Tarrés-Freixas F, Blanco I, Guallar V,
Carrillo J, Blanco J, Telenti A, Heyn H, Segalés J, Clotet B, Martinez-Picado J, Vergara-
Alert J, Izquierdo-Useros N. 2021. Siglec-1 on dendritic cells mediates SARS-CoV-2
trans-infection of target cells while on macrophages triggers proinflammatory
responses. bioRxiv 8:2021.05.11.443572.

31. Rodon J, Muñoz-Basagoiti J, Perez-Zsolt D, Noguera-Julian M, Paredes R, Mateu L,
Quiñones C, Perez C, Erkizia I, Blanco I, Valencia A, Guallar V, Carrillo J, Blanco J,
Segalés J, Clotet B, Vergara-Alert J, Izquierdo-Useros N. 2021. Identification of
Plitidepsin as Potent Inhibitor of SARS-CoV-2-Induced Cytopathic Effect After a Drug
Repurposing Screen. Front Pharmacol 12:646676.

32. Haagmans BL, van den Brand JMA, Raj VS, Volz A, Wohlsein P, Smits SL, Schipper D,
Bestebroer TM, Okba N, Fux R, Bensaid A, Solanes Foz D, Kuiken T, Baumgärtner W,
Segalés J, Sutter G, Osterhaus ADME. 2016. An orthopoxvirus-based vaccine

reduces virus excretion after MERS-CoV infection in dromedary camels. Science (80-
) 351:77–81.

33. Brustolin M, Rodon J, Rodríguez de la Concepción ML, Ávila-Nieto C, Cantero G,
Pérez M, Te N, Noguera-Julián M, Guallar V, Valencia A, Roca N, Izquierdo-Useros N,
Blanco J, Clotet B, Bensaid A, Carrillo J, Vergara-Alert J, Segalés J. 2021. Protection
against reinfection with D614- or G614-SARS-CoV-2 isolates in golden Syrian
hamster. Emerg Microbes Infect 10:797.

34. Dowall SD, Graham VA, Tipton TRW, Hewson R. 2009. Multiplex cytokine profiling
with highly pathogenic material: use of formalin solution in luminex analysis. J
Immunol Methods 348:30–35.

Figure Legends

Figure 1. Expression of hACE2 in the Col1a1-K18-hACE2 KI model. (A) Schematic

representation of insertion strategy. The original K18-hACE2 transgene was inserted into

the collagen COL1A1 locus using a recombinase mediated cassette exchange (RMCE) FLP-

FRT system in KH2 cells via blastocyst injection. pPGK-ATG-frt plasmid: vector backbone

with Ampicillin resistant gene (Amp), transcription start site (ATG) and Flippase recognition

target (Frt). Hygro: Hygromicin resistance gene. pCAGGS-FlpE: expression plasmid for FLPe

recombinase expression. (B) Relative quantification of hACE2 receptor expression to GAPDH

expression in uninfected Col1a1-K18-hACE2 ((blue empty dot, n=3) and K18-hACE2 males

(black empty square, n=3). Delta Ct values are inversely shown to facilitate interpretation.

The lower the absolute number, the higher the relative expression. Solid line and bars

represent mean and SEM. (C) Relative comparison of hACE2 receptor expression in Col1a1-

K18-hACE2 versus K18-hACE2 mice (n=3 each, males). Higher receptor expression in Col1a1-

K18-hACE2 (negative values in Y axis) is marked by blue bars, and higher expression in K18-

hACE2 (positive values in Y axis) is marked in black bars.

Figure 2. Experimental setting and progression of SARS-CoV-2 infection in Col1a1-K18-

hACE2 and K18-hACE2 mouse models. (A) Schematic representation of the experimental

setting. Knock-in Col1a1-K18-hACE2 (n=27) and transgenic K18-hACE2 mice (n=4) were

intranasally challenged with a 1000 $TCID_{50}$ dose of a B.1 SARS-CoV-2 isolate. A Col1a1-K18-

hACE2 uninfected control group (n=5) was challenged with PBS. Mice were monitored for

weight-loss and clinical signs for 14 days post infection (dpi). Euthanasia was performed at

3, 7, and 14 dpi or upon fulfilment of humane endpoint criteria, for sample and tissue

collection (n=8 per timepoint). Infections were performed in two separate experiments

between January and June 2022. Created with Biorender.com. (B) Relative body weight

follow-up referred to day 0. Col1a1-K18-hACE2 uninfected (blue empty dot), Col1a1-K18-

hACE2 infected (blue dot), K18-hACE2 infected (black square). Solid lines and bars represent

mean±SD. (C) Survival (Kaplan-Meier). All K18-hACE2 infected animals (n=4) had to be

euthanized due to endpoint criteria by 7dpi, and only three infected Col1a1-K18-hACE2 at

8, 9, and 10 dpi. No uninfected Col1a1-K18-hACE2 had to be euthanized under this criterion.

Col1a1-K18-hACE2 uninfected (blue dashed line), Col1a1-K18-hACE2 infected (blue line),

K18-hACE2 infected (black line). Statistical differences were identified using a Log-rank

(Mantel-Cox) test (<0.0001), followed by individual comparisons (**$p < 0.005$; ****$p <$

0.0001)

Figure 3. Progression of SARS-CoV-2 infection in Col1a1-K18-hACE2 KI mice. Col1a1-K18-

hACE2 (blue circles, *n=27*) and K18-hACE2+ mice (black squares, *n=4*) were inoculated with

1000 $TCID_{50}$ of a SARS-CoV-2 B.1 isolate (full shapes) or uninfected (empty shapes of each

colour, *n=5*). Animals were euthanized at 3dpi (n=8), 7dpi (n=8), 14 dpi (n=8) or upon

fulfilment of humane endpoint criteria. (A) SARS-CoV-2 viral RNA loads (copies/mL) of

oropharyngeal swab, lung, and nasal turbinate samples. Dashed line represents limit of

detection, established by 2SD of uninfected animals. Statistical differences were identified

using a Peto-Peto Left censored test (*$p < 0.05$, **$p<0.005$). (B) Viral titration of replicative

virus ($TCID_{50}$/mL) in lung samples from B.1 infected mice at different endpoints in Vero E6

cells on day 5 of culture. Titers were compared using an Independence Asymptotic

Generalized Pearson Chi-Squared Test for ordinal data (*$p < 0.05$, **$p<0.005$). (C) Lung

histopathological scoring of broncho-interstitial pneumonia (left) and SARS-CoV-2 NP

immunohistochemical (IHC) scoring (right) in both models at 3, 7, 14 dpi and endpoint.

Statistical differences were identified using an Independence Asymptotic Generalized

Pearson Chi-Squared Test for ordinal data (*p < 0.05). (D) Representative lung histology and

IHC pictures of both models at 3, 7, 14 dpi and endpoint. Images show low-power

magnification bars (200 μm).

Figure 4. Inflammatory response in the lung. The concentration of Inflammatory cytokines

in lung extracts in the knock-in and transgenic models at 3,7,14 dpi or endpoint are shown.

Col1a1-K18-hACE2 infected (blue full circles, *n=23*), uninfected (blue empty circles, n=3) and

K18-hACE2+ infected mice (black full squares, *n=4*). The bar shows Median with

interquartile range. Limit of detection for each cytokine is indicated by a dotted line.

Statistical differences were identified using a Kruskal Wallis, and a Conover's nonparametric

all-pairs comparison test (*p < 0.05, **p<0.005).

Figure 5. Progression of SARS-CoV-2 infection in brain from the Col1a1-K18-hACE2 KI mice.

Col1a1-K18-hACE2 (blue circles, *n=27*) and K18-hACE2+ mice (black squares, *n=4*) were

inoculated with 1000 $TCID_{50}$ of a B.1 SARS-CoV-2 isolate (full shapes) or uninfected (empty

shapes, *n=5*) and followed until 14 days post infection (dpi). Samples were collected at 3, 7,

14 dpi and endpoint. (A) SARS-CoV-2 viral RNA loads (copies/mL) in brain extracts. Dashed

line represents limit of detection, established by 2SD of uninfected animals. Statistical

differences were identified using a Peto-Peto Left-censored samples test with correction for

multiple comparisons. (B) Brain histopathological scoring of multifocal lymphoplasmacytic

meningo-encephalitis in brain (Left) and SARS-CoV-2 NP IHC scoring (right) in brain of both

models. Statistical differences were identified using an Independence Asymptotic

Generalized Pearson Chi-Squared Test for ordinal data (*p < 0.05; **p < 0.01). (C)

Representative images of brain Histology and IHC images of both hACE2+ mouse models.

Images show low-power magnification (top; bars: 600μm) and medium-power

magnification (bottom; bars: 300μm). Col1a1-K18-hACE2 KI mice samples shown were

collected at 7 dpi, while K18-hACE2 samples were collected at humane endpoint (6-7dpi).

Figure 6. Spider Plot of clinical data compilation at 7 dpi from both animal models.

Comparative data summary of Col1a1-K18-hACE2 and K18-hACE2 mice infection by a B.1

SARS-CoV-2 isolate. Data at 7dpi of the KI (blue; euthanized at 7 dpi) and K18-hACE2 (black;

euthanized at 6-7dpi) of different parameters was compiled and median values are shown

in each axis of the spider plot. Data ranges are adjusted for each kind of data, and shown by

concentric circles, being the outside circle the highest level. Clinical status was assessed in

different categories: respiration, appearance, weight-loss, lack of responsiveness and

neurological signs. Histology and IHQ data were semiquantitatively scored from 0-2, 0-3 and

0-3 respectively as described in Materials and Methods section. Viral loads are shown as

Log10 copies/mL, titration as Log10 TCID50 and cytokines as pg/mL.

A

B

hACE2 Receptor Expression

C

Relative hACE2 Receptor Expression

Fig 1.

Fig 2.

Fig 3.

- • Col1a1-K18-hACE2 Infected
- ○ Col1a1-K18-hACE2 Uninfected
- ■ K18-hACE2 Infected

Fig 4.

Fig 5.

Clinical Status

Neurological
Lack.of.Responsiveness
Weight.loss
Appearance
Respiratory

Brain

Brain.Histology
Brain.IHC
Viral.Load.Brain

Lung
Cytokines

MIP1b
MCP1
IP10
IL6
IFNg

Respiratory System

Viral.Load.Swab
Viral.Load.NT
Viral.Load.Lung
Titration
Lung.IHC
Lung.Histology

○ K18-hACE2
○ Col1a1-K18-hACE2

Fig 6.