The interplay between Mobilome and Resistome in *Staphylococcus aureus*

Rachel Contarin[1-2], Antoine Drapeau[2], Pauline François[2], Jean-Yves Madec[2], Marisa Haenni[2*], Emilie Dordet-Frisoni[1*]

[1]INTHERES, Université de Toulouse, INRAE, ENVT, Toulouse, France
[2]Anses – Université de Lyon, Unité Antibiorésistance et Virulence Bactériennes, Lyon, France

* These authors contributed equally to the work

Corresponding author:

Emilie Dordet-Frisoni, emilie.dordet-frisoni@inrae.fr

Marisa Haenni, marisa.haenni@anses.fr

Abstract

Antibiotic resistance genes (ARGs) in *Staphylococcus aureus* can disseminate vertically through successful clones, but also horizontally through the transfer of genes conveyed by mobile genetic elements (MGEs). The underexplored MGE/ARG associations in S*. aureus* favor the emergence of multidrug-resistant clones, posing a significant threat to human and animal health.

This study investigated the interplay between the mobilome, encompassing MGEs, and the resistome, the collection of ARGs, in more than 10,000 *S. aureus* genomes from human and animal sources. The analysis revealed a remarkable diversity of MGEs and ARGs, with plasmids and transposons being the main carriers of resistance genes. Numerous MGE/ARG associations were identified, suggesting that MGEs play a critical role in the dissemination of resistance. A high degree of similarity was observed in MGE/ARG associations between human and animal isolates, highlighting the potential for unrestricted spread of ARGs between hosts. While clonal expansion is a major driver of resistance dissemination in *S. aureus*, our results showed that MGEs and their associated ARGs can spread across different strain types (STs), favoring the evolution of these clones and their adaptation in selective environments. The high variability of MGE/ARG associations within individual STs and their spread across several STs highlight the crucial role of MGEs in shaping the *S. aureus* resistome.

Overall, this study provides valuable insights into the complex interplay between MGEs and ARGs in *S. aureus*, emphasizing the need to elucidate the mechanisms governing the epidemic success of MGEs, particularly those implicated in ARG transfer.

Importance

The research presented in this article highlights the crucial importance of understanding the interactions between mobile genetic elements (MGEs) and antibiotic resistance genes (ARGs) carried by *Staphylococcus aureus*, a versatile bacterium that can be both a harmless commensal and a dangerous pathogen for humans and animals. *S. aureus* strains represent a major threat due to their ability to rapidly acquire and disseminate ARGs. By analyzing a large dataset of *S. aureus* genomes, we

40   highlighted the substantial role of MGEs, in particular plasmids and transposons, in the dissemination

41   of ARGs within and between *S. aureus* populations, bypassing the host barrier. Given that multidrug-

42   resistant *S. aureus* strains are classified as a high-priority pathogen by global health organizations, this

43   knowledge is crucial for understanding the complex dynamics of transmission of antibiotic resistance

44   in this species.

Introduction

Staphylococcus aureus is a versatile bacterium that can be a commensal microorganism and a lethal pathogen for both humans and animals. The large range of symptoms caused by *S. aureus* in each host is the result of the intricate interplay between complex factors, including host health status, genetic composition of the *S. aureus* strain and the site of infection (1–3). The overuse and misuse of antibiotics in both human healthcare and veterinary medicine have promoted the emergence of numerous antibiotic-resistance genes (ARGs) and the dissemination of multidrug-resistant clones (4). As a result, this species has been classified by the WHO (World Health Organization) as high-priority for the development of new antibiotic targets, and by the WOAH (World Organization for Animal Health) on the surveillance list of multi-resistant pathogens. Studies on antimicrobial resistance (AMR) in *S. aureus* have primarily focused on methicillin-resistance, due to the presence of the *mecA/mecC* genes, which confer resistance to all beta-lactam antibiotics. Consequently, *S. aureus* strains are commonly classified as methicillin-resistant (MRSA) or methicillin-susceptible (MSSA) isolates (5). However, *S. aureus* can acquire multiple additional ARGs conferring resistance to all known antibiotic families, shaping the resistome of each isolate (6). In 2018, a review drew up an inventory of 46 ARGs shared between *S. aureus* isolates of human and animal sources, suggesting potential interspecies transmissions (7).

In contrast to a highly clonal core genome, up to 25% of the *S. aureus* genome is composed of accessory genes, primarily encompassing mobile genetic elements (MGEs) that contribute significantly to the adaptability and survival of these bacteria in various environmental conditions by spreading ARGs mainly occurs through horizontal gene transfer (HGT) (8, 9). MGEs are DNA fragments that can promote intra- or intercellular transfer of genetic material, and play a crucial role in the dissemination of ARGs. The staphylococcal mobilome (set of MGEs) exhibits a remarkable variability (8, 10–13). *S. aureus* MGEs include (i) plasmids (extrachromosomal genetic elements that can be conjugative, mobilizable, or non-mobilizable in *S. aureus*) (12), (ii) integrative and conjugative elements (ICEs) (MGEs that integrate into the bacterial chromosome and have the potential to excise and transfer

themselves to other bacteria via conjugation) (14, 15), (iii) insertion sequences (IS) and transposons

(Tn) (chromosomally-integrated MGEs that are often associated with other MGEs, in single or multiple

copies) (13), (iv) prophages (MGEs usually integrated into the host's chromosome, since they can

repress the phage's lytic functions) (16), and (v) genomic islands (such as the staphylococcal

chromosomal cassettes (SCC*mec*) specific to *mec*-carrying staphylococci) (13).

It is usually admitted that *S. aureus* disseminate through clonal expansion, spreading resistance genes

vertically through waves of successful lineages. While the role of MGEs, particularly plasmids, has been

extensively studied in Enterobacterales, few studies have addressed the diverse MGEs and associated

ARGs carried by *S. aureus* (8, 9, 13, 17). Likewise, no large-scale analyses have been conducted to

localize ARGs across *S. aureus* genomes from diverse sources and hosts. To fill in this knowledge gap,

this study aimed at (i) identifying the most prevalent MGEs and ARGs in 10,063 *S. aureus* genomes

from the NCBI public database, (ii) highlighting MGE/ARG associations and (iii) evaluating the role of

MGEs in the emergence and spread of antibiotic resistance in *S. aureus*.

Results

Diversity of human and animal *S. aureus* genomes

A total of 9,408 human and 655 animal *S. aureus* genomes were analyzed. Animal *S. aureus* genomes

came from 15 different hosts, mainly livestock (n=527, see porcine, bovine, avian, ovine and caprine

origins in Table S1). Data originated from all continents, with a significant proportion of *S. aureus* of

human source (Hm-SA) isolated from North America and Europe, and of *S. aureus* of animal source

(An-SA) isolated from Europe and Asia (Figure 1, Table S1). The core genome of all *S. aureus* strains

consisted of 1,716 genes, including 86 genes unique to An-SA and 43 genes unique to Hm-SA. A total

of 433 distinct sequence types (STs) were identified, but only 21 were found in >1% of either Hm-SA

and/or An-SA were identified. Among the 433 STs, 362 (three present in > 1% of Hm-SA) were unique

to Hm-SA, 34 (two present in > 1% of An-SA) to An-SA*,* and 37 (16 present in >1% of Hm-SA and An-SA)

were shared between both groups, highlighting the source-specificity of STs (Figure S1A).

The most frequently identified STs were (i) the two major livestock associated ST398 (n=216, mainly in

Europe) and pig-associated ST9 (n=125, mainly in Asia) (Figures 1A & S1B) in An-SA and (ii) ST5

(n=2,138) and ST8 (n=1,719) in Hm-SA (Figures 1B and S1B).

Mobile genetic elements arsenal in *S. aureus* genomes

Analysis of accessory genomes highlighted a remarkable diversity of MGEs including ICEs, transposons

and plasmids regardless of their human or animal sources (Figure 1, Table 1). A total of 114,845 MGEs,

or 59,768, if excluding IS, were detected (Tables 1 & S1), with IS elements, prophages, plasmids and

SCC*mec* being the most prevalent, followed by transposons and ICEs (Table 1). Composite transposons

(CTn) were the least common MGE detected in *S. aureus* genomes. Transposons and ICEs were found

in *S. aureus* genomes irrespective of their origin. In contrast, IS elements and plasmids exhibited

greater host specificity, since only 33% and 23% were respectively shared between Hm-SA and An-SA

strains (Table 1).

The MGE content varied significantly depending on STs. Plasmids, IS and SCC were present in all major

STs (found in >1% of isolates), regardless of their sources. However, specific MGEs could be absent

from certain STs, such as transposons that were not identified in ST22 genomes (Figure S2A & B). The

mobilome was also diverse among isolates of a same ST, particularly in their plasmid and transposon

content, as observed for An-SA ST398 and Hm-SA ST8 (Figure 1).

A wide diversity was also observed for each MGE (Figure 2). Among plasmids, Rep1 was predominant

in both An-SA (43%) and Hm-SA (29%) genomes, while others segregated depending on the source,

like RepA_N and RepL in Hm-SA (16%) versus Rep_trans in An-SA (24%) (Figure 2). Plasmids also

frequently harbored multiple *rep* genes (37% of all Hm-SA plasmids possessing multiple *rep* genes and

23% in An-SA). For transposons, the Tn*554* family was prevalent (30%) in Hm-SA, the Tn*558* family was

widespread (27%) in An-SA and the Tn*552* was equally distributed (about 20% in Hm-SA and An-SA)

(Figure 2). Concerning ICEs, the ICE*6013* family prevailed in Hm-SA (44% of genomes versus 26% in An-

SA), whereas the Tn*916* family exhibited a higher prevalence in An-SA (38%, and 12% for Hm-SA)

(Figure 2). For SCC*mec*, type II(2A) and type IVa(2B) were frequent in Hm-SA (27% and 19%,

respectively), while type XII(9C2) was the most prevalent (21.5%) in An-SA (Figure 3A). Among

prophages, the most common were phi2958PVL and P282 families, present in more than 33% of Hm-

SA genomes, while the prophage StauST398 was the most abundant in An-SA (28%) (Figure 3B). For IS,

the most common families were IS*Sau6*, IS*Sau3* and IS*1272*, regardless of the source (Figure 3C). IS*Sau4*

was very frequent in Hm-SA (41%) and almost absent in An-SA (0.9%) (Figure 3C). Contrarily, IS*256* was

widespread in An-SA (36%) but rarely found in Hm-SA (9%) (Figure 3C). Finally, as observed for ISs, the

CTn containing IS*Sau6* was the most prevalent, identified in over 23% of *S. aureus* genomes, regardless

of their source (Figure 3D). The CTn with IS*Sau4* was abundant in Hm-SA genomes (11.5%) but rare in

An-SA genomes (0.5%).

S. aureus antibiotic resistance genes content

Seventy-eight different ARGs were identified, of which 50 were shared by both Hm-SA and An-SA

genomes, 27 were unique to Hm-SA, and one (*optrA*) was unique to An-SA (Figure 4, Table S1).

S. aureus genomes harbored an average of six or five different ARGs per An-SA or Hm-SA genomes,

respectively, with a maximum of 17 per Hm-SA and 15 per An-SA (Table S1). In An-SA isolates, the

lnu(B) gene was consistently found in tandem with the *lsa*(E) gene (>99%) (Figure S3). Similarly, in Hm-

SA some ARGs were frequently collocated (>90%) as *ant(9)-Ia* and *erm(A)*, *ant(6)-Ia* and *aph(3')-III*, and

mph(C) and *msr(A)* (Figure S3).

The *blaZ* and *mecA* genes, conferring resistance to beta-lactam antibiotics, were commonly found

regardless of the source and often co-harbored (n=417 in An-SA and n=5,913 in Hm-SA) (Figures 4 &

S3). The *aac(6')-aph(2'')* (aminoglycoside), *tet(M)* and *tet(L)* (tetracyclines), *dfrG* (trimethoprim)*, lnu(B)*

and lsa(E) (lincosamides) and *fexA* (phenicol) genes were more prevalent (present in >25% of genomes)

in An-SA, while the *ant(9)-Ia* and *aph(3')-III* (aminoglycosides) and *erm(A)* (macrolides) genes were

more common in Hm-SA (Figure 4). Of note, the *aac(6')-aph(2'')* gene was found in 83% of the equine

S. aureus, while it was present in less than 60% of the genomes of other animal hosts (Figure 4). Among

animals, pig isolates presented the largest number of resistance genes, with 11 genes belonging to

seven different families present in more than 40% of the genomes (Figure 4). In contrast, in bovine,

only *blaZ* and *mecA* were present in more than 40% of the genomes (Figure 4). Certain ARGs exhibited

restricted distribution patterns, as *vga(E)* present only in human-derived ST398 *S. aureus* strains or

vga(A)LC and *apmA* identified in animal-derived ST398/ST9 strains. The ARG/ST associations were

source-specific, except for *str*, *spd*, *ampA*, *erm(T)*, *dfrK* and *vga(E)*, which were predominantly (>80%)

identified in ST398 genomes in both Hm-SA and An-SA (Figures S2C & D).

Association of mobile genetic elements with antibiotic resistance genes in *S. aureus*

A Pearson correlation analysis revealed a strong positive correlation between the number of ARGs and

the number of MGEs (excluding IS) in *S. aureus* genomes (r=0.57, p<0.0001, r=0.39, p<0.0001, for Hm-

SA and An-SA, respectively). This correlation was particularly high for transposons, with R-values of

0.45 (p<0.0001) for Hm-SA and 0.60 (p<0.0001) for An-SA strains, but also for SCC elements (r=0.45,

p<0.0001; r=0.48, p<0.0001) and plasmids (r=0.24, p<0.0001; r=0.46, p<0.0001).

The majority of ARGs (78% in Hm-SA, 75% in An-SA) were associated with MGEs (Figure5, Table S2).

Plasmids (47% in An-SA and 42% in Hm-SA) were the predominant MGE harboring ARGs, followed by

SCC elements (22% in Hm-SA and 11% in An-SA) and transposons (18% in Hm-SA and 12% in An-SA)

(Figure 5, Table S2). The diversity of MGE/ARG co-occurrences was high since 21 and 12 different

combinations of MGE/ARG were identified in Hm-SA and An-SA, respectively (Figure 5). The *spd*,

cat(pC221), *qacG*, *str*, *tet(T)* and *vga(A)LC* genes were exclusively found on plasmids, while *ant(9)-Ia*,

erm(A), *fexA*, *vga(A)V* and *vga(E)* were primarily associated with transposons, regardless of the

S. aureus source (Figure 5). Several genes were associated with various types of MGEs, like *erm(B)* and

dfrK which were frequently carried by plasmids or transposons and by putative plasmid/transposon

tandem (Figure 5). *S. aureus* MGEs could be nested within each other, like plasmids or ICEs harboring

transposons, or SCC*mec* elements hosting a diverse array of other MGEs (Figure 5).

Plasmid/ARG associations. Each ARG was primarily associated with a specific Rep family (Figure 6).

Rep1 and RepA_N were the dominant plasmid families associated with ARGs, either alone or in

combination with other Rep (80%, 496/619 in Hm-SA; 78%, 164/210 in An-SA) (Figure 6). Within these,

Rep1/*aadD* (8%, n=1,603/20,299) in Hm-SA and Rep1/*tet(L)* (6.5%, n=133/2,038) in An-SA, emerged as

the most common. An-SA displayed two prevalent plasmid/multi-ARGs co-occurrences (Figure 7A). The

first involved seven ARGs *aac(6')-aph(2'')*, *aadD*, *ant(6)-Ia*, *erm(C)*, *lnu(B)*, *lsa(E)* and *tet(L)* carried by a

Rep1 plasmid (n=42). The second involved the *aadD*, *tet(L)* and *tet(T)* genes carried by an Inc18/Rep1

plasmid (n=13). In Hm-SA, the two most widespread associations were Inc18/Rep1/RepA_N (n=401)

and Inc18/RepA_N (n=363), each associated with the same six ARGs *ant(6)-Ia*, *aph(3')-III*, *blaZ, mph(C),*

msr(A) and sat-4 (Figure 7B).

Transposon/ARG associations. Each ARG was associated with one type of transposon and identical

transposon/ARG associations were identified in both human and animal genomes (Figure 6). The

Tn*554/ant(9)-Ia*, Tn*554/erm(A)* and Tn*558/fexA* associations were the most frequently detected in

Hm-SA and An-SA, respectively (Figure 6). Apart from *erm(A)* and *ant(9)-Ia* found on Tn*554* and Tn*6133*

or *erm(B)* associated with Tn*551* and Tn*917* (Figure 6), ARGs were found only on one type of

transposon. Some ARGs, such as *blaZ*, were also found associated with both transposons and ICEs

(Table S2). A strong co-occurrence between certain genes and transposons was observed, as for *lnu(G)*,

vga(A)V, *vga(E)* or *fexA* genes that were predominantly associated with transposons (Figure 6).

Transposon also harbored multiple ARGs. Notably, the Tn*554/erm(A)/ant(9)-Ia* association emerged

as the most frequent MGE/multi-ARGs association in Hm-SA (n=3,643) and the second most common

in An-SA (n=25) (Figure 7). Additionally, the Tn*6133/erm(A)/ant(9)-Ia/vga(E)* co-occurrence was

identified in both Hm-SA (n=25) and An-SA (n=11) genomes (Figure 7).

Integrative and conjugative element/ARG associations. Overall, a minority (<9% of MGE/ARG

associations) of ARGs were associated with ICEs, but associations were diverse (Table S2). The most

widespread element was Tn*916/tet(M)* (Figure 6). This gene was itself usually located on ICEs

(96%/100% in Hm-SA/An-SA) (Figure 6) and sometimes associated with other genes like *dfrG* (Figure

7).

Staphylococcal chromosomal cassette/ARG associations. The *mecA* gene was carried by several

types of SCC*mec* cassettes, while *mecC* was exclusively associated with the type XI(8E) element. In

addition to *mec* genes, 23 different ARGs were found on SCC*mec* elements and formed 97 SCC*mec*/ARG

associations in Hm-SA genomes, while nine different ARGs were found forming 12 different

SCC*mec*/ARG associations in An-SA strains (Table S2). In Hm-SA, the most common genes were *ant(9)-*

Ia and *erm(A)* genes carried by a type II(2A) SCC*mec* element (1,513 for *ant(9)-Ia* and 1,473 for *erm(A)*)

and in An-SA genomes *blaZ* was always associated with type XI(8E) cassette carrying *mecC* (n=10)

(Table S2). The SCC*mec*/multi-ARGs associations, like type Vc/*mecA/tet(K)* and type IVa/*mecA/dfrC*,

were found in both humans (n=27 and and animals even though their proportions differed depending

on the source (n=27 and 41 in Hm-SA, n=8 and 3 in An-SA) (Figure 7, Table S2).

Prophage/ARG associations. ARGs were rarely associated with prophages (0.3% in Hm-SA, 0.1% An-

SA genomes). In Hm-SA genomes, the most common prophage/ARG associations were

phi2958PVL/*dfrG* (n=27) and SPbeta_like/*aac(6')-aph(2'')* (n=23) (Table S2). This prophage was found

associated with 17 different ARGs (Table S2). Prophage/ARG occurrences were not exclusive, i.e. *blaZ*

could be associated with 10 different types of prophages. In An-SA genomes, only two associations

were identified, i.e. *aac(6')-aph(2'')*, *aph(3')-III* and *ant(6)-Ia* carried by a SPbeta-like prophage, and

dfrG associated with YMC/09/04/R1988.

Composite transposon/ARG associations. These MGEs were only associated with about 5% of ARGs

in both Hm-SA and An-SA (Table S2). IS*Sau6* was the most common CTn associated with several ARGs

(24 different ARGs in Hm-SA and 15 in An-SA genomes) (Table S2). The most frequently identified co-

occurrences were IS*Sau6*/*mecA*, harbored by a SCC*mec* element, in An-SA strains (n=103), but also

abundant in Hm-SA (n=534), even if IS*Sau6*/*aadD*, also located on SCC*mec*, was most widespread in

Hm-SA (n=547).

Discussion

Although MGEs are recognized key players in the dissemination of AMR, the association between

MGEs and ARGs has been understudied in Gram-positive bacteria, particularly in *S. aureus*. While

previous studies focused either on MGEs and/or ARGs (8, 9, 18, 19), our large-scale analysis of over

10,000 publicly available *S. aureus* genomes from both animal and human sources is, to our

knowledge, the first to reveal the intricate interplay between mobilome and resistome. Our analysis

highlighted a remarkable diversity of STs (n=433), MGEs (n=278) and ARGs (n=78) in *S. aureus,* as well

as the routes of their dissemination within the species. Our study, like all studies based on publicly

available genomes, has some limitations imposed by the bias in sequenced genomes deposited in the

NCBI database. Due to sequencing costs, only strains harboring the most critical resistance genes for

human and animal health were analyzed. This resulted in an over-representation of MRSA strains (80%

Hm-SA and 71% An-SA) and the under-representation of An-SA, except for *mecA*-positive ST398 and

ST9 isolates of porcine origin (n=143/655 An-SA, 22%) and MRSA of bovine-origin (n=81/655 An-SA,

12%). Additionally, our analyses included both complete and draft genomes, which may have resulted

in the exclusion of elements spanning several contigs. However, since our study aimed to capture the

diversity of MGEs and ARGs rather than quantify them, this bias did not significantly interfere with our

analyses.

Our study showed a wide distribution and a large diversity of ARGs in *S. aureus* of human and animal

origin, even larger than the one reported in the review by Schwartz *et al.* (7), and capable of conferring

resistance to all antibiotic families known to treat staphylococcal infections. The most widespread

genes were *blaZ* and *mecA*, as also pointed out in a recent study of AMR in publicly available *S. aureus*

genomes (20). A recent study on the resistome/mobilome association in ESKAPE pathogens highlighted

the overall high prevalence of genes conferring resistance to aminoglycosides, chloramphenicol,

trimethoprim and tetracyclines (21). Specific genes conferring resistance to these antibiotic families

were also particularly frequent in our study dedicated to *S. aureus*, and their wide dissemination is

consistent with antibiotic use in both human and veterinary medicine. Globally, our study revealed a

higher diversity of ARGs in Hm-SA and a larger set of human-specific genes. This could be explained by

the larger number of genomes analyzed, but also by the higher exposure of humans to antibiotic

selection in various environments (hospitals, farm environments or food). Only one gene, *optrA*, was

specific to An-SA. This gene, first discovered in human and animal enterococci on a conjugative plasmid

(22), and then observed in *Staphylococcus sciuri* of porcine origin (23), is conferring resistance to

phenicols. It was found on *S. aureus* chromosome near the SCC*mec*, but also on a plasmid with the *cfr*

gene (23). In our study, *optrA* was identified on the chromosome and not associated with an MGE:

nevertheless, it was located <10kb from the *fexA* gene carried by Tn*558*, an MGE/ARG association

already found on the plasmid carrying *cfr* in other staphylococci (24, 25). Besides these host

specificities, a large set of genes, among which *aph(3')-III*, *dfrC*, *fusA*, *tet(T)*, *apmA* and *vga(E)*, were

shared between An-SA and Hm-SA. This suggests that numerous ARGs have no host boundaries and

can be disseminated from human *S. aureus* to animals and vice versa.

The *S. aureus* mobilome was largely composed of IS and phages, which rarely carried ARGs, followed

by plasmids and transposons that were identified as the main carriers of ARGs for this species. Our

study also highlighted that plasmids exhibited the highest diversity among MGE, with 297 different *rep*

genes combinations often forming mosaic plasmids with multiple *rep* genes (37% in Hm-SA and 23% in

An-SA) (12, 26). Previous studies mainly reported the association of only two *rep* genes, belonging to

the same or two different plasmid families (27, 28). However, our work demonstrates that as many as

eight different *rep* genes can be found in a single plasmid. Unlike plasmids circulating in

Enterobacterales, those identified in *S. aureus* present few Rep proteins incompatibilities. This

facilitates the formation of mosaic plasmids, enabling *S. aureus* to adapt more effectively to diverse

hosts and environmental conditions (27, 29–31). Furthermore, the presence of numerous insertion

sequences (IS elements) and the rolling-circle replication (RCR) system of small plasmids (<10 kb) favors

the recombination and fusion of several small plasmids, resulting in unique large hybrids carrying

multiple *rep* genes (27, 32).

The prevailing view for a long time held that *S. aureus* possessed very few plasmids, and horizontal

gene transfer was believed to occur primarily through transduction. However, studies on Gram-

positive bacteria showed that plasmids can be mobilized by using the conjugation system of other

plasmids or ICEs present in the strain (33–35). A study in *Bacillus subtilis* showed that plasmids can be

mobilized by the *ori*T of ICE*Bs1,* an ICE of the Tn*916* family (34, 35). In *S. aureus*, plasmids can be

mobilized if they possess a relaxase or by forming co-integrates that are subsequently resolved by

"conduction" (36, 37). O'Brien *et al.* (38) showed that the conjugative plasmid pWBG749 could

mobilize the transfer of numerous small plasmids carrying an *ori*T similar to its own (38, 39). Thus,

plasmids carrying ARGs, even non-conjugative ones, could be easily transferred between cells,

highlighting their central role in the dissemination of these genes (15).

Often misclassified in the literature as transposons or plasmids (40), ICEs are also capable of HGT.

Studies on enterococci and streptococci showed that these MGEs play a crucial role in the

dissemination of ARGs (41). The well-known Tn*916* family was found in many Firmicutes, always

carrying the *tet(M)* gene (42–44). This element exhibits a similar backbone sequence between

S. aureus and streptococci (43). Within *S. aureus*, it was first described in the Mu50 strain and named

Tn*5801* (45). Our study identified another ICE, ICE*6013*, as the most prevalent family typically

harboring few resistance genes itself. However, it can carry other MGEs like Tn*552*, often associated

with the *blaZ gene*. This association of multiple MGEs with ARG was first identified in *S. aureus* ST239

(14). Despite sharing some protein similarities with the ICE*Bs1* of *Bacillus subtilis*, ICE*6013* belongs to

a distinct family (14). Other MGEs were also found nested within each other, like transposons found

within plasmids and SCC*mec* cassettes. For instance, Tn*554/ant(9)-Ia/erm(A)* was found on both Rep1

plasmids and SCC*mec* type II(2A) cassettes. The interweaving of MGEs significantly amplifies the

potential for dissemination of ARGs (14, 40).

Beyond revealing the remarkable diversity of ARGs and MGEs in *S. aureus*, our study importantly

evidenced numerous associations between these elements. The high degree of similarities between

MGE/ARG associations observed in An-SA and Hm-SA strongly suggested unrestricted dissemination

of MGEs across hosts. This finding implies that An-SA could serve as a reservoir for Hm-SA and vice

versa. Our results identified a few dominant STs within Hm-SA or An-SA populations, typically

harboring the same set of ARG/MGE associations. This observation highlights clonal expansion as a

major driver of dissemination in *S. aureus*. This is the case for *S. aureus* ST22, initially identified in

United Kingdom hospitals (46), which has become a globally disseminated epidemic clone and has

spread across Europe and worldwide, impacting both healthcare settings and the community,

including domestic animals like dogs and cats (47, 48). Due to its pandemic character, very few

variations were observed in ST22 MGE content in genomes available in public databases. Although our

study did not capture evidence of such events, the reported presence of a plasmid carrying via Tn*558*

the *cfr* and *fexA* genes in Irish ST22 isolates (25) demonstrates the potential for horizontal transfer

even within this otherwise stable clonal lineage. Besides this specific example of a rare event, our

results also evidenced more frequent transfers. Notably, within the same ST, the observed diversity of

MGEs and ARGs demonstrates that the mobilome and resistome are sources of significant genomic

plasticity, despite core genome similarity. Furthermore, our analysis revealed an intense spread of

certain MGE/ARG associations across diverse STs (Figure 6). For instance, the Rep1/*aadD* association

was found in all five most prevalent STs. Additionally, the Tn*916*/*tet(M)* is primarily associated with

ST398 in An-SA (80% of identified *tet(M)* genes are in ST398), while in Hm-SA, it was found in both

ST398 (40%) and ST239 (20%). This suggests potential host adaptation, in line with recent studies

demonstrating the role of *tet(M)*-carrying Tn*916* transposon, facilitating the jump from An-SA to Hm-

SA (49). Despite lacking their own transfer systems and residing primarily within chromosomes,

transposons also contribute to *S. aureus* genome plasticity and resistome dissemination among STs.

Our findings revealed the presence of transposons in diverse STs, like Tn*554*/*ant(9)-Ia*/*erm(A)* which

was identified in more than 12 STs, and Tn*558*/*fexA* found in 10 An-SA STs. This widespread distribution

across different epidemic clones highlights the horizontal propagation of these elements, suggesting

their important role in shaping the resistome and, more broadly the genomes of *S. aureus*.

In conclusion, this study revealed the remarkable abundance and diversity of MGEs and ARGs within

S. aureus genomes, primarily associated with plasmids and transposons. While it is known that

S. aureus disseminate through clonal waves, our results showed that these clones can also evolve

through horizontal acquisition of MGE/ARG associations that are promoting the dissemination of

resistance genes. This study provides additional evidence that An-SA can serve as reservoirs for Hm-

SA and vice versa, and that the mobilome's only limitation is its transfer efficiency. Plasmids in *S.*

aureus, as primary reservoirs of ARGs, are key players of AMR dissemination. This highlights the critical

need to elucidate the mechanisms governing the epidemic success of MGEs, particularly those

implicated in ARG transfer.

Materials and methods

Genome characterization and phylogenetic analysis

Genome assemblies (complete and draft genomes) of *S. aureus* of animal (n=1,436) and human

sources (n=11,888) were downloaded from the NCBI public database in October 2021. Quality and

completeness of assemblies were evaluated using QUAST v.5.0.2 (50) and BUSCO v.5.2.1 (51) (Figure

S4). All contigs <1000 bp were removed from the analysis. Genomes were characterized according to

their ST using MLST v.2.1.1 (52). Transposons, ICE, IS and CTn were identified using

MobileElementFinder v.1.0.3 (80% coverage and 70% identity) (53). Plasmid replicons (*rep* coding

gene) were detected using PlasmidFinder v.2.1.6 software (databases: v.2021-11-29) (60% coverage

and 90% identity) (54, 55). SCCs were identified using SCC*mec*Finder v.1.2.1 (database extended, 60%

coverage and 90% identity) (54, 56, 57). ARGs were identified using ResFinder v.4.1.7 (database:

v.2021-09-22) (60% coverage and 80% identity) (54, 58) and RGI-CARD (Comprehensive Antibiotic

Resistance Database) (59) (percentage length of the reference sequence >200). Co-occurrence was

considered when several ARGs were identified within the same strain. A phylogenetic tree was

constructed using pyMLST v.2.1.3 with default parameters (60). A neighbor-joining tree based on the

S. aureus core genome was constructed using cgMLST software (https://www.cgmlst.org/ncs) (61). To

avoid duplicates, only one genome with identical MGE and ARG content, geographical origin, collection

year and NCBI bioProject was selected in each group determined by cgMLST with a maximum distance

of 10 different alleles (Figure S4). Following this selection, a total of 9,408 genomes of Hm-SA and 655

genomes of An-SA, carrying at least one ARG and one MGE (except IS) were included in this study.

Identification and characterization of detected mobile genetic elements

Plasmids identification. PlasForest (54, 62, 63), RFPlasmid for *Staphylococcus* species (64), Plasmer

(65) and Plasflow (66) were used with default parameters to identify whether contigs were plasmidic

or chromosomal. A contig was considered plasmidic if predicted by at least three software. Due to the

fragmented nature of draft genomes, some plasmids were split into multiple contigs during assembly.

To identify contigs belonging to the same plasmid, those identified as plasmidic were analyzed using

BLASTN. Contigs were considered part of the same plasmid if they shared more than 60% identity and

coverage with the same reference plasmid sequence. Some previously unclassified contigs were

identified as plasmidic due to their similarity to known plasmids. Plasmids were classified according to

the active protein domain of the Rep protein that was encoded by *rep* gene. All *rep* genes identified

on chromosomal contigs were excluded from the analysis.

Composite transposons identification. A CTn was predicted when two IS were at a specified distance

from each other. The threshold value was 12kb according to the length of the largest CTn identified in

S. aureus (8).

Staphylococcal Chromosomal Cassettes identification. Boundaries of SCC was fixed with *orfX* gene

and the end was adjusted based on the size of the cassette type (https://www.sccmec.org/). SCCs

found scattered across multiple contigs were adjusted based on gene order (synteny) to define their

limits. SCC*mec*Finder v.1.2.1 did not determine the cassette type for some SCC elements, considered

as "indeterminate". SCC lacking *mecA* or *mecC* genes were classified "SCC" instead of SCC*mec*.

Prophages identification. These elements were identified using PHASTEST, last accessed October

2023 (67–69). Only prophages with a "complete" score were retained. Prophages were named

according to the bacteriophage exhibiting the closest similarity, determined by both sequence identity

and the number of common genes.

Integrative and Conjugative Elements identification. In addition to MobileElementFinder v.1.0.3

detection, a BLASTP analysis was performed against a custom database created from literature and

ICEBerg 2.0 database (15, 45, 70, 71). This database included conserved components of the two main

ICE families, Tn*916* and ICE*6013*, identified based on a previous study (70) (Figure S5A). Only conserved

elements meeting the following criteria were considered as ICEs: proximity between elements,

number of conserved elements on the same contig and integrity of the gene (Figure S5B). ICEs

boundaries were extended by 10kb at each end of the ICE, reflecting the average size of Staphylococci

ICE (approximately 30kb).

Identification of associations between antibiotic resistance genes and mobile genetic

elements

To assess the correlation between the number of ARGs and the number of each MGE within a genome,

the Pearson correlation coefficient was calculated using GraphPad Prism 9.5.1 (GraphPad Software

Inc., CA, USA). MGE/ARG associations were identified through co-localization analysis when the MGE

region entirely overlapped the ARG region. This rule was applied only for transposons, ICE, SCC,

prophages and CTn. For plasmids, an ARG was considered associated when it was localized on a

plasmidic contig (see above). For plasmidic contig lacking *rep* gene, associated ARGs were considered

carried by a plasmid of non-determinate (ND) type. Unclassified contigs identified as associated with

SCC, prophage, ICE were reclassified as chromosomal. When multiple ARGs were found within the

same MGE from a single strain, the MGE was considered to harbor multiple ARGs.

Acknowledgments

This work was supported by PhD grant from INRAE (French National Research Institute for Agriculture,

Food and the Environment) and ANSES (French National Agency for Food, Environmental and

Occupational Health and Safety). We are grateful to the Genotoul bioinformatics platform Toulouse

Midi-Pyrenees, France for providing support and storage resources. We address a special thanks to

Marie-Stephane Trotard from Genotoul, for his expertise, advice and installation of all bioinformatics

software. We thank Gaëtan Durand for the python scripts that enabled us to analyze a large volume of

data.

Table

Table 1. Prevalence of mobile genetic elements (MGEs) identified in *S. aureus* genomes of animal (An-SA) and human (Hm-SA) origin.

	Plasmid		Transposon		ICE		SCC		Phage		IS		CTn		All MGEs		All MGEs (except IS)	
Sources	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**	**Hm**	**An**
Quantity of MGEs identified	12,084	929	6,581	499	5,287	436	471	7,518	867	19,124	2,873	52,204	5,585	387	108,383	6,462	56,179	3,589
Mean of MGEs per strain	1.5	1.7	1.4	1.2	1.1	1.2	1	1	2.1	1.6	5.6	4.5	1.7	1.9	11.5	9.9	6.0	5.5
Maximum of MGEs per strain	8	6	7	3	4	3	1	1	6	5	58	41	17	11	78	57	23	16
Strains containing MGEs (%tot genomes)*	8,064 (86%)	532 (81%)	4,707 (50%)	420 (64%)	4,751 (50%)	363 (55%)	7,518 (80%)	471 (72%)	9,090 (97%)	554 (85%)	9,286 (99%)	636 (97%)	3,368 (36%)	206 (31%)	9,408 (100%)	655 (100%)	9,408 (100%)	655 (100%)
MGEs specific to sources	203	25	2	0	0	0	12	1	30	4	35	8	10	2	292	40	257	32
MGEs shared between sources (%MGEs shared)*	69 (23%)		9 (82%)		3 (100%)		18 (58%)		51 (60%)		21 (33%)		10 (45%)		181 (35%)		160 (36%)	

411 *ICE: Integrative and Conjugative Element; SCC: Staphylococcal chromosomal cassette; IS: Insertion Sequence; CTn: Composite transposon.*
412 ** Number of MGE/total number of genomes (in %)*
413 ***Number of MGE shared between sources/total number of type of MGEs (in %)*

References

1. DeLeo FR, Chambers HF. 2009. Reemergence of antibiotic-resistant *Staphylococcus aureus* in the genomics era. J Clin Invest 119:2464–2474. https://doi.org/10.1172/JCI38226.

2. Weems Jr JJ. 2001. The many faces of *Staphylococcus aureus* infection. Postgrad Med 110:24–36. https://doi.org/10.3810/pgm.2001.10.1042.

3. Peton V, Le Loir Y. 2014. *Staphylococcus aureus* in veterinary medicine. Infect Genet Evol 21:602–615. https://doi.org/10.1016/j.meegid.2013.08.011.

4. Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Aguilar GR, Gray A, Han C, Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G, Hamadani BHK, Kumaran EAP, McManigal B, Achalapong S, Agarwal R, Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Babin F-X, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Berkley JA, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalheiro C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Chiurchiù S, Chowdhury F, Donatien RC, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe S, Day NPJ, Luca MD, Dokova K, Dramowski A, Dunachie SJ, Bich TD, Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garcia C, Garrett D, Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Hsia Y, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Jenney AWJ, Khorana M, Khusuwan S, Kissoon N, Kobeissi E, Kostyanev T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Lim K, Limmathurotsakul D, Loftus MJ, Lunn M, Ma J, Manoharan A, Marks F, May J, Mayxay M, Mturi N, Munera-Huertas T, Musicha P, Musila LA, Mussi-Pinhata MM, Naidu RN, Nakamura T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Ochoa TJ, Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Ounchanum P, Pak GD, Paredes JL, Peleg AY, Perrone C, Phe T, Phommasone K, Plakkal N, Ponce-de-Leon A, Raad M, Ramdin T, Rattanavong S, Riddell A, Roberts T, Robotham JV, Roca A, Rosenthal VD, Rudd KE, Russell N, Sader HS, Saengchan W, Schnall J, Scott JAG, Seekaew S, Sharland M, Shivamallappa M,

Sifuentes-Osornio J, Simpson AJ, Steenkeste N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong

A, Thwaites G, Tigoi C, Turner C, Turner P, Doorn HR van, Velaphi S, Vongpradith A, Vongsouvath

M, Vu H, Walsh T, Walson JL, Waner S, Wangrangsimakul T, Wannapinij P, Wozniak T, Sharma

TEMWY, Yu KC, Zheng P, Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi M.

2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. The

Lancet 399:629–655. https://doi.org/10.1016/S0140-6736(21)02724-0.

5.   Lakhundi S, Zhang K. 2018. Methicillin-Resistant *Staphylococcus aureus*: Molecular

Characterization, Evolution, and Epidemiology. Clin Microbiol Rev 31:e00020-18.

https://doi.org/10.1128/CMR.00020-18.

6.   Wright GD. 2007. The antibiotic resistome: the nexus of chemical and genetic diversity. 3. Nat

Rev Microbiol 5:175–186. https://doi.org/10.1038/nrmicro1614.

7.   Schwarz S.Feßler AT.Loncaric I.Wu C.Kadlec K.Wang Y.Shen J. 2018. Antimicrobial Resistance

among Staphylococci of Animal Origin. Microbiol Spectr 6:10.1128/microbiolspec.arba-0010-

2017. https://doi.org/10.1128/microbiolspec.arba-0010-2017.

8.   Firth N, Jensen SO, Kwong SM, Skurray RA, Ramsay JP. 2018. Staphylococcal Plasmids,

Transposable and Integrative Elements. Microbiol Spectr 6:10.1128/microbiolspec.gpp3-0030–

2018. https://doi.org/10.1128/microbiolspec.gpp3-0030-2018.

9.   Alibayov B, Baba-Moussa L, Sina H, Zdeňková K, Demnerová K. 2014. *Staphylococcus aureus*

mobile genetic elements. Mol Biol Rep 41:5005–5018. https://doi.org/10.1007/s11033-014-

3367-3.

10. Frost LS, Leplae R, Summers AO, Toussaint A. 2005. Mobile genetic elements: the agents of open

source evolution. 9. Nat Rev Microbiol 3:722–732. https://doi.org/10.1038/nrmicro1235.

11.   Siefert JL. 2009. Defining the Mobilome, p. 13–27. *In* Gogarten, MB, Gogarten, JP, Olendzenski,

462   LC (eds.), Horizontal Gene Transfer: Genomes in Flux. Humana Press, Totowa, NJ.

463   https://doi.org/10.1007/978-1-60327-853-9_2.

464   12.   Haaber J, Penadés JR, Ingmer H. 2017. Transfer of Antibiotic Resistance in *Staphylococcus*

465   *aureus*. Trends Microbiol 25:893–905. https://doi.org/10.1016/j.tim.2017.05.011.

466   13.   Partridge SR, Kwong SM, Firth N, Jensen SO. 2018. Mobile Genetic Elements Associated with

467   Antimicrobial   Resistance.   Clin   Microbiol   Rev   31:10.1128/cmr.00088-17.

468   https://doi.org/10.1128/cmr.00088-17.

469   14.   Smyth DS, Robinson DA. 2009. Integrative and Sequence Characteristics of a Novel Genetic

470   Element,   ICE*6013*,   in   *Staphylococcus   aureus*.   J   Bacteriol   191:5964–5975.

471   https://doi.org/10.1128/jb.00352-09.

472   15.   Sansevere EA, Robinson DA. 2017. Staphylococci on ICE: Overlooked agents of horizontal gene

473   transfer. Mob Genet Elem 7:1–10. https://doi.org/10.1080/2159256X.2017.1368433.

474   16.   Deghorain M, Melderen LV. 2012. The Staphylococci Phages Family: An Overview. Viruses

475   4:3316. https://doi.org/10.3390/v4123316.

476   17.   Malachowa N, DeLeo FR. 2010. Mobile genetic elements of *Staphylococcus aureus*. Cell Mol Life

477   Sci CMLS 67:3057–3071. https://doi.org/10.1007/s00018-010-0389-4.

478   18.   Jensen SO, Lyon BR. 2009. Genetics of antimicrobial resistance in *Staphylococcus aureus*. Future

479   Microbiol 4:565–582. https://doi.org/10.2217/fmb.09.30.

480   19.   Lowy FD. 2003. Antimicrobial resistance: the example of *Staphylococcus aureus*. J Clin Invest

481   111:1265–1273. https://doi.org/10.1172/JCI200318535.

482   20.   Pennone V, Prieto M, Álvarez-Ordóñez A, Cobo-Diaz JF. 2022. Antimicrobial Resistance Genes
483   Analysis of Publicly Available *Staphylococcus aureus* Genomes. 11. Antibiotics 11:1632.
484   https://doi.org/10.3390/antibiotics11111632.

485   21.   Botelho J, Cazares A, Schulenburg H. 2023. The ESKAPE mobilome contributes to the spread of
486   antimicrobial resistance and CRISPR-mediated conflict between mobile genetic elements.
487   Nucleic Acids Res 51:236–252. https://doi.org/10.1093/nar/gkac1220.

488   22.   Wang Y, Lv Y, Cai J, Schwarz S, Cui L, Hu Z, Zhang R, Li J, Zhao Q, He T, Wang D, Wang Z, Shen Y,
489   Li Y, Feßler AT, Wu C, Yu H, Deng X, Xia X, Shen J. 2015. A novel gene, *optrA*, that confers
490   transferable resistance to oxazolidinones and phenicols and its presence in *Enterococcus*
491   *faecalis* and *Enterococcus faecium* of human and animal origin. J Antimicrob Chemother
492   70:2182–2190. https://doi.org/10.1093/jac/dkv116.

493   23.   Fan R, Li D, Wang Y, He T, Feßler AT, Schwarz S, Wu C. 2016. Presence of the *optrA* Gene in
494   Methicillin-Resistant *Staphylococcus sciuri* of Porcine Origin. Antimicrob Agents Chemother
495   60:7200–7205. https://doi.org/10.1128/aac.01591-16.

496   24.   Jiang F, Kong Z, Liu K, Cheng C, Jiang T, Ma P, Li R. 2022. Phenotypic and genotypic
497   characterisation of linezolid-resistant coagulase-negative staphylococci possessing *cfr*-carrying
498   plasmid. J Glob Antimicrob Resist 28:226–232. https://doi.org/10.1016/j.jgar.2022.01.008.

499   25.   Shore AC, Lazaris A, Kinnevey PM, Brennan OM, Brennan GI, O'Connell B, Feßler AT, Schwarz S,
500   Coleman DC. 2016. First Report of *cfr*-Carrying Plasmids in the Pandemic Sequence Type 22
501   Methicillin-Resistant *Staphylococcus aureus* Staphylococcal Cassette Chromosome *mec* Type IV
502   Clone. Antimicrob Agents Chemother 60:3007–3015. https://doi.org/10.1128/AAC.02949-15.

503   26.   Freitas AR, Coque TM, Novais C, Hammerum AM, Lester CH, Zervos MJ, Donabedian S, Jensen
504   LB, Francia MV, Baquero F, Peixe L. 2020. Human and Swine Hosts Share Vancomycin-Resistant

Enterococcus faecium CC17 and CC5 and *Enterococcus faecalis* CC2 Clonal Clusters Harboring Tn*1546* on Indistinguishable Plasmids. J Clin Microbiol 49:925–931. https://doi.org/10.1128/jcm.01750-10.

27. Kwong SM, Ramsay JP, Jensen SO, Firth N. 2017. Replication of Staphylococcal Resistance Plasmids. Front Microbiol 8:2279. https://doi.org/10.3389/fmicb.2017.02279.

28. Jensen LB, Garcia-Migura L, Valenzuela AJS, Løhr M, Hasman H, Aarestrup FM. 2010. A classification system for plasmids from enterococci and other Gram-positive bacteria. J Microbiol Methods 80:25–43. https://doi.org/10.1016/j.mimet.2009.10.012.

29. Douarre P-E, Mallet L, Radomski N, Felten A, Mistou M-Y. 2020. Analysis of COMPASS, a New Comprehensive Plasmid Database Revealed Prevalence of Multireplicon and Extensive Diversity of IncF Plasmids. Front Microbiol 11.

30. Carattoli A. 2009. Resistance Plasmid Families in Enterobacteriaceae. Antimicrob Agents Chemother 53:2227–2238. https://doi.org/10.1128/aac.01707-08.

31. Villa L, García-Fernández A, Fortini D, Carattoli A. 2010. Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. J Antimicrob Chemother 65:2518–2529. https://doi.org/10.1093/jac/dkq347.

32. Novick RP. 1989. Staphylococcal Plasmids and Their Replication. Annu Rev Microbiol 43:537–563. https://doi.org/10.1146/annurev.mi.43.100189.002541.

33. Wassenaar TM, Ussery DW, Ingmer H. 2016. The *qacC* Gene Has Recently Spread between Rolling Circle Plasmids of Staphylococcus, Indicative of a Novel Gene Transfer Mechanism. Front Microbiol 7:1528. https://doi.org/10.3389/fmicb.2016.01528.

526   34.   Lee CA, Thomas J, Grossman AD. 2012. The *Bacillus subtilis* Conjugative Transposon ICE*Bs1*

527   Mobilizes Plasmids Lacking Dedicated Mobilization Functions. J Bacteriol 194:3165–3172.

528   https://doi.org/10.1128/JB.00301-12.

529   35.   Showsh SA, Andrews RE. 1999. Analysis of the Requirement for a pUB110 mob Region during

530   Tn*916*-Dependent Mobilization. Plasmid 41:179–186. https://doi.org/10.1006/plas.1999.1398.

531   36.   Projan SJ, Novick R. 1988. Comparative analysis of five related staphylococcal plasmids. Plasmid

532   19:203–221. https://doi.org/10.1016/0147-619X(88)90039-X.

533   37.   Macrina FL, Archer GL. 1993. Conjugation and Broad Host Range Plasmids in Streptococci and

534   Staphylococci, p. 313–329. *In* Clewell, DB (ed.), Bacterial Conjugation. Springer US, Boston, MA.

535   https://doi.org/10.1007/978-1-4757-9357-4_12.

536   38.   O'Brien FG, Yui Eto K, Murphy RJT, Fairhurst HM, Coombs GW, Grubb WB, Ramsay JP. 2015.

537   Origin-of-transfer sequences facilitate mobilisation of non-conjugative antimicrobial-resistance

538   plasmids   in   *Staphylococcus   aureus*.   Nucleic   Acids   Res   43:7971–7983.

539   https://doi.org/10.1093/nar/gkv755.

540   39.   O'Brien FG, Ramsay JP, Monecke S, Coombs GW, Robinson OJ, Htet Z, Alshaikh FAM, Grubb WB.

541   2015. *Staphylococcus aureus* plasmids without mobilization genes are mobilized by a novel

542   conjugative   plasmid   from   community   isolates.   J   Antimicrob   Chemother   70:649–652.

543   https://doi.org/10.1093/jac/dku454.

544   40.   Sansevere EA, Luo X, Park JY, Yoon S, Seo KS, Robinson DA. 2017. Transposase-Mediated

545   Excision, Conjugative Transfer, and Diversity of ICE*6013* Elements in *Staphylococcus aureus*. J

546   Bacteriol 199:10.1128/jb.00629-16. https://doi.org/10.1128/jb.00629-16.

41. Dechêne-Tempier M, de Boisséson C, Lucas P, Bougeard S, Libante V, Marois-Créhan C, Payot S.

2024. Virulence genes, resistome and mobilome of *Streptococcus suis* strains isolated in France.

Microb Genomics 10:001224. https://doi.org/10.1099/mgen.0.001224.

42. Roberts AP, Mullany P. 2011. Tn916-like genetic elements: a diverse group of modular mobile

elements   conferring   antibiotic   resistance.   FEMS   Microbiol   Rev   35:856–871.

https://doi.org/10.1111/j.1574-6976.2011.00283.x.

43. León-Sampedro R, Novais C, Peixe L, Baquero F, Coque TM. 2016. Diversity and Evolution of the

Tn*5801-tet(M)*-Like Integrative and Conjugative Elements among Enterococcus, Streptococcus,

and   Staphylococcus.   Antimicrob   Agents   Chemother   60:1736–1746.

https://doi.org/10.1128/AAC.01864-15.

44. Clewell DB, Weaver KE, Dunny GM, Coque TM, Francia MV, Hayes F. 2014. Extrachromosomal

and Mobile Elements in Enterococci: Transmission, Maintenance, and Epidemiology, *In* Gilmore,

MS, Clewell, DB, Ike, Y, Shankar, N (eds.), Enterococci: From Commensals to Leading Causes of

Drug Resistant Infection. Massachusetts Eye and Ear Infirmary, Boston.

45. Soge OO, Beck NK, White TM, No DB, Roberts MC. 2008. A novel transposon, Tn*6009*, composed

of a Tn*916* element linked with a *Staphylococcus aureus mer* operon. J Antimicrob Chemother

62:674–680. https://doi.org/10.1093/jac/dkn255.

46. Cury J, Jové T, Touchon M, Néron B, Rocha EP. 2016. Identification and analysis of integrons and

cassette   arrays   in   bacterial   genomes.   Nucleic   Acids   Res   44:4539–4550.

https://doi.org/10.1093/nar/gkw319.

47. Moodley A, Espinosa-Gongora C, Nielsen SS, McCarthy AJ, Lindsay JA, Guardabassi L. 2012.

Comparative Host Specificity of Human- and Pig- Associated *Staphylococcus aureus* Clonal

Lineages. PLoS ONE 7:e49344. https://doi.org/10.1371/journal.pone.0049344.

48. Toleman MS, Watkins ER, Williams T, Blane B, Sadler B, Harrison EM, Coll F, Parkhill J, Nazareth

B, Brown NM, Peacock SJ. 2017. Investigation of a Cluster of Sequence Type 22 Methicillin-

Resistant *Staphylococcus aureus* Transmission in a Community Setting. Clin Infect Dis Off Publ

Infect Dis Soc Am 65:2069–2077. https://doi.org/10.1093/cid/cix539.

49. Ba X, Matuszewska M, Kalmar L, Fan J, Zou G, Corander D, Raisen CL, Li S, Li L, Weinert LA, Tucker

AW, Grant AJ, Zhou R, Holmes MA. 2023. High-Throughput Mutagenesis Reveals a Role for

Antimicrobial Resistance- and Virulence-Associated Mobile Genetic Elements in *Staphylococcus*

aureus   Host   Adaptation.   Microbiol   Spectr   11:e04213-22.

https://doi.org/10.1128/spectrum.04213-22.

50. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome

assemblies.   Bioinforma   Oxf   Engl   29:1072–1075.

https://doi.org/10.1093/bioinformatics/btt086.

51. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and

Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of

Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol 38:4647–4654.

https://doi.org/10.1093/molbev/msab199.

52. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén

T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-

sequenced bacteria. J Clin Microbiol 50:1355–1361. https://doi.org/10.1128/JCM.06094-11.

53. Johansson MHK, Bortolaia V, Tansirichaiya S, Aarestrup FM, Roberts AP, Petersen TN. 2021.

Detection of mobile genetic elements associated with antibiotic resistance in *Salmonella*

enterica using a newly developed web tool: MobileElementFinder. J Antimicrob Chemother

76:101–109. https://doi.org/10.1093/jac/dkaa390.

54.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+:

594   architecture and applications. BMC Bioinformatics 10:421. https://doi.org/10.1186/1471-2105-

595   10-421.

596   55.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup

597   F, Hasman H. 2014. *In Silico* Detection and Typing of Plasmids using PlasmidFinder and Plasmid

598   Multilocus   Sequence   Typing.   Antimicrob   Agents   Chemother   58:3895–3903.

599   https://doi.org/10.1128/aac.02412-14.

600   56.   International Working Group on the Classification of Staphylococcal Cassette Chromosome

601   Elements (IWG-SCC). 2009. Classification of staphylococcal cassette chromosome *mec*

602   (SCC*mec*): guidelines for reporting novel SCC*mec* elements. Antimicrob Agents Chemother

603   53:4961–4967. https://doi.org/10.1128/AAC.00579-09.

604   57.   Kondo Y, Ito T, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, Hiramatsu K. 2007. Combination

605   of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid

606   identification system for *mec*, *ccr*, and major differences in junkyard regions. Antimicrob Agents

607   Chemother 51:264–274. https://doi.org/10.1128/AAC.00165-06.

608   58.   Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe RL, Rebelo

609   AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G, Bender JK, Stingl K,

610   Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H, Pinholt M, Anjum MF, Duggett

611   NA, Kempf I, Nykäsenoja S, Olkkola S, Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S,

612   Ragimbeau C, Lund O, Aarestrup FM. 2020. ResFinder 4.0 for predictions of phenotypes from

613   genotypes. J Antimicrob Chemother 75:3491–3500. https://doi.org/10.1093/jac/dkaa345.

614   59.   Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA, Edalatmand A, Petkau A,

615   Syed SA, Tsang KK, Baker SJC, Dave M, McCarthy MC, Mukiri KM, Nasir JA, Golbon B, Imtiaz H,

616   Jiang X, Kaur K, Kwong M, Liang ZC, Niu KC, Shan P, Yang JYJ, Gray KL, Hoad GR, Jia B, Bhando T,

Carfrae LA, Farha MA, French S, Gordzevich R, Rachwalski K, Tu MM, Bordeleau E, Dooley D,

Griffiths E, Zubyk HL, Brown ED, Maguire F, Beiko RG, Hsiao WWL, Brinkman FSL, Van Domselaar

G, McArthur AG. 2023. CARD 2023: expanded curation, support for machine learning, and

resistome prediction at the Comprehensive Antibiotic Resistance Database. Nucleic Acids Res

51:D690–D699. https://doi.org/10.1093/nar/gkac920.

60.   Biguenet A, Bordy A, Atchon A, Hocquet D, Valot B. 2023. Introduction and benchmarking of

pyMLST: open-source software for assessing bacterial clonality using core genome MLST.

Microb Genomics 9. https://doi.org/10.1099/mgen.0.001126.

61.   Leopold SR, Goering RV, Witten A, Harmsen D, Mellmann A. 2020. Bacterial Whole-Genome

Sequencing Revisited: Portable, Scalable, and Standardized Analysis for Typing and Detection of

Virulence and Antibiotic Resistance Genes. J Clin Microbiol 52:2365–2370.

https://doi.org/10.1128/jcm.00262-14.

62.   Pradier L, Tissot T, Fiston-Lavier A-S, Bedhomme S. 2021. PlasForest: a homology-based random

forest classifier for plasmid detection in genomic datasets. BMC Bioinformatics 22:349.

https://doi.org/10.1186/s12859-021-04270-w.

63.   Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F,

Wilczynski B, de Hoon MJL. 2009. Biopython: freely available Python tools for computational

molecular biology and bioinformatics. Bioinforma Oxf Engl 25:1422–1423.

https://doi.org/10.1093/bioinformatics/btp163.

64.   van der Graaf-van Bloois L, Wagenaar JA, Zomer AL. 2021. RFPlasmid: predicting plasmid

sequences from short-read assembly data using machine learning. Microb Genomics 7:000683.

https://doi.org/10.1099/mgen.0.000683.

65.   Zhu Q, Gao S, Xiao B, He Z, Hu S. 2023. Plasmer: an Accurate and Sensitive Bacterial Plasmid

Prediction Tool Based on Machine Learning of Shared k-mers and Genomic Features. Microbiol

Spectr 11:e04645-22. https://doi.org/10.1128/spectrum.04645-22.

66.   Krawczyk PS, Lipinski L, Dziembowski A. 2018. PlasFlow: predicting plasmid sequences in

metagenomic data using genome signatures. Nucleic Acids Res 46:e35.

https://doi.org/10.1093/nar/gkx1321.

67.   Wishart DS, Han S, Saha S, Oler E, Peters H, Grant JR, Stothard P, Gautam V. 2023. PHASTEST:

faster than PHASTER, better than PHAST. Nucleic Acids Res 51:W443–W450.

https://doi.org/10.1093/nar/gkad382.

68.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster

version of the PHAST phage search tool. Nucleic Acids Res 44:W16-21.

https://doi.org/10.1093/nar/gkw387.

69.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. Nucleic

Acids Res 39:W347-352. https://doi.org/10.1093/nar/gkr485.

70.   Guglielmini J, Quintais L, Garcillán-Barcia MP, de la Cruz F, Rocha EPC. 2011. The repertoire of

ICE in prokaryotes underscores the unity, diversity, and ubiquity of conjugation. PLoS Genet

7:e1002222. https://doi.org/10.1371/journal.pgen.1002222.

71.   Liu M, Li X, Xie Y, Bi D, Sun J, Li J, Tai C, Deng Z, Ou H-Y. 2019. ICEberg 2.0: an updated database

of bacterial integrative and conjugative elements. Nucleic Acids Res 47:D660–D665.

https://doi.org/10.1093/nar/gky1123.

Figures

Figure 1. Genomic diversity of *S. aureus* genomes of animal (An-SA) (A) and human (Hm-SA) (B) sources. The phylogenetic trees represent 655 An-SA and 9,408 Hm-SA. This tree based on cgMLST distance was constructed using Neighbor-Joining method and visualized using iTOL v6 (https://itol.embl.de). From outer circle to inner circle, host, location, sequence type (ST), Staphylococcal Chromosomal Cassette (SCC) (green), Integrative and Conjugative Element (ICE) (pink), transposon (Tn) (purple) and plasmid (yellow) were represented. The correspondence between colors and each type of host, location, or ST were indicated below the figure. Only STs with a frequency >1% in Hm-SA and/or in An-SA, were represented. Genomes with no associated ST and/or no location were annotated as "not determined" (ND). As they were present in up to 90% of the analyzed genomes, IS and prophages were not represented.

Figure 2. Diversity and occurrence of the most prevalent mobile genetic elements (MGEs) families (plasmids, transposons and integrative and conjugative elements (ICEs)) among *S. aureus* of human (Hm-SA, black) and animal (An-SA, white) origins. Bar chart of the percentage of genomes carrying MGE which corresponds to the number of genome containing at least one MGE out of the total number of genomes analyzed (Hm-SA n=9,408 and An-SA n=655). Only types of MGEs identified in >1% of Hm-SA and/or An-SA genomes are represented in the x-axis.

Figure 3. Occurrence and diversity of Staphylococcal Chromosomal Cassettes (SCC) (A), prophages (B), insertion sequence (IS) (C) and composite transposons (CTn) (D) detected families in *S. aureus* of human (Hm-SA, black) and animal (An-SA, white) sources. Bar chart of the percentage of genomes carrying MGE which corresponds to the number of genomes carrying at least one MGE out of the total

number of genomes analyzed (Hm-SA n=9,408 and An-SA n=655). For SCC and CTn only MGEs

identified in >1% of Hm-SA and/or An-SA genomes were represented. For prophages and IS only MGEs

identified in >5% of Hm-SA and/or An-SA genomes were depicted.

Figure 4. Occurrence of antibiotic resistance genes (ARGs) in *S. aureus* genomes. *S. aureus* hosts were

indicated on the left of the Heatmap, detected ARGs on the bottom and the total number of genomes

analyzed for each host on the right. Hosts grouped under the label 'others' correspond to dolphins

(n=4), bats (n=4), fishes (n=2), cervid (n=1), shrimp (n=1) and monkey (n=1). The bar at the bottom of

the Heatmap corresponds to the families of antibiotics with TMP for trimethoprim and 'others' for

nucleoside (*sat-4*) and mupirocin (*mupA)* families. The percentages of genomes containing ARGs were

obtained by dividing the number of genomes possessing the resistance gene by the total number of

genomes on the top right of the Heatmap. These percentages are represented by different shades of

red color, as shown at the bottom of the Heatmap. Only ARGs with a frequency >0.1% in Hm-SA and/or

in An-SA, were represented. Genes exclusively present in Hm-SA are indicated by an asterisk.

Figure 5. Associations of antibiotic resistance genes (ARGs) and mobile genetic elements (MGEs) in

***S. aureus* genomes of animal (An-SA) (A) and human (Hm-SA) sources (B).** Bar chart representing the

percentage of associations between ARGs and MGEs, with each association assigned a color and

indicated in the legend. When an ARG is not associated, the percentage is represented in gray. Only

the ARGs with a frequency >0.1% in Hm-SA and/or An-SA, were depicted. The percentage of ARG

corresponds to the number of ARG associated with MGE or a combination of MGEs divided by the total

number of this ARG, identified in An-SA or Hm-SA. Tn: transposon; ICE: Integrative and Conjugative

Element; SCC: Staphylococcal chromosomal cassette; CTn: composite transposon.

Figure 6. Co-occurrences of antibiotic resistance genes (ARGs) with plasmids, transposons and

integrative and conjugative elements (ICEs) in *S. aureus* genomes of animal (An-SA) (A) and human

(Hm-SA) (B) sources depending on sequence type (ST). Each ARG, MGE and ST were represented by a

node that is linked by a flow when associated. The color of the ARG nodes corresponds to the type of

MGE with which it is associated: pink for ICE, yellow for plasmid, purple for transposon and brown for

plasmid and transposon. The thickness of the flow is proportional to the percentage of ARG associated

with the MGE and the percentage of MGE associated with ST. This percentage is calculated by dividing

the number of ARG associated with the MGE by the total number of these ARGs identified in An-SA or

Hm-SA. Only ARGs with a frequency >0.1% in Hm-SA and/or in An-SA, and association frequencies of

MGE/ARG >10% were represented. For the MGE/ST association, only the 6 most abundant STs in both

Hm-SA and/or in An-SA were represented. The maximum thickness of the flow corresponds to 100%,

as demonstrated by Tn*916*/*tet(M)* for (A) and Inc18,Rep1/*tet(T)* for (B). Sankey diagrams were created

using SankeyMATIC (http://sankeymatic.com).

Figure 7. Co-occurrences between multiple antibiotic resistance genes (ARGs) and mobile genetic

elements (MGEs) in *S. aureus* of animal (An-SA) (A) and human (Hm-SA) (B) sources. Associations

were represented by links that indicate that ARGs were found within the same genome on the same

MGE. The thickness of the line is proportional to the number of genomes carrying this co-occurrence.

The thickest lines correspond to Rep1/*aac(6')-aph(2'')/aadD/ant(6)-Ia/erm(C)/lnu(B)/lsa(E)/tet(L)*

association identified in 42 An-SA genomes and Tn*554*/*ant(9)-Ia/erm(A)* co-occurrence in 3,643 Hm-

SA genomes. Only associations found in at least two An-SA genomes, or 10 Hm-SA genomes were

shown. SCC: Staphylococcal chromosomal cassette; CTn: Composite transposon; ICE: Integrative and

Conjugative Element.

A

Host
Location
ST
SCC
ICE
Tn
Plasmid

Tree scale: 100 ——

Location

■ Europe	■ North America	■ Asia	■ South America	■ Oceania	■ Africa	■ ND

Host

■ Porcine	■ Bovine	■ Avian	■ Equine	■ Canine	■ Rodent	■ Ovine	■ Caprine
■ Feline	■ Fish	■ Dolphin	■ Bat	■ Cervid	■ Shrimp	■ Monkey	

B

Location
ST
SCC
ICEs
Tn
Plasmid

ST

■ ST5	■ ST8	■ ST22	■ ST398	■ ST105	■ ST239	■ ST30
■ ST45	■ ST36	■ ST1	■ ST59	■ ST121	■ ST15	■ ST9
■ ST97	■ ST612	■ ST3597	■ ST130	■ ST164	■ ST1687	■ ND

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.