Title: Reevaluating the Neural Noise Hypothesis in Dyslexia: Insights from EEG and 7T MRS Biomarkers

Agnieszka Glica[1], Katarzyna Wasilewska[1], Julia Jurkowska[2], Jarosław Żygierewicz[2], Bartosz Kossowski[3], Katarzyna Jednoróg[1*]

[1]Laboratory of Language Neurobiology, Nencki Institute of Experimental Biology, Polish Academy of Sciences, Pasteur 3 Street, 02-093 Warsaw, Poland

[2]Faculty of Physics, University of Warsaw, Pasteur 5 Street, 02-093 Warsaw, Poland

[3]Laboratory of Brain Imaging, Nencki Institute of Experimental Biology, Polish Academy of Sciences, Pasteur 3 Street, 02-093 Warsaw, Poland

*Corresponding Author: Katarzyna Jednoróg, k.jednorog@nencki.edu.pl, Laboratory of Language Neurobiology, Nencki Institute of Experimental Biology, Polish Academy of Sciences, Pasteur 3 Street, 02-093 Warsaw, Poland, +48 22 5892 392

Keywords: dyslexia, neural noise, excitation-inhibition balance, glutamate, gamma-aminobutyric acid, GABA

CRediT: conceptualization: K.J., B.K.; investigation: A.G., K.W., J.J.; data curation: A.G., K.W., J.J.; formal analysis: A.G., K.W.; methodology: K.J., B.K., J.Ż.; project administration: K.J.; funding acquisition: K.J.; supervision: K.J., B.K., J.Ż.; visualization: A.G., K.W., K.J.; writing – original draft: A.G., K.W., K.J.; writing – review & editing: K.J., J.Ż., B.K., J.J.

Competing interests: The authors declare no competing interests.

Abstract

The neural noise hypothesis of dyslexia posits an imbalance between excitatory and inhibitory

(E/I) brain activity as an underlying mechanism of reading difficulties. This study provides

the first direct test of this hypothesis using both EEG power spectrum measures in 120 Polish

adolescents and young adults (60 with dyslexia, 60 controls) and glutamate (Glu) and gamma-

aminobutyric acid (GABA) concentrations from magnetic resonance spectroscopy (MRS) at

7T MRI scanner in half of the sample. Our results, supported by Bayesian statistics, show no

evidence of E/I balance differences between groups, challenging the hypothesis that cortical

hyperexcitability underlies dyslexia. These findings suggest alternative mechanisms must be

explored and highlight the need for further research into the E/I balance and its role in

neurodevelopmental disorders.

INTRODUCTION

According to the neural noise hypothesis of dyslexia, reading difficulties stem from an imbalance between excitatory and inhibitory (E/I) neural activity (Hancock et al., 2017). The hypothesis predicts increased cortical excitation leading to more variable and less synchronous neural firing. This instability supposedly results in disrupted sensory representations and impedes phonological awareness and multisensory integration skills, crucial for learning to read (Hancock et al., 2017). Yet, studies testing this hypothesis are lacking.

The non-invasive measurement of the E/I balance can be derived through assessment of glutamate (Glu) and gamma-aminobutyric acid (GABA) neurotransmitters concentration via magnetic resonance spectroscopy (MRS) (Finkelman et al., 2022) or through various E/I estimations from the electroencephalography (EEG) signal (Ahmad et al., 2022).

MRS measurements of Glu and GABA yielded conflicting findings. Higher Glu concentrations in the midline occipital cortex correlated with poorer reading performance in children (Del Tufo et al., 2018; Pugh et al., 2014), while elevated Glu levels in the anterior cingulate cortex (ACC) corresponded to greater phonological skills (Lebel et al., 2016). Elevated GABA in the left inferior frontal gyrus was linked to reduced verbal fluency in adults (Nakai and Okanoya, 2016), and increased GABA in the midline occipital cortex in children was associated with slower reaction times in a linguistic task (Del Tufo et al., 2018). However, notable null findings exist regarding dyslexia status and Glu levels in the ACC among children (Horowitz-Kraus et al., 2018) as well as Glu and GABA levels in the visual and temporo-parietal cortices in both children and adults (Kossowski et al., 2019).

Both beta (~13-28 Hz) and gamma (> 30 Hz) oscillations may serve as E/I balance indicators (Ahmad et al., 2022), as greater GABA-ergic activity has been associated with greater beta power (Jensen et al., 2005; Porjesz et al., 2002) and gamma power or peak

frequency (Brunel and Wang, 2003; Chen et al., 2017). Resting-state analyses often reported

nonsignificant beta power associations with dyslexia (Babiloni et al., 2012; Fraga González et

al., 2018; Xue et al., 2020), however, one study indicated lower beta power in dyslexic

compared to control boys (Fein et al., 1986). Mixed results were also observed during tasks.

One study found decreased beta power in the dyslexic group (Spironelli et al., 2008), while

the other noted increased beta power relative to the control group (Rippon and Brunswick,

2000). Insignificant relationship between resting gamma power and dyslexia was reported

(Babiloni et al., 2012; Lasnick et al., 2023). When analyzing auditory steady-state responses,

the dyslexic group had a lower gamma peak frequency, while no significant differences in

gamma power were observed (Rufener and Zaehle, 2021). Essentially, the majority of studies

in dyslexia examining gamma frequencies evaluated cortical entrainment to auditory stimuli

(Lehongre et al., 2011; Marchesotti et al., 2020; Van Hirtum et al., 2019). Therefore, the

results from these tasks do not provide direct evidence of differences in either gamma power

or peak frequency between the dyslexic and control groups.

The EEG signal comprises both oscillatory, periodic activity, and aperiodic activity,

characterized by a gradual decrease in power as frequencies rise (1/f signal) (Donoghue et al.,

2020). Recently recognized as a biomarker of E/I balance, a lower exponent of signal decay

(flatter slope) indicates a greater dominance of excitation over inhibition in the brain, as

shown by the simulation models of local field potentials, ratio of AMPA/GABA$_a$ synapses in

the rat hippocampus (Gao et al., 2017) and recordings under propofol or ketamine in

macaques and humans (Gao et al., 2017; Waschke et al., 2021). However, there are also

pharmacological studies providing mixed results (Colombo et al., 2019; Salvatore et al.,

2024). Nonetheless, the 1/f signal has shown associations with various conditions putatively

characterized by changes in E/I balance, such as early development in infancy

(Schaworonkow and Voytek, 2021), healthy aging (Voytek et al., 2015) and

neurodevelopmental disorders like ADHD (Ostlund et al., 2021), autism spectrum disorder

(Manyukhina et al., 2022) or schizophrenia (Molina et al., 2020). Despite its potential

relevance, the evaluation of the 1/f signal in dyslexia remains limited to one study, revealing

flatter slopes among dyslexic compared to control participants at rest (Turri et al., 2023),

thereby lending support to the notion of neural noise in dyslexia.

Here, we examined both EEG (1/f signal, beta, and gamma oscillations during both

rest and a spoken language task) and MRS (Glu and GABA) biomarkers of E/I balance in

participants with dyslexia and age-matched controls. The neural noise hypothesis predicts

flatter slopes of 1/f signal, decreased beta and gamma power, and higher Glu concentrations

in the dyslexic group. Furthermore, we tested the relationships between different E/I

measures. Flatter slopes of 1/f signal should be related to higher Glu level, while enhanced

beta and gamma power to increased GABA level.

RESULTS

No evidence for group differences in the EEG E/I biomarkers

We recruited 120 Polish adolescents and young adults – 60 with dyslexia diagnosis

and 60 controls matched in sex, age, and family socio-economic status. The dyslexic group

scored lower in all reading and reading-related tasks and higher in the Polish version of the

Adult Reading History Questionnaire (ARHQ-PL) (Bogdanowicz et al., 2015), where a higher

score indicates a higher risk of dyslexia (see Table 1). Although all participants were within

the intellectual norm, the dyslexic group scored lower on the IQ scale (including nonverbal

subscale only) than the control group. However, the Bayesian statistics did not provide

evidence for the difference between groups in the nonverbal IQ.

Table 1.

Demographic and Behavioral Characteristics of the Entire Sample of 120 Participants

	DYS 28 F, 32 M		CON 28 F, 32 M		*t* (df)	*p*	Cohen's *d*	BF$_{10}$
	M	*SD*	*M*	*SD*				
Demographics								
Age	19.41	3.18	19.54	2.96	0.25 (118)	.806	0.05	0.20
Mother's education (years)	17.20	3.36	16.58	2.28	-1.19 (103.89)	.235	-0.22	0.37
Father's education (years)	16.12[a]	3.10[a]	17.13[a]	3.27[a]	1.71 (114)	.091	0.32	0.73
IQ	103.56[b]	11.83[b]	111.12	10.43	3.70 (117)	**< .001**	0.68	75.31
Nonverbal IQ (scaled score)	10.40	2.94	11.62	2.57	2.42 (118)	**.017**	0.44	2.63
ARHQ-PL	51.50	9.70	25.47	8.00	-16.04 (113.87)	**< .001**	-2.93	>10000
Reading and reading-related tasks								
Words/min	108.38	20.93	134.57	13.29	8.18 (99.90)	**< .001***	1.49	>10000
Pseudowords/min	56.75	14.16	83.43	17.04	9.33 (118)	**< .001***	1.70	>10000
RAN objects (s)	32.12	5.11	28.70	4.43	-3.92 (118)	**< .001***	-0.72	149.93
RAN colors (s)	35.83	6.82	31.18	5.73	-4.04 (118)	**< .001***	-0.74	229.96
RAN digits(s)	19.32	4.61	16.25	2.94	-4.34 (100.28)	**< .001***	-0.79	642.86
RAN letters (s)	22.70	4.53	19.68	3.16	-4.23 (105.42)	**< .001***	-0.77	433.23
Reading comprehension (s)	64.47	20.13	43.72	9.63	-7.20 (84.66)	**< .001***	-1.32	>10000
Phoneme deletion (% correct)	76.41	24.68	91.47	9.07	4.44 (74.66)	**< .001***	0.81	898.25
Spoonerisms phonemes (% correct)	54.29	35.42	82.74	22.06	5.28 (98.78)	**< .001***	0.96	>10000
Spoonerisms syllables	46.94	30.61	73.06	23.98	5.20 (111.62)	**< .001***	0.95	>10000

(% correct)

Orthographic awareness (accuracy/time)	0.33	0.13	0.53	0.14	8.12 (118)	**< .001***	1.48	>10000
Perception speed (sten score)	3.32	2.04	4.50	1.67	3.48 (118)	**< .001***	0.64	38.71
Digits forward	5.53	1.64	6.98	1.95	4.40 (118)	**< .001***	0.80	792.55
Digits backward	5.25	1.49	7.33	2.25	5.99 (102.59)	**< .001***	1.09	>10000

Note. DYS – dyslexic group; CON – control group; F – females, M – males. BF_{10} – Bayes Factor indicating ratio of the likelihood of an alternative hypothesis (H1) to a null hypothesis (H0). ARHQ-PL – Polish version of the Adult Reading History Questionnaire. RAN – rapid automatized naming. Boldface indicates statistical significance at *p* < .05 level (uncorrected).
*Significance after Bonferroni correction for 14 planned comparisons for reading and reading-related tasks;
[a]*n* = 58 (two participants did not provide information about the father's education);
[b]*n* = 59 (one participant did not attempt a verbal subtest of the scale, thus we were not able to calculate overall IQ)

The EEG signal was recorded at rest and during a spoken language task, where

participants listened to a sentence and had to indicate its veracity. In the initial step of the

analysis, we analyzed the aperiodic (exponent and offset) components of the EEG spectrum.

The exponent reflects the steepness of the EEG power spectrum, with a higher exponent

indicating a steeper signal; while the offset represents a uniform shift in power across

frequencies, with a higher offset indicating greater power across the entire EEG spectrum

(Donoghue et al., 2020).

Due to a technical error, the signal from one person (a female from the dyslexic group)

was not recorded during most of the language task and was excluded from the analyses.

Hence, the results are provided for 119 participants – 59 in the dyslexic and 60 in the control

group.

First, aperiodic parameter values were averaged across all electrodes and compared

between groups (dyslexic, control) and conditions (resting state, language task) using a 2x2

repeated measures ANOVA. Age negatively correlated both with the exponent ($r = -.27$, $p =$

.003, $BF_{10} = 7.96$) and offset ($r = -.40$, $p < .001$, $BF_{10} = 3174.29$) in line with previous

investigations (Cellier et al., 2021; McSweeney et al., 2021; Schaworonkow and Voytek,

2021; Voytek et al., 2015), therefore we included age as a covariate. Post-hoc tests are

reported with Bonferroni corrected p-values.

For the mean exponent, we found a significant effect of age ($F(1,116) = 8.90$, $p =$

.003, $\eta^2_p = .071$, $BF_{incl} = 10.47$), while the effects of condition ($F(1,116) = 2.32$, $p = .131$, η^2_p

$= .020$, $BF_{incl} = 0.39$) and group ($F(1,116) = 0.08$, $p = .779$, $\eta^2_p = .001$, $BF_{incl} = 0.40$) were not

significant and Bayes Factor did not provide evidence for either inclusion or exclusion.

Interaction between group and condition ($F(1,116) = 0.16$, $p = .689$, $\eta^2_p = .001$, $BF_{incl} = 0.21$)

was not significant and Bayes Factor indicated against including it in the model.

For the mean offset, we found significant effects of age ($F(1,116) = 22.57$, $p < .001$,

$\eta^2_p = .163$, $BF_{incl} = 1762.19$) and condition ($F(1,116) = 23.04$, $p < .001$, $\eta^2_p = .166$, $BF_{incl} >$

10000) with post-hoc comparison indicating that the offset was lower in the resting state

condition ($M = -10.80$, $SD = 0.21$) than in the language task ($M = -10.67$, $SD = 0.26$, $p_{corr} <$

.001). The effect of group ($F(1,116) = 0.00$, $p = .964$, $\eta^2_p = .000$, $BF_{incl} = 0.54$) was not

significant while Bayes Factor did not provide evidence for either inclusion or exclusion.

Interaction between group and condition was not significant ($F(1,116) = 0.07$, $p = .795$, $\eta^2_p =$

.001, $BF_{incl} = 0.22$) and Bayes Factor indicated against including it in the model.

Next, we restricted analyses to language regions and averaged exponent and offset

values from the frontal electrodes corresponding to the left (F7, FT7, FC5) and right inferior

frontal gyrus (F8, FT8, FC6), as well as temporal electrodes, corresponding to the left (T7,

TP7, TP9) and right superior temporal sulcus, STS (T8, TP8, TP10) (Giacometti et al., 2014;

Scrivener and Reader, 2022). A 2x2x2x2 (group, condition, hemisphere, region) repeated

measures ANOVA with age as a covariate was applied. Power spectra from the left STS at

rest and during the language task are presented in Figure 1A and C, while the results for the

exponent, offset, and beta power are presented in Figure 1B and D.

143
Figure 1. *Selected results for the EEG E/I balance biomarkers. (A) Power spectral densities averaged across 3*
electrodes (T7, TP7, TP9) corresponding to the left superior temporal sulcus (STS) separately for dyslexic (DYS)
and control (CON) groups at rest and (C) during the language task. (B) Plots illustrating results for the
exponent, offset, and the beta power from the left STS electrodes at rest and (D) during the language task.

For the exponent, there were significant effects of age ($F(1,116) = 14.00$, $p < .001$, η^2_p

$= .108$, $BF_{incl} = 11.46$) and condition $F(1,116) = 4.06$, $p = .046$, $\eta^2_p = .034$, $BF_{incl} = 1.88$),

however, Bayesian statistics did not provide evidence for either including or excluding the

condition factor. Furthermore, post-hoc comparisons did not reveal significant differences

between the exponent at rest ($M = 1.51$, $SD = 0.17$) and during the language task ($M = 1.51$,

$SD = 0.18$, $p_{corr} = .546$). There was also a significant interaction between region and group,

although Bayes Factor indicated against including it in the model ($F(1,116) = 4.44$, $p = .037$,

$\eta^2_p = .037$, $BF_{incl} = 0.25$). Post-hoc comparisons indicated that the exponent was higher in the

frontal than in the temporal region both in the dyslexic ($M_{frontal} = 1.54$, $SD_{frontal} = 0.15$, $M_{temporal}$

= 1.49, $SD_{temporal}$ = 0.18, p_{corr} < .001) and in the control group ($M_{frontal}$ = 1.54, $SD_{frontal}$ = 0.17,

$M_{temporal}$ = 1.46, $SD_{temporal}$ = 0.20, p_{corr} < .001). The difference between groups was not

significant either in the frontal (p_{corr} = .858) or temporal region (p_{corr} = .441). The effects of

region ($F(1,116)$ = 1.17, p = .282, η^2_p = .010, BF_{incl} > 10000) and hemisphere ($F(1,116)$ =

1.17, p = .282, η^2_p = .010, BF_{incl} = 12.48) were not significant, although Bayesian statistics

indicated in favor of including them in the model. Furthermore, the interactions between

condition and group ($F(1,116)$ = 0.18, p = .673, η^2_p = .002, BF_{incl} = 3.70), and between

region, hemisphere, and condition ($F(1,116)$ = 0.11, p = .747, η^2_p = .001, BF_{incl} = 7.83) were

not significant, however Bayesian statistics indicated in favor of including these interactions

in the model. The effect of group ($F(1,116)$ = 0.12, p = .733, η^2_p = .001, BF_{incl} = 1.19) was not

significant, while Bayesian statistics did not provide evidence for either inclusion or

exclusion. Any other interactions were not significant and Bayes Factor indicated against

including them in the model. Since Bayes Factor suggested the inclusion of the

condition*group interaction in the model, we further conducted Bayesian *t*-tests to determine

whether this was driven by differences between control and dyslexic groups in either

condition. The results, however, supported the null hypothesis in both the resting state

condition (M_{DYS} = 1.51, SD_{DYS} = 0.16, M_{CON} = 1.50, SD_{CON} = 0.19, BF_{10} = 0.22) and during

the language task (M_{DYS} = 1.52, SD_{DYS} = 0.17, M_{CON} = 1.51, SD_{CON} = 0.19, BF_{10} = 0.20).

In the case of offset, there were significant effects of condition ($F(1,116)$ = 20.88, p <

.001, η^2_p = .153, BF_{incl} > 10000) and region ($F(1,116)$ = 6.18, p = .014, η^2_p = .051, BF_{incl} >

10000). For the main effect of condition, post-hoc comparison indicated that the offset was

lower in the resting state condition (M = -10.88, SD = 0.33) than in the language task (M = -

10.76, SD = 0.38, p_{corr} < .001), while for the main effect of region, post-hoc comparison

indicated that the offset was lower in the temporal (M = -10.94, SD = 0.37) as compared to the

frontal region (M = -10.69, SD = 0.34, p_{corr} < .001). There was also a significant effect of age

($F(1,116) = 20.84$, $p < .001$, $\eta^2_p = .152$, $BF_{incl} = 0.23$) and interaction between condition and

hemisphere, ($F(1,116) = 4.35$, $p = .039$, $\eta^2_p = .036$, $BF_{incl} = 0.21$), although Bayes Factor

indicated against including these factors in the model. Post-hoc comparisons for the

condition*hemisphere interaction indicated that the offset was lower in the resting state

condition than in the language task both in the left ($M_{rest} = -10.85$, $SD_{rest} = 0.34$, $M_{task} = -10.73$,

$SD_{task} = 0.40$, $p_{corr} < .001$) and in the right hemisphere ($M_{rest} = -10.91$, $SD_{rest} = 0.31$, $M_{task} = -$

10.79, $SD_{task} = 0.37$, $p_{corr} < .001$) and that the offset was lower in the right as compared to the

left hemisphere both at rest ($p_{corr} < .001$) and during the language task ($p_{corr} < .001$). The

interactions between region and condition ($F(1,116) = 1.76$, $p = .187$, $\eta^2_p = .015$, $BF_{incl} >$

10000), hemisphere and group ($F(1,116) = 1.58$, $p = .211$, $\eta^2_p = .013$, $BF_{incl} = 1595.18$),

region and group ($F(1,116) = 0.27$, $p = .605$, $\eta^2_p = .002$, $BF_{incl} = 9.32$), as well as between

region, condition, and group ($F(1,116) = 0.21$, $p = .651$, $\eta^2_p = .002$, $BF_{incl} = 2867.18$) were not

significant, although Bayesian statistics indicated in favor of including them in the model. The

effect of group ($F(1,116) = 0.18$, $p = .673$, $\eta^2_p = .002$, $BF_{incl} < 0.00001$) was not significant

and Bayesian statistics indicated against including it in the model. Any other interactions were

not significant and Bayesian statistics indicated against including them in the model or did not

provide evidence for either inclusion or exclusion. Since Bayes Factor suggested the inclusion

of hemisphere*group, region*group and region*condition*group interactions in the model,

we further conducted Bayesian t-tests to determine whether this was driven by differences

between control and dyslexic groups. The results indicated in favor of the null hypothesis both

in the left ($M_{DYS} = -10.78$, $SD_{DYS} = 0.38$, $M_{CON} = -10.80$, $SD_{CON} = 0.36$, $BF_{10} = 0.20$) and right

hemisphere ($M_{DYS} = -10.83$, $SD_{DYS} = 0.32$, $M_{CON} = -10.87$, $SD_{CON} = 0.36$, $BF_{10} = 0.24$) as well

as in the frontal ($M_{DYS} = -10.68$, $SD_{DYS} = 0.34$, $M_{CON} = -10.71$, $SD_{CON} = 0.34$, $BF_{10} = 0.21$) and

temporal regions ($M_{DYS} = -10.92$, $SD_{DYS} = 0.36$, $M_{CON} = -10.96$, $SD_{CON} = 0.38$, $BF_{10} = 0.22$).

Similarly, results for the region*condition*group interaction, indicated in favor of null

hypothesis both in frontal (M_{DYS} = -10.75, SD_{DYS} = 0.31, M_{CON} = -10.77, SD_{CON} = 0.32, BF_{10} =

0.21) and temporal electrodes at rest (M_{DYS} = -10.98, SD_{DYS} = 0.34, M_{CON} = -11.01, SD_{CON} =

0.36, BF_{10} = 0.22) as well as in frontal (M_{DYS} = -10.61, SD_{DYS} = 0.37, M_{CON} = -10.64, SD_{CON} =

0.37, BF_{10} = 0.21) and temporal electrodes during the language task (M_{DYS} = -10.87, SD_{DYS} =

0.39, M_{CON} = -10.90, SD_{CON} = 0.41, BF_{10} = 0.22).

212   Then, we analyzed the aperiodic-adjusted brain oscillations. Since the algorithm did

not find the gamma peak (30-43 Hz) above the aperiodic component in the majority of

participants, we report the results only for the beta (14-30 Hz) power. We performed a similar

regional analysis as for the exponent and offset with a 2x2x2x2 (group, condition,

hemisphere, region) repeated measures ANOVA. However, we did not include age as a

covariate, as it did not correlate with any of the periodic measures. The sample size was 117

(DYS n = 57, CON n = 60) since in 2 participants the algorithm did not find the beta peak

above the aperiodic component in the left frontal electrodes during the task.

220   The analysis revealed a significant effect of condition ($F(1,115)$ = 8.58, p = .004, η^2_p =

.069, BF_{incl} = 5.82) with post-hoc comparison indicating that the beta power was greater

during the language task (M = 0.53, SD = 0.22) than at rest (M = 0.50, SD = 0.19, p_{corr} =

.004). There were also significant effects of region ($F(1,115)$ = 10.98, p = .001, η^2_p = .087,

BF_{incl} = 23.71), and hemisphere ($F(1,115)$ = 12.08, p < .001, η^2_p = .095, BF_{incl} = 23.91). For

the main effect of region, post-hoc comparisons indicated that the beta power was greater in

the temporal (M = 0.52, SD = 0.21) as compared to the frontal region (M = 0.50, SD = 0.19,

p_{corr} = .001), while for the main effect of hemisphere, post-hoc comparisons indicated that the

beta power was greater in the right (M = 0.52, SD = 0.20) than in the left hemisphere (M =

0.51, SD = 0.20, p_{corr} < .001). There was a significant interaction between condition and

region ($F(1,115)$ = 12.68, p < .001, η^2_p = .099, BF_{incl} = 55.26) with greater beta power during

the language task as compared to rest significant in the temporal (M_{rest} = 0.50, SD_{rest} = 0.20,

$M_{task} = 0.55$, $SD_{task} = 0.24$, $p_{corr} < .001$), while not in the frontal region ($M_{rest} = 0.49$, $SD_{rest} =$

0.18, $M_{task} = 0.51$, $SD_{task} = 0.22$, $p_{corr} = .077$). Also, greater beta power in the temporal as

compared to the frontal region was significant during the language task ($p_{corr} < .001$), while

not at rest ($p_{corr} = .283$). The effect of group ($F(1,115) = 0.05$, $p = .817$, $\eta^2_p = .000$, $BF_{incl} <$

0.00001) was not significant and Bayes Factor indicated against including it in the model.

Any other interactions were not significant and Bayesian statistics indicated against including

them in the model or did not provide evidence for either inclusion or exclusion. Descriptive

statistics for EEG results separately for dyslexic and control groups are provided in Table S1

in the Supplementary Material.

Additionally, building upon previous findings which demonstrated differences in

dyslexia in aperiodic and periodic components within the parieto-occipital region (Turri et al.,

2023), we have included analyses for the same cluster of electrodes in the Supplementary

Material. However, in this region, we also did not find evidence for group differences either

in the exponent, offset or beta power.

No evidence for group differences in Glu and GABA+ concentrations in the left STS

The MRS voxel was placed in the left STS, in a region showing highest activation for

both visual and auditory words (compared to control stimuli) localized individually in each

participant, based on an fMRI task. Sample voxel localization and sample MRS spectrum are

presented in Figure 2A and 2B, while overlap of the MRS voxel placement across participants

and MRS spectra separately for the dyslexic and control groups are presented in Figure 2C

and 2D. We decided to analyze the neurometabolites' levels derived from the left STS, as this

region is consistently related to functional and structural differences in dyslexia across

languages (Yan et al., 2021). Moreover, the neural noise hypothesis of dyslexia identifies

perisylvian areas as being affected by increased glutamatergic signaling, and directly predicts

associations between Glu and GABA levels in the superior temporal regions and phonological

skills (Hancock et al., 2017).

Due to financial and logistical constraints, 59 out of the 120 recruited subjects,

selected progressively as the study unfolded, were examined with MRS. Subjects were

matched by age and sex between the dyslexic and control groups. Due to technical issues and

to prevent delays and discomfort for the participants, we collected 54 complete sessions.

Additionally, four datasets were excluded based on our quality control criteria (linewidth > 20

Hz and visual inspection), and three GABA+ estimates exceeded the selected CRLB threshold

(> 20%). Ultimately, we report 50 estimates for Glu (21 participants with dyslexia) and 47 for

GABA+ and Glu/GABA+ ratios (20 participants with dyslexia). Demographic and behavioral

characteristics for the subsample of 47 participants are provided in the Table S2. For

comparability with previous studies in dyslexia (Del Tufo et al., 2018; Pugh et al., 2014) we

report Glu and GABA as a ratio to total creatine (tCr).

For each metabolite, we performed a separate univariate ANCOVA with the effect of

group being tested and voxel's gray matter volume (GMV) as a covariate (see Figure 2E). For

the Glu analysis, we also included age as a covariate, due to negative correlation between

variables ($r = -.35$, $p = .014$, $BF_{10} = 3.41$). The analysis revealed significant effect of GMV

($F(1,46) = 8.18$, $p = .006$, $\eta^2_p = .151$, $BF_{incl} = 12.54$), while the effects of age ($F(1,46) = 3.01$,

$p = .090$, $\eta^2_p = .061$, $BF_{incl} = 1.15$) and group ($F(1,46) = 1.94$, $p = .170$, $\eta^2_p = .040$, $BF_{incl} =$

0.63) were not significant and Bayes Factor did not provide evidence for either inclusion or

exclusion.

Conversely, GABA+ did not correlate with age ($r = -.11$, $p = .481$, $BF_{10} = 0.23$), thus

age was not included as a covariate. The analysis revealed a significant effect of GMV

($F(1,44) = 4.39$, $p = .042$, $\eta^2_p = .091$, $BF_{incl} = 1.64$), however Bayes Factor did not provide

evidence for either inclusion or exclusion. The effect of group was not significant ($F(1,44) =$

0.49, $p = .490$, $\eta^2_p = .011$, $BF_{incl} = 0.35$) although Bayesian statistics did not provide evidence

for either inclusion or exclusion.

Also, Glu/GABA+ ratio did not correlate with age ($r = -.05$, $p = .744$, $BF_{10} = 0.19$),

therefore age was not included as a covariate. The results indicated that the effect of GMV

was not significant ($F(1,44) = 0.95$, $p = .335$, $\eta^2_p = .021$, $BF_{incl} = 0.43$) while Bayes Factor did

not provide evidence for either inclusion or exclusion. The effect of group was not significant

($F(1,44) = 0.01$, $p = .933$, $\eta^2_p = .000$, $BF_{incl} = 0.29$) and Bayes Factor indicated against

including it in the model.

Following a recent study examining developmental changes in both EEG and MRS E/I

biomarkers (McKeon et al., 2024), we calculated an additional measure of Glu/GABA+

imbalance, computed as the absolute residual value from the linear regression of Glu

predicted by GABA+ with greater values indicating greater Glu/GABA+ imbalance. Alike the

previous work (McKeon et al., 2024), we took the square root of this value to ensure a normal

distribution of the data. This measure did not correlate with age ($r = -.05$, $p = .719$, $BF_{10} =$

0.19); thus, age was not included as a covariate. The results indicated that the effect of GMV

was not significant ($F(1,44) = 0.63$, $p = .430$, $\eta^2_p = .014$, $BF_{incl} = 0.37$) while Bayes Factor did

not provide evidence for either inclusion or exclusion. The effect of group was not significant

($F(1,44) = 0.74$, $p = .396$, $\eta^2_p = .016$, $BF_{incl} = 0.39$) although Bayesian statistics did not

provide evidence for either inclusion or exclusion. Descriptive statistics for MRS results

separately for dyslexic and control groups are provided in Table S1 in the Supplementary

Material.

Figure 2. *Selected results for the MRS E/I balance biomarkers. (**A**) Sample MRS voxel localization. (**B**) Sample MRS spectrum with individual total creatine (tCr), Glu and GABA+ contributions plotted below. (**C**) Group results (CON > DYS) from the fMRI localizer task for words compared to the control stimuli (p < .05 FWE cluster threshold) and overlap of the MRS voxel placement across participants. (**D**) MRS spectra separately for the dyslexic (DYS) and control (CON) groups. (**E**) Plots illustrating results for the Glu, GABA+, Glu/GABA+ ratio and the Glu/GABA+ imbalance.*

Correspondence between Glu and GABA+ concentrations and EEG E/I biomarkers is

limited

Next, we investigated correlations between Glu and GABA+ concentrations in the left

STS and EEG biomarkers of E/I balance. Semi-partial correlations were performed (Table 2)

to control for confounding variables – for Glu the effects of age and GMV were regressed, for

GABA+, Glu/GABA+ ratio and Glu/GABA+ imbalance the effect of GMV was regressed,

while for exponents and offsets the effect of age was regressed. For zero-order correlations

between variables see Table S3.

Glu negatively correlated with offset in the left STS both at rest ($r = -.38$, $p = .007$,

$BF_{10} = 6.28$; Figure 3A) and during the language task ($r = -.37$, $p = .009$, $BF_{10} = 5.05$), while

any other correlations between Glu and EEG biomarkers were not significant and Bayesian

statistics indicated in favor of null hypothesis or provided absence of evidence for either

hypothesis. Furthermore, Glu/GABA+ imbalance positively correlated with exponent at rest

both averaged across all electrodes ($r = .29$, $p = .048$, $BF_{10} = 1.21$), as well as in the left STS

electrodes ($r = .35$, $p = .017$, $BF_{10} = 2.87$; Figure 3B) although Bayes Factor provided absence

of evidence for either alternative or null hypothesis. Conversely, GABA+ and Glu/GABA+

ratio were not significantly correlated with any of the EEG biomarkers and Bayesian statistics

indicated in favor of null hypothesis or provided absence of evidence for either hypothesis.

Table 2.
Semi-partial Correlations Between MRS and EEG Biomarkers of Excitatory-Inhibitory
Balance. For Glu the Effects of Age and Gray Matter Volume (GMV) Were Regressed, for
GABA+, Glu/GABA+ Ratio and Glu/GABA+ Imbalance the Effect of GMV was Regressed,
While for Exponents and Offsets the Effect of Age was Regressed

Variable	1. r (BF_{10})	2.	3.	4.	5.	6.	7.	8.
EEG resting state								
1. Glu	–							
2. GABA+	.32*[a] (1.82)	–						
3. Glu/GABA+ ratio	-.08[a] (0.21)	-.91***[a] (>10000)	–					
4. Glu/GABA+ imbalance	.12[a] (0.25)	.31*[a] (1.63)	-.18[a] (0.37)	–				

	1	2	3	4	5	6	7	8
5. Exponent mean (rest)	$-.03_b$ (0.18)	$.04_a$ (0.19)	$-.11_a$ (0.23)	$.29*_a$ (1.21)	–			
6. Offset mean (rest)	$-.21_b$ (0.49)	$.08_a$ (0.21)	$-.17_a$ (0.35)	$.25_a$ (0.69)	$.68***_c$ (>10000)	–		
7. Exponent left STS (rest)	$-.16_b$ (0.32)	$.01_a$ (0.18)	$-.07_a$ (0.20)	$.35*_a$ (2.87)	$.68***_c$ (>10000)	$.45***_c$ (>10000)	–	
8. Offset left STS (rest)	$-.38**_b$ (6.28)	$-.10_a$ (0.23)	$.02_a$ (0.18)	$.18_a$ (0.38)	$.18*_c$ (0.80)	$.47***_c$ (>10000)	$.66***_c$ (>10000)	–
9. Beta power left STS (rest)	$-.12_b$ (0.25)	$.17_a$ (0.33)	$-.25_a$ (0.74)	$.01_a$ (0.18)	$.18*_c$ (0.80)	$.21*_c$ (1.40)	$.51***_c$ (>10000)	$.59***_c$ (>10000)
EEG language task								
5. Exponent mean (task)	$-.10_b$ (0.22)	$.06_a$ (0.20)	$-.15_a$ (0.29)	$.21_a$ (0.49)	–			
6. Offset mean (task)	$-.26_b$ (0.92)	$.09_a$ (0.22)	$-.20_a$ (0.43)	$.22_a$ (0.53)	$.72***_c$ (>10000)	–		
7. Exponent left STS (task)	$-.20_b$ (0.44)	$.01_a$ (0.18)	$-.09_a$ (0.22)	$.22_a$ (0.51)	$.65***_c$ (>10000)	$.51***_c$ (>10000)	–	
8. Offset left STS (task)	$-.37**_b$ (5.05)	$-.08_a$ (0.21)	$.00_a$ (0.18)	$.15_a$ (0.30)	$.29**_c$ (18.31)	$.55***_c$ (>10000)	$.77***_c$ (>10000)	–
9. Beta power left STS (task)	$-.19_b$ (0.42)	$.16_a$ (0.32)	$-.22_a$ (0.55)	$-.03_a$ (0.19)	$.05_c$ (0.13)	$.13_c$ (0.31)	$.47***_c$ (>10000)	$.60***_c$ (>10000)

Note. r – Pearson's correlation coefficient; BF_{10} – Bayes Factor indicating ratio of the likelihood of an alternative hypothesis (H1) to a null hypothesis (H0); mean – values averaged across all electrodes; left STS – values averaged across 3 electrodes corresponding to the left superior temporal sulcus (T7, TP7, TP9).
$***p < .001$ (uncorrected); $**p < .01$ (uncorrected); $*p < .05$ (uncorrected);
[a]$n = 47$; [b]$n = 50$; [c]$n = 119$

Figure 3. *Relationships between EEG and MRS E/I balance biomarkers. (**A**) Semi-partial correlation between offset at rest in the left superior temporal sulcus (STS) electrodes and Glu controlling for age and gray matter*

Testing the paths from neural noise to reading

The neural noise hypothesis of dyslexia predicts impact of the neural noise on reading through the impairment of 1) phonological awareness, 2) lexical access and generalization and 3) multisensory integration (Hancock et al., 2017). Therefore, we analyzed correlations between these variables, reading skills and MRS and EEG biomarkers of E/I balance. For the composite score of phonological awareness, we averaged z-scores from phoneme deletion, phoneme and syllable spoonerisms tasks. For the composite score of lexical access and generalization we averaged z-scores from objects, colors, letters and digits subtests from rapid automatized naming (RAN) task, while for the composite score of reading we averaged z-scores from words and pseudowords read per minute, and text reading time in reading comprehension task. The outcomes from the RAN and reading comprehension task have been transformed from raw time scores to items/time scores in order to provide the same direction of relationships for all z-scored measures, with greater values indicating better skills. For the multisensory integration score we used results from the redundant target effect task reported in our previous work (Glica et al., 2024), with greater values indicating a greater magnitude of multisensory integration.

Age positively correlated with multisensory integration ($r = .38$, $p < .001$, $BF_{10} = 87.98$), composite scores of reading ($r = .22$, $p = .014$, $BF_{10} = 2.24$) and phonological awareness ($r = .21$, $p = .021$, $BF_{10} = 1.59$), while not with the composite score of RAN ($r = .13$, $p = .151$, $BF_{10} = 0.32$). Hence, we regressed the effect of age from multisensory integration, reading and phonological awareness scores and performed semi-partial correlations (Table 3, for zero-order correlations see Table S4).

Phonological awareness positively correlated with offset in the left STS at rest ($r = .18$, $p = .049$, $BF_{10} = 0.77$) and with beta power in the left STS both at rest ($r = .23$, $p = .011$, BF_{10}

= 2.73; Figure 4A) and during the language task ($r = .23$, $p = .011$, $BF_{10} = 2.84$; Figure 4B),

although Bayes Factor provided absence of evidence for either alternative or null hypothesis.

Furthermore, multisensory integration positively correlated with GABA+ concentration ($r =$

$.31$, $p = .034$, $BF_{10} = 1.62$) and negatively with Glu/GABA+ ratio ($r = -.32$, $p = .029$, $BF_{10} =$

1.84), although Bayes Factor provided absence of evidence for either alternative or null

hypothesis. Any other correlations between reading skills and E/I balance biomarkers were

not significant and Bayesian statistics indicated in favor of null hypothesis or provided

absence of evidence for either hypothesis.

Table 3.
Semi-partial Correlations Between Reading, Phonological Awareness, Rapid Automatized
Naming, Multisensory Integration and Biomarkers of Excitatory-Inhibitory Balance. For
Reading, Phonological Awareness and Multisensory Integration the Effect of Age was
Regressed, for Glu the Effects of Age and Gray Matter Volume (GMV) Were Regressed, for
GABA+, Glu/GABA+ Ratio and Glu/GABA+ Imbalance the Effect of GMV was Regressed,
While for Exponents and Offsets the Effect of Age was Regressed

Variable	1. r (BF_{10})	2.	3.	4.
EEG resting state				
1. Reading	–			
2. Phonological awareness	.60***$_c$ (>10000)	–		
3. RAN	.71***$_c$ (>10000)	.48***$_c$ (>10000)	–	
4. Multisensory integration	.16$_d$ (0.41)	.25*$_d$ (2.09)	.02$_d$ (0.14)	–
5. Glu	-.03$_b$ (0.18)	-.12$_b$ (0.25)	-.05$_b$ (0.19)	.03$_b$ (0.18)
6. GABA+	-.18$_a$ (0.36)	-.06$_a$ (0.20)	-.23$_a$ (0.56)	.31*$_a$ (1.62)
7. Glu/GABA+ ratio	.07$_a$ (0.20)	.04$_a$ (0.19)	.09$_a$ (0.22)	-.32*$_a$ (1.84)
8. Glu/GABA+ imbalance	-.21$_a$ (0.47)	-.08$_a$ (0.21)	-.20$_a$ (0.44)	.17$_a$ (0.33)
9. Exponent mean (rest)	-.08$_c$ (0.17)	.10$_c$ (0.20)	-.06$_c$ (0.14)	.02$_d$ (0.14)
10. Offset mean (rest)	.06$_c$ (0.14)	.14$_c$ (0.35)	.03$_c$ (0.12)	.16$_d$ (0.38)

11. Exponent left STS (rest)	-.08$_c$ (0.16)	.06$_c$ (0.14)	-.04$_c$ (0.12)	-.04$_d$ (0.14)
12. Offset left STS (rest)	.12$_c$ (0.25)	.18*$_c$ (0.77)	.08$_c$ (0.17)	.08$_d$ (0.18)
13. Beta power left STS (rest)	.04$_c$ (0.13)	.23*$_c$ (2.73)	-.04$_c$ (0.12)	.05$_d$ (0.15)
EEG language task				
9. Exponent mean (task)	-.07$_c$ (0.16)	.13$_c$ (0.30)	-.10$_c$ (0.21)	.01$_d$ (0.14)
10. Offset mean (task)	.05$_c$ (0.13)	.14$_c$ (0.34)	.01$_c$ (0.12)	.18$_d$ (0.50)
11. Exponent left STS (task)	-.03$_c$ (0.12)	.09$_c$ (0.18)	-.07$_c$ (0.15)	-.04$_d$ (0.14)
12. Offset left STS (task)	.13$_c$ (0.28)	.18$_c$ (0.71)	.07$_c$ (0.15)	.09$_d$ (0.19)
13. Beta power left STS (task)	.07$_c$ (0.15)	.23*$_c$ (2.84)	.02$_c$ (0.12)	.15$_d$ (0.33)

Note. r – Pearson's correlation coefficient; BF_{10} – Bayes Factor indicating ratio of the likelihood of an alternative hypothesis (H1) to a null hypothesis (H0); mean – values averaged across all electrodes; left STS – values averaged across 3 electrodes corresponding to the left superior temporal sulcus (T7, TP7, TP9); ***$p < .001$ (uncorrected); **$p < .01$ (uncorrected); *$p < .05$ (uncorrected); [a]$n = 47$; [b]$n = 50$; [c]$n = 119$; [d]$n = 87$

Given that beta power correlated with phonological awareness, and considering the prediction that neural noise impedes reading by affecting phonological awareness — we examined this relationship through a mediation model. Since phonological awareness correlated with beta power in the left STS both at rest and during language task, the outcomes from these two conditions were averaged prior to the mediation analysis. Macro PROCESS v4.2 (Hayes, 2017) on IBM SPSS Statistics v29 with model 4 (simple mediation) with 5000 Bootstrap samples to assess the significance of indirect effect was employed. Since age correlated both with phonological awareness and reading, we also included age as a covariate.

The results indicated that both effects of beta power in the left STS ($b = .96$, $t(116) = 2.71$, $p = .008$, $BF_{incl} = 7.53$) and age ($b = .06$, $t(116) = 2.55$, $p = .012$, $BF_{incl} = 5.98$) on phonological awareness were significant. The effect of phonological awareness on reading was also significant ($b = .69$, $t(115) = 8.16$, $p < .001$, $BF_{incl} > 10000$), while the effects of beta power ($b = -.42$, $t(115) = -1.25$, $p = .213$, $BF_{incl} = 0.52$) and age ($b = .03$, $t(115) = 1.18$, $p = .241$, $BF_{incl} = 0.49$) on reading were not significant when controlling for phonological

awareness. Finally, the indirect effect of beta power on reading through phonological

awareness was significant (b = .66, SE = .24, 95% CI = [.24, 1.18]), while the total effect of

beta power was not significant (b = .24, t(116) = 0.61, p = .546, BF_{incl} = 0.41). The results

from the mediation analysis are presented in Figure 4D.

Although similar mediation analysis could have been conducted for the Glu/GABA+

ratio, multisensory integration, and reading based on the correlations between these variables,

we did not test this model due to the small sample size (47 participants), which resulted in

insufficient statistical power.

414
Figure 4. *Associations between beta power, phonological awareness and reading. (**A**) Semi-partial correlation*
*between phonological awareness controlling for age and beta power (in the left STS electrodes) at rest and (**B**)*
*during the language task. (**C**) Partial correlation between phonological awareness and reading controlling for*
*age. (**D**) Mediation analysis results. Unstandardized b regression coefficients are presented. Age was included in*
the analysis as a covariate. 95% CI – 95% confidence intervals; left STS – values averaged across 3 electrodes
corresponding to the left superior temporal sulcus (T7, TP7, TP9).
421

DISCUSSION

The current study aimed to validate the neural noise hypothesis of dyslexia (Hancock

et al., 2017) utilizing E/I balance biomarkers from EEG power spectra and ultra-high-field

MRS. Contrary to its predictions, we did not observe differences either in 1/f slope, beta

power, or Glu and GABA+ concentrations in participants with dyslexia. Relations between

E/I balance biomarkers were limited to significant correlations between Glu and the offset

when controlling for age, and between Glu/GABA+ imbalance and the exponent.

In terms of EEG biomarkers, our study found no evidence of group differences in the

aperiodic components of the EEG signal. In most of the models, we did not find evidence for

either including or excluding the effect of the group when Bayesian statistics were evaluated.

The only exception was the regional analysis for the offset, where results indicated against

including the group factor in the model. These findings diverge from previous research on an

Italian cohort, which reported decreased exponent and offset in the dyslexic group at rest,

specifically within the parieto-occipital region, but not the frontal region (Turri et al., 2023).

Despite our study involving twice the number of participants and utilizing a longer acquisition

time, we observed no group differences, even in the same cluster of electrodes (refer to

Supplementary Material). The participants in both studies were of similar ages. The only

methodological difference – EEG acquisition with eyes open in our study versus both eyes-

open and eyes-closed in the work by Turri and colleagues (2023) – cannot fully account for

the overall lack of group differences observed. The diverging study outcomes highlight the

importance of considering potential inflation of effect sizes in studies with smaller samples.

Although a lower exponent of the EEG power spectrum has been associated with other

neurodevelopmental disorders, such as ADHD (Ostlund et al., 2021) or ASD (but only in

children with IQ below average) (Manyukhina et al., 2022), our study suggests that this is not

the case for dyslexia. Considering the frequent comorbidity of dyslexia and ADHD (Germanò

et al., 2010; Langer et al., 2019), increased neural noise could serve as a common underlying

mechanism for both disorders. However, our specific exclusion of participants with a

comorbid ADHD diagnosis indicates that the EEG spectral exponent cannot serve as a

neurobiological marker for dyslexia in isolation. No information regarding such exclusion

criteria was provided in the study by Turri et al. (2023); thus, potential comorbidity with

ADHD may explain the positive findings related to dyslexia reported therein.

Regarding the aperiodic-adjusted oscillatory EEG activity, Bayesian statistics for beta

power, indicated in favor of excluding the group factor from the model. Non-significant group

differences in beta power at rest have been previously reported in studies that did not account

for aperiodic components (Babiloni et al., 2012; Fraga González et al., 2018; Xue et al.,

2020). This again contrasts with the study by Turri et al. (2023), which observed lower

aperiodic-adjusted beta power (at 15-25 Hz) in the dyslexic group. Concerning beta power

during task, our results also contrast with previous studies which showed either reduced

(Spironelli et al., 2008) or increased (Rippon and Brunswick, 2000) beta activity in

participants with dyslexia. Nevertheless, since both of these studies employed phonological

tasks, involved children's samples and did not account for aperiodic activity, their relevance

to our work is limited.

We could not perform analyses for the gamma oscillations since in the majority of

participants the gamma peak was not detected above the aperiodic component. Due to the 1/f

properties of the EEG spectrum, both aperiodic and periodic components should be

disentangled to analyze 'true' gamma oscillations; however, this approach is not typically

recognized in electrophysiology research (Hudson and Jones, 2022). Indeed, previous studies

that analyzed gamma activity in dyslexia (Babiloni et al., 2012; Lasnick et al., 2023; Rufener

and Zaehle, 2021) did not separate the background aperiodic activity. For the same reason, we

could not analyze results for the theta band, which often does not meet the criteria for an

oscillatory component manifested as a peak in the power spectrum (Klimesch, 1999).

Moreover, results from a study investigating developmental changes in both periodic and

aperiodic components suggest that theta oscillations in older participants are mostly observed

in frontal midline electrodes (Cellier et al., 2021), which were not analyzed in the current

study.

Remarkably, in some models, results from Bayesian and frequentist statistics yielded

divergent conclusions regarding the inclusion of non-significant effects. This was observed in

more complex ANOVA models, whereas no such discrepancies appeared in *t*-tests or

correlations. Given reports of high variability in Bayesian ANOVA estimates across repeated

runs of the same analysis (Pfister, 2021), these results should be interpreted with caution.

Therefore, following the recommendation to simplify complex models into Bayesian *t*-tests

for more reliable estimates (Pfister, 2021), we conducted follow-up Bayesian *t*-tests in every

case that favored the inclusion of non-significant interactions with the group factor. These

analyses provided further evidence for the lack of differences between the dyslexic and

control groups. Another source of discrepancy between the two methods may stem from the

inclusion of interactions between covariates and within-subject effects in frequentist ANOVA,

which were not included in Bayesian ANOVA to adhere to the recommendation for simpler

Bayesian models (Pfister, 2021).

In terms of neurometabolite concentrations derived from the MRS, we found no

evidence for group differences in either Glu, GABA+ or Glu/GABA+ imbalance in the

language-sensitive left STS. Conversely, the Bayes Factor suggested against including the

group factor in the model for the Glu/GABA+ ratio. While no previous study has localized the

MRS voxel based on the individual activation levels, nonsignificant group differences in Glu

and GABA concentrations within the temporo-parietal and visual cortices have been reported

in both children and adults (Kossowski et al., 2019), as well as in the ACC in children

(Horowitz-Kraus et al., 2018). Although our MRS sample size was half that of the EEG

sample, previous research reporting group differences in Glu concentrations involved an even

smaller dyslexic cohort (10 participants with dyslexia and 45 typical readers in Pugh et al.,

2014). Consistent with earlier studies that identified group differences in Glu and GABA

concentrations (Del Tufo et al., 2018; Pugh et al., 2014) we reported neurometabolite levels

relative to total creatine (tCr) with gray matter volume included as a covariate in the models,

indicating that the absence of corresponding results cannot be ascribed to reference

differences. Notably, our analysis of the fMRI localizer task revealed greater activation in the

control group as compared to the dyslexic group within the left STS for words than control

stimuli (see Figure 2C and the Supplementary Material) in line with previous observations

(Blau et al., 2009; Dębska et al., 2021; Yan et al., 2021).

We chose ultra-high field MRS to improve data quality (Özütemiz et al., 2023) as the

increased sensitivity and spectral resolution at 7T allows for better separation of overlapping

metabolites compared to lower field strengths. Additionally, 7T provides a higher signal-to-

noise ratio (SNR), improving the reliability of metabolite measurements and enabling the

detection of small changes in Glu and GABA concentrations. Despite the theoretical

advantages several practical obstacles should be considered, such as susceptibility artifacts

and inhomogeneities at higher field strengths that can impact data quality. Interestingly, actual

methodological comparisons (Pradhan et al., 2015; Terpstra et al., 2016) show only a slight

practical advantage of 7T single-voxel MRS when compared to optimized 3T acquisition. For

example, fitting quality yielded reduced estimates of variance in concentration of Glu in 7T

(CRLB) and slightly improved reproducibility levels for Glu and GABA (at both fields below

5%). Choosing the appropriate MRS sequence involves a trade-off between the accuracy of

Glu and GABA measurements, as different sequences are recommended for each metabolite.

J-edited MRS is recommended for measuring GABA, particularly with 3T scanners (Mullins

et al., 2014). However, at 7T, more reliable results can be obtained using moderate echo-time,

non-edited MRS (Finkelman et al., 2022). We have opted for a short-echo time sequence (28

ms), which is optimal for measuring Glu. However, this approach results in macromolecule

contamination of the GABA signal (referred to as GABA+). Moreover, as Finkelman et al.

(2022) demonstrated that TE of 80 ms was superior to a shorter echo time of 42 ms for

resolving Glu and GABA at 7T, the data reported in the present study are likely influenced by

the difficulty in resolving overlapping Glu and GABA peaks at 2.3 ppm, which could impact

our measures of the Glu/GABA+ ratio and imbalance.

Irrespective of dyslexia status, we found negative correlations between age and

exponent and offset, consistent with previous research (Cellier et al., 2021; McSweeney et al.,

2021; Schaworonkow and Voytek, 2021; Voytek et al., 2015) and providing further evidence

for maturational changes in the aperiodic components (indicative of increased E/I ratio). At

the same time, in line with previous MRS works (Kossowski et al., 2019; Marsman et al.,

2013), we observed a negative correlation between age and Glu concentrations. This suggests

a contrasting pattern to EEG results, indicating a decrease in neuronal excitation with age. We

also found a condition-dependent change in offset, with a lower offset observed at rest than

during the language task. The offset value represents the uniform shift in power across

frequencies (Donoghue et al., 2020), with a higher offset linked to increased neuronal spiking

rates (Manning et al., 2009). Change in offset between conditions is consistent with observed

increased alpha and beta power during the task, indicating elevated activity in both broadband

(offset) and narrowband (alpha and beta oscillations) frequency ranges during the language

task.

In regard to relationships between EEG and MRS E/I balance biomarkers, we

observed a negative correlation between the offset in the left STS (both at rest and during the

task) and Glu levels, after controlling for age and GMV. This correlation was not observed in

zero-order correlations (see Supplementary Material). Contrary to our predictions, informed

by previous studies linking the exponent to E/I ratio (Colombo et al., 2019; Gao et al., 2017;

Waschke et al., 2021), we found the correlation with Glu levels to involve the offset rather

than the exponent. This outcome was unexpected, as none of the referenced studies reported

results for the offset. However, given the strong correlation between the exponent and offset

observed in our study ($r = .68$, $p < .001$, $BF_{10} > 10000$ and $r = .72$, $p < .001$, $BF_{10} > 10000$ at

rest and during the task respectively) it is conceivable that similar association might be

identified for the offset if it were analyzed.

Nevertheless, previous studies examining relationships between EEG and MRS E/I

balance biomarkers (McKeon et al., 2024; van Bueren et al., 2023) did not identify a similar

negative association between Glu and the offset. Instead, one study noted a positive

correlation between the Glu/GABA ratio and the exponent (van Bueren et al., 2023), which

was significant in the intraparietal sulcus but not in the middle frontal gyrus. This finding

presents counterintuitive evidence, suggesting that an increased E/I balance, as indicated by

MRS, is associated with a higher aperiodic exponent, considered indicative of decreased E/I

balance. In line with this pattern, another study discovered a positive relationship between the

exponent and Glu levels in the dorsolateral prefrontal cortex (McKeon et al., 2024).

Furthermore, they observed a positive correlation between the exponent and the Glu/GABA

imbalance measure, calculated as the absolute residual value of a linear relationship between

Glu and GABA (McKeon et al., 2024), a finding replicated in the current work. This implies

that a higher spectral exponent might not be directly linked to MRS-derived Glu or GABA

levels, but rather to a greater disproportion (in either direction) between these

neurotransmitters. These findings, alongside the contrasting relationships between EEG and

MRS biomarkers and age, suggest that these methods may reflect distinct biological

mechanisms of E/I balance.

Evidence regarding associations between neurotransmitters levels and oscillatory

activity also remains mixed. One study found a positive correlation between gamma peak

frequency and GABA concentration in the visual cortex (Muthukumaraswamy et al., 2009), a

finding later challenged by a study with a larger sample (Cousijn et al., 2014). Similarly, a

different study noted a positive correlation between GABA in the left STS and gamma power

(Balz et al., 2016), another study, found non-significant relation between these measures

(Wyss et al., 2017). Moreover, in a simultaneous EEG and MRS study, an event-related

increase in Glu following visual stimulation was found to correlate with greater gamma power

(Lally et al., 2014). We could not investigate such associations, as the algorithm failed to

identify a gamma peak above the aperiodic component for the majority of participants. Also,

contrary to previous findings showing associations between GABA in the motor and

sensorimotor cortices and beta power (Cheng et al., 2017; Gaetz et al., 2011) or beta peak

frequency (Baumgarten et al., 2016), we observed no correlation between Glu or GABA+

levels and beta power. However, these studies placed MRS voxels in motor regions which are

typically linked to movement-related beta activity (Baker et al., 1999; Rubino et al., 2006;

Sanes and Donoghue, 1993) and did not adjust beta power for aperiodic components, making

direct comparisons with our findings limited.

Finally, we examined pathways posited by the neural noise hypothesis of dyslexia,

through which increased neural noise may impact reading: phonological awareness, lexical

access and generalization, and multisensory integration (Hancock et al., 2017). Phonological

awareness was positively correlated with the offset in the left STS at rest, and with beta power

in the left STS, both at rest and during the task. Additionally, multisensory integration showed

correlations with GABA+ and the Glu/GABA+ ratio. Since the Bayes Factor did not provide

conclusive evidence supporting either the alternative or null hypothesis, these associations

appear rather weak. Nonetheless, given the hypothesis's prediction of a causal link between

these variables, we further examined a mediation model involving beta power, phonological

awareness, and reading skills. The results suggested a positive indirect effect of beta power on

reading via phonological awareness, whereas both the direct (controlling for phonological

awareness and age) and total effects (without controlling for phonological awareness) were

not significant. This finding is noteworthy, considering that participants with dyslexia

exhibited reduced phonological awareness and reading skills, despite no observed differences

in beta power. Given the cross-sectional nature of our study, further longitudinal research is

necessary to confirm the causal relation among these variables. The effects of GABA+ and

the Glu/GABA+ ratio on reading, mediated by multisensory integration, warrant further

investigation. Additionally, considering our finding that only males with dyslexia showed

deficits in multisensory integration (Glica et al., 2024), sex should be considered as a potential

moderating factor in future analyses. We did not test this model here due to the smaller

sample size for GABA+ measurements.

Our findings suggest that the neural noise hypothesis, as proposed by Hancock and

colleagues (2017), does not fully explain the reading difficulties observed in dyslexia. Despite

the innovative use of both EEG and MRS biomarkers to assess excitatory-inhibitory (E/I)

balance, neither method provided evidence supporting an E/I imbalance in dyslexic

individuals. Importantly, our study focused on adolescents and young adults, and the EEG

recordings were conducted during rest and a spoken language task. These factors may limit

the generalizability of our results. Future research should include younger populations and

incorporate a broader array of tasks, such as reading and phonological processing, to provide

a more comprehensive evaluation of the E/I balance hypothesis. Moreover, since the MRS

data was collected only from the left STS, it is plausible that other areas might be associated

with differences in Glu or GABA concentrations in dyslexia. Nevertheless, the neural noise

hypothesis predicted increased glutamatergic signaling in perisylvian regions, specifically in

the left superior temporal cortex (Hancock et al., 2017). Furthermore, although our results do

not support the idea of E/I balance alterations as a source of neural noise in dyslexia, they do

not preclude other mechanisms leading to less synchronous neural firing posited by the

hypothesis. In this context, there is evidence showing increased trial-to-trial inconsistency of

neural responses in individuals with dyslexia (Centanni et al., 2022) or poor readers

(Hornickel and Kraus, 2013) and its associations with specific dyslexia risk genes (Centanni

et al., 2018; Neef et al., 2017). At the same time, the observed trial-to-trial inconsistency was

either present only in a subset of participants (Centanni et al., 2018), limited to some

experimental conditions (Centanni et al., 2022), or specific brain regions – e.g., brainstem in

Hornickel and Kraus (2013), left auditory cortex in Centanni et al. (2018), or left

supramarginal gyrus in Centanni et al. (2022). Also, one study analyzing behavioral and fMRI

response patterns did not find similar evidence for increased variability in dyslexia (Tan et al.,

2022). Together, these results highlight the need to explore alternative neural mechanisms

underlying dyslexia and suggest that cortical hyperexcitability may not be the primary cause

of reading difficulties.

In conclusion, while our study challenges the neural noise hypothesis as a sole

explanatory framework for dyslexia, it also underscores the complexity of the disorder and the

necessity for multifaceted research approaches. By refining our understanding of the neural

underpinnings of dyslexia, we can better inform future studies and develop more effective

interventions for those affected by this condition.

MATERIALS AND METHODS

Participants

A total of 120 Polish participants aged between 15.09 and 24.95 years (M = 19.47, SD

= 3.06) took part in the study. This included 60 individuals with a clinical diagnosis of

dyslexia performed by the psychological and pedagogical counseling centers (28 females and

32 males) and 60 control participants without a history of reading difficulties (28 females and

32 males). Since there are no standardized diagnostic norms for dyslexia in adults in Poland,

individuals were assigned to the dyslexic group based on a past diagnosis of dyslexia. All

participants were right-handed, born at term, without any reported neurological/psychiatric

diagnosis and treatment (including ADHD), without hearing impairment, with normal or

corrected-to-normal vision, and IQ higher than 80 as assessed by the Polish version of the

Abbreviated Battery of the Stanford-Binet Intelligence Scale-Fifth Edition (SB5) (Roid et al.,

2017).

655   The study was approved by the institutional review board at the University of Warsaw,

656 Poland (reference number 2N/02/2021). All participants (or their parents in the case of

657 underaged participants) provided written informed consent and received monetary

658 remuneration for taking part in the study.

Reading and Reading-Related Tasks

662   Participants' reading skills were assessed by multiple paper-pencil tasks described in

663 detail in our previous work (Glica et al., 2024). Briefly, we evaluated words and pseudowords

664 read in one minute (Szczerbiński and Pelc-Pękała, 2013), rapid automatized naming (Fecenec

665 et al., 2013), and reading comprehension speed. We also assessed phonological awareness by

666 a phoneme deletion task (Szczerbiński and Pelc-Pękała, 2013) and spoonerisms tasks

667 (Bogdanowicz et al., 2016), as well as orthographic awareness (Awramiuk and Krasowicz-

668 Kupis, 2013). Furthermore, we evaluated non-verbal perception speed (Ciechanowicz and

669 Stańczak, 2006) and short-term and working memory by forward and backward conditions

670 from the Digit Span subtest from the WAIS-R (Wechsler, 1981). We also assessed

671 participants' multisensory audiovisual integration by a redundant target effect task, which

672 results have been reported in our previous work (Glica et al., 2024).

673 **Electroencephalography Acquisition and Procedure**

674   EEG was recorded from 62 scalp and 2 ear electrodes using the Brain Products system

675 (actiCHamp Plus, Brain Products GmbH, Gilching, Germany). Data were recorded in

676 BrainVision Recorder Software (Vers. 1.22.0002, Brain Products GmbH, Gilching, Germany)

677 with a 500 Hz sampling rate. Electrodes were positioned in line with the extended 10-20

678 system. Electrode Cz served as an online reference, while the Fpz as a ground electrode. All

679 electrodes' impedances were kept below 10 kΩ. Participants sat in a chair with their heads on

a chin-rest in a dark, sound-attenuated, and electrically shielded room while the EEG was

recorded during both a 5-minute eyes-open resting state and the spoken language

comprehension task. The paradigm was prepared in the Presentation software (Version 20.1,

Neurobehavioral Systems, Inc., Berkeley, CA, www.neurobs.com).

During rest, participants were instructed to relax and fixate their eyes on a white cross

presented centrally on a black background. After 5 minutes, the spoken language

comprehension task automatically started. The task consisted of 3 to 5 word-long sentences

recorded in a speech synthesizer which were presented binaurally through sound-isolating

earphones. After hearing a sentence, participants were asked to indicate whether the sentence

was true or false by pressing a corresponding button. In total, there were 256 sentences – 128

true (e.g., "Plants need water") and 128 false (e.g., "Dogs can fly").

Sentences were presented in a random order in two blocks of 128 trials. At the

beginning of each trial, a white fixation cross was presented centrally on a black background

for 500 ms, then a blank screen appeared for either 500, 600, 700, or 800 ms (durations set

randomly and equiprobably) followed by an auditory sentence presentation. The length of

sentences ranged between 1.17 and 2.78 seconds and was balanced between true (M = 1.82

seconds, SD = 0.29) and false sentences (M = 1.82 seconds, SD = 0.32; $t(254)$ = -0.21, p =

.835; BF_{10} = 0.14). After a sentence presentation, a blank screen was displayed for 1000 ms

before starting the next trial. To reduce participants' fatigue, a 1-minute break between two

blocks of trials was introduced, and it took approximately 15 minutes to complete the task.

fMRI Acquisition and Procedure

MRI data were acquired using Siemens 3T Trio system with a 32-channel head coil.

Structural data were acquired using whole brain 3D T1-weighted image (MP_RAGE, TI =

1100 ms, GRAPPA parallel imaging with acceleration factor PE = 2, voxel resolution =

1mm^3, dimensions = 256x256x176). Functional data were acquired using whole-brain echo

planar imaging sequence (TE = 30ms, TR = 1410 ms, flip angle FA = 90°, FOV = 212 mm,

matrix size = 92x92, 60 axial slices 2.3mm thick, 2.3x2.3 mm in-plane resolution, multiband

acceleration factor = 3). Due to a technical issue, data from two participants were acquired

with a 12-channel coil (see Supplementary Material).

The fMRI task served as a localizer for later MRS voxel placement in language-

sensitive left STS. The task was prepared using Presentation software (Version 20.1,

Neurobehavioral Systems, Inc., Berkeley, CA, www.neurobs.com) and consisted of three

runs, each lasting 5 minutes and 9 seconds. Two runs involved the presentation of visual

stimuli, while the third run of auditory stimuli. In each run, stimuli were presented in 12

blocks, with 14 stimuli per block. In visual runs, there were four blocks from each category:

1) 3 to 4 letters-long words, 2) the same words presented as a false font string (BACS font)

(Vidal et al., 2017), and 3) strings of 3 to 4-long consonants. Similarly, in the auditory run,

there were four blocks from each category: 1) words recorded in a speech synthesizer, 2) the

same words presented backward, and 3) consonant strings recorded in a speech synthesizer.

Stimuli within each block were presented for 800 ms with a 400 ms break in between. The

duration of each block was 16.8 seconds. Between blocks, a fixation cross was displayed for 8

seconds. Participants performed a 1-back task to maintain focus. The blocks were presented in

a pseudorandom order and each block included 2 to 3 repeated stimuli.

MRS Acquisition and Procedure

The GE 7T system with a 32-channel coil was utilized. Structural data were acquired

using whole brain 3D T1-weighted image (3D-SPGR BRAVO, TI = 450ms, TE = 2.6ms, TR

= 6.6ms, flip angle = 12 deg, bandwidth = ±32.5kHz, ARC acceleration factor PE = 2, voxel

resolution = 1mm, dimensions = 256 x 256 x 180). MRS spectra with 320 averages were

acquired from the left STS using single-voxel spectroscopy semiLaser sequence (Deelchand

et al., 2021) (voxel size = 15 x 15 x 15 mm, TE = 28ms, TR = 4000ms, 4096 data points,

water suppressed using VAPOR). Eight averages with unsuppressed water as a reference were

collected.

To localize left STS, T1-weighted images from fMRI and MRS sessions were

coregistered and fMRI peak coordinates were used as a center of voxel volume for MRS.

Voxels were then adjusted to include only the brain tissue. During the acquisition, participants

took part in a simple orthographic task.

Statistical Analyses

EEG data

The continuous EEG signal was preprocessed in the EEGLAB (Delorme and Makeig,

2004). The data were filtered between 0.5 and 45 Hz (Butterworth filter, 4th order) and re-

referenced to the average of both ear electrodes. The data recorded during the break between

blocks, as well as bad channels, were manually rejected. The number of rejected channels

ranged between 0 and 4 ($M = 0.19$, $SD = 0.63$). Next, independent component analysis (ICA)

was applied. Components were automatically labeled by ICLabel (Pion-Tonachini et al.,

2019), and those classified with 50-100% source probability as eye blinks, muscle activity,

heart activity, channel noise, and line noise, or with 0-50% source probability as brain

activity, were excluded. Components labeled as "other" were visually inspected, and those

identified as eye blinks and muscle activity were also rejected. The number of rejected

components ranged between 11 and 46 ($M = 28.43$, $SD = 7.26$). Previously rejected bad

channels were interpolated using the nearest neighbor spline (Perrin et al., 1987, 1989).

The preprocessed data were divided into a 5-minute resting-state signal and a signal

recorded during a spoken language comprehension task using MNE (Gramfort, 2013) and

custom Python scripts. The signal from the task was cut up based on the event markers

indicating the beginning and end of a sentence. Only trials with correct responses given

between 0 and 1000 ms after the end of a sentence were included. The signals recorded during

every trial were further multiplied by the Tukey window with α = 0.01 in order to normalize signal amplitudes at the beginning and end of every trial. This allowed a smooth concatenation of signals recorded during task trials, resulting in a continuous signal derived only when participants were listening to the sentences.

The continuous signal from the resting state and the language task was epoched into 2-second-long segments. An automatic rejection criterion of +/-200 μV was applied to exclude epochs with excessive amplitudes. The number of epochs retained in the analysis ranged between 140–150 ($M = 149.66$, $SD = 1.20$) in the resting state condition and between 102–226 ($M = 178.24$, $SD = 28.94$) in the spoken language comprehension task.

Power spectral density (PSD) for 0.5-45 Hz in 0.5 Hz increments was calculated for every artifact-free epoch using Welch's method for 2-second-long data segments windowed with a Hamming window with no overlap. The estimated PSDs were averaged for each participant and each channel separately for the resting state condition and the language task. Aperiodic and periodic (oscillatory) components were parameterized using the FOOOF method (Donoghue et al., 2020). For each PSD, we extracted parameters for the 1-43 Hz frequency range using the following settings: *peak_width_limits* = [1, 12], *max_n_peaks* = infinite, *peak_threshold* = 2.0, *mean_peak_height* = 0.0, *aperiodic_mode* = 'fixed'.

Two broadband aperiodic parameters were extracted: the exponent, which quantifies the steepness of the EEG power spectrum, and the offset, which indicates signal's power across the entire frequency spectrum. We also extracted aperiodic-adjusted oscillatory power, bandwidth, and the center frequency parameters for the theta (4-7 Hz), alpha (7-14 Hz), beta (14-30 Hz) and gamma (30-43 Hz) bands. Since in the majority of participants, the algorithm did not find the peak above the aperiodic component in theta and gamma bands, we calculated the results only for the alpha and beta bands. The results for other periodic parameters than the beta power are reported in Supplementary Material.

First, exponent and offset values were averaged across all electrodes and analyzed

using a 2x2 repeated measures ANOVA with group (dyslexic, control) as a between-subjects

factor and condition (resting state, language task) as a within-subjects factor. Age was

included in the analyses as a covariate due to correlation between variables.

Next, exponent and offset values were averaged across electrodes corresponding to the

left (F7, FT7, FC5) and right inferior frontal gyrus (F8, FT8, FC6), and to the left (T7, TP7,

TP9) and right superior temporal sulcus (T8, TP8, TP10). The electrodes were selected based

on the analyses outlined by Giacometti and colleagues (2014) and Scrivener and Reader

(2022). For these analyses, a 2x2x2x2 repeated measures ANOVA with age as a covariate

was conducted with group (dyslexic, control) as a between-subjects factor and condition

(resting state, language task), hemisphere (left, right), and region (frontal, temporal) as within-

subjects factors.

Results for the alpha and beta bands were calculated for the same clusters of frontal

and temporal electrodes and analyzed with a similar 2x2x2x2 repeated measures ANOVA;

however, for these analyses, age was not included as a covariate due to a lack of significant

correlations.

Apart from the frequentist statistics, we also performed Bayesian statistics using JASP

(JASP Team, 2023). For Bayesian repeated measures ANOVA, we reported the Bayes Factor

for the inclusion of a given effect (BF_{incl}) with the 'across matched model' option, as suggested

by Keysers and colleagues (2020), calculated as a likelihood ratio of models with a presence

of a specific factor to equivalent models differing only in the absence of the specific factor.

For Bayesian t-tests and correlations, we reported the BF_{10} value, indicating the ratio of the

likelihood of an alternative hypothesis to a null hypothesis. We considered $BF_{incl/10} > 3$ and

$BF_{incl/10} < 1/3$ as evidence for alternative and null hypotheses respectively, while $1/3 < BF_{incl/10}$

< 3 as the absence of evidence (Keysers et al., 2020).

fMRI data

MRS voxel localization in the native space

The data were analyzed using Statistical Parametric Mapping (SPM12, Wellcome

Trust Centre for Neuroimaging, London, UK) run on MATLAB R2020b (The MathWorks

Inc., Natick, MA, USA). First, all functional images were realigned to the participant's mean.

Then, T1-weighted images were coregistered to functional images for each subject. Finally,

fMRI data were smoothed with a 6mm isotropic Gaussian kernel.

In each subject, the left STS was localized in the native space as a cluster in the middle

and posterior left superior temporal sulcus, exhibiting higher activation for visual words

versus false font strings and auditory words versus backward words (logical AND

conjunction) at $p < .01$ uncorrected. For 6 participants (DYS $n = 2$, CON $n = 4$), the threshold

was lowered to $p < .05$ uncorrected, while for another 6 participants (DYS $n = 3$, CON $n = 3$)

the contrast from the auditory run was changed to auditory words versus fixation cross due to

a lack of activation for other contrasts.

In the Supplementary Material, we also performed the group-level analysis of the

fMRI data (Tables S5-S7 and Figure S1).

MRS data

MRS data were analyzed using fsl-mrs version 2.0.7 (Clarke et al., 2021). Data stored

in pfile format were converted into NIfTI-MRS using spec2nii tool. We then used the

fsl_mrs_preproc function to automatically perform coil combination, frequency and phase

alignment, bad average removal, combination of spectra, eddy current correction, shifting

frequency to reference peak and phase correction.

To obtain information about the percentage of WM, GM and CSF in the voxel we used

the *svs_segmentation* with results of *fsl_anat* as an input. Voxel segmentation was performed

on structural images from a 3T scanner, coregistered to 7T structural images in SPM12, as the

latter exhibited excessive artifacts and intensity bias in the temporal regions. Next,

quantitative fitting was performed using *fsl_mrs* function. As a basis set, we utilized a

collection of 27 metabolite spectra simulated using FID-A (Simpson et al., 2017) and a script

tailored for our experiment. We supplemented this with synthetic macromolecule spectra

provided by *fsl_mrs*. In contrast to symmetrical editing technique (Edden et al., 2012), we

were unable to fully separate GABA peaks from the spectra of macromolecules. Therefore,

we use 'GABA+' as an abbreviation to indicate the potential contributions of other

compounds. Signals acquired with unsuppressed water served as water reference.

Spectra underwent quantitative assessment and visual inspection and those with

linewidth higher than 20Hz, %CRLB higher than 20%, and poor fit to the model were

excluded from the analysis (see Table S8 in the Supplementary Material for a detailed

checklist). Glu and GABA+ concentrations were expressed as a ratio to total-creatine (tCr;

Creatine + Phosphocreatine) following previous MRS studies in dyslexia (Del Tufo et al.,

2018; Pugh et al., 2014).

To analyze the metabolite results, separate univariate ANCOVAs were conducted for

Glu, GABA+, Glu/GABA+ ratio and Glu/GABA+ imbalance measures with group (control,

dyslexic) as a between-subjects factor and voxel gray matter volume (GMV) as a covariate.

Additionally, for the Glu analysis, age was included as a covariate due to a correlation

between variables. Both frequentist and Bayesian statistics were calculated. Glu/GABA+

imbalance measure was calculated as the square root of the absolute residual value of a linear

relationship between Glu and GABA+ (McKeon et al., 2024).

Data Availability Statement

Behavioral data, raw and preprocessed EEG data, 2^{nd} level fMRI data, preprocessed

MRS data and Python script for the analysis of preprocessed EEG data can be found at OSF:

https://osf.io/4e7ps/

Acknowledgments

This study was supported by the National Science Centre grant
(2019/35/B/HS6/01763) awarded to Katarzyna Jednoróg.

We gratefully acknowledge valuable discussions with Ralph Noeske from GE
Healthcare for his support in setting up the protocol for an ultra-high field MR spectroscopy
and sharing the set-up for basis set simulation in FID-A.

REFERENCES

Ahmad J, Ellis C, Leech R, Voytek B, Garces P, Jones E, Buitelaar J, Loth E, dos Santos
        FP, Amil AF, Verschure PFMJ, Murphy D, McAlonan G. (2022) **From
        mechanisms to markers: Novel noninvasive EEG proxy markers of the neural
        excitation and inhibition system in humans** *Transl Psychiatry* **12**:467.
        doi:10.1038/s41398-022-02218-z

Awramiuk E, Krasowicz-Kupis G. (2014) **Reading and spelling acquisition in Polish:
        Educational and linguistic determinants** *L1-Educ Stud Lang Lit* **14**:1–24.
        doi:10.17239/L1ESLL-2014.01.13

Babiloni C, Stella G, Buffo P, Vecchio F, Onorati P, Muratori C, Miano S, Gheller F,
        Antonaci L, Albertini G, Rossini PM. (2012) **Cortical sources of resting state
        EEG rhythms are abnormal in dyslexic children** *Clin Neurophysiol* **123**:2384–
        2391. doi:10.1016/j.clinph.2012.05.002

Baker SN, Kilner JM, Pinches EM, Lemon RN. (1999) **The role of synchrony and
        oscillations in the motor output** *Exp Brain Res* **128**:109–117.
        doi:10.1007/s002210050825

Balz J, Keil J, Roa Romero Y, Mekle R, Schubert F, Aydin S, Ittermann B, Gallinat J,
        Senkowski D. (2016) **GABA concentration in superior temporal sulcus predicts**

gamma power and perception in the sound-induced flash illusion *Neuroimage* **125**:724–730. doi:10.1016/j.neuroimage.2015.10.087

Baumgarten TJ, Oeltzschner G, Hoogenboom N, Wittsack H-J, Schnitzler A, Lange J. (2016) **Beta peak frequencies at rest correlate with endogenous GABA+/Cr concentrations in sensorimotor cortex areas** *PLoS One* **11**:e0156829. doi:10.1371/journal.pone.0156829

Blau V, van Atteveldt N, Ekkebus M, Goebel R, Blomert L. (2009) **Reduced neural integration of letters and speech sounds links phonological and reading deficits in adult dyslexia** *Curr Biol* **19**:503–508. doi:10.1016/j.cub.2009.01.065

Bogdanowicz KM, Wiejak K, Krasowicz-Kupis G, Gawron N. (2015) **Ocena przydatności kwestionariusza Adult Reading History Questionnaire do diagnozy rodzinnego ryzyka dysleksji w Polsce. [Evaluation of the usefulness of The Adult Reading History Questionnaire for identification of familial risk of dyslexia in Poland]**. *Edukacja* **1**:117–138.

Bogdanowicz M, Sajewicz-Radtke U, Radtke BM, Kalka D, Karpińska E, Bogdanowicz KM, Łockiewicz M. (2016) **Bateria metod diagnozy przyczyn niepowodzeń szkolnych u uczniów powyżej 16. roku życia. BATERIA-16 plus [B-16]. [A battery of methods for diagnosing the causes of school failure in students over 16 years of age. BATTERY-16 plus [B-16]]**. Gdańsk, Poland: Pracownia Testów Psychologicznych i Pedagogicznych [Laboratory of Psychological and Pedagogical Tests].

Brunel N, Wang X-J. (2003) **What determines the frequency of fast network oscillations with irregular neural discharges? I. Synaptic dynamics and excitation-inhibition balance** *J Neurophysiol* **90**:415–430. doi:10.1152/jn.01095.2002

Cellier D, Riddle J, Petersen I, Hwang K. (2021) **The development of theta and alpha neural oscillations from ages 3 to 24 years** *Dev Cogn Neurosci* **50**:100969. doi:10.1016/j.dcn.2021.100969

Centanni TM, Beach SD, Ozernov-Palchik O, May S, Pantazis D, Gabrieli JDE. (2022) **Categorical perception and influence of attention on neural consistency in response to speech sounds in adults with dyslexia** *Ann Dyslexia* **72**:56–78. doi:10.1007/s11881-021-00241-1

Centanni TM, Pantazis D, Truong DT, Gruen JR, Gabrieli JDE, Hogan TP. (2018) **Increased variability of stimulus-driven cortical responses is associated with genetic variability in children with and without dyslexia** *Dev Cogn Neurosci* **34**:7–17. doi:10.1016/j.dcn.2018.05.008

Chen G, Zhang Y, Li X, Zhao X, Ye Q, Lin Y, Tao HW, Rasch MJ, Zhang X. (2017) **Distinct inhibitory circuits orchestrate cortical beta and gamma band oscillations** *Neuron* **96**:1403-1418.e6. doi:10.1016/j.neuron.2017.11.033

Cheng C-H, Tsai S-Y, Liu C-Y, Niddam DM. (2017) **Automatic inhibitory function in the human somatosensory and motor cortices: An MEG-MRS study** *Sci Rep* **7**:4234. doi:10.1038/s41598-017-04564-1

Ciechanowicz A, Stańczak J. (2006) **Testy Uwagi i Spostrzegawczości: TUS: podręcznik. [Attention and Perceptiveness Tests: TUS: handbook]**. Warszawa, Poland: Pracownia Testów Psychologicznych Polskiego Towarzystwa Psychologicznego [Laboratory of Psychological Tests of the Polish Psychological Association].

Clarke WT, Stagg CJ, Jbabdi S. (2021) **FSL☐MRS: An end☐to☐end spectroscopy analysis package** *Magn Reson Med* **85**:2950–2964. doi:10.1002/mrm.28630

Colombo MA, Napolitani M, Boly M, Gosseries O, Casarotto S, Rosanova M, Brichant J-F, Boveroux P, Rex S, Laureys S, Massimini M, Chieregato A, Sarasso S. (2019) **The spectral exponent of the resting EEG indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and ketamine** *Neuroimage* **189**:631–644. doi:10.1016/j.neuroimage.2019.01.024

Cousijn H, Haegens S, Wallis G, Near J, Stokes MG, Harrison PJ, Nobre AC. (2014) **Resting GABA and glutamate concentrations do not predict visual gamma frequency or amplitude**. *Proc Natl Acad Sci* **111**:9301–9306. doi:10.1073/pnas.1321072111

Dębska A, Banfi C, Chyl K, Dzięgiel-Fivet G, Kacprzak A, Łuniewska M, Plewko J, Grabowska A, Landerl K, Jednoróg K. (2021) **Neural patterns of word processing differ in children with dyslexia and isolated spelling deficit** *Brain Struct Funct* **226**:1467–1478. doi:10.1007/s00429-021-02255-2

Deelchand DK, Berrington A, Noeske R, Joers JM, Arani A, Gillen J, Schär M, Nielsen J, Peltier S, Seraji☐Bozorgzad N, Landheer K, Juchem C, Soher BJ, Noll DC, Kantarci K, Ratai EM, Mareci TH, Barker PB, Öz G. (2021) **Across☐vendor standardization of semi☐LASER for single☐voxel MRS at 3T** *NMR Biomed* **34**:e4218. doi:10.1002/nbm.4218

Del Tufo SN, Frost SJ, Hoeft F, Cutting LE, Molfese PJ, Mason GF, Rothman DL, Fulbright RK, Pugh KR. (2018) **Neurochemistry predicts convergence of written and spoken language: A proton magnetic resonance spectroscopy study of cross-modal language integration** *Front Psychol* **9**:1507. doi:10.3389/fpsyg.2018.01507

Delorme A, Makeig S. (2004**) EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis** *J Neurosci Methods* **134**:9–21. doi:10.1016/j.jneumeth.2003.10.009

Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, Noto T, Lara AH, Wallis JD, Knight RT, Shestyuk A, Voytek B. (2020) **Parameterizing neural power spectra into periodic and aperiodic components** *Nat Neurosci* **23**:1655–1665. doi:10.1038/s41593-020-00744-x

Edden RA, Puts NA, Barker PB. (2012) **Macromolecule-suppressed GABA-edited magnetic resonance spectroscopy at 3T** *Magn Reson Med* **68**:657–661. doi:10.1002/mrm.24391

Fecenec D, Jaworowska A, Matczak A, Stańczak J, Zalewska E. (2013) **Test szybkiego nazywania (TSN). [Rapid Automatized Naming Task]**. Warszawa, Poland: Pracownia Testów Psychologicznych Polskiego Towarzystwa Psychologicznego [Laboratory of Psychological Tests of the Polish Psychological Association].

Fein G, Galin D, Yingling CD, Johnstone J, Davenport L, Herron J. (1986) **EEG spectra in dyslexic and control boys during resting conditions** *Electroencephalogr Clin Neurophysiol* **63**:87–97. doi:10.1016/0013-4694(86)90001-5

Finkelman T, Furman-Haran E, Paz R, Tal A. (2022) **Quantifying the excitatory-inhibitory balance: A comparison of SemiLASER and MEGA-SemiLASER for simultaneously measuring GABA and glutamate at 7T** *Neuroimage* **247**:118810. doi:10.1016/j.neuroimage.2021.118810

Fraga González G, Smit DJA, van der Molen MJW, Tijms J, Stam CJ, de Geus EJC, van der Molen MW. (2018) **EEG resting state functional connectivity in adult dyslexics using phase lag index and graph analysis** *Front Hum Neurosci* **12**:341. doi:10.3389/fnhum.2018.00341

Gaetz W, Edgar JC, Wang DJ, Roberts TPL. (2011) **Relating MEG measured motor cortical oscillations to resting γ-Aminobutyric acid (GABA) concentration** *Neuroimage* **55**:616–621. doi:10.1016/j.neuroimage.2010.12.077

Gao R, Peterson EJ, Voytek B. (2017) **Inferring synaptic excitation/inhibition balance from field potentials** *Neuroimage* **158**:70–78. doi:10.1016/j.neuroimage.2017.06.078

Germanò E, Gagliano A, Curatolo P. (2010) **Comorbidity of ADHD and dyslexia** *Dev Neuropsychol* **35**:475–493. doi:10.1080/87565641.2010.494748

Giacometti P, Perdue KL, Diamond SG. (2014) **Algorithm to find high density EEG scalp coordinates and analysis of their correspondence to structural and functional regions of the brain** *J Neurosci Methods* **229**:84–96. doi:10.1016/j.jneumeth.2014.04.020

Glica A, Wasilewska K, Kossowski B, Żygierewicz J, Jednoróg K. (2024) **Sex differences in low-level multisensory integration in developmental dyslexia** *J Neurosci* **44**:e0944232023. doi:10.1523/JNEUROSCI.0944-23.2023

Gramfort A. (2013) **MEG and EEG data analysis with MNE-Python** *Front Neurosci* **7**:267–267. doi:10.3389/fnins.2013.00267

Hancock R, Pugh KR, Hoeft F. (2017) **Neural noise hypothesis of developmental dyslexia** *Trends Cogn Sci* **21**:434–448. doi:10.1016/j.tics.2017.03.008

Hayes AF. (2017) **Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-Based Approach. Third Edition**. Guilford publications.

Hornickel J, Kraus N. (2013) **Unstable representation of sound: A biological marker of dyslexia** *J Neurosci* **33**:3500–3504. doi:10.1523/JNEUROSCI.4205-12.2013

Horowitz-Kraus T, Brunst KJ, Cecil KM. (2018). **Children with dyslexia and typical readers: Sex-based choline differences revealed using proton magnetic**

resonance spectroscopy acquired within anterior cingulate cortex *Front Hum Neurosci* **12**:466. doi:10.3389/fnhum.2018.00466

Hudson MR, Jones NC. (2022) **Deciphering the code: Identifying true gamma neural oscillations** *Exp Neurol* **357:**114205. doi:10.1016/j.expneurol.2022.114205

JASP Team. (2023) **JASP (Version 0.18.1)** [Computer software]

Jensen O, Goel P, Kopell N, Pohja M, Hari R, Ermentrout B. (2005) **On the human sensorimotor-cortex beta rhythm: Sources and modeling** *Neuroimage* **26**:347–355. doi:10.1016/j.neuroimage.2005.02.008

Keysers C, Gazzola V, Wagenmakers E-J. (2020) **Using Bayes factor hypothesis testing in neuroscience to establish evidence of absence** *Nat Neurosci* **23**:788–799. doi:10.1038/s41593-020-0660-4

Klimesch W. (1999) **EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis** *Brain Res Rev* **29:**169-195. doi:10.1016/s0165-0173(98)00056-3

Kossowski B, Chyl K, Kacprzak A, Bogorodzki P, Jednoróg K. (2019) **Dyslexia and age related effects in the neurometabolites concentration in the visual and temporo-parietal cortex** *Sci Rep* **9**:5096. doi:10.1038/s41598-019-41473-x

Lally N, Mullins PG, Roberts M V., Price D, Gruber T, Haenschel C. (2014) **Glutamatergic correlates of gamma-band oscillatory activity during cognition: A concurrent ER-MRS and EEG study** *Neuroimage* **85**:823–833. doi:10.1016/j.neuroimage.2013.07.049

Langer N, Benjamin C, Becker BLC, Gaab N. (2019) **Comorbidity of reading disabilities and ADHD: Structural and functional brain characteristics** *Hum Brain Mapp* **40**:2677–2698. doi:10.1002/hbm.24552

Lasnick OHM, Hancock R, Hoeft F. (2023) **Left-dominance for resting-state temporal low-gamma power in children with impaired word-decoding and without comorbid ADHD** *PLoS One* **18**:e0292330. doi:10.1371/journal.pone.0292330

Lebel C, MacMaster FP, Dewey D. (2016) **Brain metabolite levels and language abilities in preschool children** *Brain Behav* **6**:e00547. doi:10.1002/brb3.547

Lehongre K, Ramus F, Villiermet N, Schwartz D, Giraud A-L. (2011) **Altered low-gamma sampling in auditory cortex accounts for the three main facets of dyslexia** *Neuron* **72**:1080–1090. doi:10.1016/j.neuron.2011.11.002

Manning JR, Jacobs J, Fried I, Kahana MJ. (2009) **Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans** *J Neurosci* **29**:13613–13620. doi:10.1523/JNEUROSCI.2041-09.2009

Manyukhina VO, Prokofyev AO, Galuta IA, Goiaeva DE, Obukhova TS, Schneiderman JF, Altukhov DI, Stroganova TA, Orekhova E V. (2022) **Globally elevated excitation–inhibition ratio in children with autism spectrum disorder and below-average intelligence** *Mol Autism* **13**:20. doi:10.1186/s13229-022-00498-2

Marchesotti S, Nicolle J, Merlet I, Arnal LH, Donoghue JP, Giraud A-L. (2020**) Selective enhancement of low-gamma activity by tACS improves phonemic processing and reading accuracy in dyslexia** *PLoS Biol* **18**:e3000833. doi:10.1371/journal.pbio.3000833

Marsman A, Mandl RCW, van den Heuvel MP, Boer VO, Wijnen JP, Klomp DWJ, Luijten PR, Hilleke E. HP. (2013) **Glutamate changes in healthy young adulthood** *Eur Neuropsychopharmacol* **23**:1484–1490. doi:10.1016/j.euroneuro.2012.11.003

McKeon SD, Perica MI, Parr AC, Calabro FJ, Foran W, Hetherington H, Moon C-H, Luna B. (2024) **Aperiodic EEG and 7T MRSI evidence for maturation of E/I balance**

supporting the development of working memory through adolescence *Dev Cogn Neurosci* **66**:101373. doi:10.1016/j.dcn.2024.101373

McSweeney M, Morales S, Valadez EA, Buzzell GA, Fox NA. (2021) **Longitudinal age- and sex-related change in background aperiodic activity during early adolescence** *Dev Cogn Neurosci* **52**:101035. doi:10.1016/j.dcn.2021.101035

Molina JL, Voytek B, Thomas ML, Joshi YB, Bhakta SG, Talledo JA, Swerdlow NR, Light GA. (2020) **Memantine effects on electroencephalographic measures of putative excitatory/inhibitory balance in schizophrenia** *Biol Psychiatry Cogn Neurosci Neuroimaging* **5**:562–568. doi:10.1016/j.bpsc.2020.02.004

Mullins PG, McGonigle DJ, O'Gorman RL, Puts NA, Vidyasagar R, Evans CJ, Cardiff Symposium on MRS of GABA, Edden RA. (2014) **Current practice in the use of MEGA-PRESS spectroscopy for the detection of GABA** *Neuroimage* **86**:43–52. doi:10.1016/j.neuroimage.2012.12.004

Muthukumaraswamy SD, Edden RAE, Jones DK, Swettenham JB, Singh KD. (2009) **Resting GABA concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation in humans** *Proc Natl Acad Sci* **106**:8356–8361. doi:10.1073/pnas.0900728106

Nakai T, Okanoya K. (2016) **Individual variability in verbal fluency correlates with γ-aminobutyric acid concentration in the left inferior frontal gyrus** *Neuroreport* **27**:987–991. doi:10.1097/WNR.0000000000000645

Neef NE, Müller B, Liebig J, Schaadt G, Grigutsch M, Gunter TC, Wilcke A, Kirsten H, Skeide MA, Kraft I, Kraus N, Emmrich F, Brauer J, Boltze J, Friederici AD. (2017) **Dyslexia risk gene relates to representation of sound in the auditory brainstem** *Dev Cogn Neurosci* **24**:63–71. doi:10.1016/j.dcn.2017.01.008

Ostlund BD, Alperin BR, Drew T, Karalunas SL. (2021) **Behavioral and cognitive correlates of the aperiodic (1/f-like) exponent of the EEG power spectrum in adolescents with and without ADHD** *Dev Cogn Neurosci* **48**:100931. doi:10.1016/j.dcn.2021.100931

Özütemiz C, White M, Elvendahl W, Eryaman Y, Marjańska M, Metzger GJ, Patriat R, Kulesa J, Harel N, Watanabe Y, Grant A, Genovese G, Cayci Z. (2023) **Use of a commercial 7-T MRI scanner for clinical brain imaging: Indications, protocols, challenges, and solutions – A single-center experience** *AJR Am J Roentgenol* **221**: 788–804. doi:10.2214/AJR.23.29342

Perrin F, Pernier J, Bertnard O, Giard MH, Echallier JF. (1987) **Mapping of scalp potentials by surface spline interpolation** *Electroencephalogr Clin Neurophysiol* **66**:75–81. doi:10.1016/0013-4694(87)90141-6

Perrin F, Pernier J, Bertrand O, Echallier JF. (1989) **Spherical splines for scalp potential and current density mapping** *Electroencephalogr Clin Neurophysiol* **72**:184–187. doi:10.1016/0013-4694(89)90180-6

Pfister R. (2021) **Variability of Bayes Factor estimates in Bayesian analysis of variance** *Quant Meth Psychol* **17**:40–45. doi:10.20982/tqmp.17.1.p040

Pion-Tonachini L, Kreutz-Delgado K, Makeig S. (2019) **ICLabel: An automated electroencephalographic independent component classifier, dataset, and website** *Neuroimage* **198**:181–197. doi:10.1016/j.neuroimage.2019.05.026

Porjesz B, Almasy L, Edenberg HJ, Wang K, Chorlian DB, Foroud T, Goate A, Rice JP, O'Connor SJ, Rohrbaugh J, Kuperman S, Bauer LO, Crowe RR, Schuckit MA, Hesselbrock V, Conneally PM, Tischfield JA, Li T-K, Reich T, Begleiter H. (2002) **Linkage disequilibrium between the beta frequency of the human EEG and a**

GABAA receptor gene locus *Proc Natl Acad Sci* **99**:3729–3733. doi:10.1073/pnas.052716399

Pradhan S, Bonekamp S, Gillen JS, Rowland LM, Wijtenburg SA, Edden RA, Barker PB. (2015) **Comparison of single voxel brain MRS AT 3T and 7T using 32-channel head coils** *Magn Reson Imaging* **33**:1013–1018. doi:10.1016/j.mri.2015.06.003

Pugh KR, Frost SJ, Rothman DL, Hoeft F, Del Tufo SN, Mason GF, Molfese PJ, Mencl WE, Grigorenko EL, Landi N, Preston JL, Jacobsen L, Seidenberg MS, Fulbright RK. (2014) **Glutamate and choline levels predict individual differences in reading ability in emergent readers** *J Neurosci* **34**:4082–4089. doi:10.1523/JNEUROSCI.3907-13.2014

Rippon G, Brunswick N. (2000) **Trait and state EEG indices of information processing in developmental dyslexia** *Int J Psychophysiol* **36**:251–265. doi:10.1016/S0167-8760(00)00075-1

Roid GH, Sajewicz-Radtke U, Radtke BM, Lipowska M. (2017) **Skala Inteligencji Stanford-Binet, Edycja Piąta. [Stanford-Binet Intelligence Scales, Fifth Edition]**. Gdańsk, Poland: Pracownia Testów Psychologicznych i Pedagogicznych [Laboratory of Psychological and Pedagogical Tests].

Rubino D, Robbins KA, Hatsopoulos NG. (2006) **Propagating waves mediate information transfer in the motor cortex** *Nat Neurosci* **9**:1549–1557. doi:10.1038/nn1802

Rufener KS, Zaehle T. (2021) **Dysfunctional auditory gamma oscillations in developmental dyslexia: A potential target for a tACS-based intervention** *Prog Brain Res* **264**:211-232. doi:10.1016/bs.pbr.2021.01.016

Salvatore S V., Lambert PM, Benz A, Rensing NR, Wong M, Zorumski CF, Mennerick S. (2024) **Periodic and aperiodic changes to cortical EEG in response to**

pharmacological manipulation *J Neurophysiol* **131**:529–540. doi:10.1152/jn.00445.2023

Sanes JN, Donoghue JP. (1993) **Oscillations in local field potentials of the primate motor cortex during voluntary movement** *Proc Natl Acad Sci* **90**:4470–4474. doi:10.1073/pnas.90.10.4470

Schaworonkow N, Voytek B. (2021) **Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life** *Dev Cogn Neurosci* **47**:100895. doi:10.1016/j.dcn.2020.100895

Scrivener CL, Reader AT. (2022) **Variability of EEG electrode positions and their underlying brain regions: Visualizing gel artifacts from a simultaneous EEG☐fMRI dataset** *Brain Behav* **12**:e2476. doi:10.1002/brb3.2476

Simpson R, Devenyi GA, Jezzard P, Hennessy TJ, Near J. (2017) **Advanced processing and simulation of MRS data using the FID appliance (FID☐A) – an open source, MATLAB☐based toolkit** *Magn Reson Med* **77**:23–33. doi:10.1002/mrm.26091

Spironelli C, Penolazzi B, Angrilli A. (2008) **Dysfunctional hemispheric asymmetry of theta and beta EEG activity during linguistic tasks in developmental dyslexia** *Biol Psychol* **77**:123–131. doi:10.1016/j.biopsycho.2007.09.009

Szczerbiński M, Pelc-Pękała O. (2013) **Zestaw metod do diagnozy trudności w czytaniu. [A Set of Tools for Diagnosing Reading Difficulties]**. Gdańsk, Poland: Pracownia Testów Psychologicznych i Pedagogicznych [Laboratory of Psychological and Pedagogical Tests].

Tan Y, Chanoine V, Cavalli E, Anton J-L, Ziegler JC. (2022) **Is there evidence for a noisy computation deficit in developmental dyslexia?** *Front Hum Neurosci* **16**:919465. doi:10.3389/fnhum.2022.919465

Terpstra M, Cheong I, Lyu T, Deelchand DK, Emir UE, Bednařík P, Eberly LE, Öz G. (2016) **Test-retest reproducibility of neurochemical profiles with short-echo, single-voxel MR spectroscopy at 3T and 7T** *Magn Reson Med* **76**:1083–1091. doi:10.1002/mrm.26022

Turri C, Di Dona G, Santoni A, Zamfira DA, Franchin L, Melcher D, Ronconi L. (2023) **Periodic and aperiodic EEG features as potential markers of developmental dyslexia** *Biomedicines* **11**:1607. doi:10.3390/biomedicines11061607

van Bueren NER, van der Ven SHG, Hochman S, Sella F, Cohen Kadosh R. (2023) **Human neuronal excitation/inhibition balance explains and predicts neurostimulation induced learning benefits** *PLoS Biol* **21**:e3002193. doi:10.1371/journal.pbio.3002193

Van Hirtum T, Ghesquière P, Wouters J. (2019) **Atypical neural processing of rise time by adults with dyslexia** *Cortex* **113**:128–140. doi:10.1016/j.cortex.2018.12.006

Vidal C, Content A, Chetail F. (2017) **BACS: The Brussels Artificial Character Sets for studies in cognitive psychology and neuroscience** *Behav Res Methods* **49**:2093–2112. doi:10.3758/s13428-016-0844-8

Voytek B, Kramer MA, Case J, Lepage KQ, Tempesta ZR, Knight RT, Gazzaley A. (2015) **Age-related changes in 1/f neural electrophysiological noise** *J Neurosci* **35**:13257–13265. doi:10.1523/JNEUROSCI.2332-14.2015

Waschke L, Donoghue T, Fiedler L, Smith S, Garrett DD, Voytek B, Obleser J. (2021) **Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent** *Elife* **10**:e70068. doi:10.7554/eLife.70068

Wechsler D. (1981) **Wechsler Adult Intelligence Scale-Revised (WAIS-R)**. Psychological Corporation.

Wyss C, Tse DHY, Kometer M, Dammers J, Achermann R, Shah NJ, Kawohl W, Neuner I. (2017) **GABA metabolism and its role in gamma□band oscillatory activity during auditory processing: An MRS and EEG study** *Hum Brain Mapp* **38**:3975–3987. doi:10.1002/hbm.23642

Xue H, Wang Z, Tan Y, Yang H, Fu W, Xue L, Zhao J. (2020) **Resting-state EEG reveals global network deficiency in dyslexic children** *Neuropsychologia* **138**:107343. doi:10.1016/j.neuropsychologia.2020.107343

Yan X, Jiang K, Li H, Wang Z, Perkins K, Cao F. (2021) **Convergent and divergent brain structural and functional abnormalities associated with developmental dyslexia** *Elife* **10**:e69523. doi:10.7554/eLife.69523