Brain-derived exosomal hemoglobin transfer contributes to neuronal mitochondrial homeostasis under hypoxia

Zhengming Tian[1], Feiyang Jin[2], Zirui Xu[1], Yakun Gu[1], Mengyuan Guo[1], Yuning Li[1], Qianqian Shao[1], Yingxia Liu[1], Hanjiang Luo[3], Yue Wang[4], Suyu Zhang[4], Chenlu Yang[4], Xin Liu[4], Xunming Ji[1,2]*, Jia Liu[1]*

1 Beijing Institute of Brain Disorders, Laboratory of Brain Disorders, Laboratory for Clinical Medicine, Chinese Institutes for Medical Research, Ministry of Science and Technology, Collaborative Innovation Center for Brain Disorders, Beijing Advanced Innovation Center for Big Data-based Precision Medicine, Capital Medical University, Beijing, China.

2 Department of Neurosurgery, Xuanwu Hospital, Capital Medical University, Beijing, China

3 Laboratory of Neuroscience, Affiliated Hospital of Guilin Medical University, Guilin, Guangxi Proveince, China

4 BGI-Beijing, Beijing, China

*Corresponding author:

Prof. Jia Liu; Beijing Institute of Brain Disorders, Capital Medical University, 10 Xi Tou Tiao, You Anmen, Beijing 100069, China.

E-mail: liujia_19901005@163.com

Prof. Xunming Ji; Beijing Institute of Brain Disorders, Capital Medical University, 10 Xi Tou Tiao, You Anmen, Beijing 100069, China.

E-mail: jixm@ccmu.edu.cn

Abstract

Hypoxia is an important physiological stress causing nerve injuries and several brain
diseases. However, the mechanism of brain response to hypoxia remains unclear, thus
limiting the development of interventional strategies. This study conducted combined
analyses of single-nucleus transcriptome sequencing and extracellular vesicle
transcriptome sequencing on hypoxic mouse brains, described cell-cell
communication in the brain under hypoxia from intercellular and extracellular
dimensions, confirmed that hemoglobin mRNA was transferred from non-neuronal
cells to neurons, and eventually expressed. Then we further explored the role of
exosomal hemoglobin transfer *in vitro*, clarified that hypoxia promoted the transfer
and expression of exosomal hemoglobin between endothelial cells and neurons. And
we found the vital function of exosomal hemoglobin to protect against neurological
injury by maintaining mitochondrial homeostasis in neurons. In conclusion, this study
identified a novel mechanism of 'mutual aid' in hypoxia responses in the brain,
involving exosomal hemoglobin transfer, clarified the important role of exosomal
communication in the process of brain stress response, and provided a novel
interventional perspective for hypoxia-related brain diseases.

Key words: Brain-derived extracellular vesicles, Hemoglobin, Hypoxia,
Mitochondrial homeostasis, Nerve injury

1   Introduction

Extracellular vesicles (EVs) are membrane-enclosed vesicles secreted by cells into the extracellular environment, and are an important means of cell communication[1-3]. EV-mediated communication can participate in the regulation of intercellular signals in physiological and pathological conditions by transferring different contents, such as bioactive lipids, non-coding RNAs, mRNAs, and proteins, and the process is closely related to the occurrence and development of various diseases[4-6]. Current research on the biological function of EVs is mostly focused on EVs in body fluids, such as plasma and saliva; however, although their acquisition is relatively simple, such EVs have low tissue specificity[7,8]. In contrast, more in-depth studies have obtained and analyzed EVs from the extracellular environment in organs with complex structures, thus improving tissue specificity[9-11]. Nevertheless, this still fails to identify the cell type from which they are derived, making it impossible to carry out more detailed analyses of directional interactions between cells. The development of single-cell/nuclear transcriptome sequencing (snRNA-seq) technology allows us to explore the heterogeneity of cells in complex organs, and the combination of snRNA-seq and tissue EVs transcriptome sequencing (EVs-RNA-seq) can help us to clarify the source of tissue EVs and analyze the complex physiological and pathological conditions of organs, using the dual advantages of intracellular and extracellular approaches[12,13].

Hypoxia is a common stress response in the body that can mediate a series of physiological reactions, including embryonic development and bone marrow hematopoiesis. Hypoxia is also an important factor in various pathologies[14-16]. As the most O2-dependent organ, the brain accounts for 20% of the resting O2 consumption in humans and is thus highly sensitive to hypoxic stress[17-19]. Hypoxia is a key cause of many neurological diseases, including Alzheimer's disease, Parkinson's disease (PD), and other age-related neurodegenerative diseases[20-22]. Although there is known to be a close relationship between hypoxic stress and neurodegeneration, the responses of and damage to brain cells in hypoxic environments remain unclear. The application of snRNA-seq combined with tissue EVs-RNA-seq analysis can be used to comprehensively analyze the interactions among various cell types in the brain under hypoxic stress, thus helping us to understand the mechanisms of neurodegenerative injury and to identify potential therapeutic molecules for interventions.

Hemoglobin is a major protein involved in the transport and utilization of O2 in the body. It is composed of globin and heme and is mainly expressed in red blood cells[23]. Previous studies found that hemoglobin also existed in the brain, mainly in neurons, where it was involved in the regulation of oxygen homeostasis and the maintenance of mitochondrial homeostasis, and its expression level was significantly increased under hypoxic stimulation[24-27]. However, recent studies found that hemoglobin may also be involved in the development of neurodegeneration; for example, α-synuclein pathology-hemoglobin-mitochondrial aggregates were found in Lewy bodies in patients with PD[28]. In addition, neuronal hemoglobin induced dopaminergic neuron

loss and cognitive deficits in the substantia nigra in mice[29], and hemoglobin deposited
in the cerebral cortex after intracerebral hemorrhage exerted neurotoxicity and
induced neuronal death in rats[30]. In summary, the specific role of hemoglobin
expression in neurons is controversial, and the role of hypoxia in the function of
hemoglobin in neurons is still unknown.

In this study, we established a chronic hypoxia mouse model and identified one of the
brain's responses to hypoxia. Specifically, we found that non-neuronal cells in the
brain regulate neuronal hemoglobin levels via EV-communication under hypoxic
conditions, as revealed by the combined analyses of snRNA-seq and EVs-RNA-seq.
We also established an *in vitro* hypoxia neuron model to investigate the specific
molecular mechanism of neuroprotection, demonstrating that exosomal hemoglobin
helped neurons resist hypoxic damage by maintaining mitochondrial homeostasis. In
summary, this study found a new mechanism of exosomal hemoglobin delivery in the
brain 's hypoxic response ' mutual aid ', and clarified the important role of exosomal
communication in the brain 's hypoxic stress response and resistance to injury. It
provides a new research perspective for the prevention and treatment of
hypoxic-ischemic-related brain diseases.

2   Results

2.1 Hypoxia induced cognitive and motor dysfunction and neuropathological
damage

In order to clarify the damage caused by hypoxia in the brain, we established a chronic
hypoxia model in C57/BL mice using a closed and controllable hypoxic box, the mice
were divided into a control group (Con) and a hypoxic treatment group (hypoxia for 28
days, H28d) (Fig. 1A). Behavioral tests and neuropathological staining were used to
examine the effect of chronic hypoxia on neurological impairment and nerve damage.
The results of the Y maze tests showed that the alternation behavior was significantly
decreased, and the novel object recognition tests showed that the exploration time of
new objects was significantly reduced after hypoxia, suggesting that hypoxia induced
cognitive dysfunction in mice (Fig. 1B and C). To evaluate spontaneous movement, we
performed an open-field tests and recorded the motor function and cognitive function
indicators of the two groups. The movement trajectory in the open field were
significantly reduced in H28d group (Fig. 1D). Assuming that the total exercise times
were similar in both groups, the exercise distance and speed were also significantly
reduced in H28d mice (Fig. 1E). These results suggested that the motor function was
impaired after hypoxia. In addition, H28d mice spent less time exploring the central
area of the open field and crossing the central area, indicating anxiety-like behavior
(Fig. 1F).

We subsequently examined the brains in the two groups of mice by neuropathological
staining to determine if this chronic hypoxic treatment induced nerve damage. Nissl
staining showed that neurons in the hippocampal CA1 region were disordered in H28d

group, with gaps, abnormal arrangement, and darker cytoplasm compared with Con group, suggested that the hippocampus was damaged in H28d group (Fig. 1G). It was also found that the hippocampal region of H28d group showed obvious atrophy (Fig. 1H). Neuronal degeneration staining (Fluoro-Jade B staining, FJB) showed neurodegeneration in the cerebral cortex and hippocampal region in mice in H28d group (Fig. 1I). We also carried out Golgi staining to reveal the morphology and structure of neurons in the brains in the two groups of mice, and showed that the density of dendritic spines on neurons was significantly lower in H28d group (Fig. 1J and K). Above all, chronic hypoxia induced cognitive and motor dysfunction and nerve damage in mice.

2.2 Hypoxia induced transcription of hemoglobin in non-neuronal cells

To explore the molecular mechanism responsible for chronic hypoxia-induced nerve injury in the brain, we performed snRNA-seq of two groups, and visualized the distribution of gene expression in all samples using UMAP visualization and identified 13 cell types according to the expression of cell-specific markers (Fig. 2A, Fig. S1A and B). We described the proportions of cells before and after hypoxia (Fig. 2B). Differentially up- and down-regulated genes in different cell types were analyzed by gene ontology (GO) enrichment analysis (Fig. S1C to L). Among these, the number of differentially expressed genes (DEGs) in endothelial cells (Endo) was the highest after hypoxia, and most of them were up-regulated (Fig. 2C). We subsequently displayed DEGs in all cell subsets, labeled the top 10 genes, and found the mRNA expression levels of hemoglobin subunits (*Hba-a1*, *Hba-a2*, *Hbb-bs*, *Hbb-bt*) were significantly up-regulated in most non-neuronal cells after hypoxia, indicated that might be a common mechanism whereby various non-neuronal cells in the brain respond to hypoxic stress (Fig. 2D).

We further clarified the interactions among various cell types in the brain under chronic hypoxic stress using the circle diagram and heat map, to show the changes in communication among different cell types in the H28d group compared with the Con group (Fig. 2E and F). We found that the number and intensity of intercellular communications in the brain both increased after hypoxia. Above all, these results suggested that various non-neuronal cells in the brain jointly transcribed hemoglobin under chronic hypoxic stress, which might represent a common mechanism in response to hypoxic stress.

2.3 Hypoxia induced the release of non-neuronal EVs carrying hemoglobin mRNA

SnRNA-seq analysis showed that chronic hypoxia increased intercellular communication in the brain; however, the effects of hypoxia on EVs-mediated communication remain unclear. Hypoxia has been shown to be an important factor inducing EVs release by cells, suggesting that the EVs-mediated communication pathway may be an important mechanism for cells to respond to hypoxia[31]. We

therefore performed EVs-RNA-seq of two groups to analyze the mechanisms involved
in the brain's resistance to hypoxic damage from the perspective of EVs regulation (Fig.
3A). We identified EVs isolated from mouse brain tissue. Transmission electron
microscopy confirmed that the diameter of the EVs was approximately 100 nm and
displayed a characteristic cup-shaped morphology (Fig. 3B). Western blot analysis
revealed positive expression of EV markers TSG101, HSP90, and CD9, while calnexin
was not detected (Fig. 3C). Nanoparticle tracking analysis indicated that the
predominant size of the isolated EVs ranged from 50 to 200 nm, with increased EV
secretion observed in the brains following hypoxia (Fig. 3D).

After the transcriptome sequencing of EVs isolated from Con and H28d mice, we
analyzed the DEGs between these two groups (Fig. S2A and B). A total of 2257 DEGs
were identified, of which 1493 were up-regulated and 764 were down-regulated after
hypoxia. Hemoglobin mRNA was also significantly up-regulated in EVs after hypoxia,
consistent with the snRNA-seq results (Fig. 3E). We conducted gene enrichment
analysis on all identified up- and down-regulated DEGs and selected the top 5 enriched
pathways for presentation. We found that several pathways related to RNA metabolism
were significantly up-regulated, while several pathways associated with energy
metabolism were down-regulated (Fig. S2C to F, Fig. 3F). These results suggested that
energy metabolism pathways in the brain may have been damaged by hypoxia, and
various cell types in the brain may regulate RNA metabolism via the EVs pathway to
resist hypoxic injury. The protein-protein interaction (PPI) analysis yielded results
consistent with the aforementioned analysis (Fig. S2C to H). And the Gene Set
Enrichment Analysis (GSEA) showed that the function of mitochondria in the brain
was also severely impaired after hypoxia (Fig. S2I and J).

We compared the DEGs obtained from EVs-RNA-seq with the snRNA-seq, and
screened out DEGs that changed consistently within the cell and in the extracellular
environment after hypoxia. Our findings revealed the proportion of cells releasing EVs
in each group, indicating a significant increase in EVs secretion solely by brain Endo
following hypoxia (Fig. 3G). Multiple differential mRNAs were present in EVs
secreted by various types of cells after hypoxia, including mRNAs of hemoglobin
subunits, suggesting that this may be a common way for multiple cells to respond to
hypoxia via the EVs pathway (Fig. 3H). We therefore analyzed the hemoglobin mRNA
levels in EVs from the two groups, and found that exosomal mRNA levels of
hemoglobin were significantly increased after hypoxia, but only existed in EVs
secreted by non-neuronal cells (Fig.3I and J). We subsequently performed GO analysis
on all DEGs with consistent changes in both snRNA-seq and EVs-RNA-seq according
to cell types for GO analysis, and presented the common signaling pathways
accumulated in three or more cell types. Many of these pathways were related to
hemoglobin function, suggesting that changes in hemoglobin after hypoxia induction
could be a common means by which various types of cells respond to hypoxia (Fig. 3K).
Above all, chronic hypoxia can enhance EVs communication in the brain, and induce
non-neuronal cells to transcribe hemoglobin and release them in the form of EVs.

2.4 Hypoxia induced the expression of hemoglobin in neurons

Combined analysis of snRNA-seq and EVs-RNA-seq showed numerous hemoglobin mRNA existed in the intracellular and extracellular environments in the brain after hypoxia, which may provide a common mechanism for various cell types to cope with hypoxic injury. However, changes in hemoglobin protein levels after hypoxia were still unclear. We observed brain tissues from two groups of mice and found that brain tissue in the H28d group remained red after perfusion with normal saline, indicating large amounts of hemoglobin in the brain (Fig. 4A). We further explored the transcriptional changes in hemoglobin subunits in cells after hypoxia by snRNA-seq and visualized the results in a violin plot, and the mRNA levels of hemoglobin subunits were low in all cell types in the Con group, but were increased in non-neuronal cells after hypoxia (Fig. 4B). RT-qPCR confirmed that *Hba-al* mRNA levels in the brain were significantly increased after hypoxia (Fig. 4C). Western blotting and immunofluorescence also revealed that Hba protein levels were also significantly increased in the H28d group (Fig. 4D to G).

Although previous studies showed that transient hypoxia induced the expression of hemoglobin in neurons in the brain, its specific function is still unclear[26]. We therefore speculated that hemoglobin expression levels may also be increased in neurons after hypoxia. We labeled Hba and Map2 by immunofluorescence, found that the Hba fluorescence intensity in Map2-positive cells increased significantly after hypoxia, indicating that hypoxia induced hemoglobin specifically expressed and accumulated in neurons (Fig. 4H to J). SnRNA-seq and EVs-RNA-seq showed that chronic hypoxia-induced hemoglobin transcription was mainly increased in non-neuronal cells and their secreted EVs (Fig. 4B and 3J), but immunofluorescence showed that hypoxia-induced increased hemoglobin was specifically present in neurons, confirming that non-neuronal cells in the brain could dynamically regulate hemoglobin levels in neurons via the EVs pathway under hypoxia, thus helping neurons to resist the hypoxic damage.

Neuroglobin (Ngb) is a recently discovered type of oxygen-carrying globin, which under hypoxic conditions can facilitate the diffusion of oxygen into mitochondria, enhancing tissue oxygen utilization. Particularly, it plays a crucial protective role in neural tissues during cerebral ischemia and hypoxia. To elucidate whether Ngb exerts protective effects in the brains of H28d mice, we conducted an analysis of snRNA-seq data. Visualization of Ngb transcription levels revealed consistently low expression levels across various brain cell types before and after hypoxia, contrasting with the behavior of hemoglobin. Consequently, this study continued its focus on elucidating the precise molecular mechanisms underlying hemoglobin's response to hypoxia.

Mitochondria are the main organelles involved in aerobic respiration and are highly sensitive to oxygen concentrations. We therefore determined if hypoxia induced mitochondrial damage in brain. Western blotting showed that mitochondrial complex I and IV were both significantly decreased in the brain after hypoxia, through analysing

the markers of mitochondrial complex I (NDUFB8) and complex IV (citrate synthase) (Fig. 4K to M). We further detected the activity of mitochondrial complex I in the brain in both groups, and showed that its activity was significantly decreased after hypoxia (Fig. 4N). These results confirm that chronic hypoxia can damage mitochondrial activity in the brain. Above all, these results indicate that hypoxia induced non-neuronal transcriptionally increased hemoglobin is transmitted to neurons via the EVs pathway.

2.5 Hypoxia specifically increased hemoglobin transcription in endothelial cells *in vitro*

We constructed an *in vitro* hypoxia model to clarify the specific protective mechanism of exosomal hemoglobin produced in hypoxia-induced neurons on mitochondria. In view of the most significant increase in cell communication intensity and the highest transcription of hemoglobin in endothelial cells (EC) after hypoxia, we used the human brain microvascular endothelial cells (hcMEC/D3 cells) as a representative of non-neuronal cells in the brain, and the human neuroblastoma cells (SH-SY5Y cells) as a cell model of hypoxia-induced nerve injury. We established four different hypoxia-duration groups using an oxygen concentration of 1%, and detected the effects on cell viability and cytotoxicity. The viability of hcMEC/D3 cells was significantly reduced in the H4h and subsequent groups, compared with the H0h group, the cytotoxicity was significantly increased in the H12h group (Fig. 5A and B). The viability of SH-SY5Y cells was significantly reduced in the H4h and subsequent groups, compared with the H0h group, the cytotoxicity was significantly increased in the H4h group and subsequent groups as well (Fig. 5C and D).

Considering the mitochondrial damage in the mouse hypoxia models, we detected the mitochondrial status of SH-SY5Y cells before and after hypoxia. We labeled the mitochondria using MitoTracker and detected the fluorescence intensity in each group of SH-SY5Y cells by flow cytometry to determine the number of mitochondria (Fig. 5E). Fluorescence intensity in SH-SY5Y cells decreased gradually with hypoxia time, indicating that the number of mitochondria was significantly reduced by prolongation of hypoxia (Fig. 5F). The mitochondrial membrane potential is an important indicator of mitochondrial function. Its reduction will induce mitochondrial homeostasis damage in cells, which in turn induces cell damage. We therefore evaluated the effects of hypoxia by JC-1 staining, and found the mitochondrial membrane potential also significantly decreased with prolongation of hypoxia (Fig. 5G). These results indicate that severe mitochondrial damage occurred when neurons encountered hypoxic stress, which may be a key factor in hypoxic nerve injury.

To further simulate the response of non-neuronal cells to transcribed hemoglobin and EVs to neurons and express in the former research, we first detected *HBA1* mRNA levels in hcMEC/D3 and SH-SY5Y cells in each group. Level of *HBA1* mRNA was significantly up-regulated in hcMEC/D3 cells in the H4h group, and then slowly decreased to pre-hypoxia levels (Fig. 5H). However, level of *HBA1* mRNA in

SH-SY5Y cells was unaffected by hypoxia (Fig. 5I). We then isolated and analyzed
EVs from hcMEC/D3 cells under different treatment durations and showed that EVs
were significantly increased in the H8h group compared with the H0h group, indicating
that hypoxia for 8 h induced EVs release by hcMEC/D3 cells (Fig. 5J, Fig. S3A to C).

Above all, we established an experimental hypoxia method for EC and neurons *in vitro*,
and found that hypoxia treatment for 8 hours induced a significant transcription of
hemoglobin in EC, which was subsequently released into the culture medium via EVs
(Fig. 5K).

2.6 Neurons resisted hypoxic injury by expressing exosomal hemoglobin *in vitro*

Based on the results showing that hypoxia treatment for 8 h had no significant toxic
effect on hcMEC/D3 cells, and hcMEC/D3 cells showed the most significant increase
in *HBA1* transcription and the largest release of EVs, we selected conditioned medium
from H8h hcMEC/D3 cells as the donors of hemoglobin mRNA EVs in subsequent *in*
vitro experiments, with conditioned medium from H0h hcMEC/D3 cells as the control.
To determine if the EVs secreted by EC under hypoxia were taken up and utilized by
neurons, we first isolated EVs from hcMEC/D3 H8h cells by ultracentrifugation, and
co-cultured them with SH-SY5Y cells after fluorescence labeling. The results showed
that hcMEC/D3 H8h EVs were actively taken up by SH-SY5Y cells, thus providing a
basis for subsequent exploration of the neuroprotective effect of EC-derived EVs (Fig.
6A). In order to determine if EVs secreted by EC helped neurons to dynamically
regulate hemoglobin levels and resist hypoxia-induced nerve damage, we used
EC-conditioned medium exchange experiments. Cells were subjected to hypoxia for 0
(H0h) or 8 h (H8h), hypoxia for 8 h + control medium (H8h + H0h (EC)), and hypoxia
for 8 h + conditioned medium (H8h + H8h (EC)). In the H8h + H0h (EC) and H8h +
H8h (EC) groups, endothelial cell H0h (+H0h (EC)) and H8h medium (+ H8h (EC))
were exchanged into SH-SY5Y cells, followed by hypoxic treatment. We initially
demonstrated that EC-conditioned medium does not cause damage to the cell viability
and mitochondrial homeostasis of SH-SY5Y cells (Fig. S3D to G).

We detected *HBA1* mRNA levels in SH-SY5Y cells in each group. *HBA1* mRNA levels
were significantly up-regulated in the H8h + H8h (EC) group, suggesting that
conditioned medium from EC treated with hypoxia for 8 h could increase the
transcription of hemoglobin after hypoxia (Fig. 6B). We also detected the protein levels
of Hba by immunofluorescence, and showed that Hba protein levels were significantly
increased in the H8h + H8h (EC) group, confirming that H8h conditioned medium from
EC could increase hemoglobin levels in neurons (Fig. 6C to D).

We confirmed that hypoxia treatment for 8 h decreased SH-SY5Y cell viability, the
number of mitochondria, and mitochondrial membrane potential. We therefore detected
the cell and mitochondrial viability of SH-SY5Y cells supplemented with EC-derived
conditioned medium and hypoxia, according to the above experimental protocol (Fig.
6E). Treatment with H8h EC-conditioned medium improved the number of

mitochondria and mitochondrial membrane potential (Fig. 6F and G), and improved the viability of neurons after hypoxia for 8 h (Fig. 6H).

To verified the function of exosomal hemoglobin in helping neurons to resist hypoxic injury in primary neurons (PN) *in vitro*, we designed the following experiments. The results of immunofluorescence showed that hcMEC/D3 H8h EVs were actively taken up by PN (Fig. 6I). The PN were then subjected to EC-conditioned medium exchange combined with hypoxia treatment to determine if conditioned medium could increase the transcription and expression levels of hemoglobin. The mRNA and protein levels of hemoglobin were significantly up-regulated in the H8h + H8h (EC) group (Fig. 6J to L). JC-1 staining showed that the mitochondrial membrane potential was significantly higher in the H8h + H8h (EC) group than in the H8h group (Fig. 6M). MitoTracker staining showed that mitochondrial numbers were significantly increased in the H8h + H8h (EC) group compared with the H8h group (Fig. 6N). We also evaluated the viability of PN in each group by PI/Hoechst staining and showed that their cell viability was significantly reduced after 8 h of hypoxia treatment, and that this damage could be attenuated by EC-conditioned medium (Fig. 6O). Above all, we found that EC-derived EVs carrying hemoglobin mRNA could regulate hemoglobin levels in neurons under hypoxic conditions, aiding neurons in maintaining mitochondrial homeostasis and resisting hypoxic damage (Fig. 6P).

To confirm that the protective effect of the EC-derived conditioned medium in terms of regulating hemoglobin levels in neurons was derived from EVs, we removed EVs from each group of medium and repeated the above experiments (Fig. 7A). After removing EVs, the H8h EC-conditioned medium lost its abilities to improve mitochondrial homeostasis and neuronal viability under hypoxic stress (Fig. 7B to D).

We further confirmed that the protective effect of EC-derived conditioned medium in regulating hemoglobin levels in neurons was derived from *HBA1* mRNA carried by EVs, by pre-treating the EC with *HBA1* RNAi plasmid interference (Fig. 7E, Fig. S3H to J). We screened plasmids for specific knock-down of *HBA1* mRNA in hcMEC/D3 cells, and verified that the resulting conditioned medium failed to increase hemoglobin mRNA in SH-SY5Y cells (Fig. S3K). Following HBA1 RNAi treatment of EC, the conditioned medium lost its abilities to improve mitochondrial homeostasis and neuronal viability under hypoxic stress (Fig. 7F to H).

In summary, this study found that EVs containing hemoglobin mRNA secreted by EC after hypoxia treatment regulated the hemoglobin expression in neurons, thus helping neurons to resist mitochondrial dysfunction and nerve injury under hypoxia stress (Fig. 7I).

3   Discussion

The brain is the most vulnerable organ to hypoxic injury and hypoxia is associated with the occurrence and development of various central nervous system diseases;

however, the specific molecular mechanisms are still unclear. Beyond the hypoxia-inducible factor signaling pathways, intercellular communication among diverse brain cell types is imperative for mitigating stress-induced damage in hypoxic conditions. Mounting evidence suggests that hypoxia enhances intercellular communication primarily through interactions involving EVs, thereby facilitating the transmission of various cargoes with distinct functional roles[5,31]. Our research findings also indicate that hypoxia induces a significant enhancement in EVs-mediated communication among various cell types in the brain. Through the isolation and purification of tissue-derived EVs followed by transcriptomic sequencing analysis, we are able to describe changes in intercellular communication within the brain from an extracellular perspective. However, this method does not provide a comprehensive map of intracellular communication[10]. Through the combined analysis of snRNA-seq and EVs-RNA-seq, we can accurately identify the cellular origin of brain-derived EVs, thus comprehensively revealing changes in intercellular communication within the brain induced by hypoxia. This study elucidates a novel mechanism whereby various non-neuronal cells in the brain regulate hemoglobin levels in neurons through EV-communication under hypoxic stress, emphasizing the crucial role of extracellular vesicle communication in the brain's response to hypoxia.

Hemoglobin, as the main protein carrying and transporting oxygen molecules in the body, is also expressed in neurons, but its precise function remains controversial [26,32,33]. Current research suggests that hemoglobin expressed in the brains of PD patients participates in regulating intraneuronal oxygen homeostasis and maintaining mitochondrial stability[34]. However, other studies indicate that hemoglobin may exert neurotoxic effects by inducing oxidative stress and neuronal death, thereby contributing to the onset and progression of neurodegeneration[29,30]. To clarify the role of hemoglobin in the brain under hypoxic stress, this study, based on the elucidation of the mechanism by which non-neuronal cells regulate neuronal hemoglobin through extracellular vesicles, conducted in vitro experiments. These experiments revealed that increased expression of extracellular vesicle-derived hemoglobin in neurons under hypoxia helps maintain mitochondrial homeostasis by sustaining mitochondrial membrane potential, thus aiding neurons in resisting hypoxic damage. This study clarified that the expression of exosomal hemoglobin in neurons contributes to the maintenance of mitochondrial homeostasis under stress, expanded the function of hemoglobin except transporting oxygen over a long range.

However, our research also had some limitations. First, we only clarified the hypoxic protective effect of hemoglobin on neuronal mitochondria *in vitro*, and were unable to explain the inconsistency between the increase in neuronal hemoglobin expression and the decrease of mitochondrial activity observed *in vivo*. We speculated that some pathological changes may make hemoglobin, which is increased in response to hypoxia, unable to exert its mitochondrial protective effect, even more than making hemoglobin transform and exacerbating nerve damage. For example, previous studies reported large numbers of phosphorylated α-synuclein and hemoglobin aggregates in

Lewy bodies in PD patients[28], and we previously confirmed that chronic hypoxia increased α-synuclein phosphorylation[35]. We therefore speculate that hemoglobin is increased in response to hypoxia and then phosphorylated by α-synuclein binding to stay in the cytoplasm, thus losing its mitochondrial protective ability and form pathological depositions, inducing cell death. However, further studies are needed to test this hypothesis.

In summary, we used the combined analysis of snRNA-seq and EVs-RNA-seq to examine the effects of hypoxia on interactions among different cell types in the brain, and to clarify the hypoxic response mechanism of non-neuronal EVs communication pathways to dynamically regulate hemoglobin levels in neurons. We revealed that increased levels of exosomal hemoglobin in neurons may help neurons to resist hypoxic injury by protecting mitochondrial function. These results clarify the molecular mechanism by which various non-neuronal cells interact via the EVs pathway to help neurons resist hypoxic injury, and highlight the important role of hemoglobin in hypoxia-related nerve injury, thus providing a new perspective for the prevention and treatment of various hypoxic-ischemic brain diseases.

4   Materials and Methods

4.1 Construction of chronic hypoxia model

A mouse model of chronic hypoxia was established in closed controlled hypoxic chambers (Ningbo Hua Yi Ning Chuang, China). Male C57BL/6J mice aged 2–3 months underwent a 28-day hypoxic cycle. Mice were maintained in a hypoxic chamber and the oxygen concentration was gradually decreased by 1% per day, from 21% O_2 to 7% O_2, and then maintained at 7% O_2 concentration for 14 days to establish a chronic hypoxic mouse model (H28d). Mice in the control group were maintained under 21% O_2 conditions (Con). All experimental animals were raised in a specific pathogen-free barrier environment, with normal temperature, humidity, and circadian rhythm, and given sufficient water and food. The study was approved by the Ethics Committee of Beijing Institute of Brain Disorders, Capital Medical University. All animal experiments were conducted in accordance with the National Institutes of Health Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines[36].

4.2 Behavioral tests

4.2.1   Y-maze test

Y-maze tests were performed as described previously[37]. The Y-maze consisted of three arms (40 × 4 × 9.5 cm, marked as A, B, and C, respectively) at angles of 120° to the central point. Each mouse was placed in the center of the starting arm and allowed to move freely in the maze for 8 min. The test was videotaped using a Nikon camera, manually recording the order of arm entry and calculating the percentage of alternations (calculated as [actual number of alternations/maximum number of alternations] ×100%).

4.2.2 Novel object preference test

Novel object preference tests were performed as described previously[38]. The mice were pre-accustomed in an empty open-field box for 3 consecutive days (5 min each time). During training, the mice were exposed to two identical objects for 10 min. During the test, the mice were put back in the cage for 60 min before the test, a familiar object and a new object were added, and the mice were allowed to explore these freely for 5 min. The numbers of times the mice explored new and old objects were recorded using a Nikon camera. The discrimination index was calculated as ([number of new object explorations/total number of object explorations] ×100). The cages and objects were cleaned with 75% ethanol to minimize any olfactory cues during the test.

4.2.3 Open-field test

Open-field tests were performed as described previously[39]. The mice were pre-accustomed in an open-field box (40 × 40 × 40 cm) for 3 consecutive days (5 min each time). During the test, the mice were placed in the open field and allowed to explore freely for 5 min. The movement time, movement distance, number of times through the central area, and residence time in the central area were recorded using a Nikon camera. The cages and objects were cleaned with 75% ethanol to minimize any olfactory cues during the test.

In the behavioral tests, mice with cognitive and motor disorders at baseline were excluded. All behavioral videos were analyzed blindly by researchers other than the operator.

4.3 Neuropathological staining

4.3.1 Nissl staining

Mouse brain tissue sections were fixed in 70% ethanol and dehydrated in graded ethanol solutions. Brain sections were permeabilized with xylene and incubated in 1% tar violet (Solarbio, China) for 30 min, washed with distilled water, and fixed in 70% alcohol. The slices were then dehydrated again in graded ethanol solutions and sealed with neutral glue.

4.3.2 Fluoro-Jade B (FJB) staining

Mouse brain tissue sections were immersed in 100% ethanol for 3 min, followed by 70% ethanol for 1 min, ddH$_2$O for 1 min, and 0.006% potassium permanganate for 15 min. After rinsing in ddH$_2$O for 1 min, the sections were incubated in 0.001% FJB (Cell Signaling Technology, Beverly, MA, USA) staining solution for 30 min, immersed in xylene, dried, and sealed with an anti-quenching sealer containing DAPI.

4.3.3 Golgi staining

Golgi–Cox staining was performed using a Golgi–Cox OptimStain Kit (Servicebio, China) to detect potential changes in the density and characteristics of neuronal dendritic spines. According to the instructions, whole brains were immersed in the dye solution for 14 days at room temperature in the dark. The tissue was then cut into 100-μm sections and photographed under a phase contrast microscope. The types of dendritic spines were detected, and cells with clear dendritic spine resolution (>10 μm long) and straight end branches were selected for dendritic spine counting and analysis.

4.4 Single-nuclear transcriptomics analysis of brain tissue

4.4.1 Mouse brain tissue separation

Mice were euthanized by rapid neck dislocation and perfused with intracardiac pre-cooled phosphate-buffered saline (PBS). The brains were removed, dissected into 50–200-mg pieces, and quickly frozen in liquid nitrogen until the nucleus was extracted.

4.4.2 snRNA-seq

Single-cell nuclear separation was performed as described previously[40]. Briefly, the frozen tissue was homogenized twice with a homogenizer (TIANZ, China), filtered through a 40-μm cell filter, centrifuged at $500 \times g$ at 4°C for 5 min, and the nucleus was resuspended in cell suspension buffer. Cells/nuclei were counted using a blood cell counter, and the concentration was adjusted to 1500 nuclei per μL for library preparation.

The single-cell nuclear suspension was subjected to droplet generation, emulsion fragmentation, bead collection, reverse transcription, and cDNA amplification to generate a barcode library. The index library was constructed according to the manufacturer's instructions. The sequencing library was quantified using a Qubit ssDNA Assay Kit (ThermoFisher Scientific, Waltham, MA, USA) and sequenced using BGI-Beijing's DIPSEQ-T7 sequencer.

4.4.3 snRNA-seq

CellRanger software was used to quantitatively analyze the gene expression of snRNA-seq data. The results were filtered using the R package Seurat (4.0.3), with removal of cells expressing 7000 genes and cells with mitochondrial gene content >25%. Unsupervised clustering was carried out using the Seurat package and the sample data were normalized using the NormalizeData function. The first 2000 highly variable genes were screened using the FindVariableGenes function. Uniform Manifold Approximation and Projection (UMAP) was used for dimensionality reduction and subsequent clustering. According to the clustering results, the corresponding gene markers were used for cell-type annotation. CellChat (v1.4.0) was

used to infer and quantify cell–cell communication in brain cell clusters in the Con
and H28d groups. The ligand–receptor interaction mouse database in CellChatDB
(containing CellChat) was used as a reference for cell–cell interaction analysis.

4.5 Transcriptome analysis of brain tissue EVs

4.5.1   Isolation of brain tissue EVs

Brain tissue suspension after enzymatic hydrolysis was subjected to gradient
centrifugation at 300, 2000, and 10,000 $\times$ g at 4°C, and filtered through a 0.22-μm
sieve to remove all cell debris and apoptotic bodies. The collected supernatant was
ultracentrifuged at 4°C, 110,000 $\times$ g for 2 h to obtain a precipitate containing EVs.
The precipitate was resuspended in PBS and further purified using an Exosupur ®
column (Echobiotech, China). The fraction was concentrated to a final volume of 200
μL using a 100 kDa cut-off Amicon ® Ultra rotary filter (Merck, Germany). Details of
the process are provided in the EV-TRACK protocol (EV220325)[41].

4.5.2   Western blots of EVs

The collected EV supernatant was denatured in sodium dodecyl sulfate buffer and
detected by western blot. The purified products were characterized as EVs by
incubation with primary antibodies for the EV-positive markers CD9 (Abcam,
ab92726), HSP70 (Abcam, ab181606), and TSG101 (Abcam, ab125011), and the
EV-negative marker calnexin (Proteintech, 10427-2-AP).

4.5.3   Transmission electron microscopy of EVs

The collected EV supernatant was fixed with 2% paraformaldehyde solution for 20
min. Sample droplets were then incubated on a copper mesh containing a biological
support membrane for 5–10 min, washed twice with ddH$_2$O, stained with 2% uranyl
acetate for 5 min, and washed twice with ddH$_2$O. The samples were then dried
thoroughly using an infrared baking lamp, and then observed and photographed with a
transmission electron microscope (Hitachi Ltd., Tokyo, Japan).

4.5.4   Nanoparticle tracking analysis (NTA) of EVs

The purified vesicle suspension was observed using a ZetaView PMX 110 equipped
with a 405 nm laser (Particle Metrix, Meerbusch, Germany) to determine the number
and size of the separated particles. A 60-s video was recorded at a speed of 30
frames/s, and the particle motion was analyzed using NTA software (ZetaView
8.02.28).

4.5.5   Analysis of EVs-RNA-seq

A total of 250 pg to 10 ng RNA per sample was used as input material for sequencing
libraries (1–500 ng for small RNA libraries) using the SMARTer Stranded Total
RNA-Seq Kit V2 (Takara Bio USA, Inc.) following the manufacturer's

recommendations, and index codes were added to attribute sequences to each sample.

RNA per sample was used as input material for the RNA sample preparations.
Sequencing libraries were generated using a QIAseq miRNA Library Kit (Qiagen,
Frederick, MD, USA) following the manufacturer's recommendation, and index codes
were added to attribute sequences to each sample. Reverse transcription (RT) primers
with unique molecular indices were introduced to analyze the quantification of mRNA
expression during cDNA synthesis and polymerase chain reaction (PCR)
amplification. The library quality was finally assessed using an Agilent Bioanalyzer
2100 and quantitative PCR (qPCR). Clustering of the index-coded samples was
performed using an acBot Cluster Generation System using TruSeq PE Cluster
Kitv3-cBot-HS (Illumina, San Diego, CA, USA) according to the manufacturer's
instructions. After cluster generation, the library preparations were sequenced on an
Illumina Hiseq platform and paired-end reads were generated.

Raw data (raw reads, fastq format) were first processed using in-house perl scripts. In
this step, clean reads were obtained by removing reads containing adapters, poly-N
reads, and low-quality reads. At the same time, the Q20, Q30, GC-content, and
sequence duplication level of the clean data were calculated. All downstream analyses
were based on high-quality clean data. Paired-end clean reads were aligned to the
reference genome GRCh38 using TopHat2/Bowtie2. Mapped reads were used to
quantify the gene expression level and for differential expression analysis.

Stringtie was used to calculate fragments per kilobase of exon per million fragments
mapped (FPKMs, calculated based on the length of the fragments and reads count
mapped to this fragment) of coding genes in each sample. FPKMs were computed by
summing the FPKMs of transcripts in each gene group. Differential expression was
determined between the Con and H28d groups.

4.6 Combined analysis of brain tissue EV transcriptome and single-cell
transcriptome

First, the online tool CIBERSORTx for deconvolution was used to create a signature
matrix from snRNA-seq data[42]. The parameter "Min. Expression" was set to zero and
"Replicates" was set to 100. In summary, this combined analysis method was used to
clarify the origin of EVs containing different mRNAs.

4.7 Real-time PCR

Total RNA was extracted from mouse cerebral cortex using a SteadyPure universal
RNA extraction kit (Accurate Bio, China). The RNA was then reverse transcribed into
cDNA using an Evo M-MLV reverse transcription premix kit (Accurate Bio, China).
Real-time PCR was carried out using an ABI real-time PCR system (Life
Technologies, USA) and SYBR Green Pro TaqHS premix qPCR kit III (Accurate Bio,
China), according to the manufacturer's instructions. The following primers were
used:

Mouse *Actb*, Forward: 5'-GGCTGTATTCCCCTCCATCG-3',

Reverse: 5'-CCAGTTGGTAACAATGCCATGT-3';

Mouse *Hba-a1*, Forward: 5'-GCCCTGGAAAGGATGTTTGC-3',

Reverse: 5'-TTCTTGCCGTGACCCTTGAC-3';

Human *RPL19*, Forward: 5'-CTAGTGTCCTCCGCTGTGG-3',

Reverse 5'-AAGGTGTTTTTCCGGCATC-3';

Human *HBA1*, Forward: 5'-TTCTGGTCCCCACAGACTCA-3',

Reverse: 5'-CTAGTGTCCTCCGCTGTGG-3'.

4.8 Western blot

Proteins were extracted from cells or tissues using RIPA buffer (Solarbio, China), and
total protein concentrations were quantified using a Pierce™ BCA Protein Assay Kit
(ThermoFisher Scientific, Waltham, MA, USA). Samples for gel electrophoresis were
prepared by boiling in loading buffer for 10 min. The protein lysates were then
analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, and
immunoblotted onto a polyvinylidene fluoride membrane. The membrane was
blocked with 5% skim milk at room temperature for 1 h, washed with TBST, and then
incubated with the specified primary antibody at 4°C overnight. The primary
antibodies included β-actin (HuaBio, EM21002, 1:10000), Hba (Abcam, ab92492,
1:1000), NDUFB8 (Proteintech, 14794-1-AP, 1:5000) and Citrate synthase
(Proteintech, 16131-1-AP, 1:2000).After incubation, the membrane was washed three
times and then incubated with the secondary antibodyat room temperature for 1 h,
including IRDye 680RD goat anti-mouse IgG (H + L) (Licor, 926-68070), IRDye
800CW goat anti-rabbit IgG (H + L) (Licor, 926-32211). The membrane was scanned
using a detection system (Odyssey, USA) and the band intensity was normalized to
β-actin. Statistical analysis was performed using ImageJ and GraphPad software.

4.9 Immunofluorescence

Mouse brain sections were placed in citric acid buffer (pH 6.0), boiled for 10 min for
antigen retrieval, and then washed three times with PBS for 5 min each time. The
brain slices were then placed in 0.2% PBS-Triton and incubated at room temperature
for 30 min, followed by 5% bovine serum albumin–PBS solution, and then blocked at
room temperature for 2 h. The slices were then incubated overnight in a primary
antibody diluent at 4°C in the refrigerator. The following primary antibodies were
used: Hba (Abcam, ab92492; Proteintech, 14537-1-AP), MAP2 (Abcam, ab32454).
Brain slices were washed three times with PBS for 5 min each time, and then
incubated with fluorescent secondary antibodies in the dark at room temperature for 1
h. The following secondary antibodies were used: goat anti-mouse IgG (H + L) highly

cross-adsorbed secondary antibody Alexa Fluor 488 (Invitrogen/ThermoFisher), and goat anti-rabbit IgG (H + L) highly cross-adsorbed secondary antibody Alexa Fluor 594 (Invitrogen/ThermoFisher). Nuclei were stained with DAPI before mounting (Sigma, D9542). Images were observed and captured by confocal microscopy.

4.10 Cell culture

4.10.1 Primary neurons

Primary cortical neuron cultures were prepared from Sprague-Dawley rat embryos at embryonic days 16–18. The cortex was dissected and separated mechanically by blowing through a straw to create a single-cell suspension, washed, and counted. The single-cell suspension was centrifuged at $1,000 \times g$ for 5 min and resuspended in DMEM by blowing, as above. The cell suspension was passed through a 40-μm filter. The cells were diluted to the required density and then transferred to a polylysine-coated plate and cultured in a 5% CO_2 incubator at 37°C for 4 h. The DMEM was then replaced with Neurobasal medium containing B27, glutamine, and penicillin–streptomycin. Half of the medium was replaced every 3 days after cell adhesion.

4.10.2 Cell lines

The human neuroblastoma cell line (SH-SY5Y) and human brain microvascular endothelial cell line (hcMEC/D3) were purchased from the American Center for Type Culture Collection (ATCC, USA). SH-SY5Y and hcMEC/D3 cells were cultured at 37°C in DMEM supplemented with 10% fetal bovine serum, penicillin (100 U/mL), and streptomycin (100 μg/mL) in a humidified 5% CO_2/95% air incubator. The cells were subjected to hypoxia in a three-gas incubator (Ningbo Hua Yi Ning Chuang, China) rinsed with a gaseous mixture of 1% O_2, 5% CO_2, and balanced N_2. The cells were inoculated and cultured under normoxia for 12 h to adhere, and then transferred to a hypoxic incubator for a specified time. All cell culture reagents were from ThermoFisher Scientific.

4.11 Isolation of EVs from cell culture medium

EVs Isolation from cell culture medium were performed as described previously[43]. After treatment of hcMEC/D3 cells, the culture medium was collected and centrifuged at $300 \times g$ for 5 min at room temperature to remove cell debris, followed by $3000 \times g$ at 4°C for 10 min to remove apoptotic bodies, and then filtered with a 0.22-μm filter membrane. The supernatant was then centrifuged at $10,000 \times g$ for 30 min to remove large microbubbles (>500 nm). The supernatant was ultracentrifuged at $110,000 \times g$ at 4°C for 70 min, and the EVs in the precipitate were collected and re-suspended in PBS.

4.12 EV uptake detection

Dil solution (5 μM; Invitrogen/ThermoFisher) was added to PBS containing EVs and

incubated for 30 min. Excess dye in the labeled EVs was removed by ultracentrifugation at $110,000 \times g$ at 4°C for 1 h. The EVs were then washed three times by re-suspending the precipitate in PBS, and the final washing was re-suspended in PBS. PBS solution containing Dil-EVs was added to a culture dish containing SH-SY5Y cells, and Dil-EV uptake by the cells was observed under a confocal microscope after 24 h of culture.

4.13   Cell viability

Cells were inoculated into 96-well plates at a density of 2000–3000 cells per well. CCK-8 cell viability assay (New Cell and Molecular Biotech, China) was carried out as follows: DMEM (1:10) solution was added to each well of the cell plate containing cells and samples and incubated at 37°C for 3 h. The absorbance (optical density) of each well was then measured at 450 nm using a microplate reader (SpectraMax i3, Molecular Devices, USA).

4.14   Cytotoxicity detection

Cytotoxicity was measured by lactate dehydrogenase (LDH) assay (Roche, USA). The powder was dissolved in ddH$_2$O and fully mixed to prepare a catalytic solution, followed by adding 250 μL of this catalytic solution to the dyeing solution (11.25 mL) and mixing to prepare the LDH reaction solution. Cell supernatant (100 μL) from each group was then added to a new 96-well plate and 100 μL LDH reaction solution was added and incubated at room temperature in the dark for 30 min. After incubation, 50 μL of termination solution was added to each well and gently mixed for 10 min, and the absorbance of each well at 490 nm was measured using a microplate reader.

4.15   Mitochondrial viability detection

Mitochondria in the cultured cells were labeled and their activity was detected using MitoTracker™ dye (Invitrogen/ThermoFisher), and the mitochondrial membrane potential was detected using JC-1 dye (Sigma-Aldrich, USA). MitoTracker dye was diluted to 50 nM with DMEM before staining, and JC-1 dye was diluted to 10 μg/mL with DMEM before staining, and stored in the dark at 4°C until use. After removing the original medium at the designated treatment end point, the cultured cells were digested and centrifuged at $1,000 \times g$ for 5 min in 1.5 mL centrifuge tubes and then resuspended in DMEM with dye and incubated for 20 min. The dye-containing medium was then removed by centrifugation at $1,000 \times g$ for 5 min, and the cells were washed twice with PBS, and then resuspended in PBS. The fluorescence intensity of each sample was detected using a BD LSRFortessa SORP flow cytometer (Becton, Dickinson and Company, USA). Follow-up analysis was performed using FlowJo.

4.16   Cell death detection

Propidium iodide (PI)/Hoechst staining was used to detect primary neuronal cell death.

The culture medium was discarded and the cells were washed three times with PBS, followed by incubation with PI (4 mM) and Hoechst (0.5 mg/mL) (Sigma-Aldrich, USA) at 37°C for 10 min, and washed with PBS for 5 min. Images were observed and captured by confocal microscopy.

4.17   Exchange of endothelial cell-conditioned medium

Medium exchange was performed to determine if EC-conditioned medium could exert neuroprotective effects. hcMEC/D3 cells were subjected to hypoxia and the culture medium was then collected. Cell debris and apoptotic bodies were removed by gradient centrifugation and filtered with a 0.22-μm filter membrane to obtain EC-conditioned medium. The conditioned medium was added to SH-SY5Y cells that had been adherent for 12 h, and subsequent hypoxia treatment and cell detection were performed.

EV removal and medium-exchange experiments were used to determine if the EVs in EC-conditioned medium could exert neuroprotective effects. The collected conditioned medium was subjected to ultracentrifugation to remove EVs, followed by medium exchange and subsequent experiments.

HBA1 RNAi plasmid transfection combined with medium-exchange experiments was used to determine if *HBA1* mRNA in EVs from EC-conditioned medium had a neuroprotective role. The hcMEC / D3 cells were transfected with *HBA1* RNAi plasmid mixed with Lipofectamine^{TM} 3000 and P3000^{TM} (Invitrogen / Thermo Fisher) in advance for 24 hours, followed by hypoxia and conditioned medium collection, and medium exchange and subsequent experiments were performed.

4.18   Statistical analysis

Data storage, recording, and statistical analysis were carried out using Excel and GraphPad Prism 10.0 software. The image data were analyzed using ImageJ. The Shapiro–Wilk tests and the Kolmogorov-Smirnov tests were used to check the normality of the variables. Unpaired *t*-tests (two-tailed) were used to compare differences between two sets of normally distribution data with equal variance, otherwise, non-parametric tests were used. Differences among multiple groups were analyzed by one-way/two-way analysis of variance (ANOVA) followed by Tukey's *post hoc* test. Image data were analyzed by ImageJ. Data were expressed as mean ± standard error (SEM) and $P < 0.05$ was considered statistically significant.

Author Contributions

Zhengming Tian: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Visualization; Writing—original draft. **Feiyang Jin:** Investigation; Methodology. **Yakun Gu:** Methodology. **Zirui Xu:** Methodology. **Mengyuan Guo:** Methodology. **Yuning Li:** Methodology. **Qianqian Shao:** Methodology. **Yingxia Liu:** Methodology. **Hanjiang Luo:** Methodology. **Yue Wang:** Methodology. **Suyu Zhang:** Methodology. **Chenlu Yang:** Methodology. **Xin Liu:** Methodology. **Xunming Ji:** Conceptualization; Formal analysis; Funding acquisition; Investigation; Project administration; Supervision; Validation; Writing—review & editing. **Jia Liu:** Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Supervision; Validation; Writing—original draft; Writing—review & editing.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (Grant number: 32100925), the Beijing Nova Program (Grant number: Z211100002121038, 20230484436), the Chinese Institutes for Medical Research, Beijing (Grant number: CX23YQ01), Beijing-Tianjin-Heibei Basic Research Cooperation Project (Grant number: 22JCZXJ00190).

Conflict of Interests

The authors declare that they have no competing interests.

References

1.   van Niel, G., Carter, D.R.F., Clayton, A., Lambert, D.W., Raposo, G., and Vader, P.

(2022). Challenges and directions in studying cell-cell communication by extracellular

vesicles. Nat Rev Mol Cell Biol *23*, 369-382. 10.1038/s41580-022-00460-3.

2.   Liang, W., Sagar, S., Ravindran, R., Najor, R.H., Quiles, J.M., Chi, L., Diao, R.Y.,

Woodall, B.P., Leon, L.J., Zumaya, E., et al. (2023). Mitochondria are secreted in

extracellular vesicles when lysosomal function is impaired. Nat Commun *14*, 5031.

10.1038/s41467-023-40680-5.

3.   Cocozza, F., Martin-Jaular, L., Lippens, L., Di Cicco, A., Arribas, Y.A., Ansart, N.,

Dingli, F., Richard, M., Merle, L., Jouve San Roman, M., et al. (2023). Extracellular

vesicles and co-isolated endogenous retroviruses from murine cancer cells

differentially affect dendritic cells. Embo j *42*, e113590. 10.15252/embj.2023113590.

4.   Kalluri, R., and LeBleu, V.S. (2020). The biology, function, and biomedical applications

of exosomes. Science *367*. 10.1126/science.aau6977.

5.   Korvenlaita, N., Gómez-Budia, M., Scoyni, F., Pistono, C., Giudice, L., Eamen, S.,

Loppi, S., de Sande, A.H., Huremagic, B., Bouvy-Liivrand, M., et al. (2023). Dynamic

release of neuronal extracellular vesicles containing miR-21a-5p is induced by

hypoxia. J Extracell Vesicles *12*, e12297. 10.1002/jev2.12297.

6.   Fabbiano, F., Corsi, J., Gurrieri, E., Trevisan, C., Notarangelo, M., and D'Agostino,

V.G. (2020). RNA packaging into extracellular vesicles: An orchestra of RNA-binding

788   proteins? J Extracell Vesicles *10*, e12043. 10.1002/jev2.12043.

789 7. Yoshimura, A., Kawamata, M., Yoshioka, Y., Katsuda, T., Kikuchi, H., Nagai, Y.,

790   Adachi, N., Numakawa, T., Kunugi, H., Ochiya, T., and Tamai, Y. (2016). Generation

791   of a novel transgenic rat model for tracing extracellular vesicles in body fluids. Sci Rep

792   *6*, 31172. 10.1038/srep31172.

793 8. Li, W., Li, C., Zhou, T., Liu, X., Liu, X., Li, X., and Chen, D. (2017). Role of exosomal

794   proteins in cancer diagnosis. Mol Cancer *16*, 145. 10.1186/s12943-017-0706-8.

795 9. Mathieu, M., Martin-Jaular, L., Lavieu, G., and Théry, C. (2019). Specificities of

796   secretion and uptake of exosomes and other extracellular vesicles for cell-to-cell

797   communication. Nat Cell Biol *21*, 9-17. 10.1038/s41556-018-0250-9.

798 10. Zhou, B., Guo, W., Guo, L., Li, Y., Zheng, Z., Huai, Q., Tan, F., Li, Y., Xue, Q., Ying, J.,

799   et al. (2023). Single-cell RNA-sequencing data reveals the genetic source of

800   extracellular vesicles in esophageal squamous cell carcinoma. Pharmacol Res *192*,

801   106800. 10.1016/j.phrs.2023.106800.

802 11. D'Acunzo, P., Kim, Y., Ungania, J.M., Pérez-González, R., Goulbourne, C.N., and

803   Levy, E. (2022). Isolation of mitochondria-derived mitovesicles and subpopulations of

804   microvesicles and exosomes from brain tissues. Nat Protoc *17*, 2517-2549.

805   10.1038/s41596-022-00719-1.

806 12. Svensson, V., Vento-Tormo, R., and Teichmann, S.A. (2018). Exponential scaling of

807   single-cell RNA-seq in the past decade. Nat Protoc *13*, 599-604.

10.1038/nprot.2017.149.

13.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd,

Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration

of Single-Cell Data. Cell *177*, 1888-1902.e1821. 10.1016/j.cell.2019.05.031.

14.   Xiao, D., Hu, X.Q., Huang, X., Zhou, J., Wilson, S.M., Yang, S., and Zhang, L. (2013).

Chronic hypoxia during gestation enhances uterine arterial myogenic tone via

heightened oxidative stress. PLoS One *8*, e73731. 10.1371/journal.pone.0073731.

15.   Wang, N., Chen, C., Cheng, Y., Fu, Y., Zhong, Z., Yang, Y., Lv, L., Chen, H., Huang,

J., and Duan, Y. (2022). Hypoxia drives hematopoiesis with the enhancement of T

lineage through eliciting arterial specification of hematopoietic endothelial progenitors

from hESC. Stem Cell Res Ther *13*, 282. 10.1186/s13287-022-02967-0.

16.   Choudhry, H., and Harris, A.L. (2018). Advances in Hypoxia-Inducible Factor Biology.

Cell Metab *27*, 281-298. 10.1016/j.cmet.2017.10.005.

17.   Burtscher, J., Mallet, R.T., Burtscher, M., and Millet, G.P. (2021). Hypoxia and brain

aging: Neurodegeneration or neuroprotection? Ageing Res Rev *68*, 101343.

10.1016/j.arr.2021.101343.

18.   Kann, O., and Kovács, R. (2007). Mitochondria and neuronal activity. Am J Physiol

Cell Physiol *292*, C641-657. 10.1152/ajpcell.00222.2006.

19.   Silver, I., and Erecińska, M. (1998). Oxygen and ion concentrations in normoxic and

827   hypoxic brain cells. Adv Exp Med Biol *454*, 7-16. 10.1007/978-1-4615-4863-8_2.

828 20. Wang, Y.Y., Huang, Z.T., Yuan, M.H., Jing, F., Cai, R.L., Zou, Q., Pu, Y.S., Wang,

829   S.Y., Chen, F., Yi, W.M., et al. (2021). Role of Hypoxia Inducible Factor-1α in

830   Alzheimer's Disease. J Alzheimers Dis *80*, 949-961. 10.3233/jad-201448.

831 21. Mitroshina, E.V., Savyuk, M.O., Ponimaskin, E., and Vedunova, M.V. (2021).

832   Hypoxia-Inducible Factor (HIF) in Ischemic Stroke and Neurodegenerative Disease.

833   Front Cell Dev Biol *9*, 703084. 10.3389/fcell.2021.703084.

834 22. Burtscher, J., Syed, M.M.K., Lashuel, H.A., and Millet, G.P. (2021). Hypoxia

835   Conditioning as a Promising Therapeutic Target in Parkinson's Disease? Mov Disord

836   *36*, 857-861. 10.1002/mds.28544.

837 23. Gell, D.A. (2018). Structure and function of haemoglobins. Blood Cells Mol Dis *70*,

838   13-42. 10.1016/j.bcmd.2017.10.006.

839 24. Auvinen, J., Tapio, J., Karhunen, V., Kettunen, J., Serpi, R., Dimova, E.Y., Gill, D.,

840   Soininen, P., Tammelin, T., Mykkänen, J., et al. (2021). Systematic evaluation of the

841   association between hemoglobin levels and metabolic profile implicates beneficial

842   effects of hypoxia. Sci Adv *7*. 10.1126/sciadv.abi4822.

843 25. Singhal, N.K., Alkhayer, K., Shelestak, J., Clements, R., Freeman, E., and

844   McDonough, J. (2018). Erythropoietin Upregulates Brain Hemoglobin Expression and

845   Supports Neuronal Mitochondrial Activity. Mol Neurobiol *55*, 8051-8058.

846   10.1007/s12035-018-0971-6.

26.   Schelshorn, D.W., Schneider, A., Kuschinsky, W., Weber, D., Krüger, C., Dittgen, T.,

Bürgers, H.F., Sabouri, F., Gassler, N., Bach, A., and Maurer, M.H. (2009).

Expression of hemoglobin in rodent neurons. J Cereb Blood Flow Metab *29*, 585-595.

10.1038/jcbfm.2008.152.

27.   Ferrer, I., Gómez, A., Carmona, M., Huesa, G., Porta, S., Riera-Codina, M., Biagioli,

M., Gustincich, S., and Aso, E. (2011). Neuronal hemoglobin is reduced in Alzheimer's

disease, argyrophilic grain disease, Parkinson's disease, and dementia with Lewy

bodies. J Alzheimers Dis *23*, 537-550. 10.3233/jad-2010-101485.

28.   Killinger, B.A., Marshall, L.L., Chatterjee, D., Chu, Y., Bras, J., Guerreiro, R., and

Kordower, J.H. (2022). In situ proximity labeling identifies Lewy pathology molecular

interactions in the human brain. Proc Natl Acad Sci U S A *119*.

10.1073/pnas.2114405119.

29.   Santulli, C., Bon, C., De Cecco, E., Codrich, M., Narkiewicz, J., Parisse, P.,

Perissinotto, F., Santoro, C., Persichetti, F., Legname, G., et al. (2022). Neuronal

haemoglobin induces loss of dopaminergic neurons in mouse Substantia nigra,

cognitive deficits and cleavage of endogenous α-synuclein. Cell Death Dis *13*, 1048.

10.1038/s41419-022-05489-y.

30.   Wang, X., Mori, T., Sumii, T., and Lo, E.H. (2002). Hemoglobin-induced cytotoxicity in

rat cerebral cortical neurons: caspase activation and oxidative stress. Stroke *33*,

1882-1888. 10.1161/01.str.0000020121.41527.5d.

31.   Wang, X., Wu, R., Zhai, P., Liu, Z., Xia, R., Zhang, Z., Qin, X., Li, C., Chen, W., Li, J.,

868   and Zhang, J. (2023). Hypoxia promotes EV secretion by impairing lysosomal

869   homeostasis in HNSCC through negative regulation of ATP6V1A by HIF-1α. J

870   Extracell Vesicles *12*, e12310. 10.1002/jev2.12310.

871   32.   Russo, R., Zucchelli, S., Codrich, M., Marcuzzi, F., Verde, C., and Gustincich, S.

872   (2013). Hemoglobin is present as a canonical α2β2 tetramer in dopaminergic neurons.

873   Biochim Biophys Acta *1834*, 1939-1943. 10.1016/j.bbapap.2013.05.005.

874   33.   Zheng, R., Yan, Y., Pu, J., and Zhang, B. (2022). Physiological and Pathological

875   Functions of Neuronal Hemoglobin: A Key Underappreciated Protein in Parkinson's

876   Disease. Int J Mol Sci *23*. 10.3390/ijms23169088.

877   34.   Vanni, S., Zattoni, M., Moda, F., Giaccone, G., Tagliavini, F., Haïk, S., Deslys, J.P.,

878   Zanusso, G., Ironside, J.W., Carmona, M., et al. (2018). Hemoglobin mRNA Changes

879   in the Frontal Cortex of Patients with Neurodegenerative Diseases. Front Neurosci *12*,

880   8. 10.3389/fnins.2018.00008.

881   35.   Li, G., Liu, J., Guo, M., Gu, Y., Guan, Y., Shao, Q., Ma, W., and Ji, X. (2022). Chronic

882   hypoxia leads to cognitive impairment by promoting HIF-2α-mediated ceramide

883   catabolism and alpha-synuclein hyperphosphorylation. Cell Death Discov *8*, 473.

884   10.1038/s41420-022-01260-6.

885   36.   Percie du Sert, N., Hurst, V., Ahluwalia, A., Alam, S., Avey, M.T., Baker, M., Browne,

886   W.J., Clark, A., Cuthill, I.C., Dirnagl, U., et al. (2020). The ARRIVE guidelines 2.0:

Updated guidelines for reporting animal research. J Cereb Blood Flow Metab *40*,

1769-1777. 10.1177/0271678x20943823.

37. Maki, T., Choi, Y.K., Miyamoto, N., Shindo, A., Liang, A.C., Ahn, B.J., Mandeville, E.T.,

Kaji, S., Itoh, K., Seo, J.H., et al. (2018). A-Kinase Anchor Protein 12 Is Required for

Oligodendrocyte Differentiation in Adult White Matter. Stem Cells *36*, 751-760.

10.1002/stem.2771.

38. Lueptow, L.M. (2017). Novel Object Recognition Test for the Investigation of Learning

and Memory in Mice. J Vis Exp. 10.3791/55718.

39. Chow, H.M., Shi, M., Cheng, A., Gao, Y., Chen, G., Song, X., So, R.W.L., Zhang, J.,

and Herrup, K. (2019). Age-related hyperinsulinemia leads to insulin resistance in

neurons and cell-cycle-induced senescence. Nat Neurosci *22*, 1806-1819.

10.1038/s41593-019-0505-1.

40. Hodge, R.D., Bakken, T.E., Miller, J.A., Smith, K.A., Barkan, E.R., Graybuck, L.T.,

Close, J.L., Long, B., Johansen, N., Penn, O., et al. (2019). Conserved cell types with

divergent features in human versus mouse cortex. Nature *573*, 61-68.

10.1038/s41586-019-1506-7.

41. Van Deun, J., Mestdagh, P., Agostinis, P., Akay, Ö., Anand, S., Anckaert, J., Martinez,

Z.A., Baetens, T., Beghein, E., Bertier, L., et al. (2017). EV-TRACK: transparent

reporting and centralizing knowledge in extracellular vesicle research. Nat Methods *14*,

228-232. 10.1038/nmeth.4185.

42.   Newman, A.M., Steen, C.B., Liu, C.L., Gentles, A.J., Chaudhuri, A.A., Scherer, F.,

Khodadoust, M.S., Esfahani, M.S., Luca, B.A., Steiner, D., et al. (2019). Determining

cell type abundance and expression from bulk tissues with digital cytometry. Nat

Biotechnol *37*, 773-782. 10.1038/s41587-019-0114-2.

43.   Zhang, K., Li, R., Chen, X., Yan, H., Li, H., Zhao, X., Huang, H., Chen, S., Liu, Y.,

Wang, K., et al. (2023). Renal Endothelial Cell-Targeted Extracellular Vesicles Protect

the Kidney from Ischemic Injury. Adv Sci (Weinh) *10*, e2204626.

10.1002/advs.202204626.

Fig. 1. Hypoxia induced neurological dysfunction and nerve injury in mice. (A)
Schematic diagram of chronic hypoxic mouse model construction. Recognition
coefficient in Con and H28d mice in (B) Y-maze test (n = 6) and (C) new object
recognition test (n = 10). (D) Trajectories, (E) movement index statistics, including
movement distance, speed, and time (n = 10), and (F) cognitive indicators, including
length of exploration and number of crossings in the central area (n = 10) for Con and
H28d mice in open-field tests. (G) Nissl staining showing arrangement of neurons in
the hippocampus in Con and H28d mice. (H)The area of the hippocampus was counted
in each group (n = 5). (I) FJB staining showing neurodegeneration in the cerebral
cortex and hippocampus in Con and H28d mice. (J) Golgi staining showing dendritic
spines of neurons in the hippocampus in Con and H28d mice. (K)The number of
dendritic spines on a single axon was counted (n = 5). In (E) and (K), data expressed as
mean ± SEM (Welch's t-test); other data expressed as mean ± SEM (unpaired t-test).
ns: not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Con: control
group; H28d: mice treated with hypoxia for 28 days.

Fig. 2. Chronic hypoxia increased hemoglobin transcription in mouse brain. (A)
Transcription profiles of cell subsets in the brain in Con and H28d mice visualized by
UMAP. **(B)** Proportions of different cells in the brain in Con and H28d mice. **(C)**
Histogram showing numbers of differentially expressed genes (DEGs) in different cell
types in mouse brain after hypoxia. Up- and down-regulated forms displayed
separately. **(D)** DEGs in different cell types in mouse brain after hypoxia shown in
stacked volcano plots. Top 10 DEGs marked based on absolute log2FC value. **(E)**
Circle diagram showing number and intensity of intercellular communications among
different cell types in the brain in Con and H28d mice. Red indicates increase and
blue indicates decrease. **(F)** Communications among cells in the H28d and Con
groups demonstrated by combination of heat map and histogram. Con: control group;
H28d: mice treated with hypoxia for 28 days. AST: Astrocyte; Endo: Endothelial cell;
EPC: Ependymocyte; EX: Excitatory neuron; Granule: Granule cell; IN: Inhibitory
neuron; Microgila: Microgila; mt-rich: Mitochondrial-rich cell; Nend:
Neuronendocrine cell; OLI: Oligodendrocyte; OPC: Oligodendrocyte precursor cell;
Purkinje: Purkinje cell; VLMC: Vascular lepotomeningeal cell.

Fig. 3. Hypoxia increased release of EVs carrying hemoglobin mRNA in mouse brain. (A) Schematic diagram of EV extraction and transcriptome sequencing in brain tissue from Con and H28d mice. **(B)** Typical electron microscope images of brain tissue EVs (scale bar = 0.5 μm). **(C)** Western blot analysis of EVs markers TSG101, HSP70, and CD9 and EVs negative marker calnexin in EVs isolated from cell lysates and brain tissues. **(D)** NTA analysis of brain tissue EVs isolated from Con and H28d mice. **(E)** Differentially expressed genes (DEGs) carried by mouse brain tissue EVs after hypoxia displayed by volcanic maps. Hemoglobin subunits labeled. **(F)** Histogram showing enriched signaling pathways for top five up- and down-regulated DEGs in EVs-RNA-seq. **(G)** The proportion of cells releasing EVs in each group. Data expressed as mean ± SEM (two-way ANOVA). *: up-regulated, #: down-regulated. ns: not significant, **P<0.01, # P<0.05, ## P<0.01. **(H)** Histogram illustrating the frequencies of cell types secreting EVs containing the same DEG mRNAs. The mRNAs co-carried by EVs secreted by seven, eight, and nine cells are displayed. **(I)** mRNA levels of each hemoglobin subunit in brain tissue EVs from Con and H28d mice

(n = 3). Data expressed as mean ± SEM (two-way ANOVA). ***P<0.001,
****P<0.0001. **(J)** Heat map showing cell sources of EVs carrying hemoglobin subunit
mRNAs. **(K)** GO enrichment analysis of DEGs obtained by combined snRNA-seq and
EVs-RNA-seq. Signal pathways of more than three cell types displayed in a bubble
map. Con: control group; H28d: mice treated with hypoxia for 28 days. AST:
Astrocyte; Endo: Endothelial cell; EPC: Ependymocyte; EX: Excitatory neuron;
Granule: Granule cell; IN: Inhibitory neuron; Microgila: Microgila; mt-rich:
Mitochondrial-rich cell; Nend: Neuronendocrine cell; OLI: Oligodendrocyte; OPC:
Oligodendrocyte precursor cell; Purkinje: Purkinje cell; VLMC: Vascular
lepotomeningeal cell.
976

Fig. 4. Hypoxia increased of hemoglobin expression in mouse neurons. (**A**) Brains
of Con and H28d mice. (**B**) Expression levels of hemoglobin subunits in snRNA-seq in
different cell types in the brain shown by violin maps. (**C**) *Hba-a1* mRNA levels in
brain in Con and H28d mice detected by real-time qPCR (n = 5). (**D and E**) Hba
expression levels in the brain in Con and H28d mice detected by western blotting (n =
3). (**F and G**) Hba expression levels in the brain in Con and H28d mice detected by
immunofluorescence (scale bar = 20 μm, n = 6). (**H**) Expression levels of Hba and
Map2 in the brain in Con and H28d mice detected by immunofluorescence (scale bar =
20 μm, n = 5). (**I**) Fluorescence intensity of Map2 in the brain in Con and H28d mice.
(**J**) Co-localization of Hba and Map2 in the brain in Con and H28d mice. (**K to M**)
Expression levels of mitochondrial complex I marker NDUFB8 and mitochondrial
complex IV marker citrate synthase in the brain in Con and H28d mice detected by
western blotting (n = 4-5). (**N**) Activity of mitochondrial complex I in the brain in Con
and H28d mice (n = 4-5). In (**C**), (**E**) and (**G**), data expressed as mean ± SEM (Welch's
t-test); other data expressed as mean ± SEM (unpaired t-test). ns: not significant,
*P<0.05, **P<0.01, ***P<0.001. Con: control group; H28d: mice treated with hypoxia
for 28 days.

Fig. 5. Hypoxia increased hemoglobin transcription in endothelial cells and induced release of EVs. (A) Viability of hcMEC/D3 cells detected by CCK-8 assay in H0h, H4h, H8h, and H12h groups (n = 8) and **(B)** cytotoxicity in each group detected by lactate dehydrogenase (LDH) assay (n = 6). **(C)** Viability of SH-SY5Y cells in each group detected by CCK-8 assay (n =8) and **(D)** cytotoxicity in each group detected by LDH assay (n = 8). **(E-F)** Mitochondria were labeled in SH-SY5Y groups using MitoTracker staining, and the number of mitochondria in each group was analyzed and counted by flow cytometry (n = 5). **(G)** Mitochondrial membrane potential in each SH-SY5Y group was detected by JC-1 labeling and analyzed by flow cytometry (n = 5). *HBA1* mRNA levels in **(H)** hcMEC/D3 cells and **(I)** SH-SY5Y cells detected by real-time qPCR (n = 5 each). **(J)** NTA analysis of EVs in hcMEC/D3 cells treated with hypoxia for 0 and 8 h. **(K)** Mechanism diagram. In **(B)**, data expressed as mean ± SEM (Brown-Forsythe ANOVA test); other data as mean ± SEM (one-way ANOVA). ns: not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. H0, 4, 8, 12h: cells treated with hypoxia for 0, 4, 8, 12 hours.

1012

1013

Fig. 6. Hypoxia induced neuronal uptake of endothelial cell-derived EVs and
expression of hemoglobin. (A) EVs in medium from H8h hcMEC/D3 cells were
labeled with Dil dye and added to SH-SY5Y cells labeled with Calcein AM for 24 h.
Uptake of hcMEC/D3 EVs by SH-SY5Y cells was observed by confocal microscopy
(scale bar = 5 μm). **(B)** *HBA1* mRNA levels in SH-SY5Y cells detected by real-time
qPCR (n = 5). **(C and D)** Hba1 expression levels in SH-SY5Y cells detected by
immunofluorescence (scale bar = 10 μm) (n = 10). **(E)** hcMEC/D3 conditioned medium
treatment diagram. **(F)** Viability of SH-SY5Y cells (n = 8-10). **(G)** Numbers of
mitochondria in SH-SY5Y cells (n = 5). **(H)** Mitochondrial membrane potential in
SH-SY5Y cells (n = 5). **(I)** EVs in medium from H8h hcMEC/D3 cells were labeled
with Dil and then added to calcein AM-labeled primary neurons for 24 h. Uptake of
hcMEC/D3 EVs by primary neurons was observed by confocal microscopy (scale bar =
5 μm). **(J)** *Hba-a1* mRNA levels in primary neurons detected by real-time qPCR (n =
5). **(K and L)** Hba expression levels in primary neurons detected by
immunofluorescence (scale bar = 10 μm) (n = 10). **(M)** Viability of primary neurons in
each group characterized by PI/Hoechst staining and photographed by confocal
microscopy (n = 8). **(N)** Number of mitochondria in primary neurons in each group
characterized by MitoTracker staining and photographed by confocal microscopy (n =

9). **(O)** Mitochondrial membrane potential of primary neurons in each group characterized by JC-1 staining and photographed by confocal microscopy (n = 9). **(P)** Mechanism diagram. In **(L)**, data expressed as mean ± SEM (Brown-Forsythe ANOVA test); other data as mean ± SEM (one-way ANOVA). ns: not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Fig. 7. Exosomal hemoglobin helped neurons to resist hypoxic injury by maintaining neuronal mitochondrial homeostasis. (A) hcMEC/D3 conditioned medium EV removal pretreatment diagram. **(B to D)** Viability, mitochondrial number, and mitochondrial membrane potential of SH-SY5Y cells (n = 5-8). **(E)** HBA RNAi pretreatment diagram of hcMEC/D3 conditioned medium. **(F** to **H)** Viability, mitochondrial number, and mitochondrial membrane potential of SH-SY5Y cells (n = 3-8). **(I)** Mechanism diagram. In **(H)**, data expressed as mean ± SEM (Brown-Forsythe ANOVA test); other data as mean ± SEM (one-way ANOVA). ns: not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

1049

Fig. S1. SnRNA-seq analysis identified 13 cell types in the mouse brain. (A) Transcription profiles in the brain in Con and H28d mice visualized by UMAP. **(B)** Bubble map showing marker expression of each cell type in mouse brain. **(C to L)** Histogram showing enriched signaling pathways for top five up- and down-regulated DEGs in 10 kinds of cells in the brain. Con: control group; H28d: mice treated with hypoxia for 28 days. AST: Astrocyte; Endo: Endothelial cell; EPC: Ependymocyte; EX: Excitatory neuron; Granule: Granule cell; IN: Inhibitory neuron; Microgila: Microgila; mt-rich: Mitochondrial-rich cell; Nend: Neuronendocrine cell; OLI: Oligodendrocyte; OPC: Oligodendrocyte precursor cell; Purkinje: Purkinje cell; VLMC: Vascular lepotomeningeal cell.

1060

1061

Fig. S2. EVs-RNA-seq analysis showed that mitochondrial function was impaired
in the mouse brain after hypoxia. (A) Levels of gene expression in each sample. **(B)**
The average expression distribution heat map of DEGs top1000 in EVs-RNA-seq. **(C**
to D) Histogram diagram showing signaling pathways enriched in differentially up-
and down-regulated genes in EVs. **(E to F)** Cluster diagram showing signaling
pathways enriched in differentially up- and down-regulated genes in EVs. **(G to H)**
Cluster diagram showing protein–protein interaction network of up- regulated DEGs
in EVs. **(I to J)** GSEA analysis of EVs-RNA-seq.

1070

1071

Fig. S3. *HBA1* **siRNA treatment disrupts the transfer of hemoglobin mRNA in EC-conditioned medium. (A to B)** NTA analysis of EVs in hcMEC/D3 cells treated with hypoxia for 0, 4 and 12 h. **(C)** NTA analysis of EC-derived EVs. **(D)** Safety detection of endothelial cell conditioned medium. **(E)** Viability of SH-SY5Y cells (n = 8-10). **(F)** Numbers of mitochondria in SH-SY5Y cells (n = 5). **(G)** Mitochondrial membrane potential in SH-SY5Y cells (n = 5-6). **(H)** *HBA1* mRNA levels in hcMEC/D3 cells detected by real-time qPCR. **(I and J)** hcMEC/D3 cells were labeled with calcein AM and the ratio of calcein AM-positive to mCherry-positive cells carried by the plasmid was observed by confocal microscopy. The *HBA1* RNAi plasmid was transfected and screened. **(K)** *HBA1* mRNA levels in hcMEC/D3 cells detected by real-time qPCR. In **(K)**, data expressed as mean ± SEM (Brown- Forsythe ANOVA test); other data as mean ± SEM (one-way ANOVA). ns: not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

1085