Virtual Colon: Spatiotemporal modelling of metabolic interactions in a computational colonic environment

Georgios Marinos[1,2]*, Johannes Zimmermann[1,3,4,5]*, Jan Taubenheim[1], and Christoph Kaleta[1,$]

[1] Research Group Medical Systems Biology, University Hospital Schleswig-Holstein Campus Kiel, Kiel University, Kiel, Schleswig-Holstein, Germany

[2] CAU Innovation GmbH, Kiel University, Kiel, Schleswig-Holstein, Germany

[3] Antibiotic resistance group, Max Planck Institute for Evolutionary Biology, Ploen, Schleswig-Holstein, Germany

[4] Evolutionary Ecology and Genetics, Zoological Institute, Kiel University, Kiel, Schleswig-Holstein, Germany

[5] Junior Research Group Mechanisms of Microbial Metabolic Interactions, Friedrich Schiller University Jena, Jena, Thüringen, Germany

*Shared first authorship with an interchangeable order

[$]Senior and corresponding author: c.kaleta@iem.uni-kiel.de

Georgios Marinos ORCID 0000-0002-6443-7696

Johannes Zimmermann ORCID 0000-0002-5041-1954

Jan Taubenheim ORCID 0000-0001-7283-1768

Christoph Kaleta ORCID 0000-0001-8004-9514

1. Abstract

Host-microbial metabolic interactions have been recognised as an essential factor in host health and disease. Genome-scale metabolic modelling approaches have made important contributions to our understanding of the interactions in such communities. One particular such modelling approach is BacArena in which metabolic models grow, reproduce, and interact as independent agents in a spatiotemporal metabolic environment. Here, we present a modelling application of BacArena, a virtual colonic environment, which reveals spatiotemporal metabolic interactions in a computational colonic environment. This environment resembles the crypt space together with the mucus layers, the lumen and fluid dynamics. Our proof-of-principle experiments include mono-colonisation simulations of context-specific colonic cells and simulations of context-specific colonic cells with the SIHUMIx minimal model microbiome. Our simulations propose host-microbial and microbial-microbial interactions that can be verified based on the literature. Most importantly, the Virtual Colon offers visualisation of interactions through time and space, adding another dimension to the genome-scale metabolic modelling approaches. Lastly, like BacArena, it is freely available and can be easily adapted to model other spatially structured environments (http://www.github.com/maringos/VirtualColon).

2. Introduction

Host-microbial interactions are of crucial importance for host health and fitness[1]. A well-studied example of localised host-bacterial interactions is the mammalian large intestine, where the gut microbiome provides the host with nutrients that otherwise would not be available to the host, protects it from colonisation by pathogens and interacts with its immune and nervous system[2]. For instance, some colonic bacteria can ferment fibres and other components in the colonic lumen. Subsequently, they produce short-chain fatty acids (SCFAs), such as acetate, butyrate and propionate that have diverse beneficial properties for host health[3,4]. Acetate and propionate are carried and utilised metabolically outside the intestine[5], while butyrate is used locally for energy generation[6]. However, their effect on the host is not only metabolic, as there is evidence of an interaction between SCFA and the immune system[7].

However, the case of SCFA is far from exceptional[8]. Therefore, it would be very informative to systematically explore host-microbial interactions inside the colonic microenvironment. In the colon, bacteria reside and are separated from the colonic tissue by two layers of a host-produced mixture of glycoprotein and oligosaccharides, collectively called mucus, which is located in between the lumen and the colonic epithelium[9]. This environment is characterised by the limited presence of oxygen fostering the growth of anaerobic bacteria under physiological conditions[10,11]. The bacteria consume dietary compounds that are not absorbed in the small intestine along with host-derived compounds such as mucus and bile[12]. Therefore, it is not surprising that the architecture of the colonic micro-environment offers a palette of well-studied localised host-bacterial interactions, as well as unknown ones.

Attempts to model this physicochemical environment and its phenomena in the intestine exist and are informative. For instance, Cremer and colleagues developed a mathematical, quantitative model that combines the microbes, their production of SCFA

and colonic water absorption.[13]. However, a promising method to understand novel, unexplored host-microbial interactions is the modelling of the metabolism of both host and bacterial cells. Recent progress in the field of Systems Biology has enabled the creation of models of metabolism for a variety of organisms. Specifically, starting from the genomic information of individual species[14,15], it is possible to infer the possible metabolic pathways of the respective organisms and to organise them in networks of interconnected metabolic reactions, known as genome-scale metabolic models[16]. As a further step, we can apply mathematical methods, such as linear programming, to simulate the input and output of these networks. A common mathematical approach is flux balance analysis (FBA), which optimises the input and output of the models so that an objective function such as cell growth is maximised[17].

Simple metabolic models, such as the *Escherichia coli* core model with less than a hundred reactions, can be used to understand how the central metabolism of an organism operates[18]. For more advanced research, one can utilise existing curated models (e.g., the extensive, human-associated bacterial models collection AGORA2[19]) or even create own models tailored to the respective research question by employing modelling software such as gapseq and CarveMe[15,20]. Most importantly, it is possible to extend the layers of knowledge by combining datasets from different omics and databases, also beyond the strict boundaries of genome-scale metabolic modelling (e.g, see Virtual Metabolic Human database[21], MetaNetX[22–25], KEGG[26–28]). Emanating from this basis, spatiotemporal approaches that focus on simulating not only the models but also their environment through time have been developed[29]. For instance, an advanced approach by Versluis and colleagues has simulated how the metabolism of specific gut bacteria could alter based on the changes of the environment cues[30].

In this study, we build on the metabolic modelling software BacArena, which is an agent-based framework for spatial and temporal simulations of metabolic models. Specifically, it allows the design of a chemically defined environment, where the cells acting as agents can have predefined properties (e.g., speed, chemotaxis, linear or exponential growth etc.)[16,31]. We present a virtual colon-like 2D and context-specific application of BacArena that combines the host and bacterial sides into one single simulation framework. Our implementation spatially separates bacteria and host cells by representing multiple mucus layers. Furthermore, it enables a realistic simulation of host and microbial interactions, as it combines spatial-dependent diffusion phenomena that constrain substrate availability and mimic the lumen faecal stream powered by peristalsis activity. Lastly, our Virtual Colon software is freely available on GitHub so that it can easily be adapted by future users (http://www.github.com/maringos/VirtualColon).

3. Material and methods

General setup

We conducted computational colonisation experiments mimicking an experimental setting with germ-free mice originally described by Geva-Zatorsky and colleagues (henceforward the reference study)[32]. Therefore, each simulation comprised bacterial models together with host models in our colon-like in-silico setting. Furthermore, each setting simulates seven iterations of one hour. The first iteration was used for the diffusion of compounds into the environment. In total, each setting was replicated ten

116 times. The growth of each organism was calculated based on parsimonious flux balance
117 analysis[33]. The majority of colonisation experiments referred to the introduction of one
118 species (henceforward mono-colonisation). There were also three cases of specific-
119 pathogen-free mice (SPF). In that case, we employed the generic simplified human
120 intestinal bacterial community of eight species (see Table 1, SIHUMIx)[34].

Metabolic models

In the case of the mono-colonisation experiments, bacterial metabolic models
matching the bacterial strains of the reference study were used (see S1 Table)[32]. If a
genome of a strain was not available, we took the most similar genome from NCBI and
literature (see S1 Table) [35]. The genomes were utilised to create the respective bacterial
metabolic models by gapseq (version ce962ff) [15]. The medium for gap filling was the one
available in the lumen and mucus layers of the Virtual Colon (see S1 Table). The SIHUMIx
metabolic models were reconstructed using gapseq (version 0e7af55) [15] based on the
published genomic information by Becker and colleagues[34]. The gap-filling medium was
predicted by gapseq. Finally, the models were adapted to grow on the medium available in
the lumen and mucus layers of the Virtual Colon. To minimise the possibility of bacterial
models competing for nutritional resources with the human models, we introduced a new
"d" compartment into the bacterial models to account for an exclusive nutritional uptake
of compounds from the lumen.

The host metabolic models originate from an adapted version of the published
Recon 2.2 model[36]. This generic model was expanded so that a new compartment of
exchange reactions was available. This compartment contains exchange reactions ending
with "u" and is a copy of the original compartment of exchange reactions labelled "e". The
reason for this addition was to represent the flow of compounds between the bacteria in
the lumen and the colonic cells (compartment "e") and between the colonic cells and the
blood compartment (compartment "u"), following the strategy of Sahoo and Thelie[37].
Moreover, we used iMAT to integrate mouse transcriptomic data from the reference
study[32,38] into the host models[39] to build context-specific models of the respective colonic
tissue. After translating gene expression activity into reaction expression, a reaction was
assumed to be lowly expressed if it was within the first quartile of all reaction expression
values. A reaction was assumed to be highly expressed if it was in the fourth quartile
across all reaction expression values Moreover, we required a pre-set list of forced
reactions and unconstrained nutrition-related exchange reactions. This information was
used as input to determine a context-specific model for each condition. All exchange
reactions and the demand reaction for ATP "DM atp c" were available in the final models.
Detailed information on the utilised models and the procedures can be obtained from
Table 1 and the models are available online (see S1 Table and supplementary folder
'Models').

Representation of the colon

157 The computational space of the Virtual Colon was developed by extending
158 BacArena, which provides a two-dimensional grid area[31]. One bacterial or host cell was
159 located in one grid cell. The size of the Virtual Colonic lumen, the mucus layers, and the
160 structure of the colonic crypts were calculated based on anatomic measurements

converted to the dimensions of one grid cell. Specifically, thanks to a predefined crypt-like position of the host cells in the environment, two half-crypts were represented together with their common inter-cryptal space (see Fig. 1 and Table 1).

The Virtual Colon is composed of five spaces: lumen, outer mucus, inner mucus, intermediate space, host space. The inner mucus' width was equal to one-third of the total mucus one, while the outer mucus' width was equal to two-thirds[40]. The concentration of outer mucus layer compounds was found to be four times greater than that of the outer ones[41] and was used accordingly (see S2 Table for more information on design, calculation, and further references). The bacterial models were introduced into the colonic lumen and both mucus layers. The bacteria of the inner mucus were unable to move unless the inner mucus was partially degraded. Then the bacterial models could move through the grid cell and potentially pass towards the host. The bacterial models could not enter human space, but they could be located next to the human models in the intermediate space.

We started the simulations with reported abundances from the reference study [32]. The initial position of microbial cells in the Virtual Colon was random. Important additional parameters were: the arena's size and volume, the actual total volume of the murine host's caecum, colon and rectum (0.37 cm^3)[42], the bacterial input (200µl of 10^8 cfu/ml; personal communication with Kasper Dennis ((dennis_kasper@hms.harvard.edu), correspondent author of the reference study[32]). For instance, the amount of bacteria in the lumen was calculated based on these formulas:

$$= \frac{* * 0.2 * 10^8}{}$$

$$= * * \sqrt{*}$$

Similarly, the amount of bacteria in the outer and inner mucus areas was calculated. In the case of the SIHUMIx community simulations, the same approach was followed. However, an abundance factor accounting for the relative abundances of each species was used (see Table 1).[34]

The virtual host cells inherited the default human class settings of BacArena[31]. For the minimum weight and its standard deviation, the scaling factor was used to relate the area of human and bacterial cells. Rodriguez and colleagues have suggested that one crypt column is as long as 31.1 cells in the descending mouse colon [43]. Similarly, the two-crypt setting provided 70 model slots for host cells in total. Since the maximum allowed weight of 70 cells represented a maximum total weight of around all cells of a human crypt (BNID 110648 [44]). In addition to these adjustments, no movement was allowed for the models, and their growth was considered to be linear through time (see Table 1). It should be noted that while there is some debate about whether optimisation of growth is an adequate objective function for flux balance simulations of colonic cells, the biomass function includes all molecular components of cells also required for cell renewal. Furthermore, there is constant growth and shedding of colonic cells with an estimated turnover time of 3-5 days in mice[45]. Thus, in the colonic context, optimisation of growth is a valid objective function to predict metabolic fluxes.

The transport of molecules inside BacArena [31] was modelled by diffusion (see S2 Table). As discussed above, the arena was split into the following diffusion sectors: lumen, outer mucus layer, inner mucus layer, intermediate layer in-between, and blood. The utilised diffusion coefficients were calculated using a power regression function

(coefficients vs. molecular weight) for the colonic lumen and mucus layers. The molecular
weight was used for nutritional compounds. For other compounds, the default value of
glucose was assumed. The regression functions were generated using the intestinal
diffusion data of Winne and colleagues [46]. The same strategy was followed for the
restricted areas and the lumen [47]. Especially for mucus layers, the diffusion coefficients of
molecules in outer mucus were equal to ⅔ of mucus diffusion coefficients, while for inner
mucus, they were equal to ⅓ of mucus diffusion coefficients, based on the observation
that ⅓ of the length of the mucus belongs to the inner layer [40]. Furthermore, the mucus
compounds were not allowed to diffuse to represent the stability of the mucus in the
colonic environment. Oxygen could only diffuse to host areas to ensure limited
"oxygenation" of the Virtual Colon. Practically, the diffusion of compounds in the lumen
was designed to be less restricted than in mucus layers, while not possible in the
intermediate layer and in blood. At the same time, diffusion of blood molecules was
possible in the intermediate layer and in the blood.
218

Nutrition

Following the nutrition that was chosen by the authors of the reference publication
[32], the primary source of nutritional compounds was the LabDiet ® JL Rat and Mouse/Auto
6F 5K67. These compounds were matched to the dietary bacterial exchange reactions to
exclude nutritional competition between host and bacterial cells. The intestinal absorption
and the dimensions of the Virtual Colon were considered for calculating the exact
nutritional input. In addition, some components such as water, H^+, minerals, vitamins, and
mucus compounds were added as part of the lumen compartment. To represent the inflow
of nutrients from the blood circulation, a minimal collection of compounds was added into
the blood compartment. Their concentration was based on the literature (e.g., (BNID
110365 [44]) and various Human Metabolome Database entries (https://hmdb.ca/ [48–51]). To
infer these components, a shadow cost analysis based on the BacArena [31] function
"plotShadowCost" was utilised (see supplementary folder 'Definition_Blood'). A list of
compounds can be found in the S2 Table.
233
234

235
236
237 Figure 1: Schematic representation of the virtual colonic space. The coloured rectangles
238 represent the models of the microbial species and the two shades of green represent the
239 two mucus layers above the host's epithelium. Created with BioRender.com.

Table 1: Overview of properties of the Virtual Colon

Property	Information on values & units	Further comments	Reference
Grid cell dimension	0.00025 × 0.00025 cm^2	Default in BacArena	31
Y-axis size	179-grid-cell-long	Set by the developers	Not applicable
X-axis size	24-grid cell-long	Set by the developers	NA
Scaling factor for minimum weight and its standard deviation of host models	$\dfrac{132.73229}{4.42}$	A human COLO205 colon cell has a diameter of 13 µm; hence its area is equal to 132.73229 µm^2. The area of a bacterium in BacArena is, by default, 4.42 µm^2.	44 BNID 108890
Scaling factor for maximum weight of host models	$\dfrac{1000}{70}$	70 cells represent the weight of around 1000 cells of a human crypt	43,44 BNID 110648[44]
Starting amount of bacteria in the lumen	29	Set by the developers	Not applicable
Starting amount of bacteria in the outer mucus	18	Set by the developers	Not applicable
Starting amount of bacteria in the inner mucus	9	Set by the developers	Not applicable
Relative abundance of *Anaerostipes caccae* in SIHUMIx simulations	0.1263867	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	34
Relative abundance of *Bacteroides thetaiotaomicron* in SIHUMIx simulations	0.1444797	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	34
Relative abundance of *Bifidobacterium longum* in SIHUMIx simulations	0.1388008	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	34
Relative abundance of *Blautia producta* in SIHUMIx simulations	0.1385367	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	34
Relative abundance of *Clostridium butyricum* in SIHUMIx simulations	0.1065769	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	34
Relative abundance of *Clostridium ramosum* in SIHUMIx simulations	0.1263867	Based on the concentration in human faeces [log10 cells / g dry matter]	34
Relative abundance of *Escherichia coli* in SIHUMIx simulations	0.118859	Based on the concentration in human faeces [log10 cells / g dry matter]	34
Relative abundance of *Lactobacillus plantarum* in SIHUMIx simulations	0.09997359	Based on the concentration in human faeces [log10 cells / g dry matter]	34

258 Software

The simulations and analyses thereafter presented in this manuscript were
conducted in the R environment (version 4.2.1) [52]. Optimisations were generally conducted
using the BacArena application (version 1.8.2, commit euba/BacArena@dc8e591 and
euba/BacArena@008d186) [31] and sybil (version 2.2.1, commit
SysBioChalmers/sybil@58eb989) [53]. The linear programming solver was CPLEX (academic
version 22.1.0) [54], which is available through the R package cplexAPI (version 1.4.0) [55]. For
data management, the software tidyverse (version 2.0.0) [56] and especially its library dplyr
(version 1.1.4) [57] were used. For plotting, the software ggplot2 (version 3.5.1) [58], ggvenn
(version 0.1.10)[59], ggthemes (version 5.1.0) [60], scales (version 1.3.0) [61], ggalluvial
(version 0.12.5) [62], and BacArena (version 1.8.2, commit euba/BacArena@dc8e591) [31]
were utilised. The software devtools (version 2.4.5) [63] was also used to manage the R
environment.

272 4. Results

The main objective of this study was to provide a computational framework to
simulate host-microbial interactions in the mammalian colon. We first used our approach
to study host-microbiome interactions in the colon based on a previously published
dataset of single-strain colonisation experiments in germ-free mice[32]. Besides microbial
abundance data, Geva-Zatorsky and colleagues provided extensive transcriptomic datasets
of these colonisation experiments. Based on those transcriptomes, we reconstructed
context-specific models of colonic tissue for the respective mice. For our study, we used
context-specific Recon 2.2 [36] host models along with bacterial models for joint simulations
in a virtual colon.

The Virtual Colon provides detailed information on the host-microbial metabolic
interactions in space and time. To reveal those interactions in the mono-colonisation
experiments, we extracted all fluxes across all time points for each compound and for
each model. Since a positive flux shows production and a negative one shows consumption
of a compound, we calculated the sum of fluxes of all models for each compound.
Collectively, the host models produced 20 compounds and the bacterial ones produced 27
compounds (Fig. 2A). Interestingly, the host-microbial interactions seem to be very pair-
specific, since we could identify compounds that in some pairs are bacterially produced or
in other pairs by the host (eg., glycine, L-alanine; Fig. 2A). Furthermore, we observed the
host exchange of nitrogen-containing compounds, towards the bacterial models, and the
uptake of bacterially produced SCFA. Additionally, we found that although the majority of
the bacteria interacted metabolically with the host regardless of their position (e.g., see
cases with fluxes of models originated from all three layers: lumen, outer mucus, inner
mucus, Fig. 2B), we also observed cases of localised interactions (Fig. 2B).

Further, we employed the Virtual Colon to study host-bacterial interactions with
the SIHUMIx bacterial community. As a proof-of-principle, we studied the growth patterns
of the bacterial and host models. Each bacterial strain achieved different growth rates and
all the models were still in the exponential phase at the end of simulation. The host cells
did not reproduce, as expected (Fig. 3A). Regarding the different layers of Virtual Colon,
there is a variation of growth rate of the bacterial models depending on their position in
the colonic space (Fig. 3B). Inside the inner mucus layer they could only marginally spread

in comparison to the other locations due to the design of the movement barrier. The
SIHUMIx microbial community that resides in inner and outer colonic mucus also gradually
consumed the mucus compound N-acetylneuraminate. Most interestingly, the decreasing
quantity of the compound follows the growth of the bacteria models in the area. The
bacterial models of *Bacteroides thetaiotaomicron*, *Blautia producta*, and *Clostridium*
butyricum can degrade N-acetylneuraminate. (Fig. 3C).

The observation of microbial mucus degradation led to more general questions on
what compounds are consumed or produced by the metabolic models. The host models,
similarly to the mono-colonisation experiments, are context-specific Recon 2.2 models[36],
whose reconstruction was based on the transcriptomes GSM2354402, GSM2354403 and
GSM2354404 respectively. While those three models produce and consume mostly the
same compounds, they do have distinct metabolic profiles that can be attributed to the
different transcriptomes. For instance, the recon model GSM2354403 produced L-alanine,
which was taken up by the bacteria, while the recon model GSM2354404 consumed
bacterial L-alanine (Fig. 4A and 4D). However, all models consumed bacterially produced
short-chain fatty acids, such as formate, acetate, propionate, and butyrate (Fig. 4C and 4D).
Interestingly, the models exchange various metabolic compounds (e.g., water, vitamins,
amino acids, TCA-cycle compounds such as alpha-oxoglutarate).

Not only were the bacterially produced compounds taken up by the host models,
but they were also consumed by other bacterial species of the community (Fig. 5A and 5B).
For instance, apart from amino acids and carbon sources, they also exchanged metabolites
like protons, ammonium, phosphate, and hydrogen sulphide. It is possible to characterise
the metabolic role of community members. The bacteria model of *E. coli* was the most
active species in terms of production and consumption of compounds. On the other hand,
B. longum and *C. butyricum* contributed more compounds than they consumed, while the
opposite was true for *A. caccae* and *B. producta*.

Furthermore, in order to understand how the metabolic interactions change when
comparing the interactions from mono-colonisation with the SIHUMIx simulations, we
investigated which compounds are unique in each of the two settings. It can be observed
that the exchanged compounds were mostly common (Fig. 6). However, the majority of
the unique compounds in mono-colonisation experiments were nitrogen-containing
compounds. In the other two parts of the Venn diagram, we could identify amino acids,
metabolism intermediates (e.g. fumarate) and short-chain fatty acids.

343 Figure 2: **A.** Exchange of compounds which are produced by the bacterial models or
344 context-specific host models. The colour of the layers depends on the origin of the
345 producer. **B.** Exchange of compounds which are produced by the host models and are
346 consumed by the bacterial model and vice versa. To estimate the exchange of compounds
347 among the models, the sum of all fluxes across time for each compound and for each
348 model was determined. **A** positive sum denotes production, while a negative one denotes
349 consumption. To limit the numerical artefacts, the fluxes whose absolute value was less
350 than 1E-6 were considered zero.

Figure 3: A. Growth curves for the individual bacterial and host models. The data points
originate from all the replicates of the GSM2354402 SIHUMIx simulation. The individual

lines are plotted using the "loess" smoothing function. B. Variation in growth rates (unit h^-

^{1}) by location in Virtual Colon and Species. The growth rate Data points represent all

replicates from the GSM2354402 SIHUMIx simulation. C. Schematic representation of the

mucus compound N-acetylneuraminate through three-time points in a SIHUMIx-host

simulation (GSM2354402, 10th replicate). Each subplot represents the substance

concentration and bacterial population per time point in hours.

361
362

363   Figure 4: Schematic representation of compounds produced by the context-specific host
364   models (A) and consumed by the SIHUMIx bacteria (B) along with the bacterially produced

365 ones (C) that are consumed by the host models (D). Metabolites with the highest
366 interaction frequency (>= 50%) across all time points and simulations were selected to be
367 plotted for simplicity. For the full dataset, see S3 Table.

368
369 Figure 5: Schematic representation of compounds produced by the SIHUMIx bacterial
370 models (A) and consumed by other SIHUMIx bacterial models (B). Metabolites with the
371 highest interaction frequency (>= 50%) across all time points and simulations were
372 selected to be plotted for simplicity. For the full dataset, see S3 Table.
373

375 Figure 6: Schematic representation of compounds exchanged by the bacterial models and
376 host models in the mono-colonisation and SIHUMIx experiments. Only interactions with
377 bacteria from the shared genera were taken into account (Bacteroides, Bifidobacterium,
378 Clostridium, Lactobacillus). All simulation time points were included in the analysis.

Comparison of Virtual Colon with other published approaches

e	Approach	Microbes	Scope	Host	Space	Time	Mucus	Gut motility	Availability	Reference
biota	PDE model with fluid dynamics (Stokes)	Movement + predefined functions	Colon	Nutrient uptake	✓	✓	✓	✓	Matlab code https://forgemia.inra.fr/simon.labarthe/gut-microbiota	64
Pop	ODE model (Monod equation)	predefined functional groups	Colon	-	-	✓	✓	✓	R package https://github.com/HelenKettle/microPopGutCode	65
	ABM		Ileum	-	✓	✓	✓	✓	https://github.com/GutLogo/GutLogo	66
Gut he	ODE model (gLV)	25 microbes	Butyrate production	-	-	✓	-	-	Julia code https://github.com/RyanLincolnClark/DesignSyntheticGutMicrobiomeAssemblyFunction	67
lon	ABM + PDE	Movement + metabolic model	Colon	Metabolic model	✓	✓	✓	✓	R code http://www.github.com/maringos/VirtualColon	this article

al differential equation; ODE: Ordinary differential equations; gLV: generalised Lotka-Volterra; ABM: Agent-based modelling

5. Discussion

Host-bacterial metabolic interactions are considered important drivers of host
health and fitness. Therefore, one of the objectives of this study was to develop a data-
driven approach that allows us to systematically explore the compounds that are
exchanged between the host and bacterial cells. Here we present the Virtual Colon that
extends the agent-based modelling framework BacArena[16,31] to study host-microbiome
interactions in the mammalian large intestine. Specifically, it enables host-microbial
simulations, in which bacteria and host cells are spatially separated through multilayer
mucus layers. Moreover, it includes spatial-dependent diffusion that constrains substrate
availability for the models. Most importantly, it not only enables the joint simulation of
host and microbial interactions but also visualises them. While other methods have
contributed to the field either by using differential equations and/or agent-based

modelling (Table 2), the Virtual Colon, which is built on BacArena, combines them in one
software framework.

In our simulations, we utilised host models that originated from the human model
Recon 2.2[36]. For the simulations, the host models were adapted using transcriptomic
data[38] from mono-colonisation experiments of germ-free mice with the respective
species[32]. In the case of colonisation with non-specific pathogens, we used the SIHUMIx
bacterial models that comprise a generic eight-species human intestinal bacterial
community[34].

To continue with the host-bacterial interactions in the mono-colonisation
experiments, the observed usage of cysteine by some bacteria of the genus *Fusobacterium*
and *Clostridium* is in line with the literature, which proposes that cysteine is metabolised
by bacteria, leading to hydrogen sulphide[68]. Hydrogen sulphide has been characterised as
a compound with multiple effects, as in low levels, it supports colonic health, while in
excess, it is associated with inflammatory and pathological states (e.g., inflammatory
bowel disease and cancer)[68].

Additionally, the compound 4-aminobutanoate, also known as γ-aminobutyric acid
(GABA), is predicted to be produced by *Lactobacillus rhamnosus* and *Bacteroides finegoldi*
and be consumed by the host models, which is in accordance with the literature[69]. This
compound may act on the central nervous system as well as the enteric nervous system of
the host as part of the microbiota-gut-brain axis[69]. On the other hand, *Bacteroides*
finegoldi has not been reported to be a GABA producer. However, a recent article
proposed that the Bacteroides species can produce this molecule, although, in the
simulations, no other *Bacteroides* species did so[70].

Recently, a study based on metabolomics focusing on the colonic absorption of
compounds with low molecular weight in mice identified some of the computationally
predicted compounds, which were subject to metabolic exchange in our simulations:
(glycine) betaine, choline, alanine, glutamine, glycine, isoleucine, lactate, ornithine, and
short-chain fatty acids (Fig. 2A, B) [71]. Interestingly, host-derived glycine betaine and
choline can be further bacterially metabolised to trimethylamine-N-oxide and
trimethylamine, for which a linkage to atherosclerotic cardiovascular disease has been
demonstrated[72].

While mono-colonisation experiments offered a detailed overview of pairwise host-
bacterial interactions, the bacterial species found in the mammalian colon are quite
diverse. Therefore, we selected the representative SIHUMIx community to investigate how
our results change in the context of a more complex community. Our simulations showed
a wide exchange of compounds among the species as well as between bacteria and host
cells. Not surprisingly, many byproducts were found also in mono-colonisations, implying
that the repertoire of metabolic exchanges may be genera-dependent (Fig 6). The
bacterial models create a complex web of compounds, where carbohydrate and amino acid
metabolism as well as cofactors (e.g. vitamins) and byproducts are the most common end
products. This observation becomes intriguing, if one considers that carbohydrates rather
than amino acids are primarily used in bacterial energy metabolism, as amino acid
metabolism is less cost-effective[73]. Moreover, this finding suggests that mono-
colonisation experiments only partially capture interactions that can be observed in
community colonisation, as the metabolic repertoire and signature may alter significantly.

It must be stated that the predicted exchange of compounds between the host and
the bacterial models is based on the assumption that the host colonic cells can exchange
compounds from the luminal side of the colon. For instance, amino acids are known to be
absorbed in the small intestine, while their absorption fate in the large intestine is unclear.
While the colonic bacteria can metabolise the remaining amino acids, it is proposed that
some of them can be used by colonocytes [74]. In parallel, it has been shown that
colonocytes can excrete amino acids (e.g., alanine, aspartate, asparagine, isoleucine, and
valine) in both apical and basolateral directions in vitro [75]. Therefore, the observed
production or consumption of several compounds by the host is feasible and waits for
further experimental validation. These experiments should investigate the transport of
metabolites across the intestinal barrier. Such a pivotal set of experiments was performed
by Li and colleagues, who identified gut species that deplete and metabolise amino acids
throughout the intestine and therefore regulate host metabolism [76].

Furthermore, the Virtual Colon includes spatial parameters tailored to the large
intestinal environment. For instance, the diffusion of compounds inside the mucus layer is
limited and the bacteria in the inner mucus cannot move unless mucus is consumed by
community members, such as *B. thetaiotaomicron*, a known mucus-degrading species [77](Fig
3). Additionally, our results proposed that there are spatial, localised interactions,
therefore it could be suggested that the mucus layers can affect bacterial metabolism. (Fig
2B). In line with those observations, the growth rates across all bacteria in the outer mucus
layer seem to be the ones in the lumen, while the rates in the inner mucus layer are the
slowest (Fig 3B). Therefore, the simulation data suggests that our mucus both resembles
the function of the mucus as a physical barrier and to some extent as a nutrient pool for
the bacteria [9].

While the setting of the Virtual Colon recapitulates some of the observed host-
microbial functions, there is also room for further expansions in the simulation setting.
Future computational experiments may include more complex bacterial communities
along with an improved representation of the abundance of bacteria with respect to the
size of the simulated colon (e.g., the ratio of microbial abundance over the number of host
models). Currently, the virtual gut represents the host model as the same size as bacterial
models due to the technical limitations of BacArena. Another improvement could include
metabolomic data from colonic tissue and lumen so that the chemical environment is as
realistic as possible. Additionally, single-cell transcriptomics would allow the
representation of the various cell types of the host tissue. To improve the realism of the
simulations, the Virtual Colon also has the capacity to mimic the lumen faeces stream
powered by peristalsis activities, to mimic the replenishment of mucus compounds
through time or to initiate simulation assuming a non-empty colonic environment. Also it
would be interesting to investigate whether the variability observed in our simulations
recapitulates inter-individual variances in microbiome composition in mice.

In summary, our novel simulation framework, Virtual Colon, builds upon the
popular BacArena platform to create a comprehensive tool for studying host-microbe
interactions in the colon. By spatially separating bacteria and host cells and incorporating
multilayer mucus representation, the Virtual Colon enables the realistic simulation of host-
microbial interactions. This is achieved through a spatial-dependent diffusion mechanism
that constrains substrate availability and accurately mimics the faecal stream.
Furthermore, the Virtual Colon allows for the joint simulation of host and microbial
interactions, which we demonstrated for human gut microbiome communities and

personalised host models using omics data. Notably, we employed the Virtual Colon to identify species-specific mucus layer degradation and amino acid exchange between microbes and host, providing valuable insights into the intricate dynamics of the gut ecosystem. Our freely available software on GitHub invites future researchers to adapt and expand its capabilities to advance host-microbe interaction studies.

6. Authors' Contribution

Conceptualisation: GM, JZ

Methodology: GM, JZ, JT, CK

Software: GM, JZ

Formal analysis: GM

Investigation: GM, JZ, JT

Resources: CK

Data Curation: GM

Writing - Original Draft: GM, JZ

Writing - Review & Editing: All co-authors

Visualisation: GM

Supervision: CK, JZ

Project administration: GM

Funding acquisition: CK

7. Acknowledgements

CK acknowledges support from the German Research Foundation for the support through the collaborative research centre 1182 Origin and Function of Metaorganisms (sub-project A1), the research unit miTarget (FOR5042), and the excellence cluster "Precision medicine in chronic inflammation" (EXC2167) and additionally the German Ministry for Education and Research (E:Med iTREAT, support code 01ZX1902A). This research was supported in part through high-performance computing resources available at the Kiel University Computing Centre. The authors thank Dr Ahmed Samer Kadib Alban for providing them with critical opinions on the manuscript.

8. Conflict of Interest

There is nothing to disclose.

9. Supplementary Information

The software is freely available on GitHub (www.github.com/maringos/VirtualColon, commit: 0e56bda). The simulation RDS files that are related to this manuscript can be found on Figshare (www.doi.org/10.6084/m9.figshare.25997962). The supporting information including figures, scripts and datasets related to this manuscript can be found on Figshare (www.doi.org/10.6084/m9.figshare.25998073).

10.　References

1. Ruuskanen, M.O., Sommeria-Klein, G., Havulinna, A.S., Niiranen, T.J., and Lahti, L. (2021). Modelling spatial patterns in host-associated microbial communities. Env. Microbiol *23*, 2374–2388. https://doi.org/10.1111/1462-2920.15462.

2. Hou, K., Wu, Z.-X., Chen, X.-Y., Wang, J.-Q., Zhang, D., Xiao, C., Zhu, D., Koya, J.B., Wei, L., Li, J., et al. (2022). Microbiota in health and diseases. Signal Transduct. Target. Ther. *7*, 135. https://doi.org/10.1038/s41392-022-00974-4.

3. Morrison, D.J., and Preston, T. (2016). Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. Gut Microbes *7*, 189–200. https://doi.org/10.1080/19490976.2015.1134082.

4. Shimizu, H., Masujima, Y., Ushiroda, C., Mizushima, R., Taira, S., Ohue-Kitano, R., and Kimura, I. (2019). Dietary short-chain fatty acid intake improves the hepatic metabolic condition via FFAR3. Sci Rep *9*, 16574. https://doi.org/10.1038/s41598-019-53242-x.

5. Van Treuren, W., and Dodd, D. (2020). Microbial Contribution to the Human Metabolome: Implications for Health and Disease. Annu. Rev. Pathol. Mech. Dis. *15*, 345–369. https://doi.org/10.1146/annurev-pathol-020117-043559.

6. Donohoe, D.R., Garge, N., Zhang, X., Sun, W., O'Connell, T.M., Bunger, M.K., and Bultman, S.J. (2011). The Microbiome and Butyrate Regulate Energy Metabolism and Autophagy in the Mammalian Colon. Cell Metab. *13*, 517–526. https://doi.org/10.1016/j.cmet.2011.02.018.

7. Yao, Y., Cai, X., Fei, W., Ye, Y., Zhao, M., and Zheng, C. (2022). The role of short-chain fatty acids in immunity, inflammation and metabolism. Crit. Rev. Food Sci. Nutr. *62*, 1–12. https://doi.org/10.1080/10408398.2020.1854675.

8. Fogelson, K.A., Dorrestein, P.C., Zarrinpar, A., and Knight, R. (2023). The Gut Microbial Bile Acid Modulation and Its Relevance to Digestive Health and Diseases. Gastroenterology *164*, 1069–1085. https://doi.org/10.1053/j.gastro.2023.02.022.

9. Schroeder, B.O. (2019). Fight them or feed them: how the intestinal mucus layer manages the gut microbiota. Gastroenterol Rep *7*, 3–12. https://doi.org/10.1093/gastro/goy052.

10. Konjar, Š., Pavšič, M., and Veldhoen, M. (2021). Regulation of Oxygen Homeostasis at the Intestinal Epithelial Barrier Site. Int J Mol Sci *22*. https://doi.org/10.3390/ijms22179170.

11. Litvak, Y., Byndloss, M.X., and Bäumler, A.J. (2018). Colonocyte metabolism shapes the gut microbiota. Science *362*. https://doi.org/10.1126/science.aat9076.

12. O'Keefe, S.J.D. (2016). Diet, microorganisms and their metabolites, and colon cancer. Nat Rev Gastroenterol Hepatol *13*, 691–706. https://doi.org/10.1038/nrgastro.2016.165.

13. Cremer, J., Arnoldini, M., and Hwa, T. (2017). Effect of water flow and chemical environment on microbiota growth and composition in the human colon. Proc. Natl. Acad. Sci. *114*, 6438–6443. https://doi.org/10.1073/pnas.1619598114.

14. Mendoza, S.N., Olivier, B.G., Molenaar, D., and Teusink, B. (2019). A systematic assessment of current genome-scale metabolic reconstruction tools. Genome Biol *20*, 158. https://doi.org/10.1186/s13059-019-1769-1.

15. Zimmermann, J., Kaleta, C., and Waschina, S. (2021). gapseq: informed prediction of bacterial metabolic pathways and reconstruction of accurate metabolic models. Genome Biol *22*, 81. https://doi.org/10.1186/s13059-021-02295-1.

16. Jansma, J., and El Aidy, S. (2021). Understanding the host-microbe interactions using metabolic modeling. Microbiome *9*, 16. https://doi.org/10.1186/s40168-020-00955-1.

17. Orth, J.D., Thiele, I., and Palsson, B.Ø. (2010). What is flux balance analysis? Nat Biotechnol *28*, 245–248. https://doi.org/10.1038/nbt.1614.

18. Orth, J.D., Fleming, R.M.T., and Palsson, B.Ø. (2010). Reconstruction and Use of Microbial Metabolic Networks: the Core *Escherichia coli* Metabolic Model as an Educational Guide. EcoSal Plus *4*, 10.1128/ecosalplus.10.2.1.

https://doi.org/10.1128/ecosalplus.10.2.1.
19. Heinken, A., Hertel, J., Acharya, G., Ravcheev, D.A., Nyga, M., Okpala, O.E., Hogan, M.,
Magnúsdóttir, S., Martinelli, F., Nap, B., et al. (2023). Genome-scale metabolic
reconstruction of 7,302 human microorganisms for personalized medicine. Nat.
Biotechnol. *41*, 1320–1331. https://doi.org/10.1038/s41587-022-01628-0.
20. Machado, D., Andrejev, S., Tramontano, M., and Patil, K.R. (2018). Fast automated
reconstruction of genome-scale metabolic models for microbial species and
communities. Nucleic Acids Res. *46*, 7542–7553. https://doi.org/10.1093/nar/gky537.
21. Noronha, A., Modamio, J., Jarosz, Y., Guerard, E., Sompairac, N., Preciat, G.,
Daníelsdóttir, A.D., Krecke, M., Merten, D., Haraldsdóttir, H.S., et al. (2019). The Virtual
Metabolic Human database: integrating human and gut microbiome metabolism with
nutrition and disease. Nucleic Acids Res. *47*, D614–D624.
https://doi.org/10.1093/nar/gky992.
22. Ganter, M., Bernard, T., Moretti, S., Stelling, J., and Pagni, M. (2013). MetaNetX.org: a
website and repository for accessing, analysing and manipulating metabolic networks.
Bioinformatics *29*, 815–816. https://doi.org/10.1093/bioinformatics/btt036.
23. Bernard, T., Bridge, A., Morgat, A., Moretti, S., Xenarios, I., and Pagni, M. (2014).
Reconciliation of metabolites and biochemical reactions for metabolic networks. Brief.
Bioinform. *15*, 123–135. https://doi.org/10.1093/bib/bbs058.
24. Moretti, S., Martin, O., Van Du Tran, T., Bridge, A., Morgat, A., and Pagni, M. (2016).
MetaNetX/MNXref – reconciliation of metabolites and biochemical reactions to bring
together genome-scale metabolic networks. Nucleic Acids Res. *44*, D523–D526.
https://doi.org/10.1093/nar/gkv1117.
25. Moretti, S., Tran, V.D.T., Mehl, F., Ibberson, M., and Pagni, M. (2021).
MetaNetX/MNXref: unified namespace for metabolites and biochemical reactions in
the context of metabolic models. Nucleic Acids Res. *49*, D570–D574.
https://doi.org/10.1093/nar/gkaa992.
26. Kanehisa, M. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Res. *28*, 27–30. https://doi.org/10.1093/nar/28.1.27.
27. Kanehisa, M. (2019). Toward understanding the origin and evolution of cellular
organisms. Protein Sci. *28*, 1947–1951. https://doi.org/10.1002/pro.3715.
28. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M., and Ishiguro-Watanabe, M. (2023).
KEGG for taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. *51*,
D587–D592. https://doi.org/10.1093/nar/gkac963.
29. Scott, W.T., Benito-Vaquerizo, S., Zimmermann, J., Bajić, D., Heinken, A., Suarez-Diez,
M., and Schaap, P.J. (2023). A structured evaluation of genome-scale constraint-based
modeling tools for microbial consortia. PLOS Comput. Biol. *19*, e1011363.
https://doi.org/10.1371/journal.pcbi.1011363.
30. Versluis, D.M., Schoemaker, R., Looijesteijn, E., Muysken, D., Jeurink, P.V., Paques, M.,
Geurts, J.M.W., and Merks, R.M.H. (2022). A Multiscale Spatiotemporal Model Including
a Switch from Aerobic to Anaerobic Metabolism Reproduces Succession in the Early
Infant Gut Microbiota. mSystems *7*, e00446-22.
https://doi.org/10.1128/msystems.00446-22.
31. Bauer, E., Zimmermann, J., Baldini, F., Thiele, I., and Kaleta, C. (2017). BacArena:
Individual-based metabolic modeling of heterogeneous microbes in complex
communities. PLoS Comput Biol *13*, e1005544.
https://doi.org/10.1371/journal.pcbi.1005544.
32. Geva-Zatorsky, N., Sefik, E., Kua, L., Pasman, L., Tan, T.G., Ortiz-Lopez, A., Yanortsang,
T.B., Yang, L., Jupp, R., Mathis, D., et al. (2017). Mining the Human Gut Microbiota for
Immunomodulatory Organisms. Cell *168*, 928-943.e11.
https://doi.org/10.1016/j.cell.2017.01.022.
33. Lewis, N.E., Hixson, K.K., Conrad, T.M., Lerman, J.A., Charusanti, P., Polpitiya, A.D.,
Adkins, J.N., Schramm, G., Purvine, S.O., Lopez‐Ferrer, D., et al. (2010). Omic data from
evolved *E. coli* are consistent with computed optimal growth from genome‐scale
models. Mol. Syst. Biol. *6*, 390. https://doi.org/10.1038/msb.2010.47.

34. Becker, N., Kunath, J., Loh, G., and Blaut, M. (2011). Human intestinal microbiota:
Characterization of a simplified and stable gnotobiotic rat model. Gut Microbes *2*, 25–
33. https://doi.org/10.4161/gmic.2.1.14651.
35. Fritzenwanker, M., Hain, T., Kesper, D.A., Harb, H., Renz, H., and Domann, E. (2016).
Draft Genome Sequence and Complete Plasmid Sequence of *Acinetobacter lwoffii* F78,
an Isolate with Strong Allergy-Protective Properties. Genome Announc. *4*, e00685-16.
https://doi.org/10.1128/genomeA.00685-16.
36. Swainston, N., Smallbone, K., Hefzi, H., Dobson, P.D., Brewer, J., Hanscho, M., Zielinski,
D.C., Ang, K.S., Gardiner, N.J., Gutierrez, J.M., et al. (2016). Recon 2.2: from
reconstruction to model of human metabolism. Metabolomics *12*, 109.
https://doi.org/10.1007/s11306-016-1051-4.
37. Sahoo, S., and Thiele, I. (2013). Predicting the impact of diet and enzymopathies on
human small intestinal epithelial cells. Hum. Mol. Genet. *22*, 2705–2722.
https://doi.org/10.1093/hmg/ddt119.
38. Gene Expression Omnibus Accession Viewer GSE88919.
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi? acc=GSE88919
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi? acc=GSE88919.
39. Zur, H., Ruppin, E., and Shlomi, T. (2010). iMAT: an integrative metabolic analysis tool.
Bioinformatics *26*, 3140–3142. https://doi.org/10.1093/bioinformatics/btq602.
40. Gustafsson, J.K., Ermund, A., Johansson, M.E.V., Schütte, A., Hansson, G.C., and Sjövall,
H. (2012). An ex vivo method for studying mucus formation, properties, and thickness
in human colonic biopsies and mouse small and large intestinal explants. Am. J.
Physiol.-Gastrointest. Liver Physiol. *302*, G430–G438.
https://doi.org/10.1152/ajpgi.00405.2011.
41. Johansson, M.E.V., Phillipson, M., Petersson, J., Velcich, A., Holm, L., and Hansson, G.C.
(2008). The inner of the two Muc2 mucin-dependent mucus layers in colon is devoid of
bacteria. Proc. Natl. Acad. Sci. *105*, 15064–15069.
https://doi.org/10.1073/pnas.0803124105.
42. Casteleyn, C., Rekecki, A., Van Der Aa, A., Simoens, P., and Van Den Broeck, W. (2010).
Surface area assessment of the murine intestinal tract as a prerequisite for oral dose
translation from mouse to man. Lab. Anim. *44*, 176–183.
https://doi.org/10.1258/la.2009.009112.
43. De Rodriguez, M.S.B., Sunter, J.P., Watson, A.J., Wright, N.A., and Appleton, D.R.
(1978). Cell population kinetics in the mucosal crypts of the descending colon of the
mouse. Virchows Arch. B Cell Pathol. *29*, 351–361.
https://doi.org/10.1007/BF02899366.
44. Milo, R., Jorgensen, P., Moran, U., Weber, G., and Springer, M. (2010). BioNumbers—the
database of key numbers in molecular and cell biology. Nucleic Acids Res. *38*, D750–
D753. https://doi.org/10.1093/nar/gkp889.
45. Park, J., Kotani, T., Konno, T., Setiawan, J., Kitamura, Y., Imada, S., Usui, Y., Hatano, N.,
Shinohara, M., Saito, Y., et al. (2016). Promotion of Intestinal Epithelial Cell Turnover by
Commensal Bacteria: Role of Short-Chain Fatty Acids. PLOS ONE *11*, e0156334.
https://doi.org/10.1371/journal.pone.0156334.
46. Winne, D., and Verheyen, W. (1990). Diffusion coefficient in native mucus gel of rat
small intestine. J. Pharm. Pharmacol. *42*, 517–519. https://doi.org/10.1111/j.2042-
7158.1990.tb06611.x.
47. Pappenheimer, J.R., and Reiss, K.Z. (1987). Contribution of solvent drag through
intercellular junctions to absorption of nutrients by the small intestine of the rat. J.
Membr. Biol. *100*, 123–136. https://doi.org/10.1007/BF02209145.
48. Wishart, D.S., Tzur, D., Knox, C., Eisner, R., Guo, A.C., Young, N., Cheng, D., Jewell, K.,
Arndt, D., Sawhney, S., et al. (2007). HMDB: the Human Metabolome Database. Nucleic
Acids Res. *35*, D521–D526. https://doi.org/10.1093/nar/gkl923.
49. Wishart, D.S., Knox, C., Guo, A.C., Eisner, R., Young, N., Gautam, B., Hau, D.D.,
Psychogios, N., Dong, E., Bouatra, S., et al. (2009). HMDB: a knowledgebase for the
human metabolome. Nucleic Acids Res. *37*, D603–D610.

https://doi.org/10.1093/nar/gkn810.

50. Wishart, D.S., Jewison, T., Guo, A.C., Wilson, M., Knox, C., Liu, Y., Djoumbou, Y., Mandal,
R., Aziat, F., Dong, E., et al. (2012). HMDB 3.0—The Human Metabolome Database in
2013. Nucleic Acids Res. *41*, D801–D807. https://doi.org/10.1093/nar/gks1065.

51. Wishart, D.S., Feunang, Y.D., Marcu, A., Guo, A.C., Liang, K., Vázquez-Fresno, R., Sajed,
T., Johnson, D., Li, C., Karu, N., et al. (2018). HMDB 4.0: the human metabolome
database for 2018. Nucleic Acids Res. *46*, D608–D617.
https://doi.org/10.1093/nar/gkx1089.

52. R Core Team R: A Language and Environment for Statistical Computing. (R Foundation
for Statistical Computing).

53. Gelius-Dietrich, G., Desouki, A.A., Fritzemeier, C.J., and Lercher, M.J. (2013). sybil –
Efficient constraint-based modelling in R. BMC Syst. Biol. *7*, 125.
https://doi.org/10.1186/1752-0509-7-125.

54. IBM IBM ILOG CPLEX Optimization Studio.

55. Gelius-Dietrich, G. cplexAPI: R Interface to C API of IBM ILOG CPLEX.

56. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the Tidyverse. J. Open Source
Softw. *4*, 1686. https://doi.org/10.21105/joss.01686.

57. Wickham, H., François, R., Henry, L., Müller, K., and Vaughan, D. dplyr: A Grammar of
Data Manipulation.

58. Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer-Verlag New
York).

59. Linlin, Y. ggvenn: Draw Venn Diagram by "ggplot2."

60. Arnold, J.B. ggthemes: Extra Themes, Scales and Geoms for "ggplot2."

61. Wickham, H., and Seidel, D. scales: Scale Functions for Visualization.

62. Brunson, J. (2020). ggalluvial: Layered Grammar for Alluvial Plots. J. Open Source
Softw. *5*, 2017. https://doi.org/10.21105/joss.02017.

63. Wickham, H., Hester, J., Chang, W., and Bryan, J. devtools: Tools to Make Developing R
Packages Easier.

64. Labarthe, S., Polizzi, B., Phan, T., Goudon, T., Ribot, M., and Laroche, B. (2019). A
mathematical model to investigate the key drivers of the biogeography of the colon
microbiota. J. Theor. Biol. *462*, 552–581. https://doi.org/10.1016/j.jtbi.2018.12.009.

65. Kettle, H., Louis, P., and Flint, H.J. (2022). Process-based modelling of microbial
community dynamics in the human colon. J. R. Soc. Interface *19*, 20220489.
https://doi.org/10.1098/rsif.2022.0489.

66. Lin, C., Culver, J., Weston, B., Underhill, E., Gorky, J., and Dhurjati, P. (2018). GutLogo:
Agent-based modeling framework to investigate spatial and temporal dynamics in the
gut microbiome. PLOS ONE *13*, e0207072.
https://doi.org/10.1371/journal.pone.0207072.

67. Clark, R.L., Connors, B.M., Stevenson, D.M., Hromada, S.E., Hamilton, J.J., Amador-
Noguez, D., and Venturelli, O.S. (2021). Design of synthetic human gut microbiome
assembly and butyrate production. Nat. Commun. *12*, 3254.
https://doi.org/10.1038/s41467-021-22938-y.

68. Blachier, F., Beaumont, M., and Kim, E. (2019). Cysteine-derived hydrogen sulfide and
gut health: a matter of endogenous or bacterial origin. Curr. Opin. Clin. Nutr. Metab.
Care *22*, 68–75. https://doi.org/10.1097/MCO.0000000000000526.

69. Yong, S.J., Tong, T., Chew, J., and Lim, W.L. (2020). Antidepressive Mechanisms of
Probiotics and Their Therapeutic Potential. Front. Neurosci. *13*, 1361.
https://doi.org/10.3389/fnins.2019.01361.

70. Otaru, N., Ye, K., Mujezinovic, D., Berchtold, L., Constancias, F., Cornejo, F.A., Krzystek,
A., De Wouters, T., Braegger, C., Lacroix, C., et al. (2021). GABA Production by Human
Intestinal Bacteroides spp.: Prevalence, Regulation, and Role in Acid Stress Tolerance.
Front. Microbiol. *12*, 656895. https://doi.org/10.3389/fmicb.2021.656895.

71. Matsumoto, M., Ooga, T., Kibe, R., Aiba, Y., Koga, Y., and Benno, Y. (2017). Colonic
Absorption of Low-Molecular-Weight Metabolites Influenced by the Intestinal

Microbiome: A Pilot Study. PLOS ONE *12*, e0169207.
https://doi.org/10.1371/journal.pone.0169207.
72. Canyelles, M., Borràs, C., Rotllan, N., Tondo, M., Escolà-Gil, J.C., and Blanco-Vaca, F.
(2023). Gut Microbiota-Derived TMAO: A Causal Factor Promoting Atherosclerotic
Cardiovascular Disease? Int. J. Mol. Sci. *24*, 1940.
https://doi.org/10.3390/ijms24031940.
73. Oliphant, K., and Allen-Vercoe, E. (2019). Macronutrient metabolism by the human gut
microbiome: major fermentation by-products and their impact on host health.
Microbiome *7*, 91. https://doi.org/10.1186/s40168-019-0704-8.
74. Ma, N., and Ma, X. (2019). Dietary Amino Acids and the Gut☐Microbiome☐Immune Axis:
Physiological Metabolism and Therapeutic Prospects. Compr. Rev. Food Sci. Food Saf.
18, 221–242. https://doi.org/10.1111/1541-4337.12401.
75. Chen, Y., Dinges, M.M., Green, A., Cramer, S.E., Larive, C.K., and Lytle, C. (2020).
Absorptive transport of amino acids by the rat colon. Am. J. Physiol.-Gastrointest. Liver
Physiol. *318*, G189–G202. https://doi.org/10.1152/ajpgi.00277.2019.
76. Li, T.-T., Chen, X., Huo, D., Arifuzzaman, M., Qiao, S., Jin, W.-B., Shi, H., Li, X.V., Iliev, I.D.,
Artis, D., et al. (2024). Microbiota metabolism of intestinal amino acids impacts host
nutrient homeostasis and physiology. Cell Host Microbe *32*, 661-675.e10.
https://doi.org/10.1016/j.chom.2024.04.004.
77. Kostopoulos, I., Aalvink, S., Kovatcheva-Datchary, P., Nijsse, B., Bäckhed, F., Knol, J., De
Vos, W.M., and Belzer, C. (2021). A Continuous Battle for Host-Derived Glycans
Between a Mucus Specialist and a Glycan Generalist in vitro and in vivo. Front.
Microbiol. *12*, 632454. https://doi.org/10.3389/fmicb.2021.632454.
780