═══ R E V I E W C O M M O N S M A N U S C R I P T ═══

IMPORTANT:

- Manuscripts submitted to Review Commons are peer reviewed in a journal-agnostic way.
- Upon transfer of the peer reviewed preprint to a journal, the referee reports will be available in full to the handling editor.
- The identity of the referees will NOT be communicated to the authors unless the reviewers choose to sign their report.
- The identity of the referee will be confidentially disclosed to any affiliate journals to which the manuscript is transferred.

GUIDELINES:

- For reviewers: https://www.reviewcommons.org/reviewers
- For authors: https://www.reviewcommons.org/authors

CONTACT:

The Review Commons office can be contacted directly at: office@reviewcommons.org

Virtual Colon: Spatiotemporal modelling of metabolic interactions in a computational colonic environment

Georgios Marinos[1,*], Johannes Zimmermann[1,2,3*], Jan Taubenheim[1], and Christoph Kaleta[1,$]

[1] Research Group Medical Systems Biology, University Hospital Schleswig-Holstein Campus Kiel, Kiel University, Kiel, Schleswig-Holstein, Germany

[2] Antibiotic resistance group, Max Planck Institute for Evolutionary Biology, Ploen, Schleswig-Holstein, Germany

[3] Evolutionary Ecology and Genetics, Zoological Institute, Kiel University, Kiel, Schleswig-Holstein, Germany

*Shared first authorship with an interchangeable order

$Senior and corresponding author: c.kaleta@iem.uni-kiel.de

Georgios Marinos ORCID 0000-0002-6443-7696

Johannes Zimmermann ORCID 0000-0002-5041-1954

Jan Taubenheim ORCID 0000-0001-7283-1768

Christoph Kaleta ORCID 0000-0001-8004-9514

1. Abstract

Host-microbial metabolic interactions have been recognised as an essential factor in host health and disease. Genome-scale metabolic modelling approaches have made important contributions to our understanding of the interactions in such communities. One particular such modelling approach is BacArena in which metabolic models grow, reproduce, and interact as independent agents in a spatiotemporal metabolic environment. Here, we present a modelling application of BacArena, a virtual colonic environment, which reveals spatiotemporal metabolic interactions in a computational colonic environment. This environment resembles the crypt space together with the mucus layers, the lumen and fluid dynamics. Our proof-of-principle experiments include mono-colonisation simulations of context-specific colonic cells and simulations of context-specific colonic cells with the SIHUMIx minimal model microbiome. Our simulations propose host-microbial and microbial-microbial interactions that can be verified based on the literature. Most importantly, the Virtual Colon offers visualisation of interactions through time and space, adding another dimension to the genome-scale metabolic modelling approaches. Lastly, like BacArena, it is freely available and can be easily adapted to model other spatially structured environments (http://www.github.com/maringos/VirtualColon).

2. Introduction

Host-microbial interactions are of crucial importance for host health and fitness[1]. A well-studied example of localised host-bacterial interactions is the mammalian large intestine, where the gut microbiome provides the host with nutrients that otherwise would not be available to the host, protects it from colonisation by pathogens and interacts with its immune and nervous system[2]. For instance, some colonic bacteria can ferment fibres and other components in the colonic lumen. Subsequently, they produce short-chain fatty acids (SCFAs), such as acetate, butyrate and propionate that have diverse beneficial properties for host health[3,4]. Acetate and propionate are carried and utilised metabolically outside the intestine[5], while butyrate is used locally for energy generation[6]. However, their effect on the host is not only metabolic, as there is evidence of an interaction between SCFA and the immune system[7].

However, the case of SCFA is far from exceptional[8]. Therefore, it would be very informative to systematically explore host-microbial interactions inside the colonic microenvironment. In the colon, bacteria reside and are separated from the colonic tissue by two layers of a host-produced mixture of glycoprotein and oligosaccharides, collectively called mucus, which is located in between the lumen and the colonic epithelium[9]. This environment is characterised by the limited presence of oxygen fostering the growth of anaerobic bacteria under physiological conditions[10,11]. The bacteria consume dietary compounds that are not absorbed in the small intestine along with host-derived compounds such as mucus and bile[12]. Therefore, it is not surprising that the architecture of the colonic micro-environment offers a palette of well-studied localised host-bacterial interactions, as well as unknown ones.

Attempts to model this physicochemical environment and its phenomena in the intestine exist and are informative. For instance, Cremer and colleagues developed a mathematical, quantitative model that combines the microbes, their production of SCFA and colonic water absorption.[13]. However, a promising method to understand novel, unexplored host-microbial interactions is the modelling of the metabolism of both host and bacterial cells.

Recent progress in the field of Systems Biology has enabled the creation of models of
metabolism for a variety of organisms. Specifically, starting from the genomic information of
individual species[14,15], it is possible to infer the possible metabolic pathways of the respective
organisms and to organise them in networks of interconnected metabolic reactions, known as
genome-scale metabolic models[16]. As a further step, we can apply mathematical methods,
such as linear programming, to simulate the input and output of these networks. A common
mathematical approach is flux balance analysis (FBA), which optimises the input and output
of the models so that an objective function such as cell growth is maximised[17].

Simple metabolic models, such as the *Escherichia coli* core model with less than a
hundred reactions, can be used to understand how the central metabolism of an organism
operates[18]. For more advanced research, one can utilise existing curated models (e.g., the
extensive, human-associated bacterial models collection AGORA2[19]) or even create own
models tailored to the respective research question by employing modelling software such as
gapseq and CarveMe[15,20]. Most importantly, it is possible to extend the layers of knowledge
by combining datasets from different omics and databases, also beyond the strict boundaries
of genome-scale metabolic modelling (e.g, see Virtual Metabolic Human database[21],
MetaNetX[22–25], KEGG[26–28]). Emanating from this basis, spatiotemporal approaches that focus
on simulating not only the models but also their environment through time have been
developed[29]. For instance, an advanced approach by Versluis and colleagues has simulated
how the metabolism of specific gut bacteria could alter based on the changes of the
environment cues[30].

In this study, we build on the metabolic modelling software BacArena, which is an
agent-based framework for spatial and temporal simulations of metabolic models. Specifically,
it allows the design of a chemically defined environment, where the cells acting as agents can
have predefined properties (e.g., speed, chemotaxis, linear or exponential growth etc.)[16,31].
We present a virtual colon-like 2D and context-specific application of BacArena that combines
the host and bacterial sides into one single simulation framework. Our implementation spatially
separates bacteria and host cells by representing multiple mucus layers. Furthermore, it
enables a realistic simulation of host and microbial interactions, as it combines spatial-
dependent diffusion phenomena that constrain substrate availability and mimic the lumen
faecal stream powered by peristalsis activity. Lastly, our Virtual Colon software is freely
available on GitHub so that it can easily be adapted by future users
(http://www.github.com/maringos/VirtualColon).

3. Material and methods

General setup

We conducted computational colonisation experiments mimicking an experimental
setting with germ-free mice originally described by Geva-Zatorsky and colleagues
(henceforward the reference study)[32]. Therefore, each simulation comprised bacterial models
together with host models in our colon-like in-silico setting. Furthermore, each setting
simulates seven iterations of one hour. The first iteration was used for the diffusion of
compounds into the environment. In total, each setting was replicated ten times. The growth
of each organism was calculated based on parsimonious flux balance analysis[33]. The majority
of colonisation experiments referred to the introduction of one species (henceforward mono-
colonisation). There were also three cases of specific-pathogen-free mice (SPF). In that case,

we employed the generic simplified human intestinal bacterial community of eight species (see
Table 1, SIHUMIx)[34].

Metabolic models

In the case of the mono-colonisation experiments, bacterial metabolic models
matching the bacterial strains of the reference study were used (see S1 Table)[32]. If a genome
of a strain was not available, we took the most similar genome from NCBI and literature (see
S1 Table) [35]. The genomes were utilised to create the respective bacterial metabolic models
by gapseq[15]. The medium for gap filling was the one available in the lumen and mucus layers
of the Virtual Colon (see S1 Table). The SIHUMIx metabolic models were reconstructed using
gapseq[15] based on the published genomic information by Becker and colleagues[34]. The gap-
filling medium was predicted by gapseq. Finally, the models were adapted to grow on the
medium available in the lumen and mucus layers of the Virtual Colon. To minimise the
possibility of bacterial models competing for nutritional resources with the human models, we
introduced a new "d" compartment into the bacterial models to account for an exclusive
nutritional uptake of compounds from the lumen.

The host metabolic models originate from an adapted version of the published Recon
2.2 model[36]. This generic model was expanded so that a new compartment of exchange
reactions was available. This compartment contains exchange reactions ending with "u" and
is a copy of the original compartment of exchange reactions labelled "e". The reason for this
addition was to represent the flow of compounds between the bacteria in the lumen and the
colonic cells (compartment "e") and between the colonic cells and the blood compartment
(compartment "u"), following the strategy of Sahoo and Thelie[37]. Moreover, we used iMAT to
integrate human transcriptomic data from the reference study[32,38] into the host models[39] to
build context-specific models of the respective colonic tissue. After translating gene
expression activity into reaction expression, a reaction was assumed to be lowly expressed if
it was within the first quartile of all reaction expression values. A reaction was assumed to be
highly expressed if it was in the fourth quartile across all reaction expression values Moreover,
we required a pre-set list of forced reactions and unconstrained nutrition-related exchange
reactions. This information was used as input to determine a context-specific model for each
condition. All exchange reactions and the demand reaction for ATP "DM atp c" were available
in the final models. Detailed information on the utilised models and the procedures can be
obtained from Table 1 and the models are available online (see S1 Table and supplementary
folder 'Models').

Representation of the colon

The computational space of the Virtual Colon was developed by extending BacArena,
which provides a two-dimensional grid area[31]. One bacterial or host cell was located in one
grid cell. The size of the Virtual Colonic lumen, the mucus layers, and the structure of the
colonic crypts were calculated based on anatomic measurements converted to the dimensions
of one grid cell. Specifically, thanks to a predefined crypt-like position of the host cells in the
environment, two half-crypts were represented together with their common inter-cryptal space
(see Fig. 1 and Table 1).

The mucus layers and the lumen of the Virtual Colon were located on top of the crypts,
while below the crypts an area representing the submucosal space and the bloodstream was

added. The inner mucus' width was equal to one-third of the total mucus one, while the outer
mucus' width was equal to two-thirds[40]. The concentration of outer mucus layer compounds
was found to be four times greater than that of the outer ones[41] and was used accordingly
(see S2 Table for more information on design, calculation, and further references). The
bacterial models were introduced in the colonic lumen and both mucus layers. The bacteria of
the inner mucus were unable to move unless the inner mucus was partially degraded. To this
end, in each simulation step, the molecular quantities of inner mucus compounds were
checked. If the amount of at least one compound around a grid cell of the inner mucus layer
was equal to or less than the quantity of the outer layer, then the bacterial models could move
through the grid cell and pass towards the host. The bacterial models could not enter human
space, but they could be located next to the human models.

We started the simulations with reported abundances from the reference study [32]. The
initial position of microbial cells in the Virtual Colon was random. Important additional
parameters were: the arena's size and volume, the actual total volume of the murine host's
caecum, colon and rectum (0.37 cm^3)[42], the bacterial input (200µl of 10^8 cfu/ml; personal
communication with Kasper Dennis ((dennis_kasper@hms.harvard.edu), correspondent
author of the reference study[32]). For instance, the amount of bacteria in the lumen was
calculated based on these formulas:

$$Amount_{lumen} = \frac{Area_{lumen} * Volume_{arena} * 0.2 * 10^8}{Area_{total}}$$

$$Volume_{arena} = Length_{Xaxis} * Length_{Yaxis} * \sqrt{Length_{Xaxis} * Length_{Yaxis}}$$

Similarly, the amount of bacteria in the outer and inner mucus areas was calculated. In the
case of the SIHUMIx community simulations, the same approach was followed. However, an
abundance factor accounting for the relative abundances of each species was used (see Table
1).[34]

The virtual host cells inherited the default human class settings of BacArena[31]. For the
minimum weight and its standard deviation, the scaling factor was used to relate the area of
human and bacterial cells. Rodriguez and colleagues have suggested that one crypt column
is as long as 31.1 cells in the descending mouse colon [43]. Similarly, the two-crypt setting
provided 70 model slots for host cells in total. Since the maximum allowed weight of 70 cells
represented a maximum total weight of around all cells of a human crypt (BNID 110648 [44]). In
addition to these adjustments, no movement was allowed for the models, and their growth was
considered to be linear through time (see Table 1). It should be noted that while there is some
debate about whether optimisation of growth is an adequate objective function for flux balance
simulations of colonic cells, the biomass function includes all molecular components of cells
also required for cell renewal. Furthermore, there is constant growth and shedding of colonic
cells with an estimated turnover time of 3-5 days in mice[45]. Thus, in the colonic context,
optimisation of growth is a valid objective function to predict metabolic fluxes.

The transport of molecules inside BacArena [31] was modelled by diffusion (see S2
Table). The arena was split into the following diffusion sectors: lumen, outer mucus layer, inner
mucus layer, intermediate layer in-between, and blood. The utilised diffusion coefficients were
calculated using a power regression function (coefficients vs. molecular weight) for the colonic
lumen and mucus layers. The molecular weight was used for nutritional compounds. For other
compounds, the default value of glucose was assumed. The regression functions were
generated using the intestinal diffusion data of Winne and colleagues [46]. The same strategy

was followed for the restricted areas and the lumen [47]. Especially for mucus layers, the
diffusion coefficients of molecules in outer mucus were equal to ⅔ of mucus diffusion
coefficients, while for inner mucus, they were equal to ⅓ of mucus diffusion coefficients, based
on the observation that ⅓ of the length of the mucus belongs to the inner layer [40]. Furthermore,
the mucus compounds were not allowed to diffuse to represent the stability of the mucus in
the colonic environment. Oxygen could only diffuse to host areas to ensure limited
"oxygenation" of the Virtual Colon. Practically, the diffusion of compounds in the lumen was
designed to be less restricted than in mucus layers, while not possible in the intermediate layer
and in blood. At the same time, diffusion of blood molecules was possible in the intermediate
layer and in the blood.

Nutrition

Following the nutrition that was chosen by the authors of the reference publication [32],
the primary source of nutritional compounds was the LabDiet ® JL Rat and Mouse/Auto 6F
5K67. These compounds were matched to the dietary bacterial exchange reactions to exclude
nutritional competition between host and bacterial cells. The intestinal absorption and the
dimensions of the Virtual Colon were considered for calculating the exact nutritional input. In
addition, some components such as water, H^+, minerals, vitamins, and mucus compounds
were added as part of the lumen compartment. To represent the inflow of nutrients from the
blood circulation, a minimal collection of compounds was added into the blood compartment.
Their concentration was based on the literature (e.g., (BNID 110365 [44]) and various Human
Metabolome Database entries (https://hmdb.ca/ [48–51]). To infer these components, a shadow
cost analysis based on the BacArena [31] function "plotShadowCost" was utilised (see
supplementary folder 'Definition_Blood'). A list of compounds can be found in the S2 Table.

228
229
230 Figure 1: Schematic representation of the virtual colonic space. The coloured rectangles
231 represent the models of the microbial species and the two shades of green represent the two
232 mucus layers above the host's epithelium. Created with BioRender.com.

Table 1: Overview of properties of the Virtual Colon

Property	Information on values & units	Further comments	Reference
Grid cell dimension	0.00025 × 0.00025 cm^2	Default in BacArena	[31]
Y-axis size	179-grid-cell-long	Set by the developers	Not applicable
X-axis size	24-grid cell-long	Set by the developers	NA
Scaling factor for minimum weight and its standard deviation of host models	$\frac{132.73229}{4.42}$	A human COLO205 colon cell has a diameter of 13 μm; hence its area is equal to 132.73229 μm^2. The area of a bacterium in BacArena is, by default, 4.42 μm^2.	[44] BNID 108890
Scaling factor for maximum weight of host models	$\frac{1000}{70}$	70 cells represent the weight of around 1000 cells of a human crypt	[43,44] BNID 110648[44]
Starting amount of bacteria in the lumen	29	Set by the developers	Not applicable
Starting amount of bacteria in the outer mucus	18	Set by the developers	Not applicable
Starting amount of bacteria in the inner mucus	9	Set by the developers	Not applicable
Relative abundance of *Anaerostipes caccae* in SIHUMIx simulations	0.1263867	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	[34]
Relative abundance of *Bacteroides thetaiotaomicron* in SIHUMIx simulations	0.1444797	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	[34]
Relative abundance of *Bifidobacterium longum* in SIHUMIx simulations	0.1388008	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	[34]
Relative abundance of *Blautia producta* in SIHUMIx simulations	0.1385367	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	[34]
Relative abundance of *Clostridium butyricum* in SIHUMIx simulations	0.1065769	Based on the concentration in human faeces [log10 cells / g dry matter]; see text	[34]
Relative abundance of *Clostridium ramosum* in SIHUMIx simulations	0.1263867	Based on the concentration in human faeces [log10 cells / g dry matter]	[34]
Relative abundance of *Escherichia coli* in SIHUMIx simulations	0.118859	Based on the concentration in human faeces [log10 cells / g dry matter]	[34]
Relative abundance of *Lactobacillus plantarum* in	0.09997359	Based on the concentration in human faeces [log10 cells / g dry	[34]

SIHUMIx simulations		matter]	

Software

The simulations and analyses thereafter presented in this manuscript were conducted in the R environment (version 4.2.1) [52]. Optimisations were generally conducted using the BacArena application (version 1.8.2, commit euba/BacArena@dc8e591 and euba/BacArena@008d186) [31] and sybil (version 2.2.1, commit SysBioChalmers/sybil@58eb989) [53]. The linear programming solver was CPLEX (academic version 22.1.0) [54], which is available through the R package cplexAPI (version 1.4.0) [55]. For data management, the software tidyverse (version 2.0.0) [56] and especially its library dplyr (version 1.1.4) [57] were used. For plotting, the software ggplot2 (version 3.5.1) [58], ggvenn (version 0.1.10)[59], ggthemes (version 5.1.0) [60], scales (version 1.3.0) [61], ggalluvial (version 0.12.5) [62], and BacArena (version 1.8.2, commit euba/BacArena@dc8e591) [31] were utilised. The software devtools (version 2.4.5) [63] was also used to manage the R environment.

4. Results

The main objective of this study was to provide a computational framework to simulate host-microbial interactions in the mammalian colon. We first used our approach to study host-microbiome interactions in the colon based on a previously published dataset of single-strain colonisation experiments in germ-free mice[32]. Besides microbial abundance data, Geva-Zatorsky and colleagues provided extensive transcriptomic datasets of these colonisation experiments. Based on those transcriptomes, we reconstructed context-specific models of colonic tissue for the respective mice. For our study, we used context-specific Recon 2.2 [36] host models along with bacterial models for joint simulations in a virtual colon.

The Virtual Colon provides detailed information on the host-microbial metabolic interactions in space and time. To reveal those interactions in the mono-colonisation experiments, we extracted all fluxes across all time points for each compound and for each model. Since a positive flux shows production and a negative one shows consumption of a compound, we calculated the sum of fluxes of all models for each compound. Collectively, the host models produced 20 compounds and the bacterial ones produced 27 compounds (Fig. 2A). Interestingly, the host-microbial interactions seem to be very pair-specific, since we could identify compounds that in some pairs are bacterially produced or in other pairs by the host (eg., glycine, L-alanine; Fig. 2A). Furthermore, we observed the host exchange of nitrogen-containing compounds, towards the bacterial models, and the uptake of bacterially produced SCFA. Additionally, we found that although the majority of the bacteria interacted metabolically with the host regardless of their position (e.g., see cases with fluxes of models originated from all three layers: lumen, outer mucus, inner mucus, Fig. 2B), we also observed cases of localised interactions (Fig. 2B).

Further, we employed the Virtual Colon to study host-bacterial interactions with the SIHUMIx bacterial community. Fig. 3A shows how the SIHUMIx microbial community that resides in inner and outer colonic mucus gradually consumed the mucus compound N-acetylneuraminate. Furthermore, the bacteria inside the inner mucus layer could only marginally spread in comparison to the other positions due to the design movement barrier. Most interestingly, the decreasing quantity of those compounds follows the growth of the

bacteria models in the area. The bacterial models of *Bacteroides thetaiotaomicron*, *Blautia*
producta, and *Clostridium butyricum* can degrade N-acetylneuraminate (Fig. 3B).

The observation of microbial mucus degradation led to more general questions on what
compounds are consumed or produced by the metabolic models. The host models, similarly
to the mono-colonisation experiments, are context-specific Recon 2.2 models[36], whose
reconstruction was based on the transcriptomes GSM2354402, GSM2354403 and
GSM2354404 respectively. While those three models produce and consume mostly the same
compounds, they do have distinct metabolic profiles that can be attributed to the different
transcriptomes. For instance, the recon model GSM2354403 produced L-alanine, which was
taken up by the bacteria, while the recon model GSM2354404 consumed bacterial L-alanine
(Fig. 4A and 4D). However, all models consumed bacterially produced short-chain fatty acids,
such as formate, acetate, propionate, and butyrate (Fig. 4C and 4D). Interestingly, the models
exchange various metabolic compounds (e.g., water, vitamins, amino acids, TCA-cycle
compounds such as alpha-oxoglutarate).

Not only were the bacterially produced compounds taken up by the host models, but
they were also consumed by other bacterial species of the community (Fig. 5A and 5B). For
instance, apart from amino acids and carbon sources, they also exchanged metabolites like
protons, ammonium, phosphate, and hydrogen sulphide. It is possible to characterise the
metabolic role of community members. The bacteria model of *E. coli* was the most active
species in terms of production and consumption of compounds. On the other hand, *B. longum*
and *C. butyricum* contributed more compounds than they consumed, while the opposite was
true for *A. caccae* and *B. producta*.

Furthermore, in order to understand how the metabolic interactions change when
comparing the interactions from mono-colonisation with the SIHUMIx simulations, we
investigated which compounds are unique in each of the two settings. It can be observed that
the exchanged compounds were mostly common (Fig. 6). However, the majority of the unique
compounds in mono-colonisation experiments were nitrogen-containing compounds. In the
other two parts of the Venn diagram, we could identify amino acids, metabolism intermediates
(e.g. fumarate) and short-chain fatty acids.

322
323
324
325

Figure 2: A. Exchange of compounds which are produced by the bacterial models or context-
specific host models. The colour of the layers depends on the origin of the producer. B.
Exchange of compounds which are produced by the host models and are consumed by the
bacterial model and vice versa. To estimate the exchange of compounds among the models,
the sum of all fluxes across time for each compound and for each model was determined. A
positive sum denotes production, while a negative one denotes consumption. To limit the
numerical artefacts, the fluxes whose absolute value was less than 1E-6 were considered
zero.

Figure 3: A. Schematic representation of the mucus compound N-acetylneuraminate through three-time points in a SIHUMIx-host simulation (GSM2354402, 10th replicate) B. Same as A

340   with the inclusion of the SIHUMIx bacterial species. Each subplot represents the substance

341   concentration and bacterial population per time point in hours.

343
344 Figure 4: Schematic representation of compounds produced by the context-specific host
345 models (A) and consumed by the SIHUMIx bacteria (B) along with the bacterially produced

ones (C) that are consumed by the host models (D). Metabolites with the highest interaction frequency (>= 50%) across all time points and simulations were selected to be plotted for simplicity. For the full dataset, see S3 Table.

Figure 5: Schematic representation of compounds produced by the SIHUMIx bacterial models (A) and consumed by other SIHUMIx bacterial models (B). Metabolites with the highest interaction frequency (>= 50%) across all time points and simulations were selected to be plotted for simplicity. For the full dataset, see S3 Table.

Figure 6: Schematic representation of compounds exchanged by the bacterial models and host models in the mono-colonisation and SIHUMIx experiments. Only interactions with bacteria from the shared genera were taken into account (Bacteroides, Bifidobacterium, Clostridium, Lactobacillus). All simulation time points were included in the analysis.

Table 2: Comparison of Virtual Colon with other published approaches

Name	Approach	Microbes	Scope	Host	Space	Time	Mucus	Gut motility	Availability	Reference
Gut-Microbiota	PDE model with fluid dynamics (Stokes)	Movement + predefined functions	Colon	Nutrient uptake	✓	✓	✓	✓	Matlab code https://forgemia.inra.fr/simon.labarthe/gut-microbiota	[64]
MicroGutPop	ODE model (Monod equation)	predefined functional groups	Colon	-	-	-	✓	✓	R package https://github.com/HelenKettle/microPopGutCode	[65]
GutLogo	ABM		Ileum	-	-	✓	✓	✓	https://github.com/GutLogo/GutLogo	[66]
Synthetic Gut Microbiome	ODE model (gLV)	25 microbes	Butyrate production	-	-	-	✓	-	Julia code https://github.com/RyanLincolnClark/DesignSyntheticGutMicrobiomeAssemblyFunction	[67]
Virtual Colon	ABM + PDE	Movement + metabolic model	Colon	Metabolic model	✓	✓	✓	✓	R code http://www.github.com/maringos/VirtualColon	this article
PDE: Partial differential equation; ODE: Ordinary differential equations; gLV: generalised Lotka-Volterra; ABM: Agent-based modelling										

5. Discussion

Host-bacterial metabolic interactions are considered important drivers of host health
and fitness. Therefore, one of the objectives of this study was to develop a data-driven
approach that allows us to systematically explore the compounds that are exchanged between
the host and bacterial cells. Here we present the Virtual Colon that extends the agent-based
modelling framework BacArena[16,31] to study host-microbiome interactions in the mammalian
large intestine. Specifically, it enables host-microbial simulations, in which bacteria and host
cells are spatially separated through multilayer mucus layers. Moreover, it includes spatial-
dependent diffusion that constrains substrate availability for the models. Most importantly, it
not only enables the joint simulation of host and microbial interactions but also visualises them.
While other methods have contributed to the field either by using differential equations and/or

agent-based modelling (Table 2), the Virtual Colon, which is built on BacArena, combines
them in one software framework.

In our simulations, we utilised host models that originated from the human model
Recon 2.2[36]. For the simulations, the host models were adapted using transcriptomic data[38]
from mono-colonisation experiments of germ-free mice with the respective species[32]. In the
case of colonisation with non-specific pathogens, we used the SIHUMIx bacterial models that
comprise a generic eight-species human intestinal bacterial community[34].

To continue with the host-bacterial interactions in the mono-colonisation experiments,
the observed usage of cysteine by some bacteria of the genus *Fusobacterium* and *Clostridium*
is in line with the literature, which proposes that cysteine is metabolised by bacteria, leading
to hydrogen sulphide[68]. Hydrogen sulphide has been characterised as a compound with
multiple effects, as in low levels, it supports colonic health, while in excess, it is associated
with inflammatory and pathological states (e.g., inflammatory bowel disease and cancer)[68].

Additionally, the compound 4-aminobutanoate, also known as γ-aminobutyric acid
(GABA), is predicted to be produced by *Lactobacillus rhamnosus* and *Bacteroides finegoldi*
and be consumed by the host models, which is in accordance with the literature[69]. This
compound may act on the central nervous system as well as the enteric nervous system of
the host as part of the microbiota-gut-brain axis[69]. On the other hand, *Bacteroides finegoldi*
has not been reported to be a GABA producer. However, a recent article proposed that the
Bacteroides species can produce this molecule, although, in the simulations, no other
Bacteroides species did so[70].

Recently, a study based on metabolomics focusing on the colonic absorption of
compounds with low molecular weight in mice identified some of the computationally predicted
compounds, which were subject to metabolic exchange in our simulations: (glycine) betaine,
cholinem, alanine, glutamine, glycine, isoleucine, lactate, ornithine, and short-chain fatty acids
(Fig. 2A, B) [71]. Interestingly, host-derived glycine betaine and choline can be further bacterially
metabolised to trimethylamine-N-oxide and trimethylamine, for which a linkage to
atherosclerotic cardiovascular disease has been demonstrated[72].

While mono-colonisation experiments offered a detailed overview of pairwise host-
bacterial interactions, the bacterial species found in the mammalian colon are quite diverse.
Therefore, we selected the representative SIHUMIx community to investigate how our results
change in the context of a more complex community. Our simulations showed a wide
exchange of compounds among the species as well as between bacteria and host cells. Not
surprisingly, many byproducts were found also in mono-colonisations, implying that the
repertoire of metabolic exchanges may be genera-dependent (Fig 6). The bacterial models
create a complex web of compounds, where carbohydrate and amino acid metabolism as well
as cofactors (e.g. vitamins) and byproducts are the most common end products. This
observation becomes intriguing, if one considers that carbohydrates rather than amino acids
are primarily used in bacterial energy metabolism, as amino acid metabolism is less   cost-
effective[73]. Moreover, this finding suggests that mono-colonisation experiments only partially
capture interactions that can be observed in community colonisation, as the metabolic
repertoire and signature may alter significantly.

It must be stated that the predicted exchange of compounds between the host and the
bacterial models is based on the assumption that the host colonic cells can exchange
compounds from the luminal side of the colon. For instance, amino acids are known to be
absorbed in the small intestine, while their absorption fate in the large intestine is unclear.

While the colonic bacteria can metabolise the remaining amino acids, it is proposed that some
of them can be used by colonocytes [74]. In parallel, it has been shown that colonocytes can
excrete amino acids (e.g., alanine, aspartate, asparagine, isoleucine, and valine) in both apical
and basolateral directions in vitro [75]. Therefore, the observed production or consumption of
several compounds by the host is feasible and waits for further experimental validation. These
experiments should investigate the transport of metabolites across the intestinal barrier. Such
a pivotal set of experiments was performed by Li and colleagues, who identified gut species
that deplete and metabolise amino acids throughout the intestine and therefore regulate host
metabolism [76].

Furthermore, the Virtual Colon includes spatial parameters tailored to the large
intestinal environment. For instance, the diffusion of compounds inside the mucus layer is
limited and the bacteria in the inner mucus cannot move unless mucus is consumed by
community members, such as *B. thetaiotaomicron*, a known mucus-degrading species [77](Fig
3). Additionally, our results proposed that there are spatial, localised interactions, therefore it
could be suggested that the mucus layers can affect bacterial metabolism. (Fig 2B). Therefore,
the simulation data suggests that our mucus both resembles the function of the mucus as a
physical barrier and as well as a nutrient pool for the bacteria [9].

While the setting of the Virtual Colon recapitulates some of the observed host-microbial
functions, there is also room for further expansions in the simulation setting. Future
computational experiments may include more complex bacterial communities along with an
improved representation of the abundance of bacteria with respect to the size of the simulated
colon (e.g., the ratio of microbial abundance over the number of human models). Another
improvement could include metabolomic data from colonic tissue and lumen so that the
chemical environment is as realistic as possible. Additionally, single-cell transcriptomics would
allow the representation of the various cell types of the host tissue. To improve the realism of
the simulations, the Virtual Colon also has the capacity to mimic the lumen faeces stream
powered by peristalsis activities, if the related parameters are set in the simulation settings.
Most importantly, the presented approach provides users with an easily adaptable platform
that can include all of these extensions.

In summary, our novel simulation framework, Virtual Colon, builds upon the popular
BacArena platform to create a comprehensive tool for studying host-microbe interactions in
the colon. By spatially separating bacteria and host cells and incorporating multilayer mucus
representation, the Virtual Colon enables the realistic simulation of host-microbial interactions.
This is achieved through a spatial-dependent diffusion mechanism that constrains substrate
availability and accurately mimics the faecal stream. Furthermore, the Virtual Colon allows for
the joint simulation of host and microbial interactions, which we demonstrated for human gut
microbiome communities and personalised host models using omics data. Notably, we
employed the Virtual Colon to identify species-specific mucus layer degradation and amino
acid exchange between microbes and host, providing valuable insights into the intricate
dynamics of the gut ecosystem. Our freely available software on GitHub invites future
researchers to adapt and expand its capabilities to advance host-microbe interaction studies.

6. Authors' Contribution

Conceptualisation: GM, JZ
Methodology: GM, JZ, JT, CK
Software: GM, JZ

Formal analysis: GM

Investigation: GM, JZ, JT

Resources: CK

Data Curation: GM

Writing - Original Draft: GM, JZ

Writing - Review & Editing: All co-authors

Visualisation: GM

Supervision: CK, JZ

Project administration: GM

Funding acquisition: CK

7. Acknowledgements

CK acknowledges support from the German Research Foundation for the support through the
collaborative research centre 1182 Origin and Function of Metaorganisms (sub-project A1),
the research unit miTarget (FOR5042), and the excellence cluster "Precision medicine in
chronic inflammation" (EXC2167) and additionally the German Ministry for Education and
Research (E:Med iTREAT, support code 01ZX1902A). This research was supported in part
through high-performance computing resources available at the Kiel University Computing
Centre. The authors thank Dr Ahmed Samer Kadib Alban for providing them with critical
opinions on the manuscript.

8. Conflict of Interest

500 The authors declare that they have no conflict of interest.

9. Supplementary Information

503 The software is freely available on GitHub (www.github.com/maringos/VirtualColon, commit:
504 0e56bda). The simulation RDS files that are related to this manuscript can be found on
505 Figshare (www.doi.org/10.6084/m9.figshare.25997962). The supporting information including
506 figures, scripts and datasets related to this manuscript can be found on Figshare
507 (www.doi.org/10.6084/m9.figshare.25998073).

10.   References

1. Ruuskanen, M. O., Sommeria-Klein, G., Havulinna, A. S., Niiranen, T. J. & Lahti, L.

Modelling spatial patterns in host-associated microbial communities. *Env. Microbiol* **23**,

2374–2388 (2021).

2. Hou, K. *et al.* Microbiota in health and diseases. *Signal Transduct. Target. Ther.* **7**, 135

(2022).

3.   Morrison, D. J. & Preston, T. Formation of short chain fatty acids by the gut microbiota

and their impact on human metabolism. *Gut Microbes* **7**, 189–200 (2016).

4.   Shimizu, H. *et al.* Dietary short-chain fatty acid intake improves the hepatic metabolic

condition via FFAR3. *Sci Rep* **9**, 16574 (2019).

5.   Van Treuren, W. & Dodd, D. Microbial Contribution to the Human Metabolome:

Implications for Health and Disease. *Annu. Rev. Pathol. Mech. Dis.* **15**, 345–369 (2020).

6.   Donohoe, D. R. *et al.* The Microbiome and Butyrate Regulate Energy Metabolism and

Autophagy in the Mammalian Colon. *Cell Metab.* **13**, 517–526 (2011).

7.   Yao, Y. *et al.* The role of short-chain fatty acids in immunity, inflammation and

metabolism. *Crit. Rev. Food Sci. Nutr.* **62**, 1–12 (2022).

8.   Fogelson, K. A., Dorrestein, P. C., Zarrinpar, A. & Knight, R. The Gut Microbial Bile Acid

Modulation and Its Relevance to Digestive Health and Diseases. *Gastroenterology* **164**,

1069–1085 (2023).

9.   Schroeder, B. O. Fight them or feed them: how the intestinal mucus layer manages the

gut microbiota. *Gastroenterol Rep* **7**, 3–12 (2019).

10.   Konjar, Š., Pavšič, M. & Veldhoen, M. Regulation of Oxygen Homeostasis at the

Intestinal Epithelial Barrier Site. *Int J Mol Sci* **22**, (2021).

11.   Litvak, Y., Byndloss, M. X. & Bäumler, A. J. Colonocyte metabolism shapes the gut

microbiota. *Science* **362**, (2018).

12.   O'Keefe, S. J. D. Diet, microorganisms and their metabolites, and colon cancer. *Nat Rev*

Gastroenterol Hepatol **13**, 691–706 (2016).

13.   Cremer, J., Arnoldini, M. & Hwa, T. Effect of water flow and chemical environment on

microbiota growth and composition in the human colon. *Proc. Natl. Acad. Sci.* **114**,

6438–6443 (2017).

14.   Mendoza, S. N., Olivier, B. G., Molenaar, D. & Teusink, B. A systematic assessment of

current genome-scale metabolic reconstruction tools. *Genome Biol* **20**, 158 (2019).

15.   Zimmermann, J., Kaleta, C. & Waschina, S. gapseq: informed prediction of bacterial

metabolic pathways and reconstruction of accurate metabolic models. *Genome Biol* **22**,

81 (2021).

16. Jansma, J. & El Aidy, S. Understanding the host-microbe interactions using metabolic

modeling. *Microbiome* **9**, 16 (2021).

17. Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat Biotechnol* **28**,

245–248 (2010).

18. Orth, J. D., Fleming, R. M. T. & Palsson, B. Ø. Reconstruction and Use of Microbial

Metabolic Networks: the Core *Escherichia coli* Metabolic Model as an Educational

Guide. *EcoSal Plus* **4**, 10.1128/ecosalplus.10.2.1 (2010).

19. Heinken, A. *et al.* Genome-scale metabolic reconstruction of 7,302 human

microorganisms for personalized medicine. *Nat. Biotechnol.* **41**, 1320–1331 (2023).

20. Machado, D., Andrejev, S., Tramontano, M. & Patil, K. R. Fast automated reconstruction

of genome-scale metabolic models for microbial species and communities. *Nucleic Acids

Res.* **46**, 7542–7553 (2018).

21. Noronha, A. *et al.* The Virtual Metabolic Human database: integrating human and gut

microbiome metabolism with nutrition and disease. *Nucleic Acids Res.* **47**, D614–D624

(2019).

22. Ganter, M., Bernard, T., Moretti, S., Stelling, J. & Pagni, M. MetaNetX.org: a website and

repository for accessing, analysing and manipulating metabolic networks. *Bioinformatics*

29, 815–816 (2013).

23. Bernard, T. *et al.* Reconciliation of metabolites and biochemical reactions for metabolic

networks. *Brief. Bioinform.* **15**, 123–135 (2014).

24. Moretti, S. *et al.* MetaNetX/MNXref – reconciliation of metabolites and biochemical

reactions to bring together genome-scale metabolic networks. *Nucleic Acids Res.* **44**,

D523–D526 (2016).

25. Moretti, S., Tran, V. D. T., Mehl, F., Ibberson, M. & Pagni, M. MetaNetX/MNXref: unified

namespace for metabolites and biochemical reactions in the context of metabolic

models. *Nucleic Acids Res.* **49**, D570–D574 (2021).

26. Kanehisa, M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.*

28, 27–30 (2000).

27. Kanehisa, M. Toward understanding the origin and evolution of cellular organisms.

Protein Sci. **28**, 1947–1951 (2019).

28. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG

for taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res.* **51**, D587–

D592 (2023).

29. Scott, W. T. *et al.* A structured evaluation of genome-scale constraint-based modeling

tools for microbial consortia. *PLOS Comput. Biol.* **19**, e1011363 (2023).

30. Versluis, D. M. *et al.* A Multiscale Spatiotemporal Model Including a Switch from Aerobic

to Anaerobic Metabolism Reproduces Succession in the Early Infant Gut Microbiota.

mSystems **7**, e00446-22 (2022).

31. Bauer, E., Zimmermann, J., Baldini, F., Thiele, I. & Kaleta, C. BacArena: Individual-

based metabolic modeling of heterogeneous microbes in complex communities. *PLoS*

Comput Biol **13**, e1005544 (2017).

32. Geva-Zatorsky, N. *et al.* Mining the Human Gut Microbiota for Immunomodulatory

Organisms. *Cell* **168**, 928-943.e11 (2017).

33. Lewis, N. E. *et al.* Omic data from evolved *E. coli* are consistent with computed optimal

growth from genome-scale models. *Mol. Syst. Biol.* **6**, 390 (2010).

34. Becker, N., Kunath, J., Loh, G. & Blaut, M. Human intestinal microbiota: Characterization

of a simplified and stable gnotobiotic rat model. *Gut Microbes* **2**, 25–33 (2011).

35. Fritzenwanker, M. *et al.* Draft Genome Sequence and Complete Plasmid Sequence of

Acinetobacter lwoffii F78, an Isolate with Strong Allergy-Protective Properties. *Genome*

Announc. **4**, e00685-16 (2016).

36. Swainston, N. *et al.* Recon 2.2: from reconstruction to model of human metabolism.

Metabolomics **12**, 109 (2016).

37. Sahoo, S. & Thiele, I. Predicting the impact of diet and enzymopathies on human small

intestinal epithelial cells. *Hum. Mol. Genet.* **22**, 2705–2722 (2013).

38. Gene Expression Omnibus Accession Viewer. GSE88919.

39. Zur, H., Ruppin, E. & Shlomi, T. iMAT: an integrative metabolic analysis tool.

Bioinformatics **26**, 3140–3142 (2010).

40. Gustafsson, J. K. *et al.* An ex vivo method for studying mucus formation, properties, and

thickness in human colonic biopsies and mouse small and large intestinal explants. *Am.*

J. Physiol.-Gastrointest. Liver Physiol. **302**, G430–G438 (2012).

41. Johansson, M. E. V. *et al.* The inner of the two Muc2 mucin-dependent mucus layers in

colon is devoid of bacteria. *Proc. Natl. Acad. Sci.* **105**, 15064–15069 (2008).

42. Casteleyn, C., Rekecki, A., Van Der Aa, A., Simoens, P. & Van Den Broeck, W. Surface

area assessment of the murine intestinal tract as a prerequisite for oral dose translation

from mouse to man. *Lab. Anim.* **44**, 176–183 (2010).

43. De Rodriguez, M. S. B., Sunter, J. P., Watson, A. J., Wright, N. A. & Appleton, D. R. Cell

population kinetics in the mucosal crypts of the descending colon of the mouse.

Virchows Arch. B Cell Pathol. **29**, 351–361 (1978).

44. Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers—the

database of key numbers in molecular and cell biology. *Nucleic Acids Res.* **38**, D750–

D753 (2010).

45. Park, J. *et al.* Promotion of Intestinal Epithelial Cell Turnover by Commensal Bacteria:

Role of Short-Chain Fatty Acids. *PLOS ONE* **11**, e0156334 (2016).

46. Winne, D. & Verheyen, W. Diffusion coefficient in native mucus gel of rat small intestine.

J. Pharm. Pharmacol. **42**, 517–519 (1990).

47. Pappenheimer, J. R. & Reiss, K. Z. Contribution of solvent drag through intercellular

junctions to absorption of nutrients by the small intestine of the rat. *J. Membr. Biol.* **100**,

123–136 (1987).

48. Wishart, D. S. *et al.* HMDB: the Human Metabolome Database. *Nucleic Acids Res.* **35**,

D521–D526 (2007).

49. Wishart, D. S. *et al.* HMDB: a knowledgebase for the human metabolome. *Nucleic Acids*

Res. **37**, D603–D610 (2009).

50. Wishart, D. S. *et al.* HMDB 3.0—The Human Metabolome Database in 2013. *Nucleic*

Acids Res. **41**, D801–D807 (2012).

51. Wishart, D. S. *et al.* HMDB 4.0: the human metabolome database for 2018. *Nucleic*

Acids Res. **46**, D608–D617 (2018).

52. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing.

53. Gelius-Dietrich, G., Desouki, A. A., Fritzemeier, C. J. & Lercher, M. J. sybil – Efficient

constraint-based modelling in R. *BMC Syst. Biol.* **7**, 125 (2013).

54. IBM. IBM ILOG CPLEX Optimization Studio.

55. Gelius-Dietrich, G. cplexAPI: R Interface to C API of IBM ILOG CPLEX.

56. Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

57. Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. dplyr: A Grammar of

Data Manipulation.

58. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York,

2016).

59. Linlin, Y. ggvenn: Draw Venn Diagram by 'ggplot2'.

60. Arnold, J. B. ggthemes: Extra Themes, Scales and Geoms for 'ggplot2'.

61. Wickham, H. & Seidel, D. scales: Scale Functions for Visualization.

62. Brunson, J. ggalluvial: Layered Grammar for Alluvial Plots. *J. Open Source Softw.* **5**,

2017 (2020).

63. Wickham, H., Hester, J., Chang, W. & Bryan, J. devtools: Tools to Make Developing R

Packages Easier.

64. Labarthe, S. *et al.* A mathematical model to investigate the key drivers of the

biogeography of the colon microbiota. *J. Theor. Biol.* **462**, 552–581 (2019).

65. Kettle, H., Louis, P. & Flint, H. J. Process-based modelling of microbial community

dynamics in the human colon. *J. R. Soc. Interface* **19**, 20220489 (2022).

66. Lin, C. *et al.* GutLogo: Agent-based modeling framework to investigate spatial and

temporal dynamics in the gut microbiome. *PLOS ONE* **13**, e0207072 (2018).

67. Clark, R. L. *et al.* Design of synthetic human gut microbiome assembly and butyrate

production. *Nat. Commun.* **12**, 3254 (2021).

68. Blachier, F., Beaumont, M. & Kim, E. Cysteine-derived hydrogen sulfide and gut health:

a matter of endogenous or bacterial origin. *Curr. Opin. Clin. Nutr. Metab. Care* **22**, 68–75

(2019).

69. Yong, S. J., Tong, T., Chew, J. & Lim, W. L. Antidepressive Mechanisms of Probiotics

and Their Therapeutic Potential. *Front. Neurosci.* **13**, 1361 (2020).

70. Otaru, N. *et al.* GABA Production by Human Intestinal Bacteroides spp.: Prevalence,

Regulation, and Role in Acid Stress Tolerance. *Front. Microbiol.* **12**, 656895 (2021).

71. Matsumoto, M. *et al.* Colonic Absorption of Low-Molecular-Weight Metabolites

Influenced by the Intestinal Microbiome: A Pilot Study. *PLOS ONE* **12**, e0169207 (2017).

72. Canyelles, M. *et al.* Gut Microbiota-Derived TMAO: A Causal Factor Promoting

Atherosclerotic Cardiovascular Disease? *Int. J. Mol. Sci.* **24**, 1940 (2023).

73. Oliphant, K. & Allen-Vercoe, E. Macronutrient metabolism by the human gut microbiome:

major fermentation by-products and their impact on host health. *Microbiome* **7**, 91

(2019).

74. Ma, N. & Ma, X. Dietary Amino Acids and the Gut-Microbiome-Immune Axis:

Physiological Metabolism and Therapeutic Prospects. *Compr. Rev. Food Sci. Food Saf.*

18, 221–242 (2019).

75. Chen, Y. *et al.* Absorptive transport of amino acids by the rat colon. *Am. J. Physiol.-*

Gastrointest. Liver Physiol. **318**, G189–G202 (2020).

76. Li, T.-T. *et al.* Microbiota metabolism of intestinal amino acids impacts host nutrient

homeostasis and physiology. *Cell Host Microbe* **32**, 661-675.e10 (2024).

77. Kostopoulos, I. *et al.* A Continuous Battle for Host-Derived Glycans Between a Mucus

Specialist and a Glycan Generalist in vitro and in vivo. *Front. Microbiol.* **12**, 632454

(2021).

Lumen

Outer mucus

Inner mucus

Host area