Impaired Suppression of Plasma Lipid Extraction and its Partitioning Away from Muscle

by Insulin in Humans with Obesity

Christos S. Katsanos,[1,5] Lee Tran,[1] Nyssa Hoffman,[1] Lori R. Roust,[2] Eleanna De Filippis,[2]

Lawrence J. Mandarino,[6] Kailin Johnsson,[1] Marek Belohlavek,[3] and Matthew R. Buras[4]

[1]School of Life Sciences, Arizona State University, Tempe, AZ 85259, [2]College of Medicine,

[3]Department of Cardiovascular Diseases, [4]Department of Quantitative Health Sciences, and

[5]Department of Physiology and Biomedical Engineering, Mayo Clinic Arizona, Scottsdale, AZ

85259, and [6]Department of Medicine, and Center for Disparities in Diabetes, Obesity, and

Metabolism, University of Arizona College of Medicine, Tucson, AZ 85724

Correspondence: Christos S. Katsanos, PhD
16   Health Futures Center, Room 331C
17   6161 E. Mayo Blvd
18   Phoenix, AZ 85054
19   Phone: (602) 543-4254
20   E-mail: christos.katsanos@asu.edu
21   ORCID: https://orcid.org/0000-0003-3025-5752
22

Abstract

Context: Humans with obesity and insulin resistance exhibit lipid accumulation in skeletal muscle,

but the underlying biological mechanisms responsible for the accumulation of lipid in the muscle

of these individuals remain unknown.

Objective: We investigated how plasma insulin modulates the extraction of circulating

triglycerides (TGs) and non-esterified fatty acids (NEFAs) from ingested and endogenous origin

in the muscle of lean, insulin-sensitive humans (Lean-IS) and contrasted these responses to those

in humans with obesity and insulin resistance (Obese-IR).

Methods: The studies were performed in a postprandial state associated with steady-state plasma

TG concentrations. The arterio-venous blood sampling technique was employed to determine the

extraction of circulating lipids across the forearm muscle before and after insulin infusion. We

distinguished kinetics of TGs and NEFAs from ingested origin from those from endogenous origin

across muscle by incorporating stable isotope-labeled triolein in the ingested fat.

Results: Insulin infusion rapidly suppressed the extraction of plasma TGs from endogenous, but

not ingested, origin in the muscle of the Lean-IS, but this response was absent in the muscle of the

Obese-IR. Furthermore, in the muscle of the Lean-IS, insulin infusion decreased the extraction of

circulating NEFAs from both ingested and endogenous origin; however, this response was absent

for NEFAs from ingested origin in the muscle of the Obese-IR subjects.

Conclusions: Partitioning of circulating lipids away from the skeletal muscle when plasma insulin

increases during the postprandial period is impaired in humans with obesity and insulin resistance.

Humans with obesity and insulin resistance display accumulation of lipids in skeletal muscle

(1,2,3). An important factor in the equation determining lipid accumulation in muscle is the

disposal of circulating lipids into the muscle, such as non-esterified fatty acids (NEFAs) and fatty

acids released from circulating triglycerides (TGs). Although elevated circulating NEFAs in the

fasting-state have been proposed to drive lipid accumulation into the muscle of humans with

obesity and insulin resistance (4,5), several lines of evidence argue against this notion (6,7,8).

Circulating TGs constitute a source of fatty acids taken by the muscle, and current evidence shows

that muscle takes up a substantial portion, if not all, of the fatty acids liberated following extraction

of plasma TGs (9,10). This response is of particular importance in the postprandial period (i.e., fed

state) because not only TGs from endogenous origin, but also TGs from ingested origin are present

in the circulation. Postprandial lipid metabolism is complex, and it has been a challenge to

delineate between favorable and unfavorable responses, particularly in regards to how ingested fat

is handled in tissues during the postprandial period (11).

Humans with insulin resistance have increased uptake of plasma TGs by the muscle during the

postprandial period (12,13). A key step for to the uptake of fatty acids liberated from circulating

TGs is the extraction of the circulating TGs by the action of endothelial lipoprotein lipase (LPL)

in muscle capillaries. The increase in plasma insulin is a key hormonal response during the

postprandial period. In healthy individuals, insulin infusion decreases the activity of LPL in muscle

(14). Moreover, insulin sensitivity correlates directly with the insulin-mediated decrease in muscle

LPL activity (14), suggesting that the physiological effect of plasma insulin on suppressing the

extraction of plasma TGs in muscle would be less evident in the presence of insulin resistance.

Further, it is likely that plasma insulin suppresses the extraction of TGs from endogenous origin,

as there is a preferential extraction of circulating TGs from ingested origin during the postprandial

period (10).

We employed a carefully-designed experimental protocol to determine how acute increase in

plasma insulin affects the fractional extraction of plasma TGs in muscle of healthy humans and

humans with obesity and insulin resistance. Study participants ingested fat labeled with a stable

isotope tracer to discern the specific effects of plasma insulin on regulating the extraction of lipids

from ingested origin. The protocol was chosen to avoid recycling of ingested lipid in the

endogenous plasma TG pool, and because it takes at least three hours after a meal for fat from that

meal to appear in the endogenous TG pool (10,15). We hypothesized that acute hyperinsulinemia

immediately suppresses the fractional extraction of plasma TGs in the muscle of healthy, insulin-

sensitive humans, but not in humans with obesity and insulin resistance.

Materials and Methods

Study design

Seventeen volunteers, ages 18 to 40 years, took part in a study to quantify the fractional

extraction of circulating lipids across the forearm muscle. To quantify the extraction of lipids

derived specifically from ingested fat, subjects ingested whipping cream enriched with ^{13}C-

triolein. We determined muscle extraction of both ingested and endogenous plasma TGs and

NEFAs, along with that of plasma glucose, at fasting plasma insulin concentrations (Basal Study

Period) and during increased plasma insulin concentrations (Insulin Study Period; Figure 1). The

arterio-venous (A-V) sampling technique was employed to determine the extraction of circulating

substrates across the forearm muscle. The study was conducted according to Declaration of

Helsinki principles and all study procedures were carried out after obtaining approval from the

Institutional Review Board at Mayo Clinic (IRB # 12005173). Prior to participation, each study

participant gave written, informed consent. The study was registered with ClinicalTrials.gov

(NCT01860911).

Study participants

Initial screening of interested participants was performed over the phone. Individuals were

excluded from the study if they reported diabetes, history of abnormal lipid metabolism,

medications, smoking, current participation in a weight-loss regimen, extreme dietary practices,

took supplements known to affect the vasculature or lipid metabolism (i.e., cocoa, arginine,

protein, fish oil), used anabolic steroids or corticosteroids, or participated in physical activity more

than two days per week. Individuals that met the initial criteria were invited for further screening

in order to estimate their insulin sensitivity if they had body mass index (BMI) ≤ 25 kg/m^2 (i.e.,

lean subjects) or BMI ≥ 32 kg/m^2 (i.e., subjects with obesity).

Screening procedures

For the screening visit, participants arrived in the Clinical Studies Infusion Unit (CSIU) at

Mayo Clinic in Scottsdale, Arizona, in the morning after an overnight fasting period (i.e., 12

hours). The purpose of the study, the exact experimental procedures, and the associated risks were

discussed with each participant prior to obtaining written consent. Screening procedures included

medical history, standard physical examination, electrocardiogram (ECG), routine blood tests, and

urinalysis. Subjects were excluded from the study if there was evidence of acute illness, metabolic

disease such as liver or renal disease, unstable angina, congestive heart failure, plasma TG

concentration >2.3 mmol/l, or 2-h plasma glucose > 11.1 mmol/l during an oral glucose tolerance

test (OGTT). Insulin sensitivity was estimated from the responses of the plasma glucose and

insulin concentrations during the OGTT, and by calculating the Matsuda Insulin Sensitivity Index

(ISI) (16). We studied humans with obesity and insulin resistance (Obese-IR) and lean, insulin-

sensitive humans (Lean-IS) with Matsuda ISI < 4.5 and > 7.0, respectively. We studied groups of

subjects with distinct differences in insulin sensitivity in order to minimize variability in the

responses resulting from biological and technical confounding factors. Body composition,

including distinct components of body lean and fat tissues (i.e., visceral fat, forearm fat), were

evaluated using Dual-energy X-ray absorptiometry (DEXA).

Infusion study visit

Participants arrived at the CSIU for the infusion study visit at ~ 7 am after fasting for ~12

hours. Subjects were asked to maintain their typical diet starting three days prior to the infusion

day, and avoid any beverages with alcohol, as well as any form of physical activity during the

same period beyond their normal daily physical activities. Compliance with these instructions was

verbally verified when participants arrived at the CSIU, and prior to the initiation of the

experimental procedures. An intravenous (IV) line was placed in a retrograde fashion deep in an

antecubital vein draining the forearm muscle. To ensure that the selected vein drains primarily

forearm muscle, oxygen saturation in the blood drawn from this IV line had to be <60% (17); if

not, the IV line would be repositioned. Another catheter was placed under local anesthesia in the

radial artery of the opposite arm for arterial blood sampling. In addition to the arterial catheter, an

additional IV line was inserted in an antecubital vein of the same arm for infusion of insulin. All

catheters were kept patent by continuous infusion of saline.

A depiction of the experimental protocol is shown in Figure 1. Background blood samples

were obtained at time 0, prior to ingestion of whipping cream containing (per 15 ml) 5 g of fat, 1

g of carbohydrate, and 0 g of protein. Whipping cream was enriched (~4%) with $[1,1,1-^{13}C_3,$

99%]triolein (Cambridge Isotope Laboratories, Inc., MA) to differentiate the metabolism of

ingested fat in plasma TGs and NEFAs across the forearm muscle. Labeled triolein was

incorporated into the whipping cream immediately prior to each experiment by sonication for five

mins. After adding a small amount of artificial sweetener, the whipping cream was refrigerated

until it was consumed. Subjects ingested a priming bolus of whipping cream [200 mg fat$\cdot$kg^{-1}fat

free mass (FFM)], followed by ingestion of smaller boluses starting 2 hours later and every 15

mins thereafter until the end of the infusion study (Figure 1), resulting in an overall fat ingestion

of 100 mg fat$\cdot$kg FFM$^{-1}\cdot$h^{-1} during the course of the experiment. We employed an experimental

design that involves ingestion of small boluses of fat in order to avoid increase in the concentration

of plasma TGs carrying ingested TG. Lipoprotein particles carrying ingested TG have 50-fold

higher affinity for LPL than lipoproteins carrying endogenous TG (18), and in a way that can

independently inhibit clearance of lipoproteins carrying endogenous TG by substrate competition

for LPL (19). Therefore, by studying the plasma TG metabolism across muscle with no

perturbation in the plasma TG concentrations and experimentally manipulating a single variable

(i.e., plasma insulin concentration), our experimental approach allowed isolating the specific role

of plasma insulin in regulating the extraction of circulating TG across muscle.

To determine the effects of increased plasma insulin concentrations on variables of interest,

insulin was infused at a rate of 1.0 mU$\cdot$kg$^{-1}\cdot$min^{-1} (2.0 mU$\cdot$kg$^{-1}\cdot$min^{-1} for the first 10 mins) and 0.5

mU$\cdot$kg$^{-1}\cdot$min^{-1} (1.0 mU$\cdot$kg$^{-1}\cdot$min^{-1} for the first 10 mins) in the Lean-IS and Obese-IR subjects,

respectively. Insulin was infused at a lower rate in the Obese-IR subjects, in order to achieve

comparable absolute change in plasma insulin concentrations between the Obese-IR and Lean-IS

subjects during the insulin infusion period. To maintain blood glucose concentrations at the basal

levels during the insulin infusion, 20% dextrose was infused at a rate that was adjusted based on

blood glucose concentration measurements performed every five mins (i.e., hyperinsulinemic-

euglycemic clamp procedure). Two blood samples were collected five minutes apart at the end of

the Basal and Insulin infusion study periods (Figure 1) to compare the effects of increased plasma

insulin on the variables of interest. Arterial and venous blood samples were collected

simultaneously from the radial artery line and the line draining the forearm muscle, respectively.

Blood flow

We measured branchial artery blood flow using Doppler ultrasound in the Basal period and

at the end of the 30-min Insulin infusion period in a group of Lean-IS subjects (i.e., first seven

Lean-IS subjects), and by following procedures previously described (20). Branchial artery blood

flow was measured by evaluating the branchial artery diameter and blood velocity using 2D

Doppler ultrasound (CX50, Philips Medical Systems) and a linear array transducer placed over the

branchial artery. We recorded two separate 2-minute segments, which were processed at a later

time using specialized software. The following equation was used to compute blood flow

(expressed as ml·min^{-1}): Blood Flow = v·π·(d/2)^2·60, where v = mean blood velocity (cm·s^{-1}), π

= 3.14, and d = arterial diameter (cm).

Analyses of plasma samples

Blood samples for the determination of plasma TGs and NEFAs were collected in tubes

containing orlistat, which was added (i.e., 30 ug per ml of blood) to inhibit the in vitro lipolysis of

TGs (21). All tubes were kept on ice until they were centrifuged for the collection of plasma

samples, which were then stored at -80ºC until analyses. An automated blood glucose analyzer

(YSI 2300, Yellow Springs, OH) was used to measure the arterial blood glucose concentration

during the infusion study in order to obtain feedback on how to adjust the infusion rate of 20%

dextrose in response to the insulin infusion. Following the study, concentrations of plasma glucose

in arterial and venous samples were determined by averaging measurements obtained in triplicate

using the same glucose analyzer. Concentrations of plasma TG in the arterial and venous samples

were determined enzymatically using a commercially available assay (Sigma-Aldrich, St. Louis,

MO) and by calculating an average concentration based on ten replicates.

TG from ingested origin is carried in plasma chylomicron lipoproteins (TG content: 90%),

whereas that from endogenous origin is carried in plasma very-low-density-lipoproteins (VLDL;

TG content: 70%) secreted by the liver as well as low-density lipoproteins (LDL; TG content:

10%) resulting from the metabolism of VLDL (22). We distinguished muscle intravascular lipid

metabolism originating from ingested fat by analyzing the ^{13}C-oleate enrichment in the circulating

TGs and NEFAs. Analyses for the determinations of the percentage of oleate within the TG

fraction (i.e, oleate-TG), ^{13}C enrichment of oleate-TG (i.e., [^{13}C]-oleate-TG), concentrations of

individual NEFAs (i.e., myristate, palmitate, palmitoleate, palmitelaidate, stearate, oleate, elaidate,

linoleate, α-linolenate, arachidonate, eicosapentaenoate, docosahexaenoate), and ^{13}C enrichment

of oleate-NEFA (i.e., [^{13}C]-oleate-NEFA) in the arterial and venous plasma samples were

performed at the Mayo Clinic Metabolomics Core. The analyses were performed on a liquid

chromatography mass spectrometry (LC-MS) as previously described (23) using an Agilent 6460

triple quadrupole mass spectrometer coupled with a 1290 Infinity liquid chromatography system,

allowing for high levels of analytical precision measurements. TGs were isolated by solid phase

extraction as described by Bodennec et al. (24). Isolated TGs underwent basic hydrolysis and

additional extraction prior to analysis on the LC-MS. Oleate-TG was expressed as a percentage of

the sum of 12 individual fatty acids from the same fraction, while [^{13}C]-oleate-TG was expressed

as tracer-to-tracee ratio (TTR). Concentrations of NEFAs and TTR of [^{13}C]-oleate-NEFA in

plasma were measured in aliquots that did not go through TG isolation. The concentration of total

NEFAs in plasma was calculated as the sum of the measured concentrations of the individual

plasma NEFAs indicated above.

Concentrations of amino acids (alanine, arginine, asparagine, glutamine, glycine,

isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tryptophan, tyrosine,

valine) in the arterial plasma samples were determined using high-performance liquid

chromatography as we have described previously (25). The concentrations of plasma insulin

(Alpco Diagnostics Cat# 80-INSHU-E01.1, RRID:AB_2801438), apolipoprotein C-II (APOC-II)

(Abcam Cat# ab168549, RRID:AB_3107089) and apolipoprotein C-III (APOC-III) (Abcam Cat#

ab154131, RRID:AB_3107090) in the arterial plasma samples were measured using commercially

available ELISA assays.

Calculations

The concentration of oleate-TG in arterial and venous plasma samples was calculated by

multiplying the measured percentage of oleate in the plasma TGs by the corresponding measured

concentration of plasma TGs. The concentration of plasma [^{13}C]-oleate-TG was calculated by

multiplying the [^{13}C]-oleate-TG TTR by the corresponding plasma oleate-TG concentration. The

concentration of unlabeled-oleate-TG in plasma was calculated as the difference between the

concentration of plasma oleate-TG and that of plasma [^{13}C]-oleate-TG. Arterial and venous plasma

concentrations of [^{13}C]-oleate-NEFA were calculated by multiplying the [^{13}C]-oleate-NEFA TTR

by the corresponding plasma oleate-NEFA concentration. The concentration of unlabeled-oleate-

NEFA was calculated as the difference between the concentration of plasma oleate-NEFA and that

of plasma [^{13}C]-oleate-NEFA.

We calculated the A-V concentration differences of the plasma metabolites of interest

across the muscle to describe metabolite uptake (i.e., positive A-V concentration difference) or

release (i.e., negative A-V concentration difference) from the muscle. Fractional extraction of the

metabolites of interest across the muscle was calculated as the A-V plasma metabolite

concentration difference divided by the arterial plasma metabolite concentration, and expressed as

a percentage.

Statistics

The normality of the data was evaluated using the Shapiro-Wilk test in conjunction with a

visual examination of the corresponding Q-Q plots. A logarithmic transformation was applied to

datasets that displayed a non-normal distribution prior to performing statistical analysis. Unpaired

t-test was used to assess differences in subject characteristics. A two-way (insulin x group)

repeated-measures analysis of variance (ANOVA) was employed to examine the main effects of

plasma insulin concentrations and obesity/insulin-resistant state on the parameters of interest.

Pairwise comparisons were performed using the Bonferroni multiple comparisons test. All tests

were two-sided and $P \leq 0.05$ was considered statistically significant. The APOC-II and APOC-III

data were analyzed using a one-sided test, based on the expectation of a specific directional effect

(26), as higher concentrations of these apolipoproteins are found alongside insulin resistance and

elevated plasma TGs, as observed in our subjects with obesity (27,28). Correlations between

variables of interest were evaluated using the Pearson correlation coefficient (r). Data are reported

as mean ± standard error of the mean (SEM). All statistical analyses were performed using

GraphPad Prism Software (version 8.4.3). To enhance the interpretation of our results and to

facilitate integration of our data in similar research, we employed a web-based effect size

calculator (29) to compute effect size values (i.e., Cohen's d) for the study's primary end-points.

Results

Anthropometric and clinical characteristics of the Lean-IS and Obese-IR subjects are presented

in Table 1. By design, the two groups differed in BMI and insulin sensitivity.

Insulin

Plasma insulin concentrations across the study periods are shown in Table 2. Plasma insulin

concentrations were not different between Fasting and Basal study periods in either group, but they

were higher ($P < 0.05$) in the Obese-IR than the Lean-IS subjects in both study periods. When

compared to the Basal study period, plasma insulin concentrations increased ($P < 0.05$) in both the

Lean-IS and the Obese-IR during the insulin infusion. As expected based on our study design to

achieve comparable absolute changes between groups in the concentrations of plasma insulin

during the Insulin infusion, the delta change from Basal in the plasma insulin concentrations

($\mu IU \cdot ml^{-1}$) was not statistically significantly different between the Lean-IS and Obese-IR groups

(49 ± 3 vs 58 ± 9).

Glucose

Plasma glucose concentrations are shown in Table 2. There was no statistically significantly

difference in plasma glucose concentrations between the Fasting and Basal study periods in either

group, nor were there any statistically significantly differences between the groups in any of these

study periods. Also, during the insulin infusion, plasma glucose concentrations were not

statistically significantly different from Basal in either the Obese-IR or the Lean-IS (Figure 2A).

Plasma glucose fractional extraction (%) across muscle was not statistically significantly

different between groups in the Basal period (Lean-IS: 5.0 ± 0.8; Obese-IR: 2.6 ± 0.4), but it

increased from Basal in response to the Insulin infusion in the Lean-IS (18.2 ± 2.6; $d = 2.26$), but

not in the Obese-IR (7.4 ± 1.9; $d = 1.28$) (Figure 2B). The A-V difference in the concentrations of

plasma glucose (Table 3) was not statistically significantly different between groups at Basal, but

it increased ($P < 0.05$) in response to the Insulin infusion in the Lean-IS ($d = 2.09$), but not in

Obese-IR ($d = 1.30$).

Amino acids

Plasma total amino acid concentrations ($umol \cdot L^{-1}$) were higher ($P < 0.05$) in the Obese-IR

(2652 ± 121) than the Lean-IS (2282 ± 85) at Basal, including the concentrations of the branched

chain amino acids (BCAAs; 631 ± 28 vs 487 ± 27). During the insulin infusion, the concentration

($umol \cdot L^{-1}$) of plasma total amino acids was not statistically significantly different from Basal in

either the Obese-IR (2560 ± 126) or Lean-IS (2261 ± 68), but that of plasma BCAA was lower (P

< 0.05) compared to Basal for both the Obese-IR (582 ± 34) and the Lean-IS (463 ± 27).

Triglycerides

Plasma total TG concentrations were not statistically significantly different between the

Fasting, Basal and Insulin infusion periods within either group. Also, the concentrations of neither

the unlabeled-oleate-TG or the [^{13}C]-oleate-TG were statistically significantly different between

the Basal and Insulin infusion periods. In terms of [^{13}C]-oleate-TG enrichment, there was a main

effect for insulin, but not for group (Table 4); however, there were no statistically significantly

differences in either arterial or venous plasma [^{13}C]-oleate-TG enrichment in response to the

insulin infusion in neither group.

Plasma total TG fractional extraction (%) in the Basal period was lower ($P < 0.05$) in the

Obese-IR (3.3 ± 0.8) than the Lean-IS (8.8 ± 1.9; $d = 1.22$). In response to the insulin infusion, the

fractional extraction (%) of plasma total TG across muscle decreased ($P < 0.05$) when compared

to Basal in the Lean-IS (3.4 ± 0.9; $d = -1.18$), but not in the Obese-IR (2.8 ± 0.6; $d = -0.23$). At

Basal, the fractional extraction of plasma total TGs in muscle across all subjects correlated

statistically significantly and inversely with the concentration of plasma BCAA ($r = -0.52$), but

not with that of plasma total amino acids ($r = -0.42$). However, during insulin infusion, there was

no statistically significant correlation between the delta changes in the fractional extraction of

circulating TGs and the plasma concentrations of BCAA or total amino acids.

The fractional extraction (%) of unlabeled-oleate-TG across the muscle in the Basal Period

was lower ($P < 0.05$) in the Obese-IR (4.0 ± 0.9; $d = 1.15$;) than the Lean-IS (9.9 ± 2.3). The

fractional extraction of unlabeled-oleate-TG across muscle decreased from Basal in response to

the insulin infusion in the Lean-IS (3.3 ± 1.7; $d = -3.31$), but not in the Obese-IR (5.1 ± 1.0; $d =$

1.22) (Figure 3A). In the Basal period, and in contrast to the lower ($P < 0.05$) fractional extraction

of unlabeled-oleate-TG in the muscle of the Obese-IR, the fractional extraction (%) of [^{13}C]-oleate-

TG was not statistically significantly different between the Lean-IS and the Obese-IR (29.5 ± 13.7

vs 22.0 ± 13.6; $d = 0.18$). In response to the insulin infusion, the fractional extraction (%) of plasma

[^{13}C]-oleate-TG in muscle did not change from Basal in either the Lean-IS (31.2 ± 12.4; $d = 0.13$)

or the Obese-IR (-4.7 ± 20.9; $d = -1.51$) (Figure 3B).

At Basal, A-V differences in plasma total TG concentrations were not statistically significantly

different between groups ($d = -0.13$). Although not statistically significant, the decrease in A-V

differences in plasma total TG concentrations in response to insulin infusion showed a larger effect

size in the Lean-IS ($d = -1.08$) compared to the Obese-IR ($d = -0.39$). Similarly, although also not

statistically significant, larger effect size was observed for the decrease in A-V differences in

plasma concentrations of unlabeled oleate-TG in response to insulin infusion in the Lean-IS ($d = -$

0.93) than the Obese-IR ($d = -0.07$). The decreases in A-V differences in plasma concentrations of

[^{13}C]-oleate-TG in response to insulin infusion were not statistically significantly different, with

effect sizes of $d = -1.46$ and $d = -1.02$ in the Lean-IS and Obese-IR, respectively.

Non-esterified fatty acids

The concentrations of plasma NEFA in arterial plasma are shown in Table 2. Plasma NEFA

concentrations were not statistically significantly different either between the Fasting and Basal

study periods within each group, or between the groups during these study periods. Insulin infusion

decreased ($P < 0.05$) the concentrations of total plasma NEFA in arterial plasma in both the Lean-

IS and the Obese-IR. Moreover, during the insulin infusion, the concentration of total NEFAs in

arterial plasma was lower ($P < 0.05$) in the Lean-IS than the Obese-IR. Insulin infusion decreased

($P < 0.05$) the concentration of unlabeled-oleate NEFAs in arterial plasma in both groups, but this

decrease was greater ($P < 0.05$) in the Lean-IS. Insulin infusion decreased ($P < 0.05$) the

concentration of [^{13}C]-oleate-NEFAs in the Lean-IS, only. In terms of [^{13}C]-oleate-NEFA

enrichment, there were main effects for both group and insulin in arterial plasma, but only for

insulin in venous plasma (Table 4).

The fractional extraction (%) of total plasma NEFAs in the Basal period was not statistically

significantly different between the Lean-IS and Obese-IR (3.6 ± 4.2 vs 3.6 ± 3.8; $d = 0.01$). In

response to the insulin infusion, the fractional extraction of total plasma NEFAs became negative

(from positive at Basal) in both groups indicating a release of NEFAs from muscle, and this

response was statistically significantly different in the Lean-IS ($d = -4.05$), but not in the Obese-

IR ($d = -2.25$).

Comparable findings to that of the fractional extraction of plasma total NEFA in the muscle

were obtained with respect to the fractional extraction of the unlabeled oleate within the plasma

NEFA pool. The fractional extraction (%) of plasma unlabeled-oleate-NEFAs across muscle in the

Basal period was nto statistically significantly different between the Obese-IR (5.5 ± 3.8) and

Lean-IS (5.6 ± 4.9) subjects ($d = 0.01$). In response to the insulin infusion, the fractional extraction

of plasma unlabeled-oleate NEFAs decreased in the Lean-IS (-58.6 ± 23.4; $d = -3.81$) but not in

the Obese-IR (-7.2 ± 6.4; $d = -2.39$) (Figure 3C). The fractional extraction (%) of plasma [^{13}C]-

oleate-NEFA in muscle in the Basal period was not statistically significantly different between the

Lean-IS and the Obese-IR (17.4 ± 6.4 vs -1.13 ± 14.7; $d = 0.58$). In response to the insulin infusion,

the fractional extraction (%) of plasma [^{13}C]-oleate-NEFA in muscle decreased from Basal in the

Lean-IS (-8.9 ± 8.1; $d = -3.59$) but not in the Obese-IR (-3.0 ± 11.7; $d = -0.14$) (Figure 3D).

Although the A-V difference in plasma total NEFA concentrations was not statistically

significantly different between groups at Basal, the A-V difference decreased ($P < 0.05$) in

response to the insulin infusion in both the Lean-IS ($d = -1.27$) and the Obese-IR ($d = -0.62$). Also,

the A-V differences in the concentrations of unlabeled-oleate within the plasma NEFA pool

decreased ($P < 0.05$) in response to the insulin infusion in both the Lean-IS ($d = -1.14$) and Obese-

IR ($d = -0.73$). However, the A-V difference in the concentrations of [^{13}C]-oleate within the plasma

NEFA pool in response to the insulin infusion decreased ($P < 0.05$) in the Lean-IS ($d = -1.21$) but

not in the Obese-IR ($d = 0.03$).

Blood flow

The branchial artery blood flow (ml·min^{-1}) during the insulin infusion was not statistically

significantly different from Basal (107 ± 8 vs 102 ± 12; $d = 0.71$).

Apolipoproteins CII and CIII

The concentrations of plasma APOC-II (mg·L^{-1}) in the fasting state were not statistically

significantly different between the Lean-IS (31 ± 6) and Obese-IR (23 ± 3) ($d = 0.52$). However,

the concentrations of plasma APOC-III (mg·L^{-1}) were higher ($P < 0.05$) in the Obese-IR (36 ± 5)

than the Lean-IS (26 ± 2.0) ($d = 0.98$).

Discussion

Our findings show that the postprandial increase in plasma insulin acutely suppresses the

fractional extraction of circulating TGs in muscle. However, this response is absent in the muscle

of humans with obesity and insulin resistance. As a result, during postprandial hyperinsulinemia,

the muscle of individuals with obesity/insulin resistance are exposed to higher amounts of fatty

acids released from circulating TGs as they pass through the muscle, a situation compounded by

their already elevated plasma TG concentrations of endogenous origin. Additionally, the greater

fractional extraction of circulating NEFAs under these conditions in individuals with

obesity/insulin resistance suggests increased plasma fatty acid disposal into their muscles. These

responses may facilitate lipid accumulation in the muscles of humans with obesity/insulin

resistance during the postprandial period. This notion is supported by a recent study conducted

under experimental conditions comparable to this study, which found that insulin infusion

increased muscle diacylglycerol content in individuals with insulin resistance (30).

The fractional extraction of plasma TGs in muscle is mediated by the activity of intravascular

LPL. When assessed *in vitro*, the activity of LPL in muscle is lower in humans with obesity and

insulin resistance (31). Our findings confirm this observation *in vivo* by showing lower fractional

extraction of circulating TGs in muscle of humans with obesity/insulin resistance. The mechanisms

that sustain lower fractional extraction of plasma TGs (i.e., lower LPL activity) in the non-insulin

stimulated state in the muscle of humans with obesity/insulin resistance are currently unknown. At

Basal, and when evaluated across subjects, we found an inverse correlation between plasma BCAA

concentrations and fractional extraction of plasma TGs in muscle. However, a mechanism linking

higher plasma BCAA concentrations to lower TG hydrolysis (i.e., lower LPL activity) in muscle

remains to be identified. On the other hand, there is evidence for lower LPL gene expression in

the muscle of humans with obesity (32). Also, we found that the plasma concentrations of APOC-

III were higher in our subjects with obesity/insulin resistance. Because APOC-III directly inhibits

LPL from hydrolyzing TGs (33), lower fractional extraction of plasma TGs in the muscle of

humans with obesity/insulin resistance at the basal state in our study may be sustained by increased

levels of APOC-III in the circulation of these individuals. It is important to emphasize that the

physiological significance of any given fractional extraction of plasma TGs in muscle must be

evaluated in the context of the prevailing plasma TG concentrations. In our study, the plasma TG

concentrations were ~2.3 times higher in the subjects with obesity/insulin resistance. Because of

that, the absolute amount of TGs extracted by the muscle in the basal, non-insulin-stimulated, state

would be comparable between subjects with obesity/insulin resistance and lean controls, as

demonstrated by comparable A-V differences in the concentrations of plasma TGs.

Our study provides novel evidence that the physiological role of plasma insulin is to acutely

(i.e., within minutes) decrease the extraction of plasma TGs in muscle. Because the

hyperinsulinemia observed during the postprandial period does not increase the gene expression

of LPL in muscle (13), it is expected that the increase in plasma insulin in the present study

downregulated the activity, but not the content, of LPL in muscle. In fact, hormones and dietary

factors are the primary regulators of the activity of the LPL (34,35). At this time, we can only

speculate about the mechanisms by which the activity of LPL in muscle is downregulated by an

increase in plasma insulin. One potential mechanism is the stimulation of nitric oxide production

in the muscle vasculature by plasma insulin (36). Several lines of evidence show that nitric oxide

reduces the activity of LPL (37,38,39), likely through nitration of LPL (40). However, such an

LPL-mediated mechanism remains be elucidated *in vivo* in humans, to explain the decrease in the

fractional extraction of plasma TGs we observed in muscle in response to acute hyperinsulinemia

in our lean/insulin-sensitive subjects.

From a physiological perspective, a decrease in the fractional extraction of plasma TGs in

muscle when plasma insulin levels rise - such as during the postprandial period - indicates that TG

metabolism is partitioned away from the muscle during the postprandial period in healthy, insulin

sensitive humans. However, the absence of a comparable decrease in the fractional extraction of

plasma TGs in the muscle of individuals with obesity/insulin resistance by increased plasma

insulin, coupled with higher plasma TG concentrations in these individuals, exposes their muscles

to greater amounts of fatty acids liberated from circulating TGs as they pass through the muscle

during the postprandial period. Given that muscle takes up a large amount, if not all, of the fatty

acids liberated following the local extraction of plasma TGs (9,10), the inability to decrease the

postprandial fractional extraction of circulating TGs in the muscle of individuals with

obesity/insulin resistance can contribute to lipid accumulation in the muscle of these individuals

during the postprandial period.

Our findings, which show that the fractional extraction of plasma NEFAs in skeletal muscle is

suppressed by the rise in plasma insulin, are consistent with previous research that demonstrated

that insulin reduces the fractional extraction of plasma NEFAs in the heart muscle (41). The failure

of subjects with obesity/insulin resistance to suppress the fractional extraction of plasma NEFAs

from both endogenous and ingested origin during hyperinsulinemia may result from the higher

availability of these plasma NEFAs during the insulin infusion period and/or an increased content

of CD36 fatty acid transporters, which transport fatty acids into muscle (42), in the sarcolemma of

muscle cells of humans with obesity (43). The lack of suppression of fractional extraction of

plasma NEFAs during insulin infusion in our subjects with obesity/insulin resistance, combined

with their inability to decrease the plasma NEFA concentrations to the same extent as the lean

controls, suggests greater NEFA disposal to muscle, likely contributing to lipid accumulation in

the muscle of subjects with obesity/insulin resistance during the postprandial period.

Conclusions

We demonstrate for the first time a direct physiological role of the plasma insulin

increase during the postprandial period in partitioning plasma lipids away from skeletal muscle.

This is achieved by inhibiting the fractional extraction of plasma TGs from endogenous (but not

ingested) origin, and by reducing the fractional extraction of circulating NEFAs from both

endogenous and ingested origin. More importantly, we show that these plasma insulin-mediated

responses are either absent or reduced in individuals with obesity and insulin resistance. Thus,

the capacity of humans with obesity and insulin resistance to redirect circulating lipids away

from the skeletal muscle when plasma insulin rises during the postprandial period is impaired.

Acknowledgments

We gratefully acknowledge the assistance of the subject recruitment team and the nursing

staff in the Clinical Studies Infusion Unit at Mayo Clinic in Scottsdale, Arizona for their support

in conducting the studies. We also thank the subjects for their participation and commitment to the

study procedures.

Funding

The study was supported by American Diabetes Association grant # 7-12-CT-40 and the Mayo

Clinic Metabolomics Resource Core through grant number U24DK100469 from the NIH/NIDDK

and the Mayo Clinic CTSA grant UL1TR000135 from the National Center for Advancing

Translational Sciences.

Data Availability

Original data generated and analyzed during this study are included in this published article.

References

1. Sinha R, Dufour S, Petersen KF, et al. Assessment of skeletal muscle triglyceride content by (1)H nuclear magnetic resonance spectroscopy in lean and obese adolescents: relationships to insulin sensitivity, total body fat, and central adiposity. Diabetes. 2002;51:1022-1027.

2. Stinkens R, Goossens GH, Jocken JW, Blaak EE. Targeting fatty acid metabolism to improve glucose metabolism. Obes Rev. 2015;16:715-757.

3. Coen PM, Goodpaster BH. Role of intramyocelluar lipids in human health. Trends Endocrinol Metab. 2012;23:391-398.

4. Corcoran MP, Lamon-Fava S, Fielding RA. Skeletal muscle lipid deposition and insulin resistance: effect of dietary fatty acids and exercise. Am J Clin Nutr. 2007;85:662-677.

5. Boden G. Obesity, insulin resistance and free fatty acids. Curr Opin Endocrinol Diabetes Obes. 2011;18:139-143.

6. Karpe F, Dickmann JR, Frayn KN. Fatty acids, obesity, and insulin resistance: time for a reevaluation. Diabetes. 2011;60:2441-2449.

7. Kelley DE, Goodpaster B, Wing RR, Simoneau JA. Skeletal muscle fatty acid metabolism in association with insulin resistance, obesity, and weight loss. Am J Physiol. 1999;277:E1130-1141.

8. Cao C, Koh HE, Van Vliet S, et al. Increased plasma fatty acid clearance, not fatty acid concentration, is associated with muscle insulin resistance in people with obesity. Metabolism. 2022;132:155216.

9. Evans K, Burdge GC, Wootton SA, Clark ML, Frayn KN. Regulation of dietary fatty acid entrapment in subcutaneous adipose tissue and skeletal muscle. Diabetes. 2002;51:2684-2690.

10. Bickerton AS, Roberts R, Fielding BA, et al. Preferential uptake of dietary Fatty acids in

adipose tissue and muscle in the postprandial period. Diabetes. 2007;56:168-176.

11. Katsanos CS. Dietary Fatty Acid Spillover in Plasma: A Marker of Good Health, Bad

Health, or Irrelevant? J Clin Endocrinol Metab. 2018;103:5-6.

12. van der Kolk BW, Goossens GH, Jocken JW, Blaak EE. Altered skeletal muscle fatty acid

handling is associated with the degree of insulin resistance in overweight and obese humans.

Diabetologia. 2016;59:2686-2696.

13. van Hees AM, Jans A, Hul GB, Roche HM, Saris WH, Blaak EE. Skeletal muscle fatty acid

handling in insulin resistant men. Obesity (Silver Spring). 2011;19:1350-1359.

14. Kiens B, Lithell H, Mikines KJ, Richter EA. Effects of insulin and exercise on muscle

lipoprotein lipase activity in man and its relation to insulin action. J Clin Invest.

1989;84:1124-1129.

15. Heath RB, Karpe F, Milne RW, Burdge GC, Wootton SA, Frayn KN. Selective partitioning

of dietary fatty acids into the VLDL TG pool in the early postprandial period. J Lipid Res.

2003;44:2065-2072.

16. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance

testing: comparison with the euglycemic insulin clamp. Diabetes Care. 1999;22:1462-1470.

17. Miles JM, Park YS, Walewicz D, et al. Systemic and forearm triglyceride metabolism: fate

of lipoprotein lipase-generated glycerol and free fatty acids. Diabetes. 2004;53:521-527.

18. Xiang SQ, Cianflone K, Kalant D, Sniderman AD. Differential binding of triglyceride-rich

lipoproteins to lipoprotein lipase. J Lipid Res. 1999;40:1655-1663.

19. Chait A, Ginsberg HN, Vaisar T, Heinecke JW, Goldberg IJ, Bornfeldt KE. Remnants of the

Triglyceride-Rich Lipoproteins, Diabetes, and Cardiovascular Disease. Diabetes.

2020;69:508-516.

20. Green D, Cheetham C, Reed C, Dembo L, O'Driscoll G. Assessment of brachial artery blood

flow across the cardiac cycle: retrograde flows during cycle ergometry. J Appl Physiol

(1985). 2002;93:361-368.

21. Labbe SM, Croteau E, Grenier-Larouche T, et al. Normal postprandial nonesterified fatty

acid uptake in muscles despite increased circulating fatty acids in type 2 diabetes. Diabetes.

2011;60:408-415.

22. Cox RA, Garcia-Palmieri MR. Cholesterol, Triglycerides, and Associated Lipoproteins. In:

Walker HK, Hall WD, Hurst JW, eds. Clinical Methods: The History, Physical, and

Laboratory Examinations. 3rd ed. Boston1990.

23. Persson XM, Blachnio-Zabielska AU, Jensen MD. Rapid measurement of plasma free fatty

acid concentration and isotopic enrichment using LC/MS. J Lipid Res. 2010;51:2761-2765.

24. Bodennec J, Koul O, Aguado I, Brichon G, Zwingelstein G, Portoukalian J. A procedure for

fractionation of sphingolipid classes by solid-phase extraction on aminopropyl cartridges. J

Lipid Res. 2000;41:1524-1531.

25. Tran L, Kras KA, Hoffman N, et al. Lower fasted-state but greater increase in muscle

protein synthesis in response to elevated plasma amino acids in obesity. Obesity (Silver

Spring). 2018;26:1179-1187.

26. Murphy R. On the use of one-sided statistical tests in biomedical research. Clin. Exp.

Pharmacol. Physiol. 2018;45:109-114.

27. Qamar A, Khetarpal SA, Khera AV, Qasim A, Rader DJ, Reilly MP. Plasma apolipoprotein

C-III levels, triglycerides, and coronary artery calcification in type 2 diabetics. Arterioscler

Thromb Vasc Biol. 2015;35:1880-1888.

28. Hiukka A, Fruchart-Najib J, Leinonen E, Hilden H, Fruchart JC, Taskinen MR. Alterations

of lipids and apolipoprotein CIII in very low density lipoprotein subspecies in type 2

diabetes. Diabetologia. 2005;48:1207-1215.

29. Wilson DB. Practical meta-analysis effect size calculator (Version 2023.11.27). 2023;

https://campbellcollaboration.org/research-resources/effect-size-calculator.html. Accessed

December 17, 2023, 2023.

30. McKenna CF, Stierwalt HD, Zemski Berry KA, et al. Intramuscular diacylglycerol

accumulates with acute hyperinsulinemia in insulin-resistant phenotypes. Am J Physiol

Endocrinol Metab. 2024;327:E183-E193.

31. Pollare T, Vessby B, Lithell H. Lipoprotein lipase activity in skeletal muscle is related to

insulin sensitivity. Arterioscler Thromb. 1991;11:1192-1203.

32. Boucher P, Ducluzeau PH, Davelu P, et al. Expression and regulation by insulin of low-

density lipoprotein receptor-related protein mRNA in human skeletal muscle. Biochim

Biophys Acta. 2002;1588:226-231.

33. Larsson M, Allan CM, Jung RS, et al. Apolipoprotein C-III inhibits triglyceride hydrolysis

by GPIHBP1-bound LPL. J Lipid Res. 2017;58:1893-1902.

34. Doolittle MH, Ben-Zeev O, Elovson J, Martin D, Kirchgessner TG. The response of

lipoprotein lipase to feeding and fasting. Evidence for posttranslational regulation. J Biol

Chem. 1990;265:4570-4577.

35. Wang H, Eckel RH. Lipoprotein lipase: from gene to obesity. Am J Physiol Endocrinol

Metab. 2009;297:E271-288.

36. Muniyappa R, Montagnani M, Koh KK, Quon MJ. Cardiovascular actions of insulin.

Endocr. Rev. 2007;28:463-491.

37. Picard F, Kapur S, Perreault M, Marette A, Deshaies Y. Nitric oxide mediates endotoxin-

induced hypertriglyceridemia through its action on skeletal muscle lipoprotein lipase.

FASEB J. 2001;15:1828-1830.

38. Uchida Y, Tsukahara F, Ohba K, et al. Nitric oxide mediates down regulation of lipoprotein

lipase activity induced by tumor necrosis factor-alpha in brown adipocytes. Eur. J.

Pharmacol. 1997;335:235-243.

39. Ricart-Jane D, Casanovas A, Jane N, et al. Nitric oxide and the release of lipoprotein lipase

from white adipose tissue. Cell. Physiol. Biochem. 2008;22:525-530.

40. Casanovas A, Carrascal M, Abian J, Lopez-Tejero MD, Llobera M. Lipoprotein lipase is

nitrated in vivo after lipopolysaccharide challenge. Free Radic Biol Med. 2009;47:1553-

1560.

41. Ng CK, Soufer R, McNulty PH. Effect of hyperinsulinemia on myocardial fluorine-18-FDG

uptake. J. Nucl. Med. 1998;39:379-383.

42. Glatz JFC, Nabben M, Luiken J. CD36 (SR-B2) as master regulator of cellular fatty acid

homeostasis. Curr Opin Lipidol. 2022;33:103-111.

43. Bonen A, Parolin ML, Steinberg GR, et al. Triacylglycerol accumulation in human obesity

and type 2 diabetes is associated with increased rates of skeletal muscle fatty acid transport

and increased sarcolemmal FAT/CD36. FASEB J. 2004;18:1144-1146.

Figure Legends

Figure 1. Depiction of the experimental protocol employed to determine the postprandial

extraction of plasma triglycerides and non-esterified fatty acids prior to (Basal) and following

insulin infusion (Insulin). $[1,1,1-^{13}C_3]$triolein-enriched whipping cream was ingested at the time

points indicated in the form of small boluses to induce steady-state concentrations of ingested

and overall plasma triglycerides. Arterio-venous (A-V) blood samples were collected from a

radial artery and a vein draining forearm muscle.

Figure 2. Concentrations of plasma glucose (**A**) and forearm muscle fractional extraction of

plasma glucose (**B**) prior to (Basal) and following insulin infusion (Insulin) in lean, insulin

sensitive humans (Lean – IS) and humans with obesity and insulin resistance (Obese - IR). Mean

$\pm$ SEM are shown in (A) and individual data points in (B). Data were analyzed by a two-way

ANOVA followed by pairwise comparisons using the Bonferroni multiple comparisons test.

***$P \leq 0.001$

Figure 3. Forearm muscle fractional extraction of plasma endogenously-produced triglycerides

(TG) (i.e., Unlabeled-oleate-TG) (**A**) and non-esterified fatty acids (NEFA) (i.e., Unlabeled-

oleate-NEFA) (**C**) as well as ingested TG (i.e., $[^{13}C]$-oleate-TG) (**B**) and NEFA (i.e., $[^{13}C]$-

oleate-NEFA) (**D**) prior to (Basal) and following insulin infusion (Insulin) in lean, insulin

sensitive humans (Lean – IS) and humans with obesity and insulin resistance (Obese - IR).

Individual data points are shown. Data were analyzed by a two-way ANOVA followed by

pairwise comparisons using the Bonferroni multiple comparisons test. *$P \leq 0.05$, **$P \leq 0.01$.

Table 1. Subject characteristics

	Lean - IS	Obese - IR
n (F/M)	9 (2/7)	8 (3/5)
Age (years)	24.6 ± 1.6	27.3 ± 3.0
Weight (kg)	70.6 ± 3.4	107.7 ± 7.2***
BMI (kg/m^2)	22.4 ± 0.8	35.0 ± 2.2***
Lean body mass (kg)	55.7 ± 2.3	65.0 ± 4.7
Forearm lean mass (kg)	0.99 ± 0.07	0.99 ± 0.14
Forearm fat mass (kg)	0.20 ± 0.03	0.56 ± 0.08***
Forearm fat mass (%)	15.7 ± 2.5	35.3 ± 5.5**
Body fat mass (%)	17.5 ± 1.8	36.6 ± 3.6***
Android/Gynoid Fat	0.29 ± 0.03	0.62 ± 0.07***
Visceral Fat Mass (g)	287 ± 68	714 ± 135*
Fasting plasma glucose (mmol·L^{-1})	4.6 ± 0.1	5.0 ± 0.2*
Fasting plasma insulin (μIU·ml^{-1})	2.9 ± 0.3	11.7 ± 1.4***
HOMA-IR	0.6 ± 0.1	2.6 ± 0.3***
Matsuda-ISI	11.5 ± 1.2	3.1 ± 0.3***
HbA1c (mmol·mol^{-1})	33.5 ± 1.6	34.0 ± 1.4
HbA1c (%)	5.2 ± 0.1	5.3 ± 0.1
TSH (mIU·l^{-1})	1.3 ± 0.3	1.5 ± 0.3
Plasma triglycerides (mmol·L^{-1})	0.8 ± 0.1	1.6 ± 0.2**
Total plasma cholesterol (mmol·L^{-1})	3.8 ± 0.2	4.1 ± 0.2
Plasma HDL-Cholesterol (mmol·L$^-$	1.4 ± 0.1	1.0 ± 0.1***
Plasma LDL-Cholesterol (mmol·L^{-1})	2.0 ± 0.3	2.4 ± 0.2

Values are mean ± SEM. Lean-IS, lean, insulin sensitive subjects; Obese-IR,

subjects with obesity and insulin resistance; BMI, body mass index;

Android/Gynoid fat, ratio of android fat mass to gynoid fat mass; HOMA-IR,

homeostatic model assessment of insulin-resistance; Matsuda-ISI, Matsuda

Insulin Sensitivity Index; HbA1c, glycated hemoglobin; TSH, thyroid-stimulating

hormone; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

*$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, statistically significantly different from Lean-IS.

Table 2. Concentration of insulin and metabolites in arterial plasma during fasting, the basal period and insulin infusion

	Fasting	Basal	Insulin	ANOVA (*P Value*)		
				Interaction	Group	Insulin
Insulin (μIU·ml^{-1})				*0.37*	*< 0.001*	*< 0.001*
Lean-IS	4.0 ± 0.6	4.4 ± 0.7	53.3 ± 3.3			
Obese-IR	15.2 ± 1.2	16.8 ± 3.1	74.9 ± 9.1			
Glucose (mmol·L^{-1})				*0.42*	*0.45*	*0.82*
Lean-IS	4.4 ± 0.1	4.5 ± 0.1	4.4 ± 0.1			
Obese-IR	4.6 ± 0.2	4.6 ± 0.2	4.6 ± 0.1			
Total-TG (μmol·L^{-1})				*0.76*	*0.02*	*0.31*
Lean-IS	585 ± 68	642 ± 91	589 ± 95			
Obese-IR	1297 ± 280	1437 ± 338	1384 ± 329			
Unlabeled-Oleate-TG (μmol·L^{-1})				*0.78*	*0.05*	*0.07*
Lean-IS	N/A	218 ± 35	198 ± 36			
Obese-IR	N/A	493 ± 135	479 ± 132			
[^{13}C]-oleate-TG (μmol·L^{-1})				*0.24*	*0.21*	*0.16*
Lean-IS	N/A	1.5 ± 0.4	1.6 ± 0.3			
Obese-IR	N/A	2.8 ± 1.1	3.5 ± 1.5			
Total-NEFA (μmol·L^{-1})				*0.17*	*0.07*	*<0.001*
Lean-IS	418 ± 72	430 ± 36	101 ± 12			
Obese-IR	444 ± 53	496 ± 48	281 ± 38			
Unlabeled-oleate-NEFA				*0.10*	*0.02*	*<0.001*
Lean-IS	N/A	159 ± 14	30 ± 3			
Obese-IR	N/A	174 ± 20	89 ± 13			
[^{13}C]-oleate-NEFA (μmol·L^{-1})				*0.01*	*0.55*	*0.42*
Lean-IS	N/A	0.96 ± 0.17	0.52 ± 0.07			
Obese-IR	N/A	0.51 ± 0.08	0.76 ± 0.22			

614   Values are mean ± SEM; Lean-IS, lean, insulin sensitive subjects; Obese-IR, subjects with

615   obesity and insulin resistance; TG, triglycerides; NEFA, non-esterified fatty acids.

Table 3. Arterio-venous differences of metabolites in the basal period and during insulin infusion

	Basal	Insulin	ANOVA (*P value*)		
			Interaction	Group	Insulin
Glucose (mmol·L^{-1})			*0.03*	*0.01*	*<0.001*
Lean-IS	0.23 ± 0.04	0.81 ± 0.13			
Obese-IR	0.12 ± 0.02	0.35 ± 0.09			
Total-TG (µmol·L^{-1})			*0.59*	*0.46*	*0.03*
Lean-IS	55 ± 15	18 ± 6			
Obese-IR	63 ± 27	40 ± 14			
Unlabeled-oleate-TG (µmol·L^{-1})			*0.18*	*0.20*	*0.09*
Lean-IS	21 ± 7	6 ± 3			
Obese-IR	26 ± 11	24 ± 7			
^{13}C-Oleate-TG (µmol·L^{-1})			*0.60*	*0.62*	*0.12*
Lean-IS	0.58 ± 0.23	0.29 ± 0.16			
Obese-IR	0.96 ± 0.61	0.40 ± 0.48			
Total-NEFA (µmol·L^{-1})			*0.04*	*0.50*	*<0.001*
Lean-IS	16.5 ± 17.8	-45.5 ± 14.6			
Obese-IR	14.8 ± 17.7	-13.8 ± 14.4			
Unlabeled-oleate-NEFA (µmol·L^{-1})			*0.06*	*0.53*	*<0.001*
Lean-IS	8.7 ± 8.2	-16.4 ± 6.1			
Obese-IR	8.1 ± 6.4	-4.3 ± 5.3			
^{13}C-Oleate-NEFA (µmol·L^{-1})			*0.04*	*0.53*	*0.05*
Lean-IS	0.18 ± 0.07	-0.06 ± 0.06			
Obese-IR	0.01 ± 0.06	0.01 ± 0.06			

616   Values are mean ± SEM; Lean-IS, lean, insulin sensitive subjects; Obese-IR, subjects with obesity

617   and insulin resistance; TG, triglycerides; NEFA, non-esterified fatty acids.

Table 4. Isotopic enrichment of plasma triglycerides and non-esterified fatty acids with [¹³C]-oleate in arterial and venous plasma in the basal period and during insulin infusion

| | Basal | Insulin | ANOVA (*P value*) | | |
			Interaction	Group	Insulin
TG (TTR) - Arterial			*0.55*	*0.32*	*0.02*
Lean-IS	0.007 ± 0.001	0.008 ± 0.001			
Obese-IR	0.005 ± 0.001	0.007 ± 0.002			
TG (TTR) - Venous			*0.78*	*0.69*	*<0.01*
Lean-IS	0.005 ± 0.001	0.006 ± 0.002			
Obese-IR	0.004 ± 0.001	0.006 ± 0.001			
NEFAs (TTR) - Arterial			*<0.01*	*<0.01*	*<0.001*
Lean-IS	0.006 ± 0.001	0.017 ± 0.001			
Obese-IR	0.003 ± 0.001	0.008 ± 0.002			
NEFA (TTR) - Venous			*0.06*	*0.06*	*<0.001*
Lean-IS	0.005 ± 0.001	0.012 ± 0.001			
Obese-IR	0.003 ± 0.001	0.007 ± 0.002			

618   Values are tracer-to-tracee ratio (TTR) mean ± SEM; Lean-IS, lean, insulin sensitive subjects; Obese-

619   IR, subjects with obesity and insulin resistance; TG, triglycerides; NEFA, non-esterified fatty acids.

Figure 1

Figure 2

Figure 3