B4 Raf-like MAPKKK RAF24 regulates *Arabidopsis thaliana* flowering time through HISTONE MONO-UBIQUITINATION 2

[1]Qiaomu Li, [1]Le Wang, [1]Maria Camila Rodriguez Gallo, [1,2]Devang Mehta, [1]Sabine Scandola, [1]Mohana Talasila and [1,3#]Uhrig RG.

[1]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada.
[2]Department of Biosystems, Katholieke Universiteit Leuven, Leuven, Belgium
[3]Department of Biochemistry, University of Alberta, Edmonton, AB, Canada.

Corresponding Author
R. Glen Uhrig
Department of Biological Sciences
University of Alberta
11455 Saskatchewan Drive
Edmonton, Alberta
Canada
T6G 2E9

Keywords: *Arabidopsis thaliana* Flowering, Quantitative Phosphoproteomics, RAF24, HISTONE MONO-UBIQUITINATION 2 (HUB2) Interactome.

Running Title: Loss of RAF24 leads to a decrease in phosphorylated HUB2 resulting in early flowering.

ABSTRACT

The timing of flowering is a critical agronomic trait governed by a number of external cues. Despite our genetic understanding of flowering time being well established, we have a limited understanding of how these signals are transmitted to different flowering genes through protein phosphorylation. Here, we characterize a novel B4 Raf-like MAPKKK protein kinase called RAF24, whose mutation results in an early flowering phenotype. Comparative analysis to related B4 Raf-like MAPKKKs indicates that RAF24 unqiuely affects flowering time, while phosphoproteome analyses found RAF24 impacts the phosphorylation status of proteins involved in distinct flowering pathways. In particular, we found the RING-type ubiquitin ligase HISTONE MONO-UBIQUITINATION 2 (HUB2) to possess the largest phosphorylation change in *raf24* deficient plants relative to wild-type Arabidopsis and that RAF24 suppresses ligase activity of HUB2 in order to maintain approriate levels of H2Bub1. Furthermore, we found that RAF24 regulates HUB2 phosphorylation through subclass I and III SUCROSE NON-FERMENTING KINASE 2 (SnRK2) protein kinases; known substrates of B4 RAF-like MAPKKKs. Lastly, using a combination of phospho-mimetic and -ablative plant lines, we validate the importance of HUB2 phosphorylation at S^{314} in regulating flowering time. Collectively, our findings implicate RAF24 as a higher-order flowering regulator, while further implicating HUB2 as a centerpiece of flowering regulation.

INTRODUCTION

As sessile organisms, land plants are able to perceive and adapt to constantly changing external conditions, such as temperature, light and photoperiod. This ability enables them to adjust their developmental transitions, such as flowering time, in response to diel rhythms or seasonal changes (Mouradov *et al.*, 2002; Creux & Harmer, 2019; Bao *et al.*, 2020). Premature flowering affects the overall fitness of plants, thereby negatively influencing crop yield and productivity (Gaudinier & Blackman, 2020). To avoid this, plants employ a series of complex regulatory networks. To date, our knowledge of flowering rests largely at the transcriptional level, with numerous transcription factors having been identified to fine-tune flowering time. Ultimately, these networks culminate with *FLOWERING LOCUS C* (FLC), a major flowering repressor, directly regulating the expression of floral integrators, *FLOWERING LOCUS T* (FT) and *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1* (SOC1), to suppress premature flowering (Lee & Lee, 2010; Deng *et al.*, 2011; Bao *et al.*, 2020).

Post-translational modifications (PTMs), such as protein phosphorylation and ubiquitination, work synchronously with transcriptional changes to control flowering time (Linden & Callis, 2020). Protein phosphorylation is directed by protein kinases, which transfer the phosphate group from ATP to Ser, Thr and Tyr residues of a protein substrate (Ubersax & Ferrell, 2007), resulting in a multitude of outcomes, including: altered enzymatic activity, protein stability and subcellular localization (Humphrey *et al.*, 2015). For instance, calcium-dependent protein kinase CPK28 directly phosphorylates PLANT U-BOX 25 (PUB25) and PUB26 to upregulate its ubiquitin ligase activity and ensure immune homeostasis (Wang *et al.*, 2018a). So far, several kinases have been related to flowering time control (Ogiso *et al.*, 2010; Chen *et al.*, 2020; Sanagi *et al.*, 2021; Li *et al.*, 2021), including SHAGGY-like kinase 12 (SK12), which was recently resolved to phosphorylate the core flowering regulator CONSTANS (CO) causing rapid degradation of CO by the 26S proteasome, thereby repressing flowering (Chen et al., 2020).

In addition to phosphorylation, protein ubiquitination is also associated with flowering. Protein ubiquitination is mediated by a multi-enzymatic cascade. This involves E1 enzymes, which activate ubiquitin molecules that are then passed to E2 enzymes to form an E2-ubiquitin intermediate. Lastly, E3 enzymes direct the transfer of an ubiquitin moiety from E2 enzymes to specifically recognized protein targets, leading to ubiquitination of substrates (Vierstra, 2009). Interestingly, a number of E2s and E3s have been shown to possess regulatory functions in flowering development. For example, the interaction of RING-type ubiquitin ligases UBC1/2, with HISTONE MONOUBIQUITINATION1 (HUB1) and HUB2 to mono-ubiquitinate histone B2 (H2B) proteins and regulate flowering time (Cao *et al.*, 2008; Gu *et al.*, 2009). Absence of ubiquitin-modified H2B (H2Bub1) has been suggested to decrease FLC levels leading to early flowering phenotypes in *hub1* and *hub2* mutants (Cao *et al.*, 2008; Gu *et al.*, 2009). H2Bub1 is also known to modulate the circadian expression of genes (Himanen *et al.*, 2012; Woloszynska *et al.*, 2019), and to induce light-responsive genes (Bourbousse *et al.*, 2012), in addition to modulating the expression of auxin biosynthesis genes (Zhang *et al.*, 2021). Most recently, HUB1 and 2 have been shown to interact with SPEN3 and RNA-binding KH domain-containing

protein (KHD1), to influence the formation of anti-sense COOLAIR transcripts (Woloszynska *et al.*, 2019). Collectively, these findings indicate HUB2 maintains a complex, multifaceted connection to flowering time. Despite HUB2 being extensively linked to multiple biological processes in plants, how HUB2 is modulated in plants remains unknown. Interestingly, the human orthologs of HUB1/2, Ring Finger Protein 20/40 (RNF20/40), are found to be phosphorylated by the ATAXIA-TELANGIECTASIA MUTATED (ATM) protein kinase, which modulates H2Bub1 levels, suggesting that phosphorylation may affect the ubiquitin ligase activity of HUB2 (Moyal *et al.*, 2011).

Recently, a subfamily of Arabidopsis B4 Raf-like MAPKKKs (RAF18, RAF20 and RAF24) were reported to phosphorylate sucrose non-fermenting-1 (SNF1)-related kinases (SnRK2s) (MAPK Group, 2002; Lozano-Juste et al., 2020). Upon osmotic stress conditions, RAF18, RAF20 and RAF24 strongly phosphorylate subclass I SnRK2s, which then phosphorylate and activate VARICOSE (VCS) to modulate the mRNA population and plant stress response (Lin et al., 2020; Soma et al., 2020; Fàbregas et al., 2020). In fact, Lin et al also found that RAF24 was able to weakly phosphorylate SnRK2.6, one of the members from subclass III (Lin et al., 2020). On the other hand, some RAFs have been found to be involved in plant growth and fitness. For example, CTR1/RAF1 was characterized as a negative regulator of ethylene hormone signaling (Ju et al., 2012), while EDR1/RAF2 has been implicated in fine-tuning plant immunity (Zhao et al., 2014). In addition, RAF22 and RAF36 were reported to negatively regulate abiotic stress responses and BHP/RAF27 was shown to be a regulator of stomatal opening (Hayashi et al., 2017; Kamiyama et al., 2021; Sun et al., 2022).

Although RAF18, 20 and 24 have been assigned roles in osmotic stress response, other biological roles for these RAFs remain unknown. In this study, we report that RAF24 functions as a flowering time repressor that operates independently of its close relatives; RAF18 and RAF20. Our combined use of genetic, molecular and biochemical analyses coupled with quantitative phosphoproteomics, reveals that RAF24, through select subclass I and III SnRKs, regulates HUB2 to influence flowering time. Moreover, we reveal that the ability of RAF24 to fine-tune flowering seems to be dependent on FLC and FT, but not SOC1. We also find that loss of RAF24 results in changes in both HUB2 ubiquitin ligase activity and its protein interactome. Finally, using a combination of phospho-mimetic (HUB2^{S314D}) and -ablative (HUB2^{S314A}) mutations, we show that phosphorylation of HUB2 at S^{314} suppresses the early flowering phenotype of the *raf24-2* mutant. Taken together, we reveal RAF24 to be a protein kinase whose function is to repress flowering by influencing HUB2 ligase activity and its interactions with other proteins.

Results
Loss of RAF24 causes early flowering in Arabidopsis
The Arabidopsis subgroup B Raf-like MAPKKKs include seven members. Among these, clade B4 (RAF18, RAF20 and RAF24) are phylogenetically related (Supplemental Figure 1; (MAPK Group, 2002; Lozano-Juste *et al.*, 2020)). Previously, RAF18, RAF20 and RAF24 were found to be phosphorylated by subclass I SnRK2 kinases in response to osmotic stress conditions

(Stecker *et al.*, 2014; Soma *et al.*, 2017, 2020; Lin *et al.*, 2020). Further, diel phosphoproteomic
analyses found RAF18 and RAF24, but not RAF20, to possess a significantly increased
phosphorylation status in rosette leaves at the end of day (ED) compared to the end of night
(EN), suggesting that B4 clade RAFs may also have additional roles independent of osmotic
stress signaling (Supplemental Figure 1) (Uhrig *et al.*, 2019). To better understand potential
alternative roles for these RAFs, we acquired homozygous T-DNA insertion lines for *RAF18,*
RAF20 and *RAF24*, which included *raf18-1, raf20-1* and *raf24-2*, respectively. Here, we found
that only *raf24-2* mutants displayed early flowering compared to wild-type *Arabidoipsis thaliana*
Col-0 (WT Col-0) and related *raf18 / 20* mutants (Figure 1B - C; Supplemental Figure 2A - B).
Furthermore, *RAF24* exhibited high transcriptional expression in leaf tissues compared to *RAF18*
and *RAF20*; both of which are more abundant in the roots (Soma *et al.*, 2020).
To further confirm the early flowering phenotype of *raf24-2*, we acquired two additional T-
DNA insertional lines, *raf24-4* and *raf24-5* and assessed their time to flowering (Figure 1A).
Here, we found *raf24-4* and *raf24-5* loss-of-function lines to also exhibit an early flowering
phenotype (Figure 1D - E). Upon comparison, *raf24-2* showed the strongest early flowering
phenotype, which corresponded to its lower *RAF24* mRNA levels relative to *raf24-4* and *raf24-5*
(Supplemental Figure 2C). In addition to modulating flowering time, we also found that *raf24*
mutants have a significantly smaller leaf area compared to WT Col-0 plants, indicating that
RAF24 likely has additional developmental roles beyond flowering time regulation
(Supplemental Figure 3).
158
Quantitative phosphoproteomic analysis of *raf24-2* mutant reveals changes in flowering
related proteins.
To elucidate the underlying molecular mechanisms by which RAF24 modulates flowering,
we conducted a quantitative phosphoproteomic analysis of *raf24-2*. Here, we found a decrease in
VCS (AT3G13300) phosphorylation, which is consistent with the previously described role of
RAF24 in the SnRK2s-VCS module (Soma *et al.*, 2017; Kawa *et al.*, 2020). We also found
several flowering-related proteins to be significantly less phosphorylated in the *raf24-2* mutant
compared to WT Col-0 (Table 1; Supplemental Data 1). These included proteins involved in
regulating *FLC* expression, such as: VERNALIZATION INDEPENDENCE 4 (VIP4;
AT5G61150), ARABIDOPSIS SF1 HOMOLOG (SF1; AT5G51300) and HISTONE MONO-
UBIQUITINATION 2 (HUB2) (Zhang & van Nocker, 2002; Lee *et al.*, 2020). In addition,
several FLC-independent flowering phosphoproteins also exhibited a decreased phosphorylation
status in *raf24-2* plants (Table 1). These included: BRI1-EMS-SUPPRESSOR 1 (BES1;
AT1G19350), a positive regulator of photoperiodic flowering through the promotion of FT
expression and FLOWERING BHLH 3 (FBH3; AT1G51140), a flowering regulator that
promotes the expression of CO (Wang *et al.*, 2019; Sanagi *et al.*, 2021). Taken together, this
suggests that RAF24 may have wide-ranging, higher-order control over flowering signaling.
Among these flowering-related phosphoproteins, phosphorylation of HUB2 (AT1G55250) at site
S^{314} exhibited the most dramatic decrease in abundance, being below detection in *raf24-2*

relative to WT Col-0 (Table 1). Consequently, we hypothesized that RAF24 plays a key role in
regulating flowering time through modulation of HUB2 phosphorylation at S^{314}.
180
RAF24 and HUB2 co-localize to the nucleus.
To better understand the molecular role of RAF24, we next investigated the subcellular
localization of RAF24 through the transient expression of c-terminal YFP-tagged RAF24
(RAF24-YFP) in *Nicotiana benthamiana* leaf epidermal cells. Consistent with previous
observations (Soma *et al.*, 2020), RAF24-YFP is predominantly localized to the cytoplasm
(Supplemental Figure 4), with smaller quantities of RAF24-YFP found to localize to the nuclei
(Supplemental Figure 4). Next, we then transiently co-expressed RAF24-YFP and mCherry-
HUB2 in the *N. benthamiana* to test if they co-localize in the plant cell. Here, we see that a major
proportion of RAF24-YFP co-localizes with mCherry-HUB2 in the cell (Figure 2).
190
Genetic analysis reveals RAF24 and HUB2 control flowering time and the expression of
FLC* and *FT
As a result of previous studies demonstrating *hub2* deficient plants to possess an early
flowering phenotype (Cao *et al.*, 2008; Gu *et al.*, 2009), we next explored the potential of a
genetic interaction between RAF24 and HUB2 by generating double loss-of-function plant lines.
Under a 16:8 photoperiod, *raf24-2* and *hub2-1* single mutants flower earlier than Col-0 as
expected, while *raf24-2 hub2-1* double mutants were found to flower slightly earlier than *hub2-*
1, but not *raf24-2* (Figure 3A - B). However, upon evaluating the same mutant lines under a
12:12 photoperiod, we found that the *raf24-2 hub2-1* plants flower dramatically earlier than both
parental single mutants (Figure 3C - D). Collectively, this indicates that RAF24 and HUB2 act
synergistically to influence flowering time and that to some extent, this genetic interplay between
HUB2 and RAF24 is photoperiod dependent. The additive effects of *raf24-2 hub2-1* in
accelerating flowering time also indicates likely intersections with the other flowering targets
observed to possess reduced phosphorylation in the *raf24-2* mutant.
HUB2 is known to regulate flowering pathways by modulating *FLC* transcription through the
monoubiquitination of H2B (Cao *et al.*, 2008; Gu *et al.*, 2009). Therefore, we hypothesize that
RAF24 may also affect FLC mRNA levels by modulating HUB2 phosphorylation to fine-tune
flowering time. To assess this, we examined *FLC* expression levels in *raf24-2*. Consistent with
the accelerated flowering time phenotype, *FLC* transcription is significantly decreased in *raf24-2*
mutants relative to WT Col-0 (Figure 4A). Simultaneously, we also assessed the expression
levels of *FT* and *SOC1*, downstream floral integrators whose expression levels are directly
targeted by FLC. Here, we find that *FT* expression is significantly higher, while *SOC1*
expression remains unchanged, indicating that RAF24 regulates floral development via FLC and
FT (Figure 4A). We next validated that flowering time repression by RAF24 is FLC and FT
dependent by crossing *raf24-2* with *flc-6* and *ft-10* mutants, respectively. Here, *raf24-2 flc-6*
double mutants flowered just as early as each single mutant plants, respectively, suggesting that
the loss of *FLC* was epistatic to *RAF24* and the premature flower development in *raf24-2* was

due to *FLC* function (Figure 4B & C). Conversely, *raf24-2 ft-10* plants flowered an average of
7.2 days earlier than *ft-10.* Collectively, this suggests that RAF24 modulates floral development
in a FLC and FT dependent manner (Figure 4B & C).
221
RAF24 suppresses mono-ubiquitination of H2B by HUB2
HUB2 is responsible for H2B mono-ubiquitination, which has been suggested to promote
expression of flowering genes (Cao *et al.*, 2008; Gu *et al.*, 2009). In humans, the phosphorylation
of RNF20/40 (the human homolog of HUB1/HUB2) (Moyal *et al.*, 2011) promotes H2Bub1 and
activation of DNA repair responses. Therefore, we hypothesized that RAF24 may regulate the
phosphorylation status of HUB2 in order to modulate H2Bub1 levels, thereby influencing FLC
expression and the control of flowering time. To test this, we performed immunoblotted WT Col-
0 and *raf24-2* plants constitutively over-expressing HUB2 using anti-H2Bub1 and -H2B
antibodies (Figure 5A & B). Here, we found a higher accumulation of H2Bub1 in *raf24-2* plants
over-expressing a tandem affinity purification (TAP)-tagged HUB2 construct (TAP-HUB2),
suggesting that RAF24 suppresses H2B mono-ubiquitination by phosphorylating HUB2 (Figure
5A & B). At the same time, we find that *raf24-2 hub2-1* and *hub2-1* plants maintain no H2Bub1
as expected, validating the effectiveness of the anti-H2Bub1 antibodies (Supplemental Figure 5).
Surprisingly, we find that TAP-HUB2/Col-0 plants also display an early flowering phenotype
when compared to WT Col-0 as previously reported (Woloszynska *et al.*, 2019) (Figure 5C),
suggesting HUB2 likely operates as an integration point for multiple flowering related pathways,
with optimal HUB2 levels required for fine-tuning flowering time. Moreover, over-expression of
HUB2 in *raf24-2* induces earlier flowering than in the WT Col-0 (Figure 5C), implying that
accumulation of H2Bub1 may also trigger inappropriate flowering time. Collectively, this
evidence indicates that RAF24 operates upstream of HUB2 to suppress H2Bub1 in order to fine-
tune FLC expression and appropriately regulate flowering time.
243
RAF24 affects protein-protein binding of HUB2.
By using RT-qPCR assays, we found that loss of *RAF24* does not affect the expression of
either HUB1 or HUB2 (Supplemental Figure 6; Supplemental Data 2), suggesting that RAF24
impacts HUB2 at the protein level, potentially through protein-protein interactions. As
previously reported, monoubiquitnation of H2B is suggested to rely on interactions between
UBC1/2 and HUB1/2 as well as the binding between HUB2 and H2B (Liu *et al.*, 2007; Cao *et
al.*, 2008). To better understand this, we performed affinity purification mass spectrometry (AP-
MS) experimentation using TAP-HUB2 constructs expressed in either WT Col-0 or *raf24-2*
(Supplemental Data 3). Here, we quantified all known interactors of HUB2, except for UBC1/2,
in both backgrounds, however, a direct interaction between HUB2 and UBC1/2 has only been
shown using yeast 2-hybrid and has not been validated *in vivo*. Importantly, we find the
interaction between TAP-HUB2 and HUB1 as unchanging when quantified in the presence or
absence of RAF24, suggesting that the phosphorylation of HUB2 is not vital for heterodimer
formation between HUB1 and HUB2 (Figure 6A). However, TAP-HUB2 is able to

immunoprecipitate significantly more H2B (AT5G22880) in *raf24-2* plants than that of WT Col-0, implying that RAF24-induced phosphorylation of HUB2 dampens its affinity for H2B as a substrate, suggesting a mechanism by which RAF24 suppresses H2Bub1.

Next, we queried our HUB2 interactome data for changes in the abundance of other known HUB2 interactors. Here we found a quantifiable decrease in the pulldown of the HUB2 interactor SPEN3 (AT1G27750), which is also known to affect flowering time (Woloszynska *et al.*, 2019). Compared to WT Col-0, where a functional RAF24 is present, SPEN3 went undetected in TAP-HUB2 pulldowns from *raf24-2* plants, while equal amounts of the protein KHD1 (AT1G51580) (Woloszynska *et al.*, 2019) was found in pulldowns from both WT Col-0 and *raf24-2* (Figure 6A) Supplemental Data 3). Further assessment of *SPEN3* and *KHD1* expression levels in both WT Col-0 and *raf24-2* by qRT-PCR found no change in their expression level (Supplemental Figure 6), confirming that absence of RAF24 affects the interaction between RAF24 and SPEN3, but not expression levels of *SPEN3*. Interestingly, SPEN3 is an RNA binding protein that is suggested to promote *COOLAIR* transcript formation, a non-coding transcript that can repress *FLC* expression (Li *et al.*, 2015; Eom *et al.*, 2018). Using RT-qPCR we find a significant reduction in total *COOLAIR* transcripts in *raf24-2* mutants, suggesting that loss of RAF24 not only impairs the association of HUB2 with SPEN3, but it also has a measurable effect on *COOLAIR* mRNA abundance (Figure 6 A, B&C; Supplemental Data 2).

Apart from these known flowering-related protein interactors of HUB2, we also found several characterized flowering regulators, including VERNALIZATION INDEPENDENCE 3 (VIP3; AT4G29830), HISTONE ACETYLTRANSFERASE OF THE MYST FAMILY 1/2 (HAM1/2; AT5G64610/ AT5G09740) and GLYCINE RICH PROTEIN 7/8 (GRP7/8; AT2G21660/AT4G39260), were pulled down by TAP-HUB2. (Figure 6D; Supplemental Data 3). Many candidate binding partners regulate flowering time in an FLC-dependent manner, but in different signaling pathways. VIP3 and EARLY FLOWERING 7 (ELF7; AT1G79730) being part of *Arabidopsis* relative of yeast RNA POLYMERASE II ASSOCIATED FACTOR 1 (PAF1) complex, modulate vernalization-responsive flowering time via FLC (Zhang *et al.*, 2003; He *et al.*, 2004; He, 2009; Cao *et al.*, 2015), further supporting the previous hypothesis that HUB2 and PAF1 play roles in the same pathway (Wright *et al.*, 2011). In addition, MULTICOPY SUPPRESSOR OF IRA1 4 (MSI4/FVE; AT2G19520) acts as a primary repressor of FLC in autonomous flowering signaling (Kim *et al.*, 2004; Sun *et al.*, 2021). Together, this suggests that in addition to affecting H2Bub1 levels, HUB2 may play additional regulatory roles in FLC expression by interacting with different binding partners upon upstream environmental cues. On the other hand, SHORT VEGETATIVE PHASE (SVP; AT2G22540), which serves as a transcriptional repressor of flowering in concert with FLC, is also found as a candidate interactor of HUB2 and has been found to be controlled by the 26S proteasome (Lee *et al.*, 2013; Jin & Ahn, 2021). However, unlike the human HUB1/2 ubiquitin ligase orthologs RNF20/40, HUB2 has not been found to polyubiquitinate substrates for proteasomal degradation (Jeon *et al.*, 2020). Therefore, future studies could investigate whether HUB2 polyubiquitinates SVP and if this affects its control over flowering initiation.

SnRK2s interact with and phosphorylate HUB2 *in planta*.

From our TAP-HUB2 pull-down data, we also found a number of protein kinases that possibly interact with HUB2 (Figure 7A). In particular, SnRK2s and CASEIN KINASE II (CK2), which have each been implicated in the regulation of flowering time (Figure 7A) (Ogiso *et al.*, 2010; Mulekar & Huq, 2012; Wang *et al.*, 2013). Interestingly, HUB2 was previously found to be a direct phosphorylation target of SnRK2.6 (Wang et al., 2020). Given that SnRK2s are phosphorylated and activated by RAF24 (Lin et al., 2020; Soma et al., 2020; Fàbregas et al., 2020), we hypothesized that RAF24 may impact HUB2 phosphorylation through the SnRK2s. To test this, we first confirmed which SnRK2s interact with HUB2. Using a bimolecular fluorescence (BiFC) assay through transient co-expression of nYFP-HUB2 with each of the *Arabidopsis* SnRK2s, we find that fluorescence was only reconstituted with SnRK2.2-cYFP, SnRK2.4-cYFP or SnRK2.6-cYFP (Figure 7B; Supplemental Figure 7). These results suggest that SnRK2.2, SnRK2.4 and SnRK2.6 can interact with HUB2 *in vivo*, corroborating previous findings that SnRK2.4 and SnRK2.6 are directly phosphorylated by RAF24 (Lin et al., 2020; Soma et al., 2020). We next focused on the relationship between SnRK2.4 and SnRK2.6 with HUB2. First, we conducted split-luciferase complementation assays to further confirm interactions between HUB2-SnRK2.4 and -SnRK2.6. Here, we detect a significantly larger luciferase signal when co-expressing HUB2 with SnRK2.4 and SnRK2.6, versus SnRK2.8 (Figure 7C). Next, we tested if SnRK2.4 and SnRK2.6 phosphorylate HUB2 by conducting an *in vitro* ADP-Glo protein kinase activity assay. We found that HUB2 assays conducted with SnRK2.4-His$_6$ or SnRK2.6-His$_6$ displayed significantly higher relative light units (RLUs), implicating SnRK2.4 and SnRK2.6 in the phosphorylation of HUB2 *in vitro* (Figure 7D). Lastly, we examined if SnRK2.4 and SnRK2.6 phosphorylates HUB2 *in vivo*. To this end, we co-expressed the TAP-HUB2 and SnRK2s-YFP in *Nicotiana benthamiana*. Upon isolating TAP-HUB2, we found HUB2 to possess increased phosphorylation when co-expressed with SnRK2.4 or SnRK2.6, compared to that of SnRK2.8 (Figure 7E). Collectively, these results indicate that SnRK2.4, in addition to SnRK2.6, physically interact with, and phosphorylates, HUB2.

Phosphorylation of HUB2 at S^{314} represses flowering and ligase activity.

Lastly, to understand if phosphorylation of HUB2 at S^{314} translates to control of flowering time, we next created phospho-mimetic (HUB2^{S314D}) and phospho-ablative (HUB2^{S314A}) expressing transgenic *raf24-2* plants. We observed HA-HUB2^{S314D} expressing *raf24-2* plants to possess a significant delay in flowering time relative to both the HA-HUB2^{S314A} and HA-HUB2^{S314S} *raf24-2* plants (Figure 8A&B). The delayed flowering phenotype indicates that phosphorylation of HUB2 at S^{314} in the *raf24-2* background directly influences the flowering time of Arabidopsis. In addition, we observe higher H2Bub1 levels in HA-HUB2^{S314A} than HA-HUB2^{S314D} in the *raf24-2* background (Figure 8C), further confirming that phosphorylation of HUB2 at S^{314} suppresses its ligase activity.

DISCUSSION

Recent research has resolved roles for RAF-like MAPKKK kinases in abscisic acid (ABA) (Lin *et al.*, 2021; Kamiyama *et al.*, 2021), osmotic signaling (Lin *et al.*, 2020; Soma *et al.*, 2020), auxin signaling (Kuhn et al., 2024) and embryonic development (Wang *et al.*, 2018b), with functions in flowering time remaining undefined. Here we identify a novel function for RAF24 in the control of flowering, which diverges from its shared roles with other B4 RAF MAPKKK kinases in osmotic stress responses where their phosphorylation of SnRK2 kinases facilitates down-stream signaling (Lin *et al.*, 2020; Soma *et al.*, 2020) and auxin signaling (Kuhn et al., 2024). Our identification of an independent role for RAF24 can be explained by its divergent phylogenetic relationship to its most closely related orthologs RAF18 and RAF20 and by their differential gene expression profiles, where we find RAF24 to be expressed in rosette leaves, versus a predominant root expression profile for RAF18 and RAF20 (MAPK Group, 2002; Soma *et al.*, 2020). This suggests an intrinsic difference between the function of different B4 RAF MAPKKK protein kinases as it pertains to developmental regulation versus stress responses. In particular, in the early onset of abiotic stress, plants may employ several kinases to rapidly transduce stress signals, however, a single protein might be sufficient to induce certain changes in plant development.

RAF24 sits high in flowering-time signaling, and may crosstalk with brassinosteroid and/or abiotic stress signalling.

Here we find evidence that suggests RAF24 regulates flowering initiation through a number of pathways. First, phosphoproteomic analysis of the *raf24-2* mutant found several phosphoproteins related to flowering are potential RAF24 substrates (Table 1). Among these, VIP4 is responsible for vernalization-gated flowering time, while BES1 and FBH3 are involved in the photoperiod flowering regulation (Zhang & van Nocker, 2002; Ito *et al.*, 2012; Wang *et al.*, 2019)(Table 1). These results suggest RAF24 might gate signal transduction of multiple flowering pathways. In addition, RAF24 also may play roles in brassinosteroid (BR) signaling. Several BR-related phosphoproteins, BES1, BRI1 SUPPRESSOR 1 (BSU1)-LIKE 1 (BSL1/2; AT4G03080 / AT1G08420) and BES1/BZR1 HOMOLOG 4 (BEH4; AT1G78700), displayed significantly lower phosphopeptide abundance in *raf24-2* phosphoproteomic analysis (Table1). The potential involvement of RAF24 in BR signaling may explain why *raf24-2* mutants exhibited smaller rosette compared to WT, as BR is known to regulate cell expansion (Haubrick & Assmann, 2006; Zhiponova et al., 2013) (Supplemental Figure 3). In fact, BR signaling is also associated with flowering time (Li et al., 2010; Li et al., 2018). Therefore, RAF24 may mediate flowering via crosstalk with BR signalling by directly or indirectly modulating phosphorylation status of BR-related proteins. Furthermore, we find that loss of RAF24 leads to lower HUB2 phosphorylation at S^{314} through SnRK2.4 and SnRK2.6, which directly interact with, and phosphorylate, HUB2. Considering that RAF24 phosphorylates both SnRK2.4 and SnRK2.6, and that SnRK2.6 has been reported to control flowering time in plants (Wang *et al.*, 2013; Chong *et al.*, 2022), our results indicate that RAF24 phosphorylates SnRK2.4 and SnRK2.6 to impact

HUB2 phosphorylation status and thereby fine tune flowering time. Furthermore, this RAF24-SnRK2s-HUB2 signalling module indicates RAF24 sits high in flowering initiation pathways. This differs from the flowering kinases SK12 and CPK32 described to date, which directly phosphorylate key flowering-time proteins, CO and FCA, respectively (Chen *et al.*, 2020; Li *et al.*, 2022). Recently, SnRK2.6 in tomato (*Solanum lycopersicum*) was reported to promote flowering time upon drought stress conditions (Chong *et al.*, 2022). Since HUB2 has been related to abiotic stress tolerance (Zhou *et al.*, 2017; Ma *et al.*, 2019; Chen *et al.*, 2019), it may be interesting to investigate whether RAF24-SnRK2s-HUB2 module mediates flowering time following exposure to abiotic stress stimulus. Therefore, further research can be conducted on roles of RAF24 in mediated flowering time under varying hormone or abiotic stress conditions.

Appropriate flowering time requires H2Bub1 homeostasis, which depends on tightly regulated HUB2 ligase activity by phosphorylation

Similar to its human ortholog, we find HUB2 to also be a phosphoprotein in plants, with its phosphorylation status involved in controlling HUB2 functionality and maintaining appropriate H2Bub1 levels (Moyal *et al.*, 2011). So far, H2Bub1 homeostasis has been linked to flowering time by multiple studies (Cao *et al.*, 2008; Schmitz *et al.*, 2009; Woloszynska *et al.*, 2019), with *hub2* plants displaying lower H2Bub1 levels and an earlier flowering time (Cao *et al.*, 2008). Conversely, loss of UBIQUITIN-SPECIFIC PROTEASE 26 (UBP26) results in an accumulation of H2Bub1, and also early flowering initiation (Schmitz et al., 2009). Therefore, we propose that precise phospho-mediate regulation of H2Bub1 levels are required in order to avoid premature flowering. Our results support this model, as loss of RAF24 leads to a significant increase in H2Bub1 levels, therefore accelerating flowering initiation. Similarly, in the *raf24-2* background, HA-HUB2^{S314A} had increased accumulation of H2Bub1 and accelerated flowering relative to HA-HUB2^{S314D}, indicating that phosphorylation of HUB2 acts to suppresses the overaccumulation of H2Bub1 to avoid premature flowering. However, HUB2^{S314D} did not completely rescued the early flowering phenotype compared to the *raf24* mutant, as an overall increase in HUB2 levels also initiates premature flowering (Woloszynska *et al.*, 2019). However, HA-HUB2^{S314D} did complement the early flowering phenotype in *raf24* plants expressing comparable levels of HUB2 (HA-HUB2^{S314S}).

To ensure H2Bub1 homeostasis, HUB2 ligase activity needs to be tightly regulated. In Arabidopsis, activity of ubiquitin ligases has been found to be modulated by protein kinases, such as SnRK2s (Ding *et al.*, 2015a,b). For example, SnRK2.6 phosphorylates RING-type CHY ZINC-FINGER AND RING PROTEIN1 (CHYR1) to enhance its ligase activity in ABA and drought response signaling (Ding *et al.*, 2015b). In addition, SnRK2.6 also phosphorylates PUB25/PUB26 in response to freezing treatment, therefore increasing its E3 ligase activity (Ding *et al.*, 2015a). Likewise, here we show that SnRK2.6, as well as SnRK2.4, interacts with and phosphorylates HUB2. So far, H2Bub1 is the only characterized substrate of HUB2 (Liu et al., 2007). Therefore, it is reasonable to use changes in H2Bub1 levels to reflect the ligase activity of HUB2. Correspondingly, we observed higher H2Bub1 levels in the absence of RAF24,

indicating that RAF24 functions to suppress HUB2 ligase activity. Interestingly, H2Bub1 levels are known to be influenced by osmotic stress in Arabidopsis (Chen *et al.*, 2019), and by ABA treatment in rice (Ma *et al.*, 2019); both biological responses related to RAF24 (Lin *et al.*, 2020; Soma *et al.*, 2020). Since SnRK2.4 and SnRK2.6 are also activated by osmotic stress and ABA, respectively, it is possible that changes of H2Bub1 levels in respose to abiotic stress, are in part, controlled by different SnRK2s and their phosphorylation of HUB2.

In addition to exerting roles in flowering pathway, HUB2 is also known to regulate many other biological processes, but specific protein interactors of HUB2 in these biological contexts have not been determined. Our TAP pulldown results reveal an array of candidate HUB2 interactors involved in diverse signaling pathways linked to a wide range of biological processes (Supplemental Data 3). Beyond flowering, HUB2 also functions in plant innate immunity (Hu *et al.*, 2014; Zou *et al.*, 2014). Here, we find immunity-related proteins MODIFIER OF SNC1, 2 (MOS2; AT1G33520), MODIFIER OF SNC1,4 (MOS4; AT3G18165), PECTIN METHYLESTERASE 17 (PME17; AT2G45220) and RECOGNITION OF PERONOSPORA PARASITICA 13 (RPP13; AT3G46530) present as candidate binding partner for HUB2 (Bittner-Eddy *et al.*, 2000; Rose *et al.*, 2004; Zhang *et al.*, 2005; Monaghan *et al.*, 2009; Del Corpo *et al.*, 2020). In addition, HUB2 affects the dynamics of microtubules (Hu *et al.*, 2014), and here we find multiple microtubule-associated proteins MICROTUBULE-ASSOCIATED PROTEINS 65-1 (MAP65-1; AT5G55230) and MAP65-6 (AT2G01910) as candidate HUB2 interactors (Mao *et al.*, 2005; Lucas *et al.*, 2011). Moreover, RESPONSIVE TO DEHYDRATION 21A (RD21A; AT1G47128), SNW/SKI-INTERACTING PROTEIN (SKIP; AT1G77180) and ABA INSENSITIVE 8 (ABI8; AT3G08550) are also present as candidate HUB2 interactors, implicating HUB2 in abiotic stress resistance (Brocard-Gifford *et al.*, 2004; Ma *et al.*, 2019; Chen *et al.*, 2019; Liu *et al.*, 2020; Zhang *et al.*, 2022), with RD21A having been reported to be polyubiquitinated (Kim & Kim, 2013). Thus, it is possible that HUB2 may also be involved in the polyubiquitination of RD21A for 26S proteasomal degradation. Lastly, H2Bub1 implicated in circadian regulation through candidate HUB2 interactors BASIC LEUCINE ZIPPER 63 (bZIP63; AT5G28770) and GRP7 (AT2G21660)(Heintzen *et al.*, 1997; Staiger *et al.*, 2003; Frank *et al.*, 2018). Further studies based on such protein-protein interaction information could help advance our insights into HUB2's diverse roles in regulating plant development and fitness.

In summary, we propose a model whereby RAF24 functions to repress flowering initiation by modulating the phosphorylation of HUB2 through SnRK2.4 and SnRK2.6 (Figure 9A). In the presence of RAF24, HUB2 phosphorylation is maintained, which in turn optimizes HUB2ub1 levels and maintains effective transcriptional activation of FLC (Figure 9A). Upon the loss of RAF24 HUB2 is no longer phosphorylated, resulting in an over-accumulation of H2Bub1, and reduced transcriptional activation of FLC (Figure 9A). In addition, we find that RAF24 acts upstream of a variety of diverse flowering time pathways, suggesting a wide-ranging role in fine-tuning floral development in response to different internal or external cues that goes beyond HUB2 (Figure 9B). Moreover, the diverse set of HUB2 candidate interactors found here

458 indicates that HUB2 may function as part of a broad series of protein complexes that impact
459 flowering initiation (Figure 9B), each of which warrants further investigation in future studies.
460
461

METHODS
Plant Growth and Phenotyping Assays
Plant Materials and Growth Conditions - All Arabidopsis seeds used in this study were based
on Col-0 ecotype. T-DNA insertion lines, *raf18-1*, *raf20-1*, *raf24-2*, *raf24-4*, *raf24-5*, *hub2-1*,
flc-6 as well as *ft-10* were obtained from The Arabidopsis Information Center and confirmed by
genotyping evidence. The *raf24-2*, *hub2-1* were crossed to generate *raf24-2 hub2-1* double
mutants. Similarly, *raf24-2*, *flc-6* and *ft-10* mutants were crossed to create *raf24-2 flc-6* and
raf24-2 ft-10 double mutants. All genotyping primers are listed in Supplemental Table 1.
Arabidopsis seeds were stratified under 4°C and dark conditions for 3 days. Then all seeds were
moved to a 16:8 or 12:12 light : dark photoperiod at 22 °C with a light intensity of 100
µE m−2 s−2 unless otherwise specified.
Leaf area measurement - Each chamber was equipped with a single-board Raspberry Pi 3 B+
computers and ArduCam Noir Cameras (OV5647 1080p) with a motorized IR-CUT filter and
two infrared LEDs. Pictures of plants were taken every 15 min over seven days (between 14-d
and 20-d post-seed imbibition). Using the open-source software package PlantCV
(https://plantcv.readthedocs.io/), the areas of each plant in the picture collection were extracted
and quantified as described (Gehan *et al.*, 2017).
479

Molecular Cloning and Transgenic Plant Generation
N-terminal TAP-HUB2G plants were created using the 35S::pKNGSrhino-HUB2G (TAP-
HUB2) construct (Woloszynska *et al.*, 2019). *Agrobacterium tumefaciens* GV3101 were
propegated in 2ml liquid culture consisting of LB media containing 50 µg/µl Spectinomycin, 25
µg/µl Rifampicillin and 40 µg/µl Gentamicin at 28°C 150 rpm for 48 h followed by additional 24
h sub-culturing at 1:20 dilution with the same antibiotic regiment. *Agrobacterium* was then
pelleted and re-suspended in 40 ml 5% (w/v) sucrose, 10 ml LB and 3% (v/v) Silwet L-77
immediately prior to transfection. Floral dipping was performed as previously described (Clough
& Bent, 1998). The transgenic seeds were selected on 0.5x MS addition with 50 µg/µl
Kanamycin plates and screened under compound microscope (Olympus BX51 Fluorescence
Microscope®) for fluorescence, prior to confirmation by western immunoblotting. Subcellular
localization assays utilized RAF24 and HUB2 cloned into pK7YWG2 and pEearlygate104
plasmids, respectively. For BiFC assays, the full-length coding sequence of each SnRK2 and
HUB2 were cloned into pB5GWcY and pB5nYGW constructs, respectively. Lastly, HUB2
phospho-mimetic and phospho-ablative over-expressing lines were created using site-directed
mutagenized 35S::pKNGSrhino-HUB2G (Supplemental Data 2), followed by insertion into the
binary vector pEarlygate201. After transformation into GV3101, *raf24-2* plants were transfected
to generate to HA-HUB2^{S314}, HA-HUB2^{S314A}, or HA-HUB2^{S314D} plant lines.

Quantitative Phosphoproteome Analysis

Protein extraction, quantification, digestion and phosphopeptide enrichment were performed as previously described (Uhrig *et al.*, 2019). Dissolved peptides were injected using an Easy-nLC 1000 system (Thermo Scientific) and separated on 50 cm ES803 Easy-Spray PepMap C18 Column (Thermo Scientific). Peptides were eluted using the following gradient of solvent B (0.1% (v/v) FA in 80% (v/v) ACN): 0–50 min, 0–25% B; 50–60 min, 25–32% B; 60–70 min, 32–98% B, at a flow rate of 300 nL/min at 50^{o}C. Mass spectra were acquired using an Orbitrap Hybrid mass spectrometer (Thermo Scientific) operated in data-dependent acquisition mode as previously described (Uhrig *et al.*, 2019). Raw data were analyzed using MaxQuant software version 2.0.3.0 (https://www.maxquant.org/; (Cox & Mann, 2008; Tyanova *et al.*, 2016)) using default settings, which included: fixed modifications (cysteine carbamidomethylation; C), variable modifications (serine, threonine, and tyrosine phosphorylaton; pSTY & methionine oxidation; M$_{ox}$) and peptide, protein and PSM false discovery rate of 1%. Data outputs were processed using Perseus software version 1.6.15.0 (https://www.maxquant.org/; (Tyanova *et al.*, 2016) and subsequently filtered for significantly changing phosphopeptides (students t-test; *p-value* < 0.05) with a data completeness of ~65% (5 / 8 samples) and a PTM-site probability score ≥ 0.75.

Affinity Purification Mass Spectrometry (AP-MS)

Positive TAP-HUB2 in Col-0 (TAP-HUB2/Col-0) and *raf24*-2 (TAP-HUB2/*raf24-2*) seedlings were grown for 14 d and harvested. Proteins were extracted using 50 mM HEPES-KOH (pH7.5), 100 mM NaCl, 0.1% (v/v) NP-40, 50 mM NaPPi, 1 mM Na$_3$VO$_4$, 2 mM PMSF, 1x Roche cOmplete™ EDTA-free protease tablet and 1x Roche PhosSTOP™ phosphatase inhibitor cocktail tablet (Roche). Extracts were clarified at 20,000 x g for 20 minutes at 4^{o}C and then measured by Bradford Assay (Bradford, 1976). Extracts were then incubated with custom magnetic IgG-coupled beads (Cytiva) end-over-end for 1 h at 4^{o}C to bind TAP-HUB2. IgG-coupled beads were then washed 3x for 5 min with 0.1% (v/v) NP-40 and 4 washes with no NP-40, followed by elution using 100 mM Glycine pH 3.0 neutralized with 1M Tris-HCl. TAP-HUB2 pull-downs were then reduced (10mM Dithiothreitol; 45min @ 37^{o}C), alkylated (55 mM Iodoacetamide; 1h @ 23^{o}C) and digested in solution with 1:100 trypsin (V5113; Promega). Peptides were acidified to 0.5% (v/v) trifluoroacetic acid (TFA; Sigma), dried, dissolved in 3% (v/v) ACN / 0.1% (v/v) TFA and desalted as previously described (Uhrig *et al.*, 2019). Differences in the Col-0 and raf24-2 HUB2-protein interactome were quantified by BoxCarDIA LC-MS/MS analysis as previously described (Mehta et al., 2022). Acquired data was then analyzed using Spectronaut v14 (Biognosys AG) using default settings. Candidate HUB2 interactor abundance was then normalized against HUB2 levels to obtain relative fold-changes in abundance between each genotypic background.

Co-localization and Bimolecular fluorescence complementation (BiFC) analysis

RAF24-YFP, mCherry-HUB2, nYFP-HUB2, SnRK2s-cYFP as well as PRP4Kc-cYFP were transformed into *Agrobacterium* through electroporation. Positive colonies were selected, cultured and harvested in infiltration buffer (10 mM MES pH 5.6, 10 mM MgCl$_2$, 100 µM acetosyringone) as described (Schütze *et al.*, 2009). Cell suspensions were diluted to OD = 0.8 to 1.0 and incubated in dark for at least 2 hours at room temperature before injecting 4-week-old *N. benthamiana* plants. Infected plants were grown for 2 - 3d, with infiltrated sections analyzed with a Leica SP8 STED or ZEISS LSM 710 facilities.

Split-Luciferase Complementation Assay

 Split-Luciferase complementation assays were conducted based on previously reported (Chen et al., 2008). *A. tumefaciens* GV3101 harboring cLUC-HUB2 (pGWB-CLUC) and SnRK2s-nLUC (pGWB-NLUC) were infiltrated into *N. benthamiana* as described above. Three days after infiltration, leaf discs were collected and placed into 96-well plate. Then 1 mM D-luciferin (Cayman, 25836) was applied into each well and incubated for 10-15 minutes in dark before luminescence measurement with the SPARK® multimode microplate reader (TECAN).

Recombinant Protein Expression And in Vitro ADP-Glo Kinase Assays

 His$_6$-HUB2 (pDEST17) and SnRK2s-His$_6$ (pET-DEST42) were used for protein expression. Briefly, BL21-CodonPlus cells containing His$_6$-HUB2 and SnRK2.4-His$_6$ were grown at 4 ^{o}C for 48 hours, while SnRK2.6-His$_6$ and SnRK2.8-His$_6$ were grown at room temperature for 24 h. Lysis was performed by French Press, with whole cell extracts cleared by centrifuged at 20000 x g at 4 ^{o}C degree for 30 minutes. Supernatants were retained and passed by miracloth, then incubated with His$_6$-tag Purification Resin (Roche, 05893682001) at 4 ^{o}C for 1 hour. This was followed by three washes buffer A (50 mM HEPES pH 7.5, 500 mM NaCl, 1% (v/v) Tween-20) and three washes buffer B (50 mM HEPES pH 7.5, 150 mM NaCl. Proteins were eluted (50 mM HEPES, 150 mM NaCl, 300 mM imidazole) and concentrated using 10000 MWCO concentrator spin-columns (Millipore). Final protein concentrations were measured using Bradford's reagent (Thermo Fisher).

 Phosphorylation of HUB2 by SnRK2s was examined with ADP-Glo Protein Kinase Activity Assay Kit (Promega, V9101) with slight modifications (Kathania *et al.*, 2022). 2 µg of His$_6$-HUB2 was mixed with or without 1 ug of SnRK2s-His$_6$ in kinase reaction buffer (50 mM Tris-HCl pH 7.5, 20 mM MgCl$_2$, 100 µM ATP, 2 mM DTT and 1 mM EGTA) and incubated at 30 ^{o}C for 1 hour and at room temperature for 2 hours. Assays were stopped by adding ADP-Glo reagents and incubation for 40 minutes in the dark. Kinase activity detection reagent was then added and kept at room temperature for 30 minutes, luminescence signals were quantified with a SPARK® microplate reader (TECAN).

In Vivo **Phosphorylation Assays**

 Agrobacterium harboring TAP-HUB2G and SnRK2s-GFP (pB7YWG2) were co-expressed in *N. benthamiana.* Leaves were collected and ground using a mortar and pestle after 3 d, then

extracted with protein extraction buffer (50 mM HEPES pH 7.5, 150 mM NaCl, 1 mM EDTA, 10% (v/v) glycerol, 2% (w/v) PVPP, 100 mM DTT, 0.1% (v/v) NP-40, 1 mM PMSF, 1x Roche cOmplete™ EDTA-free protease tablet and 1x Roche PhosSTOP™ phosphatase inhibitor cocktail tablet (Roche)) end-over-end for 30 minutes. Protein extracts were then incubated with custom magnetic IgG-coupled beads (Cytiva) for 1 hour at 4 ^{o}C and washed five times (50 mM HEPES pH 7.5, 100 mM NaCl). Beads were then mixed with 60 µL Tris-HCl pH 8.0 and 20 µL of 4x Laemmli Sample Buffer (BIO-RAD) for 10 minutes at 95 ^{o}C. Standard procedures were performed to visualize the TAP-HUB2, SnRK2s-GFP as well as phosphorylated HUB2 proteins by using anti-PAP (1:1000, SIGMA, P1291), anti-GFP (1:2000, ChromoTek, PABG1) and anti-phosphoserine antibodies (1:250, abcam, ab9332), respectively.

Western Immunoblot Analysis

Protein Extraction and Preparation – 10-day-old Arabidopsis seedlings were homogenized (2010 Geno/Grinder®) and then incubated with extraction buffer (4% (w/v) SDS, 50 mM Tris-HCl pH 8.0) for 30 minutes in an end-to-end rotator at 4 ^{o}C. Next, supernatants were retained after a 30-minute spin down with 16,000 x g at 4 ^{o}C. Protein estimation was conducted with Pierce BCA Protein Assay kit (Thermo Scientific). Finally, protein samples were mixed with 4x Laemmli Sample Buffer (BIO-RAD) for 5 minutes at 95 ^{o}C or 10 minutes at 65 ^{o}C.

Western Immunoblotting – Samples were loaded onto 7.5% or 15% SDS-PAGE gels and electro-transferred with Trans-Blot® SD Semi-Dry Transfer Cell (BIO-RAD) at 18 V for 45 minutes. After blocking 1 hour with 5% (w/v) non-fat milk, PVDF membranes were incubated overnight with anti-H2B (1:2500, abcam, ab1790), anti-H2Bub1 (1:1000, Cell Signaling, #5546), anti-GFP (1:2000, ChromoTek, PABG1), anti-HA(1:1000, Invitrogen, 26183), anti-H3 (1:2000, Invitrogen, PA5-16183) or anti-PAP (1:1000, SIGMA, P1291) followed by a 2-hour incubation with secondary antibodies, anti-rabbit (1:10000, Invitrogen, G-21234) or anti-mouse (1:5000, Invitrogen, G-21040).

Quantitative PCR Analysis

RNA extractions - RNA was extracted from 4 biological replicate pools of 2-week-old Arabidopsis seedling. 100 mg of seedling tissue was harvested and ground using Geno/Grinder® for 30 seconds at 1200 rpm. Total RNA was extracted using a modified TRIzol based protocol (Chomczynski & Sacchi, 1987). 1 ml of TRI Reagent® (Sigma-Aldrich, T9424) was added per 100 mg of tissue and incubated for 10 minutes at room temperature. Extracellular material was removed by centrifuging at 13000 x g for 10 minutes. Supernatant was acquired and combined with 200 µl of chloroform. Mixture was then inverted variously for 15 seconds and then left at RT for 3 minutes. Phase separation was achieved by centrifuging at 13000 x g for 15 minutes at 4 ^{o}C. The aqueous phase was transferred to new tubes. RNA was precipitated by adding 500 µl of 100% (v/v) isopropanol and then left to incubate for 10 min at RT. Sample was centrifuged at 13000 x g for 15 minutes at 4 ^{o}C to acquire RNA pellet. Supernatant was removed and the pellet was washed with 1 ml of 75% (v/v) ethanol. Sample was centrifuged once again at 7500 x g for 5

minutes at 4 $^{\circ}$C. Pellet was dried and subsequently resuspended in nuclease-free water. Total RNA was quantified by NanoDrop ND 1000 spectrophotometer (Thermo Scientific). Quality of RNA was assessed by visualizing 28S and 18S bands on a 1% (w/v) agarose gel stained with SYBRsafe (Thermo Fisher Scientific). Genomic DNA was removed using DNAse I kit (Thermo Fisher) followed by reverse transcription using RevertAid kit (ThermoFisher Scientific Inc) with 2 ug of starting RNA.

Transcript quantification - RT-qPCR was performed with 3 technical replicates per biological replicate using a proprietary mix developed, and distributed by the Molecular Biology Service Unit (MBSU), in the Department of Biological Science at the University of Alberta. The SYBR mix contains Tris (pH 8.3), KCl, $MgCl_2$, Glycerol, Tween 20, DMSO, dNTPs, ROX as a normalizing dye, SYBR Green (Molecular Probes, Life Technologies) as the detection dye, and an antibody inhibited Taq polymerase. A QuantStudio (TM) 6 Flex System was used, with PP2A3 (AT1G13320) deployed as the reference for normalization. Transcript abundance was measured using delta CT method. Primer sequences are listed in Supplemental Data 2.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Mieke Van Lijsebettens (Center for Plant Systems Biology, Belgium) for providing the 35S::pKNGSrhino-HUB2G construct, to acknowledge Dr. Hugo Zheng and Dr. Jiaqi Sun (McGill University) for sharing BiFC vectors and 35S::mCherry vectors, as well as thank Dr. Gitta Coaker (University of California, Davis) and Dr. Sonia Gazzarrini (University of Toronto) for providing SnRK2s clones. Further thanks go to Jack Moore of the Alberta Mass Spectrometry and Proteomics facility for mass spectrometry assistance.

DATA AVAILABILITY

All raw proteomics data can be found in PRoteomics IDEntification database (PRIDE; https://www.ebi.ac.uk/pride/) PXD038688.

AUTHOR CONTRIBUTIONS

QL conducted phenotypic analysis and characterized phospho-mutants, in addition to performing all WB, confocal microscopy, phosphorylation assays as well as designing experiments. LW generated all seed sources and performed phenotypic analysis, TAP-pulldown experiments, proteomic experimentation and qPCR analyses unless otherwise stated. MR performed all COOLAIR qPCR analysis. MT performed protein purificaiton and enzyme assays. DM assisted with experimental design and analysis. SS performed phenotypic analyses relating to plant surface area measurements and processing with PlantCV. RGU conceived of the study, while QL and RGU wrote the manuscript with editorial input from DM.

TABLES

Table 1: Uniquely differentially phosphophorylated proteins in *raf24-2*. Quantitative phosphoproteome analysis of *raf18-1*, *raf20-1* and *raf24-2* was performed. Proteins involved in flowering and exhibiting a phosphorylation status decrease unique to *raf24-2* are described. Arabidopsis gene identifier, common name, gene description, predicted localization (SUBA5; https://suba.live/), identified phosphorylation site and corresponding references of research involving each protein are presented.

FIGURES

Figure 1. Protein kinase RAF24 regulates flowering time. (A) Schematic genomic diagram of *RAF24* loci. Arrows indicate T-DNA insertion positions; exon 1 of *raf24-4*, exon1 and 2 of *raf24-5* as well as exon 5 and 6 of *raf24-2* mutants. Exons are denoted by black boxes. (B) Early flowering time observed in *raf24* mutants but not in related *raf18* and *raf20* mutants (Supplemental Figure 1). (C) Quantification of flowering time by days to bolting and rosette leaf number at bolting. (D) and (E) Quantification of early flowering in *raf24* mutants by days to bolting and rosette leaf number at bolting. Different letters indicate significant differences among genotypes (one-way ANOVA *p-value < 0.05*).

Figure 2. RAF24 co-localizes with HUB2. RAF24-YFP and mCherry-HUB2 were transiently expressed in *N. benthamiana* leaves (See materials and methods). Fluorescence signals were visualized by confocal microscopy 2 days after injection. Scale bars = 20 μm.

Figure 3. Genetic interaction analysis of HUB2 and RAF24. (A) and (B) Flowering time comparison of *raf24-2*, *hub2-1* and *raf24-2 hub2-1* double mutants relative to WT. 29-day-old plants were photographed under 16:8 photoperiodic conditions and flowering initiation was scored by days to bolting and rosette leaf number at bolting. Calculations were conducted based on more than 12 plants of each genotype. (C) and (D) Flowering time comparison of *raf24-2*, *hub2-1* and *raf24-2 hub2-1* double mutants relative to WT. 56-day-old plants were photographed after being grown under a 12:12 photoperiod. Flowering time was scored by days to bolting and rosette leaf number at bolting. Letters indicate significant differences between genotypes (one-way ANOVA *p-value < 0.05*).

Figure 4. RAF24 fine-tunes flowering time through FLC and FT. (A) Transcriptional expression of *FLC*, *FT* and *SOC1* in 14d old WT and *raf24-2* plants. Stars denote a Student's t-test *p-value ≤ 0.05* (*) and *p-value ≤ 0.01* (**). (B) and (C) Flowering time comparison of *raf24-2 flc-6* and *raf24-2 ft-10* double mutants. 29-day-old plants (upper) and 46-day-old plants (bottom) were photographed (B), with flowering time determined by days to bolting (upper) and rosette number at bolting (bottom) (C). Letters indicate significant differences between genotypes (one-way ANOVA *p-value < 0.05*).

Figure 5. RAF24 suppress the monoubiquitination of H2B by HUB2. (A) and (B) Loss of RAF24 increases H2Bub1 levels. Immunoblotting analysis of 10d-old WT and *raf24-2* plants constitutively over-expressing TAP-HUB2. 50 μg of protein samples were loaded in each lane. Biological replicates (n=3) were examined and quantified with ImageJ software, with levels of H2Bub1 normalized to H2B levels. (C) and (D) Over-expression of TAP-HUB2 in *raf24-2* also leads to an early flowering phenotype. Two independent lines of 29d-old WT and *raf24-2* plants (denoted by the number) constitutively expressing TAP-HUB2 were grown and imaged (C), with flowering time determined by days to bolting (upper) and rosette number at bolting (bottom) (D). Letters indicate significant differences between genotypes (one-way ANOVA *p-value < 0.05*).

Figure 6. Loss-of-RAF24 impacts the interaction of HUB2 with H2B and SPEN3. (A) TAP-HUB2 pull-downs from 14d-old WT and *raf24-2* plants revealed altered association of known HUB2 interactors (n = 4). Quantification values were normalized to the abundance of TAP-HUB2 in either the TAP-HUB2/WT or TAP-HUB2/*raf24-2* backgrounds, respectively. (B) and (C) Transcript abundance of total COOLAIR in 14d-old WT and *raf24-2* mutants. Levels of target transcripts were normalized to PP2A (n=4; see material and methods). Stars denote a Student's t-test *p-value ≤ 0.05* (*) and *p-value ≤ 0.01* (**). (D) Schematic of quantified flowering-related interactors detected in TAP-HUB2 pull-down experiments (Supplemental Data 3). Known interactors were represented by colored bubbles with a black line. Novel flowering-related candidate HUB2-interactors were shown as bold text with a dash line.

Figure 7. SnRK2.4 and SnRK2.6 interacts with and phosphorylates HUB2. (A) TAP-HUB2 pulldowns identified multiple protein kinases, including flowering-related SUCROSE NONFERMENTING 1-RELATED KINASE 2 (SnRK2) and CAESIN KINASE 2 (CK2) kinases. (B) Bi-molecular fluorescence complementation analysis validates interactions between HUB2 with SnRK2.4 and SnRK2.6 *in planta*. Co-infiltration of nYFP-HUB2 and SnRK2-cYFP constructs alongside PRP4Kc-cYFP (negative control) was performed using *N. benthamiana* leaves. Fluorescence signals were visualized with confocal microscopy 2 to 3d after infiltration (n > 10 images per set; Scale bars = 10 μm). (C) HUB2 interacts with SnRK2.4 and SnRK2.6, but not SnRK2.8 in luciferase complementation assays. Pairs of SnRK2s-nLUC and cLUC-HUB2 constructs were co-expressed using *N. benthamiana* leaves. After 3d incubation, luciferase intensity was quantified (n = 22 plants; Stars (*) denote one-way ANOVA *p-value = 0.0001* (***) or *p-value ≤ 0.0001* (****). (D) Phosphorylation of recombinant HUB2 by SnRK2.4 and SnRK2.6 *in vitro*. Relative HUB2 phosphorylation when mixed with recombinant SnRK2s or water (negative control). ADPglo luciferase signals were quantified after 3 h (n = 3; Stars denote one-way ANOVA *p-value < 0.05* (*) and *p-value ≤ 0.0001* (****)). (E) Phosphorylation of HUB2 by SnRK2.4 and SnRK2.6 *in vivo*. Co-expression of TAP-HUB2 and either YFP-tagged SnRK2.4, 2.6 or 2.8 was conducted through infiltrated *N. benthamiana* leaves. Immunoblotting analysis of each co-expression set was performed with anti-TAP, anti-GFP as well as anti-Ser IgG. 50 μg of input protein samples were loaded in each lane.

Figure 8. Phosphorylation of HUB2 at S^{314} controls flowering time. (A) and (B) Comparison of flowering time in *raf4-2* plants constitutively over-expressing HA-HUB2 with no (S314S), phospho-ablative (S314A) and phospho-mimetic (S314D) mutations (A). Flowering time was determined by days to bolting and rosette number at bolting (B). Letters indicate significant differences between genotypes (one-way ANOVA *p-value* < 0.05). (C) Immunoblot analysis of *raf24-2* plants constitutively over-expressing phospho-ablative (HA-HUB2^{S314A}), phospho-mimetic (HA-HUB2^{S314D}), which finds HA-HUB2^{S314D} maintains lower H2Bub1 levels. 50 µg of protein samples were loaded in each lane. Equal amounts of HA-HUB over-expression in the *raf24-2* background was observed across the lines depicted here (Supplemental Figure 7).

Figure 9. Roles of RAF24 in regulating HUB2 and flowering time. (A) In WT, the presence of RAF24 phosphorylates SnRK2s to induce phosphorylation of HUB2, ensuring appropriately tuned HUB2-ligase activity. This in turn optimizes HUB2ub1 levels to maintain proper transcriptional activation of FLC expression and flowering time (Figure 8A). Upon the loss of RAF24, SnRK2 fails to phosphorylate HUB2, causing an over-accumulation of H2Bub1 and reduced transcriptional activation of FLC expression. This results in earlier flowering (B) Both the phosphoproteomic and HUB2-TAP pulldowns find that RAF24 acts upstream of a wide variety of floral regulators, including BES1 and FBH3 as well as HUB2 to fine-tune flower time. Identification of both known HUB2 interactors (with straight line) and new, candidate HUB2 interactors (with a dash line) related to flowering time indicates that RAF24 possesses versatile functions / mechanisms in controlling flowering time.

SUPPLEMENTAL FIGURES

Supplemental Figure 1. Distinct phopsho-motifs of B4 like RAFs are phosphorylated in response to diel or osmotic stress changes. Relative abundance and retrieved from published phosphoproteomic analysis.

Supplemental Figure 2. Loss-of-RAF24 induces premature flowering. (A) and (B) Early flowering time of *raf24* mutants but not *raf18* and *raf20* mutants. 56-day-old plants were photographed under 12:12 photoperiodic conditions and flowering time scored by days to bolting and rosette leaf number at bolting. Letters indicate significant differences between genotypes (one-way ANOVA *p-value < 0.05*). (C) Relative expression analysis of *RAF24* in each *raf24* mutant. Transcript levels of *RAF24* were quantified using 14d-old *raf24-2*, *raf24-4* and *raf24-5* mutants with P1 and P2 primer pairs amplifying the 5' and 3'ends of RAF24, respectively (Supplemental Data 2). Data were normalized to UBC21 expression levels and then relative to *RAF24* transcriptional levels in WT background.

Supplemental Figure 3. Loss-of-RAF24 displays a smaller leaf area. (A) and (B) The leaf area of 29-days post-imbibition WT and *raf24-2* plants was measured as described (see materials and methods), with *raf24-2* plants displaying reduced leaf area. Plants were grown and photographed under either 16:8 (upper panel) or 12:12 (bottom panel) photoperiodic conditions.

Supplemental Figure 4. Localization of RAF24-YFP in *N. benthamiana* leaves. Fluorescence signal of RAF24-YFP was detected by confocal microscopy 3d after infiltration. Scale bars = 20 μm.

Supplemental Figure 5. Immunoblot analysis of H2Bub1 levels across *hub2* and *raf24-2/hub2* plants. Immunoblot analysis was performed using 10-day-old plants. 50 μg of protein samples were loaded in each lane. Abundance of H3 proteins serves as the loading control.

Supplemental Figure 6. Quantitative PCR analysis of known HUB2 interactors in WT and raf24-2 plants. Transcript quantification was performed using 14d-old WT and *raf24-2* plants. Data indicates that no statistically relevant change in transcript levels were found between WT and *raf24-2* genotypes. Student's t-test *p-value ≤ 0.05*.

Supplemental Figure 7. HUB2 specifically interacts with SnRK2.2, SnRK2.4 and SnRK2.6. *N. benthamiana* leaves were co-infiltrated with nYFP-HUB2 and each cYFP-SnRK2s as per the materials and methods. Infiltration of nYFP-HUB2 alone served as a negative control along with PRP4Kc-cYFP (Figure 7). Fluorescence signals were visualized with confocal microscopy 3d after injection. n > 10 images were acquired for each set. Scale bars = 20 μm.

Supplemental Figure 8. Immunoblotting of HA-HUB2 proteins levels when expressed in the *raf24-2* background. 10d-old plants were harvested and analyzed by immunoblot. Ponceau staining serves as a loading control. 50 μg of protein samples were loaded in each lane. Equal amounts of HUB2 over-expression is observed across plants expressing plants expressing either HA-HUB2^{S314}, HA-HUB2^{S314A}, or HA-HUB2^{S314D}.

SUPPLEMENTAL DATA
Supplemental Data 1: Phosphoproteome data
Supplemental Data 2: Primer sets used in the study
Supplemental Data 3: HUB2-TAP pulldown data
Supplemental Data 4:SnRK2 subclasses

REFERENCES
Bao S, Hua C, Shen L, Yu H. **2020**. New insights into gibberellin signaling in regulating flowering in Arabidopsis. *Journal of Integrative Plant Biology* **62**: 118–131.

Bittner-Eddy PD, Crute IR, Holub EB, Beynon JL. **2000**. RPP13 is a simple locus in
Arabidopsis thaliana for alleles that specify downy mildew resistance to different avirulence
determinants in Peronospora parasitica. *The Plant Journal* **21**: 177–188.
Bourbousse C, Ahmed I, Roudier F, Zabulon G, Blondet E, Balzergue S, Colot V, Bowler
C, Barneche F. **2012**. Histone H2B monoubiquitination facilitates the rapid modulation of gene
expression during Arabidopsis photomorphogenesis. *PLoS Genetics* **8**: e1002825.
Bradford MM. **1976**. A rapid and sensitive method for the quantitation of microgram quantities
of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**: 248–254.
Brocard-Gifford I, Lynch TJ, Garcia ME, Malhotra B, Finkelstein RR. **2004**. The
Arabidopsis thaliana ABSCISIC ACID-INSENSITIVE8 encodes a novel protein mediating
abscisic acid and sugar responses essential for growth. *The Plant Cell* **16**: 406–421.
Cao Y, Dai Y, Cui S, Ma L. **2008**. Histone H2B monoubiquitination in the chromatin of
FLOWERING LOCUS C regulates flowering time in Arabidopsis. *The Plant Cell* **20**: 2586–
2602.
Cao Y, Wen L, Wang Z, Ma L. **2015**. SKIP Interacts with the Paf1 Complex to Regulate
Flowering via the Activation of FLC Transcription in Arabidopsis. *Molecular Plant* **8**: 1816–
1819.
Chen H, Feng H, Zhang X, Zhang C, Wang T, Dong J. **2019**. An Arabidopsis E3 ligase
HUB2 increases histone H2B monoubiquitination and enhances drought tolerance in transgenic
cotton. *Plant Biotechnology Journal* **17**: 556–568.
Chen Y, Song S, Gan Y, Jiang L, Yu H, Shen L. **2020**. SHAGGY-like kinase 12 regulates
flowering through mediating CONSTANS stability in Arabidopsis. *Science Advances* **6**:
eaaw0413.
Chomczynski P, Sacchi N. **1987**. Single-step method of RNA isolation by acid guanidinium
thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* **162**: 156–159.
Chong L, Xu R, Huang P, Guo P, Zhu M, Du H, Sun X, Ku L, Zhu J-K, Zhu Y. **2022**. The
tomato OST1-VOZ1 module regulates drought-mediated flowering. *The Plant Cell* **34**: 2001–
2018.
Clough SJ, Bent AF. **1998**. Floral dip: a simplified method for *Agrobacterium*-mediated
transformation of *Arabidopsis thaliana*. *The Plant Journal* **16**: 735–743.
Del Corpo D, Fullone MR, Miele R, Lafond M, Pontiggia D, Grisel S, Kieffer-Jaquinod S,
Giardina T, Bellincampi D, Lionetti V. **2020**. AtPME17 is a functional Arabidopsis thaliana
pectin methylesterase regulated by its PRO region that triggers PME activity in the resistance to
Botrytis cinerea. *Molecular Plant Pathology* **21**: 1620–1633.
Cox J, Mann M. **2008**. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature Biotechnology*
26: 1367–1372.
Creux N, Harmer S. **2019**. Circadian rhythms in plants. *Cold Spring Harbor Perspectives in*
Biology **11**.

Deng W, Ying H, Helliwell CA, Taylor JM, Peacock WJ, Dennis ES. **2011**. FLOWERING LOCUS C (FLC) regulates development pathways throughout the life cycle of Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 6680–6685.

Ding Y, Li H, Zhang X, Xie Q, Gong Z, Yang S. **2015a**. OST1 kinase modulates freezing tolerance by enhancing ICE1 stability in Arabidopsis. *Developmental Cell* **32**: 278–289.

Ding S, Zhang B, Qin F. **2015b**. Arabidopsis RZFP34/CHYR1, a Ubiquitin E3 Ligase, Regulates Stomatal Movement and Drought Tolerance via SnRK2.6-Mediated Phosphorylation. *The Plant Cell* **27**: 3228–3244.

Eom H, Park SJ, Kim MK, Kim H, Kang H, Lee I. **2018**. TAF15b, involved in the autonomous pathway for flowering, represses transcription of FLOWERING LOCUS C. *The Plant Journal* **93**: 79–91.

Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D, Yochikawa A, Cano-Ramirez DL, Chembath A, *et al.* 2018. Circadian Entrainment in Arabidopsis by the Sugar-Responsive Transcription Factor bZIP63. *Current Biology* **28**: 2597–2606.e6.

Gaudinier A, Blackman BK. **2020**. Evolutionary processes from the perspective of flowering time diversity. *The New Phytologist* **225**: 1883–1898.

Gehan MA, Fahlgren N, Abbasi A, Berry JC, Callen ST, Chavez L, Doust AN, Feldman MJ, Gilbert KB, Hodge JG, *et al.* 2017. PlantCV v2: Image analysis software for high-throughput plant phenotyping. *PeerJ* **5**: e4088.

Gu X, Jiang D, Wang Y, Bachmair A, He Y. **2009**. Repression of the floral transition via histone H2B monoubiquitination. *The Plant Journal* **57**: 522–533.

He Y. **2009**. Control of the transition to flowering by chromatin modifications. *Molecular Plant* **2**: 554–564.

He Y, Doyle MR, Amasino RM. **2004**. PAF1-complex-mediated histone methylation of FLOWERING LOCUS C chromatin is required for the vernalization-responsive, winter-annual habit in Arabidopsis. *Genes & Development* **18**: 2774–2784.

Heintzen C, Nater M, Apel K, Staiger D. **1997**. AtGRP7, a nuclear RNA-binding protein as a component of a circadian-regulated negative feedback loop in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences of the United States of America* **94**: 8515–8520.

Himanen K, Woloszynska M, Boccardi TM, De Groeve S, Nelissen H, Bruno L, Vuylsteke M, Van Lijsebettens M. **2012**. Histone H2B monoubiquitination is required to reach maximal transcript levels of circadian clock genes in Arabidopsis. *The Plant Journal* **72**: 249–260.

Hu M, Pei B-L, Zhang L-F, Li Y-Z. **2014**. Histone H2B monoubiquitination is involved in regulating the dynamics of microtubules during the defense response to Verticillium dahliae toxins in Arabidopsis. *Plant Physiology* **164**: 1857–1865.

Humphrey SJ, James DE, Mann M. **2015**. Protein phosphorylation: A major switch mechanism for metabolic regulation. *Trends in Endocrinology and Metabolism* **26**: 676–687.

Ito S, Song YH, Josephson-Day AR, Miller RJ, Breton G, Olmstead RG, Imaizumi T. **2012**.
FLOWERING BHLH transcriptional activators control expression of the photoperiodic
flowering regulator CONSTANS in Arabidopsis. *Proceedings of the National Academy of*
Sciences of the United States of America **109**: 3582–3587.

Jeon M, Jeong G, Yang Y, Luo X, Jeong D, Kyung J, Hyun Y, He Y, Lee I. **2023**.
Vernalization-triggered expression of the antisense transcript COOLAIR is mediated by CBF
genes. *eLife* **12**.

Jeon YG, Lee JH, Ji Y, Sohn JH, Lee D, Kim DW, Yoon SG, Shin KC, Park J, Seong JK, *et*
***al*. 2020**. RNF20 functions as a transcriptional coactivator for ppary by promoting ncor1
degradation in adipocytes. *Diabetes* **69**: 20–34.

Jin S, Ahn JH. **2021**. Regulation of flowering time by ambient temperature: repressing the
repressors and activating the activators. *The New Phytologist* **230**: 938–942.

Kamiyama Y, Hirotani M, Ishikawa S, Minegishi F, Katagiri S, Rogan CJ, Takahashi F,
Nomoto M, Ishikawa K, Kodama Y, *et al*. 2021. Arabidopsis group C Raf-like protein kinases
negatively regulate abscisic acid signaling and are direct substrates of SnRK2. *Proceedings of*
the National Academy of Sciences of the United States of America **118**.

Kathania M, Kumar R, Lenou ET, Basrur V, Theiss AL, Chernoff J, Venuprasad K. **2022**.
Pak2-mediated phosphorylation promotes RORγt ubiquitination and inhibits colonic
inflammation. *Cell reports* **40**: 111345.

Kawa D, Meyer AJ, Dekker HL, Abd-El-Haliem AM, Gevaert K, Van De Slijke E,
Maszkowska J, Bucholc M, Dobrowolska G, De Jaeger G, *et al*. 2020. SnRK2 Protein
Kinases and mRNA Decapping Machinery Control Root Development and Response to Salt.
Plant Physiology **182**: 361–377.

Kim H-J, Hyun Y, Park J-Y, Park M-J, Park M-K, Kim MD, Kim H-J, Lee MH, Moon J,
Lee I, *et al*. 2004. A genetic link between cold responses and flowering time through FVE in
Arabidopsis thaliana. *Nature Genetics* **36**: 167–171.

Kim JH, Kim WT. **2013**. The Arabidopsis RING E3 ubiquitin ligase AtAIRP3/LOG2
participates in positive regulation of high-salt and drought stress responses. *Plant Physiology*
162: 1733–1749.

Kuhn A, Roosjen M, Mutte S, Dubey SM, Carrasco VPC, Boeren S, Monzer A, Koehorst J,
Kohchi T, Nishihama R, Fendrych M, Sprakel J, Friml J, Weijers D. 2024. RAF-like protein
kinases mediate a deeply conserved, rapid auxin response. *Cell* 187: 130-148.e17.

Lee KC, Chung KS, Lee HT, Park J-H, Lee JH, Kim J-K. **2020**. Role of Arabidopsis Splicing
factor SF1 in Temperature-Responsive Alternative Splicing of FLM pre-mRNA. *Frontiers in*
Plant Science **11**: 596354.

Lee J, Lee I. **2010**. Regulation and function of SOC1, a flowering pathway integrator. *Journal of*
Experimental Botany **61**: 2247–2254.

Lee JH, Ryu H-S, Chung KS, Posé D, Kim S, Schmid M, Ahn JH. **2013**. Regulation of
temperature-responsive flowering by MADS-box transcription factor repressors. *Science* **342**:
628–632.

Li X, Chen L, Yao L, Zou J, Hao J, Wu W. **2022**. Calcium-dependent protein kinase CPK32 mediates calcium signaling in regulating Arabidopsis flowering time. *National science review* **9**: nwab180.

Li P, Tao Z, Dean C. **2015**. Phenotypic evolution through variation in splicing of the noncoding RNA COOLAIR. *Genes & Development* **29**: 696–701.

Lin Z, Li Y, Wang Y, Liu X, Ma L, Zhang Z, Mu C, Zhang Y, Peng L, Xie S, *et al.* 2021. Initiation and amplification of SnRK2 activation in abscisic acid signaling. *Nature Communications* **12**: 2456.

Lin Z, Li Y, Zhang Z, Liu X, Hsu C-C, Du Y, Sang T, Zhu C, Wang Y, Satheesh V, *et al.* 2020. A RAF-SnRK2 kinase cascade mediates early osmotic stress signaling in higher plants. *Nature Communications* **11**: 613.

Linden KJ, Callis J. **2020**. The ubiquitin system affects agronomic plant traits. *The Journal of Biological Chemistry* **295**: 13940–13955.

Liu Y, Koornneef M, Soppe WJJ. **2007**. The absence of histone H2B monoubiquitination in the Arabidopsis hub1 (rdo4) mutant reveals a role for chromatin remodeling in seed dormancy. *The Plant Cell* **19**: 433–444.

Liu Y, Wang K, Cheng Q, Kong D, Zhang X, Wang Z, Wang Q, Xie Q, Yan J, Chu J, *et al.* 2020. Cysteine protease RD21A regulated by E3 ligase SINAT4 is required for drought-induced resistance to Pseudomonas syringae in Arabidopsis. *Journal of Experimental Botany* **71**: 5562–5576.

Lozano-Juste J, Alrefaei AF, Rodriguez PL. **2020**. Plant osmotic stress signaling: mapkkks meet snrk2s. *Trends in Plant Science* **25**: 1179–1182.

Lucas JR, Courtney S, Hassfurder M, Dhingra S, Bryant A, Shaw SL. **2011**. Microtubule-associated proteins MAP65-1 and MAP65-2 positively regulate axial cell growth in etiolated Arabidopsis hypocotyls. *The Plant Cell* **23**: 1889–1903.

Ma S, Tang N, Li X, Xie Y, Xiang D, Fu J, Shen J, Yang J, Tu H, Li X, *et al.* 2019. Reversible Histone H2B Monoubiquitination Fine-Tunes Abscisic Acid Signaling and Drought Response in Rice. *Molecular Plant* **12**: 263–277.

Mao T, Jin L, Li H, Liu B, Yuan M. **2005**. Two microtubule-associated proteins of the Arabidopsis MAP65 family function differently on microtubules. *Plant Physiology* **138**: 654–662.

MAPK Group. **2002**. Mitogen-activated protein kinase cascades in plants: a new nomenclature. *Trends in Plant Science* **7**: 301–308.

Mehta D, Scandola S, Uhrig RG. 2022. BoxCar and Library-Free Data-Independent Acquisition Substantially Improve the Depth, Range, and Completeness of Label-Free Quantitative Proteomics. *Anal Chem* 94:793-802.

Monaghan J, Xu F, Gao M, Zhao Q, Palma K, Long C, Chen S, Zhang Y, Li X. **2009**. Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant innate immunity. *PLoS Pathogens* **5**: e1000526.

Mouradov A, Cremer F, Coupland G. **2002**. Control of flowering time: interacting pathways
as a basis for diversity. *The Plant Cell* **14 Suppl**: S111–30.
Moyal L, Lerenthal Y, Gana-Weisz M, Mass G, So S, Wang S-Y, Eppink B, Chung YM,
Shalev G, Shema E, *et al.* 2011. Requirement of ATM-dependent monoubiquitylation of histone
H2B for timely repair of DNA double-strand breaks. *Molecular Cell* **41**: 529–542.
Mulekar JJ, Huq E. **2012**. Does CK2 affect flowering time by modulating the autonomous
pathway in Arabidopsis? *Plant Signaling & Behavior* **7**: 292–294.
Ogiso E, Takahashi Y, Sasaki T, Yano M, Izawa T. **2010**. The role of casein kinase II in
flowering time regulation has diversified during evolution. *Plant Physiology* **152**: 808–820.
Rose LE, Bittner-Eddy PD, Langley CH, Holub EB, Michelmore RW, Beynon JL. **2004**.
The maintenance of extreme amino acid diversity at the disease resistance gene, RPP13, in
Arabidopsis thaliana. *Genetics* **166**: 1517–1527.
Sanagi M, Aoyama S, Kubo A, Lu Y, Sato Y, Ito S, Abe M, Mitsuda N, Ohme-Takagi M,
Kiba T, *et al.* 2021. Low nitrogen conditions accelerate flowering by modulating the
phosphorylation state of FLOWERING BHLH 4 in Arabidopsis. *Proceedings of the National*
Academy of Sciences of the United States of America **118**.
Schmitz RJ, Tamada Y, Doyle MR, Zhang X, Amasino RM. **2009**. Histone H2B
deubiquitination is required for transcriptional activation of FLOWERING LOCUS C and for
proper control of flowering in Arabidopsis. *Plant Physiology* **149**: 1196–1204.
Schütze K, Harter K, Chaban C. **2009**. Bimolecular fluorescence complementation (BiFC) to
study protein-protein interactions in living plant cells. *Methods in Molecular Biology* **479**: 189–
202.
Soma F, Mogami J, Yoshida T, Abekura M, Takahashi F, Kidokoro S, Mizoi J, Shinozaki
K, Yamaguchi-Shinozaki K. **2017**. ABA-unresponsive SnRK2 protein kinases regulate mRNA
decay under osmotic stress in plants. *Nature Plants* **3**: 16204.
Soma F, Takahashi F, Suzuki T, Shinozaki K, Yamaguchi-Shinozaki K. **2020**. Plant Raf-like
kinases regulate the mRNA population upstream of ABA-unresponsive SnRK2 kinases under
drought stress. *Nature Communications* **11**: 1373.
Staiger D, Zecca L, Wieczorek Kirk DA, Apel K, Eckstein L. **2003**. The circadian clock
regulated RNA-binding protein AtGRP7 autoregulates its expression by influencing alternative
splicing of its own pre-mRNA. *The Plant Journal* **33**: 361–371.
Stecker KE, Minkoff BB, Sussman MR. **2014**. Phosphoproteomic analyses reveal early
signaling events in the osmotic stress response. *Plant Physiology* **165**: 1171–1187.
Sun D, Li Y, Ma Z, Yan X, Li N, Shang B, Hu X, Cui K, Koiwa H, Zhang X. **2021**. The
epigenetic factor FVE orchestrates cytoplasmic SGS3-DRB4-DCL4 activities to promote
transgene silencing in Arabidopsis. *Science Advances* **7**.
Tyanova S, Temu T, Cox J. **2016**. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. *Nature Protocols* **11**: 2301–2319.
Ubersax JA, Ferrell JE. **2007**. Mechanisms of specificity in protein phosphorylation. *Nature*
Reviews. Molecular Cell Biology **8**: 530–541.

Uhrig RG, Schläpfer P, Roschitzki B, Hirsch-Hoffmann M, Gruissem W. **2019**. Diurnal
changes in concerted plant protein phosphorylation and acetylation in Arabidopsis organs and
seedlings. *The Plant Journal* **99**: 176–194.

Vierstra RD. **2009**. The ubiquitin-26S proteasome system at the nexus of plant biology. *Nature*
Reviews. Molecular Cell Biology **10**: 385–397.

Wang F, Gao Y, Liu Y, Zhang X, Gu X, Ma D, Zhao Z, Yuan Z, Xue H, Liu H. **2019**. BES1-
regulated BEE1 controls photoperiodic flowering downstream of blue light signaling pathway in
Arabidopsis. *The New Phytologist* **223**: 1407–1419.

Wang J, Grubb LE, Wang J, Liang X, Li L, Gao C, Ma M, Feng F, Li M, Li L, *et al.* **2018a**.
A regulatory module controlling homeostasis of a plant immune kinase. *Molecular Cell* **69**: 493–
504.e6.

Wang B, Liu G, Zhang J, Li Y, Yang H, Ren D. **2018b**. The RAF-like mitogen-activated
protein kinase kinase kinases RAF22 and RAF28 are required for the regulation of
embryogenesis in Arabidopsis. *The Plant Journal* **96**: 734–747.

Wang P, Xue L, Batelli G, Lee S, Hou Y-J, Van Oosten MJ, Zhang H, Tao WA, Zhu J-K.
2013. Quantitative phosphoproteomics identifies SnRK2 protein kinase substrates and reveals
the effectors of abscisic acid action. *Proceedings of the National Academy of Sciences of the*
United States of America **110**: 11205–11210.

Woloszynska M, Le Gall S, Neyt P, Boccardi TM, Grasser M, Längst G, Aesaert S,
Coussens G, Dhondt S, Van De Slijke E, *et al.* **2019**. Histone 2B monoubiquitination complex
integrates transcript elongation with RNA processing at circadian clock and flowering regulators.
Proceedings of the National Academy of Sciences of the United States of America **116**: 8060–
8069.

Wright DE, Wang C-Y, Kao C-F. **2011**. Flickin' the ubiquitin switch: the role of H2B
ubiquitylation in development. *Epigenetics* **6**: 1165–1175.

Zhang Y, Cheng YT, Bi D, Palma K, Li X. **2005**. MOS2, a protein containing G-patch and
KOW motifs, is essential for innate immunity in Arabidopsis thaliana. *Current Biology* **15**:
1936–1942.

Zhang L, Luo P, Bai J, Wu L, Di D-W, Liu H-Q, Li J-J, Liu Y-L, Khaskheli AJ, Zhao C-M,
et al. **2021**. Function of histone H2B monoubiquitination in transcriptional regulation of auxin
biosynthesis in Arabidopsis. *Communications Biology* **4**: 206.

Zhang H, Ransom C, Ludwig P, van Nocker S. **2003**. Genetic analysis of early flowering
mutants in Arabidopsis defines a class of pleiotropic developmental regulator required for
expression of the flowering-time switch flowering locus C. *Genetics* **164**: 347–358.

Zhang H, van Nocker S. **2002**. The VERNALIZATION INDEPENDENCE 4 gene encodes a
novel regulator of FLOWERING LOCUS C. *The Plant Journal* **31**: 663–673.

Zhang Q, Zhang W, Wei J, Gao Z, Guan J, Cui Z, Wang X. **2022**. SKIP Regulates ABA
Signaling through Alternative Splicing in Arabidopsis. *Plant & Cell Physiology* **63**: 494–507.

Zhou S, Chen Q, Sun Y, Li Y. **2017**. Histone H2B monoubiquitination regulates salt stress-
induced microtubule depolymerization in Arabidopsis. *Plant, Cell & Environment*.

Zhou H, Low TY, Hennrich ML, van der Toorn H, Schwend T, Zou H, Mohammed S,
Heck AJR. **2011**. Enhancing the identification of phosphopeptides from putative basophilic
kinase substrates using Ti (IV) based IMAC enrichment. *Molecular & Cellular Proteomics* **10**:
M110.006452.
Zou B, Yang D-L, Shi Z, Dong H, Hua J. **2014**. Monoubiquitination of histone 2B at the
disease resistance gene locus regulates its expression and impacts immune responses in
Arabidopsis. *Plant Physiology* **165**: 309–318.
1059
1060

A

RAF24: AT2G35050

B

WT *raf18-1* *raf20-1* *raf24-2*

C

Days to bolting

WT *raf18-1* *raf20-1* *raf24-2*

Rosette leaf number

WT *raf18-1* *raf20-1* *raf24-2*

D

WT *raf24-2* *raf24-4* *raf24-5*

E

Days to bolting

WT *raf24-2* *raf24-4* *raf24-5*

Rosette leaf number

WT *raf24-2* *raf24-4* *raf24-5*

A

HA-HUB2S314S   HA-HUB2S314A   HA-HUB2S314D

raf24-2

B

C

Table 1: Uniquely differentially phosphophorylated proteins in *raf24-2*

Leading Protein	Common name	Description	SUBA5	Phosphorylation site	Pathway	Reference
AT3G13300	VCS	VARICOSE	Cytosol	pSer664	Osmotic stress signaling	(Soma et al., 2017)
AT1G55250	HUB2	HISTONE MONO-UBIQUITINATION 2	Nucleus	pSer314	Flowering pathway	(Cao et al., 2008; Woloszynska et al., 2019; Gu et al., 2009)
AT5G61150	VIP4	VERNALIZATION INDEPENDENCE 4	Nucleus	pSer600 pSer605	Autonomous flowering pathway	(Zhang and van Nocker, 2002)
AT5G51300	SF1	SPLICING FACTOR 1	Nucleus	pSer77	Thermosensory flowering pathway	(Lee *et al.*, 2020)
AT1G51140	FBH3	FLOWERING BHLH 3	Nucleus	pSer259	Photoperiodic flowering pathway; Osmotic stress signaling	(Ito *et al.*, 2012)
AT1G08420	BSL2	BRI1 SUPPRESSOR 1 (BSU1)-LIKE 2	Nucleus	pSer973	BR signaling	(Kim et al., 2016)
AT4G03080	BSL1	BRI1 SUPPRESSOR 1 (BSU1)-LIKE 1	Nucleus	pSer491	BR signaling	(Kim et al., 2016; Kim et al., 2018)
AT1G19350	BES1	BRI1-EMS-SUPPRESSOR 1	Nucleus	pTyr175	Photoperiodic flowering pathway; BR signaling	(Wang *et al.*, 2019)
AT1G78700	BEH4	BES1/BZR1 HOMOLOG 4	Nucleus	pSer310 pSer312	BR signaling	(Galstyan and Nemhauser, 2019)

subcellular localization
pSer = phosphorylated serine
pTyr = phosphorylated tyrosine