On the potential for GWAS with phenotypic population means and allele-frequency data (popGWAS)

Markus Pfenninger [1,2,3]*

[1] Dept. Molecular Ecology, Senckenberg Biodiversity and Climate Research Centre, Georg-Voigt-Str. 14-16, D-60325 - Frankfurt am Main, Germany
[2] LOEWE Centre for Translational Biodiversity Genomics, Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, D-60325 – Frankfurt am Main, Germany
[3] Institute for Molecular and Organismic Evolution, Johannes Gutenberg University, Johann-Joachim-Becher-Weg 7, D-55128 – Mainz, Germany

*Corresponding author
Correspondence: Markus.Pfenninger@senckenberg.de

ABSTRACT

It is vital to understand the genomic basis of differences in ecologically important traits if we are to understand the impact of global change on biodiversity and enhance our ability for targeted intervention. This study explores the potential of a novel genome-wide association study (GWAS) approach for identifying loci underlying quantitative polygenic traits in natural populations, based on phenotypic population means and genome-wide allele frequency data as obtained e.g. by PoolSeq approaches. Extensive population genetic forward simulations demonstrate that the approach is generally effective for oligogenic and moderately polygenic traits and relatively insensitive to low heritability. However, applicability is limited for highly polygenic architectures and pronounced population structure. The required sample size is moderate with very good results being obtained already for a few dozen populations scored. When combined with machine learning for feature selection, the method performs very well in predicting population means. The data efficiency of the method, particularly when using pooled sequencing and bulk phenotyping, makes GWAS studies more accessible for research in biodiversity genomics. Moreover, in a direct comparison to individual based GWAS, the proposed method performed consistently better with regard to the number of true positive loci identified and prediction accuracy. Overall, this study highlights the promise of popGWAS for dissecting the genetic basis of complex traits in natural populations.

Keywords: biodiversity population genomics, molecular trait basis

Introduction

A major goal as well as a major challenge in evolutionary biology is to understand how genes influence traits, i.e. the genotype-phenotype link, (Brandes et al., 2022; Uffelmann et al., 2021). The difficulties in achieving this goal are primarily due to the fact that the heritable variation of many, if not most, relevant phenotypes is determined by small contributions from many genetic loci (Sella & Barton, 2019). Such complex traits are usually influenced by a few dozen genes that are mechanistically directly involved in their expression, but often also by numerous, if not almost all, other genes as well as the environment in the widest sense (Boyle et al., 2017).

Genome-wide association studies (GWAS) are commonly used to link complex phenotypic traits to their genomic basis (Brandes et al., 2022; Visscher et al., 2012). However, given the complexity of causal mechanisms and the small effects of individual loci, often only a small fraction of the genetic variation underlying phenotypic variance is identified, despite the considerable logistic effort in terms of the number of phenotyped and genotyped individuals (Brandes et al., 2022; Visscher et al., 2017). As a result, accurate predictions of phenotypes from genomic data are still quite limited and there is currently no other strategy than to keep increasing sample sizes (Brandes et al., 2022). This is a problem in the medical sciences (Shendure et al., 2019), but the greater challenge for science and society probably lies in addressing the global biodiversity crisis. It would be highly desirable to have affordable methods to accurately understand the genomic basis of relevant traits and predict (non-model) species responses to all aspects of global change (Bernatchez et al., 2023; Waldvogel et al., 2020).

GWAS with wild populations has been advocated for some time (Santure & Garant, 2018). However, despite recent progress in high throughput, automated phenotyping (Dunker et al., 2022; Tills et al., 2023; Xie & Yang, 2020), the advances of biodiversity genomics in obtaining high quality reference genomes for almost every species (Exposito-Alonso et al., 2020; Formenti et al., 2022) and the possibility to gain cost-effective genome-wide population data (Czech, Peng, Spence, Lang, Bellagio, Hildebrandt, Fritschi, Schwab, Rowan, & Weigel, 2022; Schlötterer et al., 2014), relatively few empirical studies are currently available [citations]. This gap between the possibilities and actual practical application in biodiversity conservation (Heuertz et al., 2023; Hogg, 2023) is probably as much due to the still existing logistic and financial challenges as to a lack of data- and resource-efficient methods.

Here, I explore the potential of a new GWAS approach using phenotypic population means and genome-wide allele-frequency data. The rationale behind the approach is straightforward. If a quantitative polygenic trait has an additive genetic component, an individual's phenotypic trait value should at least roughly correlate with the number of trait-increasing alleles at the underlying loci (Uffelmann et al., 2021). Consequently, it was theoretically expected (Orr, 1998; Pritchard & Di Rienzo, 2010) and empirically shown (Turchin et al., 2012) that trait-increasing alleles will tend to have greater frequencies in the population with higher mean trait values, compared to the population with a lower trait mean. When examining populations with a range of different phenotypic trait means, we may therefore expect that the allele frequencies at the trait-affecting loci show a linear or at least steady relation with the observed trait means (Barton, 1999). I hypothesise here that this predicted relation can be exploited to distinguish potentially causal loci (and the linked variation) from loci not associated with the focal trait. In case of a successful evaluation, the major advantages of the proposed approach would be the reduced sequencing effort by the possibility to use pooled population samples (PoolSeq) and

the opportunity to use bulk phenotyping (e.g. by satellite imaging, flow-cytometry, etc.) on
traits for which individual phenotyping is difficult or tedious.

The most important assumption for the approach is obviously that observed population
differences in the focal trait means have at least partially a genetic basis. Since the environment
has usually an effect on the phenotype (Sella & Barton, 2019), total phenotypic variance should
be adjusted for known fixed environmental effects, because this increases the fraction of
variance due to genetic factors (Visscher et al. 2008). Predicting additive genetic values with
even higher accuracy can be achieved by taking into account GxE interactions through repeated
phenotypic measurements of the same individuals under different environmental conditions,
e.g. by time series (Visscher et al. 2008). I assumed therefore that environmental influence on
the phenotypic trait variance among populations has been statistically removed as much as
possible (Harpak & Przeworski, 2021). Similarly important is the assumption that the genetic
variance of the focal quantitative trait can be adequately described by an additive model. Both
empirical and theoretical evidence suggests that this is indeed the case for most complex traits
(Hill et al., 2008). Even though epistatic interactions are wide spread (Mackay, 2014), Sella and
Barton (Sella & Barton, 2019) argue that the marginal allelic effects on quantitative traits are
well approximated by a simple additive model.

The aims of this study were i) to understand whether and under which circumstances
the hypothesised pattern of a linear relation between the population allele frequencies at
causal loci and the phenotypic population means of the respective trait emerges, ii) to
evaluate the influence of population genetic parameters of typical natural systems and the
experimental design on the likelihood of identifying causal loci underlying an additive
quantitative trait, in particular to elucidate the limits of the approach with regard to genetic
architecture and population structure, iii) to compare proposed method with the
performance of an individual-based GWAS, iv) to explore the possibilities for statistical
genomic prediction of phenotypic population means from the allele frequencies at
identified loci, and v) to evaluate the statistical power of the method for a realistic range of
effective genome sizes. I used individual-based population genomic forward simulations
and machine learning approaches (minimum entropy feature selection) for prediction and
utilised an information theory-based framework for evaluation of the proposed method.

Material and methods

Rationale of the method

Expectation of a positive correlation between quantitative trait loci allele frequencies and
phenotypic population means.

Consider a biallelic, codominant system for the additively heritable component of a
quantitative trait with n loci contributing to the trait. In this system, all loci contribute
equally to the phenotypic trait, with one allele per locus making a greater contribution than
the other. The phenotypic trait value x of an individual can then be determined by simply
adding up the number of trait-increasing alleles (g with values of 0, 1 or 2) over all n
quantitative trait loci (QTL) and multiplying this sum with a scaling constant k:

(1) $x = k \times (g_1 + g_2 + \cdots + g_n)$

When adding more individuals, the phenotypic population trait mean is defined as the
mean of the row sums:

(2)

$$
\begin{array}{cccccc}
 & QTL_1 & \cdots & \cdots & QTL_n & \text{individual trait value}\\
Ind_1 & g_{11} & g_{12} & \cdots & g_n & x_1 = k \times \sum_{l=1}^{n} g_{1l}\\
Ind_2 & g_{21} & \cdots & \cdots & \cdots & \cdots\\
\cdots & \cdots & \cdots & \cdots & \cdots & \cdots\\
Ind_m & g_{m1} & \cdots & \cdots & g_{mn} & x_m = k \times \sum_{l=1}^{n} g_{1l}
\end{array}
$$

$$\text{population trait mean}$$
$$\bar{x} = \frac{\sum_{i=1}^{m} x_i}{m}$$

128 The columns of this matrix can be used to calculate the population allele frequency (AF)
129 of the trait increasing allele for each QTL.

130 (3)

$$
\begin{array}{cccccccc}
 & QTL_1 & \cdots & \cdots & QTL_n & & & \text{individual trait value}\\
Ind_1 & g_{11} & g_{12} & \cdots & g_n & & & x_1 = k \times \sum_{l=1}^{n} g_{1l}\\
Ind_2 & g_{21} & \cdots & \cdots & \cdots & & & \cdots\\
\cdots & \cdots & \cdots & \cdots & \cdots & & & \cdots\\
Ind_m & g_{m1} & \cdots & \cdots & g_{mn} & & & x_m = k \times \sum_{l=1}^{n} g_{1l}
\end{array}
$$

$$AF \quad AF_1 = \frac{\sum_{i=1}^{m} g_i}{2m} \quad \cdots \quad \cdots \quad AF_1 = \frac{\sum_{i=1}^{m} g_i}{2m} \qquad \overline{AF} = \frac{\sum_{i=1}^{n} AF_i}{n} \quad \sim \quad \bar{x} = \frac{\sum_{i=1}^{m} x_i}{m}$$

with labels *mean QTL AF* and *population trait mean*.

The mean population allele frequency at the QTL loci is thus directly proportional to the
phenotypic population trait mean. This relationship remains unchanged even if the
individual locus contributions are not identical, with some loci contributing more or less to
the phenotypic trait value. In this case, a scaling vector is required to weigh the individual
locus contributions to individual trait values, and those of the AFs to the population trait
mean. Since the AFs are by definition bounded by zero and one, the population trait mean is
minimal when the allele frequencies of the trait-increasing allele at all QTL are zero and
maximal when all QTL AFs are one. This proportionality links the individual genotypes and
the AFs at the QTL linearly with the population trait mean.
If we extend this to a set of populations and order them with decreasing phenotypic
population means, we can be sure that the mean QTL AFs of the populations will also be
ordered in decreasing sequence:

(4)

$$
\begin{array}{ccccc}
 & Pop_1 & Pop_2 & \cdots & Pop_n\\
\text{population trait mean} & \bar{x_1} > & \bar{x_2} > & \cdots & \bar{x_n}\\
 & \sim & \sim & \cdots & \sim\\
\textit{mean QTL AF} & \overline{AF_1} > & \overline{AF_2} > & \cdots & \overline{AF_n}\\
QTL_1 & AF_{11} & AF_{21} & \cdots & AF_{11}\\
QTL_2 & AF_{12} & AF_{22} & \cdots & \cdots\\
\cdots & \cdots & \cdots & \cdots & \cdots\\
QTL_m & AF_{1m} & \cdots & \cdots & AF_{nm}
\end{array}
$$

The answer to whether the allele frequencies in every row i.e. at every contributing
locus can be used to predict the population trait mean depends on whether the expected
covariance between these two vectors is positive:

(5) $\quad E[cov(QTL_m, \text{population trait mean})] > 0$

where

(6) $\text{cov}[\text{QTL}_m, \text{population trait mean}] = E[\sum_{i=1}^{n}(AF_{im} - E[AF_m]) * (\bar{x}_1 - E[\bar{X}])]$

with $\bar{X}$ representing the grand mean over all populations. As all elements in the QTL
matrix are positive, they inherently tend to contribute positively to their column means.
Therefore, AF larger than the overall AF mean at his locus tend to be on the left side of the
population closest to the overall phenotypic mean in the ordered matrix above. Conversely,
AF smaller than the locus AF mean are rather on the right. This leads intuitively to a
positive expected covariance between each row and the column mean, in particular if the
number of populations becomes large. Conversely, the AF at (unlinked) loci not
contributing to the phenotypic population trait mean have an expectation of zero.

Testing the expectations of a detectable linear relation between allele frequencies and
phenotypic population means
I tested these general expectations and the effect of different scaling vectors for the
effect size distribution of QTL with a first set of simulations. I generated a matrix of size *n x*
m populated with random AF between zero and 1. To avoid stochastic effects due to sample
size, the number of populations *n* was fixed at 10,000. The number of QTL *m* was varied
from oligogenic to highly polygenic (10, 20, 50, 100, 200, 500, 1000, 2000, 5000). Three
different distributions of loci effects were tested, i) a flat distribution with all loci
contributing equally, ii) a mildly decreasing exponential function and iii) a steeply
decreasing exponential function with few loci contributing much and many very little
(Supplemental Figure 1).
Each of the *m* columns was used to calculate the phenotypic population mean of the
respective population by adding up the AF multiplied with the respective locus weight. The
resulting *n* phenotypic population means were then correlated to the *n* AF of each of the *m*
loci and the resulting *m* Pearson correlation coefficients (i.e. the standardized covariance)
recorded. From these, mean and standard deviation were calculated and tested, whether
they conform to a normal distribution (scipy.stats.normaltest). Furthermore, a second
matrix of identical size was populated with random AF, and the correlation of these non-
contributing loci to the population means derived from the QTL matrix was computed. The
simulations were repeated 10 times in every possible parameter combination and the
results averaged (Supplemental Script 1).

Individual based Wright-Fisher forward model used for simulation

A Wright-Fisher individual forward genetic simulation model was used to investigate the
potential of a genome-wide association study based on the means of a population trait and
population allele frequency data. As this model formed the basis for all further simulations,
its general structure is first described.
In the simulation, all loci were assumed to be unlinked, thus representing haplotypes in
LD rather than single SNPs. (Visscher et al., 2017). For each simulation run, the initial allele
frequencies for all loci in the total population were randomly drawn from a beta function
with parameters $\alpha = \beta = 0.5$ in a range 0.1 to 0.9. To generate a hermaphroditic and diploid
individual, two alleles were randomly drawn with a probability based on their frequency at
the respective locus, and the resulting genotype at this locus was recorded. This process
was repeated for all loci. As a result, each individual was represented by a vector of biallelic
genotypes (*AA* = 0, *Aa*;*aA* = 1, *aa* = 2). To model a quantitative, fully additive trait, a
variable number of loci were assigned as quantitative trait loci (QTL). In addition, a much
larger number of neutral loci was modelled.

Genetic architecture of the quantitative trait
The continuous trait value was measured in arbitrary units. The allele (*A*) at each QTL
had no effect on the individual's trait value, resulting in a completely homozygous *A*
individual at the QTL having a phenotypic trait value of 0. The alternative allele (*a*) added a
locus-specific value to the trait. Two distributional extremes have been considered for the
allelic effects on the trait value: i) a uniform distribution where each locus contributes 0.5
units to the trait. An individual that is completely homozygous for the alternative allele *a*
therefore had a trait value equal to the number of QTL, ii) an exponential distribution with
few loci having large effects and many having very small effects, scaled such that the
maximum possible trait value was also equal to the number of QTL (see Supplementary
Figure 2A). To model the effect of phenotyping errors, unaccounted environmental
influence, and/or the unspecific contribution of the genomic background to the trait, a
random value drawn from a Gaussian distribution with a mean of zero and selectable
standard deviation between 0.1 and 3 was added to the genetically determined phenotype
value of each individual. The phenotypic value of each individual's trait was determined by
summing the allelic effects of all genotypes at all QTL loci plus the random value and the
result recorded.
It was assumed that all trait variation was based on standing genetic variation.
Mutations were not considered because new or low frequency mutations do not
substantially influence phenotypic population means and their AFs. In addition, mutations
are rare and mutations affecting a particular functional trait even rarer. Therefore,
restricting the simulations to standing genetic variation seemed reasonable.

Reproduction and selection
Subpopulations in each run were created from the same initially drawn random allele
frequency array, mimicking a common descent. Due to sampling variance, the realised allele
frequencies and thus the mean subpopulation trait value differed from the initial
frequencies of the total (ancestral) population. A subpopulation always comprised 500
adult individuals.
For reproduction, two random individuals were chosen with replacement from the adult
population. The genotype of an offspring individual at a locus was determined by randomly
choosing one of the two alleles from each designated parent at this locus. Each parent
fostered n_juv offspring; therefore, 2 x n_juv were produced in each mating. After
reproduction, the parental generation was discarded to prevent overlapping generations.
Each generation had N/2 matings, resulting in an offspring population of N*n_juv
individuals. This assured that the genotypes in each subpopulation were in Hardy-
Weinberg equilibrium after the first generation.
Because the offspring population was much larger than the the size of the adult
population, it was necessary to reduce it. This was achieved by a combination of 'hard'
natural selection and random mortality. An individual's survival to the adult stage was
determined by the absolute deviation of its phenotypic trait value from a pre-specified
selective trait optimum for the respective subpopulation. This selective trait optimum for a
subpopulation was determined by adding a random value taken from a Gaussian
distribution with a mean of zero and a standard deviation of 2.5 to the initial population
mean. An individual's survival probability was determined by an exponential decline
function with strength *s* (the exponent of the function, see Supplementary Figure 2B).
Individuals were randomly selected one by one from the offspring population, the distance
of their phenotype to the selective optimum calculated and their survival probability

calculated. A respectively biased coin was then tossed to determine their fate. This process
was repeated until the adult population size was reached, and any remaining offspring
individuals were indiscriminately discarded. If the phenotypic mean of the subpopulation
was close to or at the selective optimum (see below), this process resulted in stabilising
selection. If the population was away from the optimum, rapid directed selection towards
the optimum was observed, depending on the strength of selection.

Population structure

For the assessment of the effect of population structure, each subpopulation received in
every generation a certain number of migrants randomly chosen from all other
subpopulations. Both drift and selection towards different trait optima led to variation in
population trait means among the subpopulations.
Based on the model described above, I considered scenarios were subpopulations with
quantitative phenotypic population differences in mean for the trait in question were
screened from a larger total population. Although the population trait mean differences in
the simulation of this scenario were created by drift and local adaptation, any other source
of heritable phenotypic population differentiation, such as maladaptation, introgression, or
e.g. in the case of managed species, human choice, may also be the non-exclusive reasons for
differentiation in population means. The range of phenotypic variation among the
subpopulations was not predetermined, but an emergend feature of the simulation
parameters.
After evolving the subpopulations for the desired number of generations, phenotypic
trait means, and genome-wide allele frequencies were recorded. While the phenotypic
means for each subpopulation was calculated over all individuals, the allele frequencies
were estimated in a PoolSeq (Kofler et al., 2011) like fashion from subsamples of 50
individuals. The range of phenotypic trait means of the population sample was recorded.
Trait heritability was determined in the last generation by regressing the phenotypic values
of the offspring against the mean of their respective parents (Lynch & Walsh, 1998). As
measure for population subdivision due to drift, F_{ST} among all subpopulations was
calculated from the variance of the true allele frequencies (Wright, 1949).

Simulated scenarios

Influence of natural system and experimental design factors

In a first set of simulations, I explored the influence of factors inherent to the natural
system and the experimental design on population GWAS performance (Table 1). As factors
of the natural system, I assumed characteristics that are beyond control of the researcher,
such as heritability of the trait and its genetic architecture (number of QTL, distribution of
allele trait contribution). While the degree of population differentiation and range of
phenotypic differentiation are also inherent to the organism studied, the choice of samples
may allow a certain control over these parameters. The number of subpopulations screened
is clearly a study design decision (Table 1).

Table 1. Simulation parameters, their abbreviations, values used in simulations, their
biological meaning and whether the parameter is a feature of the natural system
under scrutiny or under the control of the researcher.

Parameter	Abbreviation	Values in the simulation	Biological meaning	Degree of knowledge in natural systems/under the control of study design

Number of subpopulations scored for phenotypic population means and genome wide allele frequencies	n_pop	12, 24 36, 48, 60	-	Full control
Number of quantitative trait loci contributing to the focal trait	n_qtl	30, 50, 70, 110, 500	Genetic architecture of the trait	*A priori* unknown
Distribution of allelic effects on the focal trait	allelic_contr	Flat, exponential	Genetic architecture of the trait	*A priori* unknown
Standard deviation of random phenotypic variation added to individuals	env-var	0.1, 1, 2, 3	Heritability of the trait	*A priori* unknown
Number of migrants received in each subpopulation per generation	mig	0, 5, 10, 50, 100	Population structure	Partial control

A genetic trait architecture of 30, 50, 70, 110 and 500 loci, flat and exponential allelic
effect distributions, as well as phenotypic plasticity coefficients of 0.1, 1, 2 and 3 were
applied (Table 1). Selection strength was fixed at 0.5 (Supplemental Figure 2B). The
number of immigrants reaching and reproducing in each subpopulation in each generation
was varied between 5, 10, 50 and 100 individuals. Simulations were run for 30 generations
among 12, 24, 36, 48 and 60 subpopulations of 500 individuals each (Table 1). For this set
of simulations, 1000 neutral loci and a fixed outlier threshold (upper 5% quantile, either 21
or 22 outlier loci, respectively) were applied. Each possible parameter combination was run
in five replicates, resulting in 4000 simulation runs.

The effect of each parameter on PPV was assessed with ANOVA over all simulations,
grouped after the respective parameter classes. The relative influence of the number of
populations, QTL loci, distribution of allelic contributions, trait heritability, phenotypic
range and population subdivision on the proportion of TPL among the outlier loci was
determined with a General Linearized Model (GLM).

Hierarchical population structure
The effect of a hierarchical population structure on popGWAS performance was assessed
with a restricted parameter set (Table 2). Two, three and four regions were simulated, each
consisting of 12 subpopulations. While there was gene-flow among the populations within
the regions as described above, no gene-flow among regions was simulated. The initial
degree of differentiation among regions was determined by first drawing one set of random
(beta distributed) allele frequencies as described above. To start each independent region
with slightly, but not completely different allele frequencies, a random variate from a
normal distribution with mean 0 and either 0.05, 0.1 or 0.2 as standard deviation was then
added to each of the initially drawn allele frequencies (Table 2). The simulations were then
run as described above. The hierarchical population structure was not explicitly taken into
account in statistical analysis but performed as described above.

Table 2. Simulation parameters for hierarchical population structure, their
abbreviations, values used in simulations, their biological meaning and whether the
parameter is a feature of the natural system under scrutiny or under the control of
the researcher.

Parameter	Abbreviation	Values in the simulation	Biological meaning	Degree of knowledge in natural systems/under the control of study design
Number of regions with 12 subpopulations each	n_reg	2,3,4	-	Partial control
Number of quantitative trait loci contributing to the focal	n_qtl	30, 70, 110	Genetic architecture of the trait	*A priori* unknown

trait				
Distribution of allelic effects on the focal trait	allelic_contr	Flat, exponential	Genetic architecture of the trait	*A priori* unknown
Standard deviation of random phenotypic variation added to individuals	env_var	2	Heritability of the trait	*A priori* unknown
Number of migrants received in each subpopulation per generation	mig	5, 50	Gene-flow	Partial control

Population based GWAS (popGWAS)

Assuming a linear relation between the phenotypic (sub)population means, and the
population allele frequencies of the causal loci on the other, I calculated an ordinary linear
regression between these two variables for all loci in the genome for each simulation. I used
the resulting $-\log_{10}$ p value as measure of regression fit and effect size. I recorded the
number of true positive loci (TPL) among the loci beyond a predefined outlier threshold
(2% quantile). As GWAS performance measures, the true positive rate (TPR = recall,
sensitivity, discovered proportion of all QTL), positive predictive value (PPV = precision,
proportion of TPL among outliers considered) and false discovery rate (1 – PPV = FDR =
proportion of false positive loci among outliers considered, type I error) were calculated.
This was performed for all simulated scenarios.

Individual GWAS (iGWAS)

In addition, a traditional linear GWAS on a random sample of individuals from all
subpopulations was performed for all simulation scenarios (termed iGWAS hereafter). The
number of individuals was chosen such that the PoolSeq sequencing effort for the same
simulation was matched, assuming that a 15X mean coverage is appropriate for accurate
genotyping. A linear regression of the individuals' genotypes at each locus as predictors for
the phenotypes of each individual in the sample was calculated. The resulting -log10p
values were treated in the same way as described for popGWAS above.

Genomic prediction and validation

The loci identified by popGWAS were used to devise a statistical genomic prediction
model to obtain a score that uses observed allele frequencies at the identified loci to predict
the mean population phenotype of unmeasured populations. To remove remaining
uninformative or redundant loci, I applied feature selection, which is particularly suitable
for bioinformatic data sets that contain many features but comparatively few data points.
The minimum entropy feature selection (MEFS) technique uses mutual information to
measure the dependence between each feature and the target variable. For a given number
of features (k), the data set of the allele frequencies at selected outlier loci and the
respective phenotypic population means was repeatedly randomly divided in training
(80%) and test set (20%), a multiple regression model fitted and the r^2-fit of the test sets to
the predicted phenotypes recorded. The best model for the current k was recorded and the
process repeated for all k in a range between 2 and the number of selected loci – 1. Finally,
the best model (i.e. highest r^2) among all k was chosen as best prediction model. MEFS was
implemented with the Python module scikit-learn 1.3.2 (Pedregosa et al., 2011)
The performance of the selected best prediction model for each run was tested with
independent data. Ten additional subpopulations were created and evolved under the same
parameters as part of the metapopulation of the initial set of populations and their mean
population phenotypes calculated as described above. Then the allele frequencies at the
predictive loci as identified by the best prediction model were extracted and phenotypic

prediction scores according to the best prediction model calculated. The performance of the
statistical genomic prediction was then evaluated by calculating the Pearson correlation
coefficient r between the observed mean population phenotypes and the phenotypic
prediction scores for the ten validation populations (Supplemental Script 2).
Genomic prediction and validation were performed in the same manner for iGWAS. Here
it was the goal to predict the individual phenotypes of a sample not used to train the model
with the genotypes of the loci identified by iGWAS as predictors.

Method performance with realistic genome sizes

Whether and which proportion of TPL, i.e. causal loci can be expected to be reliably
identified with the proposed method depends crucially on the total number of loci screened
as this number determines the length and size of the distributional tail of random
associations of neutral loci with the mean population phenotypes. The number of effectively
independently evolving loci in a population depends on genome size, effective population
size (including all factors that affect it locally and globally) and recombination rate
(Chakraborty, 1981; Taylor & Higgs, 2000). There are hardly any empirical estimates in the
literature, but dividing typical genome sizes by typical mean genome-wide LD ranges
suggested that a few tens of thousands to a few hundreds of thousands of independent loci
per genome is a realistic range for a large number of taxa (see Supplemental Table 1). I have
therefore considered 1,000, 5,000, 10,000, 30,000, 50,000 and 100,000 independent neutral
loci for samples of 12, 24, 36, 48 and 60 populations with a restricted set of parameters
(number of QTL and allelic contribution). As the true number of QTL underlying a trait is
rarely *a priori* known, I considered 10, 30, 50, 70 and 110 QTL loci in this analysis. I
therefore recorded the number of TPL found in sets of loci with the absolutely highest 10,
30, 50 and 100 -$\log_{10}p$ values, as well as outlier proportions of 0.0001, 0.001, 0.01, 0.02,
0.05 and 0.1 of the total number of loci in the respective simulation. As above, all
simulations were run in all possible parameter combinations with five replicates each
(Supplemental Script 3).
I analysed the performance of the method in an Area Under the Curve – Receiver
Operator Curve (AUC-ROC) and – Precision, Recall (AUC-PR) framework as suggested by
Lotterhos et al. (Lotterhos et al., 2022). For each combination of effective genome size and
number of population scored, mean TPR, PPV and FDR were calculated over all replicates
and parameter combinations for the respective set of simulations. The maximum F1 score
(Rijsbergen, 1979), the harmonic mean of the precision and recall representing both in one
metric, was used in addition to identify the optimal number of outliers to select.
All simulations were implemented in Python 3.11.7 (Van Rossum & Drake, 2009) and run
under pypy 3.10 (Team, 2019), the respective scripts can be found in the Supplementary
Material (Scripts 1-3). General statistical tests were performed with R (R Core Team, 2013).

<div align="center"># Results</div>

Allele frequencies at QTL loci co-vary positively with the population trait mean

The mean correlation coefficient between all QTL and the respective phenotypic
population means was positive in every parameter combination and in every single
simulation (Table 3).

Table 3. Expected mean Pearson's correlation coefficients between QTL AF and
phenotypic population means for three different locus contribution distributions and
varying number of QTL.

n_QTL	Flat	Mildly exponential	Strongly exponential
1	1	1	1
10	0.322	0.307	0.165
20	0.227	0.218	0.085
50	0.142	0.116	0.035
100	0.101	0.064	0.017
200	0.070	0.033	0.008
500	0.045	0.013	0.003
1000	0.031	0.006	0.001
2000	0.023	0.003	0.001
5000	0.014	0.001	0.000

The expected mean correlation coefficient decreased with increasing number of
contributing QTL (Figure 1). This decay was best described by a negative exponential
function of the form *number of QTL$^{-1/x}$* with x ranging from 1.26 in case of the strongly
unbalanced locus contributions to 2 for the flat distribution.

Figure 1. Plot of the expected correlation coefficient between individual QTL loci and
the population trait mean in dependence of the number of QTL loci. The dashed black
line gives the upper 95% confidence interval for non-contributing, neutral loci.

The distribution of the correlation coefficients did not deviate from a normal
distribution for the flat locus contribution distribution, while it did for all other parameters.
The correlation coefficients for the non-contributing loci had an expectation of zero and a
mean standard deviation of 0.01, regardless of the number of loci.

Influence of simulation parameters on parameters of the simulated populations

In an initial set of simulations, the effect of the number of generations simulated and the
number of migrants on the degree of adaptation and population structure was simulated. It
turned out that selection/drift equilibrium was reached after less than 10 generations. As
expected, more gene-flow led to less perfect local adaptation (Supplementary Figure 3A).
The time to reach the equilibrium F_{ST} among subpopulations strongly depended on the level
of gene-flow. While it was swiftly reached within 50 generations for 20 or more migrants
per generation, it would have taken much longer for less gene-flow (Supplementary Figure

3B). However, popGWAS performance was governed by the F_{ST} at the time of analysis and
not by selection-migration-drift equilibrium (Supplemental Figure 3C). To keep
computation times for the individual based model within reasonable bounds, all following
simulations were run for 30 generations.

In the second set of simulations with 1000 neutral loci and an outlier threshold of the
most extreme 2% -$\log_{10}p$ values, the permutation of all parameters with five replicates
yielded 3884 completed independent simulation runs. The 116 missing simulations to the
expected 4000 runs were due to one or more subpopulations going extinct during the
simulation.

The number of migrants strongly influenced the population structure ($r^2 = 0.990$). F_{ST}
estimates decreased exponentially(Supplemental Figure 4A). Heritability of the trait
depended strongly on the environmental variation parameter ($r^2 = 0.788$, Supplemental
Figure 4B). It decreased on average by 0.18 per unit standard deviation, with variations of
up to 0.05 even among runs with identical parameters. Trait heritability estimates ranged
from 0.21 to 1.02.

Factors influencing the proportion of detected true positive loci among outliers

Over all simulations in the first set of scenarios, on average about 12.1 (mean proportion
0.52) true positive loci (TPL) were among the highest 2% outliers. The TPL values ranged
between none (0) and 23 (1.0); the 25 percentile was 6 (0.28), the 75 percentile 19 (0.81).
This exceeded in >90% of cases random expectations, when excluding the highly polygenic
case (n_qtl = 500), this proportion rose to more than 93%. No p-value inflation was
observed; the mean slope of qq-plots was 1.27 (s.d. 0.23), the distribution ranged between
1 and 1.99 (Supplemental Figure 4A). Supplemental Figure 5 illustrates exemplarily the
results of three individual simulation runs with different parameter sets.

442   Figure 2. Effect of simulation parameters and emergent features on the proportion of
443   identified true positive loci. A) Phenotypical plasticity parameter as a proxy for
444   heritability. B) Number of QTL. C) Distribution of allelic contributions to phenotypic
445   trait. D) Number of reproducing immigrants per generation. E) Number of
446   populations scored for population phenotypic means and allele frequencies. F) Range
447   of population phenotypic means as an emergent feature.

The environmental variability parameter had no significant effect on PPV (F = 0.13, p = 0.939, Figure 2A).. The number of QTL showed a systematic effect on mean PPV (ANOVA F = 19.4, p = 1.32 x 10⁻¹⁵, Figure 2B). The distribution of allelic effects on the trait showed a huge effect on the mean PPV (mean equal contribution = 0.81, mean exponential = 0.32, F = 3409 p = 0, Figure 2C). The relation of mean proportion of detected TPL and number of migrants was approximately linear. The more migrants per generation, the higher the probability to detect more contributing loci. This effect was strong (F = 112, p = 2.35 x 10⁻⁷⁰) were observed. The number of populations screened had a similarly strong effect on proportion of TPL among the selected loci (F = 117,4 p = 4.70 x 10⁻⁹⁵). The values ranged from a mean PPV of 0.35 (s.d. = 0.21) with 12 populations to over 0.63 (s.d. = 0.32) with 60 populations. Given the chosen threshold, a diminishing return was observed above 36 populations sampled (Figure 2E). The phenotypic range in a simulation run had a low (r =

0.11, p = 3.02 x 10^{-11}), yet significantly positive effect on detection of TPL. The realised
range of population trait means in the simulations covered on average 16% (range = 0.1-
57%) of the possible range. An increase of one unit in range increased the proportion of
TPL by 0.02 (Figure 2F).

When jointly considering the effect of all parameters on PPV in a GLM, it turned out that
all had a significant effect (Table 4). Their relative influence increased from phenotypic
range (r^2 = 0.03) over heritability(r^2 = 0.06), number of QTL (r^2 = 0.07), F_{ST} (r^2 = 0.19),
number of populations (r^2 = 0.20) to the number of allelic contributions, that had by far the
greatest influence (r^2 = 0.45). In total, the parameters explained 66.4% of variance.

Table 4. Generalised Linear Model of factors influencing the proportion of TPL among
outliers (PPV) in simulations.

Factor	Coefficient	Std.err.	t	p	r^2
Constant	0.33	0.02	18.00	<2e-16	
range_pheno	0.01	0.00	4.02	0.00006	0.03
heritability	-0.14	0.01	-10.10	<2e-16	0.06
n_qtl	0.00	0.00	-10.63	<2e-16	0.07
F_{ST}	-5.72	0.19	-29.84	<2e-16	0.19
n_pop	0.01	0.00	28.47	<2e-16	0.20
allelic_contr	0.43	0.01	73.35	<2e-16	0.45

Hierarchical population structure

The introduction of a hierarchical population structure in simulations with a restricted
parameter set did not change much with regard to the relative influence of the parameters
(Supplemental Figure 6 A-F). The initial divergence parameter had no effect on PPV. The
number of QTL loci (F = 12.13, p = 7.38 x 10^{-6}), the distribution of allelic contribution (F =
888, p = 7.73 x 10^{-109}) and the number migrants per generation (F = 18.7, p = 1.88 x 10^{-5})
had a significant effect, just like the number of regions, going along with a growing number
of subpopulations (F = 4.06, p = 0.02). The final overall F_{ST} had a significant influence (p =
2.1 x 10^{-5}), albeit with a relatively low effect size (r = -0.20).

Minimum Entropy Feature Selection and statistical phenotype prediction

Minimum Entropy Feature Selection (MEFS) removed on average 8.72 (range = 2-14,
s.d. = 4.14) loci, corresponding to a proportion of 0.38 (s.d. = 0.19) from the initally chosen
outlier set. The procedure removed on average a larger proportion of FP than TPL (mean
difference 0.14, t = -19.9, p = 6.7 x 10^{-79}). This increased the proportion of TPL in the final
prediction set on average by 0.05 (range = -0.23-0.48, s.d. = 0.09) to a mean of 0.66 (range
= 0-1, s.d. = 0.29, Figure 3).

Figure 3. Effect of Minimum Entropy Feature Selection (MEFS) on the proportion of
TPL and FP in the selected set.

The predictive accuracy of the SNP loci sets selected by MEFS was on average r = 0.61 (s.d. = 0.48). It ranged from -0.96 to 1.0. The distribution was highly skewed with 75% being higher than 0.37, the median was found at 0.86 and still 25% being higher than 0.97 (Supplemental Figure 4B).

The accuracy of mean population phenotype prediction depended linearly on the number of TPL in the prediction set ($r^2 = 0.28$, p = 0), with any additional TPL increasing the correlation coefficient by 0.05. Inversely, the accuracy of prediction decreased with a rising number of FP, but even with a considerable number of FP in the prediction set, accurate prediction was possible in a large number of cases. Overall, the prediction accuracy increased with increasing proportions of TPL among the prediction set, although even 100% TPL in the prediction set did not guarantee a highly accurate prediction (r > 0.8) in all cases.

Figure 4. Influence of simulation parameters on the accuracy of statistical population mean phenotype prediction. A) Phenotypic plasticity parameter as proxy for heritability. B) Distribution of allelic trait contributions. C) Number of trait-underlying QTLs. D) Generation of independent evolution as proxy for population structure. E) Number of populations scored. F) Proportion of TPL in the prediction loci set after MEFS.

As the prediction accuracy depended on the proportion of selected TPL, their relation to
the individual simulation parameters was very similar to the results described in the
previous section, except for the distribution of allelic contributions where the prediction
accuracy was better for loci with exponential effects (Figure 4A-F). The number of
populations screened was the most important factor. With 36 or more populations

screened, 97.8% of simulations showed a prediction accuracy of 0.8 or better, independent
of the other simulation parameters applied. In a GLM with all factors simultaneously
considered, the proportion of TPL selected had the largest influence on prediction accuracy
($r^2 = 0.28$), followed by F_{ST} (0.19), the number of QTL (0.18), distribution of allelic
contributions (0.17), the number of populations screened (0.11). Heritability had only a
minor influence on the prediction accuracy (0.06), while the range of phenotypes was not
relevant (Table 5).

Table 5. Generalised Linear Model of factors influencing the accuracy of statistical
phenotypic population mean prediction.

Factor	Coefficient	Std.err.	t	p	r^2
Constant	0.69	0.04	17.31	<2e-16	
range_pheno	0.01	0.00	1.61	0.109	0.02
heritability	-0.16	0.03	-5.65	1.77E-08	0.06
n_pop	0.00	0.00	10.05	<2e-16	0.11
allelic_contr	-0.21	0.01	-15.93	<2e-16	0.17
n_qtl	0.00	0.00	-18.62	<2e-16	0.18
F_{ST}	-7.63	0.42	-18.04	<2e-16	0.19
PPV after MEFS	0.51	0.02	24.50	<2e-16	0.28

Performance comparison to individual GWAS

In every single simulation, iGWAS performed worse than popGWAS. The approach based
on individuals found on average 7.42 TPLs (PPV 0.31), compared to 12.15 (PPV 0.53) with
popGWAS (Suppl. Figure). This difference in means of 4.73 loci was highly significant in a
pairwise t-test (t >3000 , p = 0). The results for hierarchically structured population were
similar (mean difference 5.7 loci, paired t-test t = 19.62, p = 1.6 x 10^{-62}). In more than 70%
of simulations, it was not possible to calculate a valid phenotpic prediction model for
individuals with the identified loci. For the remaining cases, the mean predictive accuracy
for the individual's phenotype was 0.079.

Method performance with realistic effective genome sizes

The values for AUC-ROC ranged between 0.067 and 0.833, for AUC-PR between 0.013
and 0.730. There was an interaction between the effective genome size and number of
populations scored. According to both AUC measures, the method performed best, when the
number of populations scored was high and the genome small (Figure 5). An at least
satisfactory (> 0.66 for AUC-ROC and > 0.53 for AUC-PR) overall performance was
observed for 24 populations for the smallest genomes considered (1,000), for 36
populations up to 30,000 independent loci and for genome sizes up to 100,000 for 48 and
60. The similar values in both statistics and the plots suggested that there are diminishing
returns for samples larger that about 48 populations. Moreover, closer inspection of the
corresponding plots (Figure 6) suggested that for samples of 48 and 60 populations, an
optimal ratio between TPL and FPL exists for approximately the 25 highest outlier loci,
independent of genome size. For combinations with good performance, the maximum F1
score suggested that choosing the 30 highest outlier provided the optimal compromise
between maximising TPR and minimising FPR (Supplemental Figure 7).

550
551
552
Figure 5. Heat-map of AUC-ROC (area under the curve – receiver operator characteristics) and AUC-PR (area under the curve – precision recall) in relation to effective genome size and number of populations scored.

AUC-ROC

effective genome size	number of populations scored				
	12	24	36	48	60
1000	0.391	0.663	0.769	0.817	0.833
5000	0.198	0.430	0.677	0.746	0.776
10000	0.112	0.459	0.623	0.700	0.754
30000	0.094	0.375	0.668	0.668	0.697
50000	0.096	0.359	0.539	0.644	0.696
100000	0.067	0.339	0.528	0.616	0.670

AUC-PR

effective genome size	number of populations scored				
	12	24	36	48	60
1000	0.250	0.514	0.644	0.709	0.730
5000	0.125	0.430	0.552	0.623	0.654
10000	0.060	0.338	0.508	0.587	0.644
30000	0.038	0.265	0.464	0.553	0.582
50000	0.032	0.238	0.422	0.532	0.577
100000	0.013	0.182	0.378	0.497	0.551

554
555
556
557

Figure 6. AUC-ROC and AUC-PR for a range of effective genome sizes. In the left column are the plots of AUC-ROC, i.e. FDR on the x-axis versus TPR on the y-axis. The right column shows AUC-PR plots, i.e. TPR on the x-axis versus PPV on the y-axis. The dotted lines indicate the threshold for a random effect.

Discussion

This study used extensive forward simulations to explore the potential of a novel GWAS
approach utilising phenotypic population means and genome-wide allele-frequency data to
identify loci potentially underlying the population differences in quantitative polygenic
traits and predict the mean phenotypes of unmeasured populations. While the approach
seems to be generally useful in a wide range of cases, there are also clear limits to its
applicability.

General validity of the underlying assumptions

The first major goal of the study was understanding the conditions under which the
hypothesised pattern of a linear relation between the population allele frequencies at causal
loci and the phenotypic population means of the respective trait emerges. The initial
simulations demonstrated that the expectation for the covariance of random population
"allele frequencies" at contributing quantitative trait loci (QTL) and the respective
population trait mean is consistently positive when an additive model applies. This is an
inherent consequence of the common dependence of both variables on the QTL genotypes
of the individuals in a population, as demonstrated in (3). Additive models seem to be an
appropriate statistical approximation for most quantitative traits at population level (Hill et
al., 2008), despite the description of many epistatic interactions on the molecular level
(Moore & Williams, 2005).
The relation appeared to be largely independent of the distribution shape of locus
contributions to the trait. While in the case of equal contributions i.e. a flat distribution, the
correlation coefficients of individual loci are themselves a random variate, normally
distributed around the expected mean. As the distribution becomes increasingly skewed,
locus contribution becomes predictive of the correlation to the trait. Loci contributing more
to the trait and thus accounting for more of the phenotypic variance will likely have a higher
correlation of their allele frequencies to the population mean. Conversely, the expectation
for non-contributing loci is approaching zero. Therefore, it is principally possible to exploit
the correlation between allele frequencies and population trait means for the identification
of loci underlying an additive quantitative trait.
However, some statistical limitations became obvious. Firstly, as the number of QTL
increases, the expected mean correlation coefficients become so small that they are likely to
be indistinguishable from the tail of the zero-centered normal distribution of non-
contributing loci, even with an unrealistically high number of samples. Consequently, the
method for identifying QTL by the positive covariance of their allele frequencies with the
population trait means is *a priori* more suited for oligogenic to moderately polygenic traits.
Secondly, the number of QTL and the distribution of locus contributions may influence the
statistical identifiability of individual QTL. In particular, loci that contribute only minimally
to the trait or that fall by chance below the expected mean correlation coefficient may
overlap with the tail of the distribution of non-contributing loci.
These predictions remind of similar conditions for the contribution of different QTL
architectures to phenotypic adaptation described by (Höllinger et al., 2023). They assert that
phenotypic adaptation of oligogenic traits is achieved by detectable allele frequency shifts
at some but not very many loci, while adaptation in highly polygenic traits is rather
achieved by subtle perturbations of standing variation, with respective consequences for

their detectability. Just as expected here, they stress the importance of stochastic effects
that may lead to apparently heterogeneous locus contributions (Höllinger et al., 2023).

Interestingly, deviation from a selection-drift-migration equilibrium were not an
impediment for detecting contributing loci. Best possible adaptation was regularly achieved
within less than 10 generations. True to theoretical expectations, higher gene-flow among
populations led to less perfect local adaptation for a given selective strength (Brady et al.,
2019). Drift-migration balance was not reached within a reasonable number of generations.
This was, however, not relevant for popGWAS performance, which was mainly driven by
the degree of population differentiation, everything else being equal. This insensitivity to
equilibrium conditions is an advantage of the method as natural populations tend not to be
in equilibrium (Müller et al., 2022). Rather, natural populations are constantly tracking
variable selective optima (Pfenninger & Foucault, 2022; Rudman et al., 2021), which leads
to intermediate allele frequencies at the loci under such a selective regime (Höllinger et al.,
2023).

Limiting factors for the identification of contributing loci in natural settings

The Wright-Fisher forward simulations of a quantitative trait in a subdivided population
with realistic properties and sample sizes largely confirmed the theoretical expectations. In
particular when a sufficient number of populations was scored (in this case more than 36),
a large proportion of true positive loci could be reliably identified, with the exception of a
few parameter combinations. The genetic architecture of the trait was an important
predictor for the ability to identify causal loci. While loci underlying oligogenic and
moderately polygenic traits could be fairly reliably identified, the highly polygenic scenario
tested (500 loci) performed poorly. A higher proportion of TPL was identified when the
locus contribution to the trait was identical than in the case of an exponential trait
contribution distribution This was likely due to the tendency of higher correlations
between higher contributing loci and the trait, which allowed the loci from the tail of the
distribution to vary freely, making them indistinguishable from non-contributing loci.

The influence of mean heritability was on GWAS performance was not marked. Even
down to trait heritability estimates of 0.3, the success rate was only slightly reduced. This
effect may be attributed to the averaging of phenotypes and genotypes across multiple
individuals, which is likely to mitigate the inherent noise associated with individual data
(Johri et al., 2022; Stinchcombe & Hoekstra, 2008). This finding is consistent with
observations by (Zhang et al., 2018), who employed pooled data for GWAS. From a practical
standpoint, the findings suggest that inevitable errors in phenotyping, which can
compromise GWAS performance on individuals (Barendse, 2011), are likely to be less
problematic when using the mean measured over many individuals. Furthermore, this
finding indicates that the failure to entirely remove non-additive variance from the analysis
does not necessarily compromise the method's ability to reliably identify trait-associated
loci.

From a statistical perspective, it was anticipated that the range of phenotypic population
means would influence the identification of true positive loci to some extent, given that a
larger range of phenotypic means is inherently associated with on average larger allele-
frequency differences among populations. The choice of populations with a large range of
environmentally unexplained variance is therefore crucial. It is, however, important to
emphasise that the underlying causes of the observed differences in trait means among
populations are not of primary concern. These may be attributed to local adaptation, but

also to maladaptation, human choice, or other factors. Likewise, increasing the number of
populations screened increased the statistical power of the approach. However, it seemed
that increasing the number of samples led beyond a certain threshold to diminishing
returns in statistical power gain.

Effect of population structure on the detection of contributing loci

It is long-since known that population structure can have a confounding effect on GWAS
studies (Marchini et al., 2004). Confounding associations between two-allelic SNPs and a
phenotype may easily arise in iGWAS, because phenotypically more similar individuals may
be also genetically more similar because they share common ancestry as members of the
same family/population/clade. In addition, a quantitative or qualitative phenotype (y) in
iGWAS is related to a genotype (x) either by a linear or logistic model – with exactly three
possible values of x. This leads inherently to low statistical power, requiring a very high
number of individuals (O'Connor, 2021) and extensive corrections for population structure
to obtain a reliable statistical association (Sul et al., 2018). popGWAS follows a principally
different approach; the possible values of x are allele frequencies in different populations
and thus limited only by sampling effort. This gives the approach a strong inherent
statistical advantage, which might be also the reason that the method does not suffer from
p-value inflation.
Population structure can nevertheless interfere with reliable associations in popGWAS,
albeit for different reasons. A pronounced population structure ($F_{ST} > \sim 0.07$) was a indeed
major factor impeding reliable identification of true positive loci, even with a high number
of samples and regardless of the reasons for the structure. However, this was probably
more due to distinct evolutionary trajectories in independently evolving populations. If the
evolutionary trajectory of the populations analysed is sufficiently independent, either by
drift and/or differential availability of adaptive mutations, similar phenotypes may have a
differential genomic basis, allowing at best to identify any common variants. The genetic
redundancy of polygenic traits can lead to evolution of the same phenotypes from different
genomic bases (de Vladar & Barton, 2014; Kaneko & Furusawa, 2006), even if evolving from
the same ancestral population (Barghi et al., 2019, 2020; Pfenninger et al., 2015). If
different loci in different populations are causal for the observed phenotypic differences, a
linear relation between population means and allele frequencies is not to be expected. It is
therefore important that the allele frequencies in the studied populations are correlated
either by recent common descent and/or recurrent gene-flow, i.e. that the population
structure between the population scored is weak (Mathieson, 2021).
By relating allele frequencies with mean phenotypes false associations may also arise, if
the trait under scrutiny is effectively neutral and thus the constituting loci follow the overall
drift pattern or the ecological differences causing the phenotypic differences are associated
to an isolation-by-distance pattern. In this case, the matrix of mean phenotypic population
differences (or Q_{ST}) should show a strong positive correlation with the overall F_{ST} matrix
among populations. A trait showing this easily testable pattern may therefore *a priori* not
be suitable for analysis with the method. To avoid such a situation, it is recommended to
test for (the absence of) a correlation between genome-wide genetic distance and
differences in phenotypic means (e.g. by a Mantel's test).

Accurate statistical genomic prediction in a wide range of conditions

Genomic prediction is deemed to be one of the major tools for the mitigation of climate
change on biodiversity (Aguirre-Liguori et al., 2021; Bernatchez et al., 2023; Capblancq et al.,
2020; Waldvogel et al., 2020). One of the major findings of the presented approach was the
highly accurate prediction of the phenotypic population means from genomic data above a
certain number of populations screened. Given the limited performance of genomic
prediction in a medical context (Visscher et al., 2017) this was initally quite surprising.
However, it should be *a priori* easier to predict the differences in population means
compared to differences among individuals for the following reasons. The difference in
means among subpopulations was mostly governed by only a couple of loci, because the
range of phenotypic population means extended over only a relatively small part of the
theoretically possible range. This applies to the simulations but most likely also in nature it
is improbable to encounter populations that are (nearly) fixed for the respectively
alternative alleles over all or even most trait contributing loci and show an accordingly
extended phenotypic range. Consequently, already the allele frequency differences of a
couple of loci actually need to follow a linear model to explain most of the variance among
phenotypic population means – the rest is free to vary and behave (almost) like neutral loci.
Even the fit of the highest outlier loci to a linear model was far from perfect as was expected
from the genetic redundancy of a polygenic trait architecture (Barghi et al., 2020).
However, as long as these important loci are the same over the majority of subpopulations
(either due to common descent or gene-flow), accurate predictions of population trait
means from these loci is therefore possible. Such loci likely have an intermediate frequency,
because alleles of very low or very high frequency (i.e. present only in few or alomost all
individuals with not necessarily extreme phenotypes) by definition cannot have a large
impact on the phenotypic population mean, even when of large effect. Conversely, the
individual phenotype can be mainly determined by rare alleles of large effects. Therefore, a
locus whose AF is important for the difference in means among populations might have
rather low predictive power for the individual, because as a locus of likely intermediate
frequency, all genotypes are necessarily realised with high probability within a population.
The accurate prediction of individual phenotypes thus likely requires much more loci than
the prediction of the phenotypic population mean.

Finally, it is crucial to distinguish between two major goals of GWAS: i) find the loci
functionally underlying biologically relevant variation and ii) the accurate prediction of
untyped instances. As (Shmueli, 2010) pointed out, the goals of explaining a phenomenon
on the one hand and predicting it on the other are related but not identical. In particular it is
important to note that an accurate prediction model does not need to contain all features
that are necessary for a comprehensive functional description of the system (Shmueli,
2010). For accurate prediction, it is not even necessary that all features in the prediction
model have actually a functional role, as long as they are statistically strongly associated
(Shmueli, 2010). While the results have shown that prediction tends to become more
accurate if more truly functional loci are included, it is therefore not necessary that the
prediction model includes all or even most of the functional loci.

Contrary to its application in medicine or selective breeding (Wray et al., 2019), however,
accurate prediction of population responses is in many instance of the current biodivsity
crisis probably more important than the prediction of individual phenotypes. Within the
limits outlined above, the proposed method delivered in many instances very accurate
predictions (r > 0.8 in > 90% of cases for more than 36 populations) of population mean

phenotypes. It should be noted, however, that the prediction is statistical in the sense that it
produces a prediction score (de Los Campos et al., 2018) that correlates with the mean
population phenotype and is explictely not a comprehensive functional quantitative genetic
model of the trait Just like with any other genomic prediction (Kachuri et al., 2024), this
limits the transferability of the prediction to other, more distantly related lineages or
species.

Reducing the false positive rate before prediction is in any case advisable, as it proved to
be the most important factor of prediction success with independent data. The application
of a Machine learning approach, in this case Minimum Entropy Feature Selection (MEFS),
prior to prediction effectively reduced the already low false positive rate among the initially
selected loci further. Whether other approaches would yield even better results remains to
be tested. Other population genetic parameters influenced prediction success in a very
similar fashion as the initial true positive rate. One notable exception was distribution of
locus contributions. While true positive loci were more reliably identified from a flat
distribution, prediction worked better when many loci of large effect were among the
prediction set, most likely because these loci contribute more to phenotypic variance
among populations (Jain & Stephan, 2015).

Typical genome sizes of real species are no obstacle

The perhaps most important challenge was showing that the proposed method has
enough statistical power to distinguish at least a part of the unknown, but likely relatively
small number of QTL reliably from the large number of non-contributing loci in real
genomes of real species. The evaluation of method performance with AUC-ROC and AUC-PR,
as recommended recently (Lotterhos et al., 2022), showed a satisfactory performance even
for genomes with moderately high effective sizes, provided a sufficiently high number of
populations is screened. In particular restricting the selection of potentially causal QTL on a
few dozen of the highest outliers promises to yield very low false positive rates. However,
as the success rate depended very much on the biological characteristics like genome size,
population structure, LD and so on, of the investigated taxon, no general recommendations
can be given here, except to carefully take them into account. As shown above, already a
limited number of true positive loci may be sufficient for reliable genomic prediction.

Practical considerations

The proposed method finds rather genomic regions or haplotypes associated to the trait
in question than directly causal SNPs. However, this is true for most GWAS methods (Wang
et al., 2010) and therefore fine-mapping and inference of causal processes remain to be
done (Wallace, 2021). In practice, this requires that regions with high SNP outlier density
need to be collapsed to haplotypes prior to further analysis. Knowledge on the local LD-
structure, mean haplotype length, respectively recombination landscape can aid haplotype
identification (Flister et al., 2013). Recently developed machine learning approaches makes
such information available for pooled data (Adrion et al., 2020).
The possibly largest advantage of the proposed method is therefore its data efficiency, if
pooled sequencing is applied. Because the Pool-Seq approach (Schlötterer et al., 2014) yields
highly accurate estimates of genome-wide allele frequencies at SNP sites (Czech, Peng,
Spence, Lang, Bellagio, Hildebrandt, Fritschi, Schwab, Rowan, & consortium, 2022) the
necessary sequencing effort is marginal compared to individual based approaches
(Ziyatdinov et al., 2021). This is mainly due to the relatively reduced costs for library

preparations necessary that, depending on the genome size, often make up a large share of
or even exceed the actual sequencing costs in individual analysis. Moreover, the popGWAS
method out-performed iGWAS in all cases, if the same sequencing effort is applied as a
benchmark.

The proposed method makes GWAS studies accessible for the usual funding in the field
of biodiversity. Pooled sequencing for GWAS has been proposed (Yang et al., 2015) and
applied (Giorello et al., 2023; Kumar et al., 2022; Pfenninger et al., 2021) with extreme
phenotypes. Automated large scale phenotyping is a very active field of research, with
satellite or other remote sensing approaches, flow cytometry, bulk measurements of
metabolic rates, metabolomes, transcriptomes, pooled HPLC or mass spectrometry,
automated video-based phenotyping, mass CT scanning provides already access to large
amounts of data. Sampling the required number of demes or subpopulations can pose a
challenge for certain taxa, however, e.g. biodiversity monitoring schemes or museum
collections may offer readily access to respective samples. Application of the method to a
real world data set from European beech on phenological traits (Pfenninger et al. in prep),
showed that such an approach can be sucessfully performed with moderate logistic effort.
Many of the identified loci by popGWAS were previously implicated in phenological traits
and yielded excellent predictive power in predicting the mean phenotypes of independent
populations (Pfenninger et al. in prep).

Conclusion

This study demonstrated the potential of the proposed GWAS approach for biodiversity
genomics. As with any GWAS method, its usefulness depends on the study goals. If it is the
aim to infer all loci in a genome contributing to a given quantitative trait and to quantify
their relative influence, probably no existing (single) GWAS method and no experimental
design can achieve this. If it is the goal to accurately predict individual quantitative
phenotypes with the inferred genomic basis, e.g. for medical or breeding purposes, the
suggested method may not lead very far, as detailed above. Nevertheless, it can add a
substantial number of candidate loci that are not easily found by other methods. However, if
it is important to to infer the major contributing loci for quantitative differences in
ecologically relevant traits observed among natural populations, popGWAS can be a
valuable tool. By carefully considering the factors influencing its performance and
addressing the limitations, this method holds some promise in identifying the genetic basis
of complex traits and performing accurate statistical predictions of phenotypic population
means from genomic data in natural populations.

Acknowledgements

The author wants to thank Bob O'Hara, Barbara Feldmeyer and the reviewers for helpful
comments on the manuscript.

Funding

The author declares that he has received no specific funding for this study.

Conflict of interest disclosure

The author declares that he complies with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Supplementary information including the Python code used for the simulations is available at https://**10.5281/zenodo.11562472**

References

Adrion, J. R., Galloway, J. G., & Kern, A. D. (2020). Predicting the landscape of recombination using deep learning. *Molecular Biology and Evolution*, *37*(6), 1790–1808.

Aguirre-Liguori, J. A., Ramírez-Barahona, S., & Gaut, B. S. (2021). The evolutionary genomics of species' responses to climate change. *Nature Ecology & Evolution*, *5*(10), 1350–1360.

Barendse, W. (2011). The effect of measurement error of phenotypes on genome wide association studies. *BMC Genomics*, *12*(1), 232. https://doi.org/10.1186/1471-2164-12-232

Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: A unifying framework to understand positive selection. *Nature Reviews Genetics*, *21*(12), 769–781.

Barghi, N., Tobler, R., Nolte, V., Jakšić, A. M., Mallard, F., Otte, K. A., Dolezal, M., Taus, T., Kofler, R., & Schlötterer, C. (2019). Genetic redundancy fuels polygenic adaptation in Drosophila. *PLoS Biology*, *17*(2), e3000128.

Barton, N. H. (1999). Clines in polygenic traits. *Genetics Research*, *74*(3), 223–236.

Bernatchez, L., Ferchaud, A.-L., Berger, C. S., Venney, C. J., & Xuereb, A. (2023). Genomics for monitoring and understanding species responses to global climate change. *Nature Reviews Genetics*, 1–19.

Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An expanded view of complex traits: From polygenic to omnigenic. *Cell*, *169*(7), 1177–1186.

Brady, S. P., Bolnick, D. I., Angert, A. L., Gonzalez, A., Barrett, R. D., Crispo, E., Derry, A. M., Eckert, C. G., Fraser, D. J., & Fussmann, G. F. (2019). Causes of maladaptation. *Evolutionary Applications*, *12*(7), 1229–1242.

Brandes, N., Weissbrod, O., & Linial, M. (2022). Open problems in human trait genetics. *Genome Biology*, *23*(1), 131. https://doi.org/10.1186/s13059-022-02697-9

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., & Keller, S. R. (2020). Genomic prediction of (mal) adaptation across current and future climatic landscapes. *Annual Review of Ecology, Evolution, and Systematics*, *51*, 245–269.

Chakraborty, R. (1981). The distribution of the number of heterozygous loci in an individual in natural populations. *Genetics*, *98*(2), 461.

Czech, L., Peng, Y., Spence, J., Lang, P., Bellagio, T., Hildebrandt, J., Fritschi, K., Schwab, R., Rowan, B., & Weigel, D. (2022). Efficient analysis of allele frequency variation from whole-genome pool-sequencing data. *Population, Evolutionary, and Quantitative Genetics Conference (PEQG 2022)*, 99. https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item_3474009

Czech, L., Peng, Y., Spence, J. P., Lang, P. L., Bellagio, T., Hildebrandt, J., Fritschi, K., Schwab, R., Rowan, B. A., & consortium, G. (2022). Monitoring rapid evolution of plant populations at scale with Pool-Sequencing. *BioRxiv*, 2022–02.

de Los Campos, G., Vazquez, A. I., Hsu, S., & Lello, L. (2018). Complex-trait prediction in the
era of big data. *Trends in Genetics*, *34*(10), 746–754.

de Vladar, H. P., & Barton, N. (2014). Stability and response of polygenic traits to stabilizing
selection and mutation. *Genetics*, *197*(2), 749–767.

Dunker, S., Boyd, M., Durka, W., Erler, S., Harpole, W. S., Henning, S., Herzschuh, U., Hornick,
T., Knight, T., Lips, S., Mäder, P., Švara, E. M., Mozarowski, S., Rakosy, D., Römermann, C.,
Schmitt-Jansen, M., Stoof-Leichsenring, K., Stratmann, F., Treudler, R., … Wilhelm, C.
(2022). The potential of multispectral imaging flow cytometry for environmental
monitoring. *Cytometry Part A*, *101*(9), 782–799. https://doi.org/10.1002/cyto.a.24658

Exposito-Alonso, M., Drost, H., Burbano, H. A., & Weigel, D. (2020). The Earth BioGenome
project: Opportunities and challenges for plant genomics and conservation. *The Plant*
Journal, *102*(2), 222–229. https://doi.org/10.1111/tpj.14631

Flister, M. J., Tsaih, S.-W., O'Meara, C. C., Endres, B., Hoffman, M. J., Geurts, A. M., Dwinell, M.
R., Lazar, J., Jacob, H. J., & Moreno, C. (2013). Identifying multiple causative genes at a
single GWAS locus. *Genome Research*, *23*(12), 1996–2002.

Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C.,
Crottini, A., Godoy, J. A., & Höglund, J. (2022). The era of reference genomes in
conservation genomics. *Trends in Ecology & Evolution*, *37*(3), 197–202.

Giorello, F. M., Farias, J., Basile, P., Balmelli, G., & Da Silva, C. C. (2023). Evaluating the
potential of XP-GWAS in Eucalyptus: Leaf heteroblasty as a case study. *Plant Gene*, *36*,
100430.

Harpak, A., & Przeworski, M. (2021). The evolution of group differences in changing
environments. *PLoS Biology*, *19*(1), e3001072.

Heuertz, M., Carvalho, S. B., Galindo, J., Rinkevich, B., Robakowski, P., Aavik, T., Altinok, I.,
Barth, J. M., Cotrim, H., & Goessen, R. (2023). The application gap: Genomics for
biodiversity and ecosystem service management. *Biological Conservation*, *278*, 109883.

Hill, W. G., Goddard, M. E., & Visscher, P. M. (2008). Data and theory point to mainly additive
genetic variance for complex traits. *PLoS Genetics*, *4*(2), e1000008.

Hogg, C. J. (2023). Translating genomic advances into biodiversity conservation. *Nature*
Reviews Genetics, 1–12.

Höllinger, I., Wölfl, B., & Hermisson, J. (2023). A theory of oligogenic adaptation of a
quantitative trait. *Genetics*, *225*(2), iyad139. https://doi.org/10.1093/genetics/iyad139

Jain, K., & Stephan, W. (2015). Response of polygenic traits under stabilizing selection and
mutation when loci have unequal effects. *G3: Genes, Genomes, Genetics*, *5*(6), 1065–
1074.

Johri, P., Aquadro, C. F., Beaumont, M., Charlesworth, B., Excoffier, L., Eyre-Walker, A.,
Keightley, P. D., Lynch, M., McVean, G., & Payseur, B. A. (2022). Recommendations for
improving statistical inference in population genomics. *PLoS Biology*, *20*(5), e3001669.

Kachuri, L., Chatterjee, N., Hirbo, J., Schaid, D. J., Martin, I., Kullo, I. J., Kenny, E. E., Pasaniuc,
B., Yuji 29, P. R. M. in D. P. (PRIMED) C. M. W. G. A. P. L. 20 C. M. P. 21 C. D. V. 22 23 D. Y.
24 W. Y. 19 25 26 Z. H. 27 28 Z., & Witte, J. S. (2024). Principles and methods for
transferring polygenic risk scores across global populations. *Nature Reviews Genetics*,
25(1), 8–25.

Kaneko, K., & Furusawa, C. (2006). An evolutionary relationship between genetic variation
and phenotypic fluctuation. *Journal of Theoretical Biology*, *240*(1), 78–86.

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., Kosiol, C., &
Schlötterer, C. (2011). PoPoolation: A toolbox for population genetic analysis of next
generation sequencing data from pooled individuals. *PloS One*, *6*(1), e15925.

Kumar, S., Deng, C. H., Molloy, C., Kirk, C., Plunkett, B., Lin-Wang, K., Allan, A., & Espley, R. (2022). Extreme-phenotype GWAS unravels a complex nexus between apple (*Malus domestica*) red-flesh colour and internal flesh browning. *Fruit Research*, *2*(1), 1–14. https://doi.org/10.48130/FruRes-2022-0012

Lotterhos, K. E., Fitzpatrick, M. C., & Blackmon, H. (2022). Simulation Tests of Methods in Evolution, Ecology, and Systematics: Pitfalls, Progress, and Principles. *Annual Review of Ecology, Evolution, and Systematics*, *53*(1), 113–136. https://doi.org/10.1146/annurev-ecolsys-102320-093722

Lynch, M., & Walsh, B. (1998). *Genetics and analysis of quantitative traits* (Vol. 1). Sinauer Sunderland, MA.

Mackay, T. F. (2014). Epistasis and quantitative traits: Using model organisms to study gene–gene interactions. *Nature Reviews Genetics*, *15*(1), 22–33.

Marchini, J., Cardon, L. R., Phillips, M. S., & Donnelly, P. (2004). The effects of human population structure on large genetic association studies. *Nature Genetics*, *36*(5), 512–517.

Mathieson, I. (2021). The omnigenic model and polygenic prediction of complex traits. *The American Journal of Human Genetics*, *108*(9), 1558–1563.

Moore, J. H., & Williams, S. M. (2005). Traversing the conceptual divide between biological and statistical epistasis: Systems biology and a more modern synthesis. *BioEssays*, *27*(6), 637–646. https://doi.org/10.1002/bies.20236

Müller, R., Kaj, I., & Mugal, C. F. (2022). A nearly neutral model of molecular signatures of natural selection after change in population size. *Genome Biology and Evolution*, *14*(5), evac058.

O'Connor, L. J. (2021). The distribution of common-variant effect sizes. *Nature Genetics*, *53*(8), 1243–1249.

Orr, H. A. (1998). Testing natural selection vs. Genetic drift in phenotypic evolution using quantitative trait locus data. *Genetics*, *149*(4), 2099–2104.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830.

Pfenninger, M., & Foucault, Q. (2022). *Population genomic time series data of a natural population suggests adaptive tracking of fluctuating environmental changes* (p. in press). Integrative and Comparative Biology. https://doi.org/10.1101/2020.06.16.154054

Pfenninger, M., Patel, S., Arias-Rodriguez, L., Feldmeyer, B., Riesch, R., & Plath, M. (2015). Unique evolutionary trajectories in repeated adaptation to hydrogen sulphide-toxic habitats of a neotropical fish (*Poecilia mexicana*). *Molecular Ecology*, *24*(21), 5446–5459. https://doi.org/10.1111/mec.13397

Pfenninger, M., Reuss, F., Kiebler, A., Schönnenbeck, P., Caliendo, C., Gerber, S., Cocchiararo, B., Reuter, S., Blüthgen, N., & Mody, K. (2021). Genomic basis for drought resistance in European beech forests threatened by climate change. *Elife*, *10*, e65532.

Pritchard, J. K., & Di Rienzo, A. (2010). Adaptation–not by sweeps alone. *Nature Reviews Genetics*, *11*(10), 665–667.

R Core Team, R. (2013). *R: A language and environment for statistical computing*.

Rijsbergen, C. van. (1979). *Information retrieval*. Butterworth-Heinemann. https://dl.acm.org/doi/abs/10.5555/539927

Rudman, S. M., Greenblum, S. I., Rajpurohit, S., Betancourt, N. J., Hanna, J., Tilk, S., Yokoyama, T., Petrov, D. A., & Schmidt, P. (2021). *Direct observation of adaptive tracking on*

ecological timescales in Drosophila (p. 2021.04.27.441526). bioRxiv.
https://doi.org/10.1101/2021.04.27.441526

Santure, A. W., & Garant, D. (2018). Wild GWAS—association mapping in natural
populations. *Molecular Ecology Resources*, *18*(4), 729–738.
https://doi.org/10.1111/1755-0998.12901

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals—
Mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics*,
15(11), 749–763.

Sella, G., & Barton, N. H. (2019). Thinking about the evolution of complex traits in the era of
genome-wide association studies. *Annual Review of Genomics and Human Genetics*, *20*,
461–493.

Shendure, J., Findlay, G. M., & Snyder, M. W. (2019). Genomic medicine–progress, pitfalls,
and promise. *Cell*, *177*(1), 45–57.

Shmueli, G. (2010). *To explain or to predict?* https://projecteuclid.org/journals/statistical-
science/volume-25/issue-3/To-Explain-or-to-Predict/10.1214/10-STS330.short

Stinchcombe, J. R., & Hoekstra, H. E. (2008). Combining population genomics and
quantitative genetics: Finding the genes underlying ecologically important traits.
Heredity, *100*(2), 158–170.

Sul, J. H., Martin, L. S., & Eskin, E. (2018). Population structure in genetic studies:
Confounding factors and mixed models. *PLoS Genetics*, *14*(12), e1007309.

Taylor, C. F., & Higgs, P. G. (2000). A population genetics model for multiple quantitative
traits exhibiting pleiotropy and epistasis. *Journal of Theoretical Biology*, *203*(4), 419–
437.

Team, T. P. (2019, December 28). *PyPy*. PyPy. https://www.pypy.org/

Tills, O., Holmes, L. A., Quinn, E., Everett, T., Truebano, M., & Spicer, J. I. (2023). Phenomics
enables measurement of complex responses of developing animals to global
environmental drivers. *Science of the Total Environment*, *858*, 159555.

Turchin, M. C., Chiang, C. W. K., Palmer, C. D., Sankararaman, S., Reich, D., & Hirschhorn, J. N.
(2012). Evidence of widespread selection on standing variation in Europe at height-
associated SNPs. *Nature Genetics*, *44*(9), 1015–1019. https://doi.org/10.1038/ng.2368

Uffelmann, E., Huang, Q. Q., Munung, N. S., De Vries, J., Okada, Y., Martin, A. R., Martin, H. C.,
Lappalainen, T., & Posthuma, D. (2021). Genome-wide association studies. *Nature
Reviews Methods Primers*, *1*(1), 59.

Van Rossum, G., & Drake, F. L. (2009). *Introduction to python 3: Python documentation
manual part 1*. CreateSpace. https://dl.acm.org/doi/abs/10.5555/1592885

Visscher, P. M., Brown, M. A., McCarthy, M. I., & Yang, J. (2012). Five years of GWAS
discovery. *The American Journal of Human Genetics*, *90*(1), 7–24.

Visscher, P. M., Wray, N. R., Zhang, Q., Sklar, P., McCarthy, M. I., Brown, M. A., & Yang, J.
(2017). 10 years of GWAS discovery: Biology, function, and translation. *The American
Journal of Human Genetics*, *101*(1), 5–22.

Waldvogel, A.-M., Feldmeyer, B., Rolshausen, G., Exposito-Alonso, M., Rellstab, C., Kofler, R.,
Mock, T., Schmid, K., Schmitt, I., & Bataillon, T. (2020). Evolutionary genomics can
improve prediction of species' responses to climate change. *Evolution Letters*, *4*(1), 4–
18.

Wallace, C. (2021). A more accurate method for colocalisation analysis allowing for multiple
causal variants. *PLoS Genetics*, *17*(9), e1009440.

Wang, K., Dickson, S. P., Stolle, C. A., Krantz, I. D., Goldstein, D. B., & Hakonarson, H. (2010).
Interpretation of association signals and identification of causal variants from genome-
wide association studies. *The American Journal of Human Genetics*, *86*(5), 730–742.
Wray, N. R., Kemper, K. E., Hayes, B. J., Goddard, M. E., & Visscher, P. M. (2019). Complex
trait prediction from genome data: Contrasting EBV in livestock to PRS in humans:
genomic prediction. *Genetics*, *211*(4), 1131–1141.
Wright, S. (1949). THE GENETICAL STRUCTURE OF POPULATIONS. *Annals of Eugenics*,
15(1), 323–354. https://doi.org/10.1111/j.1469-1809.1949.tb02451.x
Xie, C., & Yang, C. (2020). A review on plant high-throughput phenotyping traits using UAV-
based sensors. *Computers and Electronics in Agriculture*, *178*, 105731.
Yang, J., Jiang, H., Yeh, C.-T., Yu, J., Jeddeloh, J. A., Nettleton, D., & Schnable, P. S. (2015).
Extreme-phenotype genome-wide association study (XP-GWAS): A method for
identifying trait-associated variants by sequencing pools of individuals selected from a
diversity panel. *The Plant Journal*, *84*(3), 587–596.
Zhang, W., Liu, A., Albert, P. S., Ashmead, R. D., Schisterman, E. F., & Mills, J. L. (2018). A
pooling strategy to effectively use genotype data in quantitative traits genome-wide
association studies. *Statistics in Medicine*, *37*(27), 4083–4095.
https://doi.org/10.1002/sim.7898
Ziyatdinov, A., Kim, J., Prokopenko, D., Privé, F., Laporte, F., Loh, P.-R., Kraft, P., & Aschard, H.
(2021). Estimating the effective sample size in association studies of quantitative traits.
G3, *11*(6), jkab057.
1030
1031