On the potential for GWAS with phenotypic population means and allele-frequency data (popGWAS)

Pfenninger Markus[1,2,3]*

[1] Dept. Molecular Ecology, Senckenberg Biodiversity and Climate Research Centre, Georg-Voigt-Str. 14-16, D-60325 - Frankfurt am Main, Germany
[2] LOEWE Centre for Translational Biodiversity Genomics, Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, D-60325 – Frankfurt am Main, Germany
[3] Institute for Molecular and Organismic Evolution, Johannes Gutenberg University, Johann-Joachim-Becher-Weg 7, D-55128 – Mainz, Germany

*Corresponding author
Correspondence: Markus.Pfenninger@senckenberg.de

ABSTRACT

This study explores the potential of a novel genome-wide association study (GWAS) approach for identifying loci underlying quantitative polygenic traits in natural populations. Extensive population genetic forward simulations demonstrate that the approach is generally effective for oligogenic and moderately polygenic traits and relatively insensitive to low heritability, but applicability is limited for highly polygenic architectures and pronounced population structure. The required sample size is moderate with very good results being obtained already for a few dozen populations scored. The method performs well in predicting population means even with a moderate false positive rate. When combined with machine learning for feature selection, this rate can be further reduced. The data efficiency of the method, particularly when using pooled sequencing, makes GWAS studies more accessible for research in biodiversity genomics. Overall, this study highlights the promise of this popGWAS approach for dissecting the genetic basis of complex traits in natural populations.

Keywords: biodiversity population genomics, molecular trait basis

Introduction

A major goal as well as a major challenge in evolutionary biology is to understand how genes influence traits, i.e. the genotype-phenotype link, (Brandes et al., 2022; Uffelmann et al., 2021). The difficulties in achieving this goal are primarily due to the fact that the heritable variation of many, if not most, relevant phenotypes is determined by small contributions from many genetic loci (Sella & Barton, 2019). Such complex traits are usually influenced by a few dozen genes that are mechanistically directly involved in their expression, but often also by numerous, if not almost all, other genes as well as the environment in the widest sense (Boyle et al., 2017).

Genome-wide association studies (GWAS) are commonly used to link complex phenotypic traits to their genomic basis (Brandes et al., 2022; Visscher et al., 2012). However, given the complexity of causal mechanisms and the small effects of individual loci, often only a small fraction of the genetic variation underlying phenotypic variance is often identified, despite the considerable logistic effort in terms of the number of phenotyped and genotyped individuals (Brandes et al., 2022; Visscher et al., 2017). As a result, accurate predictions of phenotypes from genomic data are still quite limited and there is currently no other strategy than to keep increasing sample sizes (Brandes et al., 2022). This is a problem in the medical sciences (Shendure et al., 2019), but the greater challenge for science and society probably lies in addressing the global biodiversity crisis. It would be highly desirable to have affordable methods to accurately understand the genomic basis of relevant traits and predict (non-model) species responses to all aspects of global change (Bernatchez et al., 2023; Waldvogel et al., 2020).

GWAS with wild populations has been advocated for some time (Santure & Garant, 2018). However, despite recent progress in high throughput, automated phenotyping (Dunker et al., 2022; Tills et al., 2023; Xie & Yang, 2020), the advances of biodiversity genomics in obtaining high quality reference genomes for almost every species (Exposito-Alonso et al., 2020; Formenti et al., 2022) and the possibility to gain cost-effective genome-wide population data (Czech, Peng, Spence, Lang, Bellagio, Hildebrandt, Fritschi, Schwab, Rowan, & Weigel, 2022; Schlötterer et al., 2014), relatively few empirical studies are currently available. This gap between the possibilities and actual practical application in biodiversity conservation (Heuertz et al., 2023; Hogg, 2023) is probably as much due to the still existing logistic and financial challenges as to a lack of data- and resource-efficient methods.

Here, I explore the potential of a new GWAS approach using phenotypic population means and genome-wide allele-frequency data. The rationale behind the approach is straightforward. If a quantitative polygenic trait has an additive genetic component, an individual's phenotypic trait value should at least roughly correlate with the number of trait-increasing alleles at the underlying loci (Uffelmann et al., 2021). Consequently, it was theoretically expected (Orr, 1998; Pritchard & Di Rienzo, 2010) and empirically shown (Turchin et al., 2012) that trait-increasing alleles will tend to have greater frequencies in the population with higher mean trait values, compared to the population with a lower trait mean. When examining populations with a range of different phenotypic trait means, we may therefore expect that the allele frequencies at the trait-affecting loci show a linear or at least steady relation with the observed trait means (Barton, 1999). I hypothesise here that this predicted relation can be exploited to distinguish potentially causal loci (and the linked variation) from loci not associated with the focal trait. In case of a successful evaluation, the major advantages of the proposed approach would be the reduced sequencing effort by the possibility to use pooled population samples (PoolSeq) and the opportunity to use bulk

phenotyping (e.g. by satellite imaging, flow-cytometry, etc.) on traits for which individual
phenotyping is difficult or tedious.

The most important assumption for the approach is obviously that observed population
differences in the focal trait means have at least partially a genetic basis. Since the
environment has usually an effect on the phenotype (Sella & Barton, 2019), total phenotypic
variance should be adjusted for known fixed environmental effects, because this increases
the fraction of variance due to genetic factors (Visscher et al. 2008). Predicting additive
genetic values with even higher accuracy can be achieved by taking into account GxE
interactions through repeated phenotypic measurements of the same individuals under
different environmental conditions, e.g. by time series (Visscher et al. 2008). I assumed
therefore that environmental influence on the phenotypic trait variance among populations
has been statistically removed as much as possible. Similarly important is the assumption
that the genetic variance of the focal quantitative trait can be adequately described by an
additive model. Both empirical and theoretical evidence suggests that this is indeed the case
for most complex traits (Hill et al., 2008). Even though epistatic interactions are wide spread
(Mackay, 2014), Sella and Barton (Sella & Barton, 2019) argue that the marginal allelic effects
on quantitative traits are well approximated by a simple additive model.

The aims of this study were i) to understand whether and under which circumstances
the hypothesised pattern of a linear relation between the population allele frequencies at
causal loci and the phenotypic population means of the respective trait emerges, ii) to
evaluate the influence of population genetic parameters of typical natural systems and the
experimental design on the likelihood of identifying causal loci underlying an additive
quantitative trait, in particular to elucidate the limits of the approach with regard to genetic
architecture and population structure, iii) to explore the possibilities for statistical genomic
prediction of phenotypic population means from the allele frequencies at the identified loci,
and iv) to evaluate the statistical power of the method for a realistic range of effective
genome sizes. I used individual-based population genomic forward simulations and
machine learning approaches (minimum entropy feature selection) for prediction and
utilised an information theory-based framework for evaluation of the proposed method.

Material and methods

Expectation of a positive correlation between quantitative trait loci allele frequencies and
phenotypic population means.

Consider a biallelic, codominant system for the additively heritable component of a
quantitative trait with n loci contributing to the trait. In this system, all loci contribute
equally to the phenotypic trait, with one allele per locus making a greater contribution than
the other. The phenotypic trait value x of an individual can then be determined by simply
adding up the number of trait-increasing alleles (g with values of 0, 1 or 2) over all n
quantitative trait loci (QTL) and multiplying this sum with a scaling constant k:

(1) $x = k \times (g_1 + g_2 + \cdots + g_n)$

When adding more individuals, the phenotypic population trait mean is defined as the
mean of the row sums:

(2)

$$\begin{array}{cccccc} & QTL_1 & \cdots & \cdots & QTL_n & \textit{individual trait value} \\ Ind_1 & g_{11} & g_{12} & \cdots & g_n & x_1 = k \times \sum_{l=1}^{n} g_{1l} \\ Ind_2 & g_{21} & \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ Ind_m & g_{m1} & \cdots & \cdots & g_{mn} & x_m = k \times \sum_{l=1}^{n} g_{1l} \end{array}$$

$$\text{population trait mean}$$
$$\bar{x} = \frac{\sum_{i=1}^{m} x_i}{m}$$

124   The columns of this matrix can be used to calculate the population allele frequency (AF)
125   of the trait increasing allele for each QTL.

126   (3)

$$\begin{array}{cccccccc} & QTL_1 & \cdots & \cdots & QTL_n & & & \textit{individual trait value} \\ Ind_1 & g_{11} & g_{12} & \cdots & g_n & & & x_1 = k \times \sum_{l=1}^{n} g_{1l} \\ Ind_2 & g_{21} & \cdots & \cdots & \cdots & & & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & & & \cdots \\ Ind_m & g_{m1} & \cdots & \cdots & g_{mn} & & & x_m = k \times \sum_{l=1}^{n} g_{1l} \end{array}$$

$$\begin{array}{cccccccc} & & & & & \textit{mean QTL AF} & & \textit{population trait mean} \\ AF & AF_1 = \frac{\sum_{i=1}^{m} g_i}{2m} & \cdots & \cdots & AF_1 = \frac{\sum_{i=1}^{m} g_i}{2m} & \overline{AF} = \frac{\sum_{i=1}^{n} AF_i}{n} & \sim & \bar{x} = \frac{\sum_{i=1}^{m} x_i}{m} \end{array}$$

The mean population allele frequency at QTL loci is thus directly proportional to the
phenotypic population trait mean. This relationship remains unchanged even if the
individual locus contributions are not identical, with some loci contributing more or less to
the phenotypic trait value. In this case, a scaling vector is required to weigh the individual
locus contributions to individual trait values, and those of the AFs to the population trait
mean. Since the AFs are by definition bounded by zero and one, the population trait mean is
minimal when the allele frequencies of the trait-increasing allele at all QTL are zero and
maximal when all QTL AFs are one. This proportionality links the individual genotypes and
the AFs at the QTL linearly with the population trait mean.
If we extend this to a set of populations and order them with decreasing phenotypic
population means, we can be sure that the mean QTL AFs of the populations will also be
ordered in decreasing sequence:

(4)

$$\begin{array}{ccccc} & Pop_1 & Pop_2 & \cdots & Pop_n \\ \text{population trait mean} & \bar{x_1} > & \bar{x_2} > & \cdots & \bar{x_n} \\ & \sim & \sim & \cdots & \sim \\ \textit{mean QTL AF} & \overline{AF_1} > & \overline{AF_2} > & \cdots & \overline{AF_n} \\ QTL_1 & AF_{11} & AF_{21} & \cdots & AF_{11} \\ QTL_2 & AF_{12} & AF_{22} & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ QTL_m & AF_{1m} & \cdots & \cdots & AF_{nm} \end{array}$$

The answer to whether the allele frequencies in every row i.e. at every contributing
locus can be used to predict the population trait mean depends on whether the expected
covariance between these two vectors is positive:

(5) $E[\text{cov}(QTL_m, \text{population trait mean})] > 0$

where

(6) $\text{cov}[\text{QTL}_m, \text{population trait mean}] = \text{E}[\sum_{i=1}^{n}(\text{AF}_{im} - \text{E}[\text{AF}_m]) * (\bar{x}_l - \text{E}[\bar{X}])]$

with $\bar{X}$ representing the grand mean over all populations. As all elements in the QTL
matrix are positive, they inherently tend to contribute positively to their column means.
Therefore, AF larger than the overall AF mean at his locus tend to be on the left side of the
population closest to the overall phenotypic mean in the ordered matrix above. Conversely,
AF smaller than the locus AF mean are rather on the right. This leads intuitively to a
positive expected covariance between each row and the column mean, in particular if the
number of populations becomes large. Conversely, the AF at (unlinked) loci not
contributing to the phenotypic population trait mean have an expectation of zero.
I tested these general expectations and the effect of different scaling vectors for the
effect size distribution of QTL with a first set of simulations. I generated a matrix of size n x
m populated with random AF between zero and 1. To avoid stochastic effects due to sample
size, the number of populations n was fixed at 10,000. The number of QTL m was varied
from oligogenic to highly polygenic (10, 20, 50, 100, 200, 500, 1000, 2000, 5000). Three
different distributions of loci effects were tested, i) a flat distribution with all loci
contributing equally, ii) a mildly decreasing exponential function and iii) a steeply
decreasing exponential function with few loci contributing much and many very little
(Supplemental Figure 1).
Each of the m columns was used to calculate the phenotypic population mean of the
respective population by adding up the AF multiplied with the respective locus weight. The
resulting n phenotypic population means were then correlated to the n AF of each of the m
loci and the resulting m Pearson correlation coefficients (i.e. the standardized covariance)
recorded. From these, mean and standard deviation were calculated and tested, whether
they conform to a normal distribution (scipy.stats.normaltest). Furthermore, a second
matrix of identical size was populated with random AF, and the correlation of these non-
contributing loci to the population means derived from the QTL matrix was computed. The
simulations were repeated 10 times in every possible parameter combination and the
results averaged (Supplemental Script 1).

Individual based Wright-Fisher forward model

A Wright-Fisher individual forward genetic simulation model was used to investigate the
potential of a genome-wide association study based on the means of a population trait and
population allele frequency data. In the simulation, all loci were assumed to be unlinked,
thus representing haplotypes in LD rather than single SNPs. (Visscher et al., 2017). For each
simulation run, the initial allele frequencies for all loci in the total population were
randomly drawn from a range of 0.1 to 0.9. To generate a hermaphroditic and diploid
individual, two alleles were randomly drawn with a probability based on their frequency at
the respective locus, and the resulting genotype at this locus was recorded. This process
was repeated for all loci. As a result, each individual was represented by a vector of biallelic
genotypes ($AA = 0$, $Aa;aA = 1$, $aa = 2$). To model a quantitative, fully additive trait, a
variable number of loci were assigned as quantitative trait loci (QTL). In addition, a much
larger number of neutral loci was modelled.

Genetic architecture of the quantitative trait

The continuous trait value was measured in arbitrary units. The allele (A) at each QTL
had no effect on the individual's trait value, resulting in a completely homozygous A

individual at the QTL having a phenotypic trait value of 0. The alternative allele (*a*) added a
locus-specific value to the trait. Two distributional extremes have been considered for the
allelic effects on the trait value: i) a uniform distribution where each locus contributes 0.5
units to the trait. An individual that is completely homozygous for the alternative allele *a*
therefore had a trait value equal to the number of QTL, ii) an exponential distribution with
few loci having large effects and many having very small effects, scaled such that the
maximum possible trait value was also equal to the number of QTL (see Supplementary
Figure 2A). To model the effect of phenotyping errors, unaccounted environmental
influence (*i.e.* phenotypic plasticity), and/or the unspecific contribution of the genomic
background to the trait, a random value drawn from a Gaussian distribution with a mean of
zero and selectable standard deviation between 0.1 and 3 could be added to the genetically
determined phenotype value of each individual. The phenotypic value of each individual's
trait was determined by summing the allelic effects of all genotypes at all QTL loci plus the
random value and the result recorded.

Reproduction and selection

Subpopulations in each run were created from the same initially drawn random allele
frequency array, mimicking a common descent. Due to sampling variance, the realised allele
frequencies and thus the mean subpopulation trait value differed from the initial
frequencies of the total (ancestral) population. A subpopulation always comprised 500
adult individuals.
Each subpopulation was reproduced at least once to obtain a genotype distribution in
Hardy-Weinberg equilibrium. For reproduction, two random individuals were chosen with
replacement from the adult population. The genotype of an offspring individual at a locus
was determined by randomly choosing one of the two alleles from each designated parent
at this locus. Each parent fostered n_juv offspring; therefore, 2 x n_juv were produced in
each mating. After reproduction, the parental generation was discarded to prevent
overlapping generations. Each generation had N/2 matings, resulting in an offspring
population of N*n_juv individuals.
Because the offspring population was much larger than the the size of the adult
population, it was necessary to reduce it. This was achieved by a combination of 'hard'
natural selection and random mortality. An individual's survival to the adult stage was
determined by the absolute deviation of its phenotypic trait value from a pre-specified
selective trait optimum for the respective subpopulation. This selective trait optimum for a
subpopulation was determined by adding a random value taken from a Gaussian
distribution with a mean of zero and a standard deviation of 2.5 to the initial population
mean. An individual's survival probability was determined by an exponential decline
function with strength *s* (the exponent of the function, see Supplementary Figure 2B).
Individuals were randomly selected one by one from the offspring population and their
survival probability calculated. A respectively biased coin was then tossed to determine
their fate. This process was repeated until the adult population size was reached, and any
remaining offspring individuals were indiscriminately discarded. If the phenotypic mean of
the subpopulation was close to or at the selective optimum (see below), this process
resulted in stabilising selection. If the population was away from the optimum, rapid
directed selection towards the optimum was observed, depending on the strength of
selection. For the assessment of the effect of population structure, the subpopulations could
evolve in complete isolation from each other for a predetermined number of generations

(2-50). This introduced random genetic drift among the populations at both QTL and
neutral loci. Both drift and selection towards different trait optima led to variation in
population trait means among the subpopulations.

Simulation scenarios

I considered scenarios were subpopulations with quantitative phenotypic population
differences in mean for the trait in question were screened from a larger total population.
Although the population trait mean differences in the simulation of this scenario were
created by drift and local adaptation, any other source of heritable phenotypic population
differentiation, such as maladaptation, introgression, or e.g. in the case of managed species,
human choice, may also be the reason for differentiation in population means. The range of
phenotypic variation among the subpopulations was not predetermined, but an emergend
feature of the simulation parameters.
After evolving the subpopulations for the desired number of generations, phenotypic
trait means, and genome-wide allele frequencies were recorded. While the phenotypic
means for each subpopulation was calculated over all individuals, the allele frequencies
were estimated in a PoolSeq (Kofler et al., 2011) like fashion from subsamples of 50
individuals. The range of phenotypic trait means of the population sample was recorded.
Trait heritability was determined in the last generation by regressing the phenotypic values
of the offspring against the mean of their respective parents (Lynch & Walsh, 1998). As
measure for population subdivision due to drift, F_{ST} among all subpopulations was
calculated from the variance of the true allele frequencies (Wright, 1949).

Population GWAS

Assuming a linear relation between the phenotypic (sub)population means, and the
population allele frequencies of the causal loci on the other, I calculated an ordinary linear
regression between these two variables for all loci in the genome. I used the resulting -$\log_{10}$
p value as measure of regression fit and effect size. I recorded the number of true positive
loci (TPL) among the loci beyond a predefined outlier threshold. As GWAS performance
measures, the true positive rate (TPR = recall, sensitivity, discovered proportion of all
QTL), positive predictive value (PPV = precision, proportion of TPL among outliers
considered) and false discovery rate (FDR = proportion of false positive loci among outliers
considered, type I error) were calculated.

Influence of natural system and experimental design factors

In a first set of simulations, I explored the influence of factors inherent to the natural
system and the experimental design on population GWAS performance. As factors of the
natural system, I assumed characteristics that are beyond control of the researcher, such as
heritability of the trait and its genetic architecture (number of QTL, distribution of allele
trait contribution). While the degree of population differentiation and range of phenotypic
differentiation are also inherent to the organism studied, the choice of samples may allow a
certain control over these parameters. The number of subpopulations screened is clearly a
study design decision (Table 1).

Table 1. Simulation parameters, their abbreviations, values used in simulations, their
biological meaning and whether the parameter is a feature of the natural system
under scrutiny or under the control of the researcher.

Parameter	Abbreviation	Values in the simulation	Biological meaning	Degree of knowledge in natural systems/under the control of study design
Number of subpopulations scored for phenotypic population means and genome wide allele frequencies	n_pop	12, 24 36, 48, 60	-	Full control
Number of quantitative trait loci contributing to the focal trait	n_qtl	30, 50, 70, 110, 500	Genetic architecture of the trait	*A priori* unknown
Distribution of allelic effects on the focal trait	allelic_contr	Flat, exponential	Genetic architecture of the trait	*A priori* unknown
Standard deviation of random phenotypic variation added to individuals	pheno_plast	0.1, 1, 2, 3	Heritability of the trait	*A priori* unknown
Number of generations of independent evolution of the subpopulations	gen	2, 5, 10, 30, 50	Population structure	Partial control

A genetic trait architecture of 30, 50, 70, 110 and 500 loci, flat and exponential allelic
effect distributions, as well as phenotypic plasticity coefficients of 0.1, 1, 2 and 3 were
applied. Selection strength was fixed at 0.5 (Suppl. Fig. 1B). Simulations were run for 2, 5,
10, 30 and 50 generations among 12, 24, 36, 48 and 60 subpopulations of 500 individuals
each. For this set of simulations, 1000 neutral loci and a fixed outlier threshold (upper 5%
quantile, either 21 or 22 outlier loci, respectively) were applied. Each possible parameter
combination was run in five replicates, resulting in 5000 simulation runs.

The effect of each parameter on PPV was assessed with ANOVA over all simulations,
grouped after the respective parameter classes. The relative influence of the number of
populations, QTL loci, distribution of allelic contributions, trait heritability, phenotypic
range and population subdivision on the proportion of TPL among the outlier loci was
determined with a General Linearized Model (GLM).

Genomic prediction and validation

The loci identified by GWAS were used to devise a statistical genomic prediction model
to obtain a score that uses observed allele frequencies at the identified loci to predict the
mean population phenotype of unmeasured populations. To remove remaining
uninformative or redundant loci, I applied feature selection, which is particularly suitable
for bioinformatic data sets that contain many features but comparatively few data points.
The minimum entropy feature selection (MEFS) technique uses mutual information to
measure the dependence between each feature and the target variable. For a given number
of features (k), the data set of the allele frequencies at selected outlier loci and the
respective phenotypic population means was repeatedly randomly divided in training
(80%) and test set (20%), a multiple regression model fitted and the r^2-fit of the test sets to
the predicted phenotypes recorded. The best model for the current k was recorded and the
process repeated for all k in a range between 2 and the number of selected loci – 1. Finally,
the best model (i.e. highest r^2) among all k was chosen as best prediction model. MEFS was
implemented with the Python module scikit-learn 1.3.2 (Pedregosa et al., 2011)

The performance of the selected best prediction model for each run was tested with
independent data. Ten additional populations were created under the same parameters as
the initial set of populations and their mean population phenotypes calculated as described
above. Then the allele frequencies at the predictive loci as identified by the best prediction
model were extracted and phenotypic prediction scores according to the best prediction
model calculated. The performance of the statistical genomic prediction was then evaluated

by calculating the Pearson correlation coefficient r between the observed mean population
phenotypes and the phenotypic prediction scores for the ten validation populations
(Supplemental Script 2).

Method performance with realistic genome sizes

Whether and which proportion of TPL, i.e. causal loci can be expected to be reliably
identified with the proposed method depends crucially on the total number of loci screened
as this number determines the length and size of the distributional tail of random
associations of neutral loci with the mean population phenotypes. The number of effectively
independently evolving loci in a population depends on genome size, effective population
size (including all factors that affect it locally and globally) and LD structure (Chakraborty,
1981; Taylor & Higgs, 2000). There are hardly any empirical estimates in the literature, but
dividing typical genome sizes by typical mean genome-wide LD ranges suggested that a few
tens of thousands to a few hundreds of thousands of independent loci per genome is a
realistic range for a large number of taxa (see Supplemental Table 1). I have therefore
considered 1,000, 5,000, 10,000, 30,000, 50,000 and 100,000 independent neutral loci for
samples of 12, 24, 36, 48 and 60 populations with a restricted set of parameters (number of
QTL and allelic contribution). As the true number of QTL underlying a trait is rarely *a priori*
known, I considered 10, 30, 50, 70 and 110 QTL loci in this analysis. I therefore recorded
the number of TPL found in sets of loci with the absolutely highest 10, 30, 50 and 100 -
$\log_{10}p$ values, as well as outlier proportions of 0.0001, 0.001, 0.01, 0.02, 0.05 and 0.1 of the
total number of loci in the respective simulation. As above, all simulations were run in all
possible parameter combinations with five replicates each (Supplemental Script 3).
I analysed the performance of the method in an Area Under the Curve – Receiver
Operator Curve (AUC-ROC) and – Precision, Recall (AUC-PR) framework as suggested by
Lotterhos et al. (Lotterhos et al., 2022). For each combination of effective genome size and
number of population scored, mean TPR, PPV and FDR were calculated over all replicates
and parameter combinations for the respective set of simulations. The maximum F1 score
(Rijsbergen, 1979) was used in addition to identify the optimal number of outliers to select.
All simulations were implemented in Python 3.11.7 (Van Rossum & Drake, 2009) and run
under pypy 3.10 (Team, 2019), the respective scripts can be found in the Supplementary
Material (Scripts 1-3). General statistical tests were performed with R (R Core Team, 2013).

342   Results

343   Allele frequencies at QTL loci co-vary positively with the population trait mean

344   The mean correlation coefficient between all QTL and the respective phenotypic
345   population means was positive in every parameter combination and in every single
346   simulation (Table 2).

347   Table 2. Expected mean Pearson's correlation coefficients between QTL AF and
348   phenotypic population means for three different locus contribution distributions and
349   varying number of QTL.

n_QTL	Flat	Mildly exponential	Strongly exponential
1	1	1	1
10	0.322	0.307	0.165
20	0.227	0.218	0.085
50	0.142	0.116	0.035
100	0.101	0.064	0.017

200	0.070	0.033	0.008
500	0.045	0.013	0.003
1000	0.031	0.006	0.001
2000	0.023	0.003	0.001
5000	0.014	0.001	0.000

The expected mean correlation coefficient decreased with increasing number of
contributing QTL (Figure 1). This decay was best described by a negative exponential
function of the form *number of QTL$^{-1/x}$* with x ranging from 1.26 in case of the strongly
unbalanced locus contributions to 2 for the flat distribution.

Figure 1. Plot of the expected correlation coefficient between individual QTL loci and
the population trait mean in dependence of the number of QTL loci.

The distribution of the correlation coefficients did not deviate from a normal
distribution for the flat locus contribution distribution, while it did for all other parameters.
The correlation coefficients for the non-contributing loci had an expectation of zero and a
mean standard deviation of 0.01, regardless of the number of loci.

Influence of simulation parameters on parameters of the simulated populations

In the second set of simulations with 1000 neutral loci and an outlier threshold of the
most extreme 5% -$\log_{10}$p values, the permutation of all parameters with five replicates
yielded 4984 completed independent simulation runs. The 16 missing to the expected 5000
runs were due to one or more subpopulations going extinct during the simulation. In total,
more than 30 billion individuals were simulated.
The number of independently evolving generations strongly influenced the population
structure ($r^2 = 0.996$). F_{ST} estimates increased on average by 0.0018 per additional
generation, with large variation. Resulting F_{ST} values ranged between 0.012 and 0.111
(Supplemental Figure 3A). Heritability of the trait depended strongly on the plasticity
parameter ($r^2 = 0.942$, Supplemental Figure 3B). It decreased on average by 0.24 per unit
standard deviation, with variations of up to 0.05 even among runs with identical
parameters. Trait heritability estimates ranged from 0.16 to 1.02.

Factors influencing the proportion of detected true positive loci among outliers

Over all simulations in the first set with 1000 neutral loci, on average about 14.1 (mean
proportion 0.62) true positive loci (TPL) were among the highest 5% outliers. The TPL
values ranged between none (0) and 23 (1.0); the 25 percentile was 10 (0.38), the 75
percentile 20 (0.90). This exceeded in >95% of cases random expectations, when excluding
the highly polygenic case (n_qtl = 500), this proportion rose to more than 99%.

Figure 2. Effect of simulation parameters and emergent features on the proportion of
identified true positive loci. A) Phenotypical plasticity parameter as a proxy for
heritability. B) Number of QTL. C) Distribution of allelic contributions to phenotypic
trait. D) Generations of independent evolution as proxy for population structure. E)
Number of populations scored for population phenotypic means and allele
frequencies. F) Range of population phenotypic means as an emergent feature.

The phenotypic plasticity parameter had a significant effect on PPV (F = 8.37, p = 1.51 x
10^{-5}), however, as the data plot already indicated (Figure 2A), this was due to the drop of
the mean in the class with the lowest heritability only (from about 0.64 in the other classes
to 0.59), as indicated by Dunn's post-hoc test (z statistic > 2.8 and p below 4.8 x 10^{-4} in all
comparisons with class 3). The number of QTL showed a systematic effect on mean PPV
(ANOVA F = 226, p = 6.42 x 10^{-177}, Figure 2B). This was mainly due to the highly polygenic
class; while the mean PPV for all different QTL numbers up to 110 was above 62.5%, it was
as low as 39.8% for 500 QTL (z statistic > 17 and p below 2.1 x 10^{-69} in all comparisons).
The distribution of allelic effects on the trait showed a moderate but highly significant effect
on the mean PPV (mean equal contribution = 0.67, mean exponential = 0.57, F = 154,5 p =
6.58 x 10^{-35}, Figure 2C). The relation of mean proportion of detected TPL and population
structure was non-linear. Both very weak (2 generations) and strong population (30+
generations) structure led to a relatively lower proportion of TPL (0.62 and 0.46,
respectively, Figure 2D), while for intermediate values TPL proportions of 0.72 (5
generations) and 0.70 (10 generations) were observed. The by far strongest effect on
proportion of TPL among the selected loci had the number of populations screened (F =
514, p = 0). The values ranged from a mean PPV of 0.35 (s.d. = 0.18) with 12 populations to
over 0.78 (s.d. = 0.23) with 60 populations. Given the chosen threshold, a diminishing
return was observed above 36 populations sampled (Figure 2E). The phenotypic range in a
simulation run had a moderate (r = 0.42, p = 5.36 x 10^{-203}), yet significantly positive effect
on detection of TPL. The realised range of population trait means in the simulations covered
on average 15.4% (range = 0.1-56%) of the possible range. An increase of one unit in range
increased the proportion of TPL by 0.07 (Figure 2F).
When jointly considering the effect of all parameters on PPV in a GLM, it turned out that
all had a significant effect (Table 2). Their relative influence increased from F_{ST} (r^2 = 0.008)
over distribution of allelic trait contribution (r^2 = 0.013), heritability (r^2 = 0.024), the
number of populations (r^2 = 0.100), phenotypic range (r^2 = 0.130) to the number of QTLs,
that had by far the greatest influence (r^2 = 0.343). In total, the parameters explained 61.8%
of variance.

Table 2. Generalised Linear Model of factors influencing the proportion of TPL among
outliers (PPV) in simulations.

Factor	Coefficient	Std.err.	t	p	r^2
Constant	0.243	0.022	10,859	3.86E-23	
F_{ST}	-2.954	0.150	-19,749	2.21E-79	0.008
allelic_contr	0.153	0.009	16,122	6.46E-53	0.013
heritability	0.042	0.019	21,964	2.81E-02	0.024
n_pop	0.006	0.000	22,089	8.01E-99	0.100
range_pheno	0.037	0.002	19,492	2.33E-77	0.130
n_qtl	-0.001	0.304	-46,314	0.00E+00	0.343

Minimum Entropy Feature Selection and statistical phenotype prediction

Minimum Entropy Feature Selection (MEFS) removed on average 8.72 (range = 2-14,
s.d. = 4.14) loci, corresponding to a proportion of 0.38 (s.d. = 0.19) from the statistically
chosen initial outlier set. The procedure removed on average a larger proportion of FP than
TPL (mean difference 0.14, t = -19.9, p = 6.7 x 10^{-79}). This increased the proportion of TPL

in the final prediction set on average by 0.05 (range = -0.23-0.48, s.d. = 0.09) to a mean of
0.66 (range = 0-1, s.d. = 0.29, Figure 2).

Figure 3. Effect of Minimum Entropy Feature Selection (MEFS) on the proportion of
TPL and FP in the selected set.

The predictive accuracy of the SNP loci sets selected by MEFS was on average r = 0.58
(s.d. = 0.44). It ranged from -0.95 to 1.0. The distribution was highly skewed with 75%
being higher than 0.30, the median was found at 0.76 and still 25% being higher than 0.94
(Supplemental Figure 3).
The accuracy of mean population phenotype prediction depended linearly on the
number of TPL in the prediction set (r^2 = 0.28, p = 0), with any additional TPL increasing
the correlation coefficient by 0.05 (Supplemental Figure 4A). Inversely, the accuracy of
prediction decreased with a rising number of FP, but even with a considerable number of
FP in the prediction set, accurate prediction was possible in a large number of cases
(Supplemental Figure 4B). Overall, the prediction accuracy increased with increasing
proportions of TPL among the prediction set, although even 100% TPL in the prediction set
did not guarantee a highly accurate prediction (r > 0.8) in all cases (Supplemental Figure
4C).

Figure 4. Influence of simulation parameters on the accuracy of statistical population
mean phenotype prediction. A) Phenotypic plasticity parameter as proxy for
heritability. B) Distribution of allelic trait contributions. C) Number of trait-
underlying QTLs. D) Generation of independent evolution as proxy for population
structure. E) Number of populations scored. F) Proportion of TPL in the prediction
loci set after MEFS.

As the prediction accuracy depended on the proportion of selected TPL, their relation to the individual simulation parameters was very similar to the results described in the previous section (Figure 4A-F). The number of populations screened was the most important factor. With 36 or more populations screened, 97.8% of simulations showed a prediction accuracy of 0.8 or better, independent of the other simulation parameters applied. In a GLM with all factors simultaneously considered, the proportion of TPL selected had the largest influence on prediction accuracy ($r^2 = 0.48$), followed by the number of QTL (0.34), the range of mean phenotypes (0.13), the number of populations screened (0.10). Heritability, distribution of allelic contributions and F_{ST} had only a minor influence on the prediction accuracy ($<= 0.02$, Table 3).

460
461 Table 3. Generalised Linear Model of factors influencing the accuracy of statistical phenotypic population mean prediction.

Factor	Coefficient	Std.err.	t	p	r^2
Constant	-0.0304	0.02	-14183	0.1562	
F_{ST}	-0.1981	0.16	-12778	0.2014	0.01
allelic_contr	0.1611	0.01	19	0.0000	0.01
heritability	0.0216	0.02	1265	0.2059	0.02
n_pop	0.0014	0.00	45782	0.0482	0.10
range_pheno	0.0104	0.00	56485	0.0002	0.13
n_qtl	-0.0009	0.03	-30692	0.0000	0.34
prop_TPL_FS	0.7761	0.02	34676	0.0000	0.48

Method performance with realistic effective genome sizes

The values for AUC-ROC ranged between 0.067 and 0.833, for AUC-PR between 0.013
and 0.730. There was an interaction between the effective genome size and number of
populations scored. According to both AUC measures, the method performed best, when the
number of populations scored was high and the genome small (Figure 5). An at least
satisfactory (> 0.66 for AUC-ROC and > 0.53 for AUC-PR) overall performance was
observed for 24 populations for the smallest genomes considered (1,000), for 36
populations up to 30,000 independent loci and for genome sizes up to 100,000 for 48 and
60. The similar values in both statistics and the plots suggested that there are diminishing
returns for samples larger that about 48 populations. Moreover, closer inspection of the
corresponding plots (Figure 6) suggested that for samples of 48 and 60 populations, an
optimal ratio between TPL and FPL exists for approximately the 25 highest outlier loci,
independent of genome size. For combinations with good performance, the maximum F1
score suggested that choosing the 30 highest outlier provided the optimal compromise
between maximising TPR and minimising FPR (Supplemental Figure 6).

Figure 5. Heat-map of AUC-ROC (area under the curve – receiver operator
characteristics) and AUC-PR (area under the curve – precision recall) in relation to
effective genome size and number of populations scored.

Figure 6. AUC-ROC and AUC-PR for a range of effective genome sizes. In the left
column are the plots of AUC-ROC, i.e. FDR on the x-axis versus TPR on the y-axis. The
right column shows AUC-PR plots, i.e. TPR on the x-axis versus PPV on the y-axis. The
dotted lines indicate the threshold for a random effect.

Discussion

This study used extensive forward simulations to explore the potential of a novel GWAS
approach utilising phenotypic population means and genome-wide allele-frequency data to
identify loci potentially underlying quantitative polygenic traits. While the approach seems
to be generally useful in a wide range of cases, there are also clear limits to its applicability.

General validity of the underlying assumptions

The initial simulations demonstrated that the expectation for the covariance of random
population "allele frequencies" at contributing quantitative trait loci (QTL) and the
respective population trait mean is consistently positive when an additive model applies.
This is an inherent consequence of the common dependence of both variables on the QTL
genotypes of the individuals in a population, as demonstrated in (3). Additive models seems
to be an appropriate statistical approximation for most quantitative traits at population
level (Hill et al., 2008), despite the description of many epistatic interaction on the molecular
level (Moore & Williams, 2005).

The relation appeared to be largely independent of the distribution shape of locus
contributions to the trait. While in the case of equal contributions i.e. a flat distribution, the
correlation coefficients of individual loci are themselves a random variate, normally
distributed around the expected mean. As the distribution becomes increasingly skewed,
locus contribution becomes predictive of the correlation to the trait. Loci contributing more
to the trait and thus accounting for more of the phenotypic variance will likely have a higher
correlation of their allele frequencies to the population mean. Conversely, the expectation
for non-contributing loci is zero. Therefore, it is principally possible to exploit the
correlation between allele frequencies and population trait means for the identification of
loci underlying an additive quantitative trait. However, some statistical limitations became
obvious. Firstly, as the number of QTL increases, the expected mean correlation coefficients
become so small that they are likely to be indistinguishable from the tail of the zero-
centered normal distribution of non-contributing loci, even with an unrealistically high
number of samples. Consequently, the method for identifying QTL by the positive
covariance of their allele frequencies with the population trait means is *a priori* more suited
for oligogenic to moderately polygenic traits. Secondly, the number of QTL and the
distribution of locus contributions may influence the statistical identifiability of individual
QTL. In particular, loci that contribute only minimally to the trait or that fall by chance
below the expected mean correlation coefficient may overlap with the tail of the
distribution of non-contributing loci.

These predictions remind of similar conditions for the contribution of different QTL
architectures to phenotypic adaptation described by (Höllinger et al., 2023). They assert that
phenotypic adaptation of oligogenic traits is achieved by detectable allele frequency shifts
at some but not very many loci, while adaptation in highly polygenic traits is rather
achieved by subtle perturbations of standing variation, with respective consequences for
their detectability. Just as expected here, they stress the importance of stochastic effects
that may lead to apparently heterogeneous locus contributions (Höllinger et al., 2023).

Limiting factors in natural settings

The Wright-Fisher forward simulations of a quantitative trait in a subdivided population
with realistic properties and sample sizes largely confirmed the theoretical expectations. In
particular when a sufficient number of populations was scored (>60), a large proportion of
true positive loci could be reliably identified, with the exception of a few parameter
combinations. The genetic architecture of the trait was an important predictor for the
ability to identify causal loci. The most important other factor was the genetic trait
architecture. While the loci underlying oligogenic and moderately polygenic traits could be
fairly reliably identified, the highly polygenic scenario tested (500 loci) performed poorly.
The difference between the two tested locus contribution distributions was not very

pronounced. This was likely due to the tendency of higher correlations between higher
contributing loci and the trait, which ensured the inclusion of a substantial proportion of
true positive loci in the selected outliers under a wide range of conditions.

The influence of mean heritability was similarly not marked. Even down to trait
heritability estimates of 0.3, the success rate was only slightly reduced. This effect may be
attributed to the averaging of phenotypes and genotypes across multiple individuals, which
is likely to mitigate the inherent noise associated with individual data (Johri et al., 2022;
Stinchcombe & Hoekstra, 2008). This finding is consistent with observations by (Zhang et al.,
2018), who employed pooled data for GWAS. From a practical standpoint, the findings
suggest that inevitable errors in phenotyping, which can compromise GWAS performance
on individuals (Barendse, 2011), are likely to be less problematic when using the mean
measured over many individuals. Furthermore, this finding indicates that the failure to
entirely remove non-additive variance from the analysis does not necessarily compromise
the method's ability to reliably identify trait-associated loci.

From a statistical perspective, it was anticipated that the range of phenotypic population
means would influence the identification of true positive loci to some extent, given that a
larger range of phenotypic means is inherently associated with on average larger allele-
frequency differences among populations. The choice of populations with a large range of
environmentally unexplained variance is therefore crucial. It is, however, important to
emphasise that the underlying causes of the observed differences in trait means among
populations are not of primary concern. These may be attributed to local adaptation, but
also to maladaptation, human choice, or other factors. Likewise, increasing the number of
populations screened increased the statistical power of the approach. However, it seemed
that increasing the number of samples led to diminishing returns in statistical power gain
beyond a certain threshold.

A pronounced population structure ($F_{ST} > \sim 0.07$) was a major factor impeding reliable
identification of true positive loci, even with a high number of samples. This is probably due
to distinct evolutionary trajectories in independently evolving populations. The genetic
redundancy of polygenic traits can lead to evolution of the same phenotypes from different
genomic bases(de Vladar & Barton, 2014; Kaneko & Furusawa, 2006), even if evolving from the
same ancestral population (Barghi et al., 2019, 2020; Pfenninger et al., 2015). If different
loci in different populations are causal for the observed phenotypic differences, a linear
relation between population means and allele frequencies is not to be expected. It is
therefore important that the allele frequencies in the studied populations are correlated
either by recent common descent and/or recurrent gene-flow, i.e. that the population
structure between the population scored is weak (Mathieson, 2021).

A situation where the overall genetic distance and the phenotypic differences are
correlated, e.g. if an environmental gradient is correlated to the geographic distance
between populations (IBD) and the trait value is an adaptation to this gradient, should as
well be prone to produce false positives. To avoid such a situation, it is recommended to
test for (the absence of) a correlation between genome-wide genetic distance and
differences in phenotypic means (e.g. by a Mantel's test).

Accurate statistical genomic prediction in a wide range of conditions

Genomic prediction is deemed to be one of the major tools for the mitigation of climate
change on biodiversity (Aguirre-Liguori et al., 2021; Bernatchez et al., 2023; Capblancq et al.,
2020; Waldvogel et al., 2020). Contrary to its application in medicine or selective breeding

(Wray et al., 2019), however, accurate prediction of population responses is probably more
important than the prediction of individual phenotypes. However, there is no theoretical
obstacle, why the identified loci could not be used for individual genomic phenotype
prediction, but this was not investigated here. Within the limits outlined above, the
proposed method delivered very accurate predictions ($r > 0.8$) of population mean
phenotypes. It should be noted, however, that the prediction is statistical in the sense that it
produces a prediction score (de Los Campos et al., 2018) that correlates with the mean
population phenotype and not the phenotype itself. Just like with any other genomic
prediction (Kachuri et al., 2024), this limits the transferability of the prediction to other,
more distantly related lineages or species.

Reducing the false positive rate is in any case advisable, as it proved to be the most
important factor of prediction success with independent data. The application of a Machine
learning approach, in this case Minimum Entropy Feature Selection (MEFS), prior to
prediction reduced the already low false positive rate among the initially selected loci
further. Other, comparable methods, such as e.g. , likely perform comparably or even better.
Other factors influenced prediction success in a very similar fashion as the true positive
rate. One notable exception was distribution of locus contributions. While true positive loci
were more reliably identified from a flat distribution, prediction worked better when many
loci of large effect were among the prediction set, most likely because these loci contribute
more to phenotypic variance (Jain & Stephan, 2015).

Typical genome sizes of real species are no obstacle

The perhaps most important challenge was showing that the proposed method has
enough statistical power to distinguish at least a part of the unknown, but likely relatively
small number of QTL reliably from the large number of non-contributing loci in real
genomes of real species. The evaluation of method performance with AUC-ROC and AUC-PR,
as recommended recently (Lotterhos et al., 2022), showed a satisfactory performance even
for genomes with moderately high effective sizes, provided a sufficiently high number of
populations is screened. In particular restricting the selection of potentially causal QTL on a
few dozen of the highest outliers promises to yield very low false positive rates. As shown
above, already a limited number of true positive loci may be sufficient for reliable genomic
prediction.

Practical considerations

The proposed method finds rather genomic regions or haplotypes associated to the trait
in question than directly causal SNPs. However, this is true for most GWAS methods (Wang
et al., 2010) and therefore fine-mapping and inference of causal processes remain to be
done (Wallace, 2021). In practice, this requires that regions with high SNP outlier density
need to be collapsed to haplotypes prior to further analysis. Knowledge on the local LD-
structure, mean haplotype length, respectively recombination landscape can aid haplotype
identification (Flister et al., 2013). Recently developed machine learning approaches makes
such information available for pooled data (Adrion et al., 2020).
The possibly largest advantage of the proposed method is its data efficiency, if pooled
sequencing is applied. Because the Pool-Seq approach (Schlötterer et al., 2014) yields highly
accurate estimates of genome-wide allele frequencies at SNP sites (Czech, Peng, Spence,
Lang, Bellagio, Hildebrandt, Fritschi, Schwab, Rowan, & consortium, 2022) the necessary

sequencing effort is marginal compared to individual based approaches (Ziyatdinov et al.,
2021). This makes GWAS studies accessible for the usual funding in the field of biodiversity.
Pooled sequencing for GWAS has been proposed (Yang et al., 2015) and applied (Giorello et
al., 2023; Kumar et al., 2022; Pfenninger et al., 2021) with extreme phenotypes. What is now
required is the application of the method to a real-world data set, a work which is in
progress.

Conclusion

634 This study demonstrated the potential of the proposed GWAS approach for biodiversity
635 genomics. By carefully considering the factors influencing its performance and addressing
636 the limitations, this method can be a valuable tool for identifying the genetic basis of
637 complex traits in natural populations.

Acknowledgements

639 The author wants to thank Bob O'Hara and Barbara Feldmeyer for comments on the
640 manuscript.

Funding

642 The author declares that he has received no specific funding for this study.

Conflict of interest disclosure

644 The author declares that he complies with the PCI rule of having no financial conflicts of
645 interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

647 Supplementary information including the Python code used for the simulations is
648 available at https://**10.5281/zenodo.11562472**

References

Adrion, J. R., Galloway, J. G., & Kern, A. D. (2020). Predicting the landscape of recombination
using deep learning. *Molecular Biology and Evolution*, *37*(6), 1790–1808.
Aguirre-Liguori, J. A., Ramírez-Barahona, S., & Gaut, B. S. (2021). The evolutionary genomics
of species' responses to climate change. *Nature Ecology & Evolution*, *5*(10), 1350–1360.
Barendse, W. (2011). The effect of measurement error of phenotypes on genome wide
association studies. *BMC Genomics*, *12*(1), 232. https://doi.org/10.1186/1471-2164-12-
232
Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: A unifying
framework to understand positive selection. *Nature Reviews Genetics*, *21*(12), 769–781.
Barghi, N., Tobler, R., Nolte, V., Jakšić, A. M., Mallard, F., Otte, K. A., Dolezal, M., Taus, T.,
Kofler, R., & Schlötterer, C. (2019). Genetic redundancy fuels polygenic adaptation in
Drosophila. *PLoS Biology*, *17*(2), e3000128.
Barton, N. H. (1999). Clines in polygenic traits. *Genetics Research*, *74*(3), 223–236.

Bernatchez, L., Ferchaud, A.-L., Berger, C. S., Venney, C. J., & Xuereb, A. (2023). Genomics for
monitoring and understanding species responses to global climate change. *Nature
Reviews Genetics*, 1–19.
Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An expanded view of complex traits: From
polygenic to omnigenic. *Cell*, *169*(7), 1177–1186.
Brandes, N., Weissbrod, O., & Linial, M. (2022). Open problems in human trait genetics.
Genome Biology, *23*(1), 131. https://doi.org/10.1186/s13059-022-02697-9
Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., & Keller, S. R. (2020).
Genomic prediction of (mal) adaptation across current and future climatic landscapes.
Annual Review of Ecology, Evolution, and Systematics, *51*, 245–269.
Chakraborty, R. (1981). The distribution of the number of heterozygous loci in an individual
in natural populations. *Genetics*, *98*(2), 461.
Czech, L., Peng, Y., Spence, J., Lang, P., Bellagio, T., Hildebrandt, J., Fritschi, K., Schwab, R.,
Rowan, B., & Weigel, D. (2022). Efficient analysis of allele frequency variation from
whole-genome pool-sequencing data. *Population, Evolutionary, and Quantitative
Genetics Conference (PEQG 2022)*, 99.
https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item_3474009
Czech, L., Peng, Y., Spence, J. P., Lang, P. L., Bellagio, T., Hildebrandt, J., Fritschi, K., Schwab, R.,
Rowan, B. A., & consortium, G. (2022). Monitoring rapid evolution of plant populations at
scale with Pool-Sequencing. *BioRxiv*, 2022–02.
de Los Campos, G., Vazquez, A. I., Hsu, S., & Lello, L. (2018). Complex-trait prediction in the
era of big data. *Trends in Genetics*, *34*(10), 746–754.
de Vladar, H. P., & Barton, N. (2014). Stability and response of polygenic traits to stabilizing
selection and mutation. *Genetics*, *197*(2), 749–767.
Dunker, S., Boyd, M., Durka, W., Erler, S., Harpole, W. S., Henning, S., Herzschuh, U., Hornick,
T., Knight, T., Lips, S., Mäder, P., Švara, E. M., Mozarowski, S., Rakosy, D., Römermann, C.,
Schmitt-Jansen, M., Stoof-Leichsenring, K., Stratmann, F., Treudler, R., … Wilhelm, C.
(2022). The potential of multispectral imaging flow cytometry for environmental
monitoring. *Cytometry Part A*, *101*(9), 782–799. https://doi.org/10.1002/cyto.a.24658
Exposito-Alonso, M., Drost, H., Burbano, H. A., & Weigel, D. (2020). The Earth BioGenome
project: Opportunities and challenges for plant genomics and conservation. *The Plant
Journal*, *102*(2), 222–229. https://doi.org/10.1111/tpj.14631
Flister, M. J., Tsaih, S.-W., O'Meara, C. C., Endres, B., Hoffman, M. J., Geurts, A. M., Dwinell, M.
R., Lazar, J., Jacob, H. J., & Moreno, C. (2013). Identifying multiple causative genes at a
single GWAS locus. *Genome Research*, *23*(12), 1996–2002.
Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C.,
Crottini, A., Godoy, J. A., & Höglund, J. (2022). The era of reference genomes in
conservation genomics. *Trends in Ecology & Evolution*, *37*(3), 197–202.
Giorello, F. M., Farias, J., Basile, P., Balmelli, G., & Da Silva, C. C. (2023). Evaluating the
potential of XP-GWAS in Eucalyptus: Leaf heteroblasty as a case study. *Plant Gene*, *36*,
100430.
Heuertz, M., Carvalho, S. B., Galindo, J., Rinkevich, B., Robakowski, P., Aavik, T., Altinok, I.,
Barth, J. M., Cotrim, H., & Goessen, R. (2023). The application gap: Genomics for
biodiversity and ecosystem service management. *Biological Conservation*, *278*, 109883.
Hill, W. G., Goddard, M. E., & Visscher, P. M. (2008). Data and theory point to mainly additive
genetic variance for complex traits. *PLoS Genetics*, *4*(2), e1000008.
Hogg, C. J. (2023). Translating genomic advances into biodiversity conservation. *Nature
Reviews Genetics*, 1–12.

Höllinger, I., Wölfl, B., & Hermisson, J. (2023). A theory of oligogenic adaptation of a
quantitative trait. *Genetics*, *225*(2), iyad139. https://doi.org/10.1093/genetics/iyad139
Jain, K., & Stephan, W. (2015). Response of polygenic traits under stabilizing selection and
mutation when loci have unequal effects. *G3: Genes, Genomes, Genetics*, *5*(6), 1065–
1074.
Johri, P., Aquadro, C. F., Beaumont, M., Charlesworth, B., Excoffier, L., Eyre-Walker, A.,
Keightley, P. D., Lynch, M., McVean, G., & Payseur, B. A. (2022). Recommendations for
improving statistical inference in population genomics. *PLoS Biology*, *20*(5), e3001669.
Kachuri, L., Chatterjee, N., Hirbo, J., Schaid, D. J., Martin, I., Kullo, I. J., Kenny, E. E., Pasaniuc,
B., Yuji 29, P. R. M. in D. P. (PRIMED) C. M. W. G. A. P. L. 20 C. M. P. 21 C. D. V. 22 23 D. Y.
24 W. Y. 19 25 26 Z. H. 27 28 Z., & Witte, J. S. (2024). Principles and methods for
transferring polygenic risk scores across global populations. *Nature Reviews Genetics*,
25(1), 8–25.
Kaneko, K., & Furusawa, C. (2006). An evolutionary relationship between genetic variation
and phenotypic fluctuation. *Journal of Theoretical Biology*, *240*(1), 78–86.
Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., Kosiol, C., &
Schlötterer, C. (2011). PoPoolation: A toolbox for population genetic analysis of next
generation sequencing data from pooled individuals. *PloS One*, *6*(1), e15925.
Kumar, S., Deng, C. H., Molloy, C., Kirk, C., Plunkett, B., Lin-Wang, K., Allan, A., & Espley, R.
(2022). Extreme-phenotype GWAS unravels a complex nexus between apple (*Malus
domestica*) red-flesh colour and internal flesh browning. *Fruit Research*, *2*(1), 1–14.
https://doi.org/10.48130/FruRes-2022-0012
Lotterhos, K. E., Fitzpatrick, M. C., & Blackmon, H. (2022). Simulation Tests of Methods in
Evolution, Ecology, and Systematics: Pitfalls, Progress, and Principles. *Annual Review of
Ecology, Evolution, and Systematics*, *53*(1), 113–136. https://doi.org/10.1146/annurev-
ecolsys-102320-093722
Lynch, M., & Walsh, B. (1998). *Genetics and analysis of quantitative traits* (Vol. 1). Sinauer
Sunderland, MA.
Mackay, T. F. (2014). Epistasis and quantitative traits: Using model organisms to study
gene–gene interactions. *Nature Reviews Genetics*, *15*(1), 22–33.
Mathieson, I. (2021). The omnigenic model and polygenic prediction of complex traits. *The
American Journal of Human Genetics*, *108*(9), 1558–1563.
Moore, J. H., & Williams, S. M. (2005). Traversing the conceptual divide between biological
and statistical epistasis: Systems biology and a more modern synthesis. *BioEssays*, *27*(6),
637–646. https://doi.org/10.1002/bies.20236
Orr, H. A. (1998). Testing natural selection vs. Genetic drift in phenotypic evolution using
quantitative trait locus data. *Genetics*, *149*(4), 2099–2104.
Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,
Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in
Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830.
Pfenninger, M., Patel, S., Arias-Rodriguez, L., Feldmeyer, B., Riesch, R., & Plath, M. (2015).
Unique evolutionary trajectories in repeated adaptation to hydrogen sulphide-toxic
habitats of a neotropical fish (*Poecilia mexicana*). *Molecular Ecology*, *24*(21), 5446–
5459. https://doi.org/10.1111/mec.13397
Pfenninger, M., Reuss, F., Kiebler, A., Schönnenbeck, P., Caliendo, C., Gerber, S., Cocchiararo,
B., Reuter, S., Blüthgen, N., & Mody, K. (2021). Genomic basis for drought resistance in
European beech forests threatened by climate change. *Elife*, *10*, e65532.

Pritchard, J. K., & Di Rienzo, A. (2010). Adaptation–not by sweeps alone. *Nature Reviews Genetics*, *11*(10), 665–667.

R Core Team, R. (2013). *R: A language and environment for statistical computing*.

Rijsbergen, C. van. (1979). *Information retrieval*. Butterworth-Heinemann. https://dl.acm.org/doi/abs/10.5555/539927

Santure, A. W., & Garant, D. (2018). Wild GWAS—association mapping in natural populations. *Molecular Ecology Resources*, *18*(4), 729–738. https://doi.org/10.1111/1755-0998.12901

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals—Mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics*, *15*(11), 749–763.

Sella, G., & Barton, N. H. (2019). Thinking about the evolution of complex traits in the era of genome-wide association studies. *Annual Review of Genomics and Human Genetics*, *20*, 461–493.

Shendure, J., Findlay, G. M., & Snyder, M. W. (2019). Genomic medicine–progress, pitfalls, and promise. *Cell*, *177*(1), 45–57.

Stinchcombe, J. R., & Hoekstra, H. E. (2008). Combining population genomics and quantitative genetics: Finding the genes underlying ecologically important traits. *Heredity*, *100*(2), 158–170.

Taylor, C. F., & Higgs, P. G. (2000). A population genetics model for multiple quantitative traits exhibiting pleiotropy and epistasis. *Journal of Theoretical Biology*, *203*(4), 419–437.

Team, T. P. (2019, December 28). *PyPy*. PyPy. https://www.pypy.org/

Tills, O., Holmes, L. A., Quinn, E., Everett, T., Truebano, M., & Spicer, J. I. (2023). Phenomics enables measurement of complex responses of developing animals to global environmental drivers. *Science of the Total Environment*, *858*, 159555.

Turchin, M. C., Chiang, C. W. K., Palmer, C. D., Sankararaman, S., Reich, D., & Hirschhorn, J. N. (2012). Evidence of widespread selection on standing variation in Europe at height-associated SNPs. *Nature Genetics*, *44*(9), 1015–1019. https://doi.org/10.1038/ng.2368

Uffelmann, E., Huang, Q. Q., Munung, N. S., De Vries, J., Okada, Y., Martin, A. R., Martin, H. C., Lappalainen, T., & Posthuma, D. (2021). Genome-wide association studies. *Nature Reviews Methods Primers*, *1*(1), 59.

Van Rossum, G., & Drake, F. L. (2009). *Introduction to python 3: Python documentation manual part 1*. CreateSpace. https://dl.acm.org/doi/abs/10.5555/1592885

Visscher, P. M., Brown, M. A., McCarthy, M. I., & Yang, J. (2012). Five years of GWAS discovery. *The American Journal of Human Genetics*, *90*(1), 7–24.

Visscher, P. M., Wray, N. R., Zhang, Q., Sklar, P., McCarthy, M. I., Brown, M. A., & Yang, J. (2017). 10 years of GWAS discovery: Biology, function, and translation. *The American Journal of Human Genetics*, *101*(1), 5–22.

Waldvogel, A.-M., Feldmeyer, B., Rolshausen, G., Exposito-Alonso, M., Rellstab, C., Kofler, R., Mock, T., Schmid, K., Schmitt, I., & Bataillon, T. (2020). Evolutionary genomics can improve prediction of species' responses to climate change. *Evolution Letters*, *4*(1), 4–18.

Wallace, C. (2021). A more accurate method for colocalisation analysis allowing for multiple causal variants. *PLoS Genetics*, *17*(9), e1009440.

Wang, K., Dickson, S. P., Stolle, C. A., Krantz, I. D., Goldstein, D. B., & Hakonarson, H. (2010). Interpretation of association signals and identification of causal variants from genome-wide association studies. *The American Journal of Human Genetics*, *86*(5), 730–742.

Wray, N. R., Kemper, K. E., Hayes, B. J., Goddard, M. E., & Visscher, P. M. (2019). Complex
trait prediction from genome data: Contrasting EBV in livestock to PRS in humans:
genomic prediction. *Genetics*, *211*(4), 1131–1141.

Wright, S. (1949). THE GENETICAL STRUCTURE OF POPULATIONS. *Annals of Eugenics*,
15(1), 323–354. https://doi.org/10.1111/j.1469-1809.1949.tb02451.x

Xie, C., & Yang, C. (2020). A review on plant high-throughput phenotyping traits using UAV-
based sensors. *Computers and Electronics in Agriculture*, *178*, 105731.

Yang, J., Jiang, H., Yeh, C.-T., Yu, J., Jeddeloh, J. A., Nettleton, D., & Schnable, P. S. (2015).
Extreme-phenotype genome-wide association study (XP-GWAS): A method for
identifying trait-associated variants by sequencing pools of individuals selected from a
diversity panel. *The Plant Journal*, *84*(3), 587–596.

Zhang, W., Liu, A., Albert, P. S., Ashmead, R. D., Schisterman, E. F., & Mills, J. L. (2018). A
pooling strategy to effectively use genotype data in quantitative traits genome-wide
association studies. *Statistics in Medicine*, *37*(27), 4083–4095.
https://doi.org/10.1002/sim.7898

Ziyatdinov, A., Kim, J., Prokopenko, D., Privé, F., Laporte, F., Loh, P.-R., Kraft, P., & Aschard, H.
(2021). Estimating the effective sample size in association studies of quantitative traits.
G3, *11*(6), jkab057.

825