Intra-genomic genes-to-genes correlation enables genome representation

Jingjie Chen[1,9]   Xuchuan Ma[2,9]   Junwei He[1]   Yingxia Wang[3]   Yuji Ren[4]

Li Qi[7]   Liuyang Song[7]   Lin Ruan[1]   Cun Fan[1]   Jun Jiang[1]

Yongjia Bao[1]   Tao Huang[1]   Jingbo Cheng[1]   Xing Liu[5]   Fang Chen[6]

Yichen Huang[6]   Na Zhao[7,8,*]   Chunming Guo[1,*]

[1] Yunnan Key Laboratory of Cell Metabolism and Disease, Center for Life Sciences, School of Life Sciences, Yunnan University, Kunming 650500, China.

[2] Food Microbiology, Wageningen University & Research, Wageningen, the Netherlands.

[3] The Department of Pathology, First affiliated Hospital of Kunming Medical University, Kuming, People's Republic of China.

[4] The Department of Clinical Laboratory, First affiliated Hospital of Kunming Medical University, Kuming, People's Republic of China.

[5] Department of Urology, Children's Hospital of Chongqing Medical University, 136 Zhongshan Road, Chongqing 400014, PR China.

[6] Department of Urology, Shanghai Children's Hospital, School of medicine, Shanghai Jiao Tong University.

[7] Key Laboratory in Software Engineering of Yunnan Province, Yunnan University, Kunming, People's Republic of China.

[8] Big Data Research Center, University of Electronic Science and Technology of China, Chengdu, People's Republic of China.

[9] These authors contributed equally.

[*] Correspondence: Na Zhao <zhaona@ynu.edu.cn>, Chunming Guo <chunmingguo@ynu.edu.cn>

Abstract The pan-genome consists of core genes shared by all members of a taxonomy and accessory genes found in only a subset, holding the keys to advancing our understanding of evolution and tackling medical challenges. Here, we discovered a strong intra-genomic correlation among bacterial genes within each of *Escherichia coli*, *Listeria monocytogenes*, *Staphylococcus aureus*, and *Campylobacter jejuni*. With a convolutional neural network assisted core genome knock-out simulation, we found that different combinations of fewer than 20 highly variable core genes could recover the sub-species type classified by complete core genome with accuracy >95%. This observation led us to test the genes-to-genes predictability: with more than 52,000 assemblies from each species, combinations of highly variable core genes could predict the sequence variants of other core genes (average accuracy >94%) within the same genome and could also predict sequence variants (average accuracy >91%) as well as the presence (average AUROC >0.91) of some accessory genes. Furthermore, combinations of highly variable core genes could also predict multiple antibiotic resistances (AUROC >0.80) in large published datasets of *E. coli*, *S. aureus*, and *Mycobacterium tuberculosis*. Collectively, we propose that genes within the same genome can strongly correlate with each other. Therefore, the strain phylogeny and the stauts of other genes could be uniformly represented by combinations of highly variable core genes, which could further represent certain phenotypes including *in vitro* resistance.

Key Words: Core-Accessory Genome, Gene-Gene Correlation, Genome Prediction, Resistance Prediction.

Introduction

The bacterial pan-genome consists of core genes shared by all strains within a taxonomic unit and accessory genes found only in a subset. Core genes, characterized by their conservative presence, hold functions for basic survival, including but

not limited to DNA metabolism, protein synthesis, and energy metabolism. Core genome is a well established concept
proven experimentally using comparative proteomics; it defines bacterial lifestyle and the expression of core genes was
relatively inflexible with respect to culture conditions[1]. Single-copy core genome based phylogeny is a frequently used
method for strain subtyping, becoming increasingly achievable with the growing availability of assemblies and decreasing
sequencing costs. Within-species single-copy core genome, containing hundreds to more than thousands of genes, can
classify strains into different core genome subtypes (cgSTs) at a high level of resolution while ensuring interlaboratory
reproducibility[2,3]. The accessory genome, which includes genes that can be lost and acquired, facilitates the acquisition
of functions. These genes include toxin genes, virulence factors (VFs), and antibiotic resistance genes (ARGs)[4–7]. The
function acquisition partially explained the rapid adaptation of pathogens to new environments[8]. The accessory genome,
as a gigantic pool of different functions, has been employed as input for machine learning based resistance prediction for
certain antibiotics without a *priori* knowledge of mechanisms[9].

Regarding to relationship among genes, previous studies[10–12] had found that large amount of bacterial genes could display
co-occurrence or avoidance patterns with one another, and some studies on *Listeria monocytogenes* had presented a strong
correlation between core genome based sublineage and accessory gene presence[13–16]. In *L. monocytogenes*, internalin A
(*InlA*) is an example of gene-gene sequence variants correlation. Compared with original full-length InlA sequence variant,
truncated InlA is more specifically present in certain clonal complexes classified by seven housekeeping genes based Multi-
Locus Sequence Typing (MLST)[13,14,17–21]. However, studies exploring the relationships among genes in the pan-genome,
particularly on a large scale and including correlation of both presence and sequence variants, were notably missing.

Herein, our study characterized the correlation among genes in the pan-genome by evaluating the genes-to-genes pre-
dictability. Status of different core and accessory genes, including their exact sequence and presence, were employed as
input to infer the status of other genes in the same genome using convolutional nerual network (CNN) and also certain
phenotypes such as *in vitro* antibiotics resistance. A large amount genes was involved, and high average predictability was
achieved over the dataset of three bacterial phyla. Thus we believe this predictability is generalizable and will provide
novel insights for pan-genome study.

Results

VFs and ARGs correlate with cgSTs in 137 *E. coli* strains

To study genomic features held by *E. coli* that caused Urinary Tract Infection (UTI), ninety one strains were collected
from patients in The First Affiliated Hospital of Kunming Medical College. Their assembled genomes were re-annotated
together with 46 reference assemblies (Supplementary Table 1). A customized ortholog clustering pipeline OrthoSLC
was employed to build gene clusters. For a given set of assemblies, gene clusters were categorized into accessory gene
clusters (not shared by all assemblies) and core gene clusters (shared by all assemblies). If a given core gene cluster had
one gene from each of the assemblies, the cluster is defined as 'strict core' gene cluster (StCGC); if any assembly had
more than one gene in a core gene cluster, the core gene cluster is defined as 'surplus core' gene cluster (Supplementary
Fig 1). The multi-sequence alignment (MSA) of 2170 StCGCs, were concatenated for phylogeny computation. Based
on reference strains, the phylogeny was classified into 5 cgSTs with known references[3] and 1 cgST without; this newly
identified cgST was named as 'F' in the study (Figure 1a, Supplementary Data 1). All reference strains were in the same
cgSTs as classified in the previous study, and *E. coli* collected from UTI patients were more frequently present in cgST
B2 (53 out of 91)[3,22–26]. To interrogate whether B2 strains possessed specific VFs not carried by other cgSTs, the VFs
of total 137 genomes were annotated, and more VFs of category adherence were detected in B2 strains (Supplementary
Fig 2, Supplementary Data 1). This might indicate a better ability of urinary tract colonization of B2 strains which were
more frequently isolated from UTI patients. Interestingly, some VFs were almost exclusively present only in certain cgSTs.
For example, a fimbriae gene *cfaD/cfaE* was exclusively present in B1 strains (17/17 in B1, and 0/120 in non-B1 strains).
Another group of genes, *shuY*, *shuA*, *shuS*, *chuT*, *chuX*, and *chuS* exhibited an absolute absence in cgST B1 and A (0/41
in B1 and A, 96/96 in other cgSTs) (Supplementary Table 2 for gene name and VFDB[27] query correspondence).

To further examine the correlation between cgST and VFs, strains were divided into 2 groups, cgST B2 (n = 68) and
non-B2 (n = 69). Principal Coordinate Analysis (PCoA) was applied using genome VFs carriage status, including presence
of VFs and sequence variant of these present VFs which was reflected by similarity against reference sequences in VFDB[27]
(Figure 1b). The B2 and non-B2 groups exhibited a significant difference and a higher F value in Bray-curtis result than in
Jaccard result, which suggested the cgST not only correlated with VFs presence but also correlated with sequence similarity
against references. To find the VFs most significantly correlating with B2 or non-B2, the VFs status was used for B2 or
non-B2 classification by Lasso Regression (LR) models, and VFs overlapping with StCGCs were excluded from input. The

training was performed 50 times and each re-train started with randomly selected samples (70% from each cgST) as the
training set. Test set AUROC of 1.00 was achieved in all 50 times of re-trains. By extracting the average weights that LR
models gave to each VF over 50 re-trains, all VFs with an average weight of more than zero were displayed in Figure 1c. The
result showed that *espX4* as a Type III secretion system (TTSS) secreted effector, exhibited exclusive (69/69) presence in
non-B2 cgST. Similarly, two homologs of TTSS secreted effector *espL1* from *E. coli* CB9615 and EDL933 (Supplementary
Table 2) showed almost exclusive (68/69) presence in non-B2 cgST. By targeted PCR (Figure 1d) and Sanger sequencing,
those bands with correct length present in non-B2 genomes turned out to be corresponding sequences. These findings
indicated a correlation between VF presence and cgST.

Besides VFs, ARGs were annotated using Resistance Gene Identifiers (RGI) and CARD database[28] (Supplementary Data
1). The PCoA on ARGs status showed a significant difference between B2 and non-B2, and a higher F value in Bray-
curtis result than in Jaccard result could suggest that RGI score as an indirect reflection of sequence variant of ARGs
is correlated with cgST (Figure 1b). Using ARGs status as input, the re-train based cross validation in predicting cgST
of each strain returned two genes *marA* and *leuO* (Figure 1c). Both genes were present in almost all analyzed strains
(136/137 for *marA* and 135/137 for *leuO*), and the RGI score of these two genes strongly correlated with cgST. *marA*
was only scored 2 (68/68) in non-B2 strains and *leuO* was almost exclusively scored 2 in B2 strains (66/69). Hence, the
sequence variants of some ARGs were specifically associated to cgST with a pronounced manner.

Fig 1: **Both VF and ARG status correlate with cgSTs of UAEC. | (a) Single-copy core genome based phylogeny of 46 reference (cyan strain name) and 91 isolated (red name) E. coli strains.** The clade of each cgST was stained differently and corresponding cgST names are in the color bar below. Bootstrap values lower than 95 are shown. **(b) PCoA of VFs (first row) and ARGs (second row) using weighted (Bray-Curtis, first column) and unweighted (Jaccard, second column) algorithms.** The permanova based significance test between B2 (red dot) and non-B2 (purple triangle) strains is available on the bottom of each plot. **(c) Heatmap of genes specific to certain cgST selected by random re-trained LR models.** Color gradient in the VFs plot indicates the sequence similarity (<80% was regarded as absence of gene) between a corresponding gene of each strain and VFDB reference. *espL1-C* and *espL1-E* stand for *espL1* gene from *E. coli* CB9615 and EDL933 (Supplementary Table 2), respectively. RGI score of 0 stands for no resistance (not equivalent to gene absence); 1 stands for a strict hit, meaning potential resistance; and 2 stands for a perfect hit, indicating resistance. Scores were determined based on protein sequence similarity with reference and SNP site check by RGI[28]. The color on x axis stands for the cgST of each strain as in a. **(d) PCR based gene presence confirmation.** *espX4* had original length of 1581 bp in VFDB[27] and its PCR target region is ~1500 bp as indicated in sidebar; *espL1* original length is 1898 bp and target region is ~1400 bp as indicated in sidebar.

Core-to-accessory genome correlation is present in multiple phyla

To confirm whether the correlation is present in more cgSTs and is ubiquitous in bacteria kingdom, we included 4 species, *E. coli*, *L. monocytogenes*, *S. aureus*, and *C. jejuni*. The four species were selected based on diversity and assembly availability consideration. *L. monocytogenes* and *S. aureus* are Gram positive while *E. coli* and *C. jejuni* are Gram negative. They originated from three Phyla: *Pseudomonadota*, *Bacillota*, and *Campylobacterota*. And these two *Bacillota* species *L. monocytogenes* and *S. aureus* are from different families, *Listeriaceae* and *Staphylococcaceae*, respectively. 65,840 *E. coli*, 52,090 *L. monocytogenes*, 54,817 *S. aureus* and 54,905 *C. jejuni* assemblies (Supplementary Data 2) were included in the analysis as Species Assembly Collection (SAC). To obtain sufficient assemblies to study more cgSTs, we adopted a strategy of pseudo-labeling based dataset construction (Figure 2a). In the workflow, Dataset I of 600 to 900 assemblies were randomly selected from the SAC for each species, and a random forest (RF) model trained with Dataset I (Supplementary Data 3) was employed to predict the cgST of all members in SAC inputting with VF status of assemblies (Supplementary Data 2). The result of pseudo-labels was utilized to assign 200 assemblies to each cgST of Dataset II (Supplementary Data 3), including some cgST references selected from Dataset I. The result of Dataset II construction is shown in Figure 2b for *E. coli* and Supplementary Fig 3-5 for the other 3 species. The identification of cgST in Dataset II were referenced by strains selected from Dataset I (Supplementary Table 3-6). In Dataset II, 478 StCGCs for *E. coli*, 1,082 for *L. monocytogenes*, 672 for *S. aureus* and 522 for *C. jejuni* were constructed (Supplementary Data 7). The assignment of genomes from SAC to cgSTs showed a high accuracy. Besides cgST A and B in *S. aureus*, all other cgSTs among species of Dataset II were assigned 200 ± 5 assemblies. Therefore, these results indicate a correlation between cgST and VF status.

With evenly distributed cgST and sufficient assemblies in Dataset II (Supplementary Data 7), the accessory genes that most significantly correlate with certain cgSTs was analyzed. By using VF or ARG status (Supplementary Data 3 and 4) as input, 50 re-trains were applied on binary classification of each cgST using LR models. Notably, genes overlapping with StCGCs were excluded from prediction input. Genes with high average weights in trained models were displayed in Supplementary Fig 6. The average weights of all VFs or ARGs could be seen in Supplementary Data 5. The results in Supplementary Fig 6 suggested that cgST could be predicted using VF status with an average AUROC higher than 0.96 for all cgSTs and status of many accessory genes apparently correlated with cgST. Relatively low average AUROC (<0.90) when using ARGs as input in *L. monocytogenes* and *C. jejuni* could result from the lack of known ARGs in these species for cgST prediction. To date, *E. coli* had $>250,000$ total assemblies in NCBI database, and there were in total 643 VFs, 249 ARGs in Dataset II. In contrast, there were less than 100,000 total assemblies in NCBI assembly database for each of *L. monocytogenes*, *C. jejuni*, and *S. aureus*, and only 50, 151, and 122 VFs as well as 22, 37, and 79 ARGs were found in Dataset II, respectively. Despite these variations, correlation among cgST, VFs, and ARGs could be observed across Phyla and in more cgSTs.

Fig 2: **Accurate prediction of cgST by accessory gene status across three phyla. | (a) The schematic workflow of Dataset construction.** Pseudo-labels for all assemblies in SAC were generated by the random forest model trained using Dataset I. These pseudo-labels were used to assign ~200 assemblies to Dataset II (including reference with known cgST from Dataset I). **(b) cgST result of Dataset I (left) and II (right) of _E. coli._** cgSTs of clades were identified using reference strains (Supplementary Table 3-6). The uncolored branches in Dataset I were not classified into any cgST and not used in RF model training. The dashed branches were of large length and were manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The tables in the middle of **(b)** describes the confusion matrix which excludes the reference strains (from Dataset I) and amount of strains next to each cgST of Dataset I and II. Results for _L. monocytogenes_, _C. jejuni_, _S. aureus_ are available in Supplementary Fig 3-5.

Different sets of limited core genes recover the phylogeny with CNN

According to previous studies[10–12], certain bacterial genes had significant co-occurrence/avoidance patterns. Thus, the correlation between accessory gene status and cgST in Supplementary Fig 6 might actually result from the gene co-occurrence phenomenon. This was because some StCGCs might actually be a misclustering of two or more accessory gene clusters which had different functions (heterofunctional cluster). If a certain cgST was separated from other cgSTs because of a heterofunctional cluster, the specific accessory gene presence in this certain cgST could co-occur with another gene (Supplementary Fig 7). On the other hand, if a certain cgST was separated from other cgSTs because of a StCGC which consisted of genes with same function (homofunctional cluster), this specific accessory gene presence in certain cgSTs was different from gene co-occurrence but rather a correlation between core gene sequence variant and accessory gene presence (Supplementary Fig 7).

To differentiate our discovered correlation from gene co-occurrence, we delved into all StCGCs and attempted to find a subset of StCGCs, which were homofunctional and could still recover the complete core genome cgST. This was confronted with two challenges, the large number of subset StCGC combinations and extensive computation time of phylogeny with manual cgST identification. To overcome these obstacles, a simple Cluster Variation Level (CVL) algorithm was designed to value the extent of within-StCGC variation inferring the potential that each cluster might decide the cgST classification (Figure 3a). A 5-layer CNN as model A (Supplementary Fig 8), which could be reproducibly trained to generate high test set performance (accuracy >98%) models inputting with complete core genome, was employed to carry out cross validation based cgST recovery evaluation inputting with different combinations of StCGCs. Within each species, the samples of Dataset II were separated into three sub-groups, and the samples of each cgST were evenly distributed in three sub-groups. The cross validation was performed three-fold with every two sub-groups as training set and three re-trains with Xavier random initialization[29] on each training set, meaning a total of nine re-trains to evaluate each input combination. To input with a subset of StCGCs, the complete core genome dimension was maintained but positions of genes not in the input subset were all set to zero, as knock-out (KO) simulation. In Supplementary Fig 9 first row and Figure 3b first row, by increasing the number of StCGCs as prediction input starting from the highest CVL rank, the test set performance was compared with when complete core genome and all-zero matrix were input. We found that 6 highest CVL StCGCs for *E. coli*, 11 for *L. monocytogenes*, 10 for *C. jejuni* and 5 for *S. aureus* were needed to recover cgST based on complete core genome with test set accuracy >98% (Supplementary Data 8), and these top CVL StCGCs are defined as hub StCGCs. This outcome corresponded to the phylogenetic tree of *E. coli* constructed using hub StCGCs only (Supplementary Fig 10), in which the cgST result exhibited highly similarity (1197/1200) comparing with the phylogeny structure based on complete core genome. After function annotation with eggnogmapper[31], all these hub StCGCs turned out to be of homofunctional on protein sequence level (Supplementary Data 7). This result could primarily illustrate that cgST in Dataset II could be decided by homofunctional StCGCs. Therefore, the correlation that accessory gene status was specific to certain cgSTs did not result from gene co-occurrence.

To investigate if homofunctional StCGC based cgST recovery could also be achieved with non-hub StCGCs, their function annotation was performed: only two StCGCs in *E. coli* (the 106^{th} and 134^{th} in CVL rank) and one in *C. jejuni* (ranked 163^{th}) were heterofunctional among the four species (Supplementary Data 7). This result suggested that most StCGCs were homofunctional, so the same cross validations same as Figure 3b first row was applied to evaluate more combinations of input (one StCGC from each of *E. coli* and *L. monocytogenes* with 0 CVL value was excluded from input). In Figure 3b second row (Supplementary Data 8), *E. coli* result showed that 95.48±3.58% cross validated average accuracy of test set could still be achieved inputting with 281^{th} to 286^{th} highest CVL StCGCs. These 286 highest CVL StCGCs represented 59.96% of all *E. coli* StCGCs (least among four species), so the hub StCGCs were not the only options for recovering the cgST of the complete core genome. In Supplementary Fig 9 second row, when the window length of input was increased to 20 genes, better performance could be achieved, the test set performance of 401^{th} to 420^{th} highest CVL StCGCs reached 97.33±0.99% for *E. coli*. In Supplementary Fig 9 third row, around 100, 250, 150, 200 lowest CVL rank StCGCs for *E. coli*, *L. monocytogenes*, *C. jejuni* and *S. aureus*, was required to recover complete core gene cgST with accuracy >98% (test set). Performance of these input windows corresponds to Supplementary Fig 10-13 of *E. coli* phylogeny structures. When top StCGCs were used for phylogeny construction, the cgST result for all strains was recovered almost perfectly although the detail was not identical to the complete core genome phylogeny. When lower-ranked StCGCs were input, only certain cgSTs were still recovered well. In Supplementary Fig 13, when 100 lowest CVL *E. coli* StCGCs were input, 1192 out of 1200 were correctly assigned. These results suggested that complete core genome cgST could be recovered using many combinations of StCGCs while these StCGCs are homofunctional. Thus, cgST specific sequence variant and presence of an accessory gene did not result from gene co-occurrence.

Fig 3: **Limited core genes can accurately recover cgSTs with CNN in cross validations.** | **(a) schematic workflow:** re-annotated genomes of Dataset II (EC: *E. coli*, LM: *L. monocytogenes*, CJ: *C. jejuni*, SA: *S. aureus*) were sent to construct gene clusters where StCGCs were ranked by their CVL. The core genome was concatenated from high CVL StCGCs (genes more variable among strains) to low CVL gene clusters (genes more conserved among strains). Different KO combinations allowed the test of the number of hub genes (least top CVL genes needed to recover complete core genome cgST with >98% accuracy) as in **b** first row and Supplementary Fig 7 first row. By using different windows of available core genes in prediction (**b** second row), the ability to recover complete core genome cgST was tested on a wider range of core genome. Each box of distribution represents the performance of nine re-trains (three-fold, three random initialization each fold) of train set (red) and test set (blue). The CNN model adopted model A as in Supplementary Fig 7.

Different sets of limited core genes predict other genes in the same genome

In Supplementary Fig 6, the observation of correlation between cgST and accessory gene status was limited because the cgST was only divided into several general classes. To observe the correlation between core and accessory gene status more directly, accessory gene presence prediction was attempted using StCGCs as input and CNN model B from Supplementary Fig 8. Within each species, if there were more than 5,000 assemblies in SAC owning a certain gene and this gene was also absent from more than 5,000 assemblies of SAC, the presence predictability of this gene was evaluated. Two-fold cross validations were performed with two randomly initialized re-trains on each fold, so the test set performance of four re-trains was employed to evaluate the predictability of a single gene. To balance the computation labor and sample size, the input dimension was reduced. A top window in CVL rank list with only hub StCGCs were employed as input and two other windows were selected from the middle and lower part of CVL rank list (Figure 4a). Middle and lower windows included more genes to reach a similar sequence length of the top window. These three windows included a KO version and a complete version of input. In addition, a window using housekeeping genes was included (Supplementary Table 8).

Among the four species, all complete version of top windows reached average test set AUROC >0.91 (0.91±0.09 in *S. aureus* as the lowest of four species) in predicting accessory gene presence (Figure 4b first row, Supplementary Data 9). Notably, presence of accessory genes could not always be accurately predicted. For example, ARGs in *E. coli* had an average AUROC 0.79±0.06 using top window as input. Apart from complete top windows, similar performance could be achieved with complete middle, lower and housekeeping windows. This suggested that the presence of accessory genes could be inferred by different combinations of StCGCs. Besides the KO input of the top window in *C. jejuni*, prediction, KO input of all windows exhibited a slightly lower performance compared with their complete versions. These results suggested that inferring the presence of some accessory genes required a limited amount of core genes which could be selected from a large range.

Apart from presence, sequence variant prediction of VFs and ARGs was also tested. Any gene to be predicted had at least 2 prevalent sequence variants that were present in more than 2,000 assemblies of the SAC. Assemblies were grouped into classes owning different prevalent sequence variants of the gene to be predicted, and a class 'other' of assemblies holding all other non-prevalent sequence variants. A two-fold cross validation was performed, and each fold had two re-trains with random initialization, so a total of 4 re-trains for evaluation. In Figure 4b second row (Supplementary Data 10), all complete top windows reached average accuracy >91% (91.98±6.91% in *E. coli* as lowest among four species) and similar accuracy in other complete windows. Performance of KO input showed some decrease comparing with their complete versions. For *E. coli*, performance of top window input reached 96.90±5.85% when the class 'other' was excluded from the prediction (Supplementary Fig 14 first column). These results exhibited that there is a correlation between the core gene status and the status of accessory genes from the same genome, this correlation could be reflected by both accessory gene presence and sequence variant prediction by input of combinations of limited amount of StCGCs from a large range.

Besides the correlation between core and accessory genes, the observation in Figure 3b second row that different StCGC combinations could predict the same cgST led to a hypothesis: core genes within a same genome could predict the sequence variant of each other. To test this hypothesis, the same input of StCGC combinations in Figure 4b first and second row were employed. Notably, if the gene to be predicted is one of the input genes, the corresponding positions of the input would be set as zero. Self prediction was skipped when first gene in CVL rank was the only input and the gene to be predicted at the same time. In Figure 4b third row (Supplementary Data 10), all top windows showed an average accuracy >94% (94.47±2.35% in *C. jejuni* as lowest among four species), and similar accuracy could be achieved using other complete windows as input. For *E. coli*, the top window result increased from 94.65±2.40% to 97.61±2.19% when class 'other' was excluded (Supplementary Fig 14 first column). Notably, some prevalent sequence variants could have only one nucleotide difference while the prediction still reached high accuracy (Supplementary Fig 15, 16). These results suggested that correlation among genes was not limited between core and accessory genes, different combinations of StCGCs could infer other genes with a high accuracy.

Notably, this predictability involved a large range of genes in the pan-genome. In *E. coli* SAC dataset. By pre-calculating all input to the final fully connected layer with a pre-trained model, embedding was achieved. This reduced computation labor significantly by representing a long DNA sequence of each sample with a small vector of 688 float numbers and training only the final layer. Genes in *E. coli* dataset II with presence rate <100% were extracted from SAC, and status of these pan-genes were predicted with embedded input. Among these pan-genes, presence prediction of 3605 accessory genes with AUROC 0.91±0.09 and sequence variant prediction of 4913 pan-genes with accuracy of 91.39±7.17% (Supplementary Fig 17 second column, Supplementary Data 11). Here, we adopt the term genes-to-genes (g2g) correlation to conceptualize the observation. The g2g correlation means that the status of a given gene, including presence and sequence variants, correlates with the status of other genes from the same genome. Such correlation can explain the phenomenon that the status of some genes can be accurately predicted by other genes from the same genome.

Fig 4: **Limited core genes can predict status of other genes within the same genome with a high average accuracy.** | The workflow (**a**) included windows of similar length as input, a top window with all hub core genes each species and 2 windows using middle and lower end CVL rank StCGCs. All windows included a KO version and a complete version of input. KO input for top windows kept only the highest CVL StCGC as input; middle and lower windows kept only first n (amount of hub) genes. In addition, a window of housekeeping genes (Supplementary Table 8). Each point represented the average test set performance (two-fold, two random initialization) of four re-trains for accessory gene (Acg.) presence prediction (**b** first row) and sequence variant (Seq Vrts) prediction of VFs/ARGs as well as core genes (Cg.) (**b** second, third row). The number of genes being predicted are available in each plot. For each gene included in presence prediction (**b** first row), there should be >5,000 samples from SAC with the gene present in the assemblies while assemblies without the gene present should also be >5,000. For each gene included in sequence variant prediction (**b** second, third row), there should be at least two sequence variants owned by >2,000 assemblies in SAC. The CNN model adopted model B in Supplementary Fig 8.

245 Different sets of limited core genes predict host response and antibiotic resistance

Core genes exhibited great power in inferring other genes of the same genome, implying a potential to represent a genotype
closer to whole genome with limited parts. To test this potential, the prediction of certain phenotype was attempted using
limited core genes as input. Published datasets[9,32–35] (Supplementary Table 6) with clear antibiotics resistance labels
were selected from BV-BRC[36,37] from which the assemblies of *E. coli*, *S. aureus* and *Mycobacterium tuberculosis* were
obtained. Besides, the phenotype prediction of 91 patient-isolated *E. coli* in this study (Supplementary Data 12) was
tested, including mouse bladder infection host response and resistance to certain antibiotics. In Figure 5b (Supplementary
Data 13), all host response index had average performance lower than AUROC 0.80 and large deviation over nine re-
trains (three random initialization on each of the three folds). This low performance and larger deviation also occurred
with resistance prediction in our study. The high deviation among re-trains results from low sample size. Particularly,
by comparing performance consistency over re-trains of our study with published datasets, larger sample size showed
more consistency over re-trains (Supplementary Fig 17a). However, the large sample size did not necessarily lead to high
performance (Supplementary Fig 17b). For example, Amoxicillin-Clavulanate (AMC) and Ampicillin (AMP) in *E. coli* of
published datasets had average AUROC lower than 0.80 while there were at least 339 assemblies representing susceptible
and resistant strains. Interestingly, predictions of resistance to multiple drugs reached average AUROC >0.80 in at least
two input windows, and the prediction of some resistance exhibited AUROC >0.90, including Ciprofloxacin (CIP) and
Methicillin (MET). Particularly, CIP resistance exhibited prediction AUROC >0.90 across three species based on the data
from three different studies[9,33,35].

For AMP resistance prediction in *E. coli*, AUROC 0.70 ± 0.01 for dataset of Davies et al.[32] and 0.73 ± 0.02 for dataset of
Moradigaravand et al.[9] were achieved inputting with hub StCGCs (Figure 5b). However, Davies et al.[32] achieved AMP
prediction sensitivity 98% by simply predicting strains owning beta-lactamase genes (e.g., *OXA-1* and *TEM-30*) as AMP
resistant. Therefore, the presence prediction performance of *TEM-30* (AUROC 0.76 ± 0.01, Supplementary Data 9) corre-
sponded to the low AMP resistance performance when inputting with hub StCGCs. In contrast, the high performance
(AUROC >0.90) on CIP and MET resistance prediction in *S. aureus* and *E. coli* using hub StCGCs corresponded to
accurate prediction on key genes. The MET resistance related genes *mecR1, mecI, mecA* available in our study (Supple-
mentary Data 9) had presence prediction AUROC of 0.97 ± 0.01, 0.97 ± 0.01 and 0.87 ± 0.01, respectively, inputting with
hub StCGCs. Sequence variant prediction accuracy of *gyrA* (Supplementary Data 10) reached 0.80 ± 0.01 in *S. aureus*
and 0.95 ± 0.01 in *E. coli* (inputting with hub StCGCs), which corresponded to CIP resistance prediction performance.
Notably, the performance of Isoniazid (INH) and Rifampicin (RIF) resistance prediction using housekeeping gene input
was higher than other input windows. Since the most resistance-related genes for INH and RIF, *katG* and *rpoB* are
included in the housekeeping genes, respectively, this accurate resistance prediction should not result from a better ability
of housekeeping genes in inferring the status of other genes than other input windows. These results suggested that while
phenotype prediction is still challenging in general, but certain antibiotics resistance could be well inferred implying the
correspondence between phenotype and genotype which was represented by limited core genes.

Fig 5: **Limited core genes predict host response (only this study) and antibiotics resistance (in large published datasets and this study).** | The input windows shown as in **a**, included a complete top window of hub core genes and three KO input with three highest CVL genes kept, respectively. Both *E. coli* (EC) and *S. aureus* (SA) included 3 windows as in Figure 4, while *M. tuberculosis* (MT) includes only 2 windows. (**b**) Results of each species included a input window of housekeeping genes. Each box of distribution represents the test set performance of 9 re-trains (three-fold, three random initialization each fold). The CNN model adopted model B as in Supplementary Fig 8. Abbreviations: IBC (Intra-cellular Bacterial Community), AMC (Amoxicillin-Clavulanate), AMK (Amikacin), AMP (Ampicillin), AMX (Amoxicillin), CET (Cephalothin), CIP (Ciprofloxacin), CLI (Clindamycin), CPM (Capreomycin), CSN (Cycloserine), CTX (Cefotaxime), CTZ (Ceftazidime), CXM (Cefuroxime), EMB (Ethambutol), ERY (Erythromycin), ETH (Ethionamide), FUS (Fusidic Acid), GEN (Gentamicin), INH (Isoniazid), KAN (Kanamycin), MET (Methicillin), MXF (Moxifloxacin), NIF (Nitrofurantoin), OFL (Ofloxacin), PAS (Para-Aminosalicylic Acid), PEN (Penicillin), PZA (Pyrazinamide), RIF (Rifampicin), STR (Streptomycin), TBM (Tobramycin), TET (Tetracycline), TMP (Trimethoprim). The sample size is listed in brackets next to each phenotype predicted, format as "(positive samples / negative samples)". Positive samples were resistant to the antibiotics, or led to high edema level (top 50%), or generated IBC in mouse bladder, or led to high immune score (top 50%). Negative samples stand for those susceptible to given antibiotics, or led to low edema level (lower 50%), or no IBC in mouse bladder, or led to low immune score (lower 50%). Dashed horizontal lines stand for AUROC of 0.8 (lightgrey) and 0.9 (pink).

Discussion

We presented a novel manner in studying bacterial pan-genome by evaluating g2g predictability. The high average predictability among large amount of strains and genes demonstrated the generalizability. This g2g correlation allowed representation of strain characteristics using very limited genomic information which was tested with *in vitro* resistance prediction (Figure 5). Beyond lineage specific accessory gene presence[13,14,16,17] and gene co-occurrence[10–12], g2g correlation was further proved to involved the exact sequence of genes.

The strong correlation among genes demonstrated novel possibilities of genotyping and potential in phenotype inference. For example, *E. coli* cgST B2 could be identified by absence of *espX4* and *espL1* without whole genome sequencing and complicated bioinformatics processes. New typing strategies can be independent of reference strains, requiring a small CNN or RF model trained on a moderate scale of sample size and providing better genome representation.

Our study proposed a set of algorithms and methods to test and find limited core genes which allowed the representation of a genome. The total length of input gene sequences can be as short as <5,000 bp (housekeeping genes) which is highly practical considering the sequencing cost nowadays. By embedding with our pre-trained model, the representation of a long DNA sequence from genome can be further reduced to a vector of less than a thousand float numbers. All these efforts will bring feasibility and efficiency for the future large scale analysis.

g2g correlation is akin to the Pythagorean theorem, as knowing parts of a genome, many other parts could be predicted. So far, we could not find the tropism of predictability because high or low presence frequency of genes in SAC did not show high predictability (Supplementary Fig 14 third column) and many pan-genes lacked pathway annotation made it difficult to categorize genes without bias. We do not consider that plasmid or genomic islands can fully explain how g2g was formed. As the genes involved in g2g are too many to be held by a mobile genetic element. The most possible underlying process of g2g correlation could be gene co-evolution under environmental pressure. Facing a certain environment sometimes means facing pressure of multiple aspects. The specific status of some genes brings competitiveness over other strains such as better survival when fighting against the host immune response. Meanwhile, some core genes also adapted to specific nutrient availability of the environment. Genes did not necessarily interact with each other, but after generations in the environment, a certain combination of status of multiple genes was fixed. This could take a long time to stabilize which makes it hard to be observed.

Notably, the core genes that are available for prediction input, shall not be limited to the StCGCs provided in this study. The definition of StCGCs might exclude some soft core clusters which might also provide accurate prediction. By reducing the length difference limit in RBBH pairs[38], the amount of StCGCs could increase. Moreover, there are 2,617 and 3,194 gene clusters present in over 99% and 95% strains in Dataset II of *E. coli* suggesting a larger range of input options.

It is worth noting that k-mer patterns from either high or low CVL input combinations could predict cgST with high accuracy (Supplementary Fig 18,19), but within each cgST, k-mer patterns from different input combinations exhibited significant difference (Supplementary Fig 20). This means models had to learn different patterns for different input window. In addition, with models pre-trained on complete core genome, re-prediction using KO input exhibited a much lower accuracy (Supplementary Fig 21 first row), comparing with Figure 3b, where low CVL windows could still be trained to generate high performance models. Performance of pre-trained models was not improved even if a structure similar to Res-Net[39] was used for pre-trained models (Supplementary Fig 22 and Supplementary Fig 21 second row). Therefore, performance evaluation of certain input windows was by re-trains and cross validations.

Our resistance prediction results shown in Figure 5 brought up a notable issue that KO simulation might not return genes responsible for a certain phenotype. Since in multiple resistance predictions, we can see that similar accuracy could be achieved by different windows so we might not know which window the phenotype related gene is in. A very accurate prediction could also confuse, as we can't confirm the responsible gene is in the input window or is a gene outside the window but well inferred.

We have to realize that genes involved in this study are still a small portion of the entire pan-genome. In *E. coli* Dataset II, there were in total 72,275 gene clusters which could correspond to the previous study[4]. Surprisingly, 61,051 clusters were found in only 1 strain based on which we can expect that a lot genes lack enough presence and prevalent sequence variants in *E. coli* SAC. Therefore, the status predictability of majority of pan-genome remained unknown, and this could also explain why same cgST may have unpredictable phenotype.

Methods

E. coli isolation from UTI patients and sequencing

From First Affiliated Hospital of Kunming Medical College, middle stage urine samples from within 2 hours post micturition were collected and 100 μL were plated on 90 mm MacConkey Agar (Antu Biotechnology, Zhengzhou), 37°C for 8 hours. 1000 single colonies and more led to diagnosis of UTI patients whose samples were included. Plates were transferred under -4°C to laboratory of Center of Life Science, Yunnan University, and samples were assigned a unique identifier and subjected to re-cultivation using 20mL LB culture (Solarbio, Beijing, PM0011-500g), 37 °C over night under 220 rpm. To ensure long-term viability, 700 μL resultant bacterial cultures were combined with an equal volume of glycerol (Sangon Biotech, Shanghai, A600232-0500) and stored under appropriate conditions.

A 10μL aliquot of the culture was aseptically sampled using a sterile loop and inoculated into 5mL LB/MHB (Solarbio, Beijing, M8556) culture, followed by incubation with agitation at 220 rpm, 37°C over night. Subsequently, 100μL of this enriched culture was transferred to 5 mL fresh LB/MHB for subculture, shaked under 225 rpm, 37°C for 4 h. Serial dilution was then performed on the subcultured samples, achieving a 10^{-3} dilution factor, and the diluted cultures were plated onto 90 mm LB plates for 37°C over night.

E. coli Colonies were picked according to morphology and selected colonies were further propagated in 5mL LB/MHB to enrich cell populations for downstream analysis. Genomic DNA was extracted from these enriched cultures using a commercial DNA extraction kit (Beyotime Biotechnology, Shanghai, B518225), adhering strictly to the manufacturer's protocols to ensure high-quality DNA for sequencing purposes.

Species identification was performed using full length 16S ribosomal RNA gene sequencing. Libraries were constructed for identified *E. coli* strains to perform paired-end 150 bp sequencing using NovaSeq 6000 (Major Biotechnology, Shanghai).

Minimum Inhibitory Concentration (MIC) determination

MIC values are defined as the lowest concentration that results in a more than 90% reduction in bacterial growth, or the absence of well-defined bacterial spheres. 50μL MHB was added to the 96-well plate to dilute antibiotics, then 50μL ($1x10^5$CFU/mL) bacterial suspension was added to each well, mixed, and incubated at 37°C for 18h. Bacterial growth is determined by measuring the optical density at a wavelength of 600nm (OD600) (Thermofisher, Shanghai, A51119600). MIC test of this study included Kanamycin (BBI, Shanghai, A600286), Fosfomycin (Northeast Pharmaceutical Group, Liaoning, 4210509), Cephalothin (Solarbio, Beijing, C8500), Nitrofurantoin (Tianjin Lisheng Pharmaceutical, Tianjin, 2106015).

Mouse bladder infection

Individual colonies of strains were propagated in LB medium, followed by inoculation into fresh LB medium at a dilution ratio of 1:1000. This culture was incubated at 37°C for 16 hours to ensure optimal growth. Subsequently, the cultures were centrifuged at 4°C and 5000g for 5 minutes, with the bacterial pellet retained. The pellet was re-suspended in a suitable volume of sterile phosphate-buffered saline (Biosharp, Hefei, BL601A) to achieve the desired concentration for infection and stored at 4°C until further use.

C57BL/6J mice, specific pathogen-free (SPF) grade, females aged 7-9 weeks, were procured from the Experimental Animal Center of Yunnan University. A UTI model was established by adapting the 30-G subcutaneous injection needle with a PE10 catheter (BD Biosciences, 427401). Mice were anesthetized with isoflurane (RWD, Shenzhen, R510-22/20220304), and a 50 μL inoculum (approximately 10^8 CFU) was introduced into the bladder via the urethra using the modified injection apparatus, as per the methodology outlined by Hung et al.[40].

At 6 hours post-infection, mice were euthanized, and their bladders were harvested and fixed in 4% paraformaldehyde (PFA) (Sigma Aldrich, Shanghai, 441244-3KG) overnight. The tissues were then dehydrated, embedded in paraffin (Citotest, Jiangsu, 80200-0016), and sectioned for histological analysis. Hematoxylin and eosin (H&E) (Servicebio, Wuhan, G1005-1/G1005-2) staining was performed on the sections to facilitate the pathological assessment. Number of Intracellular Bacterial Communities (IBCs) per strain could generate was average by counting the IBCs in H&E-stained sections over 4 mouse and 2 sections each. Immune score evaluation was conducted according to the scoring criteria established by Hopkins et al.[41]. Edema index quantified using the ratio of bladder weight (mg) to body weight (g).

PCR target gene detection

Frozen stored strains YN-24, YN-9, YN-91, YN-16, YN-54, YN-66, YN-57, YN-35, YN-48, and YN-89, were streaked with LB. Incubate strains on plates at 37°C for 12 hours to promote growth. Following incubation, isolate a single colony from each plate and inoculate it into 5 mL of LB broth. Then, incubate these cultures at 37°C with a shaking speed of 220 rpm for 10 hours to ensure optimal bacterial growth.

For genomic DNA extraction, utilize TIANGEN's Bacterial DNA Kit (DP302-02, Beijing) to purify the total bacterial DNA from the cultivated strains. Adjust the concentration of the purified DNA to 10 ng/μL for subsequent polymerase chain reaction (PCR) analyses. Prepare the PCR reaction mix as follows: 5μL of 2x Rapid Taq Master Mix (Vazyme, Nanjing, P222-03), 3.2μL of ddH$_2$O, 1μL of template DNA, and 0.8μL of the designated primers. The PCR amplification protocol consists of an initial denaturation at 95°C for 3 minutes, followed by 30 cycles of denaturation at 95°C for 15 seconds, annealing at 60°C for 15 seconds, and extension at 72°C for 30 seconds, concluding with a final extension step at 72°C for 5 minutes. The specific primers used in this study are as follows: for *espX4*, forward primer CATGGTGGGTGAAGGA and reverse primer CGATTAGCTCGCCACTC; for *espL1*, forward primer CGATGCTGTCATACTGTG and reverse primer GAGAGGAACCGTGACA.

Bioinformatics

Sequencing results of lab isolated strains and downloaded SRA (Sequence Read Archive) reads, were quality controlled using fastp (v0.23.2)[42], where parameters were adjusted according to read length and adapter length. Including read pairs with enough quality, reads with only one direction passed the filter were also included in the assembly construction using SPAdes (v3.15.5)[43].

Bulk assemblies data downloading for SAC construction was performed using customized workflow blast_at_local_computer[44], which transformed GCA accession into rsync address using Biopython (v1.83)[45] and downloaded data using rsync from NCBI assembly database. The process included automated md5sum check and re-download of failed ones. The downloaded assemblies were primarily quality checked using QUAST (v5.2.0)[46] to filter those with N50 < 200,000 bp. The phenotype dataset were selected from BV-BRC[36] and assemblies were downloaded. Samples whose assembly not available from BV-BRC were obtained from SRA using sratoolkit(v3.0.5)[47].

Downloaded assemblies were randomly selected for construction of Dataset I and II. These selected assemblies were quality checked using checkM (v1.2.2)[48] Taxonomic-specific Workflow, in which marker file were generated for each species and then the completeness and contamination rate were assessed. Assemblies with completeness <95% and contamination rate >5% were excluded from construction of Dataset I and II.

All assemblies were re-annotated using Prokka (v1.14.6)[49] under fast mode which only detect genes without function annotation. The annotated DNA sequences including coding region, tRNA and non-coding RNA sequences were further clustered by customized pipeline OrthoSLC (v0.2.4). It utilize 100% similarity threshold pre-clustering and (Python or C++ built in) hash algorithm for binning to highly accelerate the all-vs-all blast, RBBH pair screening, reciprocal best finding and single linkage clustering. This allowed the gene cluster construction of 801 *L. monocytogenes* and 1200 *E. coli* assemblies could be finished within 25 and 320 minutes using 36 threads, respectively. The RBBH pair was screened using a limit of length difference >0.9, which means the shorter sequence in the pair shall not be shorter than 90% of the longer one. The clusters with exact 1 gene from each assembly were defined as StCGCs.

Multi-sequence alignment on the StCGCs were performed using Kalign (v3.3.5)[50] default mode and aligned clusters were later concatenated into core genome of each strain using customized script. The phylogeny computation utilized RAxML (v8.0.2)[51] under GTRCAT mode. 1000 rapid bootstraps were performed on the phylogenetic tree in Figure 1 while all other trees included in the study performed only 200 rapid bootstraps. The cgST identification was assisted by reference assemblies and followed 'one node one cgST' rule. All strains classified as a certain cgST should share a common node in the phylogeny structure and cgSTs could be identified from a given cgST using sub-nodes.

The VFs annotation on Dataset I, II and SAC were achieved by blastn (v2.13+) VFDB (Feb 10 2023) DNA sequences core dataset (setA)[27] against assemblies. Every best hit in the results were followed by screening that the aligned region should reach >80% similarity with reference and length difference limit between hit region and reference should be >0.8. For 91 patient isolated *E. coli*, the hit regions on assemblies were further tested coverage using bowtie2 (v2.5.0)[52]. The ARGs annotation on 137 *E. coli* in Figure 1 and Dataset II were achieved by inputing assemblies into Card (3.0.2) and Resistance Gene Identifier (RGI) (v6.0.2)[28]. And gene resistance mechanism for each gene was obtained by RGI heatmap generation

function. ARGs annotation on SAC utilized the ARGs found in Dataset II of each species, and blastn (v2.13+)[53] these
genes against each assembly, with same >%80 similarity and >0.8 length difference degree filtering.

Function annotation on each sequence of StCGCs was performed using eggNOG-mapper (v2.1.12) and eggdb (v5.0.2)[30,31],
in which DIAMOND (v2.0.11.149) was employed for searching. The function annotation of each cluster was summarized
and heterofunctional cluster was defined as cluster containing at least 2 kinds of function, where the function with least
sequences in the cluster shall have more than HE_{limit} sequences let a cluster become heterofunctional. Where:

$$HE_{\text{limit}} = \frac{0.5 \cdot S_{\text{count}}}{C_{\text{count}}}$$

S_{count} and C_{count} stand for the count of strains and count of cgSTs, in Dataset II of given species, respectively. This
set up is dependent on the even distribution of samples in cgSTs, so that if a StCGC contained a different function with
amount of sequences more than half of one cgST strain amount, the determination of one of the cgSTs had possibility to
be decided by this hterofunctional cluster. In other words, if a StCGC of *E. coli* ($S_{\text{count}} = 1200$, $C_{\text{count}} = 6$, $HE_{\text{limit}} = $
100) contained only 10 genes that were of different function from other 1190 sequences, this cluster could not differ a cgST
from other simply by its function composition.

The Cluster Variation Level (CVL) was calculated on each StCGC:

$$\text{CVL} = \log_{10} \left(\sum_{i,j}^{n} D_{ij} \cdot (\text{C}_i + \text{C}_j) + 1 \right), (i \neq j)$$

where n stands for all pairwise combinations of dereplicated unique sequence variants in a give cluster; D_{ij} stands for the
count of different sites in the alignment of unique sequence i and j; C_i and C_i stand for the amount of unique sequence i
and j in the cluster. The algorithm attempts to value the variation level within a cluster and clusters were then ranked
according to their CVL value for input window design.

To extract core genes of StCGC (Figure 4) and pan-genes (presence frequency >5% and <100% in Dataset II of *E. coli*,
Supplementary Fig 14 second column) from SAC, the most frequent sequence variant in each of gene cluster of Dataset
II were deemed as seeds. After blastn (v2.13+)[53] these seeds against SAC assemblies, each best hit in an assembly with
>0.9 and >0.8 length difference degree were accepted for core genes and pan-genes, respectively.

Gene extracted from SAC were grouped using Python3 built in hash algorithm for dereplication and sequence variant
summary. All sequences extracted was checked using customized scripts to ensure no hash collision occurred. The sequence
variant summary was later used for sequence presence, type prediction labels.

k-mer frequency was extracted using jellyfish (v2.3.0)[54]. The corrected frequency of a single k-mer was obtained by re-
scaling the frequency to a range between 0 and 1, by reduction of the lowest frequency of this k-mer among all samples
and division by the highest frequency after reduction.

Task parallelrization was achieved through customized Python3 scripts and GNU parallel[55] for efficiency.

Machine learning

LR based cross validation, which was applied in cgST prediction using k-mer, VFs pattern, and ARGs pattern as input,
was performed in R (v4.3.1)[56] with glmnet (v4.1-8)[57] package. Each re-train started with randomly selected training set
(70% of each cgST). An extra re-train would be added if a re-train did not reach convergence until wanted count of re-train
was met. Final average AUROC and average weight of each input feature were calculated to evaluate the performance
and to pick features significant for the prediction.

The RF model used in pseudo-labeling based dataset construction was R randomforest (v4.7-1.1) package[58], where the
number of trees used was 2 fold of the input dimension. The trained models for pseudo-label prediction was tested to have
>98% test set (30% samples from each cgST of Dataset I) performance before actual prediction.

The CNN models were constructed using Pytorch (v2.0.0)[59] and DNA sequences were transformed into usable matrix with
a one-hot like encoding manner as in Supplementary Fig 7. Input DNA sequences were first transformed using a one-hot
like manner. The transformation put IPUAC DNA ambiguity into considerations. Model A were for cgST prediction
which included 4 convolution layers and 4 maxpooling layers and only the first maxpooling and convolution layers had 6

channels. Model A had a deeper structure than model B due to different input size. Model A took complete core genome
size as input (~346,000 bp to 890,000 bp), therefore, we designed a deeper structure for feature extraction and dimension
reduction. While model B took partial genome (~14,000 bp to 27,000 bp) as input to predict accessory gene presence
and sequence variant of other genes. Model B had only 2 convolution layers and 2 maxpooling layers and the amount of
channels for first convolution and maxpooling layers was equal to the count of class (sequence variants) to be predicted.
The channel count in model B was set to 6 in phenotype prediction.

The evaluation of cgST and phenotype prediction using different input combinations was through 9 re-trains, containing
3 fold cross validation and 3 random initialization each fold. Each fold contained nearly equal amount of samples and
strains of each class were evenly distributed in 3 folds. The evaluation of accessory gene presence and sequence variant
prediction was evaluated using 4 re-trains, 2 fold and 2 random initialization each fold. In sequence prediction, the gene to
be predicted had at least 5,000 presence and absence in SAC, each fold contained 2,500 samples with target genes present
in their assemblies and another 2,500 samples without. In sequence variant prediction, assemblies owning the exactly
identical sequences will be grouped as owning the same sequence variant, equivalent to clustering of 100% similarity.
Genes whose sequence variant was predicted had at least 2 prevalent sequence variants that had at least 2,000 presence
in SAC. The prediction would include a class 'other' that containing all non-prevalent sequence variants, and if this class
'other' also had >2,000 presence in SAC. Each of the 2 folds of cross validation contained 1,000 samples of each sequence
variants. The performance evaluation of binary employed AUROC and multiple classification employed accuracy, which
was calculated as the correct predictions out of all predictions since even class count was ensured.

The CNN models employed the learning rate of 3×10^{-3} and 5×10^{-3} with 150 epoches in prediction using model A and B,
respectively. The batch size design for training was 200 in cgST prediction, 2,000 in accessory gene presence prediction,
and one fourth of training set size in sequence variant prediction (to adapt 1,000 more samples every 1 additional class).
For phenotype prediction, the learning rate was 1×10^{-3} with batch size of 120 for 600 epoches. To select the best model
among training epoches, the epoche with lowest test set loss was chosen for prediction of cgST, accessory gene presence,
and sequence variant prediction; and in phenotype prediction, the epoche with highest AUROC was selected. Notably,
train set performance of each epoch was represented by the first batch before gradient descent. Loss function of Cross
Entropy Loss (CrossEntropyLoss) and Binary Cross Entropy Loss (BCELoss) were employed for binary and multi-class
classifications, respectively.

The embedding was performed using two separately trained model for input of 1 ~ 6 (CVL rank) and 201 ~ 218, respectively.
Structure of model B was trained to predict StCGC with ID 10207-13847 in *E. coli* (Supplementary Data 7), which had
10 prevalent sequence variants. The pre-trained model was the on with lowest loss over 150 training epoches, and both
input of 1 ~ 6 and 201 ~ 218 were padded to the same length to use same model hyper-parameters. Using the pre-trained
models, all input were pre-calculated until the final layer to achieve DNA to vector (DNA2vec) embedding. In pan-genes
status predictions, these vectors became the input and only the final full connection layer was trained to predict each
gene.

For re-prediction based evaluation of cgST prediction accuracy using core gene combinations, pre-trained model A and
res-net like model were generated using 70% randomly selected samples from each cgST. Res-net like model starts with 2
convolutional layers for a primary dimension reduction and included ReLu and batch normalization which was not included
in model A. Both models were trained for 500 epoches and learning rate of 3×10^{-3}. While model A took batch size of 210,
res-net like model took batch size of 80 and a L2 regularization of 0.1 to entourage the learning of more features.10 Models
of model A and res-net, respectively, with >98% test set performance were kept for re-prediction. The re-prediction input
included both training and test set samples used in pre-trained model generation.

Statistics

505 Principle Coordinate Analysis (PCoA) was performed using R vegan (2.6-4)[60] package for VFs pattern, ARGs pattern
506 and k-mer pattern within cgST analysis. Significance test pairwise among groups were performed using PERMANOVA
507 algorithm with 999 permutations. Data visualization in this study was facilitated by R ggplot2 (v3.4.3)[61], ggtree (v3.10.0)[62]
508 packages.

Software, workflow management

510 The pipeline and softwares were managed using Docker (v>4.16).

Code availability

Bulk assembly downloading pipeline is available at https://github.com/xchuam/blast_at_local_computer; Gene clustering pipeline OrthoSLC (v0.2.4) is available at https://github.com/JJChenCharly/OrthoSLC; All machine learning, bioinformatics and partial visualization codes are available at https://github.com/JJChenCharly/g2g.

Data availability

The Data will be made available upon request.

Declaration of Competing Interest

The authors declare no competing interests.

Ethnicity statement

All animal experiments described in this study were conducted in accordance with the ethical standards and guidelines for the care and use of laboratory animals at the Animal Center of Yunnan University. Experiments were carried out under the supervision of Laboratory Animal Welfare Ethics Committee, Yunnan University (Ethics review number: YNU20220176). All procedures involving the collection and analysis of strains from UTI patients were carried out in compliance with the ethical standards of the institutional and/or national research committee. The study protocols were approved by the Ethics Committee of The First Affiliated Hospital of Kunming Medical College (2022L203) and The Committee on Human Subject Research and Ethics, Yunnan University (CHSRE number: CHSRE2022030). Informed consent was obtained from all individual participants included in the study.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (32070818 to C.M.G.), Natural Science Foundation of Yunnan Province of China [202001BB050005], the Xingdian talent support program of Yunnan Province. We thank Songnian Hu (University of Chinese Academy of Science) for early stage conceptualization support; Cheng Peng (Yunnan University) for bioinformatics and computation hardware support; Longyu Chen and Zirui Wang (University of Chinese Academy of Science) for machine learning consultation and computation power support.

534 References

1. Callister, S. J. *et al.* Comparative Bacterial Proteomics: Analysis of the Core Genome Concept. *PLOS ONE* **3**, e1542 (2008 2 6).

2. Schürch, A. C., Arredondo-Alonso, S., Willems, R. J. L. & Goering, R. V. Whole genome sequencing options for bacterial strain typing and epidemiologic analysis based on single nucleotide polymorphism versus gene-by-gene–based approaches. *Clinical Microbiology and Infection* **24**, 350–354 (2018).

3. Schreiber, H. L. *et al.* Bacterial virulence phenotypes of Escherichia coli and host susceptibility determine risk for urinary tract infections. *Science Translational Medicine* **9**, eaaf1283 (2017).

4. McInerney, J. O., McNally, A. & O'Connell, M. J. Why prokaryotes have pangenomes. *Nat Microbiol* **2**, 1–5 (2017).

5. Soucy, S. M., Huang, J. & Gogarten, J. P. Horizontal gene transfer: Building the web of life. *Nat Rev Genet* **16**, 472–482 (2015).

6. Robicsek, A., Jacoby, G. A. & Hooper, D. C. The worldwide emergence of plasmid-mediated quinolone resistance. *The Lancet Infectious Diseases* **6**, 629–640 (2006).

7. Rankin, D. J., Rocha, E. P. C. & Brown, S. P. What traits are carried on mobile genetic elements, and why? *Heredity* **106**, 1–10 (2011).

8. Croll, D. & McDonald, B. A. The Accessory Genome as a Cradle for Adaptive Evolution in Pathogens. *PLOS Pathogens* **8**, e1002608 (2012 4 26).

9. Moradigaravand, D. *et al.* Prediction of antibiotic resistance in Escherichia coli from large-scale pan-genome data. *PLOS Computational Biology* **14**, e1006258 (2018 12 14).

10. Hall, R. J. *et al.* Gene-gene relationships in an Escherichia coli accessory genome are linked to function and mobility. *Microbial Genomics* **7**, (2021).

11. Whelan, F. J., Hall, R. J. & McInerney, J. O. Evidence for Selection in the Abundant Accessory Gene Content of a Prokaryote Pangenome. *Molecular Biology and Evolution* **38**, 3697–3708 (2021).

12. Beavan, A. J. S., Domingo-Sananes, M. R. & McInerney, J. O. Contingency, repeatability, and predictability in the evolution of a prokaryotic pangenome. *Proceedings of the National Academy of Sciences* **121**, e2304934120 (2024).

13. Moura, A. *et al.* Whole genome-based population biology and epidemiological surveillance of Listeria monocytogenes. *Nat Microbiol* **2**, 1–10 (2016).

14. Maury, M. M. *et al.* Uncovering Listeria monocytogenes hypervirulence by harnessing its biodiversity. *Nat Genet* **48**, 308–313 (2016).

15. McNally, A. *et al.* Combined Analysis of Variation in Core, Accessory and Regulatory Genome Regions Provides a Super-Resolution View into the Evolution of Bacterial Populations. *PLOS Genetics* **12**, e1006280 (2016 9 12).

16. Freschi, L. *et al.* The Pseudomonas aeruginosa Pan-Genome Provides New Insights on Its Population Structure, Horizontal Gene Transfer, and Pathogenicity. *Genome Biology and Evolution* **11**, 109–120 (2019).

17. Maury, M. M. *et al.* Hypervirulent Listeria monocytogenes clones' adaption to mammalian gut accounts for their association with dairy products. *Nat Commun* **10**, 2488 (2019).

18. Jacquet, C. *et al.* A Molecular Marker for Evaluating the Pathogenic Potential of Foodborne Listeria monocytogenes. *The Journal of Infectious Diseases* **189**, 2094–2100 (2004).

19. Lecuit, M. *et al.* A Transgenic Model for Listeriosis: Role of Internalin in Crossing the Intestinal Barrier. *Science* **292**, 1722–1725 (2001).

20. Disson, O. *et al.* Conjugated action of two species-specific invasion proteins for fetoplacental listeriosis. *Nature* **455**, 1114–1118 (2008).

21. Ortiz, S., López-Alonso, V., Rodríguez, P. & Martínez-Suárez, J. V. The Connection between Persistent, Disinfectant-Resistant Listeria monocytogenes Strains from Two Geographically Separate Iberian Pork Processing Plants: Evidence from Comparative Genome Analysis. *Applied and Environmental Microbiology* **82**, 308–317 (2016).

22. Rijavec, M. *et al.* High Prevalence of Multidrug Resistance and Random Distribution of Mobile Genetic Elements Among Uropathogenic Escherichia coli (UPEC) of the Four Major Phylogenetic Groups. *Curr Microbiol* **53**, 158–162 (2006).

23. Starčič Erjavec, M., Rijavec, M., Križan-Hergouth, V., Fruth, A. & Žgur-Bertok, D. Chloramphenicol- and tetracycline-resistant uropathogenic Escherichia coli (UPEC) exhibit reduced virulence potential. *International Journal of Antimicrobial Agents* **30**, 436–442 (2007).

24. Piatti, G., Mannini, A., Balistreri, M. & Schito, A. M. Virulence Factors in Urinary Escherichia coli Strains: Phylogenetic Background and Quinolone and Fluoroquinolone Resistance. *Journal of Clinical Microbiology* **46**, 480–487 (2008).

25. Skjøt-Rasmussen, L. *et al.* Persisting clones of Escherichia coli isolates from recurrent urinary tract infection in men and women. *Journal of Medical Microbiology* **60**, 550–554 (2011).

26. Ejrnæs, K. *et al.* Characteristics of Escherichia coli causing persistence or relapse of urinary tract infections: Phylogenetic groups, virulence factors and biofilm formation. *Virulence* **2**, 528–537 (2011).

27. Liu, B., Zheng, D., Zhou, S., Chen, L. & Yang, J. VFDB 2022: A general classification scheme for bacterial virulence factors. *Nucleic Acids Research* **50**, D912–D917 (2022).

28. Alcock, B. P. *et al.* CARD 2020: Antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Research* **48**, D517–D525 (2020).

29. Glorot, X. & Bengio, Y. Understanding the difficulty of training deep feedforward neural networks. in *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics* 249–256 (JMLR Workshop and Conference Proceedings, 2010).

30. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Molecular Biology and Evolution* **38**, 5825–5829 (2021).

31. Huerta-Cepas, J. *et al.* eggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research* **47**, D309–D314 (2019).

32. Davies, T. J. *et al.* Reconciling the Potentially Irreconcilable? Genotypic and Phenotypic Amoxicillin-Clavulanate Resistance in Escherichia coli. *Antimicrobial Agents and Chemotherapy* **64**, 10.1128/aac.02026–19 (2020).

33. Bradley, P. *et al.* Rapid antibiotic-resistance predictions from genome sequence data for Staphylococcus aureus and Mycobacterium tuberculosis. *Nat Commun* **6**, 10063 (2015).

34. Kumar, N. *et al.* Evaluation of a fully automated bioinformatics tool to predict antibiotic resistance from MRSA genomes. *Journal of Antimicrobial Chemotherapy* **75**, 1117–1122 (2020).

35. Coll, F. *et al.* Genome-wide analysis of multi- and extensively drug-resistant Mycobacterium tuberculosis. *Nat Genet* **50**, 307–316 (2018).

36. Olson, R. D. *et al.* Introducing the Bacterial and Viral Bioinformatics Resource Center (BV-BRC): A resource combining PATRIC, IRD and ViPR. *Nucleic Acids Res* **51**, D678–D689 (2023).

37. Wattam, A. R. *et al.* Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* **45**, D535–D542 (2017).

38. Salichos, L. & Rokas, A. Evaluating Ortholog Prediction Algorithms in a Yeast Model Clade. *PLOS ONE* **6**, e18755 (2011 4 13).

39. He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition. in 770–778 (2016).

40. Hung, C.-S., Dodson, K. W. & Hultgren, S. J. A murine model of urinary tract infection. *Nat Protoc* **4**, 1230–1243 (2009).

41. Hopkins, W. J., Gendron-Fitzpatrick, A., Balish, E. & Uehling, D. T. Time Course and Host Responses to Escherichia coli Urinary Tract Infection in Genetically Distinct Mouse Strains. *Infection and Immunity* **66**, 2798–2802 (1998).

42. Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

43. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

44. Ma, X., Chen, J., Zwietering, M. H., Abee, T. & Den Besten, H. M. W. Stress resistant *rpsU* variants of *Listeria Monocytogenes* can become underrepresented due to enrichment bias. *International Journal of Food Microbiology* 110680 (2024) doi:10.1016/j.ijfoodmicro.2024.110680.

45. Cock, P. J. A. *et al.* Biopython. *Bioinformatics* **25**, 1422–1423 (2009).

46. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: Quality assessment tool for genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

47. Leinonen, R., Sugawara, H., Shumway, M. & on behalf of the International Nucleotide Sequence Database Collaboration. The Sequence Read Archive. *Nucleic Acids Research* **39**, D19–D21 (2011).

48. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

49. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).

50. Lassmann, T. & Sonnhammer, E. L. Kalign – an accurate and fast multiple sequence alignment algorithm. *BMC Bioinformatics* **6**, 298 (2005).

51. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

52. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357–359 (2012).

53. Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

54.   Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* **27**, 764–770 (2011).

55.   Tange, O. Gnu parallel-the command-line power tool. *Usenix Mag* 36(1), 42.

56.   Ihaka, R. & Gentleman, R. R: A Language for Data Analysis and Graphics. *Journal of Computational and Graphical Statistics* **5**, 299–314 (1996).

57.   Tay, J. K., Narasimhan, B. & Hastie, T. Elastic Net Regularization Paths for All Generalized Linear Models. *Journal of Statistical Software* **106**, 1–31 (2023).

58.   Breiman, L. Random Forests. *Machine Learning* **45**, 5–32 (2001).

59.   Paszke, A. *et al.* PyTorch: An Imperative Style, High-Performance Deep Learning Library. in *Advances in Neural Information Processing Systems* vol. 32 (Curran Associates, Inc., 2019).

60.   J, O. Vegan : Community ecology package. *http://vegan.r-forge.r-project.org/* (2010).

61.   Wilkinson, L. Ggplot2: Elegant Graphics for Data Analysis by WICKHAM, H. *Biometrics* **67**, 678–679 (2011).

62.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. Ggtree: An r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* **8**, 28–36 (2017).

Supplementary Material - Intra-genomic genes-to-genes correlation enables genome representation

Jingjie Chen[1,9]   Xuchuan Ma[2,9]   Junwei He[1]   Yingxia Wang[3]   Yuji Ren[4]

Li Qi[7]   Liuyang Song[7]   Lin Ruan[1]   Cun Fan[1]   Jun Jiang[1]

Yongjia Bao[1]   Tao Huang[1]   Jingbo Cheng[1]   Xing Liu[5]   Fang Chen[6]

Yichen Huang[6]   Na Zhao[7,8,*]   Chunming Guo[1,*]

[1] Yunnan Key Laboratory of Cell Metabolism and Disease, Center for Life Sciences, School of Life Sciences, Yunnan University, Kunming 650500, China.
[2] Food Microbiology, Wageningen University & Research, Wageningen, the Netherlands.
[3] The Department of Pathology, First affiliated Hospital of Kunming Medical University, Kuming, People's Republic of China.
[4] The Department of Clinical Laboratory, First affiliated Hospital of Kunming Medical University, Kuming, People's Republic of China.
[5] Department of Urology, Children's Hospital of Chongqing Medical University, 136 Zhongshan Road, Chongqing 400014, PR China.
[6] Department of Urology, Shanghai Children's Hospital, School of medicine, Shanghai Jiao Tong University.
[7] Key Laboratory in Software Engineering of Yunnan Province, Yunnan University, Kunming, People's Republic of China.
[8] Big Data Research Center, University of Electronic Science and Technology of China, Chengdu, People's Republic of China.
[9] These authors contributed equally.

[*] Correspondence: Na Zhao <zhaona@ynu.edu.cn>, Chunming Guo <chunmingguo@ynu.edu.cn>

Supplementary Tables

Supplementary Table 1: **46 reference *E. coli* genome information.** The genome included the circular chromosome and plasmids. These reference strains were used for cgST identification in Fig 1 and 2 for *E. coli* dataset.

Strain	Pathotype	cgST	BioProject(NCBI ID)
536	UPEC	B2	PRJNA16235
11128	EHEC	B1	PRJDA32513
11368	EHEC	B1	PRJDA32509
12009	EHEC	B1	PRJDA32511
55989	EAEC	B1	PRJNA33413
2009EL-2050	EHEC	B1	PRJNA81097
2009EL-2071	EHEC	B1	PRJNA81097
2011C-3493	EHEC	B1	PRJNA81095
ABU_83972	ABU	B2	PRJNA38725
APEC_O1	APEC	B2	PRJNA16718
APEC_O78	APEC	B1	PRJNA184588
ATCC_8739	Commensal	A	PRJNA18083
BL21(DE3)	Lab_Strain	A	PRJNA20713
BW2952	Lab_Strain	A	PRJNA33775
CB9615	EPEC	E	PRJNA42729
CE10	NMEC	D	PRJNA63597
CFT073	UPEC	B2	PRJNA313
clone_D_i14	UPEC	B2	PRJNA52023
clone_D_i2	UPEC	B2	PRJNA52021
DH1	Lab_Strain	A	PRJDA52077
DH10B	Lab_Strain	A	PRJNA20079
E2348_69	EPEC	B2	PRJEA32571
E24377A	ETEC	B1	PRJNA13960
EC4115	EHEC	E	PRJNA27739
ED1a	Commensal	B2	PRJNA33409
EDL933	EHEC	E	PRJNA259
H10407	ETEC	A	PRJEA42749
HS	Commensal	A	PRJNA13959
IAI39	UPEC	D	PRJNA33411
IHE3034	NMEC	B2	PRJNA43693
K12_MG1655	Commensal	A	PRJNA40075
LF82	AIEC	B2	PRJNA33825
NRG_857C	AIEC	B2	PRJNA41221
REL606	Lab_Strain	A	PRJNA18281
RM12579	EPEC	E	PRJNA68245
S88	Commensal	B2	PRJNA33375
SE11	Commensal	B1	PRJNA18057
SE15	Commensal	B2	PRJDA19053
SMS-3-5	Environmental	D	PRJNA19469
TW14359	EHEC	E	PRJNA30045
UM146	AIEC	B2	PRJNA50883
UMNK88	ETEC	A	PRJNA42137
UTI89	UPEC	B2	PRJNA16259
W	Lab_Strain	B1	PRJNA48011
W3110	Lab_Strain	A	PRJNA16351
Xuzhou21	EHEC	E	PRJNA45823

Supplementary Table 2: **Gene name and VFDB[1] query correspondence.** *espL1-C* and *espL1-E* stands for *espL1* gene from *E. coli* CB9615 and EDL933, respectively. The query accession could be used to search relavent information (related literature, ortholog, mechanisms) from VFDB.

gene	VFDB query
cfaD/cfaE	VFG034648(gb|WP_001245988)
shuY	VFG013068(gb|WP_000189360)
shuA	VFG013064(gb|WP_000089574)
shuS	VFG013063(gb|WP_001017208)
chuT	VFG012603(gb|WP_001302516)
chuX	VFG000920(gb|WP_000020034)
chuS	VFG000916(gb|WP_001017192)
espX4	VFG034866(gb|WP_000900547)

gene	VFDB query
espL1-C	VFG034756
espL1-E	VFG034754(gb|WP_000081129)

Supplementary Table 3: cgST reference strains. Strains selected in Dataset I to be cgST identification reference in phylogeny in Dataset II of *E. coli.*

ID in Dataset II	GCA	Description
EC_F_ref0	GCA_014794105.1	GCA_014794105.1_PDT000848429.1_genomic
EC_F_ref1	GCA_000459215.1	GCA_000459215.1_Esch_coli_HVH_208_4-3112292_V1_genomic
EC_F_ref2	GCA_021425225.1	GCA_021425225.1_PDT001218343.1_genomic
EC_F_ref3	GCA_021085865.1	GCA_021085865.1_PDT001197641.1_genomic
EC_F_ref4	GCA_019204805.1	GCA_019204805.1_PDT001088470.1_genomic

Supplementary Table 4: cgST reference strains. Strains selected in Dataset I to be cgST identification reference in phylogeny in Dataset II of *L. monocytogenes.*

ID in Dataset II	GCA or GI	Description
LM_A_ref0	GCA_020988225.1	GCA_020988225.1_PDT001185688.1_genomic
LM_A_ref1	GCA_003685255.1	GCA_003685255.1_PDT000365297.1_genomic
LM_A_ref2	GCA_004598375.1	GCA_004598375.1_PDT000218577.2_genomic
LM_A_ref3	GCA_009449765.1	GCA_009449765.1_PDT000421349.2_genomic
LM_A_ref4	GCA_024119015.1	GCA_024119015.1_PDT001338036.3_genomic
LM_B_ref0	GCA_004484765.1	GCA_004484765.1_PDT000065844.2_genomic
LM_B_ref1	GCA_018225825.1	GCA_018225825.1_PDT001017255.1_genomic
LM_B_ref2	GCA_028162645.1	GCA_028162645.1_PDT001589962.1_genomic
LM_B_ref3	GCA_023538895.1	GCA_023538895.1_PDT001314493.1_genomic
LM_B_ref4	GCA_009490045.1	GCA_009490045.1_PDT000454753.2_genomic
LM_C_ref0	GCA_020757795.1	GCA_020757795.1_PDT001168515.1_genomic
LM_C_ref1	GCA_000307065.1	GCA_000307065.1_ASM30706v1_genomic
LM_C_ref2	GCA_010080955.1	GCA_010080955.1_PDT000626992.1_genomic
LM_C_ref3	GCA_004472405.1	GCA_004472405.1_PDT000447650.1_genomic
LM_C_ref4	GCA_009361515.1	GCA_009361515.1_ASM936151v1_genomic
10403S	GI: 345535581	cgST D
EGD	GI: 543859294	cgST D
EGD-e	GI: 1796433495	cgST D
LO28	GI: 2173384332	cgST D
Scott_A	GI: 1796436383	cgST B

Supplementary Table 5: cgST reference strains. Strains selected in Dataset I to be cgST identification reference in phylogeny in Dataset II of *C. jejuni.*

ID in Dataset II	GCA	Description
CJ_B_ref0	GCA_018994985.1	GCA_018994985.1_PDT001074550.1_genomic
CJ_B_ref1	GCA_008248845.1	GCA_008248845.1_PDT000573498.1_genomic
CJ_B_ref2	GCA_005373965.1	GCA_005373965.1_PDT000370886.1_genomic
CJ_B_ref3	GCA_017182455.1	GCA_017182455.1_PDT000945415.1_genomic
CJ_B_ref4	GCA_010299745.1	GCA_010299745.1_PDT000563316.1_genomic
CJ_C_ref0	GCA_004822525.1	GCA_004822525.1_PDT000406612.1_genomic
CJ_C_ref1	GCA_004987965.1	GCA_004987965.1_PDT000271318.2_genomic
CJ_C_ref2	GCA_028591385.1	GCA_028591385.1_PDT001612276.1_genomic
CJ_C_ref3	GCA_005316785.1	GCA_005316785.1_PDT000266958.5_genomic
CJ_C_ref4	GCA_009188145.1	GCA_009188145.1_PDT000610439.1_genomic
CJ_D_ref0	GCA_004278335.1	GCA_004278335.1_PDT000384529.1_genomic
CJ_D_ref1	GCA_008170205.1	GCA_008170205.1_PDT000570024.1_genomic
CJ_D_ref2	GCA_025939745.1	GCA_025939745.1_ASM2593974v1_genomic
CJ_D_ref3	GCA_017239835.1	GCA_017239835.1_PDT000939616.1_genomic
CJ_D_ref4	GCA_011737495.1	GCA_011737495.1_PDT000712334.1_genomic
CJ_A_ref0	GCA_005354025.1	GCA_005354025.1_PDT000360883.1_genomic
CJ_A_ref1	GCA_025020425.1	GCA_025020425.1_PDT001409990.1_genomic
CJ_A_ref2	GCA_005156375.1	GCA_005156375.1_PDT000173386.2_genomic

ID in Dataset II	GCA	Description
CJ_A_ref3	GCA_015088835.1	GCA_015088835.1_PDT000856263.1_genomic
CJ_A_ref4	GCA_004903555.1	GCA_004903555.1_PDT000428963.1_genomic

Supplementary Table 6: cgST reference strains. Strains selected in Dataset I to be cgST identification reference in phylogeny in Dataset II of *S. aureus*.

ID in Dataset II	GCA	Description
SA_B_ref0	GCA_019716075.1	GCA_019716075.1_ASM1971607v1_genomic
SA_B_ref1	GCA_903808755.1	GCA_903808755.1_22114_8_114_genomic
SA_B_ref2	GCA_026882685.1	GCA_026882685.1_PDT001525048.1_genomic
SA_B_ref3	GCA_021482505.1	GCA_021482505.1_ASM2148250v1_genomic
SA_B_ref4	GCA_026837635.1	GCA_026837635.1_PDT001520774.1_genomic
SA_A_ref0	GCA_027466985.1	GCA_027466985.1_PDT001546625.1_genomic
SA_A_ref1	GCA_018912765.1	GCA_018912765.1_ASM1891276v1_genomic
SA_A_ref2	GCA_018998385.1	GCA_018998385.1_ASM1899838v1_genomic
SA_A_ref3	GCA_009896805.1	GCA_009896805.1_ASM989680v1_genomic
SA_A_ref4	GCA_029038185.1	GCA_029038185.1_PDT001652831.1_genomic
SA_D_ref0	GCA_026463645.1	GCA_026463645.1_PDT001498828.1_genomic
SA_D_ref1	GCA_028588125.1	GCA_028588125.1_PDT001611913.1_genomic
SA_D_ref2	GCA_026872015.1	GCA_026872015.1_PDT001523982.1_genomic
SA_D_ref3	GCA_018915145.1	GCA_018915145.1_ASM1891514v1_genomic
SA_D_ref4	GCA_020587635.1	GCA_020587635.1_ASM2058763v1_genomic
SA_C_ref0	GCA_024360785.1	GCA_024360785.1_ASM2436078v1_genomic
SA_C_ref1	GCA_027539185.1	GCA_027539185.1_PDT001556321.1_genomic
SA_C_ref2	GCA_027528505.1	GCA_027528505.1_PDT001553608.1_genomic
SA_C_ref3	GCA_020665265.1	GCA_020665265.1_ASM2066526v1_genomic
SA_C_ref4	GCA_018149675.1	GCA_018149675.1_PDT001013994.1_genomic

Supplementary Table 7: Studies and species included from the study. Studies were selected from BV-BRC[2,3] from which assemblies were obtained. Corresponding strain ID and resistance label to each antibiotics are in Supplementary Data 11.

Study	Species included
Davies et al.[4]	*E. coli*
Moradigaravand et al.[5]	*E. coli*
Bradley et al.[6]	*S. aureus* and *M. tuberculosis*
Kumar et al.[7]	*S. aureus*
Coll et al.[8]	*M. tuberculosis*

Supplementary Table 8: House keeping genes. The house keeping genes selected from published datasets. Their names, total length of target regions for *E. coli*[9], *L. monocytogenes*[10], *C. jejuni*[11], *S. aureus*[12], and *M. tuberculosis*[13–15]

Species	house keeping genes	total length
E. coli	*adk, fumC, gyrB, icd, mdh, purA, recA*	3423 bp
L. monocytogenes	*abcZ, bglA, cat, dapE, dat, idh, ihkA*	3288 bp
C. jejuni	*aspA, glnA, gltA, glyA, pgm, tkt, uncA*	3309 bp
S. aureus	*arcC, aroE, glpF, gmk, pta, tpi*	2670 bp
M. tuberculosis	*gyrA, gyrB, katG, purA, recA, rpoB, sodA*	4612 bp

Supplementary Data

Supplementary Data 1: VFs, ARGs annotation and cgST of 46 reference and 91 patient isolated *E. coli*.

cgST	VFs annotation	ARGs annotation
cgST_137_Ec.csv	VFs_137_Ec.csv	ARGs_137_Ec.csv

Supplementary Data 2: VFs and pseudo-cgST-label of SAC (Species Assembly Collection) of 4 species.

Sp.	pseudo-cgST label	VFs
E. coli	pcgST_EC_SAC.csv	VFs_EC_SAC.csv
L. monocytogenes	pcgST_LM_SAC.csv	VFs_LM_SAC.csv
C. jejuni	pcgST_CJ_SAC.csv	VFs_CJ_SAC.csv
S. aureus	pcgST_SA_SAC.csv	VFs_SA_SAC.csv

Supplementary Data 3: VF annotation and cgST of Dataset I and II of 4 species.

Sp.	cgST I	cgST II	VFs I	VFs II
E. coli	cgST_EC1.csv	cgST_EC2.csv	VFs_EC1.csv	VFs_EC2.csv
L. monocytogenes	cgST_LM1.csv	cgST_LM2.csv	VFs_LM1.csv	VFs_LM2.csv
C. jejuni	cgST_CJ1.csv	cgST_CJ2.csv	VFs_CJ1.csv	VFs_CJ2.csv
S. aureus	cgST_SA1.csv	cgST_SA2.csv	VFs_SA1.csv	VFs_SA2.csv

Supplementary Data 4: ARG annotation of Dataset II of 4 species.

Sp.	ARGs II
E. coli	ARGs_EC2.csv
L. monocytogenes	ARGs_LM2.csv
C. jejuni	ARGs_CJ2.csv
S. aureus	ARGs_SA2.csv

Supplementary Data 5: Average weight of 50 re-trains using VFs or ARGs to predict cgST in Dataset II.

Sp.	ARGs weight	VFs weight
E. coli	ARGs_wei_EC.csv	VFs_wei_EC.csv
L. monocytogenes	ARGs_wei_LM.csv	VFs_wei_LM.csv
C. jejuni	ARGs_wei_CJ.csv	VFs_wei_CJ.csv
S. aureus	ARGs_wei_SA.csv	VFs_wei_SA.csv

Supplementary Data 6: VFs name (in Fig3) and VFDB query correspondence; ARGs name (in Fig3) to CARD model name correspondence.

ARGs	VFs
ARGs_CARD.XLSX	VFs_VFDB.XLSX

Supplementary Data 7: CVL result, StCGC cluster parameters, prokka/eggnogmapper annotation result, and the most prevalent sequence variant of StCGCs of Dataset II.

Sp.	ARGs weight
E. coli	CVL_n_annotation_EC2.csv

Sp.	ARGs weight
L. monocytogenes	CVL_n_annotation_LM2.csv
C. jejuni	CVL_n_annotation_CJ2.csv
S. aureus	CVL_n_annotation_SA2.csv

Supplementary Data 8 (Fig 3): KO simulation for cgST recovery performance. EC: *E. coli*, LM: *L. monocytogenes*, CJ: *C. jejuni*, SA: *S. aureus*.

Supplementary Data 9: Accessory gene presnece prediction results. EC: *E. coli*, LM: *L. monocytogenes*, CJ: *C. jejuni*, SA: *S. aureus*.

Species	File
E. coli	presence_EC.xlsx
L. monocytogenes	presence_LM.xlsx
C. jejuni	presence_CJ.xlsx
S. aureus	presence_SA.xlsx

Supplementary Data 10: Gene sequence variant prediction results. EC: *E. coli*, LM: *L. monocytogenes*, CJ: *C. jejuni*, SA: *S. aureus*.

Species	File
E. coli	seqtype_EC.xlsx and seqtype_EC_other_excluded.xlsx
L. monocytogenes	seqtype_LM.xlsx
C. jejuni	seqtype_CJ.xlsx
S. aureus	seqtype_SA.xlsx

Supplementary Data 11: pan-gene status prediction results of *E. coli*. presence and sequence variants prediciton result. The function annotation result includs the sequence of the most prevalent sequence variant and frequency in SAC.

Supplementary Data 12: Phenotype labels for data from this study and published datasets. Phenotype_labels.xlsx.

Supplementary Data 13: Phenotype prediction result. Phenotype_prediction_result.csv.

Supplementary Data 14: *E. coli* 3 to 7-mer frequency. file name format: "input gene CVL rank"_"k-mer length"-mer.csv.

Input CVL rank	3-mer	7-mer
1-6	1-6_3-mer.csv	1-6_7-mer.csv
201-206	201-206_3-mer.csv	201-206_7-mer.csv
400-406	400-406_3-mer.csv	400-406_7-mer.csv
odd	odd_3-mer.csv	odd_7-mer.csv
even	even_3-mer.csv	even_7-mer.csv

Supplementary Figures

Supplementary Fig 1: **Categorization of gene clusters.** | Gene Clusters shared by only a subset of assemblies participated analysis are defined as accessory gene clusters. If a cluster had more number of genes than the number of assemblies participated analysis but all genes in this cluster were found in only a subset of all assemblies, the cluster is defined as accessory. Clusters shared by all assemblies are defined as core clusters. If a core cluster had one gene from each assembly, it is defined as 'strict core'. If one or more assemblies had more than one gene in a core cluster, the cluster is defined as 'surplus core'.

Supplementary Fig 2: VFs annotation of 46 reference and 91 patient isolated *E. coli*. | Plot based on Supplementary Data 1. A fragment owned by strain with similarity against reference lower than 80% was regarded as absence of the corresponding gene in the strain.

Supplementary Fig 3: Prediction of cgST by accessory gene status in *L. monocytogenes*. | cgST result of Dataset I (left) and II (right) of *L. monocytogenes*. The dashed branches were of large length and were manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The tables in the middle described the confusion matrix which did not include the reference strains (known cgST in Dataset I) and cgST distribution of Dataset I and II next to each cgST.

Supplementary Fig 4: Prediction of cgST by accessory gene status in *C. jejuni*. | cgST result of Dataset I (left) and II (right) of *C. jejuni*. The dashed branches were of large length and were manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The tables in the middle described the confusion matrix which did not include the reference strains (known cgST in Dataset I) and cgST distribution of Dataset I and II next to each cgST.

Supplementary Fig 5: Prediction of cgST by accessory gene status in *S. aureus*. | cgST result of Dataset I (left) and II (right) of *S. aureus*. The dashed branches were of large length and were manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The tables in the middle described the confusion matrix which did not include the reference strains (known cgST in Dataset I) and cgST distribution of Dataset I and II next to each cgST.

Supplementary Fig 6: VFs (a) and ARGs (b) specific to cgST were selected by LR based re-trains. | The x axis of each plot are strains in Dataset II and are arranged in the same order as in the phylogeny of Dataset II (Fig 2 and Supplementary Fig 2-4) and with same color. The text on x axis also exhibited the average AUROC of re-trains when using VFs (**a**) or ARGs (**b**) patterns to predict cgST. The color on y axis suggests the VFs category (**a**) and resistance mechanism (B) of genes. VFs (**a**) and its VFDB query correspondence could be found in Supplementary Data 6. ARGs (**b**) with simplifies model names correspond to their full name in CARD database in Supplementary Data 6. In plot B, *marA* with "*" suggests the gene was present in 2 different resistance mechanisms.

Core-to-accessory

Strain	cgST	Homofunctional (GeneX)	accessory (GeneY)
1	A	Function 1.1	**Present**
2	A	Function 1.1	**Present**
3	A	Function 1.1	**Present**
4	B	Function 1.2	Absent
5	B	Function 1.2	Absent
6	B	Function 1.2	Absent

Gene co-occurrence

Strain	cgST	Heterofunctional (GeneX)	accessory (GeneY)
1	A	Function 1	**Present**
2	A	Function 1	**Present**
3	A	Function 1	**Present**
4	B	Function 2	Absent
5	B	Function 2	Absent
6	B	Function 2	Absent

Supplementary Fig 7: Diagram of difference between core-to-accessory relationship and gene co-occurrence. | In the diagram, we show 2 possible cases that accessory genes can correlate with certain cgST. In this simplified example, 6 strains were classified into 2 different cgSTs (A and B) by a singe hypothetical core gene cluster (name as GeneX) shared by all 6 strains. However, the core gene cluster of GeneX, might be of same function genes (left), where all 6 pieces of GeneX are, for example, a kind of enzyme that catalyze the same reaction (Homofunctional). However, GeneX of Function 1.1 from cgST might exhibit better efficiency than the GeneX of Function 1.2. The GeneX from cgST A and B might also come from different function (Heterofunctional, right). In this case, the gene cluster of 6 GeneX could actually be the fusion of two accessory gene cluster. In this case, the specific presence of GeneY in cgST A, is a co-occurrence with GeneX of Function 1.

Supplementary Fig 8: Transform DNA to tensor using one-hot like encoding and applying predictions with CNN model A and B. | Input DNA sequences were first transformed using a one-hot like manner. The transformation put IPUAC DNA ambiguity into considerations. Model A were using for cgST prediction which included 4 convolution layers and 4 maxpooling layers and only the first maxpooling and convolution layers had 6 channels. Model had a deeper structure than model B due input size. Model A took complete core genome size as input (~346,000 bp to 890,000 bp), therefore, a deeper structure for feature extraction and dimension reduction. While model B took partial genome (~14,000 bp to 27,000 bp) as input to predict accessory gene presence and sequence variant of other genes. Model B had only 2 convolution layers and 2 maxpooling layers and the amount of channels for first convolution and maxpooling layers was equal to the count of class (n = count of sequence variants) to be predicted. The channel count in model B was set to 6 in phenotype prediction.

Supplementary Fig 9: Different core gene combinations can accurately recover cgSTs with CNN in cross validations. | (EC: *E. coli*, LM: *L. monocytogenes*, CJ: *C. jejuni*, SA: *S. aureus*). First row: by increasing available genes in the input with increase of ten genes (from the high CVL StCGCs), cgST recovery of >98% could be achieved with less than 20 StCGCs. Second row: By using different window (20 genes each window) of available core genes in prediction, ability to recover complete core genome cgST was tested on a wider range of core genome. Third row: by increasing available genes from the lower end from the CVL list, least needed number to recover cgST with >98% accuracy from the bottom of CVL list is exhibited. Each box represents the performance of nine re-trains (three-fold, three random initialization each fold) of train set (red) and test set (blue). The CNN model adopted model A as in Supplementary Fig 8.

Supplementary Fig 10: *E. coli* **Dataset II phylogeny constructed using subset of StCGCs.** | Left tree used gene 1 ∼ 6 and right tree used gene 1 ∼ 20 in CVL list. Dashed branch was a outlier with too large branch length manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The color for each strain is the same as in Fig 2.

Supplementary Fig 11: *E. coli* **phylogeny constructed using subset of StCGCs.** | Left tree used gene 201 ~ 206 and right tree used gene 201 ~ 220 in CVL list. Dashed branch was a outlier with too large branch length manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The color for each strain is the same as in Fig 2.

Supplementary Fig 12: *E. coli* **phylogeny constructed using subset of StCGCs.** | Left tree used gene 401 ~ 406 and right tree used gene 401 ~ 420 in CVL list. Dashed branch was a outlier with too large branch length manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The color for each strain is the same as in Fig 2.

Supplementary Fig 13: *E. coli* **phylogeny constructed using subset of StCGCs.** | The phylogeny was constructed using last 100 genes (378 ~ 477) in CVL list. Dashed branch was a outlier with too large branch length manually shortened to help visualization. The nodes with bootstrap value ≥ 95 are marked with circles and those with bootstrap value >80 and <95 were marked with triangles. The color for each strain is the same as in Fig 2.

Supplementary Fig 14: Predict gene status using limited core genes, embedded core genes, and correlation between frequency and predictability. | First column: the prediction of sequence variant (Seq Vrts) of VFS/ARGs and core genes (Cg.) were tested when only the prevalent sequence variants were predicted (excluding the class 'other'). **Second column**: with embedded input, the gene status prediction of a large range of genes from the pan-genome was tested and achieved a high average predictability. **Third column**: predictability of genes in the pan-genome was test for correlation with genes frequency (presence rate) in SAC in which gene status predictability do not show correlation with gene frequency.

Predicting 7th using 1th ~ 6th (CVL rank in *E.coli*)

Confusion matrix (sample size per class = 1000)

Pairwise distance (mismatch + gap)

Test Set 1 — Random initialization 1

	sv0	sv1	sv2	sv3	sv4	sv5	other
sv0	1000	0	0	0	0	0	0
sv1	0	987	7	3	0	2	1
sv2	0	3	947	21	0	21	8
sv3	0	1	14	939	0	29	17
sv4	0	0	1	0	999	0	0
sv5	0	2	11	40	0	924	23
other	9	5	103	67	3	25	788

Test Set 2 — Random initialization 1

	sv0	sv1	sv2	sv3	sv4	sv5	other
sv0	999	0	0	0	0	0	1
sv1	0	984	8	4	0	3	1
sv2	0	3	935	18	0	18	26
sv3	0	1	12	910	0	48	29
sv4	0	0	0	0	997	0	3
sv5	0	0	14	57	0	896	33
other	11	4	78	45	1	33	828

Test Set 1 — Random initialization 2

	sv0	sv1	sv2	sv3	sv4	sv5	other
sv0	999	0	0	0	0	0	1
sv1	0	969	12	7	0	9	3
sv2	0	0	899	28	0	29	44
sv3	0	2	6	955	0	23	14
sv4	0	0	0	0	998	0	2
sv5	0	0	9	71	0	905	15
other	7	5	71	63	4	43	807

Test Set 2 — Random initialization 2

	sv0	sv1	sv2	sv3	sv4	sv5	other
sv0	999	0	0	0	0	0	1
sv1	0	981	4	2	0	7	6
sv2	0	1	953	10	0	18	18
sv3	0	1	1	946	0	21	31
sv4	0	0	0	0	998	0	2
sv5	0	0	3	42	0	931	24
other	3	5	68	46	6	31	841

Pairwise distance (mismatch + gap)

	sv1	sv2	sv3	sv4	sv5
sv0	15	16	14	28	15
sv1	-	1	1	31	2
sv2	-	-	2	30	3
sv3	-	-	-	32	1
sv4	-	-	-	-	33

Parameters of 7th Cluster

length		distance	
min	max	min	max
1011	1011	1	33

Frequency in SAC

sv0	7301
sv1	5516
sv2	9191
sv3	4379
sv4	4401
sv5	6299
Sum	37,087

Supplementary Fig 15: **Results of predicting *E.coli* [th] (CVL rank) gene using 1[th] to 6[th] and different between sequence variants.** | The confusion matrix of each four re-trains (**left**), two random initialization within each of two-fold rows, had rows as actual class and columns as predicted class where each class has 1,000 samples. The table on **top right** indicated the pariwise distance (mismatches + gaps) among all prevalent sequence variants (sv). In **bottom right**, the table indicated that all prevalent st has same length therefore the minimum (min) distance is 1 nucleotide difference over all pairwise distances; this table also indicated the frequency of each prevalent st in the SAC.

Predicting 12th using 1th ~ 6th (CVL rank in *E.coli*)

Confusion matrix (sample size per class = 1000)

Random initialization 1

Test Set 1

	sv0	sv1	sv2	sv3	sv4	sv5	sv6	other
sv0	1000	0	0	0	0	0	0	0
sv1	0	987	0	0	0	0	0	13
sv2	0	0	1000	0	0	0	0	0
sv3	0	1	0	957	0	0	0	42
sv4	0	0	0	2	998	0	0	0
sv5	0	0	0	1	0	998	0	1
sv6	0	0	0	1	0	0	996	3
other	27	50	6	70	6	6	1	834

Test Set 2

	sv0	sv1	sv2	sv3	sv4	sv5	sv6	other
sv0	998	0	0	0	0	0	0	2
sv1	0	969	0	2	0	0	0	29
sv2	0	0	1000	0	0	0	0	0
sv3	0	1	1	954	0	0	0	44
sv4	0	0	1	1	998	0	0	0
sv5	0	0	0	0	0	999	0	1
sv6	0	0	0	1	0	0	998	1
other	27	49	6	64	6	7	0	841

Random initialization 2

Test Set 1

	sv0	sv1	sv2	sv3	sv4	sv5	sv6	other
sv0	999	0	0	0	0	0	0	1
sv1	0	981	0	0	0	0	0	19
sv2	0	0	999	1	0	0	0	0
sv3	0	6	0	940	0	0	0	54
sv4	0	0	0	0	999	0	1	0
sv5	0	0	0	0	0	1000	0	0
sv6	0	0	0	1	0	0	998	1
other	25	39	8	67	6	5	2	848

Test Set 2

	sv0	sv1	sv2	sv3	sv4	sv5	sv6	other
sv0	1000	0	0	0	0	0	0	0
sv1	0	982	0	4	1	0	0	13
sv2	0	0	999	1	0	0	0	0
sv3	0	6	0	952	1	0	0	41
sv4	0	0	0	0	1000	0	0	0
sv5	0	0	0	0	0	996	0	4
sv6	0	0	0	1	0	0	998	1
other	25	42	7	73	9	2	1	841

Pairwise distance (mismatch + gap)

	sv1	sv2	sv3	sv4	sv5	sv6
sv0	52	53	52	53	44	54
sv1	-	19	18	19	51	20
sv2	-	-	1	2	55	3
sv3	-	-	-	1	54	2
sv4	-	-	-	-	55	3
sv5	-	-	-	-	-	56

Parameters of 12th Cluster

length		distance	
min	max	min	max
1419	1419	1	56

Frequency in SAC

sv	
sv0	6664
sv1	5928
sv2	3357
sv3	6792
sv4	2255
sv5	4411
sv6	3077
Sum	32,484

Supplementary Fig 16: Results in predicting *E.coli* 12^{th} (CVL rank) gene using 1^{th} to 6^{th} and different between sequence variants. | The confusion matrix of each four re-trains (**left**), two random intialization within each of two-fold rows, had rows as acutal class and columns as predicted class where each class has 1,000 samples. The table on **top right** indicated the pariwise distance (mismatches + gaps) among all prevalent sequence variants (sv). In **bottom right**, the table indicated that all prevalent st has same length therefore the minimum (min) distance is 1 nucleotide difference over all pairwise distances; this table also indicated the frequency of each prevalent st in the SAC.

Supplementary Fig 17: Sample size available in training correlates with consistent test set cross validation performance (a) but not with better prediction accurary (b). | x axis of both **a** and **b** is the minimum class size, as the minimum number among positive and negative sample size (Fig 5). **(a)** y axis represents the mean standard deviation of all input windows of a phenotype prediction. **(b)** y axis represents the mean resistance prediction performance of all input windows of a phenotype prediction.

Supplementary Fig 18: *E. coli* **cgST specific k-mers selected by LR based re-trains.** | 3-mer (first row) and 7-mer (second row) frequency were extracted from Dataset II of *E. coli*. These k-mer patterns from StCGCs CVL rank window of 1 to 6 (first column), 201 to 206 (second column), and 401 to 406 (third column) were used to predict stain's cgST where re-train were performed 20 times for each cgST. Strains on the x axis were stained the same color as in Fig 2 with cgST corresponding color. Number on the x axis is the average AUROC over 20 re-trains in predicting the cgST.

Supplementary Fig 19: *E. coli* **cgST specific k-mers selected by LR based re-trains.** | 3-mer (first row) and 7-mer (second row) frequency were extracted from Dataset II of *E. coli*. And these k-mer patterns from StCGCs rank list with odd (first column) and even (second column) rank were used to predict stain's cgST where re-train were performed 20 times for each cgST. Strains on the x axis were stained the same color as in Fig 2 with cgST corresponding color. Number on the x axis is the average AUROC over 20 re-trains in predicting the cgST.

Supplementary Fig 20: *E. coli* **3-mer difference obetween core gene combinations within each cgST.** | Each plot is the PCoA of 3-mer patterns of genes from CVL window of 1 to 6 (red dot), 201 to 206 (purple triangles), and 401 to 406 (blue square). All group pairwise significance test using permanova in each plot resulted in P < 0.001 significance.

Supplementary Fig 21: *E. coli* **re-prediction performance by input of core genome subsets.** | two groups of models were pre-trained using complete core genome for cgST prediction, where one group used structure of model A (first row) in Supplementary Fig 8 and a Res-Net like structure (Supplementary Fig 21) for another group (second row). Ten pre-trained models from each group performed re-prediction using different subset of input. By using CVL window length of 6 (first column), 20 (second column), and 100 (thrid column) genes, re-prediction shows low accuracy while high test set performance models could be trained using same input combinations.

Supplementary Fig 22: *E. coli* **Res-Net like structure used for re-prediction test in Supplementary Fig 21.** | This customized structure input dimension is the same as complete core genome of *E. coli* Dataset II and applied 2 convolutional layer first to achieve certain dimension reduction. The model subsequently performs a Res-Net computation with 3 blocks and each block includes a cascade of 3 sections. Each section would perform residue connection, batch normalization and ReLu. Downsampling were performed between each block and a final fully connected layer helped softmax for cgST prediction.

References

1. Liu, B., Zheng, D., Zhou, S., Chen, L. & Yang, J. VFDB 2022: A general classification scheme for bacterial virulence factors. *Nucleic Acids Research* **50**, D912–D917 (2022).
2. Olson, R. D. *et al.* Introducing the Bacterial and Viral Bioinformatics Resource Center (BV-BRC): A resource combining PATRIC, IRD and ViPR. *Nucleic Acids Res* **51**, D678–D689 (2023).
3. Wattam, A. R. *et al.* Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* **45**, D535–D542 (2017).
4. Davies, T. J. *et al.* Reconciling the Potentially Irreconcilable? Genotypic and Phenotypic Amoxicillin-Clavulanate Resistance in Escherichia coli. *Antimicrobial Agents and Chemotherapy* **64**, 10.1128/aac.02026–19 (2020).
5. Moradigaravand, D. *et al.* Prediction of antibiotic resistance in Escherichia coli from large-scale pan-genome data. *PLOS Computational Biology* **14**, e1006258 (2018 12 14).
6. Bradley, P. *et al.* Rapid antibiotic-resistance predictions from genome sequence data for Staphylococcus aureus and Mycobacterium tuberculosis. *Nat Commun* **6**, 10063 (2015).
7. Kumar, N. *et al.* Evaluation of a fully automated bioinformatics tool to predict antibiotic resistance from MRSA genomes. *Journal of Antimicrobial Chemotherapy* **75**, 1117–1122 (2020).
8. Coll, F. *et al.* Genome-wide analysis of multi- and extensively drug-resistant Mycobacterium tuberculosis. *Nat Genet* **50**, 307–316 (2018).
9. Wirth, T. *et al.* Sex and virulence in Escherichia coli: An evolutionary perspective. *Mol Microbiol* **60**, 1136–1151 (2006).
10. Ragon, M. *et al.* A new perspective on Listeria monocytogenes evolution. *PLoS Pathog* **4**, e1000146 (2008).
11. Dingle, K. E. *et al.* Multilocus sequence typing system for Campylobacter jejuni. *J Clin Microbiol* **39**, 14–23 (2001).
12. Enright, M. C., Day, N. P. J., Davies, C. E., Peacock, S. J. & Spratt, B. G. Multilocus Sequence Typing for Characterization of Methicillin-Resistant and Methicillin-Susceptible Clones ofStaphylococcus aureus. *Journal of Clinical Microbiology* **38**, 1008–1015 (2000).
13. Pitondo-Silva, A., Santos, A. C. B., Jolley, K. A., Leite, C. Q. F. & Darini, A. L. da C. Comparison of three molecular typing methods to assess genetic diversity for *Mycobacterium Tuberculosis*. *Journal of Microbiological Methods* **93**, 42–48 (2013).
14. O'Toole, R. F. Development of a New Genome-Wide MLST Scheme for High-Resolution Typing of Diverse Mycobacterium tuberculosis Complex Strains. *eBioMedicine* **34**, 6–7 (2018).
15. Chitale, P. *et al.* A comprehensive update to the Mycobacterium tuberculosis H37Rv reference genome. *Nat Commun* **13**, 7068 (2022).