Host population structure and rare dispersal events drive leptospirosis transmission patterns among *Rattus norvegicus* in Boston, Massachusetts, US

Short Title: Brown rat population dynamics drive leptospirosis transmission in Boston, US

Nathan E. Stone[1], Camila Hamond[2], Joel Clegg[1], Ryelan F. McDonough[1], Reanna M. Bourgeois[1], Rebecca Ballard[1], Natalie B. Thornton[1], Marianece Nuttall[1], Hannah Hertzel[3], Tammy Anderson[2], Ryann N. Whealy[1,4], Skylar Timm[1,4], Alexander K. Roberts[1,4], Verónica Barragán[5], Wanda Phipatanakul[6], Jessica H. Leibler[7], Hayley Benson[3], Aubrey Specht[3], Ruairi White[3], Karen LeCount[2], Tara N. Furstenau[4], Renee L. Galloway[8], Nichola J. Hill[9], Joseph D. Madison[9,10], Viacheslav Y. Fofanov[1,4], Talima Pearson[1], Jason W. Sahl[1], Joseph D. Busch[1], Zachary Weiner[8], Jarlath E. Nally[11], David M. Wagner[1¶*], and Marieke H. Rosenbaum[3¶*]

[1]The Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ, US

[2]National Veterinary Services Laboratories, APHIS, United States Department of Agriculture, Ames, IA, US

[3]Department of Infectious Disease and Global Health, Cummings School of Veterinary Medicine, Tufts University, Grafton, MA, US

[4]School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, AZ, US

[5] Universidad San Francisco de Quito, Colegio de Ciencias Biologicas y Ambientales, Quito, EC

[6]Division of Allergy & Immunology, Boston Children's Hospital, Boston, MA, US

[7]Department of Environmental Health, Boston University School of Public Health, Boston, MA US

[8]Bacterial Special Pathogens Branch, Centers for Disease Control and Prevention, Atlanta, GA, US

[9]University of Massachusetts Boston, Biology Department, Boston, MA, US

[10]Center for Conservation Genomics, Smithsonian's National Zoo & Conservation Biology Institute,

Washington, DC, US

[11]Infectious Bacterial Diseases Research Unit, ARS, United States Department of Agriculture, Ames,

IA, US

[¶]These authors contributed equally to this work.

* Corresponding authors

Marieke.Rosenbaum@tufts.edu

Dave.Wagner@nau.edu

Keywords: Leptospirosis, urban setting, whole genome, *Rattus norvegicus*, population genetics

Formatted for PLoS Biology

No maximum word count

Figures (4)

Tables (1)

Supplementary Figures (7)

Supplementary Tables (9)

Supplementary Files (1)

Author Email Addresses

Nathan E. Stone: Nathan.Stone@nau.edu

Camila Hamond: Camila.Hamond@usda.gov

Joel Clegg: jrc665@nau.edu

Ryelan F. McDonough: Ryelan.McDonough@nau.edu

Reanna M. Bourgeois: reannamb21@gmail.com

Rebecca Ballard: Rebecca.Ballard@nau.edu

Natalie B. Thornton: nbt39@nau.edu

Marianece Nuttall mkn75@nau.edu

Hannah Hertzel: hrhertzel@gmail.com

Tammy Anderson: tammy.m.anderson@usda.gov

Ryann N. Whealy: rnw42@nau.edu

Skylar Timm: sab832@nau.edu

Alexander K. Roberts: akr348@nau.edu

Verónica Barragán: vbarragan@usfq.edu.ec

Wanda Phipatanakul: wanda.phipatanakul@childrens.harvard.edu

Jessica H. Leibler: jleibler@bu.edu

Hayley Benson: haylbenson@gmail.com

Aubrey Specht: aubrey.g.specht@gmail.com

Ruairi White: ruairi.g.white@gmail.com

Karen LeCount: karen.j.lecount@usda.gov

Tara N. Furstenau: Tara.Furstenau@nau.edu

Renee L. Galloway: zul0@cdc.gov

Nichola J. Hill: Nichola.Hill@umb.edu

Joseph D. Madison: josephdmadison7@gmail.com

Viacheslav Y. Fofanov: viacheslav.fofanov@nau.edu

Talima Pearson: talima.pearson@nau.edu

Jason W. Sahl: Jason.Sahl@nau.edu

Joseph D. Busch: Joseph.Busch@nau.edu

Zachary Weiner: xxd7@cdc.gov

Jarlath E. Nally: jarlath.nally@usda.gov

David M. Wagner: Dave.Wagner@nau.edu

Marieke H. Rosenbaum: Marieke.Rosenbaum@tufts.edu

Abstract

Leptospirosis (caused by pathogenic bacteria in the genus *Leptospira*) is prevalent worldwide but more common in tropical and subtropical regions. Transmission can occur following direct exposure to infected urine from reservoir hosts, such as rats, or a urine-contaminated environment, which then can serve as an infection source for additional rats and other mammals, including humans. The brown rat, *Rattus norvegicus*, is an important reservoir of leptospirosis in urban settings. We investigated leptospirosis among brown rats in Boston, Massachusetts and hypothesized that rat dispersal in this urban setting influences the movement, persistence, and diversity of *Leptospira*. We analyzed DNA from 328 rat kidney samples collected from 17 sites in Boston over a seven-year period (2016-2022); 59 rats representing 12 of 17 sites were positive for *Leptospira*. We used 21 neutral microsatellite loci to genotype 311 rats and utilized the resulting data to investigate genetic connectivity among sampling sites. We generated whole genome sequences for 28 *Leptospira* isolates obtained from frozen and fresh tissue from some of the 59 *Leptospira*-positive rat kidneys. When isolates were not obtained, we attempted *Leptospira* genomic DNA capture and enrichment, which yielded 14 additional *Leptospira* genomes from rats. We also generated an enriched *Leptospira* genome from a 2018 human case in Boston. We found evidence of high genetic structure and limited dispersal among rat populations that is likely influenced by major roads and/or other unknown dispersal barriers, resulting in distinct rat population groups within the city; at certain sites these groups persisted for multiple years. We identified multiple distinct phylogenetic clades of *L. interrogans* among rats, with specific clades tightly linked to distinct rat populations. This pattern suggests *L. interrogans* persists in local rat populations and movement of leptospirosis in this urban rat community is driven by rat dispersal. Finally, our genomic analyses of the 2018 human leptospirosis case in Boston suggests a link to rats as the source. These findings will be useful for guiding rat control and human leptospirosis mitigation efforts in this and other urban settings.

Introduction

Rats (*Rattus norvegicus and R. rattus*) are highly invasive species globally, persisting on all continents (except Antarctica) and in most habitats (1, 2). They cause devastating impacts to natural ecosystems (3), and are considered a serious public health threat (4, 5). *Rattus norvegicus* (the brown rat) is ubiquitous in urban environments (6, 7) and has been documented to carry/transmit (either directly or indirectly) an abundance of bacterial and viral pathogens in urban ecosystems, including *Leptospira* spp., *Bartonella* spp., *Borrelia* spp., *Yersinia pestis*, *Streptobacillus moiliformis*, antibiotic-resistant *Staphylococcus* spp., *Shigella* spp., *Campylobacter* spp., *Escherichia coli*, *Rickettsia* spp., *Salmonella* spp., influenza A virus, and Seoul hantavirus (8-11).

Multiple genetic studies utilizing various methods have assessed *R. norvegicus* population structure, movements, and dispersal patterns in urban settings in both temperate and tropical climates (2, 12-17), consolidated in a recent review (6). Despite this somewhat limited body of research, findings are generally uniform across all cities assessed (Vancouver, Canada; New York City, US; New Orleans, US; Baltimore, US; Salvador, Brazil; and Hauts-de-Seine, France): 1) individual *R. norvegicus* utilize small home ranges, display site fidelity, and display philopatry to their birth site; and 2) dispersal events are more common in adult rats (male or female), but rare overall, and often inhibited by major roads and rivers, which act as dispersal barriers.

Leptospirosis is the most globally widespread bacterial zoonoses resulting in over 1 million human infections annually and nearly 60,000 deaths (18,19). Brown rats are asymptomatic, chronic carriers of pathogenic *Leptospira* spp. (20-23) and the most important known source of human leptospirosis infections (24). Leptospires colonize and replicate in the proximal renal tubules of rats (20-22), with transmission occurring after leptospires are shed through urine, either via direct contact with the urine or indirectly via environmental or fomite contamination (24). Globally, >30% of *R. norvegicus* are estimated to be infected with *Leptospira* spp. (25).

Understanding how rat movements relate to *Leptospira* transmission in urban settings is of

vital importance to guide the rational design of disease mitigation strategies. A recent study of rat

population structure and pathogen density in an urban setting (Vancouver) suggested *L. interrogans*

prevalence was associated with rat population connectivity (16), wherein related rats sampled from

neighboring blocks shared the same pathogen status (*i.e.*, infected/not infected). Our study builds

upon this finding by pairing fine-scale population genetic analyses of rats in another urban city

(Boston, US) with fine-scale genomic analyses of the *Leptospira* they carry and transmit.

Historically, the acquisition of *Leptospira* genomes from reservoir (*e.g.*, rats) and incidental

(*e.g.*, humans) hosts has relied on the ability to obtain cultured isolates, which is inherently

challenging (26). Obtaining isolates from human leptospirosis cases is further inhibited by the

administration of antibiotics used to treat the disease (27), after which isolation of live leptospires is

almost impossible. As such, genomic analyses of infecting strains are lacking, and these data are

critical for improved understanding of leptospirosis ecology and epidemiology. We implemented two

new culture-independent techniques for the genetic/genomic characterization of infecting *Leptospira*

spp. in rats and humans in Boston: DNA capture and enrichment (28) and amplicon sequencing

(AmpSeq; described herein). Our goal was to gain a deeper understanding of the nuanced *Leptospira*

transmission network among urban *R. norvegicus* and potential spillover into humans. Our findings

are useful for guiding rat control and human leptospirosis mitigation efforts in urban settings.

Results

lipL32 qPCR in rodent samples

In total, 59 *R. norvegicus* kidney samples and one fresh urine sample (derived from an animal

that also had a positive kidney) were *lipL32* PCR positive, revealing an overall *Leptospira* infection

rate of 18.0% (59 of 328 animals) in this *R. norvegicus* dataset (**S1 Table**). *Leptospira*-positive

kidneys included 37 frozen samples (17.6%; 37/210) yielding an average Ct value of 28.77 ± 5.58,

and 23 fresh samples from 22 rats (18.6%; 22/118) yielding an average Ct value of 27.1 ± 6.06. Four

bycatch mice and one squirrel were *lipL32* PCR negative (**S2 Table**). Fifty-six of the infected *R.*

norvegicus were detected among 20/33 collections representing 12/17 sampling sites (**Fig 1 and S1**

Table) and positive rats were detected in all sampling years [2016=3/42 (7.1%), 2017=6/50 (12.0%),

2018=16/68 (23.5%), 2019=9/42 (21.4%), 2020=3/8 (37.5%), 2021=14/70 (20.0%), 2022=8/48

(16.7%)]. Precise locations were unknown for infected *R. norvegicus* collected in 2020 (samples B1,

B2, and B4; **S1 and S2 Tables**). Infected rats were more commonly adults (58/59) than juveniles

(1/59) (X^2 =60.68, p <0.001), and a higher proportion of infected rats were collected in fall (0.277)

and winter (0.283) compared to spring (0.196) and summer (0.123); a chi-square test indicated a non-

random association between *Leptospira* infection and season (X^2 =10.33, p <0.05). We also detected

a significant association between *Leptospira* infection and site (X^2 =26.58, p <0.05); however, this

association is likely driven by capture of a single rat from site 4 that was infected, and 3/5 infected

rats captured at site 15 (60%); all other sites ranged from 0-33.3% positivity. No other significant

associations were observed between other sampling variables (sex, sampling year, genetic group) and

the detection of *Leptospira*.

Fig 1. *Rattus norvegicus* population assignment. STRUCTURE population assignment of 311 *R.*

norvegicus with collections color coded according to genetic group (S1 Table); likely dispersal

barriers and major open spaces are indicated with black bars and green shading, respectively. A)

Inferred migrants are represented by one-sided arrows; shared genetic groups among collections from

sites 13, 14, and 16, and separately sites 9 and 10, are indicated by thin lines; and *lipL32* PCR

positivity at 12 of 17 sites is indicated with asterisks. Two migrants (sites 2 and 6) were both *lipL32*

PCR positive and are denoted by asterisks on one-sided arrows. B) Each individual *R. norvegicus*

(n=311) is assigned a probability of membership (Q ≥0.75) to one of 12 genetic groups

(S3 Table). Collection ID is indicated on the X-axis and probability (Q) on

the Y-axis; each thin vertical line represents the Q value for an individual rat. The map in panel A

was created using ArcGIS software by Esri. ArcGIS and Arc-Map are the intellectual property of Esri

and are used herein under license. Copyright Esri. All rights reserved. For more information about

Esri software, please visit www.esri.com. Basemap: Light Gray Canvas Base v2

https://www.arcgis.com/home/item.html?id=291da5eab3a0412593b66d384379f89f.

Culture

Twenty-eight isolates were obtained from *lipL32* PCR positive kidneys, including 15 from 37

frozen samples and 13 from 22 fresh samples. Nearly half of the collections (15/33) are represented

by these isolates as well as all sampling years (2016=1, 2017=2, 2018=6, 2019=3, 2020=3, 2021=7,

2022=6; **S2 Table**). The frozen samples were originally cultured in both T80/T40/LH and HAN

media, which were incubated at 29 and 37°C, respectively. However, it became evident that they

were heavily contaminated, and we observed that the contaminants grew better in HAN compared to

T80/T40/LH. Therefore, use of T80/T40/LH at 29°C was prioritized. Fresh kidneys were positive by

culture in both HAN and T80/40/LH media.

Serotyping of *Leptospira* isolates

All isolates had high agglutination titers with reference antiserum specific for serogroup

Icterohaemorrhagiae (**S2 Table**). Isolates were completely seronegative (at 1:100 or above) (29) for

any other serogroup tested, except for three samples: MAR_57 had a titer of 1:102,400 against

Icterohaemorrhagiae and 1:100 against Pyrogenes; Cad103 had a titer of 1:51,200 against

Icterohaemorrhagiae, 1:200 against Djasiman, and 1:100 against Hardjo; and R304 had a titer of

1:52,400 against Icterohaemorrhagiae and 1:100 against Pyrogenes. We consider all three of these

isolates to be serogroup Icterohaemorrhagiae as it is not uncommon to see cross reactions with other

serogroups (30).

Evaluation of virulence

203   Intraperitoneal inoculation of all hamsters with 10^8 leptospires derived from a frozen *R.*

204 *norvegicus* kidney (strain R47) resulted in acute disease requiring all hamsters to be euthanized

205 four days post-infection. Liver and kidney samples from each infected hamster were positive by both

206 culture and *lipL32* qPCR; Ct values for positive liver and kidney samples were on average

207 19.04 ± 0.63 and 23.64 ± 0.75, respectively.

Rat MSAT results

MSAT marker validation

Twenty-one microsatellite (MSAT) markers were suitable for use in this population genetic

analyses of *R. norvegicus*. Only one marker (D12Rat76) deviated from Hardy-Weinberg equilibrium

in one collection (8f) at the Bonferroni corrected $a = 0.000595$ ($a = 0.05/84$ tests) and no signatures of

linkage disequilibrium were detected. Null alleles, large allelic dropout, or scoring errors due to

stuttering also were not detected, although we did detect excessive homozygosity in two markers

(D12Rat76 and D5Rat83) for collection 2c and one marker in four other collections (D5Rat33 in 5b,

D10Mit5 in 6g1, D7Rat13 in 12a, and D1Cebr3 in 12ab). No single marker displayed excessive

homozygosity in more than one collection. To check for allele scoring errors, we replicated PCRs for

all 21 loci across 94 *R. norvegicus* DNAs (30.2% of the final dataset) and detected an allele scoring

error rate of 0.23% (9 errors from 3,948 possible allele calls).

Population genetic analyses: determination of genetic groups and population assignment using STRUCTURE

The delta-K method estimated 2 and 12 genetic groups as the most likely number of populations sampled in the 311 individual rats from 33 collections/17 sites (**S1A Fig**). The K of 12 is more consistent with what is known about the philopatric nature of *R. norvegicus* (6, 12) and physical distances among our collection sites. Variance among the L(K) values is very low at $K =12$ but rapidly increases beyond this number (**S1B Fig**); this phenomenon is discussed in the STRUCTURE manual and can be expected to occur after the real K is reached (31). Furthermore, the higher delta-K value at $K =2$ could simply be that $K =1$ is consistently a very poor solution and the model improved dramatically between $K =1$ and 2. The 12 genetic groups were each assigned an arbitrary color in STRUCTURE (**Fig 1**) and that color was retained across all analyses, figures, and tables. Thus, we refer to the genetic groups by the following color designations: Blue, Brown, Evergreen, Green, Light Blue, Orange, Pink, Purple, Red, Steel, Tan, Yellow, and admixed (indicated by the color White) (**Fig 1 and S1 Table**). To aid in visualizing patterns across space and time, collections were color coded according to the most dominant genetic group (≥50%) present in that collection (**defined in S1 Table and displayed in Figs 1 and 2**). In total, 247 rats were assigned to one of these genetic groups at Q ≥0.75 (**see Materials and Methods**), whereas 64 rats were assigned to the admixed group (**S3 Table**). Due to repeated sampling over multiple years at sites 1, 5, 6, 8, 9, 10, 12, and 17, we detected stable rat populations at 7/8 sites for up to five years: site 1 (Blue group: 2016-2018), site 5 (Light Blue group: 2016-2017), site 6 (Tan group: 2018-2022 and Yellow group: 2021-2022), site 8 (Red group: 2017-2018 and Steel group: 2021-2022), site 9 (Brown group: 2021-2022), site 12 (Green group: 2016-2017), and site 17 (Pink group: 2018-2019); as well as population turnover/invasion

followed by persistence at two sites: site 6 (Tan group in 2018 to Yellow and Tan groups from 2020-

2021) and site 8 (Red group from 2017-2018 to Steel group from 2021-2022) (**Fig 1**).

Fig 2. *Leptospira interrogans* serogroup Icterohaemorrhagiae whole genome phylogeny. A

maximum likelihood phylogeny of 28 *Leptospira interrogans* serogroup Icterohaemorrhagiae isolate

genomes from rats, two publicly available *L. interrogans* serogroup Icterohaemorrhagiae serovar

Copenhageni complete genomes (GenBank accession# GCA_002073495.2 and GCA_010978155.1),

and 13 enriched genomes (12 rats and 1 human) based upon a concatenated SNP alignment of 127

positions out of a core genome size of 2,123,028 nucleotides (nts). The phylogeny was rooted with

reference genome *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni strain Fiocruz

L1-130 (GenBank accession# GCA_000007685.1). Four major clades were observed (1-4 in white

text) and bootstrap values are indicated at specific nodes. The color of the genome ID corresponds to

the rat genetic groups in **Fig 1**; black text indicates that no genetic group was assigned. One inferred

migrant (indicated with a black arrow) from site 6c was trapped at site 2c and carried *Leptospira*

interrogans identical to its source collection (see R335-Migrant in clade 1).

Using Q values ≥0.75, we detected potential *R. norvegicus* migrations as

indicated by the trapping of a rat assigning to a distinct genetic group among a collection of rats

belonging to a different genetic group; any highly admixed individuals were not treated as migrants

due to the potential for mis-assignment. In total, we detected eleven such migration events, seven of

which were adult rats (5 females, 2 males) and four were juveniles (1 female, 3 males). Rat

movement distances ranged from ~0.8-5.3 km. Migrations were suspected from collection 17c or 17d

to collection 14d (1 juvenile female, Q =0.93); collection 6c to collection 2c (2 juveniles males, Q

=0.91 and 0.95, and 1 adult male, Q =0.76); collection 13d, 14d, or 16d (all belonging to the

Evergreen genetic group) to collection 15d (1 adult female, Q =0.81); collection 5a or 5b to

collection 12a (1 adult female, Q =0.92); collection 9g1, 9g2, or 10c (Brown genetic group) to

collection 6g1 (1 adult male, Q =0.76); collection 15d to collection 10d (1 adult female, Q =0.97);

collection 12a or 12b to collection 5a (1 juvenile male, Q =0.82); collection 8b or 8c to collection 1b

(1 adult female, Q =0.83); and collection 13d, 14d, or 16d (Evergreen genetic group) to collection 7d

(1 adult female, Q =0.87) (**Fig 1 and S3 Table**). Two migrants (R335 from collection 2c and Cad89

from 6g1) were *lipL32* positive (**Fig 1 and S3 Table**). Some suspected migrations could also result

from home range overlap among neighboring collections.

Population genetic analyses: genetic structure

Principal Coordinates Analysis (PCoA) generally supported the 12 *R. norvegicus* genetic

groups identified by STRUCTURE and indicated two genetic groups (Blue and Steel) may be more

genetically isolated from the other ten groups (**S2 Fig**). The collections comprising these two genetic

groups are geographically separated from each other and the other genetic groups by major roadways

(**Figs 1 and 3**). This PCoA analysis of standardized genetic distances mirrors the geographic

distribution of these genetic groups (**Fig 1**), suggesting a relationship between *R. norvegicus* genetic

and geographic distance in this study system.

Fig 3. *Rattus norvegicus* trap locations. Map indicating 64 trap locations (white circles) at 17 sites

(numbers), likely dispersal barriers (black bars), and major open spaces (green shading). Circle size

corresponds to the number of total rat samples collected from each trap location. This map was

created using ArcGIS software by Esri. ArcGIS and Arc-Map are the intellectual property of Esri and

are used herein under license. Copyright Esri. All rights reserved. For more information about Esri

software, please visit <u>www.esri.com</u>. Basemap: Light Gray Canvas Base v2

<u>https://www.arcgis.com/home/item.html?id=291da5eab3a0412593b66d384379f89f</u>.

Fairly high estimates of global F_{ST} were observed in the FSTAT analysis, with θ =0.184 (95%

CI =0.155, 0.214). Of 66 pairwise F_{ST} estimates among 12 *R. norvegicus* genetic groups, all were

significant at *a* =0.05 and only 16/66 were not significant at the Bonferroni corrected *a* =0.000758.

That said, pairwise F_{ST} values were high among the 66 pairwise comparisons, ranging from 0.0826-

0.2922 (**S4 Table**). All significant *p*-values (16/16) were associated with the Purple, Red, Orange,

and Brown genetic groups, a result in agreement with the pairwise analysis of similarities (ANOSIM)

(see below). Likewise, an Analysis of molecular variance (AMOVA) generated a moderate yet

significant PhiPT value of 0.312 (p <0.001), revealing that ~31% of the molecular variance in this

dataset was present among *R. norvegicus* genetic groups (*i.e.,* intersite variation), whereas 69% was

present within them (*i.e.,* intrasite variation).

Population genetic analyses: isolation by distance and fine-scale genetic structure

To explore signatures of isolation by distance, we used the RELATE function in PRIMER

v5.2.9 and the standardized genetic distances and geographic locations (latitude and longitude) of the

247 individual *R. norvegicus* forming the 12 genetic groups (**see Materials and Methods**). The

Spearman's rank correlation coefficient (Rho) was calculated as 0.232 (p <0.001), identifying a

statistically significant relationship between *R. norvegicus* genetic distance and geographic distance.

The global ANOSIM R statistic was 0.921 (p <0.0001), documenting that some of the 12 genetic

groups formed distinct geographic groups. Most of the pairwise ANOSIM comparisons among the 12

genetic groups (61 of 66) were significant even with a very conservative Bonferroni corrected *a*

=0.000758, indicating that most of the 12 genetic groups formed distinct geographic groups. Only

5/66 pairwise comparisons were not significant at a =0.000758 (Brown-Orange, Orange-Purple,

Purple-Red, Pink-Purple, and Purple-Evergreen), suggesting the Purple group has a genetic

connection with three other groups (Red, Pink, and Evergreen).

Consistent with the signal of *R. norvegicus* genetic isolation by distance across all sites,

microsatellite data revealed clear evidence of fine-scale spatial genetic structure within collection

sites. First, the spatial autocorrelation analysis yielded correlation values significantly greater than

zero in the first six distance classes (0-100 m out to 0-600 m), suggesting that, on average, relatives

are generally living in close proximity (**S3 Fig**). This is supported by the significantly high value of

average relatedness of individuals within collections, at 0.313 (95% CI = 0.268, 0.357), which

indicates relatedness ranging from half (0.25) to full (0.5) siblings (**S4 Table**). Therefore, dispersing

R. norvegicus would need to travel >600 m, on average, to move beyond sites occupied by relatives.

Finally, F_{IS} values were only positive in three genetic groups [Blue (0.11), Yellow (0.078), and Red

(0.006)], suggesting inbreeding within collections at sites 1 and 2 (Blue), and perhaps site 6 (Yellow)

(**Fig 1 and S4 Table**).

Leptospira genomics and genotyping

In total, 52 of 59 *lipL32* PCR positive *R. norvegicus* and one of two positive human samples

were assigned to *L. interrogans* (86.9%) using the genotyping methods described herein (**S2 Table**).

Whole genome analyses revealed all isolates derived from frozen or fresh kidneys (n=28) and all

successful enrichments (n=15) were *L. interrogans* (**S2 Table**) and most like other serogroup

Icterohaemorrhagiae serovar Copenhageni/Icterohaemorrhagiae genomes (**S4 Fig**). Enriched

genomes for samples R47 and R316, which were included as genomic analysis controls, were

identical to their isolate genome counterparts (**Fig 2**). Phylogenomic analysis of all 28 isolates and 12

enrichments [the latter selected due to average breadth and depth of coverage values >63% and >18x,

respectively (**S5 Table**)], revealed four distinct clades that largely associate with the genetic groups

of the *R. norvegicus* from which they were derived (**Fig 2**). For example, 14 of 15 *L. interrogans*

genomes assigning to clade 1 were obtained from rats belonging to the Yellow, Tan, and Green

genetic groups, all occurring in or near Boston Common Garden, and the Tan and Yellow groups co-

occur at site 6. However, the Tan group, the most prominent in Boston Common Garden in 2018,

was replaced or modified as indicated by the emergence of the Yellow group in 2021 and 2022.

Likewise, clade 3 contains mostly *L. interrogans* genomes from the Red and Steel genetic groups (6

of 7). Both genetic groups contain rats from the same site (8) but different collections and years (8b

and 8c =Red, 8f and 8fg =Steel), perhaps indicating the removal of one rat genetic group (Red) and

replacement with another (Steel). Similarly, clade 2 contains mostly *L. interrogans* genomes from

rats belonging to the Evergreen genetic group (3 of 5); this clade also contains the *L. interrogans*

genome derived from a human leptospirosis case in 2018 (MA-2). Finally, clade 4 contains only *L.*

interrogans genomes from rats belonging to the Pink genetic group, comprised of the most

geographically isolated collections (17c and 17d) (**Fig 1**). Our phylogenetic placement of two lower

coverage enriched genomes (R289 and R292; **see S5 Table**) placed them into clade 1 among other *L.*

interrogans genomes derived from rats from collection 6c and belonging to the Tan genetic group

(**S5 Fig**).

Our higher resolution [3,280,516 nucleotides (nts)] phylogenomic comparison of *L.*

interrogans clade 2 revealed the human derived genome (MA-2) from a 2018 leptospirosis case

differed by only two single nucleotide polymorphisms (SNPs) from three *L. interrogans* genomes

obtained from rats collected in 2019, 2021, and 2022, respectively, when comparing a core genome

of >3.2 mb (**Fig 4**). The four *L. interrogans* genomes comprising this clade were from rats belonging

to the Evergreen genetic group or admixed group and were collected at sites 9 (collections 9f and

9g1) and 16 (collection 16d) (**Fig 1 and S1 Table**).

Fig 4. *Leptospira interrogans* serogroup Icterohaemorrhagiae clade 2 whole genome phylogeny.
Maximum likelihood phylogeny of four *R. norvegicus* and one human derived *L. interrogans*
genomes from clade 2 (**see Fig 2**) based upon the concatenated SNP alignment of 33 positions out of
a core genome of 3,280,516 nucleotides. The color of the genome ID corresponds to the rat genetic
group in **Fig 1**; black text indicates that no genetic group was assigned.

Our phylogenomic analysis of 24 whole genome assemblies representing all 18 P1 clade
pathogenic *Leptospira* spp. placed the 24 assemblies into 18 unique species clades, as expected (**S6A**
Fig). All nine isolate genomic DNAs subjected to AmpSeq analysis were accurately placed into their
respective species clades (**S6B Fig**), which were conserved in the AmpSeq phylogeny despite the
large reduction in size of the SNP alignment used to infer it (WGS =53,132 vs. AmpSeq =375 nts)
and the core genome included in the analyses (WGS =167,849 vs. AmpSeq =1,195 nts) (**S6 Fig**). We
note the proportion of variable sites (variable sites/core genome) represented in both the whole
genome and AmpSeq phylogenies are essentially equal (WGS =0.317 vs. AmpSeq =0.314), which
likely accounts for the similar branch lengths separating the species clades. The average number of
reads that mapped for each isolate was 44,994 ± 12,348 and the number of AmpSeq loci that
amplified for each isolate ranged from 22 for *L. kmetyi* to 39 each for *L. interrogans* and *L.*
santarosai (**S6 Table**).

Of the 20 *R. norvegicus* and one human sample subjected to AmpSeq
analysis, 13 rats yielded adequate PCR amplification and sequencing
coverage (≥7 loci with >10x reads) of *Leptospira* loci to confidently assign a species
classification of *L. interrogans* (**S6 Table**). Nine displayed enough coverage (≥25 loci
with >10x reads) to assign within species phylogenetic placement to the

L. interrogans serogroup Icterohaemorrhagiae serovar Copenhageni/Icterohaemorrhagiae clade (**S7 Fig**). The remaining seven rats and one human sample were not assigned a species classification due to inadequate PCR amplification and/or sequencing coverage (≤ 2 loci with >10x reads): two rats (R66 and R108) had no mapped loci with >10x reads, two (R324 and R379) had one mapped locus, and four samples (Cad1, MAR_36, MAR_5 and MA-1) had two mapped loci (**S6 Table**). Regardless, the few loci that did amplify for samples R324, R379, Cad1, MAR_36, MAR_5, and MA-1 all mapped to *L. interrogans* references except for MAR_5 and Cad1, which had one and two loci, respectively that mapped to *L. borgpetersenii* (**data not shown**). The *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 gDNA control amplified robustly at all 42 loci (1,444,384 assigned reads in total) and was accurately phylogenetically placed among other *L. interrogans* serovar Copenhageni strains (**S7 Fig**).

Discussion

Our population genetic analyses of *R. norvegicus* in Boston revealed the existence of distinct rat populations for periods of up to five years, most likely due to low dispersal rates among rat populations and inbreeding. Our analyses revealed significant relatedness among individuals within a genetic group and, furthermore, that related individuals are generally living within close proximity to each other (**S3 Fig**), an observation consistent with other studies of *R. norvegicus* home ranges, dispersal, and site fidelity in urban settings (6, 7, 14, 17, 32-35). In an analysis of gene flow among *R. norvegicus* in two US, one Canadian, and one Brazilian city, Combs et al., found that rats were highly related to each other within 500 m (12). Our findings in Boston were similar, as rats would need to travel, on average, >600 m to encounter an unrelated individual. Although dispersal events

were observed in this study (**Fig 1 and S1 and S3 Tables**), they were rare [only 11/311 rats (3.5%)

were presumptive migrants], and this low rate is consistent with the observed patterns of strong

genetic structure and stability of genetic groups over time (**Fig 1**).

The 11 instances of inferred migrations included seven trapped adults (5 females, 2 males)

and four juveniles (1 female, 3 males) (**S1 and S3 Tables**). Rat migration distances ranged from

~0.8-5.3 km in Boston, but because of the methodology we used to assign rats to specific collections

for population genetic analyses (**see Materials and Methods**), short-distance migrations (< 0.5 km)

would only be detected among adjacent sites separated by major roads (*e.g.,* sites 8 and 9 in **Fig 3**).

Although short-distance migrations are thought to be more common (6, 7, 36), long-distance

migrations of up to 11.5 km have been documented in brown rats within urban settings (14) and

could be anthropogenically facilitated (6, 37). We detected a juvenile female that migrated over 5.3

km from site 17 (Pink group) to site 14 (Evergreen group) (**Fig 1**). Since both sites are connected by

a large road and subway line, and adjacent to the same large river, it is possible human mediated

transport on a car, truck, train, or water vessel facilitated migration. We also detected migration of

three males (2 juveniles, 1 adult) from site 6 (Boston Common Garden) to site 2 in 2017 (**Fig 1 and

S1 and S3 Tables**). These migration events were <1 km and all three rats (R335, R336, and R337)

were collected on the same day (30-Nov-2018) at the same trap location (Government Center-1) (**S1

and S2 Tables**). We are unsure if this finding represents a group migration or if it is the result of a

common dispersal throughway between sites 6 and 2, or some other unknown dispersal mechanism.

To our knowledge, group migrations have not been described previously in the literature. In general,

our observed rate of inferred migrations (3.5%) was similar to other studies of *R. norvegicus*

population genetics in urban ecosystems; Gardner-Santana et al., reported a migration rate of 4.7% in

Baltimore, US (14), and Kajdacsi et al., reported 6.8% in Salvador, Brazil (15).

Major roads were likely the largest contributor to low rates of migration of *R. norvegicus*

among neighboring sites in Boston (**Figs 1 and 3**). We observed several examples of major roads

acting (and smaller roads failing to act) as dispersal barriers among three intensely sampled sites (6,

8, and 9). We compared the dispersal dynamics within Boston Common Garden (site 6: sampled

2016-2022) that spans ~0.83 km and is bisected by a 3-lane road with those between Melnea Cass

Blvd SE (site 8: sampled 2017-2022) and Melnea Cass Blvd N (site 9: sampled 2021-2022), which

are directly adjacent to each other but separated by an 8-lane road (**Figs 1 and 3**). The 3-lane road

bisecting the northeastern and southwestern portions of Boston Common Garden did not act as a

dispersal barrier for *R. norvegicus* as both the Tan and Yellow groups were detected ubiquitously

throughout the park. In contrast, the 8-lane road separating sites 8 and 9 prevented admixing among

them and migration between them (**Fig 1**), despite the sites being separated by only ~0.25 km.

Roadways have frequently been described as significant barriers to rat dispersal (6, 7, 17, 35, 38, 39)

and larger roadways deter dispersal more than smaller ones (40). We also observed a pattern of

genetic connectivity among sites separated by >1 km (see Evergreen group; sites 13, 14, and 16 in

Fig 1) but connected by a large park complex containing green spaces, open water, and a network of

river underpasses (**Figs 1 and 3**). Taken together, these results suggest different dispersal dynamics

in commercial/residential areas (shorter movements) vs. park settings (longer movements) in this

urban ecosystem, likely associated with road size and access to dispersal throughways. These

observations are in concert with previous findings of *R. norvegicus* movement patterns in urban

ecosystems (6, 12, 13, 41).

Rattus norvegicus population turnover occurred at two sites (6 and 8), followed by robust

expansion of the new/invading population (**Fig 1**). In Boston Common Garden we observed a large

population in 2018 (collection 6c: Tan group) that was mostly displaced by the emergence of the

Yellow group in 2021 (collection 6f). Both genetic groups remained (6g1 and 6g2) at the conclusion

of sampling in 2022 although the Yellow group was most often detected in 2021 and 2022 (**Fig 1B**).

We also observed genetic signatures of admixture among the Tan and Yellow genetic groups

(individuals assigning to both groups or neither) in the later collections (6f and 6g2 in **Fig 1B**), which

could suggest these genetic groups are in the process of merging. Complete population turnover

occurred at site 8 between October 2018 (Red group) and June 2021 (Steel group) (**Fig 1**). A rat

abatement campaign took place at site 8 but did not begin until 5-Nov-2021, suggesting population

turnover from the Red to Steel group occurred before that campaign began. Furthermore, the Steel

group persisted throughout the campaign (collections 8fg and 8g in **Fig 1**). It is unknown what

occurred at site 8 between October 2018 and June 2021 that led to the population change, but our

results suggest that rat abatement is unlikely to lead to population turnover unless complete

eradication can be achieved, as populations can rapidly rebound to pre-intervention sizes once

interventions cease (42-45). However, if an intervention resulted in removal of dominant individuals

followed by immigration of other individuals into this open niche, population turnover could occur

(7, 42). We suspect an open niche once recolonized would result in the rapid expansion of a new

genetic group due to rapid urban rat reproduction rates (up to five litters per year) (7) paired with the

tendency of *R. norvegicus* to breed with individuals from their same genetic group, a result of

dispersal barriers, site fidelity, and small home ranges (6, 12).

Leptospira interrogans persists locally in *R. norvegicus* populations in Boston. We detected

pathogenic *Leptospira* DNA in 18.0% of rats representing 20/33 collections and 12/17 sampling sites

(**Fig 1 and S1 Table**) with 98.3% of infected rats classified as adults. To put these number into

context, several other studies that examined *Leptospira* carriage in *R. norvegicus* in urban ecosystems

reported 7-82% in the highly endemic tropical climate city of Salvador, Brazil (46-48), 11.1% in the

temperate city of Vancouver, Canada (49), and 35.7% in New Orleans, US (50). Furthermore, a 2019

review of *Leptospira* carriage in rats globally reported an overall carriage rate in *R. norvegicus* of

30.3% (25), with higher carriage rates associated with tropical climates. Of the studies that assessed

the relationship between *Leptospira* positivity and age or sexual maturity, all reported significantly

higher proportions of positive adults than juveniles (47-49). Using a suite of molecular methods, we

determined the *Leptospira* species circulating in *R. norvegicus* in Boston was *L. interrogans* and

most similar to other serogroup Icterohaemorrhagiae genomes (**S4 and S7 Figs**). The 28 cultured

isolates were confirmed to belong to serogroup Icterohaemorrhagiae via the microscopic

agglutination test (MAT) (**S2 Table**), whereas the samples subjected to enrichment (n=14) and

AmpSeq (n=9) could only be inferred based upon phylogenetic placement (**Fig 2 and S4, S5, and S7

Figs**). Detection of only *L. interrogans* serogroup Icterohaemorrhagiae in this rat dataset was not

surprising. Indeed, the exclusive detection of *L. interrogans* serogroup Icterohaemorrhagiae in *R.*

norvegicus from the US was documented in a recent review (25). In that same article, it was also

reported that the most frequently identified *Leptospira* in rats globally (in 36 of 43 countries) belong

to serogroup Icterohaemorrhagiae. To that point, in the broad phylogenetic framework of *L.*

interrogans, the genomes from Boston *R. norvegicus* formed a monophyletic clade with four other *L.*

interrogans serogroup Icterohaemorrhagiae genomes from Saint Kitts (SK1), Brazil (203 and L1-

130), and Malaysia (898) (**S4 Fig**); this analysis included diverse *L. interrogans* representing >9

serovars. Importantly, monophyletic groupings like this can be teased apart with clade-focused whole

genome analysis, which was demonstrated by Santos et al., in a comprehensive study of 67 *L.*

interrogans serogroup Icterohaemorrhagiae isolates originating from >15 countries across the globe

(51). At that scale, they describe the detection of 1,072 SNPs and 258 indels among the 67 isolates,

and phylogenetic clusters that broadly associated with geographic locations. Likewise, by using

clade-focused whole genome analysis we identified >100 SNPs and four distinct clades among *L.*

interrogans serogroup Icterohaemorrhagiae genomes from *R. norvegicus* in Boston (**Fig 2 and S5

Fig**).

Distinct clades of *L. interrogans* correspond to specific geographic locations and rat

populations in Boston and persist locally through time despite population turnover (**Fig 2**). Rats from

sites 6 and 12 exclusively carried *L. interrogans* from clade 1 (**Fig 2 and S5 Fig**), the virulence of

which was validated in the hamster model of leptospirosis, despite being separated by >1.3 km and

belonging to three different rat genetic groups (Tan, Yellow, and Green); at site 6 the rat population

shifted from the Tan to Yellow group between 2018 and 2021 (**Fig 1**). Furthermore, collections from

sites 6 and 12 spanned seven years (2016-2022) (**S1 Table**). This finding suggests connectivity

among these sites and/or that stable rat populations at these sites maintain endemic clades of *L.*

interrogans over time. A greenbelt containing a public walkway connects these two sites almost

directly, and although eight roads bisect this greenbelt, none exceed 4-lanes and six are only 2-lanes,

suggesting movement between these sites is possible (40). It is also important to consider that

although these two sites contain three distinct genetic groups, the Green group from site 12 was

sampled in 2016 and 2017, whereas the Tan and Yellow groups from site 6 were sampled in 2018-

2022. It is therefore possible that the Green group no longer existed beyond 2017 and that if rats were

trapped at site 12 in 2018-2022, they would resemble the Tan and Yellow genetic groups, or –

alternatively – that the Green group was present at site 6 prior to 2018. A single rat was trapped at

site 6 in 2016 (Cad18) and although it was not assigned to a genetic group due to Q <0.75 (*i.e.,*

admixed), the most probable assignment was to the Green group with Q =0.358 (**S3 Table**). Under

either scenario, the exclusive detection of clade 1 *Leptospira* at sites 6 and 12 over a period of seven

years suggests a connection between them. Likewise, *L. interrogans* from clade 2 was exclusively

found in rats from sites 9 and 16 (**Fig 2 and S5 Fig**). These sites were separated by nearly 2.5 km and

the samples were collected over different years; the site 16 rats were trapped in 2019 and the site 9

rats in 2021-2022. As with clade 1, we see two distant locations sharing a single *L. interrogans*

genotype, suggesting a connection between them, but in this case, there is no obvious dispersal

throughway linking them. That said, we see evidence of long-distance migrations throughout this

study system (**Fig 1**), so it is plausible that a *Leptospira*-positive migrating rat from site 16 to site 9

(during or post 2019) led to the establishment of the clade 2 genotype at site 9 that we observed in

2021-2022. Population turnover of the rats with persistence of the same *Leptospira* genotype was

also observed at site 8 (**Fig 2 and S5 Fig**) where the rats displayed a genetic change from Red to

Steel between 2018 and 2021, yet the *L. interrogans* clade 3 genotype remained the same. Finally, *L.*

interrogans clade 4 contained only rats from site 17 collected in 2018-2019. Site 17 is the most

geographically distant site in this study being >4.5 km away from its nearest geographic neighbor,

site 16 (**Figs 1 and 3**), so it is not surprising to see an exclusive clade of *L. interrogans* among rats

from site 17. These results provide a clearer picture of *Leptospira* transmission among rat

populations and provide compelling evidence for local persistence of distinct *L. interrogans*

serogroup Icterohaemorrhagiae genotypes among *R. norvegicus* in Boston.

Leptospira genotypes can be dispersed between populations by migrating rats**.** We detected

the migration of a *Leptospira*-positive adult rat (R335) from site 6 to site 2 in 2018 (**Fig 1**) and the

genetic signature of the *Leptospira* it carried (clade 1) further supported this assignment (**Fig 2**).

Importantly, this event clearly illustrates a mechanism by which *Leptospira* genotypes could be

transferred between rat populations. A *Leptospira*-positive migrant (Cad89) that moved from site 9 to

site 6 (**S3 Table and Fig 1**) was also detected. Unfortunately, a genome was not obtained from this

sample so fine-scale genomic assignment was not possible. However, AmpSeq assigned it to the *L.*

interrogans serogroup Icterohaemorrhagiae clade (**S7 Fig**). Regardless, it illustrates movement of

Leptospira via rat migrations between sites 9 and 6 and it is therefore reasonable to assume that

migration of a *Leptospira*-positive rat could have occurred in the opposite direction (site 6 to 9),

ultimately resulting in a rat trapped at site 9 carrying a *L. interrogans* clade 1 genotype. We also

observed single rat populations that carried diverse *L. interrogans* that assigned to multiple locations

in the phylogeny (*e.g.*, 15d, 9g1, 8f) (**Fig 2**), suggestive of movement of *L. interrogans* between rat

populations. For example, a single rat trapped at site 9 (Cad91) contained *L. interrogans* from clade 1

(**Fig 2**). Although this rat was not assigned to a genetic group the most probable assignment was to

the Brown group (Q =0.589) (**S3 Table**). We see evidence of rat mediated *L. interrogans* movement

at site 8 where three *L. interrogans* genotypes were recovered (**Fig 2**). Because the clade 3 genotype

was the most common at site 8 and maintained over three years despite the turnover of the rat

population from the Red to Steel group, we suggest the clade 3 *L. interrogans* is established at that

site, whereas the other two genotypes (n=1 each) were the result of *Leptospira*-positive rat migrations

into site 8. Finally, we found clade 3 *L. interrogans* in a rat from site 13 that clearly assigned to the

Evergreen genetic group (**Fig 1**); clade 3 genotypes were otherwise only found in rats from site 8

(**Fig 2**). Taken together, these data provide strong support for *R. norvegicus* migrations being a

mechanism by which *L. interrogans* serogroup Icterohaemorrhagiae genotypes are transferred among

rat populations in Boston, an important consideration when designing rodent control interventions or

construction, as both can drive long-distance migrations and intraspecific antagonism (42) and

human interventions may increase rat population pathogen prevalence (52).

The mechanism(s) by which identical *Leptospira* genomes becomes established in rat

populations at these sites is unclear. *Rattus norvegicus* are well known chronic and asymptomatic

carriers of *L. interrogans* serogroup Icterohaemorrhagiae (20-22), and the transmission cycle via

carriage of *Leptospira* in renal tubules, excretion through urine, environmental contamination,

ultimately leading to direct or indirect exposure to other rats as well as other susceptible mammals,

including humans, is well established (24). During chronic carriage in the rat, one might expect to

observe the accumulation of mutations over time corresponding to an inherent *L. interrogans*

mutation rate (to our knowledge, this is unknown). Indeed, we do observe this pattern across certain

rats (R269, R289, R292 in clade 1; R365, R366 in clade 2; R336-19 in clade 4) (**Fig 2 and S5 Fig**),

presumably the result of long-term chronic infection where *L. interrogans* was in a replicative state.

More often, however, we observed identical genomes from rats spanning up to a seven-year period,

suggesting a dormant or very low replication state of *L. interrogans* in this urban ecosystem. A

possible mechanism for this could be biofilm formation (53) in the renal tubules of the rats (54).

However, laboratory analysis of *R. norvegicus* infected with *L. interrogans* serovar Copenhageni

revealed they shed leptospires continuously (10^5 -10^7 leptospires per mL) in their urine for the

duration of the experiment (159 days) (22), suggesting that in this laboratory setting there was no

such dormant state. Alternatively, *Leptospira* biofilm formation could take place in the environment

(55, 56). It is important to note that because we are only analyzing the shared genomic space among

all the rat derived *L. interrogans* genomes (core genome >2.1 million nts) there could be mutations or

acquired/lost pan genomic content outside of the core genome that we did not detect. That said, if

replication were occurring at a constant rate, we would still expect to observe mutations

accumulating in the core-genome. Additionally, when we increased our resolution for clade 2 to >3.2

million nts (**Fig 4**), we observed the same identical genomes shared among three rats (MAR-47,

Cad86, R367). Furthermore, our genomic analysis of clade 2 also included a human derived *L.*

interrogans genome (MA-2 in **Figs 2 and 4**). This genome was enriched from urine DNA obtained

from a human leptospirosis case in 2018 and only differs by two SNPS (core genome >3.2 million

nts) from *L. interrogans* genomes obtained from three rats at sites 9 and 16 in 2019 and 2022 (**Fig 4**).

Due to the clonality of these *L. interrogans* genomes derived from two host species [human

(incidental) and rat (reservoir)] and collected over a five-year period (2018-2022) paired with the

knowledge that rats are recognized globally as carriers of serogroup Icterohaemorrhagiae [while

other hosts (*e.g.*, dogs) are incidental carriers], we suggest the most likely explanation for the human

infection is exposure to a fomite or the environment that was contaminated by rats from sites 9 or 16

or another *R. norvegicus* population carrying the same *L. interrogans* clade 2 genotype in the city.

That said, any attempts to link this human case to a specific site should be interpreted with caution

given that *L. interrogans* dispersal between sites does occur (**Figs 1 and 2**).

This is the first study to pair fine-scale population genetic analyses of *R. norvegicus* in an

urban ecosystem with genomic analysis of the *L. interrogans* they harbor. This analysis was possible

due to improvements in culturing techniques (described herein) enabling high rates of *Leptospira*

recovery (47.5%) paired with advances in culture independent molecular capabilities (28) that

together enabled the acquisition of 40 *L. interrogans* genomes from rats (out of 59 *lipL32* PCR

positives) and one from a human. Importantly, these genomes were derived not just from fresh

samples, but also frozen and archived samples (tissues and DNA), thereby greatly expanding the

types of samples that could be included in these genomic analyses. A particularly salient outcome

was the acquisition of a human derived *L. interrogans* genome from trace amounts of archived DNA

(22.4 ng total). Although not much is known about the origin of this urine sample beyond it

originating from a human leptospirosis case in Boston in 2018 due to patient privacy rights, our

analysis placed it among rat derived genomes collected exclusively at sites 9 and 16 (**Fig 4**),

suggesting the source of infection was localized to a relatively small geographic area within Boston

(**Fig 1**) that was accessible to rats from both sites.

We have previously applied AmpSeq to assess the genetic diversity of *lipL32* sequences from

pathogenic leptospires [original design described by Stone et al., (56)]. AmpSeq is based upon

traditional PCR methods, Illumina short-read sequencing (both technologies that are accessible to

researchers and clinicians across the world) and requires <10 μL of a *Leptospira*-positive clinical

DNA sample. Furthermore, it is relatively inexpensive on a per-sample basis due to the ability to

simultaneously sequence hundreds of samples on a single Illumina run. For this study, we

implemented and assessed an expanded *Leptospira* AmpSeq panel that is capable of providing

Leptospira species identification among all 69 currently described and validated *Leptospira* spp.

from the P1, P2, S1, and S2 clades (57) (**S1 File**), as well as sub-species phylogenetic placement

within *L. interrogans* (*i.e.*, serogroup associated clades) (**S7 Fig**). Although we did not extensively

assess sub-species placement among other P1 *Leptospira* spp., we expect a similar outcome for *L.*

kirschneri, L. noguchii, L. alexanderi, L. santarosai, L. weilii, L. borgpetersenii, and *L. mayottensis*

due to the high number of discriminating loci (<35) expected to amplify in each species (**S1 File**).

Furthermore, our phylogenetic analysis of P1 clade pathogenic *Leptospira* spp. via sequences

generated with AmpSeq suggests sub-species placement (**S6B Fig**), even with as few as 22

amplicons (**S6 Table**). Importantly, the AmpSeq method can identify mixtures of *Leptospira* spp.

within a single sample (56), which is an under-characterized aspect of *Leptospira* ecology. Molecular

advances such as these will greatly facilitate an increased understanding of mixed infections and their

potential role in leptospirosis infection and disease ecology. Thus, we suggest it could serve as an

applicable tool for species and sub-species identification of circulating and clinically relevant

leptospires in regions throughout the world.

640 Conclusion

641   In this study, we paired population genetic analyses of *R. norvegicus* in an urban setting with

genomic analysis of the infecting leptospires they transmit to gain a deeper understanding of the

nuanced transmission patterns of *Leptospira* among rats and spillover into humans. We found that

rats in Boston displayed strong signals of population structure, likely associated with behavioral

norms and dispersal barriers. We also found the *Leptospira* carried by rats in Boston was *L.*

interrogans and most similar to other serogroup Icterohaemorrhagiae isolates, and that there was

fine-scale genetic structure within that phylogenetic group that associated with distinct rat

populations. We found evidence that long-distance dispersal events lead to the movement of unique

L. interrogans genotypes throughout the city, but these events were rare. Our analysis identified

highly clonal genomes of *L. interrogans* that persisted within certain populations of rats for up to

seven years. Finally, our genomic analysis of *L. interrogans* obtained from a human leptospirosis

case using DNA capture and enrichment strongly suggests a link to rats as the source and provides a

likely geographic origin within Boston where this transmission event might have occurred, although

we cannot rule out the possibility that the same genotype of *L. interrogans* was also carried by rats in

an unsampled location in Boston.

Materials and Methods

Study location and rodent trapping

The City of Boston, Massachusetts (125.4 sq. km), in the eastern US has an estimated

population of ~650,000 people (US Census Bureau 2020 Decennial Census) with a temperate climate

(cold winters and hot, humid summers). *Rattus norvegicus* collections between July 2016 and

November 2020 were conducted in collaboration with the City of Boston's Inspectional Services

Department (COB-ISD), as described previously (10, 11).

For rats trapped between 2021-2022, 6-10 traps were placed at each site (see site definitions

below) at least 15 meters apart in locations with evidence of recent rat activity (*i.e.*, near burrows or

along runs with evidence of fresh rat droppings/urine). Rats were trapped using a commercial lethal

snap trap (T-Rex, Bell Labs, Windsor, WI, US) fixed to a remote sensing device (RATMO®, Arctic

Systems, Copenhagen, DK) with zip-ties. Traps were baited with meat jerky (Slim-Jim original

flavor, Conagra Brand, Chicago, IL, US) and placed inside a weighted, locked commercial bait

station (Protecta EVO Express Bait Station, Bell Labs, Windsor, WI, US). When rats were trapped,

the remote sensing device alerted the study team in real time for prompt retrieval and reset of the

trap. Traps were placed in the evening, attended from dusk until dawn, and deactivated and shut

down the following morning. Trapping was conducted in accordance with protocols reviewed and

approved by the Cummings School of Veterinary Medicine IACUC (G2020-126).

Rat carcasses were stored in a cooler with icepacks until necropsies were performed within

six hours of trapping. Rats were weighed, sexed, dipped in 70% ethanol, and placed in dorsal

recumbency. Sterile tissue scissors and forceps were used to reflect the abdominal skin and a new set

of sterile instruments were used to open the abdomen and aseptically harvest the kidneys. For

samples collected in 2021-2022, one fresh kidney was placed into 50 mL conical tubes containing 10

mL liquid HAN medium (26) and shipped overnight at ambient temperature to the National Centers

for Animal Health (NCAH) in Ames, IA. Archived frozen kidneys (2016-2020) were stored at -80°C

until they were shipped overnight from Tufts University to NCAH on dry ice.

684   *lipL32* qPCR detection of *Leptospira* in rodent samples

We screened for the presence of *Leptospira* DNA in DNA extracted from 210 frozen kidneys,

118 fresh kidneys, and one fresh urine sample (kidney was also evaluated from this individual) from

328 *R. norvegicus* rats collected in Boston from 2016-2022; four kidneys from bycatch mice and one

kidney from a gray squirrel were also included. Kidneys were transferred to a 150 mm petri dish, the

capsule (fibrous outer layer of the kidney) removed, and then placed in a 710 mL Whirl-Pak bag

(World Bioproducts, LLC, Madison, Wisconsin, US) containing 10 mL of liquid HAN, and

immediately macerated manually. DNA was extracted using 500 μL of kidney macerate or 200 μL of

urine, and the Maxwell RSC Purefood Purification Pathogen kit (Promega Corporation, Madison,

Wisconsin, US). We screened all samples for the presence of pathogenic *Leptospira* spp. using a

TaqMan® (Thermo Fischer Scientific, Waltham, MA, US) qPCR assay specific for the *lipL32* gene

as previously described, using primer set LipL32-47Fd and LipL32-301Rd with LipL32-189P probe

(58-60). To control for qPCR inhibitors, we added TaqMan® exogenous Internal Positive Control to

the master mix to confirm DNA amplification, detect false negatives, and qualitatively detect

presence of amplification inhibitory substances. All samples were assayed in triplicate and

considered positive when duplicate or triplicates were positive with Ct values <40 (58, 60). Using

chi-square tests of independence, we examined associations between *Leptospira* positive rats and six

sampling variables: sex, maturity (juvenile vs. adult), sampling season [spring (19-Mar to 20-Jun),

summer (20-Jun to 22-Sept), fall (22-Sept to 21-Dec), and winter (21-Dec to 19-Mar)], year, site, and

rat genetic group (assigned herein).

Culture

Remaining tissue from frozen and fresh kidneys positive by *lipL32* qPCR was used for

culture. Macerate from frozen kidneys (200 μL) was inoculated into 5 mL of semisolid T80/40/LH

media (61) and incubated at 29°C. Two 10-fold serial dilutions were made from the fresh kidney

macerate into liquid HAN media in 3% CO_2 (26) and T80/40/LH media (61) and incubated at 37°C

and 29°C, respectively. Cultures were maintained for up to six months and checked periodically by

dark-field microscopy for the presence of leptospires.

Serotyping of isolates

714   *Leptospira* isolates were serotyped by the MAT with a panel of polyclonal rabbit reference

715   antisera representing 13 serogroups: Australis, Autumnalis, Ballum, Bataviae, Canicola,

716   Grippotyphosa, Hebdomadis, Icterohaemorrhagiae, Mini, Pomona, Pyrogenes, Sejroe, and Tarassovi.

717   The serogroup of each isolate was assigned according to the reference antiserum yielding the highest

718   agglutination titer.

Evaluation of virulence

One rodent isolate (designated strain R47) was propagated in liquid HAN medium at 37°C in

3% CO_2 and evaluated for virulence by intraperitoneal injection of 10^8 leptospires in a final volume

of 1 mL into one group of four female golden Syrian hamsters (*Mesocricetus auratus*). Negative

control hamsters were inoculated with media alone. Hamsters were monitored daily for clinical signs

of disease, including weight loss, lethargy, bloody discharge from the nose or urogenital tract, and

sudden death; and were euthanized when moribund. At time of euthanasia, liver and kidney tissue

were harvested for culture and *lipL32* qPCR. These experiments were conducted in accordance with

protocols as reviewed and approved by the Animal Care & Use Committee at the NADC and as

approved by USDA institutional guidelines.

Rattus norvegicus collection definition

Rattus norvegicus were trapped over seven years (2016-2022) at 64 trap locations (**S7 Table**)

within 17 geographic sites in Boston (**Fig 3**). Site numbers were assigned from east

to west and defined as trapping grids separated by ≤0.5 km. Exceptions

to this were trapping locations ≤0.5 km apart but separated by major

roadways, which are known dispersal barriers for *R. norvegicus* and *R. rattus* (6);

these were considered separate sites. A large city park complex (~0.83 km northeast to southwest)

comprised of two smaller parks [Boston Common (northeast) and Boston Public Garden (southwest)]

was considered a single site and is referred to hereafter as Boston Common Garden. We grouped

individual rats into 33 collections based upon site ID and sampling date, whereby samples collected

>4 months apart were considered unique collections. These criteria were selected based upon the ~3

months needed to reach sexual maturity for *R. norvegicus* (62) and previous work describing genetic

relatedness among *R. norvegicus* in urban settings with distances of up to 0.5 km (12). Thus,

nomenclature for collections is as follows: site is indicated by numerals ranging from 1-17 ordered

from east to west followed by letters corresponding to year of collection where a=2016, b=2017,

c=2018, d=2019, e=2020, f=2021, and g=2022. If multiple collections were obtained within a year, a

second numeral follows the letter designation (*e.g.,* 9g2 indicates the second collection at site 9 in

2022). These definitions resulted in 33 collections from 17 sites (**S1 Table**).

Three sites were sampled intensely between August 2021 and August 2022: Boston Common

Garden (site 6), Melnea Cass Blvd SE (site 8), and Melnea Cass Blvd N (site 9) (**Fig 3**). Boston

Common Garden is a heavily landscaped urban park system with primarily mowed grass ground

cover and concrete foot paths. The southwest and northeast portions are bisected by a 3-lane road,

and the southwest portion is the site of a historical municipal burial ground. The two Melnea Cass

Blvd sites are heavily developed urban areas of mixed commercial and residential use and separated

by an 8-lane road. These three sites were identified in consultation with the COB-ISD as having

active rat populations and burrow systems. They were selected for intensive sampling to facilitate a

deeper understanding of the relationship between rat population genetics and leptospirosis

transmission in parks versus mixed commercial/residential environments, and to assess how a major

road may impact dispersal among neighboring rat populations, which we hypothesize will impact the

movement of pathogenic *Leptospira*. Furthermore, we used these sites to assess how genetic

signatures change over time across stable rat populations (sites 6 and 9) and the effects of a rat

control intervention (baiting and trapping) at site 8 that began 5 November 2021.

Rattus norvegicus microsatellite development and genotyping

Marker selection, optimization, and multiplexing

Twenty-one MSAT markers were utilized for population genetic analyses of *R. norvegicus*

samples (**Table 1**). All markers have been previously published (14, 15, 17, 32, 63) and were

selected for use in this study based upon: 1) geographic regions where previously utilized (*i.e.*,

eastern US cities as a proxy for presumed genetic compatibility with our Boston study site), 2)

number of described alleles in other study systems (prioritizing markers with multiple alleles), and 3)

successful use among diverse populations of *R. norvegicus* as inferred by inclusion in multiple

studies from diverse western hemisphere countries (*i.e.,* US and Brazil). The 21 markers were

optimized using a panel of six *R. norvegicus* DNA samples from different sites in Boston, one

Sprague Dawley rat gDNA control (Zyagen, San Diego, CA, US), and a non-template control

(molecular grade water).

We generated multi-locus genotypes for 328 rat kidney DNAs using these 21 MSAT markers

amplified in nine PCR reactions (six multiplexes and three singles, **Table 1**). All PCRs were carried

out in 10 µL volumes containing the following reagents (given in final concentrations): 10-20 ng of

DNA template, 1x PCR buffer, 2.5 mM MgCl2, 0.2 mM dNTPs, 0.8 U Platinum Taq polymerase

(Invitrogen, Carlsbad, CA, US), and 0.1-0.5 µM of each primer pair (see **Table 1**). Thermocycling

used the following conditions: 10 minutes at 95°C to release the Platinum Taq® antibody, followed

by 38 cycles of 60 s at 94°C, 30 s at the annealing temperature (Ta) and 30 s at 72°C. The Ta,

dilution, multiplex, and pooling schemes for each marker are provided in **Table 1**. Diluted PCR

products were electrophoresed on an ABI 3130xl sequencer with LIZ®-1200 size standard (Applied

Biosystems, Foster City, CA, US) and analyzed using GeneMarker v2.6.7 (SoftGenetics, State

College, PA, US).

Table 1. *Rattus norvegicus* MSAT marker primers and conditions.

Marker	T_a	Mix	Primer pair conc. (µM)	Post-PCR dilution	Dye	A	Allele sizes (bp)	Citation and primer sequence (5'-3')
D12Rat76	60	1	0.5	1/50	NED	10	86, 88, 90, 92, 94, 96, 98, 100, 102, 104	[1] F-TGCCTTTTAAAATGATGTGCA R-ATTGGCAATGCACTCATGTG
D1Wox23	60	1	0.3	1/50	VIC	10	186, 190, 194, 198, 202, 206, 210, 214, 218, 222	[2] F-TCTGACCCATACTTGTACTTTGC R-AATTTCTGCCTCTTTTTCTCAG
D7Rat97	60	1	0.3	1/50	6FAM	11	172, 174, 176, 178, 180, 184, 186, 188, 192, 196, 198	[3] F-CAAGTTTTCCTCTGCCCAAG R-GCTGTCATTCCACTGGGTTT
D11Mgh5	60	2	0.4	1/75	6FAM	9	216, 226, 228, 230, 232, 234, 236, 238, 248	[1] F-CAGCTCTAATTCCAGAAAGGTTT R-GAATCGATTGACAGATGTCTGTG
D3Cebr3	60	2	0.3	1/75	PET	5	163, 165, 167, 177, 181	[4] F-CAGGGAATGCAGAAGATACAG R-GTGGCTTTAGGACTCTGGAG

	T_a					A		
D6Cebr1	60	2	0.4	1/75	NED	7	220, 224, 226, 230, 232, 234, 236	[4] F-TGGTTTGGTTGGGGAGAA R-GTGCTGTCAGGGAAAGATGTA
D4Wox7	55	3	0.1	1/50	VIC	5	130, 134, 138, 142, 146	[2] F-GATAGCATAAAATCCCTAGAGGTT R-TCGATTTATCTGAAACCATCAC
D6Wox2	55	3	0.3	1/50	6FAM	10	96, 100, 104, 108, 116, 120, 124, 128, 132, 136	[2] F-CCAGTCCATACTTATCCATCTG R-CATTTAGATAGGTGATAGATTCAG
D1Cebr9	55	3	0.3	1/50	6FAM	6	261, 263, 265, 267, 269, 271	[4] F-GGATTTGGCTCCCTTTAAG R-CAGTAACTCTGGTTCATGTACTCC
D7Rat13	64	4	0.4	1/25	NED	12	128, 144, 146, 148, 150, 152, 154, 156, 158, 160, 162, 164	[5] F-GACTTCTGCTACACGCCACA R-CAGCCCTAGAAGGAAATGCA
D5Rat83	64	4	0.4	1/25	VIC	11	164, 168, 170, 178, 184, 186, 190, 192, 194, 196, 200	[5] F-ACTTGGAAACAGGGAGATGG R-GGGTCTTCAGGATGGCAATGT
D12Wox1	64	4*	0.4	1/50	6FAM	8	392, 396, 400, 404, 408, 412, 416, 420	[2] F-GACATTAAGGGGTCTTCCTAAG R-TATCTTTGCAACGCTGAGG
D10Rat20	64	5	0.4	1/25	PET	6	109, 113, 115, 117, 119, 123	[1] F-AGTGATTGCCATACCTGCCT R-GAAATGGCCAGGATAAACCA
D20Rat46	64	5	0.4	1/25	6FAM	13	140, 142, 144, 148, 154, 156, 160, 162, 164, 168, 172, 174, 180	[1] F-AAGTACTGAGTGGGCTGCGT R-GGCAAAACACCAATGCCTAT
D10Mit5	64	5	0.2	1/25	NED	9	135, 137, 139, 141, 143, 145, 147, 149, 151	[1] F-TGCTGGGTGAACCAGAGAG R-CTGCCCTCCAAACCACC
D9Rat13	64	5	0.4	1/25	VIC	5	101, 109, 119, 121, 123	[5] F-CCCATCTTTACACCTCCCAA R-GGAAAGGAAACTGGAGGGTC
D19Wox11	66	6	0.3	1/15	NED	5	212, 216, 220, 224, 228	[2] F-CTACCCACCCATCTATTCATCC R-GTTTCCAGCACCCATGTCC
D1Cebr3	66	6	0.3	1/15	VIC	6	79, 81, 83, 105, 107, 109	[4] F-CTTGGGAGCTGGGAGTGT R-GAAGGCTGAGGTATGAAGACTG
D16Rat81	66	6	0.4	1/15	PET	7	148, 150, 152, 154, 156, 158, 160	[1] F-GAGCCTTAGCACAGTGGCTT R-GGCCCACATGTGCATGTATA
D5Rat33	66	6*	0.4	1/15	6FAM	15	102, 106, 110, 112, 116, 118, 120, 122, 124, 126, 130, 132, 134, 138, 140	[3] F-TGGAGAAAAGAAGAACCTCCA R-GTGCCCTCAGACTGAACTC
D1Cebr4	66	6*	0.4	1/30	VIC	8	278, 284, 286, 288, 290, 292, 294, 296	[4] F-GACCTCCTGCCCTTCACTG R-TGAAAAATGAATTGCTTGTG

788 [a][1] (64), [2] (65), [3] (66), [4] (67), [5] (63).
789 [b]Additional single PCRs are pooled into a corresponding mix (*i.e.*, multiplex PCR) prior to capillary
790 electrophoresis. T_a = PCR annealing temperature. A = number of observed alleles in Boston, MA
791 *Rattus norvegicus* (2016-2019 and 2021-2022).
792

793 **MSAT sample data filters**

Our final MSAT dataset included only rats with complete genotypes

at ≥19 markers; one individual sample failed to meet this criterion and

was removed. We also removed three individuals with evidence of DNA

cross contamination as indicated by the presence of >2 alleles at

multiple markers. Finally, location provenance was incomplete for 13

individuals (including all eight rats collected during 2020), which were

also removed. This resulted in a final MSAT dataset of 311 *R. norvegicus*

representing 33 collections from 17 sites collected between 2016-2019 and 2021-2022 (**S1 Table**).

MSAT marker validation analyses

We ran a series of tests on the 21 MSAT markers to validate neutrality and utility in

population genetic analyses of *R. norvegicus* from Boston using all 311 individuals

organized into 33 collections. Validations included a query for identical

multi-locus genotypes in GenAlEx (68) and a test for linkage

disequilibrium in FSTAT v2.9.4 based on the log-likelihood ratio G

statistic (69). We used MICRO-CHECKER (70) to test for null alleles, large

allelic dropouts, and scoring errors due to PCR stuttering using the

following settings: maximum expected allele size of 420 bp, a 95%

confidence interval (CI), and 1000 repetitions. We performed a test of

Hardy-Weinberg equilibrium for each marker, also in GenAlEx, but using

only collections with ≥20 rats (*i.e.*, 5b, 6c, 6g2, and 8f, **see S1 Table**). Finally, we ran

815   PCR replicates on a subset of 94 *R. norvegicus* DNAs (30.2%) from the final dataset to check for

816   allele scoring errors.

Population genetic analyses: determination of genetic groups and population assignment using STRUCTURE

We determined the most likely number of genetic groups and conducted population

assignment analyses using STRUCTURE v2.3.4 analysis (71) on the MSAT genotypes obtained from

the 311 *R. norvegicus* DNAs. We set Bayesian parameters to assume genetic admixture and

correlated allele frequencies in the study system, and to ignore collection sites. These parameters

were run at K values from 1–15, with 25,000 burn-in iterations followed by 100,000 run iterations.

Bayesian analysis was repeated ten times for each K. Resulting log(Prob K) values were analyzed

with the delta-K method (72) to estimate the most likely number of genetic populations. We assigned

rats to a specific genetic group and identified potential migrants if their probability of membership

(Q) to a specific genetic group was ≥0.75. Individual rats not meeting

these criteria were grouped broadly into an "admixed" group, regardless

of collection site (**S3 Table**). Using these results, we sorted all 311 rats according to their

predicted genetic groups [12 distinct genetic groups (n=247) and one admixed group (n=64)] for

additional population genetic analyses [~24% of individuals (64 admixed and 11 potential migrants,

see Results) were no longer grouped according to their initial collection site].

Population genetic analyses: estimates of genetic structure at two scales

We estimated overall genetic structure among Boston *R. norvegicus* using PCoA and three

different estimators of F_{ST}. For PCoA, we used standardized genetic distances among all pairs of

individuals in GenAlEx (68). We then conducted *a posteriori* F_{ST} analyses using only individuals

from the 12 genetic groups identified in STRUCTURE (n=247) but also removed juveniles (weight

<140g) (73) and migrants, which could artificially increase or decrease, respectively, F_{ST} estimates

among the 12 predicted genetic groups. Collection 11b was the single exception because only

juveniles were trapped and they appeared to form a distinct genetic group (Orange, **see Results**).

These filters resulted in a final dataset for this analysis of 119

individual rats among the 12 genetic groups; all genetic groups were

represented by ≥4 individuals (up to 19; **S3 Table**). We generated F_{ST} estimates of

theta (θ) in FSTAT version 2.9.4 using Weir and Cockerham's method (74) to estimate population

structure. Pairwise population F_{ST} values were also estimated in FSTAT and *p*-values for all

comparisons were generated using a =0.05. We performed AMOVA on this same reduced set

(n=119) using standardized genetic distances in GenAlEx with 999 permutations to estimate PhiPT

(an F_{ST} analog). The third F_{ST} estimator was a one-way global ANOSIM in PRIMER v5.2.9

(PRIMER-e, Albany, AK, NZ), conducted on a reduced set of 247 individual rats from the 12 distinct

genetic groups (the 64 individuals forming the admixed group were removed but migrants and

juveniles were included; **S3 Table**); the R statistic was generated using 10,000 permutations.

Afterwards, a post-hoc test was conducted to determine the genetic groups contributing to R, wherein

pairwise comparisons among all genetic groups were conducted. The global alpha was divided by 66

(number of pairwise comparisons) to calculate a conservative Bonferroni corrected a =0.000758.

Population genetic analyses: isolation by distance

Because *R. norvegicus* relatives often live in close proximity, we performed a fine-scale

analysis of spatial genetic structure (SGS) on all 311 rats using spatial autocorrelation in GenAlEx

with options set to SinglePOP, even distance classes (n=20), and 999 permutations. Distance class

intervals were increased in 100 m increments (0-100 m, 0-200 m, etc.) up to a maximum of 0-2,000

m. We assessed isolation by distance using the RELATE function in PRIMER v5.2.9 and

standardized genetic distances and geographic locations (latitude and longitude) of the 247 rats

forming the 12 distinct genetic groups (**S3 Table**). We generated Spearman's rank correlation

coefficient (Rho) using 10,000 permutations. Inbreeding coefficient F_{IS} was estimated for each

genetic group and global relatedness was estimated among all groups using the reduced set of 119

rats (no juveniles or migrants) in FSTAT (**S3 Table**).

Leptospira genomics and genotyping

Whole genome sequencing of isolates

Whole genome sequencing of DNA extracted from 5 mL cultures of each isolate was

performed as previously described (75, 76). Briefly, each isolate was propagated in HAN liquid

media at 29°C, which was harvested by centrifugation at 10,000 x *g* for 15 min and DNA extracted

using the Maxwell RSC Purefood Purification Pathogen kit. For Illumina library preparation, the

concentration of reconstituted genomic DNA was determined with a Qubit dsDNA HS assay using a

Qubit® 3.0 fluorometer (Invitrogen, Carlsbad, CA, US). Whole-genome sequences were obtained

(MiSeq Desktop Sequencer, 2x250 v2 paired-end chemistry and the Nextera XT DNA Library

Preparation Kit, Illumina, San Diego, California, US) per manufacturer's instructions.

Capture and enrichment of *Leptospira* DNA from rat and human samples

We attempted *Leptospira* genome capture and enrichment from 22 PCR positive rat samples

collected between 2017-2019, 20 culture negative and two culture positive (R316 and R47); the latter

served as genome analysis controls. We also attempted enrichment on two separate human samples

from Boston: one serum from 2017 (MA-1) and one concentrated urine from 2018 (MA-2). These

sample DNAs were provided by the CDC and had *lipL32* PCR Ct values of 38.03 and 31.6,

respectively. DNA from these 24 samples were subjected to pan pathogenic *Leptospira* DNA capture

and enrichment as previously described (28). Briefly, sample DNAs were diluted separately to ~4

ng/µL in a volume of 55 µL, sonicated to an average size of 285 bp using a Q800R2 sonicator

(QSonica, Newtown, CT, US), and then single indexed libraries were prepared using Agilent Sure-

Select methodology (XT-HS kit, Agilent, Santa Clara, CA, US). The libraries (some pooled) were

subjected to one or two rounds of DNA capture and enrichment (**S5 Table**) and sequenced on an

Illumina MiSeq instrument using either a MiSeq v3 600 cycle (2x300 reads) or v2 300 cycle kit

(2x150 reads). Three samples (R309, R334-19, and R379) failed to generate successful sequencing

libraries and were excluded from DNA capture and enrichment and Illumina sequencing and six

others failed to enrich *Leptospira* DNA and were excluded from sequencing (R66, R67, R108, R318,

R324 and MA-1) (**S2 and S5 Tables**). These unsuccessful attempts were associated with low

abundance samples [PCR Ct values >38 (n=2)], enzymatic inhibition [failed library preparation

(n=3)], or perhaps contained divergent or non-pathogenic species of *Leptospira* or some other

genetically similar species not compatible with our pan pathogenic *Leptospira* DNA capture and

enrichment system (n=4 including all three *Leptospira*-positive rats from collection 5b). Pooling

schemes, rounds of DNA capture and enrichment, and Illumina sequencing kits used for each sample

are provided in **S5 Table**.

Read classifications, read mapping, SNP calling, and phylogenomics

To estimate the percentage of *Leptospira* reads in enriched sequences and for taxonomic

classification of enrichments and isolates, reads were mapped against the standard Kraken database

with Kraken v2.1.2 (77). To determine breadth of coverage for enriched genomes, reads were aligned

against *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (GenBank accession#

GCA_000007685.1) with minimap2 v2.22 (78) and the per base depth of coverage was calculated

with Samtools v1.6 (79). Two *Leptospira* phylogenies were built. The first included a diverse set of

L. interrogans (representing >9 serovars) for the purpose of contextualizing Boston rat and human

derived genomic diversity within *L. interrogans*. Single nucleotide polymorphisms were identified

among 28 genomes from cultured isolates (GenBank accession numbers provided in phylogeny), 21

publicly available *L. interrogans* genomes, and 13 enriched genomes (12 rats and 1 human) by

aligning reads against reference genome *L. interrogans* serovar Copenhageni strain

Fiocruz L1-130 using minimap2 v2.22 and calling SNPs from the BAM file

with GATK v4.2.2 (80) using a depth of coverage ≥3x and a read proportion

of 0.9. SNPs that fell within duplicated regions, based on a reference

self-alignment with NUCmer v3.1 (81), were filtered from downstream

analyses. All methods were wrapped by NASP v1.2.1 (82). A maximum

likelihood phylogeny was inferred on the concatenated SNP alignments

using IQ-TREE v2.2.0.3 with default parameters (83), the "-fast" option,

and the integrated ModelFinder method (84); the phylogeny was midpoint

rooted. Second, to provide increased genomic resolution within *L.*

interrogans and assess fine scale relatedness among the Boston genomes we replaced the 21 diverse

L. interrogans genomes with two complete, publicly available *L. interrogans* serovar Copenhageni

genomes (GenBank accession# GCA_002073495.2 and GCA_010978155.1). A maximum likelihood

phylogeny was inferred on the concatenated SNP alignments using IQ-TREE v2.2.0.3 as described

above, except with 1000 bootstrap replicates; the phylogeny was rooted with *L. interrogans* serovar

Copenhageni strain Fiocruz L1-130.

Leptospira interrogans reference phylogeny and WG-FAST placement of enriched samples

Two enriched samples (R289 and R292) had lower breadth and depth of coverage than the other enrichments (**S5 Table**) and were inserted into the phylogeny with *WG-FAST* v1.2 (85) using default settings. Reads were simulated from genome assemblies with ART vMountRainier (86) using the command "-p -na -ss MSv3 -l 250 -f 75 -m 300 -s 30". Based on the position of the enriched reads in the complete *L. interrogans* phylogeny (**see Results**), simulated reads were aligned against *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 with minimap2 v2.24 (78) and SNPs were called with GATK v4.2.6.1 (80). A maximum likelihood phylogeny was inferred on the concatenated SNPs with RAxML-NG v. 1.1.0 (87) and rooted with *L. interrogans* serovar Copenhageni strain Fiocruz L1-130.

Human and rat phylogenomic comparison

To provide additional resolution among *Leptospira* genomes derived from four *R. norvegicus* and one human that grouped together within a single *L. interrogans* serovar Copenhageni clade (designated as clade 2, **see Results**), we conducted a phylogenomic analysis using NASP v1.2.1 and IQ-TREE v2.2.0.3 as described above, except *L. interrogans* serovar Copenhageni strain SK1 (GenBank accession# GCA_010978155.1) was used as the reference and root.

AmpSeq methods

We performed highly multiplexed amplicon sequencing (AmpSeq) on two subsets of rat samples that were PCR positive for *Leptospira* but not included (or were unsuccessful) in the above steps for culturing or genome enrichment. The first subset consisted of rats from 2016 and 2020-2022 that were PCR positive but culture negative (n=11); we included one culture positive rat (Cad99) to

serve as an analysis control. The final subset for AmpSeq contained any 2017-2019 positive samples

that failed DNA capture and enrichment (n=8 rats and n=1 human) (**S2 Table**).

In total, 42 primer sets were designed and validated for the *Leptospira* AmpSeq system.

Thirteen primer sets were either modified from existing primer sets (60, 88-98) targeting genes

commonly used for molecular typing or detection of *Leptospira* (*lipl32*, 16S, *secY*, *Loa22*, *OmpL1*,

flaB, LIC_11108, LFB1, *ligB*), or redesigned from MLST loci (*glmU*, *lipL41*, *icdA*, *sucA*) (99-101) to

be compatible with AmpSeq (153-415 bp in length) (**S8 Table**). We also designed 29 new primer

sets targeting loci determined, via comparative genomic analyses, to be SNP-rich and highly

discriminatory among 778 genome assemblies from 64 pathogenic and non-pathogenic *Leptospira*

spp. (**S9 Table**). Assemblies were downloaded from the NCBI RefSeq database (102, 103) using

ncbi-genome-download v0.3.1 https://github.com/kblin/ncbi-genome-download/tree/0.3.1 and

aligned to reference genome *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 using

NUCmer (104), and SNPs were identified using NASP v1.2.0 (82). The SNPs were analyzed using

VaST (105) to identify a set of targets maximizing discriminatory power among the assembled

genomes. To avoid paralogous sequence variants and loci with highly variable alleles, we only

analyzed SNPs located within a genus-wide core genome (106). To maximize amplification of these

targets across *Leptospira* `species, primers were designed with ambiguous bases`

`for variants that occurred in ≥5% of genome assemblies.`

To generate a conservative estimate of species coverage and discrimination for this AmpSeq

system, we used *in silico* PCR to predict amplification for each of the 42 loci across 69 *Leptospira*

spp. genome assemblies listed on the International Leptospirosis Society's resource webpage

(accessed 15-Jan-2024; https://leptosociety.org/resources/); these genomes represent all validated

pathogenic (P1 and P2) and saprophytic (S1 and S2) *Leptospira* spp. (57, 107). Genomes were

downloaded with the ncbi-genome-download tool (https://github.com/kblin/ncbi-genome-download)

from the RefSeq database (102) and renamed based on annotation information in the RefSeq record.

PCR primers were screened against the 69 *Leptospira* genomes with the -search_pcr2 method in

USEARCH v11.0.667 (108) with maximum number of SNP mismatches allowed in each primer set

to two, using parameters: "-strand both -maxdiffs 2 -maxamp 500". Predicted genome hits were

compiled for each primer set. This *in silico* analysis predicted amplification in all 69 *Leptospira*

species. The number of expected loci for each species varied, ranging from a single locus to all 42

loci. The pathogenic P1 clade was well represented with predicted amplification of 18-42 loci in all

20 species (**S1 File**).

AmpSeq was performed on two sample sets. The first set was used to validate the method and

consisted of nine genomic DNA extracts obtained from isolates from seven P1 clade *Leptospira* spp.

[*L. interrogans* (n=2), *L. kirschneri*, *L. santarosai*, *L. noguchii* (n=2), *L. borgpetersenii*, *L.*

alexanderi, and *L. kmetyi*]. The second set contained unknown *Leptospira* spp. and consisted of DNA

from 20 *Leptospira* PCR positive *R. norvegicus* kidney samples and one human serum sample (MA-

1) (**S2 Table**) along with *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (ATCC,

Manassas, Virginia US, catalog# BAA-1198D-5), which served as a PCR and analysis control, and

also molecular grade water as a non-template negative control. The 42 loci were arranged into four

multiplexes and PCRs were carried out in 10 µL volumes containing the following reagents (given in

final concentrations): 10-20 ng of DNA template, 1x PCR buffer, 2.5 mM MgCl2, 0.2 mM dNTPs,

0.8 U Platinum Taq polymerase (Invitrogen, Carlsbad, CA, US), and 0.1-0.4 µM of each primer pair

(**see S8 Table**). Thermocycling utilized the following conditions: 10 minutes at 95°C to release the

Platinum Taq® antibody, followed by 38 cycles of 60 s at 94°C, 30 s at 55°C and 30 s at 72°C. PCRs

were pooled together in equal amounts in a final volume of 25 µL (6.25 µL of each multiplex) and

sequence libraries were prepared for all positive samples as previously described (56, 109). Uniquely

indexed sample libraries were pooled together in equimolar amounts and sequenced in parallel on an

Illumina NextSeq instrument using a 600 cycle (2x300) P1 kit (Illumina, San Diego, CA, US).

1006

AmpSeq analysis

1007

Read mapping and classification

1008

Illumina sequencing reads for each individual sample were analyzed using the Amplicon

Sequencing Analysis Pipeline (ASAP; https://github.com/TGenNorth/ASAP), wherein reads from

each locus are mapped against a reference database using Bowtie2 and binned according to alignment

score (110); the number of paired end reads assigning to each reference are then counted. To build a

comprehensive reference database of *Leptospira* species to be used for read mapping and

classification, we implemented two strategies. First, we downloaded reference sequences from

GenBank for the thirteen previously published loci and removed redundant sequences. Second, for

the 29 new loci we bioinformatically extracted each locus from the reference set of 69 *Leptospira*

spp. genome assemblies described above. We first amplified each locus separately (using the same

PCR conditions above for the AmpSeq multiplexes) using gDNA from *L. interrogans* serovar

Copenhageni strain Fiocruz L1-130. PCR products were visualized on a 2% agarose gel to ensure

robust amplification and treated with ExoSAP-IT (Affymetrix, Santa Clara, CA, US) using 1 µL of

ExoSAP-IT and 7 µL of PCR product under the following conditions: 37°C for 15 minutes followed

by 80°C for 15 minutes. Treated products were then diluted (based on amplicon intensity) and

sequenced using forward primer UT1F-acccaactgaatggagc and reverse primer UT2R-

acgcacttgacttgtcttc (universal sequences incorporated into each unique primer pair, **see S8 Table**) in a

BigDye® Terminator v3.1 Ready Reaction Mix (Applied Biosystems, Foster City, CA, US). We

used 10 µL volumes for sequencing reactions containing the following reagents (given in final

concentrations): 5x Sequencing Buffer, 1 µL BigDye® Terminator v3.1 Ready Reaction Mix, 1 µM

primer, and 5 μL diluted PCR product. Thermocycling conditions were: 96°C for 20 seconds,

followed by 30 cycles of 96°C for 10 seconds, 50°C for 5 seconds, and 60°C for 4 minutes. Sanger

sequences were assembled, trimmed, and converted to fasta format using SeqMan Pro (DNASTAR

Lasergene, Madison, WI, USA), which were used to extract each locus (if present) from the 69

Leptospira spp. genome assemblies using BLASTN (111) and a custom python script

(https://gist.github.com/jasonsahl/2a232947a3578283f54c).

1034

Whole genome versus AmpSeq phylogenetic resolution

We compared phylogenetic resolution and tree topology among two maximum likelihood

phylogenies generated, respectively, from whole genome reads and AmpSeq reads. Using 24 genome

assemblies representing 18 P1 clade pathogenic *Leptospira* spp. (GenBank accession numbers

provided in the phylogeny), we conducted a whole genome phylogenetic analysis using NASP v1.2.1

and IQ-TREE v2.2.0.3 as described above with 1000 bootstrap replicates and *L. interrogans* serovar

Copenhageni strain L1-130 as the reference. We then conducted an identical analysis but also

included the AmpSeq generated reads from the nine *Leptospira* isolate genomic DNAs in our

validation set. Inclusion of AmpSeq generated reads reduced the core genome to only nucleotide

positions 1) included in the 42 loci, and 2) shared among the genome (n=24) and AmpSeq (n=9)

reads. Both phylogenies were midpoint rooted.

1046

1047   **Species id and within species genotyping**

1048   Rat and human DNAs that amplified at ≥7 loci (out of 42) with >10x

1049   coverage were assigned a *Leptospira* species classification. To facilitate genotyping within

1050   species for samples with high breadth of coverage (*i.e.,* 25-42 loci amplified with >10x coverage),

1051   SNPs were identified from AmpSeq reads and a reference set of 22 publicly available *L. interrogans*

complete genome assemblies (GenBank accession numbers provided in the phylogeny). AmpSeq

reads contain Illumina adaptors, universal tail sequences, and locus specific primers, all of which

must be removed prior to SNP discovery if precise phylogenetic placement is the analysis goal;

adaptors and universal tails were removed in ASAP and locus specific primers were trimmed using

cutPrimers (https://github.com/aakechin/cutPrimers) (112). Trimmed AmpSeq reads and genome

assemblies were aligned against reference genome *Leptospira kirschneri* serovar Grippotyphosa

strain RedPanda1 (GenBank accession# GCA_027563495.1) using NASP v1.2.1 (82). A maximum

likelihood phylogeny was then inferred on the concatenated SNP alignments using IQ-TREE v2.2.0.3

with the "-fast" option, default parameters (83), and the integrated ModelFinder method (84); the

phylogeny was rooted with *Leptospira kirschneri* serovar Grippotyphosa strain RedPanda1.

Acknowledgments

The authors extend their gratitude to the City of Boston's Inspectional Services Department and the

Boston Public Health Commission for their assistance. The authors thank the *Mycobacteria* &

Brucella Section from the NVSL for their technical assistance in genome sequencing. USDA is an

equal opportunity provider and employer. Mention of trade names or commercial products in this

publication is solely for the purpose of providing specific information and does not imply

recommendation or endorsement by the U.S. Department of Agriculture. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the NIH. Any

opinions, findings, and conclusions or recommendations expressed in this material are those of the

authors and do not necessarily reflect the views of NSF. The findings and conclusions in this report

are those of the authors and do not necessarily represent the official position of the Centers for

Disease Control and Prevention.

Funding Disclosure

This research was supported in part by an appointment to the Animal and Plant Health Inspection

Service (APHIS) Research Participation Program administered by the Oak Ridge Institute for

Science and Education (ORISE) through an interagency agreement between the U.S. Department of

Energy (DOE) and the U.S. Department of Agriculture (USDA). ORISE is managed by ORAU under

DOE contract number DE-SC0014664. This project was also funded in part by the National Institute

of Allergy and Infectious Diseases in the National Institutes of Health (NIH) award number

R01AI172924 (Pearson), the National Institute of Environmental Health Sciences under award

number K08ES035460 (Rosenbaum), the National Center for Advancing Translational Sciences,

NIH under award number KL2TR002545 (Rosenbaum, Hertzel, Benson, Specht), and K-24 AI

106822 (Phipatanakul). Work conducted by JDM was completed while supported by the NSF

Postdoctoral Research Fellowships in Biology Program, Award No. 1907311 (Madison).

Competing Interests

The authors declare they have no competing interests.

Author Contributions

Nathan E. Stone: Investigation, Methodology, Project administration, Supervision, Validation,

Visualization, Writing original draft, Writing review and editing

Camila Hamond: Investigation, Methodology, Validation, Writing original draft, Writing review and

editing

Joel Clegg: Investigation, Validation

Ryelan F. McDonough: Investigation, Validation

Reanna M. Bourgeois: Investigation, Validation

Rebecca Ballard: Investigation, Validation

Natalie B. Thornton: Investigation

Marianece Nuttall: Investigation

Hannah Hertzel: Investigation

Tammy Anderson: Investigation, Validation, Writing review and editing

Ryann N. Whealy: Investigation, Validation

Skylar Timm: Investigation

Alexander K. Roberts: Investigation

Verónica Barragán: Investigation

Wanda Phipatanakul: Funding acquisition, Investigation

Jessica H. Leibler: Investigation

Hayley Benson: Investigation

Aubrey Specht: Investigation

Ruairi White: Methodology, Investigation, Resources

Karen LeCount: Funding acquisition, Investigation, Resources, Supervision, Writing – review &

editing

Tara N. Furstenau: Investigation, Methodology, Writing original draft

Renee L. Galloway: Investigation, Methodology

Nichola J. Hill: Investigation, Methodology

Joseph D. Madison: Funding acquisition, Investigation, Methodology, Writing reviewing and editing

Viacheslav Y. Fofanov: Investigation, Methodology

Talima Pearson: Funding acquisition, Project administration

Jason W. Sahl: Investigation, Methodology, Supervision, Writing review and editing

Joseph D. Busch: Project administration, Supervision, Writing original draft, Writing review and

editing

Zachary Weiner: Project administration

Jarlath E. Nally: Funding acquisition, Investigation, Methodology, Project administration,

Supervision, Writing original draft, Writing review and editing

David M. Wagner: Funding acquisition, Investigation, Methodology, Project administration,

Supervision, Writing original draft, Writing review and editing

Marieke Rosenbaum: Funding acquisition, Investigation, Methodology, Project administration,

Supervision, Writing original draft, Writing review and editing

1132

References

1.   Lund M. Commensal Rodents. Rodent Pests and Their Control, 2nd Edition. 2015:19-32.
2.   Desvars-Larrive A, Pascal M, Gasqui P, Cosson JF, Benoit E, Lattard V, et al. Population genetics,
community of parasites, and resistance to rodenticides in an urban brown rat (Rattus norvegicus) population.
PLoS One. 2017;12(9):e0184015.
3.   Harper GA, Bunbury N. Invasive rats on tropical islands: Their population biology and impacts on native
species. Glob Ecol Conserv. 2015;3:607-27.
4.   Kosoy M, Khlyap L, Cosson JF, Morand S. Aboriginal and Invasive Rats of Genus *Rattus* as Hosts of
Infectious Agents. Vector-Borne Zoonot. 2015;15(1):3-12.
5.   Meerburg BG, Singleton GR, Kijlstra A. Rodent-borne diseases and their risks for public health. Crit Rev
Microbiol. 2009;35(3):221-70.
6.   Byers KA, Lee MJ, Patrick DM, Himsworth CG. Rats About Town: A Systematic Review of Rat Movement
in Urban Ecosystems. Frontiers in Ecology and Evolution. 2019;7.
7.   Feng AYT, Himsworth CG. The secret life of the city rat: a review of the ecology of urban Norway and
black rats (Rattus norvegicus and Rattus rattus). Urban Ecosyst. 2014;17(1):149-62.
8.   Sabour S, Azimi T, Nasser A, Hadi N, Mohsenzadeh A, Shariati A. A global overview of the most
important zoonotic bacteria pathogens transmitted from Rattus norvegicus to humans in urban environments.
Infect Med (Beijing). 2022;1(3):192-207.
9.   Himsworth CG, Parsons KL, Jardine C, Patrick DM. Rats, cities, people, and pathogens: a systematic
review and narrative synthesis of literature regarding the ecology of rat-associated zoonoses in urban centers.
Vector Borne Zoonotic Dis. 2013;13(6):349-59.
10.   Gerbig GR, Piontkivska H, Smith TC, White R, Mukherjee J, Benson H, et al. Genetic characterization of
Staphylococcus aureus isolated from Norway rats in Boston, Massachusetts. Vet Med Sci. 2023;9(1):272-81.

11.	Cummings CO, Hill NJ, Puryear WB, Rogers B, Mukherjee J, Leibler JH, et al. Evidence of Influenza A in Wild Norway Rats (Rattus norvegicus) in Boston, Massachusetts. Front Ecol Evol. 2019;7.

12.	Combs M, Byers KA, Ghersi BM, Blum MJ, Caccone A, Costa F, et al. Urban rat races: spatial population genomics of brown rats (Rattus norvegicus) compared across multiple cities. Proc Biol Sci. 2018;285(1880).

13.	Combs M, Puckett EE, Richardson J, Mims D, Munshi-South J. Spatial population genomics of the brown rat (Rattus norvegicus) in New York City. Mol Ecol. 2018;27(1):83-98.

14.	Gardner-Santana LC, Norris DE, Fornadel CM, Hinson ER, Klein SL, Glass GE. Commensal ecology, urban landscapes, and their influence on the genetic characteristics of city-dwelling Norway rats (Rattus norvegicus). Mol Ecol. 2009;18(13):2766-78.

15.	Kajdacsi B, Costa F, Hyseni C, Porter F, Brown J, Rodrigues G, et al. Urban population genetics of slum-dwelling rats (Rattus norvegicus) in Salvador, Brazil. Mol Ecol. 2013;22(20):5056-70.

16.	Byers KA, Booker TR, Combs M, Himsworth CG, Munshi-South J, Patrick DM, et al. Using genetic relatedness to understand heterogeneous distributions of urban rat-associated pathogens. Evol Appl. 2021;14(1):198-209.

17.	Richardson JL, Burak MK, Hernandez C, Shirvell JM, Mariani C, Carvalho-Pereira TSA, et al. Using fine-scale spatial genetics of Norway rats to improve control efforts and reduce leptospirosis risk in urban slum environments. Evol Appl. 2017;10(4):323-37.

18.	Levett PN. Leptospirosis. Clin Microbiol Rev. 2001;14(2):296-326.

19.	Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS, et al. Global Morbidity and Mortality of Leptospirosis: A Systematic Review. PLoS Negl Trop Dis. 2015;9(9):e0003898.

20.	Sterling CR, Thiermann AB. Urban rats as chronic carriers of leptospirosis: an ultrastructural investigation. Vet Pathol. 1981;18(5):628-37.

21.	Thiermann AB. The Norway Rat as a Selective Chronic Carrier of Leptospira-Icterohaemorrhagiae. J Wildlife Dis. 1981;17(1):39-43.

22.	Monahan AM, Callanan JJ, Nally JE. Proteomic analysis of *Leptospira interrogans* shed in urine of chronically infected hosts. Infect Immun. 2008;76(11):4952-8.

23.	Bonilla-Santiago R, Nally JE. Rat model of chronic leptospirosis. Curr Protoc Microbiol. 2011;Chapter 12:Unit 12E 3.

24.	Haake DA, Levett PN. Leptospirosis in humans. Curr Top Microbiol Immunol. 2015;387:65-97.

25.	Boey K, Shiokawa K, Rajeev S. Leptospira infection in rats: A literature review of global prevalence and distribution. PLoS Negl Trop Dis. 2019;13(8):e0007499.

26.	Hornsby RL, Alt DP, Nally JE. Isolation and propagation of leptospires at 37 degrees C directly from the mammalian host. Sci Rep-Uk. 2020;10(1).

27.	Musso D, La Scola B. Laboratory diagnosis of leptospirosis: a challenge. J Microbiol Immunol Infect. 2013;46(4):245-52.

28.	Stone NE, McDonough RF, Hamond C, LeCount K, Busch JD, Dirsmith KL, et al. DNA Capture and Enrichment: A Culture-Independent Approach for Characterizing the Genomic Diversity of Pathogenic Leptospira Species. Microorganisms. 2023;11(5):1282.

29.	Cole JR, Jr., Sulzer CR, Pursell AR. Improved microtechnique for the leptospiral microscopic agglutination test. Appl Microbiol. 1973;25(6):976-80.

30.	Chirathaworn C, Inwattana R, Poovorawan Y, Suwancharoen D. Interpretation of microscopic agglutination test for leptospirosis diagnosis and seroprevalence. Asian Pac J Trop Biomed. 2014;4(Suppl 1):S162-4.

31.	Pritchard J. Documentation for STRUCTURE software: Version 2.3 2003. Available from: https://web.stanford.edu/group/pritchardlab/structure_software/release_versions/v2.3.4/structure_doc.pdf.

32.	Costa F, Richardson JL, Dion K, Mariani C, Pertile AC, Burak MK, et al. Multiple Paternity in the Norway Rat, Rattus norvegicus, from Urban Slums in Salvador, Brazil. J Hered. 2016;107(2):181-6.

33.	Glass GE, Klein SL, Norris DE, Gardner LC. Multiple Paternity in Urban Norway Rats: Extended Ranging for Mates. Vector Borne Zoonotic Dis. 2016;16(5):342-8.

34.   Davis DE. The characteristics of rat populations. Q Rev Biol. 1953;28(4):373-401.

35.   Davis DE, Emlen JT, Stokes AW. Studies on Home Range in the Brown Rat. J Mammal. 1948;29(3):207-
25.

36.   Easterbrook JD, Shields T, Klein SL, Glass GE. Norway rat population in Baltimore, Maryland, 2004.
Vector Borne Zoonotic Dis. 2005;5(3):296-9.

37.   Berthier K, Garba M, Leblois R, Navascués M, Tatard C, Gauthier P, et al. Black rat invasion of inland
Sahel: insights from interviews and population genetics in south-western Niger. Biol J Linn Soc.
2016;119(4):748-65.

38.   Traweger D, Slotta-Bachmayr L. Introducing GIS-modelling into the management of a brown rat (Rattus
norvegicus Berk.) (Mamm. Rodentia Muridae) population in an urban habitat. J Pest Sci. 2005;78(1):17-24.

39.   Worth CB. Field and Laboratory Observations on Roof Rats, Rattus-Rattus (Linnaeus), in Florida. J
Mammal. 1950;31(3):293-304.

40.   Petrie GF, Todd RE. Observations in Upper Egypt on the range of excursion of the house rodents: R.
rattus and Acomys   cahirinus. Rep Notes Pub Hlth Lab Cairo. 1923;5:14-8.

41.   Glass GE. Comparative ecology and social interactions of Norway rat (Rattus norvegicus) populations
in Baltimore, Maryland. Lawrence, Kan. :: Museum of Natural History, the University of Kansas; 1989.

42.   Davis DE, Christian JJ. Population Consequences of a Sustained Yield Program for Norway Rats. Ecology.
1958;39(2):217-22.

43.   Margulis HL. Rat Fields, Neighborhood Sanitation, and Rat Complaints in Newark, New-Jersey. Geogr
Rev. 1977;67(2):221-31.

44.   Vadell MV, Cavia R, Suárez OV. Abundance, age structure and reproductive patterns of Rattus
norvegicus and Mus musculus in two areas of the city of Buenos Aires. Int J Pest Manage. 2010;56(4):327-36.

45.   Wundram IJ, Ruback RB. Urban Rats - Symbol, Symptom and Symbiosis. Hum Organ. 1986;45(3):212-
9.

46.   Costa F, Ribeiro GS, Felzemburgh RD, Santos N, Reis RB, Santos AC, et al. Influence of household rat
infestation on leptospira transmission in the urban slum environment. PLoS Negl Trop Dis. 2014;8(12):e3338.

47.   de Faria MT, Calderwood MS, Athanazio DA, McBride AJ, Hartskeerl RA, Pereira MM, et al. Carriage of
Leptospira interrogans among domestic rats from an urban setting highly endemic for leptospirosis in Brazil.
Acta Trop. 2008;108(1):1-5.

48.   Costa F, Porter FH, Rodrigues G, Farias H, de Faria MT, Wunder EA, et al. Infections by Leptospira
interrogans, Seoul virus, and Bartonella spp. among Norway rats (Rattus norvegicus) from the urban slum
environment in Brazil. Vector Borne Zoonotic Dis. 2014;14(1):33-40.

49.   Himsworth CG, Bidulka J, Parsons KL, Feng AY, Tang P, Jardine CM, et al. Ecology of Leptospira
interrogans in Norway rats (Rattus norvegicus) in an inner-city neighborhood of Vancouver, Canada. PLoS Negl
Trop Dis. 2013;7(6):e2270.

50.   Peterson AC, Ghersi BM, Riegel C, Wunder EA, Jr., Childs JE, Blum MJ. Amplification of pathogenic
Leptospira infection with greater abundance and co-occurrence of rodent hosts across a counter-urbanizing
landscape. Mol Ecol. 2021;30(9):2145-61.

51.   Santos LA, Adhikarla H, Yan X, Wang Z, Fouts DE, Vinetz JM, et al. Genomic Comparison Among Global
Isolates of L. interrogans Serovars Copenhageni and Icterohaemorrhagiae Identified Natural Genetic Variation
Caused by an Indel. Front Cell Infect Microbiol. 2018;8:193.

52.   Lee MJ, Byers KA, Donovan CM, Bidulka JJ, Stephen C, Patrick DM, et al. Effects of Culling on Leptospira
interrogans Carriage by Rats. Emerg Infect Dis. 2018;24(2):356-60.

53.   Thibeaux R, Soupe-Gilbert ME, Kainiu M, Girault D, Bierque E, Fernandes J, et al. The zoonotic pathogen
Leptospira interrogans mitigates environmental stress through cyclic-di-GMP-controlled biofilm production.
NPJ Biofilms Microbiomes. 2020;6(1):24.

54.   Santos AAN, Ribeiro PDS, da Franca GV, Souza FN, Ramos EAG, Figueira CP, et al. Leptospira interrogans
biofilm formation in Rattus norvegicus (Norway rats) natural reservoirs. PLoS Negl Trop Dis.
2021;15(9):e0009736.

55.   Davignon G, Cagliero J, Guentas L, Bierque E, Genthon P, Gunkel-Grillon P, et al. Leptospirosis: toward
a better understanding of the environmental lifestyle of. Front Water. 2023;5.
56.   Stone NE, Hall CM, Ortiz M, Hutton SM, Santana-Propper E, Celona KR, et al. Diverse lineages of
pathogenic Leptospira species are widespread in the environment in Puerto Rico, USA. PLoS Negl Trop Dis.
2022;16(5):e0009959.
57.   Vincent AT, Schiettekatte O, Goarant C, Neela VK, Bernet E, Thibeaux R, et al. Revisiting the taxonomy
and evolution of pathogenicity of the genus *Leptospira* through the prism of genomics. PLoS Negl Trop Dis.
2019;13(5):e0007270.
58.   Galloway RL, Hoffmaster AR. Optimization of LipL32 PCR assay for increased sensitivity in diagnosing
leptospirosis. Diagn Microbiol Infect Dis. 2015;82(3):199-200.
59.   Hamond C, Browne AS, de Wilde LH, Hornsby RL, LeCount K, Anderson T, et al. Assessing rodents as
carriers of pathogenic Leptospira species in the U.S. Virgin Islands and their risk to animal and public health.
Sci Rep. 2022;12(1):1132.
60.   Stoddard RA, Gee JE, Wilkins PP, McCaustland K, Hoffmaster AR. Detection of pathogenic *Leptospira*
spp. through TaqMan polymerase chain reaction targeting the LipL32 gene. Diagn Microbiol Infect Dis.
2009;64(3):247-55.
61.   Ellis WA, Montgomery J, Cassells JA. Dihydrostreptomycin treatment of bovine carriers of *Leptospira*
interrogans serovar hardjo. Res Vet Sci. 1985;39(3):292-5.
62.   Modlinska K, Pisula W. The Norway rat, from an obnoxious pest to a laboratory pet. Elife. 2020;9.
63.   Lack JB, Hamilton MJ, Braun JK, Mares MA, Van den Bussche RA. Comparative phylogeography of
invasive Rattus rattus and Rattus norvegicus in the US reveals distinct colonization histories and dispersal. Biol
Invasions. 2013;15(5):1067-87.
64.   Jacob HJ, Brown DM, Bunker RK, Daly MJ, Dzau VJ, Goodman A, et al. A genetic linkage map of the
laboratory rat, Rattus norvegicus. Nat Genet. 1995;9(1):63-9.
65.   Heiberg AC, Leirs H, Siegismund HR. Reproductive success of bromadiolone-resistant rats in absence
of anticoagulant pressure. Pest Manag Sci. 2006;62(9):862-71.
66.   Robertson BC, Gemmell NJ. Defining eradication units to control invasive pests. J Appl Ecol.
2004;41(6):1042-8.
67.   Giraudeau F, Apiou F, Amarger V, Kaisaki PJ, Bihoreau MT, Lathrop M, et al. Linkage and physical
mapping of rat microsatellites derived from minisatellite loci. Mamm Genome. 1999;10(4):405-9.
68.   Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching
and research--an update. Bioinformatics. 2012;28(19):2537-9.
69.   Goudet J. FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3.2).
2002.
70.   Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. MICRO-CHECKER: software for identifying
and correcting genotyping errors in microsatellite data. Mol Ecol Notes. 2004;4(3):535-8.
71.   Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype
data. Genetics. 2000;155(2):945-59.
72.   Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software
STRUCTURE: a simulation study. Mol Ecol. 2005;14(8):2611-20.
73.   Macdonald DW, Barrett P. Mammals of Europe. Princeton, N.J.: Princeton University Press; 1993. 312
p., 126 p. of plates p.
74.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure. Evolution.
1984;38(6):1358-70.
75.   Hamond C, Dirsmith KL, LeCount K, Soltero FV, Rivera-Garcia S, Camp P, et al. Leptospira borgpetersenii
serovar Hardjo and Leptospira santarosai serogroup Pyrogenes isolated from bovine dairy herds in Puerto Rico.
Front Vet Sci. 2022;9:1025282.
76.   Hamond C, LeCount K, Putz EJ, Bayles DO, Camp P, Goris MGA, et al. Bovine Leptospirosis Due to
Persistent Renal Carriage of Leptospira borgpetersenii Serovar Tarassovi. Front Vet Sci. 2022;9:848664.

77.	Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol.
2019;20(1):257.
78.	Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34(18):3094-100.
79.	Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format
and SAMtools. Bioinformatics. 2009;25(16):2078-9.
80.	McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res.
2010;20(9):1297-303.
81.	Delcher AL, Salzberg SL, Phillippy AM. Using MUMmer to identify similar regions in large sequence
sets. Curr Protoc Bioinformatics. 2003;Chapter 10:Unit 10 3.
82.	Sahl JW, Lemmer D, Travis J, Schupp JM, Gillece JD, Aziz M, et al. NASP: an accurate, rapid method for
the identification of SNPs in WGS datasets that supports flexible input and output formats. Microb Genom.
2016;2(8):e000074.
83.	Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm
for estimating maximum-likelihood phylogenies. Molecular biology and evolution. 2015;32(1):268-74.
84.	Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model
selection for accurate phylogenetic estimates. Nat Methods. 2017;14(6):587-9.
85.	Sahl JW, Schupp JM, Rasko DA, Colman RE, Foster JT, Keim P. Phylogenetically typing bacterial strains
from partial SNP genotypes observed from direct sequencing of clinical specimen metagenomic data. Genome
Med. 2015;7(1):52.
86.	Huang W, Li L, Myers JR, Marth GT. ART: a next-generation sequencing read simulator. Bioinformatics.
2012;28(4):593-4.
87.	Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and user-friendly
tool for maximum likelihood phylogenetic inference. Bioinformatics. 2019;35(21):4453-5.
88.	Barragan V, Chiriboga J, Miller E, Olivas S, Birdsell D, Hepp C, et al. High Leptospira Diversity in Animals
and Humans Complicates the Search for Common Reservoirs of Human Disease in Rural Ecuador. PLoS Negl
Trop Dis. 2016;10(9):e0004990.
89.	Merien F, Portnoi D, Bourhy P, Charavay F, Berlioz-Arthaud A, Baranton G. A rapid and quantitative
method for the detection of Leptospira species in human leptospirosis. FEMS Microbiol Lett. 2005;249(1):139-
47.
90.	Matthias MA, Lubar AA, Lanka Acharige SS, Chaiboonma KL, Pilau NN, Marroquin AS, et al. Culture-
Independent Detection and Identification of Leptospira Serovars. Microbiol Spectr. 2022;10(6):e0247522.
91.	Backstedt BT, Buyuktanir O, Lindow J, Wunder EA, Jr., Reis MG, Usmani-Brown S, et al. Efficient
Detection of Pathogenic Leptospires Using 16S Ribosomal RNA. PLoS One. 2015;10(6):e0128913.
92.	Iwasaki H, Chagan-Yasutan H, Leano PS, Koizumi N, Nakajima C, Taurustiati D, et al. Combined antibody
and DNA detection for early diagnosis of leptospirosis after a disaster. Diagn Microbiol Infect Dis.
2016;84(4):287-91.
93.	Benacer D, Zain SNM, Lewis JW, Khalid M, Thong KL. A duplex endpoint PCR assay for rapid detection
and differentiation of Leptospira strains. Rev Soc Bras Med Trop. 2017;50(2):239-42.
94.	Cerqueira GM, McBride AJ, Hartskeerl RA, Ahmed N, Dellagostin OA, Eslabao MR, et al. Bioinformatics
describes novel Loci for high resolution discrimination of leptospira isolates. PLoS One. 2010;5(10):e15335.
95.	Di Azevedo MIN, Borges A, Kremer F, de Melo J, Carvalho-Costa FA, Lilenbaum W. Genetic Analysis of
LigA, LipL32, Loa22, and OmpL1 Antigens from Leptospira spp. Sejroe Serogroup: How Close We Are To a
Vaccine Against Bovine Leptospirosis? Curr Microbiol. 2023;80(9):310.
96.	Ahmed A, Engelberts MF, Boer KR, Ahmed N, Hartskeerl RA. Development and validation of a real-time
PCR for detection of pathogenic *Leptospira* species in clinical materials. PLoS One. 2009;4(9):e7093.
97.	Raheliniriha S, Moseley MH, Allan KJ, Ramanohizakandrainy E, Ravaoarinoro S, Rajerison M, et al.
Leptospira in livestock in Madagascar: uncultured strains, mixed infections and small mammal-livestock
transmission highlight challenges in controlling and diagnosing leptospirosis in the developing world.
Parasitology. 2019;146(14):1707-13.

98.   Haake DA, Suchard MA, Kelley MM, Dundoo M, Alt DP, Zuerner RL. Molecular evolution and mosaicism
of leptospiral outer membrane proteins involves horizontal DNA transfer. J Bacteriol. 2004;186(9):2818-28.
99.   Ahmed N, Devi SM, Valverde Mde L, Vijayachari P, Machang'u RS, Ellis WA, et al. Multilocus sequence
typing method for identification and genotypic classification of pathogenic *Leptospira* species. Ann Clin
Microbiol Antimicrob. 2006;5:28.
100.   Boonsilp S, Thaipadungpanit J, Amornchai P, Wuthiekanun V, Bailey MS, Holden MT, et al. A single
multilocus sequence typing (MLST) scheme for seven pathogenic Leptospira species. PLoS Negl Trop Dis.
2013;7(1):e1954.
101.   Varni V, Ruybal P, Lauthier JJ, Tomasini N, Brihuega B, Koval A, et al. Reassessment of MLST schemes
for Leptospira spp. typing worldwide. Infect Genet Evol. 2014;22:216-22.
102.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference sequence (RefSeq)
database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res.
2016;44(D1):D733-45.
103.   Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI prokaryotic
genome annotation pipeline. Nucleic Acids Res. 2016;44(14):6614-24.
104.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile and open software
for comparing large genomes. Genome Biol. 2004;5(2):R12.
105.   Furstenau TN, Cocking JH, Sahl JW, Fofanov VY. Variant site strain typer (VaST): efficient strain typing
using a minimal number of variant genomic sites. BMC Bioinformatics. 2018;19(1):222.
106.   Guglielmini J, Bourhy P, Schiettekatte O, Zinini F, Brisse S, Picardeau M. Genus-wide Leptospira core
genome multilocus sequence typing for strain taxonomy and global surveillance. PLoS Negl Trop Dis.
2019;13(4):e0007374.
107.   Sykes JE, et al.,. Understanding leptospirosis: application of state-of-the-art molecular typing tools with
a One Health lens. AJVR. 2022;In Press.
108.   Edgar RC. Search and clustering orders of magnitude faster than BLAST. Bioinformatics.
2010;26(19):2460-1.
109.   Colman RE, Schupp JM, Hicks ND, Smith DE, Buchhagen JL, Valafar F, et al. Detection of Low-Level
Mixed-Population Drug Resistance in *Mycobacterium tuberculosis* Using High Fidelity Amplicon Sequencing.
Plos One. 2015;10(5).
110.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9(4):357-9.
111.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and
applications. BMC Bioinformatics. 2009;10:421.
112.   Kechin A, Boyarskikh U, Kel A, Filipenko M. cutPrimers: A New Tool for Accurate Cutting of Primers
from Reads of Targeted Next Generation Sequencing. J Comput Biol. 2017;24(11):1138-43.

1388

Supporting Information

1390   **S1 Fig. Plots from the delta-*K* method.** The hypothetical number of subpopulations (*K*) is plotted

1391   on the X-axis against A) delta-*K*, and B) The logarithm probability for each *K* [L(*K*)]. Ten

1392   independent runs in STRUCTURE were performed for each *K* value.

1393

S2 Fig. Principle Coordinates Analysis plot. Principle Coordinates Analysis plot of standardized

genetic distances illustrating genetic similarity among 311 *Rattus norvegicus* collected in

Boston (2016-2022). Each rat is color coded according to its probability

of membership (Q ≥0.75) to one of 12 genetic groups assigned using

STRUCTURE (**Table S3 and Fig 1**). The percentage of genetic variation explained by these two

axes is indicated.

S3 Fig. Spatial autocorrelation analysis graph. Spatial autocorrelation analysis graph assessing the

correlation between genetic and geographic distance (Y-axis) using twenty even distance classes

corresponding to 100 m intervals (X-axis). The blue line is the correlation value (r), and the red

dashed lines are the 95% confidence intervals (U=upper and L=lower) about the null hypothesis of

no spatial structure as determined by bootstrap resampling. Values of r above the upper 95%

confidence interval are significant.

S4 Fig. *Leptospira interrogans* whole genome phylogeny. A maximum likelihood midpoint rooted

phylogeny of 28 *Leptospira interrogans* serogroup Icterohaemorrhagiae isolate genomes and 13

enriched genomes from 40 rats and one human (bold text), together with 22 publicly available *L.*

interrogans isolate genomes representing >9 serovars (GenBank accession numbers are provided in

the figure annotations), based upon a concatenated SNP alignment of 27,678 positions out of a core

genome size of 1,186,500 nucleotides (nts).

S5 Fig. *Leptospira interrogans* serogroup Icterohaemorrhagiae WG-FAST phylogeny A

maximum likelihood phylogeny of 28 *Leptospira interrogans* serogroup Icterohaemorrhagiae isolate

genomes from rats, together with two publicly available *L. interrogans* serogroup

Icterohaemorrhagiae serovar Copenhageni complete genomes (GenBank accession#

GCA_002073495.2 and GCA_010978155.1), and 13 enriched genomes (12 rats and 1 human), based

upon a concatenated SNP alignment of 127 positions out of a core genome size of 2,123,028

nucleotides (nts). The phylogeny was rooted with reference genome *L. interrogans* serogroup

Icterohaemorrhagiae serovar Copenhageni strain Fiocruz L1-130 (GenBank accession#

GCA_000007685.1). The color of the genome ID corresponds to the rat genetic group in Fig 1; black

text indicates that no genetic group was assigned. Two enriched samples (R289 and R292) had lower

breadth and depth of coverage than the other enrichments (**see S5 Table**) and were inserted into the

rat phylogeny based upon canonical SNPs with WG-FAST.

1427

S6 Fig. Pathogenic *Leptospira* whole genome and AmpSeq phylogenies. A) Whole genome

maximum likelihood midpoint rooted phylogeny including 18 P1 clade pathogenic *Leptospira* spp.

B) compared to an AmpSeq phylogeny that also includes AmpSeq data generated from nine

Leptospira isolate gDNAs (blue text). Species specific clades were maintained (color coded) and

similar phylogenetic topologies were observed despite a large reduction in the core-genome of the

AmpSeq tree (1,195 nts) compared to the whole genome tree (167,849 nts). Bootstrap values are

indicated at specific nodes. The nine AmpSeq *Leptospira* isolate gDNAs grouped in their respective

species clades.

1436

S7 Fig. *Leptospira interrogans* AmpSeq phylogeny. Maximum likelihood phylogeny based upon an

alignment of 4,260 shared nucleotide positions using *Leptospira* AmpSeq data generated from nine

lipL32 qPCR positive *Rattus norvegicus* (bold text) along with 22 *Leptospira interrogans* isolate

genomes (GenBank accession numbers are provided in the figure annotations). The phylogeny is

rooted with *Leptospira kirschneri* serovar Grippotyphosa strain RedPanda1 (GenBank accession#

GCA_027563495.1).

1443

S1 Table. *Rattus norvegicus* collections.

S2 Table. Analyses summary and results for all rodent and human samples.

S3 Table. *Rattus norvegicus* population assignment.

S4 Table. *Rattus norvegicus* FSTAT results

S5 Table. DNA capture and enrichment setup and results.

S6 Table. *Leptospira* AmpSeq results

S7 Table. *Rattus norvegicus* trapping locations.

S8 Table. *Leptospira* AmpSeq primers and conditions

S9 Table. *Leptospira* AmpSeq primer design genomes

S1 File: Predicted amplification of 42 AmpSeq loci against 69 *Leptospira* spp.

1454

Data Availability Statement

1455

MSAT multi-locus genotypes for 311 *R. norvegicus* are provided in **S2 Table**. Whole genome

sequencing reads have been deposited in GenBank under BioProject PRJNA1080226 with SRA

accession numbers SRR28374870-SRR28374875, SRR28374877-SRR28374878, SRR28374880,

SRR28374882-SRR28374900 for isolate genomes and SRR28374876, SRR28374879,

SRR28411353-SRR28411357, SRR28411359-SRR28411365, SRR28411370 for enriched genomes.

Raw AmpSeq data for up to 42 loci are also provided in GenBank under the same BioProject with

SRA accession numbers SRR28374881, SRR28411338-SRR28411352, SRR28411358,

SRR28411366-SRR28411369 and SRR28765937- SRR28765945.

1464

Core genome: 3,280,516 nts
33 variable sites

Enrichment: 16d-2019-R366

9f-2021-MAR_47

9g1-2022-Cad86

16d-2019-R367

Enrichment: 2018-Human-MA-2

3 SNPs

GCA_010978155.1 *Leptospira interrogans* serovar Copenhageni strain SK1