Title: The Roles of Drift and Selection on Short Stamen Loss in *Arabidopsis thaliana* along an Elevational Gradient in the Spanish Pyrenees

Authors: Sophia F. Buysse[1,2,3], Samuel G. Pérez[1,2,3], Joshua R. Puzey[4], Ava Garrison[1,2,3], Gideon S. Bradburd[5], Christopher G. Oakley[6], Stephen J. Tonsor[7], F. Xavier Picó[8], Emily B. Josephs[1,2,9]*, Jeffrey K Conner[1,2,3]*

Affiliations

[1]Plant Biology Department, Michigan State University, East Lansing, MI 48824, USA

[2]Ecology, Evolution, and Behavior Program, Michigan State University, East Lansing, MI 48824, USA

[3]Kellogg Biological Station, Hickory Corners, MI 49060, USA

[4]Department of Biology, William & Mary, Williamsburg, VA 23187, USA

[5]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109, USA

[6]Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907, USA

[7]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260

[8]Departamento de Biología Evolutiva, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Sevilla 41092, Spain

[9]Plant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA

*Co-Corresponding Authors:

Jeff Conner

3700 E. Gull Lake Drive

Hickory Corners, MI 49060

connerj@msu.edu

Emily Josephs

612 Wilson Rd, Rm 166

East Lansing MI 48824 USA

Josep993@msu.edu

Author Contributions: SGP collected the phenotypic data and conceived the research with JKC. JRP oversaw the sequencing. AG and GSB did preliminary analyses and contributed to conceptual development. CGO, SJT and FXP provided seed material and guidance in project development. SFB

analyzed the data and drafted the manuscript with guidance from EBJ and JKC. JKC managed the project.
All authors provided comments on the manuscript.

Funding: Research reported was supported by the National Institute of General Medical Sciences of the
National Institutes of Health (NIH) under award R35GM137919 (awarded to GSB) and R35GM142829
(awarded to EBJ), and by National Science Foundation awards DEB 0919452 (to JKC) and DEB 2223962
(to JKC and GSB). The content is solely the responsibility of the authors and does not necessarily
represent the official views of the NIH or NSF.

Data Availability: These sequence data will be submitted to the GenBank database upon manuscript
acceptance. All phenotypic data and code will be made available on GitHub upon acceptance. GWAS
output files will be made available on figshare upon acceptance.

Conflict of Interest: The authors declare no conflicts of interest.

Acknowledgements:

We thank D. Schemske for guidance throughout the initial stages of the project, N. Zhang for library
preparation, and A. Platts for bioinformatics assistance. This manuscript was improved by comments
from Conner lab and Josephs lab members.

Abstract

Traits that have lost function sometimes persist through evolutionary time. These traits may be maintained by a lack of standing genetic variation for the trait, if selection against the trait is weak relative to drift, or if they have a residual function. To determine the evolutionary processes shaping whether nonfunctional traits are retained or lost, we investigated short stamens in 16 populations of *Arabidopsis thaliana* along an elevational cline in the Spanish Pyrenees. We found a cline in short stamen number from retention of short stamens in high elevation populations to incomplete loss in low elevation populations. We did not find evidence that limited genetic variation constrains the loss of short stamens at high elevations nor evidence for divergent selection on short stamens between high and low elevations. Finally, we identified loci associated with short stamens in the Spanish Pyrenees that are different from loci associated with variation in short stamen number across latitudes from a previous study. Overall, we did not identify the evolutionary mechanisms maintaining an elevational cline in short stamen number but did identify different genetic loci underlying the variation in short stamen along similar phenotypic clines.

Keywords (provide 3 to 6): – genetic drift, trait loss, selection, effective population size, stamen, elevational cline

Teaser text

The evolutionary mechanisms underlying loss or retention of traits that have lost function are poorly understood. Short stamens in *Arabidopsis thaliana* provide a compelling system to investigate the roles of genetic drift and selection in trait loss across latitudinal and elevational clines. This study investigates the role of drift and selection in short stamen loss in 16 populations of *A. thaliana* along an elevational gradient in the Spanish Pyrenees. An investigation of the genetic loci underlying variation in short stamen number suggests mutations in different genes may cause trait loss in similar phenotypic clines within a species.

Introduction

Traits that have lost function are often lost through evolutionary time, yet some persist. Nonfunctional
traits could be lost through direct selection against the trait (Dorken et al., 2004; Lahti et al., 2009),
correlated responses to selection on other traits caused by pleiotropy or linkage disequilibrium
(Yoshizawa et al., 2012), or the accumulation of selectively neutral mutations (Fong et al., 1995). In
contrast, nonfunctional traits may be maintained by evolutionary constraint if the nonfunctional trait is
genetically correlated with a different, functional trait (Lande, 1979; Walsh & Blows, 2009).
Nonfunctional trait loss could also be prevented if there is not enough standing genetic variation for
selection to act on the trait (Lahti et al., 2009) or if weak selection against the trait is unable to
overcome drift (Charlesworth, 2009). Both of these processes may be pronounced in species with small
effective population sizes. Determining the evolutionary processes shaping whether nonfunctional traits
are retained or lost will have broad implications for our understanding of the interplay of direct
selection, correlated responses to selection, and genetic drift (reviewed in Futuyma, 2010).

Here, we use an elevational cline in short stamen number in *Arabidopsis thaliana* to investigate how
evolutionary processes interact to shape variation in a trait that has lost function. Almost all of the 3,700
species in the Brassicaceae family have flowers with four long and two short stamens (Endress, 1992).
While the function of short stamens is unknown, maintaining the stamen length difference is likely
adaptive in wild radish, *Raphanus raphanistrum,* an outcrossing species within Brassicaceae (Conner et
al., 2003; Waterman et al., 2023). However, *A. thaliana* evolved to be almost entirely self-fertilizing from
an outcrossing ancestor between 0.5 – 1 million years ago (Durvasula et al., 2017; Tang et al., 2007). In
flowering plants, a transition from outcrossing to self-pollination and subsequent relaxation of selection
for pollination often results in a suite of trait changes, called the 'selfing syndrome', that includes
smaller flowers and reduced distance between the anthers and stigma (Sicard & Lenhard, 2011). Self-
pollination also decreases local effective population size (Caballero, 1994), increasing the effects of
genetic drift.

Short stamens in *A. thaliana* do not contribute significantly to self-fertilized seed number, but have
nonetheless been retained in most natural populations (Royer et al., 2016). In particular, southern
European populations are more likely to lose short stamens while northern European populations retain
both short stamens. Because northern European populations of *A. thaliana* have undergone repeated
bottlenecks that reduced effective population size, and thus standing genetic variation (Beck et al.,
2008; François et al., 2008; Lewandowska-Sabat et al., 2010), short stamens may be retained in these
populations because there is insufficient genetic variation for stamen number and/or weak selection
against stamens cannot overcome drift. Alternatively, the latitudinal cline of the European populations
could be maintained by local adaptation for short stamen number or another correlated trait. The same
evolutionary processes may shape elevational clines in short stamen number. High elevations were
glaciated, even at southern latitudes (Hughes et al., 2006) and while the Spanish Pyrenees do include
putative refugia regions (Médail & Diadema, 2009), prior work in the Pyrenees has shown high elevation
montane populations have less genetic variation and smaller effective population sizes than lower
elevation coastal populations (Gomaa et al., 2011; Montesinos et al., 2009). Further, because
environmental factors vary similarly with latitude and elevation, selection may result in similar
phenotypic clines. Yet, latitudinal and elevational phenotypic clines are not always parallel (Daco et al.,
2021; Kooyers et al., 2015) and when parallel phenotypic clines are identified, they may not have a
parallel genetic basis (Fulgione et al., 2022; Gamba et al., 2022).

To understand the processes shaping variation in short stamen loss, we assessed 16 populations of *A. thaliana* along an elevational gradient in the Spanish Pyrenees. We counted short stamen number in a growth-chamber common garden and sequenced multiple individuals per population to quantify genetic variation. We identified an elevational cline in short stamen number in the Spanish Pyrenees, with more stamen loss at low elevation, similar to the previously described latitudinal cline (Royer et al., 2016). We then used multiple regression, polygenic adaptation detection, and genome wide association studies to address possible mechanisms for maintaining the cline:

1) Is the short stamen loss cline maintained by variation in effective population size? Less stamen loss in populations with less genetic variation after accounting for variation due to elevation would be consistent with the hypothesis that stamen loss is constrained by a lack of genetic variation and less effective selection.

2) Is the short stamen loss cline maintained by local adaptation? Evidence for divergent selection on short stamen number between high and low elevation populations would support local adaptation maintaining the cline.

3) What genomic loci are associated with short stamen number? If we identify loci within the three previously identified QTL (Royer et al., 2016), then this would support parallel genomic evolution between the latitudinal and elevational clines in short stamen number.

Materials and Methods

Seed Collection

Population collection sites are along an elevation gradient across the Pyrenees and along the Mediterranean coast (Fig. 1A, Table S1). Seeds from five to nine individuals were haphazardly collected from multiple patches in each of 16 populations (n = 112 genotypes) in northeast Spain (Montesinos-Navarro et al., 2011). These lineages are also used in Montesinos-Navarro et al. (2011), Montesinos-Navarro et al. (2012), and Wolfe and Tonsor (2014). They include the 10 populations from Montesinos et al. (2009) and the 9 populations from Gomaa et al. (2011).

Phenotyping

To assess genetic differentiation for short stamen number, plants were grown in chamber common gardens in three blocks at Michigan State University, East Lansing, MI. Seeds were stratified at 4°C for five days before we increased temperatures to 22°C:18°C for 16-hour days at a constant 60% humidity for four weeks. After emergence, plants were thinned to one seedling per cell in a 200 cell plug tray. One to four plants from each of the 112 genotypes (median = 2, n= 230 plants) were vernalized at 4°C with 10-hour days for six weeks before returning to 22°C:18°C for 16-hour days through flowering.

On each plant, we counted stamen number (Fig. 1B) on up to three flowers at each of up to three timepoints throughout the duration of flowering to estimate short stamen number (1 to 9 flowers per plant, median = 6, n= 1,423 flowers). Population mean short stamen number was calculated as the arithmetic mean from all flowers phenotyped in a population (42 to 169 flowers per population, mean = 88.9, median = 99.5). The arithmetic population means were highly correlated with estimated marginal means for each population calculated with the *emmeans* package (Lenth, 2021) in R (v4.2.2 (R Core Team, 2021)) from a model with population as a fixed effect, random effects of flowering timepoint

nested within plant nested within genotype which is nested within population, and random effect for
block (r= 0.975, p < 0.001).

We sequenced the entire genomes of a subset of 61 genotypes (see below). The 61 genotypes include
representatives from all 16 populations (3 or 4 different genotypes per population). In the sequenced
genotypes, stamen number was counted on up to four plants per genotype (median = 2, n = 141) and up
to nine flowers per plant (median = 7, n = 971). For all analyses that incorporate both phenotypic and
SNP information, arithmetic population means were recalculated as the mean from flowers scored for
short stamen number from sequenced genotypes (18 to 118 flowers per population, median = 63). The
population means from all flowers and the population means from only the sequenced genotypes are
highly correlated (r = 0.985, p < 0.001) indicating the sequenced genotypes are representative of all
plants scored for short stamen number.

Sequencing

A subset of 61 genotypes (3-4 per population) were chosen for Ilumina paired-end whole-genome 150bp
sequencing using the WGS-Novogene platform. Nextera adapter sequences were trimmed from the raw
sequence data with Trim Galore (Krueger, 2019). We also clipped the first 15 bp of each read because
quality checks with FastQC (Andrews, 2019) and MultiQC (Ewels et al., 2016) showed an identical,
unusual, pattern in the first 10 bases of each read: GTTTTAAACT. Reads were mapped to the TAIR10
reference genome (Berardini et al., 2015) using BWA mem with default settings (Li & Durbin, 2009). The
mean mapping rate for properly paired reads was 96.6% with a mean of 15,224,790 properly paired and
mapped reads per genotype and a total of 943,936,960 mapped and paired reads in the dataset. The
median depth across genotypes was low but acceptable at 8X. There is variation in coverage, missing
data, and quality scores between genotypes, but no genotypes were excluded on this basis (Table S2).
Duplicate reads were marked with the Genome Analysis Toolkit (GATK) v4.1.4.1 (Van der Auwera &
O'Connor, 2020) MarkDuplicates Spark. After an initial round of Haplocaller and GenotypeGVCF, the
dataset was filtered with the parameters suggested by GATK best practices (Van der Auwera &
O'Connor, 2020): QD <2, FS > 60, MQ < 40, ReadPosRankSum < -8, and MQRankSum < -12.5. The filtered
file was used as a known variants file for base quality score recalibration (BQSR). Variants were then
called again with HaplotypeCaller using the GVCF flag to keep all sites before combining samples for
GenotypeGVCF with the all-sites flag to create a dataset that includes both variant and invariant sites
(116,855,685 total sites).

Testing the drift hypotheses

To estimate the effects of past drift and gene flow in each population, genome-wide pairwise nucleotide
diversity (pi) was calculated for each population as an approximate measure of effective population size
with pixy v1.0.4 (Korunes & Samuk, 2021) from a filtered dataset containing variant and invariant sites
based on the pixy protocol. We used quality filters to remove all indels and low-quality sites that met
the following criteria: quality score less than 20, mean depth less than 3, and more than 25% missing
data. The filtered dataset contains 99,202,614 sites. A peak in nucleotide diversity was observed in the
centromere of each chromosome, likely caused by fewer mapped sites (Korunes and Samuk, pers. com.).
However, excluding previously published centromeres (Clark et al., 2007) did not meaningfully change
genome-wide nucleotide diversity (r = 0.998, p <0.001), so centromeres are included in all of the
following analyses (results excluding the centromeres can be found in Figs. S5-S7).

To test if variation in effective population size is maintaining a cline in short stamen number, we used
multiple regression to identify how short stamen number was predicted by nucleotide diversity and
elevation. While elevation is correlated with climatic variables such as precipitation and temperature
(Montesinos-Navarro et al., 2011) that could be selective agents for plant traits, we used elevation
because previous work in these same 16 populations demonstrated that elevation explained 54% of the
variance in trait principal component 1 (traits include: phenology, water use efficiency, instantaneous
CO_2 and H_2O exchange, PSII quantum efficiency, and specific leaf area) while the climate PC1 only
explained 36% of the trait variation (Wolfe & Tonsor, 2014). We then visualized these results by using
the residuals of single regression models. We used *lme4* in R for all models (Bates et al., 2015).

A negative relationship between nucleotide diversity and short stamen number after correcting for
elevation would support the hypothesis that smaller effective population sizes are constraining trait loss
due to low standing variation and less effective selection. A relationship between short stamen number
and elevation after correcting for nucleotide diversity could be evidence for divergent selection because
it indicates the short stamen number cline persists beyond the variation explained by nucleotide
diversity. However, this relationship could also result from shared evolutionary history, sometimes
called shared genetic drift, if relatedness among populations is associated with elevation (Colautti & Lau,
2015). Therefore, we further investigated evidence for divergent selection by testing for local adaptation
in short stamen number.

Testing for local adaptation

We used plink v1.9 (Chang et al., 2015) to filter the all-sites output from GATK to a "variant sites only"
dataset for conducting genetic principal component analysis (PCA). The "variant sites only" dataset was
also used for a genome-wide association study (GWAS; see below). We used quality filters to remove all
indels and non-biallelic sites in addition to low quality sites that met the following criteria: minor allele
frequency less than 5%, quality score less than 25, mean depth less than 5, and more than 25% missing
data. The variant sites dataset contains 1,858,706 SNPs. Filtering parameters were chosen to maintain
the same percent nonsynonymous sites (annotated by snpEff (Cingolani et al., 2012)) as stricter
parameters while retaining more variants in the dataset. Genetic PCA was calculated for the first 20 PCs
with plink v1.9 (Chang et al., 2015) to characterize genetic relatedness within, and differentiation
between, populations. We looked for correlations between the population average PC value and
elevation for the first 4 PCs to identify population structure associated with elevation.

We tested for divergent selection across populations on short stamen number, i.e., selection for more
stamens at high elevation and/or fewer stamens at low elevation than expected due to genetic drift,
with Qpc using the *quaint* R package (Josephs et al., 2019). Qpc is an extension of Qst-Fst analysis that
uses genetic PCs to estimate additive genetic variance within and between populations (Josephs et al.,
2019), and tests for phenotypic differentiation due to selection beyond that expected from neutral
evolution. This differs from the multiple regression test for the drift hypothesis described above because
Qpc explicitly considers among population differentiation. Qpc incorporates genetic relatedness within
and among populations through a kinship matrix, rather than a single measure of population
differentiation as in Qst-Fst. Qpc then tests for excess phenotypic divergence along major axes of
genetic relatedness (principal components of the kinship matrix) rather than excess phenotypic
divergence between populations as in Qst-Fst (Josephs et al., 2019). The input kinship matrix was
generated from a random subset of 50,000 SNPs that had no missing data. This kinship matrix is slightly

different from the kinship matrix used for genetic PCA because Qpc requires a kinship matrix
standardized across all loci, not each locus individually as in plink, and *quaint* is not capable of dealing
with missing data.

Using genome wide association studies (GWAS) to identify loci associated with short stamen loss

We performed GWAS to find genomic regions associated with short stamen number. Mean short
stamen number was not normally distributed (Shapiro-Wilk w=0.85, p = 3.27×10^{-6}; Fig. S1). Arcsine
transformation made the distribution closer to normal but still skewed (Shapiro-Wilk w = 0.91, p =
2.72×10^{-4}; Fig. S1). Phenotypic and genotypic data were merged in plink v1.9 (Chang et al., 2015). GWAS
was conducted in gemma v0.98.4 with the Wald hypothesis test for each SNP against the alternate allele
and a centered kinship matrix to account for population structure (Zhou & Stephens, 2012). The output
was visualized with the *qqman* (Turner, 2018) and *ggplot2* (Wickham et al., 2019) R packages. SNPs were
further investigated if they passed a significance threshold specified using a false discovery rate (FDR) <
0.05 determined with p.adjust in R.

Additional GWAS were conducted due to the remaining skew in the arcsine-transformed data. We ran a
GWAS with untransformed mean short stamen number and another with mean short stamen number
coded as a binary trait. In the latter, genotypes with no short stamen loss (mean short stamen number =
2) were coded as controls and genotypes with any amount of short stamen loss (mean short stamen
number < 2) were coded as cases. Finally, a fourth GWAS was conducted on the subset of genotypes
that experience short stamen loss (i.e., only the genotypes with mean short stamen number < 2;
Shapiro-Wilk w=0.93, p=0.0132) to ameliorate the zero-inflated-like distribution in the other continuous
GWAS caused by excess genotypes with a mean short stamen number equal to 2 (Fig. S1).

We identified genetic regions with SNPs associated with short stamen number in at least two GWAS to
find candidate regions for further study ("shared" SNPs). We identified overlapping regions associated
with stamen loss among the four GWAS analyses by creating a 1kb window centered on each SNP that
had an FDR adjusted p-value below 0.10 in any GWAS and searching for SNPs within that window that
were also below an FDR adjusted p-value of 0.10 in at least one other GWAS. We chose a more lenient
FDR of 0.10 for this analysis because we have higher confidence SNPs are associated with short stamen
number if they pass a significance threshold in multiple analyses. We chose a window of 1kb to include
regions flanking the associated SNP, although we recognize that *A. thaliana* genes can be larger than 1kb
and that linkage disequilibrium (LD) begins to drop off around 50kb in our individuals (Fig. S2), so this is
a conservative overlap criterion. SNPs in overlapping stamen loss associated regions were investigated
with the TAIR10 genome browser.

All figures were created with *ggplot2* (Wickham et al., 2019) and *ggpubr* (Kassambara, 2023) unless
otherwise noted.

Results

Short stamen loss is more common at low elevation

Mean short stamen number, when measured in a common garden, increases with the elevation of the
source population until reaching close to two stamens around 1300m (β = 4.84×10^{-4}, p = 0.004; γ = -
4.73×10^{-7}, p = 0.137; Fig. 2A; Table S3). Thus, short stamen retention is more common in populations
from higher elevations and loss is more common at low elevations in the Spanish Pyrenees. This

elevational cline is similar to the previously identified latitudinal cline where short stamen loss is more
common at southern latitudes (Royer et al., 2016).

Nucleotide diversity decreases with elevation, but there is no evidence that variation in effective
population size maintains the cline in short stamen number

Consistent with expectations that *A. thaliana* in the Pyrenees experienced repeated founder effects
after the last glaciation, high elevation populations have less nucleotide diversity than low elevation
populations (β = -1.81x10^{-6}, p < 0.001; Fig. 2B; Table S3). The three populations with the lowest
nucleotide diversity are high elevation populations BIS, VIE, and PAN. The results are consistent with
prior findings that genetic diversity and effective population size decrease with elevation in this region
(Gomaa et al., 2011; Montesinos et al., 2009). Population nucleotide diversity is comparable to
nucleotide diversity of *A. thaliana* across the Iberian Peninsula and the European range (Alonso-Blanco
et al., 2016).

We hypothesized a negative relationship between nucleotide diversity and short stamen number if
variation in effective population size was maintaining the short stamen cline, but nucleotide diversity did
not significantly predict short stamen number when accounting for elevation (β = -138.5, p =0.167; Fig.
2D; Table S3). Further, the relationship between short stamen number and elevation is similar whether
nucleotide diversity is included or not (Fig. 2C, compare to Fig. 2A; Table S3). These results suggest the
short stamen number cline is not maintained by variation in effective population size but may be
maintained by local adaptation, correlated responses to selection, or shared evolutionary history.

There is no evidence for divergent selection on short stamen number along the elevational cline

An elevational cline in short stamen number could result from local adaptation to different optima at
high and low elevations. To demonstrate evidence of local adaptation resulting from divergent selection,
we would need to show that the cline in short stamen number is stronger than what could be generated
by neutral evolution alone. Genetic structure, here measured using genetic PCA, shows a strong pattern
of genetic differentiation by elevation; both PC1 (r = -0.75, p = 9.22x10^{-4}) and PC2 (r = -0.54, p = 0.03) are
correlated with elevation (Fig. 3A). Neither PC3 nor PC4 are correlated with elevation (Fig. S3). The BOS
population is an outlier along both PC1 and PC2 (Fig. 3A). This aligns with geographic information
because BOS is located on a northern face of the Spanish Pyrenees while the other populations are on
southern faces, and prior work identified BOS in the Northwestern genetic cluster of the Iberian
Peninsula while the other 15 populations belong to the Northeast cluster or are classified as mixed
(Castilla et al., 2020). The PCA was conducted a second time after removing the BOS population (Fig. 3B).
These results continue to show an elevational cline along PC1 (r = -0.92, p = 1.15x10^{-6}), though PC2 (r = -
0.17, p = 0.55) separates the four highest elevation populations from each other with the closest
clustering of PAN and VIE. The strong genetic differentiation across elevation we observed means that
we have low power to test for even greater differentiation in stamen loss as evidence for divergent
selection. Not surprisingly, we did not find evidence for divergent selection on short stamen number
with Qpc. Differences in short stamen number between high and low elevation populations were not
larger than could be explained by neutral evolution (Fig. 3C, p=0.257).

Few SNPs across the genome are associated with short stamen number

We conducted GWAS to identify loci associated with short stamen number. Because short stamen
number was not normally distributed (Fig. S1), we conducted GWAS on arcsine-transformed mean short
stamen number, untransformed mean short stamen number, a binary trait of whether or not individuals
lacked any short stamens, and an untransformed subset of only genotypes that lack some short
stamens. Each GWAS found different SNPs associated with the trait. The qqplot for the arcsine-
transformed GWAS shows most p-values close to the 1:1 line between Expected and Observed (Fig. 4)
but the p-values in the binary GWAS deviate early from the 1:1 line, indicating a high false positive rate
in identifying SNPS associated with any short stamen loss (Fig. S4). In accordance with this observation,
the binary GWAS had 1,707 SNPs associated with short stamen loss at FDR < 0.05 while the other
analyses only had two or three (Figs. 4, S4). Only two SNPs are associated with arcsine-transformed
short stamen number (FDR < 0.05; Fig. 4). The SNP at Chr3:2942726 is within a FAD/NAD(P)-binding
oxidoreductase family protein (AT3G09580) that localizes in the chloroplast (Tomizioli et al., 2014). The
SNP at Chr5:13458838 is not within a gene; the closest gene, PICALM3 (AT5G35200), is just over 3kb
away.

We identified top candidates for loci associated with short stamen number by identifying 1kb windows
that have SNPs associated with short stamen number in multiple GWAS ("shared" SNPs). To do so, we
calculated 1kb windows centered on all SNPs associated with short stamen number (FDR < 0.10) and
searched for other SNPs that fit the same criteria and fell within each window. Twenty SNPs were
associated with short stamen number in more than one GWAS from 9 different 1kb windows (Table S4).
None of the SNPs fall within genes with known stamen function. However, a shared SNP at
Chr3:2253161 is approximately 40kb away, thus within LD (Fig. S2), from *FHA2* (AT3G07220), a
SMAD/FHA domain containing protein involved in stamen development (Ahn et al., 2013; Gu et al.,
2020). Mutating *FHA2* proteins can cause plants to have fewer stamens, though flowers sometimes lose
long stamens and sometimes lose short stamens (Ahn et al., 2013). The two SNPs associated with
arcsine-transformed short stamen number (FDR < 0.05) are included in the shared SNPs (Fig. 4; Table
S4).

All of the SNPs associated with variation in short stamen number fall outside the 95% credible intervals
of previously identified QTL for stamen loss using recombinant inbred lines with parents from the
extremes of the latitudinal stamen loss gradient (Royer et al., 2016). The closest intersection is on
chromosome 5, where SNPs associated with short stamens are approximately 2,000kb further into the
chromosome than the previously identified stamen loss QTL. This is beyond the start of LD decay we
estimated of 50kb (Fig. S2).

Discussion

In this study, we tested for an elevation cline in short stamen loss. We found that short stamen number
increased with elevation (Fig. 2A), similar to the latitudinal cline observed by Royer et al. (2016). Short
stamen number is one of many traits that show an elevational cline in this region (Montesinos et al.,
2009; Montesinos-Navarro et al., 2011; Wolfe & Tonsor, 2014). Of these traits, days to bolting shows
similar elevational and latitudinal clines with delayed bolting at high elevations and northern latitudes
(Montesinos-Navarro et al., 2011; Stinchcombe et al., 2004). Thus, short stamen number joins a growing
database of traits with an elevational cline in the Spanish Pyrenees and a much smaller set of traits with
parallel elevational and latitudinal clines. These clines could be maintained by effective population sizes
because both high elevation populations and northern latitude populations may have undergone

repeated bottlenecks that reduced effective population size since the last glaciation (Beck et al., 2008;
François et al., 2008; Hughes et al., 2006; Lewandowska-Sabat et al., 2010). While the Pyrenees do
include putative glacial refugia regions (Médail & Diadema, 2009), none of the 16 populations analyzed
here have been classified as relict lineages that predate the last glaciation (Castilla et al., 2020).
Selection could also maintain similar latitudinal and elevational clines because environmental factors
such as temperature often vary in the same away across latitude and elevation.

We then tested if the cline in short stamen loss was maintained by variation in effective
population size if a combination of low genetic variation and less effective selection in high
elevation populations hinders a response to selection against short stamens. We found
variation in genetic diversity across elevations consistent with expectations from repeated
founder effects during range expansion after the last glaciation and/or low gene flow (Fig.
2B). However, the multiple regression suggests that variation in genetic diversity is not
maintaining the short stamen number cline (Fig. 2C and 2D). Instead, the multiple regression
suggests that elevation can explain both variation in genetic diversity and in stamen number.
It is important to note that our clines show populations with low genetic diversity maintain
two stamens. This is opposite to the hypothesis that genetic drift would cause trait loss which
has been shown in other systems (Eckert et al., 1996; Lahti et al., 2009).

Next, we tested if the cline in short stamen loss was maintained by local adaptation using Qpc, an
extension of Qst-Fst (Josephs et al., 2019). We found no evidence of local adaptation maintaining the
cline in short stamen number (Fig. 3C). One caveat to our findings is that Qst-Fst struggles to identify
selection when there is large genetic differentiation between populations or weak selection that results
in similar values for Qst and Fst (Whitlock & Guillaume, 2009). This is also potentially an issue for Qpc in
this study as there is a great deal of genetic differentiation between high and low elevation populations
(Fig. 3A). However, the Qst-Fst approach has identified local adaptation contributing to elevational
clines in leaf succulence and specific leaf area in 14 populations of *A. thaliana* from the Swiss Alps where
there is also a great deal of genetic differentiation (Luo et al., 2015). Qpc has also previously been used
to identify local adaptation in 249 natural *A. thaliana* accessions from across the European native range;
local adaptation was identified in initial size, growth rate at 16°C, and temperature response (Clauw et
al., 2022). Ultimately, reciprocal transplant studies across elevations in the field will provide the
strongest evidence for or against local adaptation.

Finally, we used GWAS to identify loci associated with short stamen number. We identified 20 SNPs
associated with short stamen number in multiple GWAS (FDR < 0.10). These SNPs lie outside the QTL for
short stamens that were previously identified in a cross between a northern European and southern
European accession (Royer et al., 2016). The effect sizes of the Royer et al. (2016) QTL (range from 0.05
to 0.15) and our loci (Table S4) are comparable, suggesting that we would be able to identify these QTL
if they segregated at intermediate frequency in the populations used here. Instead, the populations
studied here may be fixed for the southern allele identified in Royer et al (2016). However, our GWAS
effect sizes may be inflated due to winner's curse (Göring et al., 2001; Josephs et al., 2017). In addition,
the QTL identified in Royer et al. (2016) were epistatic, which could make them harder to detect through
standard GWAS approaches. Overall, our results suggest that the latitudinal cline in short stamen
number across Europe has a different genetic basis than the elevational cline in short stamen number
observed in the Pyrenees.

While the short stamen loss clines are similar between latitude and elevation, the lack of overlap
between the associated loci suggests that short stamen loss is caused by mutations at different genes in
different geographic regions. These results are consistent with other findings that the chromosome
regions underlying the same phenotypic cline can differ by region. For example, elevational clines of *A.*
thaliana in different geographic regions show delayed flowering at higher elevations but the genetic
basis of this cline varies by global region (Gamba et al., 2022). Additionally, parallel clines in flowering
time in *A. thaliana* in the Cape Verde Islands arose from mutations at different genes (FRI and FLC)
(Fulgione et al., 2022).

In conclusion, short stamen loss is occurring in low elevation populations more than in high elevation
populations. Our results suggest that retention of two short stamens in high elevation populations are
not explained by reduced effective population sizes or by local adaptation to different optima in high
and low elevation populations. Thus, the evolutionary mechanisms underlying variation in short stamen
number are excitingly complex and deserve further study. A third hypothesis is that correlated
responses to selection may maintain the cline in short stamen number (Futuyma, 2010; Lande, 1979).
However, correlated responses to selection on other locally adapted traits could result in weak evidence
for divergent selection; we did not find this. Future work should better characterize direct selection and
correlated responses to selection on short stamens in the field by measuring fitness of plants with
natural variation in short stamen number and experimental manipulation of stamens, like that done by
Royer et al. (2016), in the field. Further, refining the short stamen number QTL to candidate genes and
comparing to genetic regions underlying locally adapted traits could uncover the role of linked or
pleiotropic genes leading to correlated responses to selection. These additional studies across both
latitudinal and elevational gradients will characterize the interplay of direct selection, correlated
responses to selection, and genetic drift in trait loss and identify parallel evolutionary forces at play
across environmental contexts.

Figures:

Short Stamen

Figure 1: Short stamen number increases with elevation. A) Each point represents the seed collection
site for 16 populations of *A. thaliana* across the Spanish Pyrenees. The map coloring indicates elevation
from green at low elevation to white at high elevation. **B)** Natural variation in short stamen number.
Flowers have 0 (left), 1 (center), or 2 (right) short stamens. Photos by Frances Whalen.

Figure 2: Effective population size does not explain retention of short stamens at high elevation. A)
Mean short stamen number shows a quadratic elevational cline. **B)** Elevation strongly predicts
population mean pairwise nucleotide diversity, our measure of effective population size. **C)** The
residuals of the model of short stamen number regressed on nucleotide diversity then regressed on
elevation and **D)** the residuals of the model of short stamen number regressed on quadratic elevation
then regressed on nucleotide diversity. In C and D, statistics from the full model (r^2 = 0.50; Table S3) are
displayed on the figures. In all panels, the color of each point represents the population elevation, black
lines are the regression, and grey lines are 95% confidence intervals.

Figure 3: Genetic variation is correlated with elevation, but there is no evidence for divergent selection on short stamen number. A) Genetic PCA for PC1 and PC2 from 1,858,706 SNPs. **B)** Genetic PCA for PC1 and PC2 with BOS individuals excluded. **C)** Qpc results showing the relationship between mean short stamen number and genetic differentiation (black line, *p*=0.26) is within expectations due to neutral evolution (grey shading). The fill color of each point represents the population elevation. Panels A and C used slightly different input data (see Methods).

Figure 4. Few SNPs are associated with short stamen number. Manhattan plot **(A)** and QQ plot **(B)** for
arcsine-transformed mean short stamen number. The yellow dashed line indicates significance at p=0.05
after Bonferroni correction. Blue points are significant below a FDR of 0.10. Pink labelled points are
significant below a FDR of 0.05. Points with a green outline are shared between at least two short
stamen GWAS below FDR 0.10 (n = 20). The black bars on the x axis are Bayes 95% credible intervals for
short stamen number QTL identified by Royer et al. (2016).

References

Ahn, E.-R., Cho, H.-K., & Pai, H.-S. (2013). The forkhead-associated domain 2 (FHA2) in Arabidopsis

plays a role in plant fertility by regulating stamen development. *Planta*, *237*(4), 1015–1023.

https://doi.org/10.1007/s00425-012-1815-7

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M. M., Cao, J.,

Chae, E., Dezwaan, T. M. M., Ding, W., Ecker, J. R. R., Exposito-Alonso, M., Farlow, A., Fitz,

J., Gan, X., Grimm, D. G. G., Hancock, A. M. M., Henz, S. R. R., Holm, S., … Zhou, X. (2016).

1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell*, *166*(2),

481–491. https://doi.org/10.1016/j.cell.2016.05.063

Andrews, S. (2019). *FastQC* (0.11.9) [Computer software]. Babraham Bioinformatics.

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/JSS.V067.I01

Beck, J. B., Schmuths, H., & Schaal, B. A. (2008). Native range genetic variation in Arabidopsis thaliana

is strongly geographically structured and reflects Pleistocene glacial dynamics. *Molecular*

Ecology, *17*(3), 902–915. https://doi.org/10.1111/j.1365-294X.2007.03615.x

Berardini, T. Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., & Huala, E. (2015). The

arabidopsis information resource: Making and mining the "gold standard" annotated reference

plant genome. *Genesis*, *53*(8), 474–485. https://doi.org/10.1002/dvg.22877

Caballero, A. (1994). Developments in the prediction of effective population size. *Heredity*, *73*(6), Article

6. https://doi.org/10.1038/hdy.1994.174

Castilla, A. R., Méndez-Vigo, B., Marcer, A., Martínez-Minaya, J., Conesa, D., Picó, F. X., & Alonso-

Blanco, C. (2020). Ecological, genetic and evolutionary drivers of regional genetic differentiation

in Arabidopsis thaliana. *BMC Evolutionary Biology*, *20*(1). https://doi.org/10.1186/s12862-020-

01635-2

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-

generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), 7.

https://doi.org/10.1186/S13742-015-0047-8

Charlesworth, B. (2009). Effective population size and patterns of molecular evolution and variation.

Nature Reviews Genetics, *10*(3), 195–205. https://doi.org/10.1038/nrg2526

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., & Ruden, D.

M. (2012). A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff. *Fly*, *6*(2), 80–92. https://doi.org/10.4161/fly.19695

Clark, R. M., Schweikert, G., Toomajian, C., Ossowski, S., Zeller, G., Shinn, P., Warthmann, N., Hu, T.

T., Fu, G., Hinds, D. A., Chen, H., Frazer, K. A., Huson, D. H., Schölkopf, B., Nordborg, M.,

Rätsch, G., Ecker, J. R., & Weigel, D. (2007). Common sequence polymorphisms shaping genetic

diversity in Arabidopsis thaliana. *Science*, *317*(5836), 338–342.

https://doi.org/10.1126/science.1138632

Clauw, P., Kerdaffrec, E., Gunis, J., Reichardt-Gomez, I., Nizhynska, V., Koemeda, S., Jez, J., &

Nordborg, M. (2022). Locally adaptive temperature response of vegetative growth in Arabidopsis

thaliana. *eLife*, *11*, e77913. https://doi.org/10.7554/eLife.77913

Colautti, R. I., & Lau, J. A. (2015). Contemporary evolution during invasion: Evidence for differentiation,

natural selection, and local adaptation. *Molecular Ecology*, *24*(9), 1999–2017.

https://doi.org/10.1111/mec.13162

Conner, J. K., Rice, A. M., Stewart, C., & Morgan, M. T. (2003). Patterns and mechanisms of selection

on a family-diagnostic trait: Evidence from experimental manipulation and lifetime fitness

selection gradients. *Evolution*, *57*(3), 480–486. https://doi.org/10.1111/j.0014-

3820.2003.tb01539.x

Daco, L., Colling, G., & Matthies, D. (2021). Altitude and latitude have different effects on population

characteristics of the widespread plant Anthyllis vulneraria. *Oecologia*, *197*(2), 537–549.

https://doi.org/10.1007/s00442-021-05030-6

Dorken, M. E., Neville, K. J., & Eckert, C. G. (2004). Evolutionary vestigialization of sex in a clonal plant: Selection versus neutral mutation in geographically peripheral populations. *Proceedings of the Royal Society B: Biological Sciences*, *271*(1555), 2375–2380. https://doi.org/10.1098/rspb.2004.2875

Durvasula, A., Fulgione, A., Gutaker, R. M., Alacakaptan, S. I., Flood, P. J., Neto, C., Tsuchimatsu, T., Burbano, H. A., Picó, F. X., Alonso-Blanco, C., & Hancock, A. M. (2017). African genomes illuminate the early history and transition to selfing in *Arabidopsis thaliana. Proceedings of the National Academy of Sciences*, *114*(20), 5213–5218. https://doi.org/10.1073/pnas.1616736114

Eckert, C. G., Manicacci, D., & Barrett, S. C. H. (1996). Genetic Drift and Founder Effect in Native Versus Introduced Populations of an Invading Plant, Lythrum Salicaria (lythraceae). *Evolution*, *50*(4), 1512–1519. https://doi.org/10.1111/j.1558-5646.1996.tb03924.x

Endress, P. K. (1992). Evolution and Floral Diversity: The Phylogenetic Surroundings of Arabidopsis and Antirrhinum. *International Journal of Plant Sciences*, *153*(3), S106–S122.

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048. https://doi.org/10.1093/bioinformatics/btw354

Fong, D. W., Kane, T. C., & Culver, D. C. (1995). VESTIGIALIZATION AND LOSS OF NONFUNCTIONAL CHARACTERS. *Annual Review of Ecology and Systematics*, *26*(1), 249–268. https://doi.org/10.1146/annurev.es.26.110195.001341

François, O., Blum, M. G. B., Jakobsson, M., & Rosenberg, N. A. (2008). Demographic History of European Populations of Arabidopsis thaliana. *PLOS Genetics*, *4*(5), e1000075. https://doi.org/10.1371/journal.pgen.1000075

Fulgione, A., Neto, C., Elfarargi, A. F., Tergemina, E., Ansari, S., Göktay, M., Dinis, H., Döring, N., Flood, P. J., Rodriguez-Pacheco, S., Walden, N., Koch, M. A., Roux, F., Hermisson, J., & Hancock, A. M. (2022). Parallel reduction in flowering time from de novo mutations enable

evolutionary rescue in colonizing lineages. *Nature Communications*, *13*(1), 1461.

https://doi.org/10.1038/s41467-022-28800-z

Futuyma, D. J. (2010). EVOLUTIONARY CONSTRAINT AND ECOLOGICAL CONSEQUENCES.

Evolution, *64*(7), 1865–1884. https://doi.org/10.1111/J.1558-5646.2010.00960.X

Gamba, D., Lorts, C., Haile, A., Sahay, S., Lopez, L., Xia, T., Kulesza, E., Elango, D., Kerby, J., Yifru,

M., Bulafu, C. E., Wondimu, T., Glowacka, K., & Lasky, J. R. (2022). *The genomics and*

physiology of abiotic stressors associated with global elevation gradients in Arabidopsis thaliana

(p. 2022.03.22.485410). bioRxiv. https://doi.org/10.1101/2022.03.22.485410

Gomaa, N. H., Montesinos-Navarro, A., Alonso-Blanco, C., & Picó, F. X. (2011). Temporal variation in

genetic diversity and effective population size of Mediterranean and subalpine Arabidopsis

thaliana populations. *Molecular Ecology*, *20*(17), 3540–3554. https://doi.org/10.1111/j.1365-

294X.2011.05193.x

Göring, H. H. H., Terwilliger, J. D., & Blangero, J. (2001). Large Upward Bias in Estimation of Locus-

Specific Effects from Genomewide Scans. *The American Journal of Human Genetics*, *69*(6),

1357–1369. https://doi.org/10.1086/324471

Gu, B.-W., Tan, L.-M., Zhang, C.-J., Hou, X.-M., Cai, X.-W., Chen, S., & He, X.-J. (2020). FHA2 is a

plant-specific ISWI subunit responsible for stamen development and plant fertility. *Journal of*

Integrative Plant Biology, *62*(11), 1703–1716. https://doi.org/10.1111/jipb.12945

Hughes, P. D., Woodward, J. C., & Gibbard, P. L. (2006). Quaternary glacial history of the Mediterranean

mountains. *Progress in Physical Geography: Earth and Environment*, *30*(3), 334–364.

https://doi.org/10.1191/0309133306pp481ra

Josephs, E. B., Berg, J. J., Ross-Ibarra, J., & Coop, G. (2019). *Detecting Adaptive Differentiation in*

Structured Populations with Genomic Data and Common Gardens.

https://doi.org/10.1534/genetics.118.301786

Josephs, E. B., Stinchcombe, J. R., & Wright, S. I. (2017). What can genome-wide association studies tell

us about the evolutionary forces maintaining genetic variation for quantitative traits? *New*

Phytologist, *214*(1), 21–33. https://doi.org/10.1111/nph.14410

Kassambara, A. (2023). *ggpubr: "ggplot2" Based Publication Ready Plots* (R package version 0.6.0)

[Computer software]. <https://CRAN.R-project.org/package=ggpubr>.

Kooyers, N. J., Greenlee, A. B., Colicchio, J. M., Oh, M., & Blackman, B. K. (2015). Replicate altitudinal

clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual Mimulus

guttatus. *New Phytologist*, *206*(1), 152–165. https://doi.org/10.1111/nph.13153

Korunes, K. L., & Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and divergence in

the presence of missing data. *Molecular Ecology Resources*. https://doi.org/10.1111/1755-

0998.13326

Krueger, F. (2019). *Trim Galore* (0.6.5) [Computer software]. Babraham Bioinformatics.

https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

Lahti, D. C., Johnson, N. A., Ajie, B. C., Otto, S. P., Hendry, A. P., Blumstein, D. T., Coss, R. G.,

Donohue, K., & Foster, S. A. (2009). Relaxed selection in the wild. *Trends in Ecology and*

Evolution, *24*(9), 487–496. https://doi.org/10.1016/j.tree.2009.03.010

Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution , Applied to Brain: Body Size

Allometry. *Evolution*, *33*(1), 402–416.

Lenth, R. V. (2021). *emmeans: Estimated marginal Means, aka Least Squares Means* (R package version

1.6.2-1) [Computer software]. https://CRAN.R-project.org/package=emmeans

Lewandowska-Sabat, A. M., Fjellheim, S., & Rognli, O. A. (2010). Extremely low genetic variability and

highly structured local populations of Arabidopsis thaliana at higher latitudes. *Molecular*

Ecology, *19*(21), 4753–4764. https://doi.org/10.1111/j.1365-294X.2010.04840.x

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Luo, Y., Widmer, A., & Karrenberg, S. (2015). The roles of genetic drift and natural selection in

quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana. *Heredity*,

114(2), 220–228. https://doi.org/10.1038/hdy.2014.89

Médail, F., & Diadema, K. (2009). Glacial Refugia Influence Plant Diversity Patterns in the

Mediterranean Basin. *Journal of Biogeography*, *36*(7), 1333–1345.

Montesinos, A., Tonsor, S. J., Alonso-Blanco, C., & Picó, F. X. (2009). Demographic and genetic

patterns of variation among populations of Arabidopsis thaliana from contrasting native

environments. *PLoS ONE*, *4*(9). https://doi.org/10.1371/journal.pone.0007213

Montesinos-Navarro, A., Picó, F. X., & Tonsor, S. J. (2012). CLINAL VARIATION IN SEED TRAITS

INFLUENCING LIFE CYCLE TIMING IN *ARABIDOPSIS THALIANA*. *Evolution*, *66*(11),

3417–3431. https://doi.org/10.1111/j.1558-5646.2012.01689.x

Montesinos-Navarro, A., Wig, J., Xavier Pico, F., & Tonsor, S. J. (2011). Arabidopsis thaliana

populations show clinal variation in a climatic gradient associated with altitude. *New Phytologist*,

189(1), 282–294. https://doi.org/10.1111/j.1469-8137.2010.03479.x

R Core Team. (2021). *R: A language and environment for statistical computing* [Computer software]. R

Foundation for Statistical Computing. https://www.R-project.org/.

Royer, A. M., Kremer, C., George, K., Pérez, S. G., Schemske, D. W., & Conner, J. K. (2016).

Incomplete loss of a conserved trait: Function, latitudinal cline, and genetic constraints.

Evolution, *70*(12), 2853–2864. https://doi.org/10.1111/evo.13096

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: A model for studying the genetic and

evolutionary basis of morphological adaptation in plants. *Annals of Botany*, *107*(9), 1433–1443.

https://doi.org/10.1093/aob/mcr023

Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., Purugganan,

M. D., & Schmitt, J. (2004). A latitudinal cline in flowering time in *Arabidopsis thaliana*

modulated by the flowering time gene *FRIGIDA*. *Proceedings of the National Academy of*

Sciences, *101*(13), 4712–4717. https://doi.org/10.1073/pnas.0306401101

Tang, C., Toomajian, C., Sherman-Broyles, S., Plagnol, V., Guo, Y. L., Hu, T. T., Clark, R. M.,

Nasrallah, J. B., Weigel, D., & Nordborg, M. (2007). The evolution of selfing in Arabidopsis

thaliana. *Science*, *317*(5841), 1070–1072.

https://doi.org/10.1126/SCIENCE.1143153/SUPPL_FILE/TANG.SOM.REVISION.1.PDF

Tomizioli, M., Lazar, C., Brugière, S., Burger, T., Salvi, D., Gatto, L., Moyet, L., Breckels, L. M., Hesse,

A.-M., Lilley, K. S., Seigneurin-Berny, D., Finazzi, G., Rolland, N., & Ferro, M. (2014).

Deciphering Thylakoid Sub-compartments using a Mass Spectrometry-based Approach *.

Molecular & Cellular Proteomics, *13*(8), 2147–2167. https://doi.org/10.1074/mcp.M114.040923

Turner, S. D. (2018). qqman: An R package for visualizing GWAS results using Q-Q and manhattan

plots. *Journal of Open Source Software*, *3*(25), 731. https://doi.org/10.21105/JOSS.00731

Van der Auwera, G., & O'Connor, B. (2020). *Genomics in the Cloud: Using Docker, GATK, and WDL in*

Terra (1st Edition). O'Reilly Media.

Walsh, B., & Blows, M. W. (2009). Abundant genetic variation + strong selection = multivariate genetic

constraints: A geometric view of adaptation. *Annual Review of Ecology, Evolution, and*

Systematics, *40*, 41–59. https://doi.org/10.1146/annurev.ecolsys.110308.120232

Waterman, R., Sahli, H., Koelling, V. A., Karoly, K., & Conner, J. K. (2023). Strong evidence for

positive and negative correlational selection revealed by recreating ancestral variation. *Evolution*,

77(1), 264–275. https://doi.org/10.1093/evolut/qpac001

Whitlock, M. C., & Guillaume, F. (2009). Testing for Spatially Divergent Selection: Comparing QST to

FST. *Genetics*, *183*(3), 1055–1063. https://doi.org/10.1534/genetics.108.099812

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,

Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J.,

Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of*

Open Source Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wolfe, M. D., & Tonsor, S. J. (2014). Adaptation to spring heat and drought in northeastern Spanish

Arabidopsis thaliana. *New Phytologist*, *201*(1), 323–334. https://doi.org/10.1111/nph.12485

Yoshizawa, M., Yamamoto, Y., O'Quin, K. E., & Jeffery, W. R. (2012). Evolution of an adaptive

627   behavior and its sensory receptors promotes eye regression in blind cavefish. *BMC Biology*,

628   *10*(108). https://doi.org/10.1186/1741-7007-11-82

629   Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association studies.

630   *Nature Genetics 2012 44:7*, *44*(7), 821–824. https://doi.org/10.1038/ng.2310

631